

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/125860/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Liu, XM, Liu, Y, Yu, S, Jiang, LM, Song, B and Chen, X 2019. Potential immunomodulatory effects of SCAP on Treg conversion in tissue regeneration for regenerative endodontic treatment. *International Endodontic Journal* 52 (12) , pp. 1758-1767. 10.1111/iej.13197

Publishers page: <http://dx.doi.org/10.1111/iej.13197>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Figure S1 Primary culture and characterization of SCAP

(a) Morphologically, SCAP was spindle-shaped under the inverted phase contrast microscope. (b) SCAP formed Alizarin Red S positive mineralized nodes after osteogenic induction. (c) SCAP formed Oil Red O positive lipid clusters after adipogenic induction. (d) Flow cytometric analysis showed SCAP expressed mesenchymal stem cell surface markers (CD73, CD90, CD105 and CD146), but failed to express hematopoietic stem cell surface markers (CD31 and CD34). Scale bars = 100 μm (a, b); and 200 μm (c).