

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/126657/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Lewis, Catrin , Roberts, Neil, Gibson, Samuel and Bisson, Jonathan 2020. Dropout from psychological therapies for post-traumatic stress disorder (PTSD) in adults: systematic review and meta-analysis. *European Journal of Psychotraumatology* 11 (1) , 1709709. 10.1080/20008198.2019.1709709

Publishers page: <http://dx.doi.org/10.1080/20008198.2019.1709709>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Word count: 4281

Number of figures: 2

Number of tables: 4

(Editorial advice will be taken on which to include)

# Dropout from psychological therapies for post-traumatic stress disorder (PTSD) in adults: systematic review and meta-analysis

Catrin Lewis Ph.D. (Research Associate)<sup>1</sup>, Neil P. Roberts D.Clin.Psy. (Consultant Clinical Psychologist)<sup>1,3</sup>, Samuel Gibson (Psychology Placement Student)<sup>1</sup>,  
Jonathan I. Bisson D.M. (Professor in Psychiatry)<sup>1</sup>

<sup>1</sup>National Centre for Mental Health (NCMH), Division of Psychological Medicine and Clinical Neurosciences, Cardiff University School of Medicine, UK

<sup>2</sup>Cardiff Traumatic Stress Service, Cardiff & Vale University Health Board, UK

<sup>3</sup>Directorate of Psychology and Psychological Therapies, Cardiff & Vale University Health Board, UK

Work conducted at the Division of Psychological Medicine and Clinical Neurosciences, Cardiff University School of Medicine, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ

**Correspondence to:** Dr. Catrin Lewis – [LewisCE7@Cardiff.ac.uk](mailto:LewisCE7@Cardiff.ac.uk); Division of Psychological Medicine and Clinical Neurosciences, Cardiff University School of Medicine, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ

**Funding:** This work was unfunded.

## **Abstract**

**Background:** Despite the established efficacy of psychological therapies for post-traumatic stress disorder (PTSD) there has been little systematic exploration of dropout rates.

**Objective:** To ascertain rates of dropout across different modalities of psychological therapy for PTSD and to explore potential sources of heterogeneity.

**Method:** A systematic review of dropout rates from randomised controlled trials (RCTs) of psychological therapies was conducted. The pooled rate of dropout from psychological therapies was estimated and reasons for heterogeneity explored using meta-regression.

**Results:** The pooled rate of dropout from RCTs of psychological therapies for PTSD was 16% (95% CI 14% - 18%). There was evidence of substantial heterogeneity across studies. We found evidence that psychological therapies with a trauma-focus were significantly associated with greater dropout. There was no evidence of greater dropout from therapies delivered in a group format; from studies that recruited participants from clinical services rather than via advertisements; that included only military personnel/veterans; that were limited to participants traumatised by sexual traumas; that included a higher proportion of female participants; or from studies with a lower proportion of participants who were University educated.

**Conclusions:** Dropout rates from recommended psychological therapies for PTSD are high and this appears to be particularly true of interventions with a trauma focus. There is a need to further explore reasons for dropout and to look at ways of increasing treatment retention.

**Keywords:** Post-traumatic stress disorder; trauma; psychological; therapy; dropout; review

## Introduction

Post-Traumatic Stress Disorder (PTSD) is a debilitating psychiatric disorder with a lifetime prevalence of approximately 8% [1]. In addition to the requirement of exposure to a major traumatic event, the diagnostic criteria for PTSD specify the presence of symptoms including re-experiencing the traumatic event; avoiding reminders of the trauma; alterations in arousal and reactivity; and changes in cognition and mood [2].

Despite decades of research converging on support for the efficacy of psychological therapy for PTSD [3-5], we know remarkably little regarding dropout from these interventions [6-9]. Many psychological therapies have been applied to the treatment of PTSD and these have fundamentally different components and proposed active ingredients [10]. It follows that these variations may have some influence on differential rates of dropout. Despite this likelihood, there have been few attempts to systematically determine dropout rates from the psychological therapies commonly applied to the treatment of PTSD.

Among the evidence-based therapies for PTSD, a major distinction can be drawn between the therapies that focus on the traumatic event and those that aim to reduce traumatic stress symptoms without directly targeting the trauma memory or related thoughts, with the strongest evidence for the effect of those with a trauma-focus [3-5]. Trauma-focused Cognitive Behaviour Therapy (CBT) and Eye Movement Desensitisation and Reprocessing (EMDR) are currently recommended as first line interventions for PTSD [11-13]. These trauma-focused psychological therapies rely on confrontation of traumatic images, which can be difficult to tolerate and may result in the potential for greater dropout [14, 15]. Psychological therapies omitting a role for trauma-focused work may be more tolerable, potentially leading to better retention. However, there is evidence that the absence of a trauma-focus results in poorer outcomes [3-5].

The issue of treatment tolerability and symptom exacerbation resulting from trauma-focused psychological therapies has been one of contention in the literature [14, 16, 17]. It is uncertain whether dropout rates vary as a function of treatment modality or whether those with a trauma-focus are associated with poorer retention. To date, a small number of meta-analyses have compared drop-out rates across different modalities of psychological therapy for PTSD [4, 17-19]. One of these studies reported no differences between therapies with and without exposure-work, however the review is now dated and includes a far smaller number of studies than currently available [17]. Another review reported a trend towards greater dropout from exposure based treatment, but did not analyse this statistically [4]. A more recent review reported that dropout was not associated with trauma-focus, however studies comparing trauma-focused CBT to waitlist or usual care control groups were excluded, restricting the review to 42 studies [18]. A more recent review found no difference in dropout rates from therapies that included exposure work in comparison to those that did not, but the review only included twenty studies of US military veterans [19].

The aim of the current review was to ascertain rates of dropout across different modalities of psychological therapy and to determine whether some psychological therapies (especially those with a trauma-focus) were associated with higher rates of dropout than others. Since there is no agreed definition of dropout, we took the number of participants that had left the study at the point of post-treatment assessment as a proxy-indicator of dropout in order to allow the inclusion of data from a maximal number of studies. We also aimed to explore potential sources of heterogeneity among the included studies. Our overarching goal was to contribute to a refined understanding of dropout from psychological therapies for PTSD that will inform the development of treatment protocols that maximise retention.

## Method

### Selection Criteria

Data on drop-out was extracted from studies that had been identified for a review of the efficacy of psychological therapies for adults with PTSD, which was undertaken as part of an update of the International Society for Traumatic Stress Studies (ISTSS) Treatment Guidelines [12]. Both reviews had the same inclusion criteria. RCTs of any defined psychological therapy aimed at the reduction of PTSD-symptoms in comparison with a control group (e.g. usual care / waiting list); other psychological therapy; or psychosocial intervention (e.g. psychoeducation / relaxation training), were included. At least 70% of study participants were required to be diagnosed with PTSD with a duration of three-months or more, according to DSM or ICD criteria determined by clinician diagnosis or an established diagnostic interview. This review considered studies of adults aged 18 or over, only. There were no restrictions based on symptom-severity or trauma-type. The diagnosis of PTSD was required to be primary and studies of comorbid PTSD and substance use disorder were excluded, but there were no other restrictions based on co-morbidity. Studies were only included if they reported data on the number of participants that had dropped out of the study by the point of post-treatment assessment. If multiple studies reported data on the same participants, dropout data was only included once. We also excluded RCTs of single-session interventions.

### Search Strategy

A search was conducted by the Cochrane Collaboration, which updated a previously published Cochrane review with the same inclusion criteria, which was published in 2013 [3]. The updated search aimed to identify all RCTs related to the prevention and treatment of PTSD, published from January 2008 to the 31st May 2018, using the search terms PTSD or posttrauma\* or post-trauma\* or "post trauma\*" or "combat disorder\*" or "stress disorder\*". The searches included results from PubMed, PsycINFO, Embase and the Cochrane database of randomised trials. This produced a group of papers related to the psychological treatment of PTSD in adults. We checked reference lists of the

included studies. We searched the World Health Organization's, and the US National Institutes of Health's trials portals to identify additional unpublished or ongoing studies. We contacted experts in the field with the aim of identifying unpublished studies and studies that were in submission. A complementary search of the Published International Literature on Traumatic Stress (PILOTS) was also conducted.

### Data Extraction

Study characteristics and dropout data were extracted by two reviewers independently and in duplicate, using a form that had been pre-piloted. Since there is no agreed definition of dropout, taking the number of participants that had left the study at the point of post-treatment assessment allowed the inclusion of data from a maximal number of studies. Study authors were contacted to obtain missing data. Therapy classifications were agreed with the ISTSS treatment guidelines committee and posted on the ISTSS website to allow comment from the membership. Reasons for dropout and adverse events were not universally available or consistently reported by studies and it was not therefore possible to extract or meta-analyse this data.

### Risk of Bias Assessment

All included studies were assessed for risk of bias at study level, using Cochrane criteria [20]. This included: (1) sequence allocation for randomisation (the methods used for randomly assigning participants to the treatment arms and the extent to which this was truly random); (2) allocation concealment (whether or not participants or personnel were able to foresee allocation to a specific group); (3) assessor blinding (whether the assessor was aware of group allocation); (4) incomplete outcome data (whether missing outcome data was handled appropriately); (5) selective outcome reporting (whether reported outcomes matched with those that were pre-specified); and (6) any other notable threats to validity (for example, premature termination of the study). Two researchers

independently assessed each study and any conflicts were discussed with a third researcher with the aim of reaching a unanimous decision.

### Data Synthesis

Meta-analyses of proportion were conducted using the *metaprop* command in STATA version 13.1 [21]. The *metaprop* command pools proportions and uses the score statistic and the exact binomial method to compute 95% confidence intervals [22]. Data were pooled across all active psychological therapies. Sub-group analyses were also conducted to determine the dropout rate for each psychological therapy. A random effects model was chosen due to the heterogeneity across studies in terms of the inclusion and exclusion criteria of the studies; the populations from which the samples were drawn; the nature and duration of therapy; the predominant trauma type; and the mean age of participants.

Heterogeneity was assessed using both the  $I^2$  statistic (which indicates the proportion of the variance that is due to heterogeneity [23]) and visual inspection of the forest plots. To explore potential sources of heterogeneity, meta-regression was performed using the *metareg* function of STATA version 13.1 [21]. Meta-regression assesses the association between study-level variables and the effect size [22]. It was hypothesised that a number of study-level variables would result in higher rates of drop-out, these being: therapies having a trauma-focus (due to the possibility of these therapies being difficult for some participants to tolerate); therapies being delivered in a group-format (since drop out from group therapies has been found to be greater than from therapies delivered on an individual basis [18]); recruitment from clinical services rather than through advertisements (due to the likelihood of more severe symptoms and a possible tendency for these participants to be less motivated to engage in treatment); whether or not the participants were selected from military/veteran populations (due a greater likelihood of complex or severe PTSD); whether the trauma experienced by participants was sexual (due to the possibility of therapy being more difficult to tolerate); and the percentage of participants who were University educated (due to the possibility



that more educated participants are better able to grasp the concepts involved in therapy). To explore the possibility of publication bias, we constructed a funnel plot using data on dropout from all active therapy groups.

## **Results**

The original Cochrane review included 70 RCTs. The update search identified 5500 potentially eligible studies published since 2008. Abstracts were reviewed and full text copies obtained for 203 potentially relevant studies. Forty-four new RCTs met inclusion criteria for the review of efficacy that informed the ISTSS treatment guidelines [12], which resulted in a total of 114 RCTs that reported sufficient data on efficacy for inclusion in that review. Forty-six of the identified studies met the eligibility criteria and reported data on dropout, resulting in 115 studies for inclusion in this review. Figure 1 presents a flow diagram for study selection.

[FIGURE 1 HERE]

## **Study Characteristics**

Study characteristics are summarised in table 1. Twenty-eight defined psychological therapies were evaluated. Eight of these were broadly categorized as CBT with a Trauma Focus (CBT-T) delivered on an individual basis: Brief Eclectic Psychotherapy (BEP); Cognitive Processing Therapy (CPT); Cognitive Therapy (CT); Narrative Exposure Therapy (NET); Prolonged Exposure (PE); Reconsolidation of Traumatic Memories (RTM); Virtual Reality Exposure Therapy (VRE) and CBT-T (not based on a specific model). Thirteen other therapies delivered to individuals were evaluated: EMDR; CBT without a Trauma Focus; Present Centred Therapy (PCT); Supportive Counselling; Written Exposure Therapy; Observed and Experiential Integration (OEI); Interpersonal Psychotherapy; Psychodynamic Psychotherapy; REM Desensitisation; Emotional Freedom Technique (EFT); Dialogical Exposure Therapy (DET); Internet-based CBT; and Relaxation Training. There were six different types of group

therapy: Group CBT-T; Group Present Centred Therapy (PCT); Group and Individual CBT-T; Group Stabilising Treatment; Group Interpersonal Therapy; Group Supportive Counselling. There were also RCTs of couples CBT-T. There were six types of control group: psychoeducation; couples psychoeducation; internet-based psychoeducation; waitlist; treatment as usual; and minimal attention/symptom monitoring.

The number of randomised participants ranged from 10 to 360. Studies were conducted in Australia (9), Canada (2), China (2), Denmark (1), Germany (5), Iran (2), Israel (1), Italy (2), Japan (1), the Netherlands (5), Norway (1), Portugal (1), Romania (1), Rwanda (1), Spain (1), Sweden (3), Switzerland (1), Thailand (1), Turkey/Syria (1), Uganda (2), UK (10) and USA (62). Participants were traumatised by military trauma (27 studies), sexual assault or rape (11 studies), war/persecution (4 studies), road traffic accidents (6 studies), earthquakes (2 studies), childhood abuse (3 studies), political detainment (1 study), terrorism (2 studies), physical assault (2 studies), domestic abuse (4 studies), medical diagnoses/emergencies (4 studies), genocide (1 study) and organised violence (3 studies). The remainder included individuals traumatised by various different traumatic events. There were 27 studies of females only and 10 of only males; the percentage of females in the remaining studies ranged from 1.75% to 96%. The percentage with a University education ranged from 4% to 90%.

### Risk of Bias

Risk of bias assessments for the included studies are summarised in table 2. Fifty-two studies reported a method of sequence allocation judged to pose a "low" risk of bias; five reported a method with a "high" risk of bias; the remainder reported insufficient details and were, therefore, rated as "unclear". Forty-one studies reported methods of allocation concealment representing a "low" risk of bias; two a method with a "high" risk of bias; with the remainder rated as "unclear". The outcome assessor was aware of the participant's allocation in 11 of the included studies; it was unclear whether the outcome assessor was aware of group allocation in 20 studies; with the remainder using blind-raters or self-report questionnaires delivered in a way that could not be influenced by members of the

research team. Twenty-three studies were judged as posing a "high" risk of bias in terms of incomplete outcome data; 79 studies were felt to have dealt with dropouts appropriately ("low" risk of bias); it was unclear in the remaining studies. The majority of studies failed to reference a published protocol, resulting in an 'unclear' risk of selective reporting for 75 studies; risk of bias was judged as "high" in five studies and low in the remainder. Seventy of the included studies presented a "high" risk of bias in other areas, for example, in relation to sample size, baseline imbalances between groups, or other methodological shortfalls. We could not rule out potential researcher allegiance, since treatment originators were involved in the evaluation of their own intervention in many of the included studies.

[TABLE 2 HERE]

### Dropout

Across the different modalities of psychological therapy, dropout rates from individual studies ranged from 0%-65%. The pooled dropout rate from psychological therapies for PTSD was 16% (95% CI 14 – 18;  $k = 116$ ) with substantial heterogeneity across studies ( $I^2 = 77.3\%$ ). The dropout rate for each modality of psychological therapy is presented in table 3. The heterogeneity in dropout rates indicates differences that may be predicted by the variables entered into meta-regression.

[TABLE 3 HERE]

### Meta-regression

We found evidence that psychological therapies with a trauma-focus were significantly associated with greater dropout ( $\beta = 0.069$ ; CI 0.011- 0.127;  $P = 0.021$ ; dropout rate of 18% (95% CI 15-21%) from those with a trauma focus versus 14% (95% CI 10-18%) from those without a trauma focus). There was no evidence of greater dropout from therapies delivered in a group format; from studies that recruited participants from clinical services rather than via advertisements; that included only

military personnel/veterans; that included only participants traumatised by sexual traumas; from studies with a higher proportion of female participants; or from studies with a lower proportion of participants who were University educated .

[TABLE 4 HERE]

### Publication Bias

A funnel plot, which was constructed using data on dropout from all active therapy groups, did not show evidence of publication bias.

[FIGURE 2 HERE]

## Discussion

### **Main findings**

The pooled dropout rate from psychological therapies for PTSD was 16% (95% CI 14-18%). This is of a similar magnitude to a previous meta-analysis of 42 studies, which found an average dropout rate of 18% [18] using the definition of dropout given by the included studies. This is also similar to the dropout rate of 17.5% obtained from a meta-analysis of dropout from RCTs of psychotherapy for depression [24] that defined dropout as unexpected attrition among individuals who were randomized to a treatment but failed to complete it. It was considerably lower than the pooled dropout rate of 36% found by a more recent review of twenty studies of US military veterans [19]. This was in comparison to a pooled dropout rate from studies of veterans/military personnel in this review of 18% (95% CI 15-22%). This is likely to reflect the fact that the previous review included a variety of different study designs including naturalistic studies and used the definition of dropout given by the authors of individual studies.

There was no evidence of greater dropout from therapies delivered in a group format. This contradicts the findings of earlier reviews that found group delivery to be associated with a significant increase in dropout [18, 19]. This may be the result of more recent studies evaluating interventions

that have been optimised to increase retention. There was also no evidence of significantly greater dropout from studies that recruited participants from clinical services rather than via advertisements; that included only military personnel/veterans; that included only participants traumatised by sexual traumas; that included only female participants; and from studies with a lower proportion of participants who were University educated. Research looking at factors associated with dropout have yielded inconsistent findings [9, 25, 26]. Although the findings of the current review contradict some previous studies; they are in agreement with others. Inconsistencies may be the result of difference in study type and design; the types of interventions of interest and the degree to which they are protocolised; or may vary according to the populations of interest.

We found evidence that psychological therapies with a trauma-focus were significantly associated with greater dropout. This challenges the findings of previous, far smaller, meta-analyses, which found no significant differences in dropout rates from therapies with and without a trauma-focus [17, 19]. However, one of these studies found a significant difference between PCT (a non-trauma focused intervention) and a group of therapies that had a trauma-focus [18]. Our findings may be a result of the accumulated data available from a larger number of studies. Although there are many reasons for dropout from psychological therapies, this finding suggests that difficulties tolerating trauma-focused treatment may be one of these. Adverse events such as the prolonged exacerbation of existing symptoms (for example, an increased frequency of unwanted thoughts or nightmares) or the occurrence of new symptoms (for example anger or self-blame) may lead to dropout, yet there is a surprising scarcity of research exploring the issue [27]. Psychological therapy is traditionally perceived as safe, presenting a low risk of unwanted effects [28]. In reality, the estimated rate of reported side effects is between 3% and 15%, which is of a similar magnitude to that reported for pharmacotherapy [29]. However, it is often difficult to draw a distinction between adverse events and time-limited negative experiences inherent to the process of some psychological therapies. This includes the experience of distress provocation, which is inevitable in the process of trauma-focused work.

A survey of psychologists' attitudes to trauma focused intervention found that concerns about tolerability and dropout were among the main reasons that psychologists did not use trauma focused intervention, despite the compelling evidence supporting its use [30]. However, only a small number of studies have acknowledged or explored adverse events such as symptom worsening or its influence on dropout in relation to trauma focused therapy. This is surprising, given that symptom exacerbation has long since been documented in the treatment of PTSD [31, 32]. It also limits our ability to judge how well various therapies were tolerated by PTSD sufferers. An RCT of imagery rehearsal therapy for trauma-related nightmares found that all four participants who actively withdrew from the treatment group had experienced increased negative imagery effects, suggesting a direct relationship between an inability to tolerate the treatment and subsequent dropout [33, 34]. Conversely, a study of 76 individuals found that only 9-21% of participants showed reliable symptom exacerbation, and these individuals were no more likely to drop out of treatment prematurely [35]. Similarly, an RCT comparing cognitive therapy (without a trauma focus) to imaginal exposure found that symptom worsening affected 10% of participants, with a significantly greater number of these being in the imaginal exposure group, however, this between-group difference was no longer present at follow-up and rates of dropout were similar from both groups [34].

The studies included in this review usually failed to provide information on adverse events and contained few explanations for dropout, so it is difficult to ascertain why participants dropped out. It must be acknowledged that symptom improvement is a possible reason for dropout [36]. It follows that termination of treatment for this reason would be highest from the most effective treatments (i.e. those with a trauma-focus [3-5]). More transparent reporting of dropout is required to explore this further. Whatever the cause, dropout is a major health and societal concern, which may result in individuals failing to receive optimal treatment [37, 38].

## **Strengths and limitations**

The review followed Cochrane guidelines for the identification of relevant studies; data extraction; and risk assessment [23]. A wide range of psychological therapies for PTSD were considered, which included participants from different countries and backgrounds, who had been exposed to a variety of different traumatic events. Inevitably, there were some limitations. The majority of studies included in the review excluded individuals with comorbidities of substance dependence, psychosis, and severe depression, who may be more likely to drop out of treatment prematurely, as evidenced by particularly high rates of drop out from studies of participants with co-morbid alcohol dependency [39-41]. All included studies were published, resulting in the possibility of publication bias. However, a funnel plot constructed from the data did not show evidence of this being an issue.

Since there is no agreed conceptualisation of dropout, this review extracted and meta-analysed data on the number of participants that had left the study at the point of post-treatment assessment to allow the inclusion of data from a maximal number of studies. There may have been participants who completed a full course of therapy but failed to attend the post-treatment assessment. Equally, there may have been participants who failed to complete the course of treatment but attended the post-treatment assessment nonetheless. Although this may bias our findings, there are limitations to all methods that we could have adopted to conceptualise dropout.

The review relied on RCT evidence, which is both a strength and a limitation. The methodology may have excluded some potentially high quality sources of evidence, such as large observational studies and non-randomized controlled effectiveness studies [40], which could contribute to a more accurate overall assessment of dropout. It may be the case that dropout from clinical trials underestimates the true extent of dropout in routine clinical care on the basis that study teams are motivated to retain participants and often provide incentives for the completion of treatment. Equally, participants may have been more inclined to drop out on the basis of the additional demands of participation in a trial,

such as regular completion of research assessments. However, taking a broader approach would risk diluting higher quality sources of evidence with weaker ones. A major weakness was that reasons for dropout were not reported or were poorly reported by most studies and it was not possible to systematically extract and analyse this information.

## **Research Implications**

Bringing together the available evidence on dropout has always been problematic given that there is no agreed definition and studies have conceptualised the phenomenon differently. Agreeing a definition of dropout would advance the field by encouraging the reporting of data that is comparable across trials. A previous study that compared the application of four operational definitions of dropout (therapist judgment, failure to attend the last scheduled appointment, a median-split procedure, and failure to return to therapy after the intake appointment), found that the rate ranged from 17.6% to 53.1%, depending on the definition that was used [42]. It follows that a framework to guide the standardised collection and documentation of data related to dropout including information on adverse events, is needed. There is currently no theoretical concept to guide the evaluation and reporting of dropout and adverse events that occur during psychological therapy, which is needed and would include a standardised list of reasons for dropout. A first step would be for research ethics committees to mandate that future RCTs of psychological treatments routinely collect and report standardised data on dropout, including the reasons for it. When possible, studies should also report on the severity of symptoms at the point that participants drop out from therapy and whether any adverse events occurred [17]. Systematic reviews that analyse individual patient data in relation to dropout enable the application of a standardised definition across studies and would advance the field by moving beyond looking at associations between study-level variables and dropout. Only when we have sufficient knowledge on the reasons for dropout can we be sure that patients are receiving the best possible intervention.



## Clinical implications

Although we cannot be sure that the reasons for dropout are negative, the findings point to the need for careful assessment of the suitability of patients for trauma-focused work. Since there is evidence for the effect of many different modalities of psychological therapy [11-13], the evidence-base should be used to guide shared-decision making between patient and clinician [13]. Enhancing patient choice may improve retention on the basis that individuals are self-selecting treatment approaches that hold personal appeal. Whether or not this ultimately impacts retention and treatment outcomes, requires investigation. Since PTSD is a highly heterogeneous condition [43], a greater understanding of dropout has the potential to facilitate the targeted recommendation of existing evidence-based treatments to specific sub-groups of patients. Dropout is clearly a complex phenomenon, which may be best conceptualised as having a multi-faceted aetiology that is likely to vary across different therapies and diagnostic groups. A multi-factorial approach is likely to be required to reduce dropout, such as a stepped care approach that is personalised and addresses the various barriers to remaining in treatment [44]. Phased therapies have been developed with preparatory work to improve stability before trauma-focused work [45]. This approach has been found to result in improved outcomes and greater retention in trauma-focused CBT for PTSD [46]. Another option is the introduction of peer support, which has been shown to encourage participants to re-enter treatment and subsequently achieve significant clinical improvement [47].

## References

1. Kessler, R.C., *Posttraumatic stress disorder: the burden to the individual and to society*. Journal of Clinical Psychiatry., 2000. **61**(5): p. 4-14.
2. American Psychiatric Association, *Diagnostic and Statistical Manual of Mental Disorders (5th Edition)*. 2013, (4th ed. Text Revision). Washington, DC.
3. Bisson, J., et al., *Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults (Review)*. Cochrane Database of Systematic Reviews 2013. **12**(CD003388).
4. Bradley, R., et al., *A multidimensional meta-analysis of psychotherapy for PTSD*. American journal of Psychiatry, 2005. **162**(2): p. 214-227.
5. Jonas, D., et al., *Psychological and pharmacological treatments for adults with posttraumatic stress disorder (PTSD)*. *Comparative Effectiveness Review No. 92*. 2013.

6. Foa, E., et al., *Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: Outcome at academic and community clinics*. Journal of Consulting and Clinical Psychology, 2005. **73**: p. 953-964.
7. Resick, P., et al., *A comparison of cognitive-processing therapy with prolonged exposure and a waiting condition for the treatment of chronic posttraumatic stress disorder in female rape victims*. Journal of Consulting and Clinical Psychology, 2002. **70**(4): p. 867-879.
8. Schnurr, P., Friedman MJ, Engel CC, Foa EB, Shea MT, Chow BK, Resick PA, Thurston V, Orsillo SM, Haug R, Turner C, Bernardy N, *Cognitive behavioural therapy for posttraumatic stress disorder in women*. JAMA 2007. **28**: p. 820-830.
9. Schottenbauer, M.A., et al., *Nonresponse and dropout rates in outcome studies on PTSD: review and methodological considerations*. Psychiatry, 2008. **71**(2): p. 134-168.
10. Foa, E.B., et al., *Effective treatments for PTSD: practice guidelines from the International Society for Traumatic Stress Studies*. 2008: Guilford Press.
11. American Psychological Association, *Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder (PTSD) in Adults*. 2017, American Psychological Association: Washington, DC.
12. International Society of Traumatic Stress Studies (ISTSS). *New ISTSS Prevention and Treatment Guidelines [Online]*. 2018 [Accessed 26/11/18]; Available from: <http://www.istss.org/treating-trauma/new-istss-guidelines.aspx>.
13. National Institute for Health and Care Excellence (NICE), *Post-traumatic stress disorder (NICE guideline NG116)*. Retrieved from <https://www.nice.org.uk/guidance/ng116>, 2018.
14. Tarrier, N., et al., *A randomized trial of cognitive therapy and imaginal exposure in the treatment of chronic posttraumatic stress disorder*. Journal of Consulting and Clinical Psychology, 1999. **67**(1): p. 13-18.
15. Pitman, R.K., et al., *Psychiatric complications during flooding therapy for posttraumatic stress disorder*. Journal of Clinical Psychiatry, 1991.
16. Devilly, G. and E. Foa, *Comments on Tarrier et al's (1999) study and the investigation of exposure and cognitive therapy*. Journal of Consulting and Clinical Psychology, 2001.
17. Hembree, E.A., Foa, E. B., Dorfman, N. M., Street, G. P., Kowalski, J., Tu, X., *Do patients drop out prematurely from exposure therapy for PTSD?* Journal of Traumatic Stress, 2003. **16**(6): p. 555-562.
18. Imel, Z.E., et al., *Meta-analysis of dropout in treatments for posttraumatic stress disorder*. Journal of consulting and clinical psychology, 2013. **81**(3): p. 394.
19. Goetter, E.M., et al., *A systematic review of dropout from psychotherapy for posttraumatic stress disorder among Iraq and Afghanistan combat veterans*. Journal of Traumatic Stress, 2015. **28**(5): p. 401-409.
20. Higgins, J.P., et al., *The Cochrane Collaboration's tool for assessing risk of bias in randomised trials*. BMJ, 2011. **343**: p. d5928.
21. StataCorp, *Stata Statistical Software: Release 13*. College Station, TX: StataCorp LP. . 2013.
22. Thompson, S.G. and J.P. Higgins, *How should meta-regression analyses be undertaken and interpreted?* Statistics in medicine, 2002. **21**(11): p. 1559-1573.
23. Higgins, J. and S. Green, *Cochrane Handbook for Systematic Reviews of Interventions*. The Cochrane Collaboration, 2011. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org), 2011.
24. Cooper, A.A. and L.R. Conklin, *Dropout from individual psychotherapy for major depression: A meta-analysis of randomized clinical trials*. Clinical Psychology Review, 2015. **40**: p. 57-65.
25. Taylor, S., *Outcome predictors for three PTSD treatments: exposure therapy, EMDR, and relaxation training*. Journal of Cognitive Psychotherapy, 2003. **17**(2): p. 149.
26. Bryant, R.A., et al., *Who drops out of treatment for post-traumatic stress disorder?* Clinical Psychologist, 2007. **11**(1): p. 13-15.
27. Berk, M. and G. Parker, *The elephant on the couch: side-effects of psychotherapy*. Australian and New Zealand journal of psychiatry, 2009. **43**(9): p. 787-794.

28. Nutt, D.J. and M. Sharpe, *Uncritical positive regard? Issues in the efficacy and safety of psychotherapy*. Journal of Psychopharmacology, 2008. **22**(1): p. 3.
29. Linden, M., *How to define, find and classify side effects in psychotherapy: From unwanted events to adverse treatment reactions*. Clinical Psychology and Psychotherapy, published online in Wiley Online Library, DOI:10.1002/cpp.1765, 2012.
30. Becker, C.B., C. Zayfert, and E. Anderson, *A survey of psychologists' attitudes towards and utilization of exposure therapy for PTSD*. Behaviour Research and Therapy, 2004. **42**(277-292).
31. Pitman, R., et al., *Psychiatric complications during flooding therapy for post-traumatic stress disorder*. Journal of Clinical Psychiatry, 1991. **52**: p. 17-20.
32. Tarrier, N., et al., *A randomized controlled trial of cognitive therapy and imaginal exposure in the treatment of chronic posttraumatic stress disorder*. Journal of Consulting & Clinical Psychology, 1999. **67**(13-18).
33. Krakow, B., Hollifield, M., Johnston, L., Koss, M., Schrader, R., Warner, T. D., & Prince, H. , *Imagery rehearsal therapy for chronic nightmares in sexual assault survivors with posttraumatic stress disorder: a randomized controlled trial*. JAMA, 2001. **286**(5): p. 537-545.
34. Tarrier, N., Sommerfield, C., Pilgrim, H., & Humphreys, L., *Cognitive therapy or imaginal exposure in the treatment of post-traumatic stress disorder: 12-month follow-up*. British Journal of Psychiatry, 1999. **175**: p. 571-575.
35. Foa, E.B. and F.N. Zoellner LA, Hembree EA, Alvarez-Conrad J., *Does imaginal exposure exacerbate PTSD symptoms?* J Consult Clin Psychol, 2002. **70**(4).
36. Szafranski, D.D., et al., *High rates of PTSD treatment dropout: A possible red herring?* Journal of Anxiety Disorders, 2017. **47**: p. 91-98.
37. Bryant, R., et al., *Augmenting cognitive behaviour therapy for post-traumatic stress disorder with emotion tolerance training: a randomized controlled trial*. 2014.
38. Craske, M.G., et al., *CBT intensity and outcome for panic disorder in a primary care setting*. Behavior Therapy, 2006. **37**(2): p. 112-119.
39. Zandberg, L.J., et al., *Predictors of dropout in concurrent treatment of posttraumatic stress disorder and alcohol dependence: Rate of improvement matters*. Behaviour research and therapy, 2016. **80**: p. 1-9.
40. Bothwell, L.E., et al., *Assessing the gold standard—lessons from the history of RCTs*. 2016, Mass Medical Soc.
41. Roberts, N.P., N. Jones, and J.I. Bisson, *Psychological interventions for post-traumatic stress disorder and comorbid substance use disorder*. Cochrane Database of Systematic Reviews, 2016(Issue 4. Art. No.: CD010204. DOI: 10.1002/14651858.CD010204.pub2).
42. Hatchett, G.T. and H.L. Park, *Comparison of four operational definitions of premature termination*. Psychotherapy: Theory, Research, Practice, Training, 2003. **40**(3): p. 226.
43. DiMauro, J., et al., *A historical review of trauma-related diagnoses to reconsider the heterogeneity of PTSD*. Journal of anxiety disorders, 2014. **28**(8): p. 774-786.
44. Zatzick, D., *Toward the estimation of population impact in early posttraumatic stress disorder intervention trials*. Depression and anxiety, 2012. **29**(2): p. 79-84.
45. Cloitre, M., et al., *Skills training in affective and interpersonal regulation followed by exposure: a phase-based treatment for PTSD related to childhood abuse*. Journal of consulting and clinical psychology, 2002. **70**(5): p. 1067.
46. Bryant, R., et al., *Augmenting cognitive behaviour therapy for post-traumatic stress disorder with emotion tolerance training: a randomized controlled trial*. Psychological Medicine, 2013: p. 1-8.
47. Hernandez-Tejada, M.A., S. Hamski, and D. Sánchez-Carracedo, *Incorporating peer support during in vivo exposure to reverse dropout from prolonged exposure therapy for posttraumatic stress disorder: Clinical outcomes*. The International Journal of Psychiatry in Medicine, 2017. **52**(4-6): p. 366-380.

48. Acarturk, C., et al., *The efficacy of eye movement desensitization and reprocessing for post-traumatic stress disorder and depression among Syrian refugees: Results of a randomized controlled trial*. Psychological medicine, 2016. **46**(12): p. 2583-2593.
49. Adenauer, H., et al., *Narrative exposure therapy for PTSD increases top-down processing of aversive stimuli - evidence from a randomized controlled treatment trial*. BMC neuroscience, 2011. **12**(127): p. 1-13.
50. Ahmadi, K., et al., *REM desensitization as a new therapeutic method for post-traumatic stress disorder: a randomized controlled trial*. Acta Medica Indonesiana, 2015. **47**(2): p. 111-9.
51. Akbarian, F., et al., *The effectiveness of cognitive behavioral therapy with respect to psychological symptoms and recovering autobiographical memory in patients suffering from post-traumatic stress disorder*. Neuropsychiatric disease and treatment, 2015. **11**: p. 395-404.
52. Asukai, N., et al., *Efficacy of exposure therapy for Japanese patients with posttraumatic stress disorder due to mixed traumatic events: A randomized controlled study*. Journal of traumatic stress, 2010. **23**(6): p. 744-750.
53. Beck, J., et al., *Group cognitive behavior therapy for chronic posttraumatic stress disorder: an initial randomized pilot study*. Behavior Therapy, 2009. **40**: p. 82-92.
54. Bichescu, D., et al., *Narrative exposure therapy for political imprisonment related chronic posttraumatic stress disorder and depression*. Behaviour Research and Therapy, 2007. **45**: p. 2212-2220.
55. Blanchard, E., et al., *A controlled evaluation of cognitive behavioral therapy for posttraumatic stress in motor vehicle accident survivors*. Behaviour Research and Therapy, 2003. **41**(1): p. 79-96.
56. Bradshaw, R.A., et al., *A randomized clinical trial of Observed and Experiential Integration (OEI): A simple, innovative intervention for affect regulation in clients with PTSD*. Traumatology traumatology: an international journal, 2014. **20**: p. 161-171.
57. Brom, D., R. Kleber, and P. Defares, *Brief psychotherapy for posttraumatic stress disorders*. Journal of Consulting & Clinical Psychology, 1989. **57**(5): p. 607-612.
58. Bryant, R., et al., *Imaginal exposure alone and imaginal exposure with cognitive restructuring in treatment of posttraumatic stress disorder*. Journal of Consulting and Clinical Psychology, 2003. **71**: p. 706-12.
59. Bryant, R.A., et al., *A randomized controlled effectiveness trial of cognitive behavior therapy for post-traumatic stress disorder in terrorist-affected people in Thailand*. World psychiatry, 2011. **10**(3): p. 205-209.
60. Buhmann, C., et al. *The effect of flexible cognitive-behavioural therapy and medical treatment, including antidepressants on post-traumatic stress disorder and depression in traumatised refugees: pragmatic randomised controlled clinical trial*. 2016. **208**, 252-259 DOI: 10.1192/bjp.bp.114.150961.
61. Butollo, W., et al., *A randomized controlled clinical trial of dialogical exposure therapy versus cognitive processing therapy for adult outpatients suffering from PTSD after type I trauma in adulthood*. Psychotherapy and psychosomatics, 2016. **85**(1): p. 16-26.
62. Capezzani, L., et al., *EMDR and CBT for cancer patients: Comparative study of effects on PTSD, anxiety, and depression*. Journal of EMDR practice and research, 2013. **7**: p. 134-143.
63. Carletto, S., et al., *Treating post-traumatic stress disorder in patients with multiple sclerosis: A randomized controlled trial comparing the efficacy of eye movement desensitization and reprocessing and relaxation therapy*. Frontiers in Psychology, 2016. **7**.
64. Carlson, J., et al., *Eye movement desensitization and reprocessing (EDMR) treatment for combat-related posttraumatic stress disorder*. Journal of Traumatic Stress 1998. **11**(1): p. 3-32.
65. Castillo, D.T., et al., *Group-Delivered Cognitive/Exposure Therapy for PTSD in Women Veterans: A Randomized Controlled Trial*. Psychological trauma : theory, research, practice and policy, 2016: p. No-Specified.

66. Chard, K.M., *An evaluation of cognitive processing therapy for the treatment of posttraumatic stress disorder related to childhood sexual abuse*. Journal of consulting and clinical psychology, 2005. **73**(5): p. 965.
67. Cloitre, M., et al., *Skills training in affective and interpersonal regulation followed by exposure: a phase-based treatment for PTSD related to childhood abuse*. Journal of Consulting and Clinical Psychology, 2002. **70**: p. 1067-1074.
68. Cloitre, M., et al., *Treatment for PTSD related to childhood abuse: A randomized controlled trial*. American Journal of Psychiatry, 2010.
69. Devilly, G., S. Spence, and R. Rapee, *Statistical and reliable change with eye movement desensitization and reprocessing: treating trauma within a veteran population*. Behavior Therapy, 1998. **29**: p. 435-455.
70. Devilly GJ, S.S., *The relative efficacy and treatment distress of EMDR and a cognitive-behavior trauma treatment protocol in the amelioration of posttraumatic stress disorder*. Journal of Anxiety Disorders, 1999. **13**(1-2): p. 131-157.
71. Dorrepaal, E., et al., *Stabilizing group treatment for complex posttraumatic stress disorder related to child abuse based on psychoeducation and cognitive behavioural therapy: a multisite randomized controlled trial*. Psychotherapy and psychosomatics, 2012. **81**: p. 217-225.
72. Duffy, M., K. Gillespie, and D. Clark, *Post-traumatic stress disorder in the context of terrorism and other civil conflict in Northern Ireland: Randomised Controlled Trial*. British Medical Journal, 2007. **334**: p. 1147-1150.
73. Dunne, R.L., J. Kenardy, and M. Sterling, *A randomized controlled trial of cognitive-behavioral therapy for the treatment of PTSD in the context of chronic whiplash*. The Clinical journal of pain, 2012. **28**(9): p. 755-765.
74. Echeburua, E., I. Zubizarreta, and B. Sarasua, *Psychological treatment of chronic posttraumatic stress disorder in victims of sexual aggression*. Behavior Modification, 1997. **21**: p. 433-56.
75. Ehlers, A., et al., *Cognitive therapy for post-traumatic stress disorder: development and evaluation*. Behaviour Research and Therapy, 2005. **43**: p. 413-431.
76. Ehlers, A., et al., *A randomized controlled trial of cognitive therapy, a self-help booklet, and repeated assessments as early interventions for posttraumatic stress disorder*. Archives of General Psychiatry, 2003. **60**(10): p. 1024-1032.
77. Ehlers, A., et al., *A randomized controlled trial of 7-day intensive and standard weekly cognitive therapy for PTSD and emotion-focused supportive therapy*. American journal of psychiatry, 2014. **171**: p. 294-304.
78. Falsetti, S., H. Resnick, and J. Davis, *Multiple channel exposure therapy for women with PTSD and comorbid panic attacks*. Cognitive behaviour therapy, 2008. **37**: p. 117-130.
79. Fecteau, G. and R. Nicki, *Cognitive behavioural treatment of post traumatic stress disorder after motor vehicle accident*. Behavioural & Cognitive Psychotherapy, 1999. **27**(3): p. 201-214.
80. Feske, U., *Treating Low-Income and Minority Women With Posttraumatic Stress Disorder A Pilot Study Comparing Prolonged Exposure and Treatment as Usual Conducted by Community Therapists*. Journal of interpersonal violence, 2008. **23**(8): p. 1027-1040.
81. Foa, E., et al., *Treatment of posttraumatic stress disorder in rape victims: a comparison between cognitive-behavioral procedures and counseling*. Journal of Consulting and Clinical Psychology, 1991. **59**(5): p. 715-723.
82. Foa, E., et al., *A comparison of exposure therapy, stress inoculation training, and their combination for reducing posttraumatic stress disorder in female assault victims*. Journal of Consulting and Clinical Psychology, 1999. **67**(2): p. 194-200.
83. Foa, E., et al. *Effect of Prolonged Exposure Therapy Delivered Over 2 Weeks vs 8 Weeks vs Present-Centered Therapy on PTSD Symptom Severity in Military Personnel: a Randomized Clinical Trial*. 2018. **319**, 354-364 DOI: 10.1001/jama.2017.21242.

84. Fonzo, G., et al. *PTSD Psychotherapy Outcome Predicted by Brain Activation During Emotional Reactivity and Regulation*. 2017. **174**, 1163-1174 DOI: 10.1176/appi.ajp.2017.16091072.
85. Forbes, D., et al., *A multisite randomized controlled effectiveness trial of cognitive processing therapy for military-related posttraumatic stress disorder*. *Journal of Anxiety Disorders*, 2012. **26**(3): p. 442-452.
86. Ford, J.D., K.L. Steinberg, and W. Zhang, *A randomized clinical trial comparing affect regulation and social problem-solving psychotherapies for mothers with victimization-related PTSD*. *Behavior Therapy*, 2011. **42**(4): p. 560-578.
87. Ford, J.D., et al., *Randomized clinical trial comparing affect regulation and supportive group therapies for victimization-related PTSD with incarcerated women*. *Behavior Therapy*, 2013. **44**(2): p. 262-276.
88. Galovski, T.E., et al., *Manualized therapy for PTSD: Flexing the structure of cognitive processing therapy*. *Journal of Consulting and Clinical Psychology*, 2012. **80**(6): p. 968.
89. Gamito, P., et al., *PTSD elderly war veterans: a clinical controlled pilot study*. *Cyberpsychology, Behavior and Social Networking*, 2010. **13**(1): p. 43-48.
90. Gersons BP, C.I., Lamberts RD, Van der Kolk BA. , *Randomized clinical trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder*. *Journal of Traumatic Stress* 2000. **13**(2): p. 333-347.
91. Gray, R., D. Budden-Potts, and F. Bourke, *Reconsolidation of Traumatic Memories for PTSD: A randomized controlled trial of 74 male veterans*. *Psychotherapy Research*, 2017: p. 1-19.
92. Hensel-Dittmann, D., et al., *Treatment of traumatized victims of war and torture: A randomized controlled comparison of narrative exposure therapy and stress inoculation training*. *Psychotherapy and psychosomatics*, 2011. **80**(6): p. 345-352.
93. Hinton, D., Chhean, D, Pich, V, Safren, SA, Hofmann, SG, Pollack, MH, *A randomised controlled trial of cognitive behaviour therapy for Cambodian refugees with treatment resistant PTSD and panic attacks: a cross over design*. *Journal of Traumatic Stress*, 2005. **18**: p. 617-629.
94. Hinton, D.E., et al., *Culturally adapted CBT (CA-CBT) for Latino women with treatment-resistant PTSD: A pilot study comparing CA-CBT to applied muscle relaxation*. *Behaviour research and therapy*, 2011. **49**(4): p. 275-280.
95. Hogberg, G., Pagani, M, Sundin, O, Soares, J, Aberg-Wistedt, A, Tarnell, B, Hallstrom, T. , *On treatment with eye movement desensitization and reprocessing of chronic post-traumatic stress disorder in public transportation workers - a randomized controlled trial*. *Nordic Journal of Psychiatry*, 2007. **61**(1): p. 54-60.
96. Hollifield, M., Sinclair-Lian, N., Warner, T.D., Hammerschlag, R, *Acupuncture for posttraumatic stress disorder* *The Journal of Nervous and Mental Disease* 2007. **195**(6): p. 504-513.
97. Ironson, G., Freund, B, Strauss, JL, Williams, J. , *Comparison of two treatments for traumatic stress: a community based study of EMDR and prolonged exposure*. *Journal of Clinical Psychology* 2002. **58**: p. 113-128.
98. Ivarsson, D., et al., *Guided internet-delivered cognitive behavior therapy for post-traumatic stress disorder: A randomized controlled trial*. *Internet interventions*, 2014. **1**: p. 33-40.
99. Jacob, N., et al., *Dissemination of psychotherapy for trauma spectrum disorders in postconflict settings: A randomized controlled trial in Rwanda*. *Psychotherapy and psychosomatics*, 2014. **83**: p. 354-363.
100. Jensen, J., *An investigation of eye movement desensitisation and reprocessing as a treatment for posttraumatic stress disorder symptoms of vietnam combat veterans*. *Behavior Therapy*, 1994. **25**: p. 311-325.
101. Johnson, D.M., C. Zlotnick, and S. Perez, *Cognitive behavioral treatment of PTSD in residents of battered women's shelters: Results of a randomized clinical trial*. *Journal of consulting and clinical psychology*, 2011. **79**(4): p. 542.

102. Johnson, D., et al. *Comparison of Adding Treatment of PTSD During and After Shelter Stay to Standard Care in Residents of Battered Women's Shelters: results of a Randomized Clinical Trial*. 2016. **29**, 365-373 DOI: 10.1002/jts.22117.
103. Karatzias, T., et al., *A controlled comparison of the effectiveness and efficiency of two psychological therapies for posttraumatic stress disorder: eye movement desensitization and reprocessing vs. emotional freedom techniques*. *Journal of nervous and mental disease*, 2011. **199**: p. 372-8.
104. Keane, T., Fairbank, JA, Caddell, JM, Zimering, RT, *Implosive (flooding) therapy reduces symptoms of PTSD in Vietnam combat veterans*. *Behavior Therapy*, 1989. **20**(2): p. 245-260.
105. Krupnick, J.L., et al., *Group interpersonal psychotherapy for low-income women with posttraumatic stress disorder*. *Psychotherapy Research*, 2008. **18**(5): p. 497-507.
106. Kubany, E., Hill, EE, Owens, JA *Cognitive trauma therapy for battered women with PTSD: preliminary findings*. *Journal of Traumatic Stress* 2003. **16**: p. 81-91.
107. Kubany, E., E. Hill, and J. Owens, *Cognitive trauma therapy for battered women with PTSD (CTT-BW)* *Journal of Consulting and Clinical Psychology*, 2004. **72**(1): p. 3-18.
108. Laugharne, J., et al. *Amygdala volumetric change following psychotherapy for posttraumatic stress disorder*. 2016. **28**, 312-318 DOI: 10.1176/appi.neuropsych.16010006.
109. Lee, C., Gavriel, H, Drummond, P, Richards, J, Greenwald, R, *Treatment of PTSD: stress inoculation training with prolonged exposure compared to EMDR*. *Journal of Clinical Psychology* 2002. **58**: p. 1071-1089.
110. Lewis, et al., *Internet-based guided self-help for posttraumatic stress disorder (PTSD): Randomized controlled trial*. *Depression and anxiety*, 2017. **34**(6): p. 555-565.
111. Littleton, H., et al., *The From Survivor to Thriver program: RCT of an online therapist-facilitated program for rape-related PTSD*. *Journal of anxiety disorders*, 2016. **43**: p. 41-51.
112. Litz, B.T., et al., *A randomized, controlled proof-of-concept trial of an Internet-based, therapist-assisted self-management treatment for posttraumatic stress disorder*. *American journal of psychiatry*, 2007. **164**: p. 1676-1683.
113. Marcus, S., Marquis, P, Sakai, C, *Controlled study of treatment of PTSD using EMDR in an HMO setting* *Psychotherapy Theory, Research and Practice* 1997. **34**: p. 307-315.
114. Markowitz, J., et al., *Is exposure necessary? A randomized clinical trial of interpersonal psychotherapy for PTSD*. *American journal of psychiatry*, 2015. **172**: p. 430-440.
115. Marks, I., et al., *Treatment of posttraumatic stress disorder by exposure and/or cognitive restructuring: a controlled study*. *Archives of General Psychiatry* 1998. **55**(4): p. 317-325.
116. McDonagh, A., et al., *Randomized controlled trial of cognitive-behavioural therapy for chronic posttraumatic stress disorder in adult female survivors of childhood sexual abuse* *Journal of Consulting and Clinical Psychology* 2005. **73**: p. 515-524.
117. McLay, R., et al., *A randomized, controlled trial of virtual reality-graded exposure therapy for post-traumatic stress disorder in active duty service members with combat-related post-traumatic stress disorder*. *Cyberpsychology, behavior and social networking*, 2011. **14**: p. 223-229.
118. McLay, R., et al. *A Randomized, Head-to-Head Study of Virtual Reality Exposure Therapy for Posttraumatic Stress Disorder*. 2017. **20**, 218-224 DOI: 10.1089/cyber.2016.0554.
119. Monson, C., et al., *Effect of cognitive-behavioral couple therapy for PTSD: a randomized controlled trial* *Comment in: JAMA*. 2012 Aug 15;308(7):714-6; PMID: 22893170. *Jama*, 2012. **308**: p. 700-709.
120. Monson, C., Schnurr, PP, Resick, PA, Friedman, MJ, Young-Xu, Y, Stevens, SP *Cognitive processing therapy for veterans with military related posttraumatic stress disorder*. *Journal of Consulting and Clinical Psychology* 2006. **74**: p. 898-907.
121. Morath, J., et al. *Effects of psychotherapy on DNA strand break accumulation originating from traumatic stress*. 2014. **83**, 289-297 DOI: 10.1159/000362739.

122. Mueser, K.T., Rosenburg, S.D. Xie, H., Jankowski, M. K. Bolton, E. E., Lu, W., Hamblen, J. L., Rosenburg, H. J., McHugo., Wolfe, R A *randomized controlled trial of cognitive-behavioral treatment for posttraumatic stress disorder in severe mental illness* Journal of Consulting and Clinical Psychology 76(2):259-271., 2008. **76**: p. 259-271.
123. Nacasch, N., et al., *Prolonged exposure therapy for combat- and terror-related posttraumatic stress disorder: A randomized control comparison with treatment as usual* NCT00229372. Journal of clinical psychiatry, 2011. **72**: p. 1174-1180.
124. Neuner, F., Kurreck, S., Ruf, M., Odenwald, M., Elbert, T., Schauer, M.. A randomized controlled pilot study *Can asylum seekers with posttraumatic stress disorder be successfully treated?* Cognitive Behaviour Therapy 2010. **39**(2): p. 81-91.
125. Neuner, F., et al., *Treatment of posttraumatic stress disorder by trained lay counselors in an African refugee settlement: a randomized controlled trial.* Journal of consulting and clinical psychology, 2008. **76**(4): p. 686.
126. Neuner, F., et al., *A comparison of narrative exposure therapy, supportive counselling, and psychoeducation for treating posttraumatic stress disorder in an African refugee settlement* Journal of Consulting and Clinical Psychology 2004. **72**(4): p. 579-587.
127. Nijdam, M.J., et al., *Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: randomised controlled trial.* The British Journal of Psychiatry, 2012. **200**(3): p. 224-231.
128. Pacella, M.L., et al., *The impact of prolonged exposure on PTSD symptoms and associated psychopathology in people living with HIV: A randomized test of concept.* AIDS and Behavior, 2012. **16**(5): p. 1327-1340.
129. Paunovic, N., *Exposure Inhibition Therapy as a Treatment for Chronic Posttraumatic Stress Disorder: A Controlled Pilot Study.* Psychology, 2011. **2**(6): p. 605.
130. Peniston, E.G. and P.J. Kulkosky, *Alpha-theta brainwave neurofeedback for Vietnam veterans with combat-related post-traumatic stress disorder.* Medical Psychotherapy, 1991. **4**(1): p. 47-60.
131. Power, K., McGoldrick, T, Brown, K, Buchanan, R, Sharp, D, Swanson, V et al. , *A controlled comparison of eye movement desensitisation and reprocessing versus exposure plus cognitive restructuring versus waiting list in the treatment of post-traumatic stress disorder.* Clinical Psychology and Psychotherapy, 2002. **9**: p. 229-318.
132. Rauch, S., et al., *Biological and symptom changes in posttraumatic stress disorder treatment: A randomized clinical trial.* Depression and anxiety, 2015. **32**: p. 204-212.
133. Ready, D.J., Gerardi, R.J., Backscheider, A. G., Mascaro, N., Rothbaum, B. O. , *Comparing virtual reality exposure therapy to present centred therapy with 11 US Vietnam Veterans with PTSD* Cyberpsychology, Behavior and Social Networking, 2010. **13**(1): p. 49-54.
134. Reger, G., et al. *Randomized controlled trial of prolonged exposure using imaginal exposure vs. virtual reality exposure in active duty soldiers with deployment-related posttraumatic stress disorder (PTSD).* 2016. **84**, 946-959 DOI: 10.1037/ccp0000134.
135. Resick, P.A., et al., *A Randomized Clinical Trial of Group Cognitive Processing Therapy Compared With Group Present-Centered Therapy for PTSD Among Active Duty Military Personnel.* Journal of consulting and clinical psychology journal of consulting psychology, 2015: p. [Epub ahead of print].
136. Resick, P.A., et al., *Effect of group vs individual cognitive processing therapy in active-duty military seeking treatment for posttraumatic stress disorder: A randomized clinical trial.* JAMA psychiatry, 2017. **74**(1): p. 28-36.
137. Rothbaum, B., *A controlled study of eye movement desensitization and reprocessing in the treatment of posttraumatic stress disorder sexual assault victims* Bulletin of the Menninger Clinic 1997. **61**(61): p. 317-334.



138. Rothbaum, B., Astin, MC, Marsteller, F. , *Prolonged exposure versus eye movement desensitisation and reprocessing (EMDR) for PTSD rape victims. .* Journal of Traumatic Stress 2005. **18**: p. 607-616.
139. Sautter, F.J., et al., *Efficacy of structured approach therapy in reducing PTSD in returning veterans: A randomized clinical trial.* Psychological services, 2015. **12**(3): p. 199-212.
140. Scheck, M., Schaeffer, JA, Gillette, C. , *Brief psychological intervention with traumatized young women: the efficacy of eye movemebnt desensitisation and reporcessing.* Journal of Traumatic Stress, 1998. **11**: p. 25-44.
141. Schnurr, P., Friedman, MJ, Foy, DW, Shea, MT, Hsieh, FY, Lavori, PW et al. , *Randomized trial of trauma-focused group therapy for posttraumatic stress disorder. .* Archives of General Psychiatry 2003. **60**: p. 481-489.
142. Schnurr, P., Friedman MJ, Engel CC, Foa EB, Shea MT, Chow BK, Resick PA, Thurston V, Orsillo SM, Haug R, Turner C, Bernardy N. , *Cognitive behavioural therapy for posttraumatic stress disorder in women. .* JAMA 2007. **28**: p. 820-830.
143. Schnyder, U., et al., *Brief eclectic psychotherapy for PTSD: A randomized controlled trial.* Journal of Clinical Psychiatry 2011. **72**(564-566).
144. Sloan, D., et al., *Written exposure as an intervention for PTSD: A randomized clinical trial with motor vehicle accident survivors.* Behaviour research and therapy, 2012. **50**(10): p. 627-635.
145. Sloan, D., et al., *A Brief Exposure-Based Treatment vs Cognitive Processing Therapy for Posttraumatic Stress Disorder: A Randomized Noninferiority Clinical Trial.* JAMA Psychiatry, 2018.
146. Spence, J., et al., *Randomised controlled trial of internet delivered cognitive behavioural therapy for posttraumatic stress disorder.* Depression and anxiety, 2011. **28**: p. 541-550.
147. Stenmark, H., et al., *Treating PTSD in refugees and asylum seekers within the general health care system. A randomized controlled multicenter study.* Behaviour research and therapy, 2013. **51**: p. 641-647.
148. Suris, A., et al., *A randomized clinical trial of cognitive processing therapy for veterans with PTSD related to military sexual trauma. .* Journal of traumatic stress, 2013. **26**: p. 28-37.
149. Taylor, S., Thordarson, DS, Maxfield, L, Fedoroff, IC, Lovell, K, Ogradniczuk, J. , *Comparative efficacy, speed, and adverse effects of three PTSD treatments: exposure therapy, EMDR, and relaxation training* Journal of Consulting and Clinical Psychology 2003. **71**: p. 330-338.
150. Tylee, D.S., et al., *Evaluation of the reconsolidation of traumatic memories protocol for the treatment of PTSD: a randomized, wait-list-controlled trial.* Journal of Military, Veteran and Family Health, 2017. **3**(1): p. 21-33.
151. Vaughan, K., Armstrong, MS, Gold, R, O'Connor, N, Jenneke, W, Tarrier, N, *A trial of eye movement desensitization compared to image habituation training and applied muscle relaxation in post-traumatic stress disorder.* Journal of Behavior Therapy and Experimental Psychiatry 1994. **25**: p. 283-91.
152. Wells, A., et al., *Metacognitive therapy versus prolonged exposure in adults with chronic post-traumatic stress disorder: A parallel randomized controlled trial.* Cognitive therapy and research, 2015: p. No-Specified.
153. Wells, A. and S. Sembi, *Metacognitive therapy for PTSD: A preliminary investigation of a new brief treatment.* Journal of Behavior Therapy and Experimental Psychiatry, 2004. **35**(4): p. 307-318.
154. Yehuda, R., et al., *Glucocorticoid-related predictors and correlates of post-traumatic stress disorder treatment response in combat veterans.* Interface Focus, 2014. **4**(5): p. 20140048.
155. Zang, Y., N. Hunt, and T. Cox, *Adapting narrative exposure therapy for Chinese earthquake survivors: A pilot randomised controlled feasibility study.* BMC psychiatry, 2014. **14**.
156. Zang, Y., N. Hunt, and T. Cox, *A randomised controlled pilot study: the effectiveness of narrative exposure therapy with adult survivors of the Sichuan earthquake.* BMC psychiatry, 2013. **13**(1): p. 1.

157. Zlotnick, C., et al., *An affect-management group for women with posttraumatic stress disorder and histories of childhood sexual abuse*. *Journal of traumatic stress*, 1997. **10**(3): p. 425-436.

Participants were traumatised by military combat (27 studies), sexual assault or rape (11 studies), war/persecution (8 studies), road traffic accidents (6 studies), earthquakes (2 studies), childhood sexual abuse (7 studies), political detainment (1 study), terrorism (2 studies), physical assault (2 studies), domestic violence (4 studies), trauma from a medical diagnosis/emergency (4 studies) and crime/organised violence (4 studies). The remainder (43 studies) included individuals traumatised by a variety of different traumatic events.

Study	N	Country	Intervention 1	Intervention 2	Intervention 3	Intervention 4	Population	Trauma type	% Female	% Unemployed	% University Educated
Acarturk 2016 [48]	98	Turkey/Syria	EMDR	WL			Refugees	War/Persecution	74	Unknown	4
Adenauer 2011 [49]	34	Germany	NET (CBT-T)	WL			Refugees	War/Persecution	44	Unknown	Unknown
Ahmadi 2015 [50]	48	Iran	EMDR	REM Desensitization	WL		Military Personnel/Veterans	Military Trauma	0	Unknown	33.3
Akbarian 2015 [51]	40	Iran	Group CBT-T	MC/RA			General Population	Various	79	Unknown	Unknown
Asukai 2010 [52]	24	Japan	PE (CBT-T)	TAU			General Population	Various	88	Unknown	Unknown
Beck 2009 [53]	44	USA	Group CBT-T	MC/RA			General Population	Road Traffic Accident	82	54	Unknown
Bichescu 2007 [54]	18	Romania	NET (CBT-T)	Psychoeducation			General Population	Political detainment	94	0	72
Blanchard 2003 [55]	98	USA	CBT-T	SC	WL		General Population	Road Traffic Accident	73	Unknown	Unknown
Bradshaw 2014 [56]	10	Canada	OEI	WL			General Population	Various	70	0	Unknown
Brom 1989 [57]	83	Netherlands	CBT-T	Psychodynamic Therapy	WL		General Population	Various	79	49	Unknown
Bryant 2003 [58]	58	Australia	CBT-T	SC			General Population	Various	52	Unknown	Unknown
Bryant 2011 [59]	28	Thailand	CBT-T	SC			General Population	Terrorism	96	84%	Unknown
Buhmann 2016 [60]	138	Denmark	CBT-T	WL			Refugees	Organised Violence	41	Unknown	Unknown
Buttolo 2016 [61]	148	Germany	CPT (CBT-T)	DET			General Population	Various	66	Unknown	Unknown
Capezzani 2013 [62]	21	Italy	EMDR	CBT-T			General Population	Medical Diagnoses/Emergencies	90	Unknown	Unknown
Carletto 2016 [63]	50	Italy	EMDR	Relaxation Training			General Population	Medical Diagnoses/Emergencies	81	Unknown	Unknown

<b>Carlson 1998 [64]</b>	35	USA	EMDR	Relaxation Training	TAU		Military Personnel/Veterans	Military Trauma	0	62	Unknown
<b>Castillo 2016 [65]</b>	86	USA	Group CBT-T	WL			Military Personnel/Veterans	Military Trauma	100	44	Unknown
<b>Chard 2005 [66]</b>	71	USA	CPT (CBT-T)	WL			General Population	Sexual Assault or Rape	100	Unknown	Unknown
<b>Cloitre 2002 [67]</b>	58	USA	CBT-T	WL			General Population	Child Abuse	100	24	52
<b>Cloitre 2010 [68]</b>	71	USA	CBT-T	CBT without a trauma focus			General Population	Child Abuse	100	31	Unknown
<b>Cooper 1989</b>	16	USA	EMDR	Relaxation Therapy			Military Personnel/Veterans	Military Trauma	0	Unknown	Unknown
<b>Deville 1998 [69]</b>	35	Australia	EMDR	TAU			Military Personnel/Veterans	Military Trauma	0	Unknown	Unknown
<b>Deville 1999 [70]</b>	32	Australia	EMDR	CBT-T			General Population	Various	100	Unknown	Unknown
<b>Dorrepaal 2012 [71]</b>	71	Netherlands	Group Stabilising Treatment	TAU			General Population	Child Abuse	Unknown	83	Unknown
<b>Duffy 2007 [72]</b>	58	UK	CT (CBT-T)	WL			General Population	Various	40	Unknown	Unknown
<b>Dunne 2012 [73]</b>	26	Australia	CBT-T	WL			General Population	Road Traffic Accident	50	31	73
<b>Echeburua 1997 [74]</b>	20	Spain	CBT-T	Relaxation Training			General Population	Sexual Assault or Rape	100	Unknown	20
<b>Ehlers 2005 [75]</b>	28	UK	CT (CBT-T)	WL			General Population	Various	50	25	35
<b>Ehlers 2003 [76]</b>	57	UK	CT (CBT-T)	MC/RA			General Population	Road Traffic Accident	Unknown	Unknown	Unknown
<b>Ehlers 2014 [77]</b>	91	UK	CT (CBT-T)	SC	WL		General Population	Various	58	23	26
<b>Falsetti 2008 [78]</b>	60	USA	Group CBT-T	WL			General Population	Various	100	Unknown	Unknown
<b>Fecteau 1999 [79]</b>	20	Canada	CBT-T	WL			General Population	Road Traffic Accident	70	Unknown	Unknown
<b>Feske 2008 [80]</b>	21	USA	PE (CBT-T)	TAU			General Population	Various	100	29%	90%
<b>Foa 1991 [81]</b>	45	USA	PE (CBT-T)	CBT without a trauma focus	Supportive counselling	WL	General Population	Sexual Assault or Rape	100	Unknown	Unknown
<b>Foa 1999 [82]</b>	66	USA	PE (CBT-T)	CBT without a trauma focus	WL		General Population	Sexual Assault or Rape	100	38	41
<b>Foa 2005 [6]</b>	179	USA	PE (CBT-T)	WL			General Population	Assault	100	17	34

Foa 2018 [83]	256	USA	Spaced PE (CBT-T)	PCT	MC/RA		Military Personnel/Veterans	Military Trauma	12	100	66
Fonzo 2017 [84]	66	USA	PE (CBT-T)	WL			General Population	Various	65	Unknown	Unknown
Forbes 2012 [85]	59	Australia	CPT (CBT-T)	TAU			Military Personnel/Veterans	Military Trauma	4	36	Unknown
Ford 2011 [86]	146	USA	CBT without a trauma focus	PCT	WL		General Population	Various	100	Unknown	22
Ford 2013 [87]	80	USA	Group CBT-T	Group Supportive Counselling			Incarcerated Women	Various	100	Unknown	Unknown
Galovski 2012 [88]	100	USA	CPT (CBT-T)	MC/RA			General Population	Various	69	Unknown	Unknown
Gamito 2010 [89]	10	Portugal	VRE (CBT-T)	Control Exposure	WL		Military Personnel/Veterans	Military Trauma	0	Unknown	Unknown
Gersons 2000 [90]	42	Netherlands	BEP (CBT-T)	WL			General Population	Various	Unknown	Unknown	Unknown
Gray 2017 [91]	74	USA	RTM (CBT-T)	WL			Military Personnel/Veterans	Military Trauma	0	Unknown	Unknown
Hensel-Dittmann 2011 [92]	28	Germany	NET (CBT-T)	CBT without a trauma focus			Asylum Seekers	Organised Violence	Unknown	Unknown	Unknown
Hinton 2005 [93]	40	USA	CBT-T	WL			Refugees	Genocide	60	Unknown	Unknown
Hinton 2011 [94]	24	USA	Group CBT-T	WL			General Population	Various	100	Unknown	Unknown
Hogberg 2007 [95]	24	Sweden	EMDR	WL			General Population	Various	38	Unknown	Unknown
Hollifield 2007 [96]	55	USA	Group trauma-focused CBT	WL			General Population	Various	68	Unknown	40
Ironson 2002 [97]	22	USA	EMDR	PE (CBT-T)			General Population	Various	77	Unknown	Unknown
Ivarsson 2014 [98]	62	Sweden	I-CBT	WL			General Population	Various	82	8	65
Jacob 2014 [99]	76	Rwanda	NET (CBT-T)	WL			Genocide Survivors	Genocide	92	Unknown	Unknown
Jensen 1994 [100]	25	USA	EMDR	WL			Military Personnel/Veterans	Military Trauma	0	68	Unknown

Johnson 2011 [101]	70	USA	CBT without a trauma focus	TAU			General Population	Domestic Abuse	100	73	7
Johnson 2016 [102]	60	USA	CBT without a trauma focus	TAU			General Population	Domestic Abuse	100	77	5
Karatzias 2011 [103]	46	UK	EMDR	EFT			General Population	Various	57	37	47
Keane 1989 [104]	24	USA	CBT-T	WL			Military Personnel/Veterans	Military Trauma	0	Unknown	Unknown
Krupnick 2008 [105]	48	USA	Group IPT	WL			General Population	Various	100	80	13%
Kubany 2003 [106]	37	USA	CBT-T	WL			General Population	Domestic Abuse	100	Unknown	Unknown
Kubany 2004 [107]	107	USA	CBT-T	WL			General Population	Domestic Abuse	100	Unknown	Unknown
Laugharne 2016 [108]	20	Australia	EMDR	PE (CBT-T)			General Population	Various	70	Unknown	Unknown
Lee 2002 [109]	24	Australia	CBT-T	EMDR			General Population	Various	46	Unknown	Unknown
Lewis 2017 [110]	42	UK	I-CBT	WL			General Population	Various	57	19	62
Lindauer	24	Netherlands	BEP	WL			General Population	Various	54	Unknown	Unknown
Littleton 2016 [111]	87	USA	I-CBT	I- Psychoeducation			General Population	Sexual Assault or Rape	100	Unknown	Unknown
Litz 2007 [112]	45	USA	I-CBT	I-SC			Military Personnel/Veterans	Terrorism / Military Trauma	Unknown	Unknown	Unknown
Marcus 1997 [113]	67	USA	EMDR	TAU			General Population	Various	79	Unknown	Unknown
Markowitz 2015 [114]	110	USA	IPT	PE (CBT-T)	Relaxation Therapy		General Population	Various	70	21	Unknown
Marks 1998 [115]	87	UK	PE (CBT-T)	Cognitive Restructuring	PE (CBT-T) (CBT-T)and Cognitive Restructuring	Relaxation without PE (CBT-T) (CBT-T)(CBT-T)or CR	General Population	Various	36	54	Unknown
McDonagh 2005 [116]	74	USA	PE (CBT-T)	PCT	WL		General Population	Sexual Assault or Rape	100	17	Unknown
McLay 2011 [117]	20	USA	VRE (CBT-T)	TAU			Military Personnel/Veterans	Military Trauma	5	Unknown	Unknown

McLay 2017 [118]	81	USA	VRE (CBT-T)	Control Exposure Therapy			Military Personnel/Veterans	Military Trauma	4	Unclear	Unclear
Monson 2012 [119]	20	USA	Couples CBT-T	WL			General Population	Various	25	40	Unknown
Monson 2006 [120]	60	USA	CPT (CBT-T)	WL			Military Personnel/Veterans	Military Trauma	10	Unknown	Unknown
Morath 2014 [121]	38	Germany	NET (CBT-T)	WL			Refugees	Organised Violence	32	Unknown	Unknown
Meuser 2008 [122]	108	USA	CBT-T	TAU			General Population	Various	79	Unknown	Unknown
Nacasch 2011 [123]	30	Israel	PE (CBT-T)	TAU			Military Personnel/Veterans	Military Trauma	Unknown	63	Unknown
Neuner 2010 [124]	32	Germany	NET (CBT-T)	TAU			Refugees	Torture	31	Unknown	Unknown
Neuner 2008 [125]	277	Uganda	NET (CBT-T)	SC	Monitoring		Refugees	War/Persecution	51	49	Unknown
Neuner 2004 [126]	43	Uganda	NET (CBT-T)	SC	Psychoeducation		Refugees	War/Persecution	60	28	Unknown
Nijdam 2012 [127]	140	Netherlands	BEP (CBT-T)	EMDR			General Population	Various	56	Unknown	30
Pacella 2012 [128]	66	USA	PE (CBT-T) (CBT-T)	MC/RA			General Population	Medical Diagnoses/Emergencies	37	Unknown	Unknown
Paunovic 2011 [129]	29	Sweden	CBT-T	WL			General Population	Various	63	74	11
Peniston 1991 [130]	29	USA	CBT-T	TAU			Military Personnel/Veterans	Military Trauma	Unknown	Unknown	Unknown
Power 2002 [131]	105	UK	EMDR	CBT-T	WL		General Population	Various	42	Unknown	Unknown
Rauch 2015 [132]	36	USA	PE (CBT-T) (CBT-T)	PCT			Military Personnel/Veterans	Military Trauma	9	Unknown	Unknown
Ready 2010 [133]	11	USA	VRE (CBT-T)	PCT			Military Personnel/Veterans	Military Trauma	Unknown	Unknown	Unknown
Reger 2016 [134]	162	USA	VRE (CBT-T)	PE (CBT-T)	WL		Military Personnel/Veterans	Military Trauma	4	Active duty	7
Resick 2015 [135]	108	USA	Group CBT-T	Group PCT			Military Personnel/Veterans	Military Trauma	8	0	8
Resick 2002 [7]	171	USA	CPT (CBT-T) (CBT-T)	PE (CBT-T)	Minimal Attention		General Population	Sexual Assault or Rape	100	Unknown	Unknown
Resick 2017 [136]	268	USA	CPT (CBT-T) (CBT-T)	Group CBT-T			Military Personnel/Veterans	Military Trauma	9	100	19

Rothbaum 1997 [137]	18	USA	EMDR	WL			General Population	Sexual Assault or Rape	100	19	43
Rothbaum 2005 [138]	60	USA	PE (CBT-T)	EMDR	WL		General Population	Sexual Assault or Rape	100	Unknown	Unknown
Sautter 2015 [139]	57	USA	Couples CBT without a trauma focus	Couples Psychoeducation			Military Personnel/Veterans	Military Trauma	1.75	12	75
Scheck 1998 [140]	60	USA	EMDR	SC			General Population	Various	100	Unknown	Unknown
Schnurr 2003 [141]	360	USA	Group CBT-T	Group PCT			Military Personnel/Veterans	Military Trauma	0	51	Unknown
Schnurr 2007 [142]	284	USA	PE (CBT-T) (CBT-T)	Group PCT			Military Personnel/Veterans	Military Trauma	100	38	Unknown
Schnyder 2011 [143]	30	Switzerland	BEP (CBT-T)	MC/RA			General Population	Various	46.7	Unknown	Unknown
Shemesh	60	USA	CBT-T	Psychoeducation			General Population	Medical Diagnoses/Emergencies	33	Unknown	Unknown
Sloan 2012 [144]	46	USA	WET	WL			General Population	Road Traffic Accident	Unclear	78	41
Sloan 2018 [145]	126	USA	WET	CPT (CBT-T)			General Population	Various	49	Unknown	13
Spence 2011 [146]	42	Australia	I-CBT	WL			General Population	Various	81	41	Not Clear
Stenmark 2013 [147]	81	Norway	NET (CBT-T)	TAU			Refugees	Various	31	Unknown	25
Suris 2013 [148]	86	USA	CPT (CBT-T)	PCT			Military Personnel/Veterans	Sexual Assault or Rape	85	43	16
Taylor 2003 [149]	60	USA	PE (CBT-T)	Relaxation Therapy	EMDR		General Population	Various	75	13	Unknown
Tylee 2017 [150]	30	USA	RTM (CBT-T)	WL			General Population	Military Trauma	0	Unknown	Unknown
Vaughan 1994 [151]	36	Australia	CBT-T	Relaxation Training	EMDR		General Population	Various	64	Unknown	Unknown
Wells 2015 [152]	32	UK	PE (CBT-T)	CBT without a trauma focus	WL		General Population	Various	38	6	Unknown
Wells 2012 [153]	20	UK	CBT without a trauma focus	WL			General Population	Various	55	Unknown	Unknown
Yehuda 2014 [154]	52	USA	PE (CBT-T)	MC/RA			Military Personnel/Veterans	Military Trauma	Unclear	Unknown	Unknown



Zang 2014 [155]	20	China	NET (CBT-T)	WL			General Population	Earthquake	90	Unknown	Unknown
Zang 2013 [156]	22	China	NET (CBT-T)	WL			General Population	Earthquake	77	Unknown	Unknown
Zlotnick 1997 [157]	48	USA	Group CBT-T	WL			General Population	Sexual Assault or Rape	100	Unknown	33

**BEP** = Brief Eclectic Psychotherapy

**CBT** = Cognitive Behavioural Therapy

**CBT-T** = Cognitive Behavioural Therapy with a Trauma focus

**CPT** = Cognitive Processing Therapy

**CR** = Cognitive Restructuring

**CT** = Cognitive Therapy

**DET** = Dialogical Exposure Therapy

**EFT** = Emotional Freedom Technique

**EMDR** = Eye Movement Desensitisation and Reprocessing

**I-CBT** = Internet-based Cognitive Behavioural Therapy

**I-Psychoeducation** = Internet based Psychoeducation

**IPT** = Interpersonal Psychotherapy

**I-SC** = Internet based Supportive Counselling

**MC/RA** = Medical Checks/Repeated Assessments

**NET** = Narrative Exposure Therapy

**OEI** = Observed and Experimental Integration

**PCT** = Present Centred Therapy

**PE** = Prolonged Exposure

**REM Desensitization** = Rapid Eye Movement Desensitization

**RTM** = Reconsolidation of Traumatic Memories

**SC** = Supportive Counselling

**TAU** = Treatment as Usual

**VRE** = Virtual Reality Exposure

**WET** = Written Emotion Therapy

**WL** = Waiting List

**Table 1: Characteristics of Included Studies**

	Random sequence generation	Allocation concealment	Incomplete outcome data assessment	Blinding of outcome	Selective reporting	Other sources of bias	Total no. high risk
Acarturk 2016	Low	Low	Low	Low	Low	Low	0
Adenauer 2011	Low	Low	Low	Low	High	High	2
Ahmadi 2015	Unclear	Unclear	High	Unclear	Unclear	High	2
Akbarian 2015	Low	High	Low	Low	Unclear	High	2
Asukai 2010	Low	Low	Low	Low	Unclear	High	1
Beck 2009	Unclear	Unclear	High	Low	Unclear	High	2
Bichescu 2007	High	Unclear	Low	Low	Unclear	High	2
Blanchard 2003	High	Unclear	Low	Low	Unclear	Low	1
Bradshaw 2014	Unclear	Unclear	Low	High	Unclear	High	2
Brom 1989	Unclear	Unclear	High	Unclear	Unclear	High	2
Bryant 2003	Low	Unclear	Low	Low	Low	High	1
Bryant 2011	Low	Low	Low	Low	Unclear	High	1
Buhmann 2016	Low	Low	Unclear	Low	Low	Low	0
Buttolo 2016	Unclear	Unclear	Low	Low	Unclear	High	1
Capezzani 2013	Unclear	Unclear	Low	Low	Unclear	High	1
Carletto 2016	Low	Low	High	Low	Low	Low	1
Carlson 1998	Unclear	Unclear	High	Unclear	Unclear	Low	1
Castillo 2016	Unclear	Unclear	Low	Low	Unclear	High	1
Chard 2005	Unclear	Unclear	Low	Low	Unclear	High	1
Cloitre 2002	Unclear	Unclear	Low	Low	High	Low	1

Cloitre 2010	Unclear	Low	Low	Low	Low	Low	0
Cooper 1989	High	High	High	Unclear	Low	High	4
Devilly 1998	Unclear	Unclear	High	Low	Unclear	Low	1
Devilly 1999	High	Unclear	High	Unclear	Unclear	High	3
Dorrepaal 2012	Unclear	Low	Low	Low	High	High	2
Duffy 2007	Low	Low	Low	Unclear	Low	High	1
Dunne 2012	Unclear	Unclear	Low	Unclear	Unclear	High	1
Echeburua1997	Unclear	Unclear	Low	Unclear	Unclear	High	1
Ehlers 2003	Low	Low	High	Low	Unclear	High	2
Ehlers 2005	Unclear	Unclear	Low	Low	Unclear	High	2
Ehlers 2014	Unclear	Low	Low	Low	Low	Low	0
Falsetti 2008	Unclear	Unclear	Low	Low	High	High	2
Fecteau 1999	Low	Unclear	High	Unclear	Unclear	High	2
Feske 2008	Unclear	Unclear	Low	Unclear	Unclear	High	1
Foa 1991	Unclear	Unclear	High	Low	Unclear	High	2
Foa 1999	Unclear	Unclear	Low	Low	Unclear	High	1
Foa 2005	Low	Low	Low	Low	Unclear	Low	0
Foa 2018	Low	Low	Low	Low	Low	Low	0
Fonzo 2017	Low	Unclear	Low	Unclear	Low	Low	0
Forbes 2012	Unclear	Low	Low	Unclear	Unclear	High	1
Ford 2011	Low	Low	Low	Low	Unclear	High	1
Ford 2013	Low	Low	High	Low	Unclear	High	2
Galovski 2012	Unclear	Unclear	Low	Low	Unclear	Low	0
Gamito 2010	Unclear	Unclear	Unclear	Unclear	High	High	2
Gersons 2000	Unclear	Unclear	Low	Low	Unclear	Low	0
Gray 2017	Low	Low	Unclear	Unclear	Unclear	Unclear	0
Hensel-Dittmann 2011	Low	Low	Low	Low	Unclear	Low	0



Monson 2006	Low	Low	Low	Low	Unclear	Low	0
Morath 2014	Low	Low	Unclear	Low	Low	Low	0
Meuser 2008	Low	Low	Low	Low	Unclear	High	1
Nacasch 2011	Low	Unclear	Low	Low	Low	High	1
Neuner 2004	Low	Unclear	Low	Low	Low	High	1
Neuner 2008	Unclear	Unclear	Low	Low	Unclear	Low	0
Neuner 2010	Unclear	Unclear	Low	Low	Unclear	High	1
Nijdam 2012	Unclear	Low	Low	Low	Low	Low	0
Pacella 2015	Low	Unclear	Low	Low	Unclear	Low	0
Paunovic 2011	Unclear	Unclear	Low	High	Unclear	High	2
Power 2002	Low	Low	High	Low	Unclear	Low	1
Rauch 2015	Unclear	Unclear	Low	Low	Unclear	High	1
Ready 2010	Unclear	Unclear	Unclear	Low	Unclear	High	1
Reger 2016	Low	Low	Low	Low	Unclear	Low	0
Resick 2002	Unclear	Unclear	Low	Low	Unclear	High	1
Resick 2015	Unclear	Unclear	Low	Low	Unclear	Low	0
Resick 2017	Low	Unclear	Low	Low	Low	Low	0
Rothbaum 1997	Unclear	Unclear	High	Low	Unclear	High	2
Rothbaum 2005	Unclear	Unclear	High	Low	Unclear	Low	1
Sautter 2015	Unclear	Unclear	Low	Low	Unclear	Low	0
Scheck 1998	Low	Low	High	Unclear	Unclear	High	2
Schnurr 2003	High	Unclear	Low	Low	Low	Low	1
Schnurr 2007	Low	Low	Low	Low	Low	Low	0
Shemesh	Low	Low	Unclear	Unclear	Unclear	Low	0
Sloan 2012	Low	Low	Unclear	Low	Unclear	Low	0
Sloan 2018	Low	Low	Low	Low	Low	Low	0
Spence 2011a	Low	Unclear	High	High	Low	Unclear	2
Stenmark 2013	Unclear	Unclear	Low	High	Low	High	2

Suris 2013	Unclear	Unclear	Low	Low	Low	High	1
Taylor 2003	Unclear	Unclear	Low	Low	Unclear	Low	0
Tylee 2017	Unclear	Unclear	Unclear	Low	Unclear	High	1
Vaughan 1994	Unclear	Unclear	Low	Low	Unclear	Low	0
Wells 2012	Low	Low	Low	Low	Unclear	High	1
Wells 2015	Low	Low	High	High	Unclear	High	3
Yehuda 2014	Unclear	Unclear	High	Unclear	Unclear	Unclear	1
Zang 2013	Unclear	Unclear	Low	Low	Low	High	1
Zang 2014	Low	Unclear	Low	Low	Low	High	1
Zlotnick 1997	Unclear	Unclear	High	Low	Low	High	2

Table 2: Risk of bias assessments of the included studies

	Number of studies	Mean % drop out (95% CI)	I <sup>2</sup> (%)
1. CBT-T (not based on a specific model)	25	13 (9-18)	64.41
2. Brief Eclectic Psychotherapy	3	17 (0-51)	90.40
3. Cognitive Processing Therapy (CPT)	8	30 (22-39)	75.15
4. Cognitive Therapy (CT)	6	9 (1-23)	82.72
5. Narrative Exposure Therapy	11	12 (3-26)	85.59
6. Prolonged Exposure (PE)	22	22 (16-28)	72.56
7. Reconsolidation of Traumatic Memories (RTM);	1	1 (0-8)	0.00
8. Virtual Reality Exposure (VRE)	5	18 (3-38)	76.32
9. Eye Movement Desensitisation and Reprocessing (EMDR)	21	18 (12-24)	62.13
10. CBT without a trauma focus	9	14 (7-23)	61.96
11. Present Centred Therapy (PCT)	6	20 (13-28)	40.85
12. Supportive Counselling	9	15 (3-32)	87.84
13. Observed and Experiential Integration (OEI)	1	0	Not applicable
14. Interpersonal Psychotherapy (IPT)	1	15 (6-30)	Not applicable
15. Psychodynamic Psychotherapy	1	14	Not applicable
16. REM Desensitization	1	38	Not applicable
17. Emotional Freedom Technique (EFT)	1	39	Not applicable
18. Dialogical Exposure Therapy (DET)	1	12	Not applicable
19. Internet-based CBT	3	16 (8-26)	32.12
20. Relaxation Training	8	10 (3-19)	56.80
21. Group CBT with a Trauma Focus (group CBT-T)	9	24 (16-33)	76.29
22. Group Present Centred Therapy (PCT)	3	14 (11-18)	0.00
23. Group and Individual CBT-T	1	22	Not applicable
24. Group Stabilizing Treatment	1	18	Not applicable
25. Group Interpersonal Psychotherapy	1	38	Not applicable

26. Group Supportive Counselling	1	3	Not applicable
27. Couples CBT-T	2	22 (11-36)	0.00
28. Psychoeducation	3	1 (0-7)	0.00
29. Couples Psychoeducation	3	12 (3-25)	64.00
30. Internet-based psychoeducation	1	7	Not applicable
31. Waitlist	53	11 (8-15)	65.43
32. Treatment usual	14	13 (7-19)	61.37
33. Minimal attention/symptom monitoring	8	13 (2-32)	92.30

**Table 3: Results of the meta-analyses of dropout**



Variable	$\beta$ (95% confidence intervals)	P
Trauma focus	0.069 (0.011-0.127)	0.021
Recruitment from clinical services	-0.028 (-0.087 – 0.030)	0.341
Delivered in a group format	-0.022 (-0.096 – 0.523)	0.564
Sample drawn from military population	0.032 (-0.023 – 0.087)	0.251
Sexual trauma	0.040 (-0.049 – 0.130)	0.376
% Female	0.040 (-0.049 – 0.130)	0.376
% University Educated	0.001 (-0.003 – 0.001)	0.208

**Table 4: Meta-regression of study-level variables on dropout from all active psychological therapies**

Trauma-focus coded as 0 = non-trauma focused, 1 = trauma focused; recruitment method coded as 0 = not recruited from clinical services, 1 = recruited from clinical services; delivered in a group format coded as 0 = not delivered in a group format, 1 = not delivered in a group format; sample drawn from military population coded 0 = not from a military population; 1 = from a military population; sexual trauma coded 0 = not a sexual trauma; 1 = a sexual trauma.




Figure 1: Study flow diagram






Figure 2: Funnel plot