

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/128234/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Abis, Laura, Dimitritatos, Nikolaos, Meenakshisundaram, Sankar, Freakley, Simon J. and Hutchings, Graham J. 2020. The effect of polymer addition on base catalysed glycerol oxidation using gold and gold-palladium bimetallic catalysts. *Topics in Catalysis* 63 , pp. 394-402. 10.1007/s11244-019-01212-y filefile

Publishers page: <http://dx.doi.org/10.1007/s11244-019-01212-y>
<<http://dx.doi.org/10.1007/s11244-019-01212-y>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Supplementary Information

The Effect of Polymer Addition on Base Catalysed Glycerol Oxidation Using Gold and Gold-Palladium Bimetallic Catalysts.

Laura Abis,¹ Nikolaos Dimitritatos,^{1,2} Meenakshisundaram Sankar,¹ Simon J. Freakley^{1,3*} and Graham J. Hutchings^{1*}

¹ Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff, CF10 3AT, UK.

² Department of Industrial Chemistry, Alma Mater Studiorum-University of Bologna, Viale Risorgimento, 440136, Bologna, Italy.

³ Department of Chemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

s.freakley@bath.ac.uk, hutch@cardiff.ac.uk




Figure S1 - Comparison of the glycerol oxidation activity for 1% wt AuPd/TiO₂ catalysts prepared stabiliser-free, PVA and PVP. *Reaction conditions:* glycerol 0.6 M, NaOH 1.2 M, 110 mg catalyst, 3 bar O₂, 60 °C, 1200 rpm, Gly/Au=500, total volume 10 mL. *Legend:* SF (●), PVA (▲), PVP (■); dotted lines - tartrate (TA) selectivity; dashed lines -glycerate (GA) selectivity.




Figure S2 – Glyceric acid oxidation conversion and selectivity profile for 1% wt AuPd/TiO₂-PVA catalyst. *Reaction conditions:* 110 mg catalyst, 60 °C, 3 bar O₂, 1200 rpm, GA/metal(s) = 500:1(mol), NaOH/GA = 2:1 (mol), total volume 10 mL. *Legend:* □ GA ◆ TA (bold line); ● Glycolate (GLYA) ■ Oxalate (OXA) ▲ Formate (FA) (dotted lines)

