

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/129120/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Taylor, Peter N. and Lazarus, John H. 2019. Hypothyroidism in pregnancy. *Endocrinology and Metabolism Clinics of North America* 48 (3) , pp. 547-556. 10.1016/j.ecl.2019.05.010

Publishers page: <http://dx.doi.org/10.1016/j.ecl.2019.05.010>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



CLINICS TITLE PAGE TEMPLATE

ARTICLE TITLE

Hypothyroidism in Pregnancy

AUTHOR NAMES AND DEGREES

Peter N Taylor PhD, John H Lazarus MD

Full name and degrees of each author exactly as they should appear in print. REMINDER: Each individual listed as an author is expected to have contributed to the article to a significant extent in line with ICMJE guidelines. Please see the [Clinics Authorship Guidelines](#) for more information.

AUTHOR AFFILIATIONS

Thyroid Research Group, Systems Immunity Research Institute, Cardiff University School of Medicine, Cardiff, UK.

Position, department, institution name, city, state, and country for each author listed above

AUTHOR CONTACT INFORMATION

C2 link corridor, UHW, Heath Park, Cardiff CF14 4XN

taylorpn@cardiff.ac.uk

lazarus@cardiff.ac.uk

Mailing address and email address for each author listed above

CORRESPONDING AUTHOR

taylorpn@cardiff.ac.uk

C2 link corridor, UHW, Heath Park, Cardiff CF14 4XN

Mailing address and email address for one author who will receive article proofs

DISCLOSURE STATEMENT

No conflicts of interest to disclose

Disclosure of any relationship with a commercial company that has a direct financial interest in subject matter or materials discussed in article or with a company making a competing product

KEYWORDS

Hypothyroidism

Pregnancy

Levothyroxine

Screening

TPO

Iodine

4-8 keywords to enhance online search results

KEY POINTS

- Hypothyroidism is common in women of child-bearing age **and** commonly associated with adverse obstetric and offspring outcomes.
- Pregnancy places substantial additional demands on the thyroid axis and women who have normal thyroid function prior to pregnancy may have insufficient reserve for pregnancy especially those with auto-immune thyroid disease
- Levothyroxine is the treatment for hypothyroidism; in women established on levothyroxine prior to pregnancy doses will need increasing.
- Screening for thyroid disease in pregnancy is being undertaken in some countries although this is a major debate in thyroidology at present.

3 to 5 bullet points of approximately 25 words each that summarize the main ideas of your article. Key points appear at the very beginning of your article in print and online.

SYNOPSIS

Brief summary of your article (100 to 150 words; no references or figures/tables). The synopsis appears only in the table of contents, and is often used by indexing services such as PubMed.

Thyroid hormone is essential for pregnancy and ensuring foetal development. Pregnancy also places substantial demands on the thyroid axis. Overt hypothyroidism is associated with substantial adverse obstetric and offspring outcomes and requires treatment. Borderline thyroid dysfunction (subclinical hypothyroidism and isolated hypothyroxinemia) are common in women and are associated with adverse obstetric and offspring outcomes although benefits of screening for and treating borderline thyroid function are less clear. Many women are established on thyroid hormone replacement prior to pregnancy and doses will need increasing during pregnancy. Care needs to be taken to prevent over-replacement as this may offset benefits observed for IQ and may increase the risk of ADHD symptoms. Universal thyroid screening in pregnancy is now being undertaken in several countries although it remains a matter of intense debate

Introduction

Adequate thyroid hormone is essential for maintaining a pregnancy and optimal fetal development¹. During the first half of pregnancy, the fetus is entirely dependent on maternal thyroid hormone. Thyroid disorders, particularly low thyroid function are common in women of childbearing age and are frequently encountered in antenatal clinics. It is well established that overt hypothyroidism results in substantially higher risks of adverse pregnancy and offspring outcomes and all endocrine and obstetric societies recommend treating this. Iodine is essential for thyroid hormone production so adequacy here is vital during pregnancy and in women of reproductive age. There is now increasing evidence that even borderline thyroid function and thyroid auto-immunity are also associated with adverse outcomes raising the possibility that universal thyroid screening in pregnancy should be considered². Furthermore, pregnancy places additional demands on the thyroid axis, so women with adequate thyroid hormone prior to pregnancy may fail to meet the additional demands of pregnancy.

Clinically thyroid function is assessed by measuring the pituitary hormone - thyroid stimulating hormone (TSH) and thyroid hormone levels. The complex inverse relationship between TSH and thyroid hormone results in small changes in thyroid hormone levels causing larger changes in TSH. It is essential to measure thyroid hormone levels as well as TSH in order to differentiate between overt and subclinical thyroid disease.

Iodine status and thyroid function

Iodine is essential for the synthesis of thyroid hormones³. The recommended daily iodine intake in pregnancy has recently been increased to 250 µg/day which implies a urinary iodine excretion of 150– 250 µg/day as being adequate^{4,5}. Iodine deficiency during pregnancy is associated with maternal goitre and results eventually in a reduced circulating maternal thyroxine concentration. These effects are preventable by iodine supplementation⁶ and benefits of this are observed only in both areas of severe iodine deficiency (24-h urinary iodine less than 50 µg) and also in areas of mild to moderate deficiency⁷. Children born to mothers with profound iodine deficiency show impaired neuro-intellectual development, sometimes to the extreme of cretinism in severely deficient states. These effects can be reduced by iodine administration before and even during gestation and this should be performed in areas of moderate to severe iodine deficiency³. It is noteworthy that even mild-moderate iodine deficiency in pregnant women is associated with adverse maternal effects including goitre⁸ and lower IQ in offspring⁹. Although a recent trial did not show benefit on iodine supplementation during pregnancy in areas of mild iodine deficiency¹⁰ further studies are required. Assessment of iodine status is challenging and urinary iodine remains an imperfect marker of iodine status serum thyroglobulin may become a useful biomarker of iodine status. Endocrine disruptors such as perchlorate, may exacerbate iodine deficiency and may also have a deleterious effect on offspring neurodevelopment^{11,12}.

Impact of pregnancy on the maternal thyroid status

Pregnancy places substantial demands on the maternal thyroid axis these are summarized in **Table 1 and Figure 1**. The fetal thyroid is not functional until 18-20 weeks gestation so additional maternal thyroid hormone is required to meet this demand. Additionally there is increased thyroxine binding globulin and increased thyroid hormone degradation by placental type 3 deiodinase¹³ which also increases demands. There is also increased urinary iodine excretion during pregnancy. Urinary iodine excretion in pregnancy is maximal in the first trimester followed by a decline in the second and third trimesters. It is increasingly recognized that countries may have iodine sufficiency in the general population, but be insufficient for pregnancy¹⁴. As a result some countries can be iodine sufficient in the general adult population, but inadequate for pregnant women, the UK and Russia are notable examples of this¹⁵. This iodine deficiency can result in increased risk of maternal goitre as well as hypothyroidism. The pregnancy

hormone beta hCG does stimulate the thyroid to produce thyroid hormone which will assist with meeting the increased demands on the thyroid axis in pregnancy. Recent data suggests that TPO antibody positive women may have an impaired thyroidal response to hCG¹⁶. This may explain why women with thyroid auto-immunity with positive thyroid peroxidase antibodies (TPO) have increased risk of adverse obstetric outcomes which can be independent of thyroid status.

Assessment of thyroid status during pregnancy

It is important to recognize that assessment of thyroid function in pregnancy is more complex than the general adult population. The above physiological changes in pregnancy have a substantial effect on the interpretation of thyroid function tests in pregnancy and result in a downward shift of TSH reference intervals. A pragmatic upper limit of 2.5 mU/L was previously advocated for TSH but recent data now shows this threshold to be too low¹⁷. One approach suggested in the current guidelines of the American Thyroid Association (ATA) is to set the pregnancy reference range at 0.5 mU/L and 0.4 mU/L below the upper and lower non-pregnant reference range respectively, reflecting the anticipated magnitude of the TSH drop¹⁸.

Current ATA guidelines advocate the use of pregnancy specific, local population-based reference ranges where possible;¹⁸ however it is well recognized that such data are not widely available. Assessment of thyroid function in pregnancy is also best done according to pregnancy specific reference-ranges calculated in a population of pregnant women free of key factors that interfere with thyroid function including women with known thyroid disease (including thyroid auto-immunity), use of thyroid altering medication, known iodine deficiency and high hCG states (twin pregnancies or IVF conception). However, it is also well established that normal thyroid status changes over pregnancy and accurate classification of thyroid function in pregnant women requires the use of gestational-age specific reference ranges¹⁹. There is also growing evidence that specific reference ranges may be considered based on ethnicity²⁰, BMI²¹ and parity²¹.

With respect to FT4, the most commonly used immunoassays are prone to biases in pregnancy due to changes in thyroid binding globulin and albumin concentrations^{18,22}. Total T4 is a more predictable alternative as it typically rises by 50% of baseline values attaining peak levels by 16 weeks gestation¹⁸. However Total T4 is less clearly associated with adverse outcomes so many clinicians prefer the use of FT4. More robust methods such as equilibrium dialysis, liquid chromatography-tandem mass spectrometry (LC-MS/MS), or FT4 index calculations are cumbersome to perform and not routinely used¹⁸.

Classification and epidemiology of thyroid abnormalities

Overt hypothyroidism in pregnancy is defined as a TSH level above the pregnancy specific reference range and a low level of FT4. Although women established on levothyroxine make -up around 1% of pregnancies, newly diagnosed overt hypothyroidism occurs in approximately 0.2-0.6% of pregnant women^{17,23}. Treatment is mandatory as it is associated with profound adverse obstetric and offspring development outcomes. The milder forms of low thyroid dysfunction are subclinical hypothyroidism (TSH above the population's pregnancy reference range, but normal FT4) and isolated hypothyroxinemia (**IH**) (Normal TSH with FT4 in the lowest 2.5% of the reference range) and these are more common. Subclinical hypothyroidism with an elevated TSH and normal FT4 concentrations—can occur in up to 18% of pregnancies depending on the precise definition and TSH **assay** used. Observational studies have indicated that subclinical hypothyroidism is associated with adverse pregnancy outcomes, but not offspring neuro development. In contrast IH is associated with adverse outcomes with regard to offspring IQ and behavior, but less consistent associations have been observed for pregnancy outcomes.

Overt hypothyroidism

Overt hypothyroidism has been repeatedly associated with substantial adverse impacts on obstetric outcomes and fetal neuro development. Adverse obstetric outcomes include fetal loss, premature delivery, low birthweight and pre-eclampsia^{23,24}. A large case-control study demonstrated that overt hypothyroidism resulted in a 7-point lower IQ than women with normal thyroid function²⁵. It is worth noting, that women with adequately treated hypothyroidism have no higher risk of pregnancy complications, unlike those with untreated hypothyroidism indicating that thyroid status is a clearly reversible risk factor for adverse obstetric outcomes.

Subclinical hypothyroidism

Numerous studies have identified that subclinical hypothyroidism is associated with adverse pregnancy similar to overt hypothyroidism but as to be expected risk effects are more marginal. From meta-analyses subclinical hypothyroidism the following outcomes were all statistically significant for adverse outcomes with (95%CI) provided. The odds of adverse outcomes are as follows - miscarriage (1.90 - 2.01) preterm delivery (1.20 - 1.81) growth restriction (1.54 - 3.36) pre-eclampsia (1.30 - 2.24) and gestational diabetes (1.28 - 4.33)²⁶. One problem has been with these analyses is that many studies have defined subclinical hypothyroidism differently. What is clear is that subclinical hypothyroidism does not appear to be associated with adverse neuro-behavioral outcomes in the offspring.

Isolated hypothyroxinemia

IH was originally considered to be a pregnancy-specific disease, reflecting a state of mild iodine deficiency. However, this has now been called into question as it also occurs in iodine sufficient areas. Other risk factors for IH include iron status, BMI and placental angiogenic factors¹. IH has been associated with impaired offspring developmental outcomes²⁷ and has also been associated with offspring verbal delay²⁸ autism²⁹ and ADHD³⁰. In contrast to subclinical hypothyroidism it has not been consistently associated with adverse obstetric outcomes.

Thyroid auto-immunity

Anti-thyroid peroxidase antibodies are found in around 10% of otherwise normal pregnant women. They are a marker of thyroid auto-immunity and the main risk factor for thyroid dysfunction in pregnancy and in the postpartum period. Even in euthyroid women TPOAb positivity is associated with premature delivery and fetal loss although the exact mechanism remains unclear^{31,32}. One randomized controlled trial has shown that levothyroxine administration reduced the risk of fetal loss and premature delivery in euthyroid TPOAb-positive women³³. A later study failed to replicate the benefits for fetal loss but did find a reduction in pre-term delivery³⁴. Women who are TPOAb positive are at an increased risk of thyroid-related adverse pregnancy outcomes, its interaction with higher TSH levels means it is likely that this risk is substantially increased in individuals who also have elevated TSH levels or those in the higher end of the normal range. As highlighted earlier women who are TPOAb positive have an impaired thyroidal response to beta hCG compared to women who are TPOAb negative this may explain its association with adverse obstetric outcomes and synergistic impact with higher TSH levels.

Trials of Treatment with levothyroxine

All endocrine and obstetric societies recommend treating *overt hypothyroidism* in pregnancy although there have been no recent trials in this area, but to have a control group here would be unethical. To date three large randomized controlled trials have investigated the effects of screening for and treating borderline low thyroid function in pregnancy: these are the controlled antenatal thyroid screening (CATS) study³⁵, a study by Casey *et al.*³⁶ and a recent study by Nazarpour *et al.*³⁴. These trials are summarized in **Table 2**. The Nazarpour trial focused on preterm delivery and identified that levothyroxine may reduce the risk of preterm delivery in individuals with TSH

levels > 4.0 mU/l (RR: 0.38; 95% CI: 0.15 to 0.98; p= 0.04)³⁴. Offspring IQ was not assessed in this study. The two other large randomized controlled trials CATS³⁵ and the study by Casey et al³⁷ studied the effects of screening and treating borderline low thyroid function in pregnancy on offspring IQ and pregnancy outcomes. Neither study showed any beneficial effects of treatment on offspring IQ^{35,37}. Reasons for failure to establish benefit aside from no treatment benefit, include the relatively late initiation of treatment (particularly in the Casey study) and early age of IQ assessment (particularly in the CATS study). It is worth highlighting that neurological development is crucial in the first 12 weeks of pregnancy. Follow on analysis of the CATS study revealed no apparent benefit of treatment at age 9 although this was only performed in a subset also showed no treatment benefit. Work in the follow-up CATS study also identified levothyroxine over-treatment may increase the risk of autism symptoms³⁸. More recent analysis using data linkage and the majority of the CATS cohort (to include those with normal thyroid function) identified that levothyroxine treatment significantly reduced the risk of miscarriage/still birth³⁸. Similarly, another prospective study did identify that levothyroxine reduced the risk of miscarriage and pre-term birth³³. However no benefit was observed with levothyroxine on obstetric outcomes in the Casey study although again this may be due to the relatively late initiation of treatment³⁷.

Treatment of established hypothyroidism in pregnancy

Approximately 1% of pregnant women are established on levothyroxine prior to pregnancy. In women established on levothyroxine prior to pregnancy, initial control in pregnancy is often sub-optimal³⁹ and may be associated with increased odds of foetal loss³⁹, and optimization should ideally occur prior to conception⁴⁰. Women of child-bearing age on levothyroxine should be reminded to increase their pre-conception dose of levothyroxine by 30-50% as soon as pregnancy is confirmed. One approach is to take double their usual levothyroxine dose on 2 days of the week⁴¹. It is worth noting, that dose increases are often higher in post-surgical or post-ablative hypothyroidism. Women on levothyroxine should also be aware of important drug interactions with iron supplements and proton pump inhibitors in particular which can impair levothyroxine absorption. Other drugs to be aware of increase which increase levothyroxine clearance include carbamazepine, rifampicin and valproate. If overt hypothyroidism is diagnosed for the first time in pregnancy, we recommend starting a weight-based dose of 2mcg/kg a day.

Close monitoring of thyroid function is essential in pregnancy we recommend to check thyroid function early in the first trimester and every 4-6 weeks thereafter. One should ideally aim for a TSH < 2.5mU/l in the first trimester and < 3 mU/l in later pregnancy. One should aim to ensure that the FT4 level is not too high and aim for the upper half of the reference range. There is some evidence that over-replacement with levothyroxine may increase ADHD symptoms and the relationship of FT4 and IQ is “U shaped” with both low -normal and high-normal FT4 are associated with lower IQ⁴². After delivery we suggest returning levothyroxine to pre-conception dose and to recheck thyroid function at 6 weeks post-partum.

Screening for and treating low thyroid function in pregnancy

Universal thyroid screening in pregnancy remains contentious². It is noteworthy that screening for and treating low thyroid function meets almost all the screening criteria laid down by Wilson and Jungner⁴³. In particular it is an important asymptomatic health problem, with an accepted treatment and a suitable diagnostic test. It is also economically viable⁴⁴. However, at present it is still unclear what TSH threshold should be used for treatment in pregnancy and whether there should be a differential threshold based on TPO status. The criterion 8 “there should be an agreed policy on who to treat” is therefore not met. However, as there is widespread variation in clinical practice at present, this should not necessarily prevent the introduction of universal thyroid screening and indeed regular audit and review may establish more definitive treatment thresholds. Current ATA guidance takes into account the potential for interaction by TPO antibody status in its guidance, but did not recommended for or against

universal thyroid screening¹⁸. This has not prevented some countries including Spain China and Poland from implementing universal thyroid screening. It is however accepted that targeted screening should be performed in those women at high risk for thyroid disease^{18,45}. Women deemed at high risk for thyroid disease in pregnancy include those with previous thyroid disease, a visible goitre, symptoms suggestive of hypo or hyperthyroidism, those with a family history of thyroid disease, those with a history of type 1 diabetes or other auto-immune conditions including positive thyroid antibodies, and women with a previous history of infertility, fetal loss or infertility. However this targeted approach would miss approximately one third of women with significant thyroid dysfunction⁴⁶. This high risk screening approach has been further called into question as universal thyroid screening in pregnancy appears to be cost-effective; screening solely for overt hypothyroidism also had a cost-effectiveness ratio of \$6,776/QALY (quality adjusted life-year)⁴⁴ which is favorable compared to gestational diabetes mellitus screening (\$12,078/QALY) and is substantially below the \$50,000/QALY figure used in the United States as a criterion for screening decisions. This work also indicated universal screening appears to be more cost-effective than targeted screening.

Conclusion

Despite the dramatic recent advances in our knowledge of thyroid physiology in pregnancy and the consequences of its variation, further research is still required. **Delete need In** particular prospective trials of *early* screening of thyroid function in pregnancy with both obstetric and developmental outcomes assessed. Continued advocacy of iodine sufficiency in pregnancy and assessment of endocrine disruptors are also needed. In current practice monitoring of thyroid function in early pregnancy needs urgent optimization.

Table 1 Physiological changes that influence thyroid function in pregnancy

Physiological change	Effect on thyroid function test results	Impact on interpretation of thyroid function tests
↑ Thyroxine binding globulin	↑ Serum Total T3 and T4 concentrations	Total thyroid hormone levels may be misleading -need to rely on free thyroid hormone levels
↑ human chorionic gonadotrophin secretion	↑ Free T4 and ↓ TSH	High human chorionic gonadotrophin levels may result in gestational thyrotoxicosis. This usually only requires symptomatic treatment but needs to be distinguished from pathological thyroid disease. Response possibly impaired in TPO antibody positive women
↑ Iodine excretion	↓ Thyroid hormone production in iodine deficient areas	Need to be mindful of iodine deficiency and ensure optimal intake ideally prior to conception
↑ Plasma volume	↑ T3 and T4 pool size	
Increased Type 3 5-deiodinase (inner ring deiodination) activity from the placental	↑ T3 and T4 degradation	May explain in part the increasing thyroid demand in pregnancy
Thyroid enlargement (in some women)	Increased thyroglobulin	Small goitres are common in pregnancy, but may be a sign of low thyroid function so merits thyroid function testing

Adapted from the Oxford Textbook of Diabetes and Endocrinology, Chapter 9.1 General considerations relating to thyroid disease in pregnancy by Peter N Taylor, LDKE Premawardhana, John H Lazarus

Table 2

	CATS	Casey	Nazarpour
Year	2012	2017	2018
Countries in trial	UK, Italy	USA	Iran
Number with low thyroid function randomised	794	677	366
Placebo-controlled	No	Yes	No
Gestational age at recruitment (weeks)	Median (IQR) Screening 12.3 (11.6 - 13.6) Controls 12,3 (11.6 - 13.5)	Mean (SD) Screening 16.6 (3.0) Controls 16.7 (3.0)	Mean (SD) Screening 11.4 (4.1) Controls 12.2 (4.3)
Baseline TSH mU/l	Median (IQR) Screening UK 3.8 (1.5-4.7) Screening Italy 3.1 (1.3-4.0) Controls UK 3.2 (1.2 - 4.2) Controls Italy 2,4 (1.3-3.9)	Mean (95%CI) Screening 4.5 (4.4-4.7) Control 4.3 (4.2 -4.5)	Median (IQR) Screening 3.8 (2.8-4.8)
Outcomes assessed	IQ age 3, IQ age 9 Obstetric outcomes (analysed later and only in UK participants)	Pregnancy outcomes, offspring IQ and behaviour	Preterm delivery
Benefit of levothyroxine	No benefit with regard to IQ. Potential reduction in foetal loss.	No benefits observed with regard to pregnancy outcomes and offspring IQ	May reduce pre-term delivery at TSH levels > 4.0 mU/l

Figure 1 Effects of Pregnancy on thyroid function





Figure 2 TSH FT4 and hCG levels over pregnancy.



References

- 1 Korevaar, T. I. M., Medici, M., Visser, T. J. & Peeters, R. P. Thyroid disease in pregnancy: new insights in diagnosis and clinical management. *Nature reviews. Endocrinology* **13**, 610-622, doi:10.1038/nrendo.2017.93 (2017).
- 2 Taylor, P. N. *et al.* Thyroid Screening in Early Pregnancy: Pros and Cons. *Frontiers in endocrinology* **9**, 626, doi:10.3389/fendo.2018.00626 (2018).
- 3 Zimmermann, M. B. Iodine deficiency. *Endocr Rev* **30**, 376-408, doi:10.1210/er.2009-0011 (2009).
- 4 Zimmermann, M. B., Gizak, M., Abbott, K., Andersson, M. & Lazarus, J. H. Iodine deficiency in pregnant women in Europe. *The lancet. Diabetes & endocrinology* **3**, 672-674, doi:10.1016/s2213-8587(15)00263-6 (2015).
- 5 Taylor, P., Vaidya, B., Taylor, P. & Vaidya, B. Iodine Supplementation in Pregnancy - is it time? *Clin Endocrinol (Oxf)*, doi:10.1111/cen.13065 (2016).
- 6 Zimmermann, M. B. Iodine deficiency in pregnancy and the effects of maternal iodine supplementation on the offspring: a review. *The American journal of clinical nutrition* **89**, 668S-672S, doi:10.3945/ajcn.2008.26811C (2009).
- 7 Taylor, P. N., Okosieme, O. E., Dayan, C. M. & Lazarus, J. H. Therapy of endocrine disease: Impact of iodine supplementation in mild-to-moderate iodine deficiency: systematic review and meta-analysis. *European journal of endocrinology / European Federation of Endocrine Societies* **170**, R1-R15, doi:10.1530/eje-13-0651 (2014).
- 8 Taylor, P. N., Razvi, S., Pearce, S. H. & Dayan, C. M. A review of the clinical consequences of variation in thyroid function within the reference range. *J Clin Endocrinol Metab* **98**, 3562-3571, doi:10.1210/jc.2013-1315 (2013).
- 9 Bath, S. C., Steer, C. D., Golding, J., Emmett, P. & Rayman, M. P. Effect of inadequate iodine status in UK pregnant women on cognitive outcomes in their children: results from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Lancet* **382**, 331-337, doi:10.1016/s0140-6736(13)60436-5 (2013).
- 10 Gowachirapant, S. *et al.* Effect of iodine supplementation in pregnant women on child neurodevelopment: a randomised, double-blind, placebo-controlled trial. *The lancet. Diabetes & endocrinology* **5**, 853-863, doi:10.1016/s2213-8587(17)30332-7 (2017).
- 11 Bellanger, M., Demeneix, B., Grandjean, P., Zoeller, R. T. & Trasande, L. Neurobehavioral deficits, diseases, and associated costs of exposure to endocrine-disrupting chemicals in the European Union. *J Clin Endocrinol Metab* **100**, 1256-1266, doi:10.1210/jc.2014-4323 (2015).
- 12 Taylor, P. N. *et al.* **? ref ok like this or do they want 3 authors et al** Maternal perchlorate levels in women with borderline thyroid function during pregnancy and the cognitive development of their offspring: data from the Controlled Antenatal Thyroid Study. *J Clin Endocrinol Metab* **99**, 4291-4298, doi:10.1210/jc.2014-1901 (2014).
- 13 Brent, G. A. Maternal thyroid function: interpretation of thyroid function tests in pregnancy. *Clinical obstetrics and gynecology* **40**, 3-15 (1997).
- 14 Taylor, P. N. *et al.* Global epidemiology of hyperthyroidism and hypothyroidism. *Nature reviews. Endocrinology*, doi:10.1038/nrendo.2018.18 (2018).
- 15 Taylor, P. N. *et al.* Global epidemiology of hyperthyroidism and hypothyroidism. *Nature reviews. Endocrinology* **14**, 301-316, doi:10.1038/nrendo.2018.18 (2018).
- 16 Korevaar, T. I. *et al.* Thyroid Autoimmunity Impairs the Thyroidal Response to Human Chorionic Gonadotropin: Two Population-Based Prospective Cohort Studies. *J Clin Endocrinol Metab* **102**, 69-77, doi:10.1210/jc.2016-2942 (2017).
- 17 Medici, M., Korevaar, T. I., Visser, W. E., Visser, T. J. & Peeters, R. P. Thyroid function in pregnancy: what is normal? *Clinical chemistry* **61**, 704-713, doi:10.1373/clinchem.2014.236646 (2015).
- 18 Alexander, E. K. *et al.* 2017 Guidelines of the American Thyroid Association for the Diagnosis and Management of Thyroid Disease During Pregnancy and the Postpartum. *Thyroid* **27**, 315-389, doi:10.1089/thy.2016.0457 (2017).

- 19 Stricker, R. *et al.* Evaluation of maternal thyroid function during pregnancy: the importance of using gestational age-specific reference intervals. *European journal of endocrinology / European Federation of Endocrine Societies* **157**, 509-514, doi:10.1530/eje-07-0249 (2007).
- 20 Korevaar, T. I. *et al.* Ethnic differences in maternal thyroid parameters during pregnancy: the Generation R study. *J Clin Endocrinol Metab* **98**, 3678-3686, doi:10.1210/jc.2013-2005 (2013).
- 21 Korevaar, T. I. *et al.* Stimulation of Thyroid Function by Human Chorionic Gonadotropin During Pregnancy: A Risk Factor for Thyroid Disease and a Mechanism for Known Risk Factors. *Thyroid* **27**, 440-450, doi:10.1089/thy.2016.0527 (2017).
- 22 Ross, D. S. *et al.* 2016 American Thyroid Association Guidelines for Diagnosis and Management of Hyperthyroidism and Other Causes of Thyrotoxicosis. *Thyroid* **26**, 1343-1421, doi:10.1089/thy.2016.0229 (2016).
- 23 Krassas, G. E., Poppe, K. & Glinoeer, D. Thyroid Function and Human Reproductive Health. *Endocrine Reviews* **31**, 702-755, doi:10.1210/er.2009-0041 (2010).
- 24 van den Boogaard, E. *et al.* Significance of (sub)clinical thyroid dysfunction and thyroid autoimmunity before conception and in early pregnancy: a systematic review. *Human reproduction update* **17**, 605-619, doi:10.1093/humupd/dmr024 (2011).
- 25 Haddow, J. E. *et al.* Maternal thyroid deficiency during pregnancy and subsequent neuropsychological development of the child. *N Engl J Med* **341**, 549-555, doi:10.1056/nejm199908193410801 (1999).
- 26 Velasco, I. & Taylor, P. Identifying and treating subclinical thyroid dysfunction in pregnancy: emerging controversies. *European journal of endocrinology / European Federation of Endocrine Societies* **178**, D1-d12, doi:10.1530/eje-17-0598 (2018).
- 27 Korevaar, T. I. *et al.* Association of maternal thyroid function during early pregnancy with offspring IQ and brain morphology in childhood: a population-based prospective cohort study. *The lancet. Diabetes & endocrinology* **4**, 35-43, doi:10.1016/s2213-8587(15)00327-7 (2016).
- 28 Henrichs, J. *et al.* Maternal thyroid function during early pregnancy and cognitive functioning in early childhood: the generation R study. *J Clin Endocrinol Metab* **95**, 4227-4234, doi:10.1210/jc.2010-0415 (2010).
- 29 Roman, G. C. *et al.* Association of gestational maternal hypothyroxinemia and increased autism risk. *Annals of neurology*, doi:10.1002/ana.23976 (2013).
- 30 Vermiglio, F. *et al.* Attention deficit and hyperactivity disorders in the offspring of mothers exposed to mild-moderate iodine deficiency: a possible novel iodine deficiency disorder in developed countries. *J Clin Endocrinol Metab* **89**, 6054-6060, doi:10.1210/jc.2004-0571 (2004).
- 31 Korevaar, T. I. *et al.* Hypothyroxinemia and TPO-antibody positivity are risk factors for premature delivery: the generation R study. *J Clin Endocrinol Metab* **98**, 4382-4390, doi:10.1210/jc.2013-2855 (2013).
- 32 Thangaratinam, S. *et al.* Association between thyroid autoantibodies and miscarriage and preterm birth: meta-analysis of evidence. *Bmj* **342**, d2616, doi:10.1136/bmj.d2616 (2011).
- 33 Negro, R. *et al.* Levothyroxine treatment in euthyroid pregnant women with autoimmune thyroid disease: effects on obstetrical complications. *J Clin Endocrinol Metab* **91**, 2587-2591, doi:10.1210/jc.2005-1603 (2006).
- 34 Nazarpour, S. *et al.* Effects of levothyroxine treatment on pregnancy outcomes in pregnant women with autoimmune thyroid disease. *European journal of endocrinology / European Federation of Endocrine Societies* **176**, 253-265, doi:10.1530/eje-16-0548 (2017).
- 35 Lazarus, J. H. *et al.* Antenatal thyroid screening and childhood cognitive function. *N Engl J Med* **366**, 493-501, doi:10.1056/NEJMoa1106104 (2012).
- 36 Casey, B. M. *et al.* Treatment of Subclinical Hypothyroidism or Hypothyroxinemia in Pregnancy. *N Engl J Med* **376**, 815-825, doi:10.1056/NEJMoa1606205 (2017).
- 37 Casey, B. M. *et al.* Treatment of Subclinical Hypothyroidism or Hypothyroxinemia in Pregnancy. *New England Journal of Medicine* **376**, 815-825, doi:10.1056/NEJMoa1606205 (2017).

- 38 Bartalena, L. *et al.* Meeting abstracts from the 64th British Thyroid Association Annual Meeting. *Thyroid Research* **10**, 2, doi:10.1186/s13044-016-0036-8 (2017).
- 39 Taylor, P. N. *et al.* TSH Levels and Risk of Miscarriage in Women on Long-Term Levothyroxine: A Community-Based Study. *J Clin Endocrinol Metab*, jc20141954, doi:10.1210/jc.2014-1954 (2014).
- 40 Okosieme, O. E., Khan, I. & Taylor, P. N. Preconception management of thyroid dysfunction. *Clin Endocrinol (Oxf)* **89**, 269-279, doi:10.1111/cen.13731 (2018).
- 41 Yassa, L., Marqusee, E., Fawcett, R. & Alexander, E. K. Thyroid hormone early adjustment in pregnancy (the THERAPY) trial. *J Clin Endocrinol Metab* **95**, 3234-3241, doi:10.1210/jc.2010-0013 (2010).
- 42 Korevaar, T. I. *et al.* Association of maternal thyroid function during early pregnancy with offspring IQ and brain morphology in childhood: a population-based prospective cohort study. *The lancet. Diabetes & endocrinology*, doi:10.1016/s2213-8587(15)00327-7 (2015).
- 43 Wilson, J. & Jungner, G. Principles and practice of screening for disease. Geneva: World Health Organization, 1968. *Public health papers* **34** (2011).
- 44 Dosiou, C. *et al.* Cost-effectiveness of universal and risk-based screening for autoimmune thyroid disease in pregnant women. *J Clin Endocrinol Metab* **97**, 1536-1546, doi:10.1210/jc.2011-2884 (2012).
- 45 De Groot, L. *et al.* Management of Thyroid Dysfunction during Pregnancy and Postpartum: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* **97**, 2543-2565, doi:10.1210/jc.2011-2803 (2012).
- 46 Vaidya, B. *et al.* Detection of thyroid dysfunction in early pregnancy: Universal screening or targeted high-risk case finding? *J Clin Endocrinol Metab* **92**, 203-207, doi:10.1210/jc.2006-1748 (2007).