

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/130424/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Bailey, Sarah, Boivin, Jacky , Ting, Cheong, Bailey, Christopher, Kitson-Reynolds, Ellen and Macklon, Nick
2020. Effective support following recurrent pregnancy loss: A randomized controlled feasibility and acceptability study. *Reproductive BioMedicine Online* 40 (5) , pp. 729-742. 10.1016/j.rbmo.2020.01.022

Publishers page: <http://doi.org/10.1016/j.rbmo.2020.01.022>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Participant	Ethnicity	Age	Number of previous miscarriages	Ongoing pregnancy (OP) or further miscarriage (M)at time of pregnancy
1	White British	38	3	OP
2	White British	37	3	OP
3	White British	42	6	M
4	White British	34	4	M
5	White British	31	4	M
6	White British	39	4	M
7	White British	33	4	M
8	White British	37	3	OP
9	Asian British	29	3	OP
10	White British	38	3	OP
11	White British	34	4	OP
12	White British	36	3	OP
13	White British	40	4	M
14	White British	34	3	OP

Table 2: Sample characteristics of QPE participants.