

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/133373/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Yevglevskis, Maksims, Lee, Guat L., Nathubhai, Amit, Petrova, Yoana D., James, Tony D., Threadgill, Michael D., Woodman, Timothy J. and Lloyd, Matthew D. 2018. Structure-activity relationships of rationally designed AMACR 1A inhibitors. *Bioorganic Chemistry* 79 , pp. 145-154.
10.1016/j.bioorg.2018.04.024

Publishers page: <http://dx.doi.org/10.1016/j.bioorg.2018.04.024>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Structure-activity relationships of rationally designed AMACR 1A inhibitors

Maksims Yevglevskis^a, Guat L. Lee^a, Amit Nathubhai^a, Yoana D. Petrova^a, Tony D. James^b,
Michael D. Threadgill^a, Timothy J. Woodman^a, Matthew D. Lloyd^a,

^a Drug & Target Discovery, Department of Pharmacy & Pharmacology, University of Bath, Claverton Down, Bath BA2 7AY, UK

^b Department of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, UK

ARTICLE INFO

Keywords:

α -Methylacyl-CoA racemase (AMACR, P504S)

Drug lipophilicity

Enzyme inhibitors

Rational drug design

Structure-activity relationships

ABSTRACT

α -Methylacyl-CoA racemase (AMACR; P504S) is a promising novel drug target for prostate and other cancers. Assaying enzyme activity is difficult due to the reversibility of the 'racemisation' reaction and the difficulties in the separation of epimeric products; consequently few inhibitors have been described and no structure-activity relationship study has been performed. This paper describes the first structure-activity relationship study, in which a series of 23 known and potential rational AMACR inhibitors were evaluated. AMACR was potently inhibited ($IC_{50} = 400\text{--}750\text{ nM}$) by ibuprofenoyl-CoA and derivatives. Potency was positively correlated with inhibitor lipophilicity. AMACR was also inhibited by straight-chain and branched-chain acyl-CoA esters, with potency positively correlating with inhibitor lipophilicity. 2-Methyldecanoyl-CoAs were ca. 3-fold more potent inhibitors than decanoyl-CoA, demonstrating the importance of the 2-methyl group for effective inhibition. Elimination substrates and compounds with modified acyl-CoA cores were also investigated, and shown to be potent inhibitors. These results are the first to demonstrate structure-activity relationships of rational AMACR inhibitors and that potency can be predicted by acyl-CoA lipophilicity. The study also demonstrates the utility of the colorimetric assay for thorough inhibitor characterisation.

1. Introduction

Branched-chain fatty acids (e.g. phytanic acid, pristanic acids) are common components of the human diet, and derivatives of such compounds are used as drug molecules e.g. ibuprofen [1,2]. Degradation of branched-chain fatty acids occurs as the acyl-CoA ester, and the acyl-CoA oxidases and other enzymes involved in β -oxidation have an absolute requirement for S-2-methylacyl-CoAs [3–5]. However, R-2-methylacyl-CoAs are produced from dietary and endogenous fatty acids and these cannot be immediately degraded by β -oxidation. The enzyme α -methylacyl-CoA racemase [1,2] (AMACR; P504S; E.C. 5.1.99.4) catalyses conversion of R-2-methylacyl-CoAs to a near 1:1 epimeric mixture [6,7] by a deprotonation/reprotonation reaction [7,8], probably via an enolate intermediate [9] (this reaction is referred to as "racemisation" [10]). The resulting S-2-methylacyl-CoAs are degraded by β -oxidation whilst the 2R epimers are further processed to the 2S epimers by AMACR [1,2]. AMACR also plays a key role in the *in vivo* pharmacological activation of R-ibuprofen by conversion to S-ibuprofen, via the corresponding acyl-CoA esters [1,2,11]. The S-ibuprofen resulting from this pathway exerts its anti-inflammatory effect by inhibiting


cyclooxygenase-1 and -2 [12].

AMACR protein levels are increased in prostate [13,14] and several other cancers [1,15–18]. Catalytic activity of AMACR is increased by 4- to 10-fold in prostate cancer cells [19,20], with the AMACR 1A splice variant [1,2,21–24] (possessing "racemase" activity [7,10]) showing the most significant increase in expression [19,20]. Reducing AMACR 1A levels using siRNA or shRNA approaches [19,25,26] has been shown to reduce proliferation of prostate cancer cells via a pathway which is synergistic with the use of an androgen receptor antagonist, studies which have validated AMACR 1A as a chemotherapeutic target. Some advanced prostate cancer cell lines revert from castrate-resistant (a.k.a. androgen-independent) growth to androgen-dependent growth upon knockdown of AMACR 1A [26]. Consequently, AMACR has attracted considerable interest as a prostate cancer biomarker [1,2,27] and drug target [25,28–31]. However, the lack of a convenient assay to measure AMACR activity [32,33] has severely hampered the development of AMACR inhibitors as new chemotherapeutic drugs against cancers that over-express AMACR, and consequently only a few rationally designed inhibitors of AMACR [28–30,34] or MCR [31,35] (M. tuberculosis homologue) have been reported. No systematic study of AMACR

Abbreviations: AMACR, α -methylacyl-CoA racemase; CDI, carbonyldiimidazole; DAST, (Diethylamino)sulfur trifluoride; DCC, Dicyclohexylcarbodiimide; DMAP, (dimethylamino) pyridine; MCR, 2-methylacyl-CoA racemase from *M. tuberculosis*; SAR, Structure-activity relationships; THF, tetrahydrofuran; TMSCl, chlorotrimethylsilane

Corresponding author.

E-mail address: M.D.Lloyd@bath.ac.uk (M.D. Lloyd).



Scheme 1. The colorimetric assay for AMACR 1A [32] showing elimination of 2,4-dinitrophenolate 2.

inhibitor SAR has been undertaken [32,33].

Recently, we reported a versatile continuous assay for AMACR based on the utilisation of our novel substrate 1 that can eliminate 2,4-dinitrophenolate 2, which can be monitored by absorbance at 354 nm, and unsaturated product 3 (Scheme 1). This new assay [32] was used to examine the potency of two known acyl-CoA inhibitors (N-dodecyl-N-methylcarbamoyl-CoA 4 [29] and ibuprofenoyl-CoA 5 [6]; Fig. 1) and selected known non-specific protein-modification agents [25]. This paper reports the first systematic examination of SAR for rationally designed acyl-CoA inhibitors of AMACR. Compounds investigated (Fig. 1) include those with aromatic side-chains, (5–11); Straight-chain acyl-CoA esters (12–17); Branched-chain substrates (18–21) and pro-duct 22. Analogues of known inhibitors with modified 2-methylacyl-CoA moieties (4, 23–26) were also examined. The results reveal a correlation between potency and lipophilicity of the inhibitors, consistent with observations on MCR inhibitors [35], the homologous en-zyme from *M. tuberculosis*.

2. Results and discussion

AMACR is a promising novel cancer drug target, but therapeutic development in this field has been slow due to the lack of a robust enzyme assay. Thus, the majority of studies reporting AMACR in-hibitors have largely focussed on rationally designed drugs [28–30,32,33]. In most cases, only one or a few examples of each in-hibitor type has been evaluated, and no systematic SAR study has been performed. Initial SAR studies have been carried out on reversible [31] and irreversible [35] inhibitors of MCR (the *M. tuberculosis* homologue). In addition, different research groups have used different assays during their studies, making it difficult to compare results directly. In this study, the SAR of rational AMACR inhibitors were explored using a series of acyl-CoA esters (Fig. 1). These included compounds previously tested as substrates (5–11 [6,11]; 12–17 [36]; 18–20, 22 [7,10]). Most of these compounds have not been tested as inhibitors with the exception of 5 [28,32,37,38] and 13, 15–17 (which were previously re-ported to be inactive [37,38]). Compound 21 was included as an epimer of 20, and has not been previously reported as a substrate or inhibitor (although the 3-fluoro-2-methylhexadecanoyl-CoA analogues are po-tent inhibitors [28]). Compound 24 is a synthetic intermediate to 25, and has not been previously tested as a substrate or inhibitor. Com-pounds 22, 25 and 26 are intermediates in the subsequent β -oxidation pathway [39], and have not been previously tested as substrates or inhibitors. Analogues of compounds 23, and 25 with different side-chains have been previously tested as inhibitors of AMACR or MCR [9,30]. Compound 4 was previously reported as the most potent AMACR inhibitor [29,32], and is included as an acyl-CoA core analogue.

2.1. Chemical synthesis of acyl-CoA inhibitors

(2S,3S)-3-Fluoro-2-methyldecanoyl-CoA 21 was synthesised by an analogous route to (2R,3R)-3-fluoro-2-methyldecanoyl-CoA 20 [10,28], using an Evans' auxiliary strategy (Scheme 2). Aldol-like reaction of deprotonated 27 with octanal gave the (2S,3R)-3-hydroxy-2-methyl intermediate 28. From here, alcohol 28 was activated and replaced with fluoride with inversion of configuration, using DAST to give 3-fluoro-2-methyl derivative 29. The reaction is thought to go with

inversion of stereochemical configuration (by analogy with the work of Carnell et al. [28]). Removal of the Evans' auxiliary from intermediate 29 provided the carboxylic acid 30 under mild conditions that involved in situ generation of lithium hydroperoxide. Intermediate 30 was sub-sequently converted to the CoA thioester 21 using the standard syn-thetic method with N,N'-carbonyldiimidazole [6,10,11,32,33,40]. This compound was stable in solution in the absence of AMACR, showing that the relative geometry of the α -proton and fluorine atom was syn-(anti- epimers rapidly eliminate fluoride, presumably by an E2 me-chanism [10]).

2-Methylenedecanoyl-CoA 23 was synthesised by an adaptation of the method reported by Morgenroth et al. [30] (Scheme 3). Meldrum's acid 33 was acylated with octanoic acid 32 using DCC activation; the intermediate ketone was reduced to the octyl-Meldrum's acid 31 with sodium triacetoxyborohydride generated in situ. Subsequent reaction of 31 with Eschenmoser's salt gave the 2-methylene ester 34. Base-hy-drolysis furnished the 2-methylene acid 35, which was coupled with CoA-SH by a mixed anhydride approach to give 2-methylenedecanoyl-CoA 23.


2-Methyl-3-oxodecanoyl-CoA 25 was synthesised by the method of Reen et al. [41] from 36 (Scheme 4). The ketone in 37 was protected as the cyclic acetal 36. Hydrolysis of the ester group in 36 gave the cor-responding acid 38, which was then coupled with CoA [6,10,11,32,33,40] to give 24. Acidolysis of the acetal protection pro-vided 25.

2S,3R-3-Hydroxy-2-methyldecanoyl-CoA 26 was synthesised (Scheme 5) from the acyl-Evan's auxiliary 28 (Scheme 2, vide supra) by hydrolysis with lithium hydroperoxide to give acid 39, which was converted to the CoA ester 26 using the standard procedure [6,10,11,32,33,40].

2.2. Evaluation of inhibitors


The selected AMACR inhibitors were evaluated using the colori-metric assay [32]. Incubation of active human AMACR 1A with sub-strate 1 results in production of 2,4-dinitrophenolate 2 and unsaturated product 3 (Scheme 1). Hence, enzyme activity can be determined based on measuring the absorbance of 2 at 354 nm over the assay time course. Inhibitory potency was assessed using dose–response curves to de-termine IC₅₀ values (Fig. 1).

As expected, ibuprofenoyl-CoA 5 and its derivatives 6–9 were in-hibitors of the enzyme, with most having IC₅₀ values of ca. 500 nM (Fig. 1). Variation of the structure of the side-chain in these inhibitors appeared to make little difference to inhibitory activity (as judged by IC₅₀ values), although fenoprofenoyl-CoA 6 appeared to be slightly more potent than the other examples and naproxenoyl-CoA 9 appeared to be slightly less so. All these compounds are known substrates of AMACR [6] and are predicted to behave as competitive inhibitors. Ibuprofenoyl-CoA 5 has been previously confirmed to be a competitive inhibitor of AMACR, with K_i = 60 nM [32], consistent with observa-tions of other workers on the human and rat enzymes [28,37,38]. The mandelic acid derivatives R- and S-2-hydroxy-2-phenylacetyl-CoA 10 and 11 were also modest inhibitors (Fig. 1), binding approximately ten times less strongly than compounds 5–9. Compounds 10 and 11 are not substrates of AMACR, since enzyme-catalysed α -proton exchange does not occur [11]. This result with 10 and 11 also demonstrates that in-hibitors can possess a 2-hydroxy- group in addition to the previously




Compound	IC ₅₀ (nM)	miLogP*	Previously tested as substrate?	Previously tested as Inhibitor?
4	~0.4	-2.61	No	Yes [29, 32]
5	540	-4.18	Yes [6, 29, 49]	Yes [28, 32, 37, 38]
6	400	-3.77	Yes [6, 11]	Not tested
7	590	-3.62	Yes [6]	Not tested
8	560	-4.07	Yes [6]	Not tested
9	750	-4.25	Yes [6]	Not tested
10	3.8 × 10 ³	-5.46	Not a substrate [11]	Not tested
11	2.3 × 10 ³	-5.46	Not a substrate [11]	Not tested
12	3.1 × 10 ³	-3.63	Yes [36]	Not tested
13	9.6 × 10 ³	-4.50	Yes [36]	Yes – No inhibition [37, 38]
14	1.6 × 10 ⁴	-4.98	Yes [36]	Not tested
15	>1.0 × 10 ⁵	-5.30	Poor substrate [36]	Yes – No inhibition [37, 38]
16	>1.0 × 10 ⁵	-5.38	Not a substrate [36]	Yes – No inhibition [37, 38]
17	>1.0 × 10 ⁵	-5.55	Not tested	Yes – No inhibition [37, 38]
18	930	-3.39	Yes [7]	Not tested
19	1170	-3.39	Yes [7, 36]	Not tested
20	200	-3.61	Yes [10]	Not tested†
21	300	-3.61	Not tested	Not tested†
22	180	-3.34	Not a substrate [10]	Yes – No inhibition [37, 38]
23	600	-3.56	Not tested	Not tested‡
24	4.1 × 10 ³	-4.02	Not tested	Not tested
25	360	-4.55	Not tested	Not tested¶
26	560	-4.43	Not tested	Not tested

Fig. 1. Structures for AMACR inhibitors, as measured by the colorimetric assay [32]. * Calculated miLogP values are for the acyl-CoA ester (<http://www.molinspiration.com/cgi-bin/properties>). †(2R,3R)- and (2S,3S)-3-Fluoro-2-methylhexadecanoyl-CoA were previously shown to be AMACR inhibitors [28]. ‡Several side-chain analogues of 23 reported as inhibitors [30]. ¶Binding of the 2-methylacetoacetyl-CoA enolate to MCR observed by X-ray crystallography [9]. (See above-mentioned references for further information.)



Scheme 2. Synthesis of (2S,3S)-3-fluoro-2-methyldecanoyl-CoA 21. Reagents and conditions: i. Bu₂BOTf, Pr₂NEt, octanal, CH₂Cl₂, -78 °C, 76%; ii. DAST, CH₂Cl₂, -78 °C, 35%; iii. LiOH, H₂O₂, H₂O/THF, 0 °C, 92%; iv. CDI, CH₂Cl₂, rt; v. CoA-SH Li⁺₃, 0.1 M NaHCO₃ aq/THF (1:1).



Scheme 3. Synthesis of 2-methylenedecanoyl-CoA 23. Reagents and conditions: i, DCC, DMAP, CH₂Cl₂; ii, NaBH₄, AcOH, 89% over two steps; iii, Me₂N⁺=CH₂ I⁻, MeOH, 74%; iv, NaOH, EtOH, 80%; v, NEt₃, EtOCOCl, THF; vi, aq. KHCO₃ (2.5% w/v), CoA-SH Li⁺₃, THF.

reported 2-trifluoromethyl- [28] and 2-chloro- [29] groups in place of the 2-methyl group. A wide range of aromatic inhibitor side-chains can therefore be accommodated by the enzyme, consistent with predictions made based on the MCR crystal structures [8] and biochemical data [6,11].


Acyl-CoA esters 12–17, possessing alkyl side-chains, were also assessed as inhibitors (Fig. 1). The potency of inhibition for acyl-CoA esters with side-chains of four carbons or fewer (15–17) is very weak, with low levels of inhibition (15–30%) observed even at very high in-hibitor concentrations (100 μM). Inhibition increased as alkyl chain-length increased. This behaviour is consistent with that of straight-chain acyl-CoA esters acting as substrates [36], where increased levels of α-proton exchange are observed with increasing chain length. Our results showing inhibition of AMACR by straight-chain acyl-CoA esters contrasts with the early observations of Schmitz et al., who reported that these compounds were not inhibitors of the native human and rat enzymes [37,38].

Inclusion of a 2-methyl group on the inhibitor increased potency by about 3-fold (compounds 18 and 19 compared to 12), again consistent with the finding that 2-methylacyl-CoA esters are much more efficient substrates than their straight-chain equivalents [36]. R-2-Methyldecanoyl-CoA 18 appeared to be a slightly more potent inhibitor than S-2-


methyldecanoyl-CoA 19, and this probably reflects the physiological role of AMACR in the conversion of R-2-methylacyl-CoAs to their S-2-methylacyl-CoA epimers [1,2].

2-Methyldecanoyl-CoA derivatives with more acidic α-protons are better inhibitors than their parent compounds (Fig. 1). R,R-3-Fluoro-2-methyldecanoyl-CoA 20 was about 5-fold more potent than was R-2-methyldecanoyl-CoA 18. A similar trend was observed with S,S-3-fluoro-2-methyldecanoyl-CoA 21 and S-2-methyldecanoyl-CoA 19, although both of these compounds were slightly less potent than their R-epimers. The product of the reaction, E-2-methyldec-2-enoyl-CoA 22, is also a potent inhibitor. It is therefore difficult to determine if the observed IC₅₀ values for 20 or 21 reflect the conversion of these substrates [10], product inhibition by 22 or both. These observations contrast with early studies [37,38], which suggest that 22 was not an inhibitor of AMACR.

2-Methyl-3-oxodecanoyl-CoA 25 was also a good inhibitor of AMACR. The α-proton of this compound is relatively acidic and 25 undergoes rapid non-enzymatic α-proton exchange with solvent via an enolate intermediate. It was therefore not possible to analyse the influence of 2-methyl group stereochemical configuration. The precursor 24 was a much poorer inhibitor than 25, presumably due to reduced acidity of the α-proton. It is also possible that the additional steric bulk



Scheme 4. Synthesis of 2-methyl-3-oxodecanoyl-CoA 25. Reagents and conditions: i, ethane-1,2-diol, TMSCl, CH₂Cl₂, 82%; ii, aq. NaOH/MeOH, 40%; iii, CDI, CH₂Cl₂; v, CoA-SH Li⁺₃, 0.1 M NaHCO₃ aq/THF (1:1); iv, aq. HCl, acetone.



Scheme 5. Synthesis of (2S,3R)-3-hydroxy-2-methyldecanoyl-CoA 26. Reagents and conditions: i. NaOH, H₂O₂, rt, quant.; ii. CDI, CH₂Cl₂, rt; iii. CoA-SH Li⁺, 0.1 M NaHCO₃ aq./THF (1:1).




Fig. 2. Incubation of 2S,3R-3-hydroxy-2-methyldecanoyl-CoA 26 with AMACR in buffer and ²H₂O. A. Heat-inactivated enzyme; B. live enzyme. Red circles highlight doublet for substrate 2-methyl group, showing no exchange of the α-proton had occurred (conversion to a single peak occurs on exchange to α-²H upon 'racemisation' [7,11]). Green circles denote expected position of 2-methyl singlet for the expected unsaturated product 22, showing that no elimination reaction has occurred.

at carbon-3 contributes to the lower potency of 24 compared to 25, as the 5-membered ring of 24 will be twisted out of plane relative to the aliphatic side-chain. However, AMACR is known to be able to accept substrates with diverse side-chain structures [6,28,36–38] and it is notable that ibuprofenoyl-CoA 5 and derivatives 6–9 (which have aromatic rings at the equivalent position) are potent inhibitors (Fig. 1).

2S,3R-3-Hydroxy-2-methyldecanoyl-CoA 26 is also a relatively potent inhibitor. Incubation of 26 with active AMACR did not result in an elimination reaction, as judged by the lack of a peak at δ 1.75 ppm [10] from the 2-methyl group of the anticipated product 22 (Fig. 2), probably because hydroxide is a relatively poor leaving group (water pK_a = ~14.0–15.7), compared to fluoride (HF pK_a = 3.2) [42]. Similarly, ¹H NMR analysis of the reaction products showed that 26 did not undergo α-proton exchange, and hence 26 is not a substrate of AMACR. It is notable that 22, 24 and 26 are intermediates in the branched-chain acyl-CoA β-oxidation pathway. One may speculate that these compounds could provide some regulation of AMACR by negative feedback control, and hence control entry of R-2-methylacyl-CoA esters into the β-oxidation pathway. Whether or not this is physiologically significant will depend on rates of flux through the β-oxidation pathway and whether intermediates 22, 24 and 26 are sequestered away from AMACR.

Acyl-CoAs which mimic the planar enolate intermediate are good inhibitors of AMACR (Fig. 1). E-2-Methyldec-2-enoyl-CoA 22 and 2-methylenedecanoyl-CoA 23 bind strongly due to having a planar sp²-hybridized α-carbon. This result is consistent with a previous study in which 2-methyleneacyl-CoAs were shown to be good competitive

inhibitors [2,30]. The enolate analogue N-dodecyl-N-methylcarbamoyl-CoA 4 is the best rationally designed inhibitor (IC₅₀ = 0.4 nM) reported to date [29,32]. Inhibition by 4 is ca. 2000× more potent than by 18 and 19 (Fig. 1). This high potency of 4 appears to largely result from the mimicking of the enolate intermediate by the carbamoyl moiety. It is also notable that the determined IC₅₀ value for 4 in this study (0.4 nM) is significantly lower than that previously determined by Carnell et al. (98 nM), when assayed against HEK-derived human AMACR using ibuprofenoyl-CoA 5 as substrate [29]. It appears that the colorimetric assay consistently determines higher levels of compound potency than other assays (e.g. K_i = 60 nM [32] vs. 56 μM [28,38] for an ibuprofenoyl-CoA epimeric mixture). The reasons for this discrepancy is not entirely clear, but higher apparent potency may be a consequence of using a substrate undergoing an irreversible reaction to measure activity (and hence avoiding the error introduced by the presence of the reverse reaction). Alternatively, this may be related to the extent of substrate or inhibitor micelle formation under the different assay conditions.

The determined IC₅₀ value of 4 (0.4 nM) is around half the calculated 'active' enzyme concentration in the assay [32] (based on comparison of k_{cat}/K_m values for the E. coli and HEK cell derived enzymes [29]). Hence 4 could be behaving as a tight-binding inhibitor. Compound 4 behaves as a rapidly reversible competitive inhibitor of AMACR, with a Hill coefficient of ~0.7 [32]. These observations are consistent with the zone A inhibitor behaviour described by Straus and Goldstein [43,44], i.e. the enzyme active site concentration is < 0.1 × the apparent K_i value (0.65 nM [32]). This rapidly reversible inhibition is significantly different behaviour to that observed for similar compounds (gem- carbamoyl inhibitors and N-decyl-carbamoyl-CoA) with the highly homologous bacterial enzyme MCR, where time-dependent inactivation was observed [35]. The reasons for this difference in behaviour are not entirely obvious.

2.3. Influence of the side-chain lipophilicity on inhibitor potency

AMACR is able to catalyse the 'racemisation' of substrates with structurally diverse side-chains [1,2]. The accommodation of these diverse structures is thought to be a result of non-specific binding of the side-chain by hydrophobic interactions to a methionine-rich surface [8]. Consistent with this, the MCR gem- [31] and gem- carbamate [35] inhibitors show increased potency for compounds with more hydrophobic alkyl side-chains. Consequently, we were interested to investigate whether inhibitor potency was related to the lipophilicity of the inhibitor side-chain. A plot of determined IC₅₀ values vs. calculated LogP values for the acyl-CoA inhibitor (Fig. 3) showed that high LogP values tend to produce low IC₅₀ values. Systematic trends were observed within those compounds containing aromatic side-chains (5–11), with potency positively correlating with lipophilicity. It is also notable that 10 and 11, possessing a single phenyl group side-chain are significantly less potent than compounds containing more lipophilic side-chains. This consistent behaviour tends to suggest that side-chain lipophilicity is driving potency, with the 2-hydroxy group of 10 and 11 making a smaller contribution. Systematic trends were also observed for inhibitors possessing alkyl side-chains (12–17), showing that lipophilicity is also an important determinant of potency for this series.

The enolate analogue 4 has a potency increased by ~875-fold compared to that predicted based on LogP values (measured IC₅₀ = ~0.4 nM vs. ~350 nM predicted for mLogP = 2.61) (Supplementary Information, Fig. S1), showing the effectiveness of the




Fig. 3. Correlation of inhibitor potency with lipophilicity (miLogP value; <http://www.molinspiration.com/cgi-bin/properties>). Compound numbers refer to structures shown in Fig. 1. Compounds with green numbers are those with aromatic side-chains (5–11); Compounds with blue numbers are straight-chain acyl-CoA esters and isobutanoyl-CoA (12–17); Compounds with purple numbers are 2-methyldecanoyl-CoA and 3-fluoro-2-methyldecanoyl-CoAs (18–21); Compounds with orange numbers are intermediates in the β -oxidation pathway which occurs subsequent to AMACR activity (22, 25 and 26); Compounds with red numbers are inhibitors or analogues of known inhibitors (4 and 23); Compound 24 is a synthetic intermediate to 25.

carbamate moiety in promoting inhibition. Acyl-CoA esters do not comply with Lipinski guidelines and hence AMACR inhibitors are delivered as their acid pro-drugs [28,30,32,34] which are converted to the acyl-CoA in vivo. Although 4 has very high potency, delivery as the pro-drug will be challenging because carbamates readily decarboxylate to the corresponding amine.

3. Conclusions

This is the first systematic SAR study of rationally designed AMACR inhibitors. The study illustrates that extremely diverse side-chain structures which can be accommodated. A minimal level of side-chain lipophilicity is required for efficient binding. For compounds with aromatic side-chains, a single aromatic ring results in modest inhibition whilst more than one aromatic ring or an aromatic ring with alkyl substituents results in much more potent inhibition. Similarly, a minimum of a 6-carbon alkyl chain appears to be required for reasonably efficient inhibitor binding, with increased potency resulting from addition of further $-\text{CH}_2-$ groups. Our results allow investigation of the contribution to potency made by the individual structural elements of these inhibitors.

AMACR has attracted much attention as both a novel drug target and cancer marker since its involvement in prostate cancer was reported [14,19]. However, exploitation of this discovery has been extremely limited, largely due to the absence of a suitable assay with which to test inhibitor potency [33]. This study shows that our novel colorimetric assay [32] allows quick and accurate measurement of drug potency and detailed kinetic characterisation of inhibitors. The systematic investigation of novel inhibitor SAR and therapeutic development is now possible.

4. Materials and methods

4.1. Sources of materials

Chemicals were purchased from the Sigma-Aldrich Chemical Co. or Fisher Scientific Ltd., unless otherwise stated, and were used without further purification. Reduced coenzyme A, tri-lithium salt, was

purchased from Calbiochem. Acyl-CoA esters 4 [33], 5–9 [6], 10 and 11 [11] were synthesised as previously described. Acyl-CoA esters 12–17 were purchased from Larodan Lipids. Substrates 18–20 and product 22 were synthesised as described [7,10]. Human recombinant AMACR 1A was expressed and purified and substrate 1 synthesised as previously described [32].

4.2. General experimental procedures

Thin layer chromatography was performed on Merck silica aluminium plates 60 (F254) and UV light, potassium permanganate or phosphomolybdic acid were used for visualisation. Column chromatography was performed using Fisher silica gel (particle size 35–70 μm). Purifications of acyl-CoA esters were performed by solid phase extraction using Oasis HLB 6cc (200 mg) extraction cartridges. Phosphate buffer was prepared from monobasic sodium phosphate and NaOH at the required proportions. Citric acid buffer was prepared from citric acid and NaOH at the required proportion for 0.8 M pH 4.0 buffer. The pH of aqueous solutions was measured using a Corning 240 pH meter and Corning general purpose combination electrode. The pH meter was calibrated using Fisher Chemicals standard buffer solutions (pH 4.0 – phthalate, 7.0 – phosphate, and 10.0 – borate) at either pH 7.0 and 10.0 or 7.0 and 4.0. Calibration and measurements were carried out at ambient room temperature. IR spectra were recorded on Perkin-Elmer RXI FTIR spectrometer instrument. NMR spectra were recorded on Bruker Avance III 400.04 MHz or 500.13 MHz spectrometers in D_2O , $(\text{CD}_3)_2\text{SO}$ or CDCl_3 and the solvent was used as an internal standard. Shifts are given in ppm and J values reported to ± 0.1 Hz. Multiplicities of NMR signals are described as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet. Stock concentrations of acyl-CoA esters for assays were determined using ^1H NMR. Mass spectra were recorded by ESI TOF. High resolution mass spectra were recorded in ES mode. Melting points were determined using a Gallenkamp melting point apparatus and are uncorrected. Syntheses were carried out at ambient temperature, unless otherwise specified. Solutions in organic solvents were dried over anhydrous MgSO_4 and the solvents were evaporated under reduced pressure. Aqueous solutions for biological experiments were prepared in Nanopure water of $18.2 \text{ M}\Omega\cdot\text{cm}^{-1}$ quality and were pH-adjusted with aq. HCl or NaOH.

4.3. (4S)-4-Benzyl-3-[(2S,3R)-3-hydroxy-2-methyldecanoyl]-1,3-oxazolidin-2-one (28)

(S)-(+)-4-Benzyl-3-propanoyl-2-oxazolidinone 27 (3.0 g, 12.9 mmol) in CH_2Cl_2 (30.0 mL) was cooled to -78°C . Dibutylboron triflate (1.0 M in CH_2Cl_2 , 13.0 mL, 12.9 mmol) and Pr^i_2NEt (2.3 mL, 12.9 mmol) were added and the mixture was stirred for 30 min before octanal (1.4 mL, 9.2 mmol) in CH_2Cl_2 (9.0 mL) was added dropwise. The mixture was stirred at -78°C for further 30 min and then at room temperature for 2 h. aq. Sodium phosphate buffer at pH 7.0 (100 mM, 100 mL) was added slowly to the reaction mixture. The organic layer was washed [aq. HCl (1.0 M), aq. NaHCO_3 (saturated), brine] and dried. Column chromatography (Petroleum ether/EtOAc 10:1 \rightarrow 6:1) gave 28 (2.53 g, 76%) as a colourless oil. $[\alpha]_D^{21} +51.4$ (c 0.74 in CHCl_3); IR ν_{max} 3517 (OH), 1780 (C=O), 1692 (C=O) cm^{-1} ; ^1H NMR (500.13 MHz; CDCl_3) δ 7.26–7.07 (5H, m, Ar-H), 4.65–4.55 (1H, m, 4-H), 4.16–4.05 (2H, m, 5-H), 3.89–3.80 (1H, m, 3'-H), 3.68 (1H, qd, $J = 7.0, 3.0$ Hz, 2'-H), 3.14 (1H, dd, $J = 13.0, 3.0$ Hz, CHHAr), 2.92 (1H, s, OH), 2.70 (1H, dd, $J = 13.0, 9.0$ Hz, CHHAr), 1.50–1.10 (15H, m, $6 \times \text{CH}_2$ and CH_3CH), 0.79 (3H, t, $J = 6.5$ Hz, 10'-H₃); ^{13}C NMR (125.77 MHz, CDCl_3) δ 177.23 (1'-C), 152.87 (2-C), 134.92 (Ar-C), 129.24 (Ar-C), 128.73 (Ar-C), 127.18 (Ar-C), 71.35 (3'-C), 65.96 (5-C), 54.91 (4-C), 42.02 (2'-C), 37.53 (CHHAr), 33.77 (CH_2), 31.62 (CH_2), 29.35 (CH_2), 29.05 (CH_2), 25.84 (CH_2), 22.45 (CH_2), 13.90 (CH_3CH), 10.31 (10'-C); ESI-MS m/z 384.2134 $[\text{M}+\text{Na}]^+$ ($\text{C}_{21}\text{H}_{31}\text{NNaO}_4$ requires 384.2151).

4.4. (4S)-4-Benzyl-3-[(2R,3S)-3-fluoro-2-methyldecanoyl]-1,3-oxazolidin-2-one (29)

(Diethylamino)sulfur trifluoride (0.5 mL, 3.9 mmol) in CH_2Cl_2 (10.0 mL) was added dropwise to **28** (1.4 g, 3.9 mmol) in CH_2Cl_2 (20.0 mL) at -78°C and the mixture was stirred for 2 h at this temperature. It was stirred for a further 2 h at room temperature, before being quenched with water (50 mL). The organic layer was washed (saturated aq. NaHCO_3 , brine). Column chromatography (petroleum ether/EtOAc 30:1) gave **29** (490 mg, 35%) as a colourless oil. $[\alpha]_D^{21} +49.2$ (c 0.63 in CHCl_3); IR ν_{max} 1782 (C=O), 1700 (C=O) cm^{-1} ; ^1H NMR (500.13 MHz, CDCl_3) δ_{H} 7.27–7.06 (5H, m, Ar-H), 4.76–4.57 (2H, m, 4-H and 3'-H), 4.15–3.96 (3 H, m, 5-H and 2'-H), 3.17 (1 H, dd, $J = 13.5, 3.5$ Hz, CHHAr), 2.72 (1 H, dd, $J = 13.5, 9.5$ Hz, CHHAr), 1.70–1.14 (12H, m, $6 \times \text{CH}_2$), 1.10 (3 H, d, $J = 7.0$ Hz, CH_3CH), 0.80 (3 H, t, $J = 6.5$ Hz, $10'\text{-H}_3$); ^{13}C NMR (125.77 MHz, CDCl_3) δ_{C} 174.37 (d, $J = 2.8$ Hz, $1'\text{-C}$), 153.12 (2-C), 135.25 (Ar-C), 129.47 (Ar-C), 128.93 (Ar-C), 127.37 (Ar-C), 94.89 (d, $J = 169.8$ Hz, $3'\text{-C}$), 66.19 (5-C), 55.38 (4-C), 42.03 (d, $J = 20.9$ Hz, $2'\text{-C}$), 37.85 (CHHAr), 32.04 (d,

$J = 20.9$ Hz, $4'\text{-CH}_2$), 31.79 (CH_2), 29.37 (CH_2), 29.16 (CH_2), 24.57 (d, $J = 2.8$ Hz, $5'\text{-CH}_2$), 22.64 (CH_2), 14.10 ($10'\text{-C}$) and 13.62 (d, $J = 8.5$ Hz, $2'\text{-CH}_3$); δ_{F} (470 MHz, CDCl_3) $-\text{179.67}$; ESI-MS m/z 386.2126 $[\text{M}+\text{Na}]^+$ ($\text{C}_{21}\text{H}_{30}\text{FNNaO}_3$ requires 386.2107).

4.5. (2R,3S)-3-Fluoro-2-methyldecanoic acid (30)

aq. H_2O_2 [30% (w/w), 0.7 mL] and LiOH (62 mg, 2.6 mmol) were added to **29** (472 mg, 1.3 mmol) in THF (14 mL) at 0°C . The mixture was stirred at ambient temperature for 20 h before being quenched with sat. aq. sodium sulfite (14 mL). The THF was evaporated and the mixture was acidified with aq. HCl (1.0 M) to pH 1.0 and extracted with CH_2Cl_2 (100 mL). The organic phase was washed with water (75 mL) and brine (75 mL). The combined organic layers were dried and the solvent was evaporated. Column chromatography (petroleum ether/EtOAc 5:1) gave **30** (245 mg, 92%) as a white solid. mp $64\text{--}65^\circ\text{C}$; $[\alpha]_D^{21} -7.8$ (c 0.51 in CHCl_3); IR ν_{max} 2925 (OH), 1693 (C=O) cm^{-1} ; ^1H NMR (400.04 MHz, CDCl_3) δ_{H} 10.89 (1H, br s, OH), 4.58–4.50 (1H, m, 3-H), 2.58–2.52 (1H, m, 2-H), 1.75–1.23 (12H, m, $6 \times \text{CH}_2$), 1.20 (3H, d, $J = 7.2$ Hz, CH_3CH), 0.88 (3H, t, $J = 6.8$ Hz, 10-H_3); ^{13}C NMR (100.60 MHz, CDCl_3) δ_{C} 180.02 (d, $J = 5.6$ Hz, C=O), 94.28 (d, $J = 172.2$ Hz, 3-C), 44.37 (d, $J = 22.0$ Hz, 2-C), 31.74 (d, $J = 21.1$ Hz, 4- CH_2), 31.74 (CH_2), 29.30 (CH_2), 29.12 (CH_2), 24.82 (d, $J = 3.0$ Hz, 5- CH_2), 22.61 (CH_2), 14.05 (10-C) and 12.56 (d, $J = 6.6$ Hz, 2- CH_3); ^{19}F NMR (470 MHz, CDCl_3) δ_{F} $-\text{181.94}$; ESI-MS m/z 203.1449 $[\text{M}-\text{H}]^-$ ($\text{C}_{11}\text{H}_{20}\text{FO}_2$ requires 203.1447).

4.6. (2S,3S)-3-Fluoro-2-methyldecanoyl-CoA (21)

N,N'-Carbonyldiimidazole (48 mg, 0.29 mmol) was added to (2R,3S)-3-fluoro-2-methyldecanoic acid **30** (30.0 mg, 0.15 mmol) in anhydrous CH_2Cl_2 (2 mL) and the mixture was stirred at ambient temperature for 1 h. CH_2Cl_2 (3 mL) was added to the mixture, which was washed with water (5×2 mL) and brine (2 mL) and dried. The solvent was evaporated to obtain the crude acyl-imidazole intermediate. Aq. NaHCO_3 (1.0 mL, 0.10 M) and tri-lithium CoA-SH (17.0 mg, 0.02 mmol) was added to the crude intermediate in THF (1.0 mL) and the mixture was stirred at ambient temperature for 18 h. The THF was evaporated and the residue was acidified to ca. pH 3 by addition of aq. HCl (1.0 M). The mixture was diluted with water (2.0 mL) and washed with EtOAc (3×3 mL). The crude aqueous solution was freeze-dried and purified with solid-phase extraction to give

21 (13.6 mg) as a white solid. ^1H NMR (500 MHz, D_2O) δ_{H} 8.63 (1H, s, adenosine CH), 8.38 (1H, s, adenosine CH), 6.16 (1H, d, $J = 6.0$ Hz, adenosine CH), 4.30–4.10 (2H, m, adenosine CH_2), 3.97 (1H, s, adenosine CH), 3.85–3.72 (1H, m, CoA(OCHH)), 3.58–3.47 (1H, m, CoA(OCHH)), 3.39 (2H, t, $J = 6.5$ Hz, CoA(CH_2)), 3.29 (2H, t,

$J = 6.0$ Hz, CoA(CH_2)), 3.08–2.90 (3H, m, CoA(SCH_2) and CHCH_3), 2.36 (2H, t, $J = 6.5$ Hz, CoA(CH_2)), 1.68–1.44 (2H, m, CHH and CHH), 1.39–1.12 (10H, m, $5 \times \text{CH}_2$), 1.07 (3H, d, $J = 7.0$ Hz, CHCH_3), 0.87 (3H, s, CoA(CH_3)), 0.81–0.70 (6H, m, CH_2CH_3 and CoA(CH_3)); ^{19}F NMR (470 MHz) δ_{F} $-\text{181.11}$; ESI-MS m/z 475.6220 $[\text{M}-2\text{H}]^{2-}$ ($\text{C}_{32}\text{H}_{53}\text{FN}_7\text{O}_{17}\text{P}_3\text{S}$ requires 475.6208).

4.7. 2,2-Dimethyl-5-octyl-1,3-dioxane-4,6-dione (31)

4-Dimethylaminopyridine (1.296 g, 10.6 mmol), N,N'-dicyclohexylcarbodiimide (1.0 M in CH_2Cl_2 , 11.1 mL, 11.1 mmol) and octanoic acid **32** (1.6 mL, 10 mmol, 1.0 eq.) were added to Meldrum's acid **33** (1.455 g, 10.1 mmol) in dry CH_2Cl_2 (100 mL). The mixture was stirred at ambient temperature for 40 h. The precipitate was removed by filtration and the filtrate was washed with aq. KHSO_4 (1.0 M, twice), water and brine, then dried. AcOH (6.0 mL) was added to the filtrate. To this solution, NaBH_4 (802 mg, 21.2 mmol) was added in portions during 1 h and the mixture was stirred for an additional 20 h. The evaporation residue was dissolved in Et_2O (100 mL) and washed with water (twice) and brine (100 mL). Drying and evaporation gave **31** (2.30 g, 89%) as a white solid: mp. $64\text{--}65^\circ\text{C}$ (lit. [45] mp. $65\text{--}67^\circ\text{C}$); ^1H NMR (400.04 MHz, CDCl_3) δ_{H} 3.49 (1H, t, $J = 4.9$ Hz, dioxane 5-H), 2.14–2.04 (m, 2H, octyl 1- H_2), 1.77 (3H, s, 2- CH_3), 1.75 (3H, s, 2- CH_3), 1.48–1.38 (2H, m, octyl 3- H_2), 1.37–1.19 (10H, m, octyl 4,5,6,7- H_8), 0.86 (3H, t, $J = 7.0$ Hz, octyl 8- H_3); ^{13}C NMR (125.77 MHz, CDCl_3) δ_{C} 165.6, 104.7, 46.1, 31.8, 29.5, 29.2, 29.1, 28.4, 26.9, 26.6, 26.5, 22.6, 14.1; IR (KBr disc) ν_{max} 1752 (C=O) cm^{-1} ; ESI-MS m/z 279.1577 $[\text{M}+\text{Na}]^+$ ($\text{C}_{14}\text{H}_{24}\text{NaO}_4$ requires 279.1572; 257.1732 $[\text{M}+\text{H}]^+$ ($\text{C}_{14}\text{H}_{25}\text{O}_4$ requires 257.1753).

4.8. Methyl 2-methylenedecanoate (34)

Compound **31** (2.10 g, 8.19 mmol) was dissolved in anhydrous MeOH (26 mL). Eschenmoser's salt (3.79 g, 20.5 mmol, 2.5 eq.) was added and the mixture was heated at reflux for 40 h. The solvent was evaporated. The residue, in Et_2O (100 mL), was washed with aq. KHSO_4 (1.0 M), water and brine and was dried. Evaporation and column chromatography (petroleum ether/EtOAc 10:1) gave **34** (1.10 g, 74%) as a colourless oil (lit. [46] oil): IR (neat) ν_{max} 1725 (C=O) cm^{-1} ; ^1H NMR (400.04 MHz, CDCl_3) δ_{H} 6.12–6.10 (1H, m, JCHH), 5.52–5.49 (1H, m, JCHH), 3.73 (3H, s, OCH_3), 2.31–2.23 (2H, m, 3- H_2), 1.49–1.37 (2H, m, 4- H_2), 1.34–1.19 (10H, m, 5,6,7,8,9- H_{10}), 0.86 (3H, t, $J = 7.1$ Hz, 10-H_3); ^{13}C NMR (125.77 MHz, CDCl_3) δ_{C} 167.8, 140.8, 124.4, 51.7 (two carbons), 31.8, 29.3, 29.2, 29.2, 28.3, 22.6, 14.0.

4.9. 2-Methylenedecanoic acid (35)

Methyl ester **34** (870 mg, 4.39 mmol) was stirred at 50°C for 2 h with aq. NaOH (1.0 M, 15 mL) in EtOH (57 mL), then cooled to ambient temperature and acidified to pH ca. 3. The volatile solvents were evaporated. The residue, in Et_2O (50 mL), was washed with water (twice) and brine and was dried. Evaporation and column chromatography (petroleum ether/EtOAc 3:1) gave **35** (650 mg, 80%) as a colourless oil (lit. [47] oil): IR (neat) ν_{max} 1696 (C=O) cm^{-1} ; ^1H NMR (400.04 MHz, CDCl_3) δ_{H} 12.05 (1H, br s), 6.34–6.23 (1H, m, JCHH), 5.70–5.59 (1H, m, =CHH), 2.32–2.25 (2H, m, 3- H_2), 1.54–1.40 (2H, m, 4- H_2), 1.36–1.19 (10H, m, 5,6,7,8,9- H_{10}), 0.87 (3H, t, $J = 7.0$ Hz, 10-H_3); ^{13}C NMR (125.77 MHz, CDCl_3) δ_{C} 173.0, 140.2, 126.9, 31.8, 31.4, 29.3, 29.2, 29.2, 28.3, 22.6, 14.1; ESI-MS m/z 207.1349 $[\text{M}+\text{Na}]^+$ ($\text{C}_{11}\text{H}_{20}\text{NaO}_2$ requires 207.1361; 183.1396 $[\text{M}-\text{H}]^-$ ($\text{C}_{11}\text{H}_{19}\text{O}_2$ requires 183.1385).

4.10. 2-Methylenedecanoyl-CoA (23)

Ethyl chloroformate (17 μL , 19 mg, 0.18 mmol) was added to **35** (33 mg, 0.18 mmol) and NEt_3 (25 μL , 18 mg, 0.18 mmol) in anhydrous

THF (2.0 mL) and the mixture was stirred at ambient temperature for 1 h. CoA-SH tri-lithium salt [(28 mg, 0.04 mmol) in aq. KHCO_3 (2.5%), 2.0 mL] was added and the mixture was stirred at ambient temperature for 16 h. The mixture was acidified to pH ca. 3 with aq. HCl (1.0 M) and the THF was evaporated. The solution was washed with EtOAc (5×3 mL) and the crude product was purified by SPE to give **23** (7.0 mg) as a colourless solid: ^1H NMR (500.13 MHz, D_2O): δ_{H} 8.47 (1H, s), 8.16 (1H, s), 6.07 (1H, d, $J = 7.0$ Hz), 5.96 (1H, s), 5.55 (1H, s), 4.19–4.11 (1H, m), 3.75 (1H, dd, $J = 9.8$, 5.1 Hz), 3.46 (1H, dd, $J = 9.8$, 4.8 Hz), 3.36 (2H, t, $J = 6.5$ Hz), 3.31–3.25 (2H, m), 2.95 (2H, t, $J = 6.2$ Hz), 2.33 (2H, t, $J = 6.6$ Hz), 2.16 (2H, t, $J = 7.3$ Hz), 1.30–1.22 (2H, m), 1.18–1.06 (10H, m), 0.79 (3H, s), 0.75 (3H, t, $J = 6.8$ Hz), 0.65 (3H, s); ESI-MS m/z 465.6152 $[\text{M}-2\text{H}]^{2-}$ ($\text{C}_{32}\text{H}_{52}\text{N}_7\text{O}_{17}\text{P}_3\text{S}$ requires 465.6177).

4.11. Methyl 2-(2-heptyl-1,3-dioxolan-2-yl)propanoate (**36**)

Ethane-1,2-diol (5.27 g, 84.8 mmol) was added to **37** (606 mg, 2.83 mmol) in dry CH_2Cl_2 (26 mL). Me_3SiCl (1.84 g, 2.15 mL, 17.0 mmol) was added and the mixture was stirred at ambient temperature for 3 d. Further ethane-1,2-diol (5.27 g) and Me_3SiCl (1.84 g) were added and the mixture was stirred for a further 3 d. Water (25 mL) was added and the mixture was extracted with CH_2Cl_2 (3×20 mL). The combined organic layers were washed with water (5×70 mL) and brine (70 mL) and were dried. Evaporation and column chromatography (petroleum ether/ EtOAc 10:1) gave **36** (600 mg, 82%) as a colourless oil: IR (neat) n_{max} 1740 $[\text{C}=\text{O}] \text{ cm}^{-1}$; ^1H NMR (400.04 MHz, CDCl_3) δ_{H} 4.04–3.90 (4H, m, $\text{OCH}_2\text{CH}_2\text{O}$), 3.67 (3H, s, OCH_3), 2.83 (1H, q, $J = 7.2$ Hz, CHCH_3), 1.81–1.62 (2H, m, heptyl 1-H₂), 1.42–1.20 (10H, m, heptyl 2,3,4,5,6-H₁₀), 1.18 (3H, d, $J = 7.2$ Hz, propanoate 3-H₃), 0.86 (3H, t, $J = 7.1$ Hz, heptyl 7-H₃); ^{13}C NMR (125.77 MHz, CDCl_3): δ_{C} 173.9, 111.3, 65.5, 65.4, 51.7, 46.7, 35.0, 31.7, 29.7, 29.2, 22.8, 22.6, 14.0, 12.5; ESI-MS m/z 281.1761 $[\text{M}+\text{Na}]^+$ ($\text{C}_{14}\text{H}_{26}\text{NaO}_4$ requires 281.1729), 259.1883 $[\text{M}+\text{H}]^+$ ($\text{C}_{14}\text{H}_{27}\text{O}_4$ requires 259.1909).

4.12. 2-(2-Heptyl-1,3-dioxolan-2-yl)propanoic acid (**38**)

Aq. NaOH (1.0 M, 6.7 mL, 6.7 mmol) was stirred with **36** (345 mg, 1.34 mmol) in MeOH (30 mL) at ambient temperature for 2 h, then at 65°C for 2 h. The mixture was cooled to ambient temperature and citric acid buffer (0.8 M, pH 4.0, 15 mL) was added. The mixture was extracted with CH_2Cl_2 (3×30 mL). The combined organic layers were washed with water and brine and dried. Evaporation and column chromatography (petroleum ether/ EtOAc 2:1) gave **38** (130 mg, 40%) as a colourless oil. IR (neat) n_{max} 1709 $[\text{C}=\text{O}] \text{ cm}^{-1}$; ^1H NMR (400.04 MHz, CDCl_3): δ_{H} 10.78 (1H, br s, OH), 4.10–3.94 (4H, m, $\text{OCH}_2\text{CH}_2\text{O}$), 2.83 (1H, q, $J = 7.2$ Hz, CHCH_3), 1.80–1.67 (2H, m, heptyl 1-H₂), 1.43–1.20 (10H, m, heptyl 2,3,4,5,6-H₁₀), 1.23 (3H, d, $J = 7.2$ Hz, CHCH_3), 0.87 (3H, t, $J = 7.1$ Hz, heptyl 7-H₃); ^{13}C NMR (125.77 MHz, CDCl_3): δ_{C} 176.9, 111.4, 65.4, 65.4, 46.6, 34.7, 31.7, 29.6, 29.2, 22.8, 22.6, 14.0, 12.2; ESI-MS m/z 267.1549 $[\text{M}+\text{Na}]^+$ ($\text{C}_{13}\text{H}_{24}\text{NaO}_4$ requires 267.1572), 245.1729 $[\text{M}+\text{H}]^+$ ($\text{C}_{13}\text{H}_{25}\text{O}_4$ requires 245.1753).

4.13. 2-(2-Heptyl-1,3-dioxolan-2-yl)propanoyl-CoA (**24**)

Using the same method as for **21**, **24** was prepared from **38** (31 mg, 0.13 mmol, 1.0 eq.) by sequential treatment with $\text{N,N}'$ -carbonyldiimidazole (41 mg, 0.25 mmol, 2.0 eq.) and CoA-SH tri-lithium salt (29 mg, 0.04 mmol, 0.3 eq.) to give **24** (10 mg) as a colourless solid: ^1H NMR (500.13 MHz, D_2O) δ_{H} 8.57 (1H, s), 8.30 (1H, s), 6.13 (1H, d, $J = 6.1$ Hz), 4.21–4.14 (1H, m), 3.99–3.88 (4H, m), 3.78 (1H, dd, $J = 9.6$, 4.5 Hz), 3.50 (1H, dd, $J = 9.7$, 4.5 Hz), 3.39 (2H, t, $J = 6.6$ Hz), 3.32–3.22 (2H, m), 3.12 (1H, 2 \times q, $J = 7.0$ Hz; both epimers), 2.93 (2H, t, $J = 6.2$ Hz), 2.36 (2H, t, $J = 6.6$ Hz), 1.66–1.53 (2H, m), 1.26–1.11 (8H, m), 1.08 (3H, d, $J = 7.0$ Hz, CHCH_3), 0.85 (3H,

s), 0.76 (3H, t, $J = 7.1$ Hz, CH_2CH_3), 0.72 (3H, s); ESI-MS m/z 506.6219 $[\text{M}+\text{Na}]^{2+}$ ($\text{C}_{34}\text{H}_{55}\text{N}_7\text{NaO}_{19}\text{P}_3\text{S}$ requires 506.6192), 495.6301 $[\text{M}+\text{H}]^{2+}$ ($\text{C}_{34}\text{H}_{56}\text{N}_7\text{O}_{19}\text{P}_3\text{S}$ requires 495.6282).

4.14. 2R,S-2-Methyl-3-oxodecanoyl-CoA (**25**)

Compound **24** (7 mg, 7.1 μmol) was dissolved in water (0.8 mL) and acetone (1.0 mL). Aq. HCl (1.0 M, 0.2 mL) was added and the reaction mixture was stirred overnight. The acetone was evaporated and the residue freeze-dried to give **25** (6 mg) as a colourless solid: ^1H NMR (500.13 MHz, D_2O): δ_{H} 8.49 (1H, s), 8.33 (1H, s), 6.10 (1H, d, $J = 5.8$ Hz), 4.20–4.10 (2H, m), 3.99 (1H, q, $J = 7.0$ Hz), 3.76 (1H, dd, $J = 9.7$, 5.0 Hz), 3.38–3.28 (2H, m), 3.22 (2H, t, $J = 6.3$ Hz), 2.97–2.88 (2H, m), 2.57–2.44 (2H, m), 2.30 (2H, t, $J = 6.6$ Hz), 1.41–1.32 (2H, m), 1.16 (3H, d, $J = 7.0$ Hz), 1.12–1.04 (8H, m), 0.80 (3H, s), 0.69 (3H, s), 0.68 (3H, t, $J = 7.0$ Hz); ESI-MS m/z 473.6174 $[\text{M}-2\text{H}]^{2-}$ ($\text{C}_{32}\text{H}_{52}\text{N}_7\text{O}_{18}\text{P}_3\text{S}$ requires 473.6151).

4.15. 2S,3R-3-Hydroxy-2-methyldecanoyl-CoA (**26**)

2S,3R-3-Hydroxy-2-methyldecanoic acid **39** was synthesised [48] from the Evan's auxiliary-protected acid **28** by hydrolysis with LiOH and H_2O_2 . Following the procedure used for **21**, 2S,3R-3-hydroxy-2-methyldecanoyl-CoA **26** was prepared from 2S,3R-3-hydroxy-2-methyldecanoic acid **39** (40 mg, 0.20 mmol, 1.0 eq.), CDI (64 mg, 0.40 mmol, 2.0 eq.) and CoA-SH tri-lithium salt (78 mg, 0.10 mmol, 0.5 eq.) to give **26** (12 mg) as a colourless solid: ^1H NMR (500.13 MHz, D_2O) δ_{H} 8.48 (1H, s), 8.20 (1H, s), 6.09 (1H, d, $J = 6.5$ Hz), 4.22–4.10 (1H, m), 3.81–3.70 (1H, m), 3.48 (1H, dd, $J = 9.8$, 4.8 Hz), 3.37 (2H, t, $J = 6.6$ Hz), 3.25 (2H, t, $J = 6.3$ Hz), 2.97–2.87 (2H, m), 2.74–2.68 (1H, m), 2.34 (2H, t, $J = 6.5$ Hz), 1.39–1.29 (2H, m), 1.21–1.09 (10H, m), 1.06 (3H, d, $J = 6.9$ Hz, CHCH_3), 0.81 (3H, s), 0.74 (3H, t, $J = 7.1$ Hz, CH_2CH_3), 0.68 (3H, s); ESI-MS m/z $[\text{M}-2\text{H}]^{2-}$ 474.6217 ($\text{C}_{32}\text{H}_{54}\text{N}_7\text{O}_{18}\text{P}_3\text{S}$ requires 474.6229).

4.16. Evaluation of inhibition of AMACR by test compounds

Colorimetric assays were performed as previously described [32]. Dose response curves were used to determine IC_{50} values for inhibitors. Enzyme ($4 \times$ stock, 150 μL) and inhibitor at the appropriate concentration ($4 \times$ stock, 150 μL) were incubated together in 96 well plates at ambient room temperature for 10 min. The sample was divided into three repeats of 100 μL before addition of substrate ($2 \times$ stock, $3 \times 100 \mu\text{L}$; final concentration of 40 μM in the assay) and monitored at 354 nm. Each 200 μL assay contained ca. 8 μg of total AMACR protein (0.85 μM , assuming a molecular weight of 47,146.8 Da. with one active site per monomer [7]). Final concentrations of inhibitor in the assay were 100, 33.3, 11.1, 3.7, 1.23, 0.411, 0.137 and 0.045 μM unless otherwise stated. Positive controls contained enzyme and substrate 1 only and negative controls buffer and substrate. Rates in $\text{Absorbance} \cdot \text{min}^{-1}$ were determined using Excel and converted to $\text{nmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ using the 2,4-dinitrophenoxide 2 extinction coefficient ($15,300 \text{ M}^{-1} \text{ cm}^{-1}$) [32] with the path-length (0.588 cm) determined by the plate-reader. IC_{50} values were determined using reaction rate, with the data fitted to a 4-parameter logistic using SigmaPlot 13 using Log_{10} inhibitor concentration (in μM). In some cases 2–3% (v/v) DMSO was included in assays; no significant change in enzyme activity was observed with DMSO concentrations of up to 8% (v/v) [32]. Half-volume 96 well plates were used for some inhibitors; identical IC_{50} values were obtained for standard inhibitors using both types of microtitre plate. IC_{50} values [32] for N-dodecyl-N-methylcarbamoyl-CoA **4** and ibuprofenoyl-CoA **5** were determined contemporaneously with the inhibitors described in this study, with the same batch of enzyme.

4.17. Computational analysis of potency of inhibition

Lipophilicity of acyl-CoA esters was assessed by calculation of miLogP values using the molecular properties calculator (<http://www.molinspiration.com/cgi-bin/properties>). miLogP values were calculated using molecular smiles obtained from Chemdraw Professional 15. IC₅₀ values (in nM) were plotted against the obtained LogP value using SigmaPlot 13.

Acknowledgements

This work was funded by Prostate Cancer UK (PC-UK) (S10-03 and PG14-009), a University of Bath Overseas Research Studentship and a Biochemical Society Summer Vacation studentship (2015). The authors are members of the Cancer Research @ Bath (CR@B) network.

Disclosure statement

MJ, GLL, MDT, TJW and MDL are named inventors on patent applications on the use and application of the colorimetric assay. The other authors report no conflicts of interest. The authors alone are re-sponsible for the content and writing of this article.

Appendix A. Supplementary material

Supplementary Information contains dose-response curves and other biological evaluation of inhibitors and spectroscopic characterisation data for compounds. Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bioorg.2018.04.024>.

References

- [1] M.D. Lloyd, D.J. Darley, A.S. Wierzbicki, M.D. Threadgill, α -Methylacyl-CoA racemase: An 'obscure' metabolic enzyme takes centre stage, *FEBS J.* 275 (2008) 1089–1102.
- [2] M.D. Lloyd, M. Yevlevskis, G.L. Lee, P.J. Wood, M.D. Threadgill, T.J. Woodman, α -Methylacyl-CoA racemase (AMACR): Metabolic enzyme, drug metabolizer and cancer marker P504S, *Prog. Lipid Res.* 52 (2013) 220–230.
- [3] P.P. VanVeldhoven, K. Croes, S. Asselberghs, P. Herdewijn, G.P. Mannaerts, Peroxisomal β -oxidation of 2-methyl-branched acyl-CoA esters: Stereospecific recognition of the 2S-methyl compounds by trihydroxycoprostanoyl-CoA oxidase and pristanoyl-CoA oxidase, *FEBS Lett.* 388 (1) (1996) 80–84.
- [4] K.P. Battaile, M. McBurney, P.P. Van Veldhoven, J. Vockley, Human long chain, very long chain and medium chain acyl-CoA dehydrogenases are specific for the S-enantiomer of 2-methylpentadecanoyl-CoA, *Biochim. Biophys. Acta - Lipids Lipid Metabol.* 1390 (3) (1998) 333–338.
- [5] P. Kasaragod, W. Schmitz, J.K. Hiltunen, R.K. Wierenga, The isomerase and hydratase reaction mechanism of the crotonase active site of the multifunctional enzyme (type-1), as deduced from structures of complexes with 3S-hydroxy-acyl-CoA, *FEBS J.* 280 (13) (2013) 3160–3175.
- [6] T.J. Woodman, P.J. Wood, A.S. Thompson, T.J. Hutchings, G.R. Steel, P. Jiao, M.D. Threadgill, M.D. Lloyd, Chiral inversion of 2-arylpropionyl-CoA esters by α -methylacyl-CoA racemase 1A (AMACR; P504S), *Chem. Commun.* 47 (2011) 7332–7334.
- [7] D.J. Darley, D.S. Butler, S.J. Prideaux, T.W. Thornton, A.D. Wilson, T.J. Woodman, M.D. Threadgill, M.D. Lloyd, Synthesis and use of isotope-labelled substrates for a mechanistic study on human α -methylacyl-CoA racemase 1A (AMACR; P504S), *Org. Biomol. Chem.* 7 (2009) 543–552.
- [8] P. Bhaumik, W. Schmitz, A. Hassinen, J.K. Hiltunen, E. Conzelmann, R.K. Wierenga, The catalysis of the 1,1-proton transfer by α -methyl-acyl-CoA racemase is coupled to a movement of the fatty acyl moiety over a hydrophobic, methionine-rich surface, *J. Mol. Biol.* 367 (2007) 1145–1161.
- [9] S. Sharma, P. Bhaumik, W. Schmitz, R. Venkatesan, J.K. Hiltunen, E. Conzelmann, A.H. Juffer, R.K. Wierenga, The enolization chemistry of a thioester-dependent racemase: The 1.4 Å crystal structure of a reaction intermediate complex characterized by detailed QM/MM calculations, *J. Phys. Chem. B* 116 (11) (2012) 3619–3629.
- [10] M. Yevlevskis, G.L. Lee, M.D. Threadgill, T.J. Woodman, M.D. Lloyd, The perils of rational design – unexpected irreversible elimination of inorganic fluoride from 3-fluoro-2-methylacyl-CoA esters catalysed by α -methylacyl-CoA racemase (AMACR; P504S), *Chem. Commun.* 50 (2014) 14164–14166.
- [11] M. Yevlevskis, C.R. Bowskill, C.C.Y. Chan, J.H.-J. Heng, M.D. Threadgill, T.J. Woodman, M.D. Lloyd, A study on the chiral inversion of mandelic acid in humans, *Org. Biomol. Chem.* 12 (2014) 6737–6744.
- [12] J.A. Mitchell, P. Akaraseenont, C. Thiemermann, R.J. Flower, J.R. Vane, Selectivity of nonsteroidal antiinflammatory drugs as inhibitors of constitutive and inducible cyclooxygenase, *Proc. Natl. Acad. Sci. USA* 90 (1993) 11693–11697.
- [13] J. Luo, S. Zha, W.R. Gage, T.A. Dunn, J.L. Hicks, C.J. Bennett, C.N. Ewing, E.A. Platz, S. Ferdinandusse, R.J. Wanders, J.M. Trent, W.B. Isaacs, A.M. De Marzo, α -Methylacyl-CoA racemase: A new molecular marker for prostate cancer, *Cancer Res.* 62 (2002) 2220–2226.
- [14] Z. Jiang, B.A. Woda, K.L. Rock, Y.D. Xu, L. Savas, A. Khan, G. Pihan, F. Cai, J.S. Babcock, P. Rathanaswami, S.G. Reed, J.C. Xu, G.R. Fanger, P504S – A new molecular marker for the detection of prostate carcinoma, *Am. J. Surg. Path.* 25 (2001) 1397–1404.
- [15] Z. Jiang, G.R. Fanger, B.F. Banner, B.A. Woda, P. Algate, K. Dresser, J.C. Xu, S.G. Reed, K.L. Rock, P.G. Chu, A dietary enzyme: α -methylacyl-CoA racemase/ P504S is overexpressed in colon carcinoma, *Cancer Detect. Prev.* 27 (2003) 422–426.
- [16] A.K. Witkiewicz, S. Varambally, R.L. Shen, R. Mehra, M.S. Sabel, D. Ghosh, A.M. Chinnaiyan, M.A. Rubin, C.G. Kleer, α -Methylacyl-CoA racemase protein expression is associated with the degree of differentiation in breast cancer using quantitative image analysis, *Cancer Epidemiol. Biomark. Prev.* 14 (2005) 1418–1423.
- [17] C.-F. Li, F.-M. Fang, J. Lan, J.-W. Wang, H.-J. Kung, L.-T. Chen, T.-J. Chen, S.-H. Li, Y.-H. Wang, H.-C. Tai, S.-C. Yu, H.-Y. Huang, AMACR amplification in myxofibrosarcomas: A mechanism of overexpression that promotes cell proliferation with therapeutic relevance, *Clin. Cancer Res.* 20 (23) (2014) 6141–6152.
- [18] Z. Jiang, G.R. Fanger, B.A. Woda, B.F. Banner, P. Algate, K. Dresser, J.C. Xu, P.G.G. Chu, Expression of α -methylacyl-CoA racemase (P504S) in various malignant neoplasms and normal tissues: A study of 761 cases, *Human Pathol.* 34 (2003) 792–796.
- [19] S. Zha, S. Ferdinandusse, S. Denis, R.J. Wanders, C.M. Ewing, J. Luo, A.M. De Marzo, W.B. Isaacs, α -Methylacyl-CoA racemase as an androgen-independent growth modifier in prostate cancer, *Cancer Res.* 63 (2003) 7365–7376.
- [20] C. Kumar-Sinha, R.B. Shah, B. Laxman, S.A. Tomlins, J. Harwood, W. Schmitz, E. Conzelmann, M.G. Sanda, J.T. Wei, M.A. Rubin, A.M. Chinnaiyan, Elevated α -methylacyl-CoA racemase enzymatic activity in prostate cancer, *Am. J. Pathol.* 164 (2004) 787–793.
- [21] B. Ouyang, Y.-K. Leung, V. Wang, E. Chung, L. Levin, B. Bracken, L. Cheng, S.-M. Hi, α -Methylacyl-CoA racemase spliced variants and their expression in normal and malignant prostate tissues, *Urology* 77 (249) (2011) e1–e7.
- [22] G.L. Shen-Ong, Y. Feng, D.A. Troyer, Expression profiling identifies a novel α -methylacyl-CoA racemase exon with fumarate hydratase homology, *Cancer Res.* 63 (2003) 3296–3301.
- [23] J.N. Mubiru, G.L. Shen-Ong, A.J. Valente, D.A. Troyer, Alternative spliced variants of the α -methylacyl-CoA racemase gene and their expression in prostate cancer, *Gene* 327 (2004) 89–98.
- [24] J.N. Mubiru, A.J. Valente, D.A. Troyer, A variant of α -methylacyl-CoA racemase gene created by a deletion in exon 5 and its expression in prostate cancer, *Prostate* 65 (2005) 117–123.
- [25] B.A.P. Wilson, H. Wang, B.A. Nacev, R.C. Mease, J.O. Liu, M.G. Pomper, W.B. Isaacs, High-throughput screen identifies novel inhibitors of cancer biomarker α -methylacyl-coenzyme A racemase (AMACR/P504S), *Mol. Cancer Therapeut.* 10 (5) (2011) 825–838.
- [26] K. Takahara, H. Azuma, T. Sakamoto, S. Kiyama, T. Inamoto, N. Ibuki, T. Nishida, H. Nomi, T. Ubai, N. Segawa, Y. Katsuoka, Conversion of prostate cancer from hormone independency to dependency due to AMACR inhibition: Involvement of increased AR expression and decreased IGF1 expression, *Anticancer Res.* 29 (7) (2009) 2497–2505.
- [27] P.-Y. Lin, K.-L. Cheng, J.D. McGuffin-Cawley, F.-S. Shieu, A.C. Samia, S. Gupta, M. Cooney, C.L. Thompson, C.C. Liu, Detection of alpha-methylacyl-CoA racemase (AMACR), a biomarker of prostate cancer, in patient blood samples using a nano-particle electrochemical biosensor, *Biosensors* 2 (2012) 377–387.
- [28] A.J. Carnell, I. Hale, S. Denis, R.J.A. Wanders, W.B. Isaacs, B.A. Wilson, S. Ferdinandusse, Design, synthesis, and in vitro testing of α -methylacyl-CoA racemase inhibitors, *J. Med. Chem.* 50 (2007) 2700–2707.
- [29] A.J. Carnell, R. Kirk, M. Smith, S. McKenna, L.-Y. Lian, R. Gibson, Inhibition of human α -methylacyl-CoA racemase (AMACR): a target for prostate cancer, *ChemMedChem* 8 (2013) 1643–1647.
- [30] A. Morgenroth, E.A. Urusova, C. Dinger, E. Al-Momani, T. Kull, G. Glatting, H. Fraundorfer, O. Jahn, F.M. Mottaghy, S.N. Reske, B.D. Zlatopolskiy, New molecular markers for prostate tumor imaging: A study on 2-methylene substituted fatty acids as new AMACR inhibitors, *Chem. Eur. J.* 17 (36) (2011) 10144–10150.
- [31] M. Pal, M. Khanal, R. Marko, S. Thirumalaiah, S.L. Bearn, Rational design and synthesis of substrate-product analogue inhibitors of α -methylacyl-coenzyme A racemase from Mycobacterium tuberculosis, *Chem. Commun.* 52 (13) (2016) 2740–2743.
- [32] M. Yevlevskis, G.L. Lee, A. Nathubhai, Y.D. Petrova, T.D. James, M.D. Threadgill, T.J. Woodman, M.D. Lloyd, A novel colorimetric assay for α -methylacyl-CoA racemase 1A (AMACR; P504S) utilizing the elimination of 2,4-dinitrophenolate, *Chem. Commun.* 53 (2017) 5087–5090.
- [33] M. Yevlevskis, G.L. Lee, J. Sun, S. Zhou, X. Sun, G. Kociok-Köhn, T.D. James, T.J. Woodman, M.D. Lloyd, A study on the AMACR catalysed elimination reaction and its application to inhibitor testing, *Org. Biomol. Chem.* 14 (2016) 612–622.
- [34] C. Festuccia, G.L. Gravina, A. Mancini, P. Muzi, E. Di Cesare, R. Kirk, M. Smith, S. Hughes, R. Gibson, L.-Y. Lian, E. Ricevuto, A.J. Carnell, Trifluorobupropfen in-hibits alpha-methylacyl coenzyme A racemase (AMACR/P504S), reduces cancer cell proliferation and inhibits in vivo tumor growth in aggressive prostate cancer models, *Anti-Cancer Ag. Med. Chem.* 14 (7) (2014) 1031–1041.

- [35] M. Pal, N.M. Easton, H. Yaphe, S.L. Bearne, Potent dialkyl substrate-product ana-logue inhibitors and inactivators of α -methylacyl-coenzyme A racemase from *Mycobacterium tuberculosis* by rational design, *Bioorg. Chem.* 77 (2018) 640–650.
- [36] F.A. Sattar, D.J. Darley, F. Politano, T.J. Woodman, M.D. Threadgill, M.D. Lloyd, Unexpected stereoselective exchange of straight-chain fatty acyl-CoA α -protons by human α -methylacyl-CoA racemase 1A (P504S), *Chem. Commun.* 46 (2010) 3348–3350.
- [37] W. Schmitz, C. Albers, R. Fingerhut, E. Conzelmann, Purification and character-ization of α -methylacyl-CoA racemase from human liver, *Eur. J. Biochem.* 231 (1995) 815–822.
- [38] W. Schmitz, R. Fingerhut, E. Conzelmann, Purification and properties of an α -methylacyl-CoA racemase from rat liver, *Eur. J. Biochem.* 222 (1994) 313–323.
- [39] M. Mukherji, C.J. Schofield, A.S. Wierzbicki, G.A. Jansen, R.J.A. Wanders, M.D. Lloyd, The chemical biology of branched-chain lipid metabolism, *Prog. Lipid Res.* 42 (2003) 359–376.
- [40] N.J. Kershaw, M. Mukherji, C.H. MacKinnon, T.D.W. Claridge, B. Odell, A.S. Wierzbicki, M.D. Lloyd, C.J. Schofield, Studies on phytanoyl-CoA 2-hydro-xy-lase and synthesis of phytanoyl-coenzyme A, *Bioorg. Med. Chem. Lett.* 11 (18) (2001) 2545–2548.
- [41] F.J. Reen, S.L. Clarke, C. Legendre, C.M. McSweeney, K.S. Eccles, S.E. Lawrence, F. O’Gara, G.P. McGlacken, Structure-function analysis of the C-3 position in ana-logues of microbial behavioural modulators HHQ and PQS, *Org. Biomol. Chem.* 10 (44) (2012) 8903–8910.
- [42] D.C. Harris, Quantitative Chemical Analysis, 8th Int. Ed. ed., W. H. Freeman, New York, 2010.
- [43] O.H. Straus, A. Goldstein, Zone behavior of enzymes, *J. Gen. Physiol.* 26 (1943) 559–585.
- [44] R.A. Copeland, Evaluation of Enzyme Inhibitors in Drug Discovery. A Guide for Medicinal Chemists and Pharmacologists, John Wiley & Sons, Inc., Hoboken, New Jersey, 2005.
- [45] D.M. Hrubowchak, F.X. Smith, The reductive alkylation of Meldrum’s acid, *Tetrahedron Lett.* 24 (1983) 4951–4954.
- [46] D. Seebach, R. Henning, T. Mukhopadhyay, Doubly deprotonated methyl 3-nitropropanoate, an acrylic ester d^2 -reagent, *Chem. Ber.* 115 (1982) 1705–1720.
- [47] S. Serota, J.R. Simon, E.B. Murray, W.M. Linfield, Novel synthesis of α -substituted acrylic acids, *J. Org. Chem.* 46 (1981) 4147–4151.
- [48] D.J. Grainger, D.J. Fox, New 3-aminocaprolactam derivatives useful in preparation of medicament for prevention or treatment of inflammatory disorder e.g. auto-immune disease, vascular disease, viral infections and asthma, Univ Cambridge Tech Services Ltd; Cambridge Enterprise Ltd, 2005.
- [49] W.R. Shieh, C.S. Chen, Purification and characterization of novel 2-arylpropionyl-CoA epimerases from rat-liver cytosol and mitochondria, *J. Biol. Chem.* 268 (1993) 3487–3493.