










## Rapid report

Differential nucleosome occupancy modulates alternative splicing in *Arabidopsis thaliana*

Author for correspondence:  
Naeem Syed  
Email: [naeem.syed@canterbury.ac.uk](mailto:naeem.syed@canterbury.ac.uk)

Received: 4 September 2020  
Accepted: 24 October 2020

Ibtissam Jabre<sup>1,2\*</sup> , Saurabh Chaudhary<sup>1,3\*</sup> , Wenbin Guo<sup>4</sup> ,  
Maria Kalyna<sup>5</sup> , Anireddy S. N. Reddy<sup>6</sup> , Weizhong Chen<sup>7</sup> ,  
Runxuan Zhang<sup>4</sup> , Cornelia Wilson<sup>1</sup>  and Naeem H. Syed<sup>1</sup> 

<sup>1</sup>School of Human and Life Sciences, Canterbury Christ Church University, Canterbury, CT1 1QU, UK; <sup>2</sup>School of Biosciences and Medicine, University of Surrey, Guildford, GU2 7XH, UK; <sup>3</sup>Cardiff School of Biosciences, Cardiff University, Cardiff, CF10 3AX, UK; <sup>4</sup>Computational Sciences, The James Hutton Institute, Dundee, DD2 5DA, UK; <sup>5</sup>Department of Applied Genetics and Cell Biology, University of Natural Resources and Life Sciences – BOKU, Muthgasse 18, 1190 Vienna, Austria; <sup>6</sup>Department of Biology and Program in Cell and Molecular Biology, Colorado State University, Fort Collins, CO 80523-1878, USA; <sup>7</sup>Department of Molecular Biology & Genetics, Cornell University, Ithaca, NY 14853-2703, USA

*New Phytologist* (2020)  
doi: 10.1111/nph.17062

**Key words:** alternative splicing, *Arabidopsis thaliana*, cold stress, co-transcriptional splicing, exons, nucleosome positioning.

## Summary

- Alternative splicing (AS) is a major gene regulatory mechanism in plants. Recent evidence supports co-transcriptional splicing in plants, hence the chromatin state can impact AS. However, how dynamic changes in the chromatin state such as nucleosome occupancy influence the cold-induced AS remains poorly understood.
- Here, we generated transcriptome (RNA-Seq) and nucleosome positioning (MNase-Seq) data for *Arabidopsis thaliana* to understand how nucleosome positioning modulates cold-induced AS.
- Our results show that characteristic nucleosome occupancy levels are strongly associated with the type and abundance of various AS events under normal and cold temperature conditions in *Arabidopsis*. Intriguingly, exons, alternatively spliced internal regions of protein-coding exons, exhibit distinctive nucleosome positioning pattern compared to other alternatively spliced regions. Likewise, nucleosome patterns differ between exons and retained introns, pointing to their distinct regulation.
- Collectively, our data show that characteristic changes in nucleosome positioning modulate AS in plants in response to cold.

## Introduction

Plants employ different strategies to control their transcriptional program during the daily cycles of light–dark and in response to environmental stress to confer adaptive responses (Zhu, 2016; Laloum *et al.*, 2017; Lämke & Bäurle, 2017). Recent evidence shows that alternative splicing (AS) regulation is a key gene regulatory mechanism in plants (Calixto *et al.*, 2018; Filichkin *et al.*, 2018; Jabre *et al.*, 2019). In plants and animals, AS is regulated co-transcriptionally (Brody *et al.*, 2011; Tilgner *et al.*, 2012; Li *et al.*, 2020; Zhu *et al.*, 2020). RNA polymerase II (RNAPII) speed during transcription may be affected by the

chromatin state that in turn determines AS outcomes (Alexander *et al.*, 2010; Ullah *et al.*, 2018; Zhu *et al.*, 2018). Emerging evidence shows that the chromatin environment has a strong bearing on the splicing process by modulating RNAPII processivity and splicing factors (SFs) recruitment (Nojima *et al.*, 2018; Jabre *et al.*, 2019; Kindgren *et al.*, 2019; Yu *et al.*, 2019; Li *et al.*, 2020; Zhu *et al.*, 2020). Recent native elongating transcript sequencing (NET-Seq) and global run-on sequencing (GRO-Seq) studies from mammals and *Arabidopsis thaliana* (hereafter *Arabidopsis*) show that phosphorylation of RNAPII C-terminal domain mediates interactions with the spliceosome and that RNAPII accumulation is associated with different chromatin states (Nojima *et al.*, 2018; Zhu *et al.*, 2018). Remarkably, sequencing of the chromatin-bound nascent

\*These authors contributed equally to this work.

RNAs in Arabidopsis revealed that almost all introns are spliced co-transcriptionally and the efficiency of intron removal is more robust in protein-coding genes than noncoding RNAs (Li *et al.*, 2020). Furthermore, it was also demonstrated that co-transcriptional splicing (CTS) efficiency is dependent on the number of exons but not gene length (Zhu *et al.*, 2020). Therefore, appropriate exon–intron definition may be important for CTS in Arabidopsis (Li *et al.*, 2020). For example, nucleosome occupancy was found to be higher on exons and also accompanied by a higher level of RNAPII (Chodavarapu *et al.*, 2010). Therefore, higher nucleosome occupancy on exons is intrinsically associated with exon definition, RNAPII processivity and splicing kinetics (Jabre *et al.*, 2019). Previously, it has been demonstrated that changes in RNAPII speed in both directions can influence SFs recruitment and splicing efficiency (Dujardin *et al.*, 2014; Godoy Herz *et al.*, 2019; Leng *et al.*, 2020). Therefore, nucleosome positioning may modulate different AS events and their ratios under variable growth and/or stress conditions to alter intron–exon boundaries and provide a context through which AS patterns could be modulated (Jabre *et al.*, 2019). For example, RNA interference (RNAi) lines of a chromatin remodeler gene (*ZmCHB101*) in maize showed altered nucleosome density, RNAPII elongation rate, and changes in splicing patterns under osmotic stress (Yu *et al.*, 2019). Similarly, widespread nucleosome remodeling in rice, as a result of phosphate starvation and cold stress, was associated with differential gene expression (Roy *et al.*, 2014; Zhang *et al.*, 2018). Since cold can influence the RNAPII elongation kinetics in Arabidopsis (Kindgren *et al.*, 2019), we reasoned that rapid cold-induced AS response in Arabidopsis (Calixto *et al.*, 2018) may be associated with nucleosome remodeling. Henceforth, we used cold treatment as a system of choice to investigate whether it could modulate nucleosome positioning and influence AS.

We performed RNA-Seq and micrococcal nuclease sequencing (MNase-Seq) for Columbia wild-type (Col-0) accession of Arabidopsis plants growing at normal temperature (22°C) and under cold stress (4°C) for 24 h. We observed genome-wide changes in AS and gene expression between Col-0<sub>22°C</sub> and Col-0<sub>4°C</sub> plants. Our results show that temperature-dependent differences in nucleosome positioning are sufficient to modulate different types of AS events and their abundance. Remarkably, exons (EIs), alternatively spliced internal regions of protein-coding exons (Marquez *et al.*, 2015), can also be distinguished from flanking exons by distinctive nucleosome occupancy to facilitate their recognition by the splicing machinery.

## Materials and Methods


The detailed experimental procedure is provided in the Supporting Information (Methods S1). Briefly, leaf tissues were harvested from 3 weeks old Col-0 plants grown at 22°C and 4°C for 24 h. Total RNA and nucleosome bound genomic DNA (gDNA) were extracted for Illumina paired-end sequencing using RNA extraction and Qiagen DNA kits, respectively. The raw reads generated from RNA-Seq and MNase-Seq experiments were quality checked using Trimmomatic (Bolger *et al.*, 2014). The high quality reads from RNA-Seq experiment were used to quantify the transcripts

expression using SALMON v.0.82 (Patro *et al.*, 2017) and AtRTD2-QUASI (Zhang *et al.*, 2017) as reference. Differential expressed genes (DEGs) and differential alternatively spliced (DAS) genes were identified using three-dimensional (3D)-RNA-Seq pipeline as described previously by Calixto *et al.* (2018) and Guo *et al.* (2019). Gene functional enrichment analysis was performed using DAVID v.6.8 (Huang *et al.*, 2009a,b). The gene ontology (GO) terms were assigned to DEGs and DAS genes with a false discovery rate (FDR)  $\leq 0.05$ . AS events, AS event inclusion level (percent spliced in (PSI) indicate how efficiently sequences of interest are spliced into transcripts) and the difference in this inclusion ( $\Delta$ PSI) between Col-0 grown at 22°C and 4°C were identified using SUPPA v.2.3 (Alamancos *et al.*, 2015; Trincado *et al.*, 2018). Only AS events having a Benjamini-Hochberg corrected  $P \leq 0.05$  are identified as DAS events. For MNase-Seq high quality reads were mapped to the TAIR10 Arabidopsis reference genome using BOWTIE v.1.2.2 (Langmead *et al.*, 2009) with ‘-m’ set to 1 to output only uniquely mapped reads. Improved nucleosome-positioning algorithm (iNPS) was used for accurate genome-wide nucleosome positioning as described previously by Chen *et al.* (2014). Differential nucleosome positioning (DNP) analysis was performed using DANPOS v.2.1.2 as described previously by Chen *et al.* (2013). Nucleosome signals were plotted around different genomic regions using DEEPTOOLS v.3.5.0 (Ramírez *et al.*, 2014).

## Results

### Cold-regulated DEGs and DAS genes affect different biological processes

In Arabidopsis, nucleosome positioning differentially marks promoter regions as well as exons and introns, indicating a potential link of chromatin architecture to gene expression and splicing regulation (Chodavarapu *et al.*, 2010). To investigate if cold-induced AS in Arabidopsis is regulated by nucleosome occupancy, we performed RNA-Seq and MNase-Seq of Col-0 ecotype plants before and after a shift from 22°C to 4°C for 24 h. Using the previously published 3D-RNA-Seq pipeline (Calixto *et al.*, 2018; Guo *et al.*, 2019) (Methods S1), we identified 6252 DEGs and 2283 DAS genes. Of the 6252 DEGs, 3323 were upregulated and 2929 were downregulated (Table S1). We observed that most transcriptional changes are associated with genes that do not display splicing changes (70.5% differential expressed only genes). Similarly, a large proportion of splicing changes occur in genes that are not differentially expressed (19.3% DAS only genes). Interestingly, we detected a significant overlap of DEGs and DAS genes (10.2%; hypergeometric test,  $P < 1.222e^{-39}$ ) (Fig. 1a), suggesting that cold stress modulates both transcriptional and AS responses of some genes, which is in line with previously published reports from Arabidopsis (Calixto *et al.*, 2018). Gene functional enrichment analysis of DEGs showed significant (FDR  $< 0.05$ ) enrichment in diverse biological functions, including circadian rhythm, cold stress, and photosynthesis regulation. Cellular components terms enrichment for DEGs were mainly for plasma membrane, and vacuole (Fig. S1a). DAS genes showed significant (FDR  $< 0.05$ ) enrichment in messenger RNA




**Fig. 1** Cold-induced changes in gene expression, alternative splicing, and nucleosome occupancy in *Arabidopsis thaliana*. (a) Venn diagram displaying the overlap between differentially expressed genes (DEGs) and differentially alternatively spliced (DAS) genes. (b) Histogram representing the number of DAS events detected using RNA-Seq data upon cold stress. (c) Venn diagram displaying the overlap between DAS genes and the genes detected within the differential nucleosome positioning (DNP) regions. DEGs and DAS are differentially expressed and alternatively spliced, genes, respectively. DNPs and DNPs-genes are differentially positioned nucleosomes and the genes associated with them, respectively. The  $P$  (hypergeometric test) relates to the significance of overlap. A5'SS, alternative 5' splice site; A3'SS, alternative 3' splice site; IR, intron retention events without exons; MX, mutually exclusive exons; ES, exon skipping; AF, alternative first exon; EI, exons; AL, alternative last exon.

(mRNA) processing, RNA splicing and protein phosphorylation, whereas those of cellular components and molecular functions were mainly enriched in the nucleus and mRNA/ATP/protein binding activities, respectively (Fig. S1b). To identify AS events regulated by cold stress, we analyzed RNA-Seq data using SUPPA v.2.3 (Methods S1) (Alamancos *et al.*, 2015; Trincado *et al.*, 2018). We identified 3032 cold-regulated AS events that showed significant ( $P \leq 0.05$ ) changes in the  $\Delta$ PSI index (Methods S1) distributed within different types of AS events. Changes in intron retention (IR) events are the most prevalent, followed by usage of alternative acceptor (alternative 3' splice site (A3'SS)) and alternative donor (alternative 5' splice site (A5'SS)) sites, exon skipping (ES), and EIs (Tables S2, S3; Figs 1b, S2). EIs are alternatively spliced internal regions of protein-coding exons (Marquez *et al.*, 2015; Staiger & Simpson, 2015; Sibley *et al.*, 2016). At least 6.6% of *Arabidopsis* and 3.7% of human protein-coding genes contain EIs (Marquez *et al.*, 2015; Zhang *et al.*, 2017). Due to their distinctive features, we grouped EIs separately from IR events

### Nucleosome occupancy modulates variety and abundance of AS events

To investigate how nucleosome occupancy modulates splicing, we performed nucleosome positioning and DNP analysis (Methods S1). We detected 19 233 significant (Methods S1) differentially positioned nucleosomes upon shifting plants from 22°C to 4°C for 24 h that were significantly (Methods S1) associated with 7357 genes (Tables S4–S6). Interestingly, 833 (9.5%) (hypergeometric test,  $P < 1.498e^{-24}$ ) cold-induced DAS genes displayed changes in nucleosome occupancy (Fig. 1c). We first profiled nucleosome occupancy across exons and flanking regions, where

we could detect a significant drop of nucleosome occupancy signals at 4°C around exons and flanking regions (one-tailed  $t$ -test,  $P < 0.0001$ ) (Fig. S3). Then, we sought if changes in nucleosome occupancy around the splice sites can modulate different AS events. For that, we profiled nucleosome signals of 3032 cold-regulated DAS events that showed significant ( $P \leq 0.05$ )  $\Delta$ PSI values upon cold stress. Interestingly, different DAS events displayed significant changes in nucleosome occupancy signals around the donor and acceptor sites of different AS events at 22°C and 4°C (one-way ANOVA,  $P < 0.01$ , Fig. 2a). We also could detect an overall drop of nucleosome occupancy level for all AS events upon cold stress. For example, nucleosome occupancy is relatively higher for ES event at 22°C compared to 4°C (one-tailed  $t$ -test,  $P < 0.0001$ , Fig. 2a) potentially impacting exon definition and loss of exons from different transcripts (Fig. 2a). Since nucleosome occupancy levels differentially associate with various types of AS events between plants grown at different temperatures, we sought to explore if this relationship holds true to explain the ratios of these AS events. For that, we grouped PSI values (Methods S1) for different AS events detected in plants grown at both temperature conditions into four bins and aligned nucleosome peaks 200 base pairs (bp) upstream and downstream the exon (or intron) for which the PSI value was calculated. It is notable that for ES, A3'SS and A5'SS, alternative regions with higher inclusion levels (higher PSI values) display more nucleosome occupancy across the splice sites, whereas IRs with higher inclusion levels display less nucleosome occupancy across the splice sites (one-way ANOVA,  $P < 0.001$ , Fig. 2b). Collectively, the differences in nucleosome occupancy levels detected for plants grown at different temperatures for different AS events or for the same PSI group within the same AS event show that alternative




**Fig. 2** Association of nucleosome occupancy with different alternative splicing (AS) events and their different ratios in *Arabidopsis thaliana*. (a) The association of nucleosome occupancy with different AS events. The x-axis is the position relative to the acceptor site (left) and donor site (right); the y-axis is the average of nucleosome signal for the selected genomic regions. An ANOVA was performed to detect the significance of differential nucleosome occupancy around the acceptor site ( $P = 0.015$ ) and donor site ( $P = 0.039$ ) of different AS events at 22°C, and the donor ( $P = 0.0138$ ) and the acceptor sites ( $P = 0.0196$ ) of different AS events at 4°C (b) Nucleosome profiles for different types of AS events grouped based on their percent spliced in (PSI) value. An ANOVA was performed to detect significance of differential nucleosome occupancy of different PSI groups at 22°C and 4°C, respectively; around the acceptor site of A3'SS ( $P = 0.0129$ ,  $P = 0.00112$ ), A5'SS ( $P = 0.00033$ ,  $P = 0.0112$ ), ES ( $P = 0.000129$ ,  $P = 0.00234$ ), and IR ( $P = 0.00236$ ,  $P = 0.132$ ), events. The x-axis is the position relative to the acceptor site; the y-axis is the average of nucleosome signal. ES, exon skipping; A3'SS, alternative 3'SS; A5'SS, alternative 5'SS; IR, intron retention. Constitutive exons or introns are colored in yellow, whereas exons–introns involved in the splicing event are colored in blue. Curved lines indicate a splicing event. Red arrow pointing towards differences in scaling used to plot nucleosome profiles for 22°C and 4°C. Blue arrows indicate regions with significant changes in nucleosome occupancy.

regions involved in different types of AS events may be associated with specific epigenetic features potentially influencing local splicing events and abundance of transcripts.

Nucleosome occupancy is strongly associated with negative or positive AS regulation

Next, we interrogated how nucleosome occupancy levels, for the same set of genes, differ between DAS and non-DAS genes under normal and cold conditions. For that, we profiled nucleosome occupancy levels across the exons of DAS and non-DAS genes. We

found relatively lower nucleosome occupancy for DAS compared to non-DAS exons at 22°C and as well as 4°C (one-tailed  $t$ -test,  $P < 0.0001$ , Fig. 3a). Since nucleosome occupancy globally drops under cold conditions, we sought to investigate how nucleosome occupancy would correlate with splice junctions (SJs) affected differently by cold stress. Therefore, we grouped the SJs of the AS events ( $P \leq 0.05$ ) obtained from SUPPA v.2.3 based on their  $\Delta$ PSI value to obtain positively, negatively, and unaffected SJs (Methods S1). Interestingly, cold stress positively regulates 1208 SJs, negatively regulates 1054, and leaves 673 SJs unaffected (Fig. 3b; Table S7). This data strongly support previous data showing that




**Fig. 3** Profiles of nucleosome occupancy across differentially alternatively spliced (DAS), non-DAS exons and alternatively spliced junctions in *Arabidopsis thaliana*. (a) Nucleosome profiles are plotted against the DAS and non-DAS exons with 500 bp upstream and downstream at 22°C (left) and 4°C (right), respectively. Nucleosome signals data were used in one-tailed *t* test, which confirmed that DAS exons has lower nucleosome occupancy compared to non-DAS exons at 22°C and 4°C ( $P = 4.31977E^{-16}$ ), and that nucleosome signals drop across non-DAS ( $P = 5.32124E^{-18}$ ) and DAS exons ( $P = 1.05564E^{-65}$ ) upon temperature shifts. The x-axis represents DAS/non-DAS exons scaled to 500 bp and their upstream and downstream flanking regions (500 bp); the y-axis represents the average nucleosome signal in the selected genomic regions. (b) Chart illustrating the number of alternatively spliced junctions that are unaffected, positively or negatively affected. Percentages are calculated relative to the significant ( $P \leq 0.05$ ) AS events detected by SUPPA v.2.3. (c) Average nucleosome occupancy level across donor (left) and acceptor (right) regions of all splicing junctions which are unaffected, positively, or negatively affected by cold stress. One-way ANOVA shows the significance of the differences in nucleosome occupancy for the different types of spliced junctions around the donor ( $P = 0.00813$ ; 22°C,  $P = 0.0206$ ; 4°C) and the acceptor ( $P = 0.0293$ ; 22°C,  $P = 0.00733$ , 4°C). Red arrow pointing towards differences in scaling used to plot nucleosome profiles for 22°C and 4°C. Blue arrows indicate regions with significant changes in nucleosome occupancy.

cold stress induces AS in plants (Calixto *et al.*, 2018). To profile nucleosome occupancy around these SJs, we plotted nucleosome density of negatively-affected, positively-affected, and unaffected SJs for Col-0 grown at 22°C and 4°C (Fig. 3c). Remarkably, nucleosome profiles of unaffected SJs for both, the donor and acceptor site are significantly different compared to negatively or positively affected SJs at both temperatures (one-way ANOVA,  $P < 0.01$ , Fig. 3c). Additionally, we also detected a significant association between negatively and positively affected SJs with regions associated with DNPs (Fisher's exact test,  $P < 0.001$ ). Overall, our results show that changes in nucleosome occupancy levels across intron–exon junctions and exons are likely to regulate splice site selection and subsequently modulate splicing regulation in both positive and negative manner.

### Characteristic nucleosome occupancy patterns define exons

EIs have a lower guanine-cytosine (GC) content than adjacent sequences of EI-containing exons (Marquez *et al.*, 2015).

Therefore, we asked whether differential GC content in EI sequences is associated with nucleosome occupancy to distinguish them from flanking exonic regions. To answer this, we profiled nucleosome occupancy across *c.* 2400 EIs identified in *Arabidopsis* (Marquez *et al.*, 2015; Zhang *et al.*, 2017) and 500 bp upstream and downstream from their starts and ends. We found sharp peaks of nucleosome occupancy located before the start and after the end of EIs and slightly lower occupancy in the middle of EIs, which is different from nucleosome patterns observed across exons (Fig. S3). Additionally, we detected a decrease of nucleosome occupancy across EIs under cold stress (one-tailed *t*-test,  $P < 0.0001$ , Fig. 4a). Comparison of nucleosome occupancy levels over EIs grouped into four bins according to the PSI values showed that higher EI inclusion correlates with higher nucleosome occupancy and showed variable levels under normal and cold conditions (Fig. 4b), hence pointing towards their regulation under cold stress. This pattern differs from the one observed for IRs, where IRs with higher inclusion levels display less nucleosome occupancy (Fig. 2b). Interestingly, in this respect, EIs are more similar



**Fig. 4** Nucleosome profiles across exons and their flanking regions in *Arabidopsis thaliana*. (a) Nucleosome profiles across exons and  $-500/+500$  bp flanking regions. Nucleosome signal data collected across exons were used in one-tailed *t*-test, which confirmed that nucleosome signal across exons drops significantly at  $4^{\circ}\text{C}$  ( $P = 5.51\text{E}^{-127}$ ). (b) Nucleosome profiles for exons grouped according to their percent spliced in (PSI) values. The x-axis is the position relative to exons, where Els and Ele are exon start and end, respectively; the y-axis is the average nucleosome signal. (c) Nucleosome occupancy across exons, retained introns, constitutively spliced introns, and their  $-500/+500$  bp flanking regions in each sample. One-way ANOVA has been performed to confirm the significance of the differences in nucleosome occupancy between constitutively spliced introns, exons, and IR<sup>EI</sup> at  $22^{\circ}\text{C}$  ( $P = 5.99\text{e}^{-09}$ ) and  $4^{\circ}\text{C}$  ( $P = 1\text{e}^{-09}$ ). IR<sup>EI</sup>, retained introns excluding exons, S and E, start and end of exons or retained/constitutive introns. Red arrow pointing towards differences in scaling used to plot nucleosome profiles for  $22^{\circ}\text{C}$  and  $4^{\circ}\text{C}$ . Blue arrows indicate regions with significant changes in nucleosome occupancy.

to ES, A5'SS and A3'SS events due to their exonic features (Fig. 2b). Since EIs have a higher GC content than IRs and constitutive introns (Marquez *et al.*, 2015), we compared their nucleosome profiles in normal and cold conditions. We observed that nucleosome occupancy levels are higher for EIs compared to IRs at  $22^{\circ}\text{C}$  and  $4^{\circ}\text{C}$  (one-way ANOVA,  $P < 0.0001$ , Fig. 4c). This implies that chromatin structure plays different roles in the definition and splicing of IRs and EIs. Overall, these results revealed the importance of nucleosome occupancy in defining EIs and their distinction from IRs to regulate their AS profiles under normal and cold conditions.

## Discussion

Recent evidence from *Arabidopsis* shows that transcription and splicing are largely coupled (Dolata *et al.*, 2015; Hetzel *et al.*, 2016; Ullah *et al.*, 2018; Jabre *et al.*, 2019), and that epigenetic features in plants regulate transcriptional activity and differentially mark exons and introns (Zhu *et al.*, 2018). Not surprisingly, very recent studies employing sequencing of chromatin-bound RNAs reveal

that almost all introns in *Arabidopsis* are spliced co-transcriptionally (Li *et al.*, 2020). Furthermore, RNAPII elongation speed has been found to be slower in nucleosome-rich exons allowing more time for the splicing process to take place (Chodavarapu *et al.*, 2010; Zhu *et al.*, 2018). However, how the chromatin environment influences different types of AS events and their ratios under variable growth and stress conditions remains elusive in plants. Since splicing/AS regulation is achieved by the context of the *cis*-regulatory sequences as well as the chromatin environment (Reddy *et al.*, 2013), it is important to understand the relative contributions of epigenetic landscapes. In this study, using *Arabidopsis* Col-0 ecotype plants, we demonstrate that cold-induced DAS is accompanied by changes in nucleosome occupancy levels. Although nucleosome occupancy falls globally under cold conditions in *Arabidopsis*; nonetheless, nucleosome profiles around intron–exon boundaries among different PSI groups, negatively and positively affected AS events displayed characteristic patterns. Further work is needed to understand how variable nucleosome occupancy modulates RNAPII processivity and AS in plants. However, since nucleosome occupancy and RNAPII density has a close

relationship in *Arabidopsis* (Chodavarapu *et al.*, 2010), it is likely that chromatin architecture plays a similar role in plants and animals as progesterone treated breast cancer cells displayed weaker nucleosome densities and lower RNAPII accumulation, resulting in alteration in splice site recognition and ES (Iannone *et al.*, 2015). Since histone modification modulate AS in humans (Luco *et al.*, 2010), similar mechanism also may modulate splicing variation in plants. Intriguingly, EIs also show distinctive nucleosome occupancy (Fig. 4), which may help to differentiate them from flanking exonic regions. Furthermore, despite their classification as a group of IR, EIs display different nucleosome patterns compared to retained introns; pointing to their distinct regulation.

We propose that these changes in nucleosome occupancy may provide the basic definition to exons and introns to coordinate RNAPII processivity. However, it is apparent that the splicing process is also fine-tuned by various *trans*-regulatory factors and histone modifications under variable growth and stress conditions (Kindgren *et al.*, 2019; Zhu *et al.*, 2020). Our data support this notion, and it is likely that higher nucleosome occupancy may regulate RNAPII accumulation around splice sites and enable SF recruitment to facilitate and/or modulate splicing variation. Interestingly, RNAPII elongation speed in *Arabidopsis* would be much slower after clearing a 3'SS and towards the end of an exon, and may not provide sufficient time (because of higher speed in plant introns) for RNAPII to recognize the 5'SS (Kindgren *et al.*, 2019). Furthermore, recent findings show that RNAPII accumulates upstream of the 5'SS, potentially to provide additional time/checkpoint to regulate splicing in mammals and plants (Nojima *et al.*, 2015; Kindgren *et al.*, 2019). Therefore, variation in nucleosome occupancy with an additional peak just after 5'SS (Fig. 3a) may mediate RNAPII accumulation and influence CTS (Nojima *et al.*, 2015; Kindgren *et al.*, 2019). Arguably, this is why 5'SS splicing dynamics are much more complicated and the scanning splicing machinery has to travel to the branch point/polypyrimidine tract to complete lariat formation and process 5'SS. Beggs and colleagues proposed, the initial propensity of splicing is low but increases subsequently to allow accumulation of splicing precursors to improve splicing efficiency in subsequent and/or successive reactions (Aitken *et al.*, 2011). These findings are in broad agreement with CTS in *Arabidopsis* as the CTS process is more efficient in genes with multiple introns–exons and is independent of the gene length (Zhu *et al.*, 2020). Mutations at the 3'SS and 5'SS impact transcription initiation and a mutant 3'SS reduces the first step of CTS in yeast (Aitken *et al.*, 2011). Similarly, splicing dynamics of the human beta-globin gene which fails to form lariat formation and complete 5'SS when a deletion removes the polypyrimidine tract and AG dinucleotide at the 3'SS (Reed & Maniatis, 1985). Therefore, it is tempting to speculate that nucleosome occupancy and/or histone decorations may be more important in the 5' regions of exons providing a checkpoint to the elongating RNAPII to help recognize 5'SS, form lariat and cleave at the 5'SS and 3'SS. It is evident that efficient splicing/AS is dependent on an optimum RNAPII elongation speed and any variation (slow or fast) results in changes in splicing patterns in humans and plants (Dujardin *et al.*, 2014; Godoy Herz *et al.*, 2019; Leng *et al.*, 2020).

Collectively our data points towards the importance of epigenetic features such as nucleosome occupancy for plants grown under different and recurrent growth and stress conditions.










## Acknowledgements

The authors thank the funding agencies for research support. Leverhulme Trust (RPG-2016-014); DOE Office of Science, Office of Biological and Environmental Research (BER) (DE-SC0010733); National Science Foundation and the US Department of Agriculture (ASNR); Austrian Science Fund (FWF) for MK (P26333).

## Author contributions

NHS conceived the study; IJ, SC performed all experiments; IJ, SC, WG, WC, MK, ASNR, RZ, CW and NHS contributed towards data analysis and write up. IJ and SC contributed equally to this work.

## ORCID

Saurabh Chaudhary  <https://orcid.org/0000-0003-2360-3325>  
Weizhong Chen  <https://orcid.org/0000-0001-5170-0398>  
Wenbin Guo  <https://orcid.org/0000-0002-1829-6044>  
Ibtissam Jabre  <https://orcid.org/0000-0003-3574-597X>  
Maria Kalyna  <https://orcid.org/0000-0003-4702-7625>  
Anireddy S. N. Reddy  <https://orcid.org/0000-0002-4038-4091>  
Naeem H. Syed  <https://orcid.org/0000-0003-4043-9969>  
Cornelia Wilson  <https://orcid.org/0000-0001-6584-6179>  
Runxuan Zhang  <https://orcid.org/0000-0001-7558-765X>

## Data availability

All RNA-Seq and MNase-Seq raw data generated in this work is submitted to NCBI-SRA under the accession number 'PRJNA592356'.

## References

- Aitken S, Alexander RD, Beggs JD. 2011. Modelling reveals kinetic advantages of co-transcriptional splicing. *PLoS Computational Biology* 7: e1002215.
- Alamancos GP, Pagès A, Trincado JL, Bellora N, Eyra E. 2015. Leveraging transcript quantification for fast computation of alternative splicing profiles. *RNA* 21: 1521–1531.
- Alexander RD, Innocente SA, Barrass JD, Beggs JD. 2010. Splicing-dependent RNA polymerase pausing in yeast. *Molecular Cell* 40: 582–593.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114–2120.
- Brody Y, Neufeld N, Bieberstein N, Causse SZ, Böhnlein EM, Neugebauer KM, Darzacq X, Shav-Tal Y. 2011. The in vivo kinetics of RNA polymerase II elongation during co-transcriptional splicing. *PLoS Biology* 9: e1000573.
- Calixto CPG, Guo W, James AB, Tzioutziou NA, Entizne JC, Panter PE, Knight H, Nimmo HG, Zhang R, Brown JWS. 2018. Rapid and dynamic alternative splicing impacts the *Arabidopsis* cold response transcriptome. *Plant Cell* 30: 1424–1444.

- Chen K, Xi Y, Pan X, Li Z, Kaestner K, Tyler J, Dent S, He X, Li W. 2013. DANPOS: dynamic analysis of nucleosome position and occupancy by sequencing. *Genome Research* 23: 341–351.
- Chen W, Liu Y, Zhu S, Green CD, Wei G, Han JDJ. 2014. Improved nucleosome-positioning algorithm iNPS for accurate nucleosome positioning from sequencing data. *Nature Communications* 18: 4909–4923.
- Chodavarapu RK, Feng S, Bernatavichute YV, Chen P-Y, Stroud H, Yu Y, Hetzel JA, Kuo F, Kim J, Cokus SJ *et al.* 2010. Relationship between nucleosome positioning and DNA methylation. *Nature* 466: 388–392.
- Dolata J, Guo Y, Kooweroo A, Smolinski D, Brzyzek G, Jarmoowski A, Swiezewski S. 2015. NTR1 is required for transcription elongation checkpoints at alternative exons in Arabidopsis. *EMBO Journal* 34: 544–558.
- Dujardin G, Lafaille C, de la Mata M, Marasco LE, Muñoz MJ, Le Jossic-Corcus C, Corcus L, Kornblihtt AR. 2014. How slow RNA polymerase II elongation favors alternative exon skipping. *Molecular Cell* 54: 683–690.
- Filichkin SA, Hamilton M, Dharmawardhana PD, Singh SK, Sullivan C, Ben-Hur A, Reddy ASN, Jaiswal P. 2018. Abiotic stresses modulate landscape of poplar transcriptome via alternative splicing, differential intron retention, and isoform ratio switching. *Frontiers in Plant Science* 9: 5–22.
- Godoy Herz MA, Kubaczka MG, Brzyzek G, Servi L, Krzyszton M, Simpson C, Brown J, Swiezewski S, Petrillo E, Kornblihtt AR. 2019. Light regulates plant alternative splicing through the control of transcriptional elongation. *Molecular Cell* 73: 1066–1074.
- Guo W, Tzioutziou N, Stephen G, Milne I, Calixto C, Waugh R, Brown JW, Zhang R. 2019. 3D RNA-seq—a powerful and flexible tool for rapid and accurate differential expression and alternative splicing analysis of RNA-seq data for biologists. *bioRxiv*. doi: 10.1101/656686.
- Hetzel J, Duttke SH, Benner C, Chory J. 2016. Nascent RNA sequencing reveals distinct features in plant transcription. *Proceedings of the National Academy of Sciences, USA* 113: 12316–12321.
- Huang DW, Sherman BT, Lempicki RA. 2009a. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 4: 44–57.
- Huang DW, Sherman BT, Lempicki RA. 2009b. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research* 37: 1–13.
- Iannone C, Pohl A, Papsaikas P, Soronellas D, Vicent GP, Beato M, Valcárcel J. 2015. Relationship between nucleosome positioning and progesterone-induced alternative splicing in breast cancer cells. *RNA* 21: 360–374.
- Jabre I, Reddy ASN, Kalyna M, Chaudhary S, Khokhar W, Byrne LJ, Wilson CM, Syed NH. 2019. Does co-transcriptional regulation of alternative splicing mediate plant stress responses? *Nucleic Acids Research* 47: 2716–2726.
- Kindgren P, Ivanov M, Marquardt S. 2019. Native elongation transcript sequencing reveals temperature dependent dynamics of nascent RNAPII transcription in Arabidopsis. *Nucleic Acids Research* 48: 2332–2347.
- Laloum T, Martín G, Duque P. 2017. Alternative splicing control of abiotic stress responses. *Trends in Plant Science* 23: 140–150.
- Lämke J, Bäurle I. 2017. Epigenetic and chromatin-based mechanisms in environmental stress adaptation and stress memory in plants. *Genome Biology* 18: 1–11.
- Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome biology* 10: R25.
- Leng X, Ivanov M, Kindgren P, Malik I, Thieffry A, Sandelin A, Kaplan CD, Marquardt S, Plant C, Centre S *et al.* 2020. Organismal benefits of transcription speed control at gene boundaries. *EMBO Reports* 21: e49315.
- Li S, Wang Y, Zhao Y, Zhao X, Chen X, Gong Z. 2020. Global co-transcriptional splicing in Arabidopsis and the correlation with splicing regulation in mature RNAs. *Molecular Plant* 13: 266–277.
- Luco RF, Pan Q, Tominaga K, Blencowe BJ, Pereira-Smith OM, Misteli T. 2010. Regulation of alternative splicing by histone modifications. *Science* 327: 996–1000.
- Marquez Y, Höpfler M, Ayatollahi Z, Barta A, Kalyna M. 2015. Unmasking alternative splicing inside protein-coding exons defines exitrons and their role in proteome plasticity. *Genome Research* 25: 995–1007.
- Nojima T, Gomes T, Grosso ARF, Kimura H, Dye MJ, Dhir S, Carmo-Fonseca M, Proudfoot NJ. 2015. Mammalian NET-seq reveals genome-wide nascent transcription coupled to RNA processing. *Cell* 161: 526–540.
- Nojima T, Rebelo K, Gomes T, Grosso AR, Proudfoot NJ, Carmo-Fonseca M. 2018. RNA polymerase II phosphorylated on CTD serine 5 interacts with the spliceosome during co-transcriptional splicing. *Molecular Cell* 72: 369–379.
- Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods* 14: 417–419.
- Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. 2014. DeepTools: a flexible platform for exploring deep-sequencing data. *Nucleic Acids Research* 42: W187–W191.
- Reddy ASN, Marquez Y, Kalyna M, Barta A. 2013. Complexity of the alternative splicing landscape in plants. *Plant Cell* 25: 3657–83.
- Reed R, Maniatis T. 1985. Intron sequences involved in lariat formation during pre-mRNA splicing. *Cell* 41: 95–105.
- Roy D, Paul A, Roy A, Ghosh R, Ganguly P, Chaudhuri S. 2014. Differential acetylation of histone H3 at the regulatory region of OsDREB1b promoter facilitates chromatin remodelling and transcription activation during cold stress. *PLoS ONE* 9: e100343.
- Sibley CR, Blazquez L, Ule J. 2016. Lessons from non-canonical splicing. *Nature Reviews Genetics* 17: 407–421.
- Staiger D, Simpson GG. 2015. Enter exitrons. *Genome Biology* 16: 1–3.
- Tilgner H, Knowles DG, Johnson R, Davis CA, Chakraborty S, Djebali S, Curado J, Snyder M, Gingeras TR, Guigó R. 2012. Deep sequencing of subcellular RNA fractions shows splicing to be predominantly co-transcriptional in the human genome but inefficient for lncRNAs. *Genome Research* 22: 1616–1625.
- Trincado JL, Entizne JC, Hysenaj G, Singh B, Skalic M, Elliott DJ, Eyra E. 2018. SUPPA2: Fast, accurate, and uncertainty-aware differential splicing analysis across multiple conditions. *Genome Biology* 19: 1–11.
- Ullah F, Hamilton M, Reddy ASN, Ben-Hur A. 2018. Exploring the relationship between intron retention and chromatin accessibility in plants. *BMC Genomics* 19: 21–32.
- Yu X, Meng X, Liu Y, Wang X, Wang TJ, Zhang A, Li N, Qi X, Liu B, Xu ZY. 2019. The chromatin remodeler ZmCHB101 impacts alternative splicing contexts in response to osmotic stress. *Plant Cell Reports* 38: 131–145.
- Zhang Q, Oh DH, DiTusa SF, RamanaRao MV, Baisakh N, Dassanayake M, Smith AP. 2018. Rice nucleosome patterns undergo remodeling coincident with stress-induced gene expression. *BMC Genomics* 19: 1–16.
- Zhang R, Calixto CPG, Marquez Y, Venhuizen P, Tzioutziou NA, Guo W, Spensley M, Entizne JC, Lewandowska D, Ten Have S *et al.* 2017. A high quality Arabidopsis transcriptome for accurate transcript-level analysis of alternative splicing. *Nucleic Acids Research* 45: 5061–5073.
- Zhu D, Mao F, Tian Y, Lin X, Gu L, Gu H, Qu L, Wu Y, Wu Z. 2020. The features and regulation of co-transcriptional splicing in Arabidopsis. *Molecular Plant* 13: 278–294.
- Zhu JK. 2016. Abiotic stress signaling and responses in plants. *Cell* 167: 313–324.
- Zhu J, Liu M, Liu X, Dong Z. 2018. RNA polymerase II activity revealed by GRO-seq and pNET-seq in Arabidopsis. *Nature Plants* 4: 1112–1123.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

**Fig. S1** Gene ontology (GO) term enrichment analysis of differentially expressed genes (DEGs) and differentially alternatively spliced (DAS) genes.

**Fig. S2** Distribution of the mean changes in percent spliced in (PSI) values along with the expression of different alternative splicing (AS) transcripts.



**Fig. S3** Genome-wide representation of nucleosome occupancy levels displaying a drop in nucleosome signal across exons and flanking regions.

**Method S1** Details of the experimental procedure.

**Table S1** Differentially expressed (DE) and differentially alternatively spliced (DAS) genes identified in *Arabidopsis thaliana* upon exposure to cold stress.

**Table S2** Event coordinates extracted from AtRTDv2 annotation file using SUPPA as well as the percent spliced in (PSI) value of each event in different samples.

**Table S3** Significant ( $P \leq 0.05$ ) differential splicing events for local alternative splicing events ( $\Delta$ PSI value) detected by SUPPA.

**Table S4** List of nucleosomes detected by iNPS at 22°C and 4°C on chromosomes 1–5.

**Table S5** Differential nucleosome positioning analysis performed by SUPPA and iNPS.

**Table S6** Genes associated with windows enriched with the differentially positioned nucleosomes.

**Table S7** Splice junctions (SJs) grouped into positively, negatively, and unaffected SJs.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



## About *New Phytologist*

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Foundation, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews and Tansley insights.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit [www.newphytologist.com](http://www.newphytologist.com) to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office ([np-centraloffice@lancaster.ac.uk](mailto:np-centraloffice@lancaster.ac.uk)) or, if it is more convenient, our USA Office ([np-usaoffice@lancaster.ac.uk](mailto:np-usaoffice@lancaster.ac.uk))
- For submission instructions, subscription and all the latest information visit [www.newphytologist.com](http://www.newphytologist.com)