

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/140359/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Peterson, Roseann E., Bigdeli, Tim B., Ripke, Stephan, Bacanu, Silviu-Alin, Gejman, Pablo V., Levinson, Douglas F., Li, Qingqin S., Rujescu, Dan, Rietschel, Marcella, Weinberger, Daniel R., Straub, Richard E., Walters, James T.R. , Owen, Michael J. , O'Donovan, Michael C. , Mowry, Bryan J., Ophoff, Roel A., Andreassen, Ole A., Esko, Tõnu, Petryshen, Tracey L., Kendler, Kenneth S. and Fanous, Ayman H. 2021. Genome-wide analyses of smoking behaviors in schizophrenia: findings from the Psychiatric Genomics Consortium. *Journal of Psychiatric Research* 137 , pp. 215-225. 10.1016/j.jpsychires.2021.02.027

Publishers page: <http://dx.doi.org/10.1016/j.jpsychires.2021.02.027>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Running Head: Genetic analyses of smoking behaviors among schizophrenia cases

**Genome-wide analyses of smoking behaviors in schizophrenia:
findings from the Psychiatric Genomics Consortium**

Roseann E Peterson, PhD^{1,‡}; Tim B Bigdeli, PhD^{1,2,‡}; Stephan Ripke, MD, PhD^{3,4}; Silviu-Alin Bacanu, PhD¹; Pablo V Gejman, MD^{5,6}; Douglas F Levinson, MD⁷; Qingqin S Li, PhD⁸; Dan Rujescu, MD, PhD^{10,9}; Marcella Rietschel, MD, PhD¹¹; Daniel R Weinberger, MD^{12,13}; Richard E Straub, PhD¹²; James TR Walters, MD, PhD¹⁴; Michael J Owen, MBChB, PhD¹⁴; Michael C O'Donovan, MBChB, PhD¹⁴; Bryan J Mowry, MD, FRANZCP^{15,16}; Roel A Ophoff, PhD^{17,18,19}; Ole A Andreassen, MD, PhD^{20,21}; Tõnu Esko, PhD^{22,23,24,25}; Tracey L Petryshen, PhD^{26,27,28}; Kenneth S Kendler, MD¹; Schizophrenia Working Group of the Psychiatric Genomics Consortium^a; Ayman H Fanous, MD^{1,2,29*}

¹Virginia Institute for Psychiatric and Behavioral Genetics, Department of Psychiatry, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

²Department of Psychiatry and Behavioral Sciences, State University of New York Downstate Medical Center, Brooklyn, NY, USA

³Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA

⁴Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA

⁵Department of Psychiatry and Behavioral Sciences, NorthShore University HealthSystem, Evanston, IL, USA

⁶Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL, USA

⁷Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA

⁸Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Raritan, NJ, USA

⁹Department of Psychiatry, University of Halle, Halle, Germany

¹⁰Department of Psychiatry, University of Munich, Munich, Germany

¹¹Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, University of Heidelberg, Heidelberg, Germany

¹²Lieber Institute for Brain Development, Baltimore, MD, USA

¹³Departments of Psychiatry, Neurology, Neuroscience and Institute of Genetic Medicine, Johns Hopkins School of Medicine, Baltimore, MD

¹⁴MRC Centre for Neuropsychiatric Genetics and Genomics, Institute of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK

¹⁵Queensland Brain Institute, The University of Queensland, Brisbane, Queensland, Australia

¹⁶Queensland Centre for Mental Health Research, University of Queensland, Brisbane, Queensland, Australia

¹⁷Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, CA, USA

¹⁸Department of Human Genetics, David Geffen School of Medicine, University of California, Los Angeles, CA, USA

¹⁹Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht, The Netherlands

²⁰NORMENT, KG Jebsen Centre for Psychosis Research, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

²¹Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway

²²Estonian Genome Center, University of Tartu, Tartu, Estonia

²³Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA, USA

²⁴Department of Genetics, Harvard Medical School, Boston, MA, USA

²⁵Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge, MA, USA

²⁶Center for Human Genetic Research and Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA

²⁷Department of Psychiatry, Harvard Medical School, Boston, MA, USA

²⁸The Broad Institute of MIT and Harvard, Boston MA

²⁹VA New York Harbor Healthcare System, Brooklyn NY, USA

*Corresponding author

Department of Psychiatry and Behavioral Sciences

SUNY Downstate Medical Center

450 Clarkson Ave

Brooklyn, NY 11203

‡These authors contributed equally to this work

^aFull list of consortium members appears in the supplement.

Word Count: Abstract = 245, Text = 4673, Tables/Figures = 6, References = 63.

Abstract

While 17% of US adults use tobacco regularly, smoking rates among persons with schizophrenia are upwards of 60%. Research supports a shared etiological basis for smoking and schizophrenia, including findings from genome-wide association studies (GWAS). However, few studies have directly tested whether the same or distinct genetic variants also influence smoking behavior among schizophrenia cases. Using data from the Psychiatric Genomics Consortium (PGC) study of schizophrenia (35476 cases, 46839 controls), we estimated genetic correlations between these traits and tested whether polygenic risk scores (PRS) constructed from the results of smoking behaviors GWAS were associated with schizophrenia risk or smoking behaviors among schizophrenia cases. Results indicated significant genetic correlations of schizophrenia with smoking initiation ($r_g=0.159$; $P=5.05\times 10^{-10}$), cigarettes-smoked-per-day ($r_g=0.094$; $P=0.006$), and age-of-onset of smoking ($r_g=0.10$; $P=0.009$). Comparing smoking behaviors among schizophrenia cases to the general population, we observe positive genetic correlations for smoking initiation ($r_g=0.624$, $P=0.002$) and cigarettes-smoked-per-day ($r_g=0.689$, $P=0.120$). Similarly, TAG-based PRS for smoking initiation and cigarettes-smoked-per-day were significantly associated with smoking initiation ($P=3.49\times 10^{-5}$) and cigarettes-smoked-per-day ($P=0.007$) among schizophrenia cases. We performed the first GWAS of smoking behavior among schizophrenia cases and identified a novel association with cigarettes-smoked-per-day upstream of the *TMEM106B* gene on chromosome 7p21.3 (rs148253479, $P=3.18\times 10^{-8}$, $n=3520$). Results provide evidence of a partially shared genetic basis for schizophrenia and smoking behaviors. Additionally, genetic risk factors for smoking behaviors were largely shared across schizophrenia and non-schizophrenia populations. Future research should address mechanisms underlying these associations to aid both schizophrenia and smoking treatment and prevention efforts.

Key Words: schizophrenia, genetics, GWAS, smoking initiation, cigarettes per day, pleiotropy.

Introduction

Schizophrenia is a chronic mental illness affecting nearly 1% of the world's population and is associated with considerable morbidity and mortality (McGrath et al., 2008; Simeone et al., 2015). Affected persons are at markedly increased risk for substance use disorders, particularly nicotine dependence (Hartz et al., 2014; Volkow, 2009). Currently, 17% of US adults and upwards of 60% of schizophrenia spectrum cases smoke tobacco regularly (de Leon and Diaz, 2005; Jamal et al., 2015; Volkow, 2009). Furthermore, patients tend to smoke a greater number of cigarettes, extract more nicotine per cigarette, and experience greater withdrawal symptoms than smokers in the general population (Centers for Disease Control and Prevention (CDC), 2013; Strand and Nybäck, 2005; Tidey et al., 2014), thereby increasing their risk of nicotine dependence and associated adverse medical conditions including cardiovascular disease and cancers (Olsson et al., 2015).

Decades of twin and family studies have demonstrated that schizophrenia is highly heritable (~80%) (Sullivan et al., 2003). Common genetic variants captured by genome-wide single-nucleotide polymorphism (SNP) arrays account for at least one third of variance in risk (International Schizophrenia Consortium et al., 2009; Ripke et al., 2013). A landmark genome-wide association study (GWAS) meta-analysis of schizophrenia identified 108 robustly associated loci (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014), one of which resides in a gene cluster encoding neuronal nicotinic acetylcholine receptors (nAChR) on chromosome 15q25, which has previously been shown to be associated with heaviness of smoking in the general population (Tobacco and Genetics Consortium, 2010).

Similarly, twin and family studies have consistently shown a significant genetic component to the liability of smoking behavior, with estimated heritabilities on the order of 0.50-

0.70 for smoking initiation and 0.60 for nicotine dependence among European ancestry populations (Maes et al., 2004; Vink et al., 2005). Large, population-based GWAS of smoking-related traits have yielded several putative risk variants, including an association between smoking initiation and *BDNF* on 11p14.1 (Tobacco and Genetics Consortium, 2010) and several associations for smoking quantity, most notably the previously reported 15q25 locus harboring three genes encoding nAChR subunits *CHRNA5-CHRNA3-CHRNA4*, a second locus encoding nAChRs on 8p11 in and near *CHRNA3-CHRNA6*, and variants in and near *CYP2A6-CYP2B6* on 19q13 encoding nicotine metabolizing enzymes (Thorgeirsson et al., 2010; Tobacco and Genetics Consortium, 2010).

Mechanisms underlying the schizophrenia-smoking association are not completely understood. Several mechanisms have been proposed to explain elevated tobacco use in those with schizophrenia including: 1) the self-medication hypothesis, 2) that smoking causes schizophrenia, and 3) a shared liability underlying both traits. The self-medication hypothesis posits that smoking is used as a strategy to alleviate adverse positive or negative symptoms of schizophrenia, cognitive impairments, or medication side-effects (Kumari and Postma, 2005). The shared vulnerability hypothesis postulates that factors common to both disorders (i.e. genetic, environmental) drive their co-occurrence. For example, dysfunction in nAChRs represents a common substrate for various symptoms of schizophrenia and comorbid nicotine use (Parikh et al., 2016; Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014; Tobacco and Genetics Consortium, 2010). It has been suggested that reduced expression of the $\alpha 7$ -nicotinic receptor in schizophrenia (Guillozet-Bongaarts et al., 2014; Severance and Yolken, 2008) results in reduced sensory gating inhibition as measured by paradigms such as P50 auditory-evoked potentials, prepulse inhibition, and mismatch negativity (Freedman, 2014).

Such deficits could conceivably diminish an individual's ability to keep extraneous stimuli from awareness, possibly giving rise to hallucinations and delusions(Howes and Kapur, 2009).

Additional research supports mechanisms 2 and 3(Chen et al., 2016; Gurillo et al., 2015; Kendler et al., 2015). For example, in a population-based Swedish cohort it was found that smoking prospectively predicted risk for schizophrenia in a dose-response relationship and shared familial/genetic factors accounted for a portion of the comorbidity between smoking and schizophrenia(Kendler et al., 2015).

Recent findings support a molecular genetic component underlying schizophrenia-smoking associations(Chen et al., 2016; Hartz et al., 2018, 2017) but has not been demonstrated conclusively(Brainstorm Consortium et al., 2018; B. Bulik-Sullivan et al., 2015; Gage et al., 2017; Gage and Munafò, 2015; Zheng et al., 2017). Therefore, in this study, we sought to advance the understanding of schizophrenia-smoking associations in the context of available smoking data in schizophrenia cases from the Psychiatric Genomics Consortium (PGC) study of schizophrenia (see Table 1 for study overview)(Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). First, we leverage summary statistics from genome-wide findings to estimate genetic correlations between smoking behaviors and schizophrenia. Next, using available phenotypic data on smoking initiation and smoking quantity for >5000 schizophrenia cases from 10 participating studies, we consider whether polygenic risk scores (PRS) constructed from results of the Tobacco and Genetics (TAG) consortium study of smoking behaviors(Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014; Tobacco and Genetics Consortium, 2010) can predict these same behaviors among schizophrenia cases. We perform the largest GWAS of smoking behaviors among schizophrenia cases to date. Finally, we consider whether smoking patterns among schizophrenia cases and genetic risk

factors for smoking are related to the clinical presentation of schizophrenia including age-of-onset and symptom-based positive, negative, manic, and depressive factor scores.

Methods

Ascertainment and assessment

The subsamples included in this study comprise 10 constituent sites from Stage 2 of the PGC study of schizophrenia (Table 2). Ascertainment, diagnostic assessment, genotyping, and genotype quality control have been previously described (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). Briefly, 52 samples from the US, Europe, and Australia comprising 34,241 cases, 45,604 controls, and 1,235 parent affected-offspring trios were genotyped using a number of commercial SNP genotyping platforms. These data were processed using the stringent PGC quality control procedures, followed by imputation of SNPs and insertion-deletions using the 1000 Genomes Project reference panel (UCSC hg19/NCBI 37) (1000 Genomes Project Consortium et al., 2012; Sachidanandam et al., 2010) using IMPUTE2 (Howie et al., 2011, 2009), resulting in nearly 9.5M markers for GWAS analysis.

Smoking behavior and clinical phenotypes

Smoking behavior variables were harmonized across sites. Smoking initiation was coded as positive if any of the following were endorsed: ever smoked, ever regular smoker, smoked 100 cigarettes, current smoker, former smoker, smoke 1 or more cigarettes-per-day, or nicotine dependence. Since smoking quantity data varied by site, cigarettes-smoked-per-day was centered and scaled for each cohort. To account for initiation, only those who endorsed ever smoked were included in genetic analyses of cigarettes-smoked-per-day. A summary of individual sites, their

sample sizes, and smoking measures available are presented in Table 2 and Supplementary Figures S1 and S2.

We assessed whether age-of-onset of schizophrenia or symptom-based factor scores representing dimensions of illness were associated with smoking behaviors among cases. Age-of-onset was determined retrospectively and defined the age at first diagnosis or hospitalization. Symptoms averaged over the course of illness were assessed using the Operational Criteria Checklist for Psychotic Illness and Affective Illness (OPCRIT), Positive and Negative Syndrome Scale (PANSS), Lifetime Dimensions of Psychosis Scale (LDPS), Schedules for Clinical Assessment in Neuropsychiatry (SCAN), Structured Clinical Interview for DSM (SCID), and Comprehensive Assessment of Symptoms and History (CASH). Factor analyses of constituent PGC studies identified positive, negative, manic, and depressive symptom dimensions and methodological details can be found in Ruderfer et al (Ruderfer et al., 2014). Association between smoking initiation or cigarettes-smoked-per-day and each clinical measure was assessed by logistic and linear regression, respectively, including sex, age-at-interview, and a study site indicator as covariates.

Estimation of SNP-based heritability and genetic correlation

We obtained estimates of SNP-based heritability (h^2) and genetic correlation (r_g) using the LD-score regression approach, as previously described (B. K. Bulik-Sullivan et al., 2015). Genome-wide summary statistics for schizophrenia and TAG smoking-related traits (ever-smoked, cigarettes-per-day, smoking cessation “former vs current”, log-transformed age-of-onset of smoking, logOnset) were filtered using default parameters (INFO>0.9, MAF>1%). Reference LD-scores estimated for European populations in the 1000 Genomes Project were used;

regression weights were based on common SNPs present in Hapmap Phase 3, as suggested by the developers of this approach (B. K. Bulik-Sullivan et al., 2015). We reduced potential bias in heritability estimation by reanalyzing the PGC schizophrenia with overlapping TAG samples omitted, and constraining regression intercepts to one and zero when estimating univariate heritability and genetic correlation, respectively. For the schizophrenia case-only binary trait of smoking initiation, we assumed population prevalence estimates (K) equal to the observed sample prevalence (Supplemental Table S1).

Replication of the observed r_g between schizophrenia risk and TAG traits utilized meta-analysis summary statistics for three East-Asian studies from the PGC (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014) and the *popcorn* method for estimating cross-ancestry correlations (Brown et al., 2016). We compared estimates based on European and East-Asian schizophrenia samples by assuming an approximately normal distribution for r_g and obtaining a Z-score for the difference in values.

Polygenic scoring analyses

To test for polygenic effects on smoking behaviors or schizophrenia risk, we performed risk score profiling as previously described (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). We constructed scores based on TAG results for smoking behaviors (Tobacco and Genetics Consortium, 2010). Given differences in the imputation reference panels between the TAG and PGC2 studies, we considered only overlapping SNPs with imputation INFO greater than 0.9 and minor allele frequency (MAF) greater than 1% in PGC2. Schizophrenia risk scores were generated for each study site in the PGC2 study of schizophrenia, using every other study as the training set in an iterative, "leave-one-out"

procedure. This approach ensured no overlap in training and testing samples, while offering improved power compared to subdividing the full cohort into approximate halves. For both PGC-schizophrenia and TAG-based analyses, we computed several scores based on varying P -value threshold signifying the proportion of SNPs with smaller P -values in the training set; P -value thresholds (P_t) ranged between 0.0001 and 1.0. We tested for association between smoking behaviors and schizophrenia-PRS by linear regression, adjusting for sex, age, study-site and 10 associated ancestry principal components (PCs). Association between schizophrenia risk and TAG-based scores was assessed by logistic regression, adjusting for study-site and all covariates used in the primary PGC-schizophrenia association analysis. Because controls subjects from the Molecular Genetics of Schizophrenia (MGS) study were included in the TAG study, we excluded MGS from our case-control analyses of schizophrenia; in the context of genetic correlation estimation from summary statistics, this permitted us to constrain the intercept.

Genome-wide association and replication sample

For each trait, associated ancestry PCs were identified for the full cohort by backwards-stepwise regression ($P < 0.159$), after adjusting for study site. We tested for association between SNPs and each trait by either linear or logistic regression, as implemented in PLINK v1.07 (<http://pngu.mgh.harvard.edu/~purcell/plink/>) (Purcell et al., 2007), using allelic dosages and adjusting for significant covariates including sex, age, and ancestry PCs. We performed GWAS of each trait separately for individual study sites, combining summary statistics in subsequent random-effects meta-analyses using METAL (Willer et al., 2010). We excluded all SNPs with MAF less than 0.01, average statistical imputation information (INFO) less than 0.6, absent from

more than half of total number of sub-studies, or displayed evidence of excessive heterogeneity (Cochran's test P -value < 0.05).

For replication efforts, a total of 1802 European-ancestry cases with complete phenotypic information from four independent “waves” were made available by Janssen Pharmaceuticals (Li et al., 2017; Metspalu et al., 2004). We identified independent (pairwise linkage disequilibrium $r^2 < 0.1$ within 500kb based on European 1000 Genomes Project samples), significant SNPs ($P < 10^{-5}$) from the random-effects meta-analyses of each smoking behavior phenotype considered. We tested these SNPs for association by linear or logistic regression, using ancestry PCs, sex, age, and study site indicator as covariates. Subsequent, joint meta-analyses of the combined discovery and replication samples were performed using METAL.

Results

I. Genetic correlations between schizophrenia and smoking behaviors

We first estimated the genetic correlation (r_g) between schizophrenia (35,476 cases, 46,839 controls) and each smoking-related trait from TAG (Table 3). The estimated genetic correlation between schizophrenia and smoking initiation in the general population was positive and highly significant ($r_g=0.159$, 95% CI:[0.108,0.210]; $P=5.05 \times 10^{-10}$); significant positive relationships between schizophrenia and both cigarettes-smoked-per-day ($r_g=0.094$, 95% CI:[0.027,0.161]; $P=0.006$) and age-of-onset of smoking ($r_g=0.100$, 95% CI:[0.026,0.174]; $P=0.009$) were also seen; a nominal association between schizophrenia and smoking cessation was found ($r_g=-0.076$, 95% CI:[-0.145,-0.007]; $P=0.032$). We sought to replicate the observed genetic correlations using an independent cohort of East-Asian schizophrenia cases ($n=1836$) and controls ($n=3383$).

No statistically significant cross-ancestry correlations were observed (Table 3) but confidence

intervals overlapped with the results from the European ancestry cohorts. Results for the East-Asian cohort should be considered tentative until replication can be performed.

II. Association of smoking behavior polygenic risk scores with schizophrenia risk

We evaluated the predictive ability of polygenic scores based on TAG results for smoking behaviors as applied to the PGC study of schizophrenia (35,476 cases, 46,839 controls, Figure 1). Genome-wide scores for smoking initiation (“ever/never smoked”) were higher among cases ($P_T < 0.3$, $\beta=0.014$, 95%CI:[0.010, 0.017], $P=4.94 \times 10^{-15}$), explaining 0.14% of the variance in schizophrenia risk. Scores based on independent SNPs significant at $P_T < 10^{-5}$ in TAG for cigarettes-smoked-per-day were also significantly higher among schizophrenia cases compared to controls ($\beta=0.026$, 95%CI:[0.014, 0.038], $P=3.58 \times 10^{-5}$), explaining 0.04% of the variance in schizophrenia risk. Although, this effect was attenuated at more inclusive P -value thresholds. Genome-wide scores based on TAG results for age-of-initiation of smoking were not associated with schizophrenia status ($P>0.056$). Scores based on TAG results for smoking cessation (“former vs current”) were significant, though only for $P_T < 10^{-4}$ and were in the expected negative direction of effect ($P_T < 10^{-5}$, $\beta=-0.130$, 95%CI:[-0.222, -0.038], $P=0.006$).

We further investigated significant polygenic score associations in order to determine if they were driven by the chromosome 15q25 locus that has been independently associated with both schizophrenia risk and smoking quantity in the general population. SNPs from the 15q25 locus were removed from TAG polygenic scores (up to 171 SNPs depending on P_T) and were re-tested for association. Results remained robust for TAG-smoking initiation polygenic scores predicting schizophrenia ($P_T < 0.3$, $\beta=0.014$, 95%CI:[0.010, 0.017], $P=1.13 \times 10^{-13}$) but associations with schizophrenia were attenuated for TAG-cigarettes-smoked-per-day ($P_T < 10^{-5}$).

⁵, $\beta=0.015$, 95%CI:[-0.025, 0.054], $P=0.469$), and smoking cessation ($P_T < 10^{-5}$, $\beta=-0.086$, 95%CI:[-0.182, 0.011], $P=0.081$). These results suggest that the association seen between TAG-smoking initiation polygenic scores and schizophrenia were not due to confounding with the 15q25. However, associations of TAG-cigarettes-smoked-per-day and TAG-Cessation scores with schizophrenia were largely driven by this locus.

III. Smoking behavior among schizophrenia cases

The average smoking initiation rate across all cohorts was 72.9% and ranged from 52.6 to 77.3% (Table 2, Figure S1). Among schizophrenia cases that smoke 29.5% smoked more than a pack per day. Figure S2 displays prevalence of smoking quantity by cohort.

III.1 Heritability of smoking behavior among schizophrenia cases

We applied the LD-score regression method to directly estimate SNP-based heritability (SNP- h^2) from GWAS summary statistics for smoking behaviors among schizophrenia cases. For neither smoking initiation nor cigarettes-smoked-per-day did observed inflation of genome-wide test statistics indicate confounding by population stratification, as indicated by regression intercept values close to one (0.998 and 0.999). Among schizophrenia cases, the SNP-based heritability of smoking initiation was estimated as 0.219 (95% CI:[-0.001,0.439]; $P=0.051$; $n=5255$); the corresponding estimate for cigarettes-smoked-per-day was 0.0917 (95% CI:[-0.096,0.280]; $P=0.340$; $n=3370$). That neither estimate was robustly statistically significant likely reflects the modest sample size. Although the SNP-based heritability point estimates for smoking behavior among schizophrenia cases were larger than the general population, their confidence intervals were overlapping (general population: smoking initiation SNP- $h^2=0.075$ (95% CI:[0.063,0.088], cigarettes-smoked-per-day SNP- $h^2=0.056$ (95% CI:[0.030,0.083])).³³

III.2 Genetic correlations between smoking behaviors among schizophrenia cases and the general population

We estimated the r_g to determine the magnitude of genetic overlap of smoking behaviors between schizophrenia cases ($n_{\text{smoking initiation}}=5255$, $n_{\text{cigarettes-smoked-per-day}}=3370$) and the general population (TAG). We observed a significant positive genetic correlation for smoking initiation ($r_g=0.624$, 95% CI: [0.228, 1.020]; $P=0.002$). Though, a positive relationship for cigarettes-smoked-per-day was not statistically significant ($r_g=0.689$, 95% CI: [-0.179, 1.557]; $P=0.120$). Given the small sample size schizophrenia cases with data on smoking behaviors, these analyses are considered exploratory and require replication.

III.3 Association of smoking behavior polygenic risk scores with smoking behavior among schizophrenia cases

We considered whether TAG scores for smoking initiation and cigarettes-smoked-per-day could predict smoking behaviors among schizophrenia subjects (Figure 1). TAG-based scores for smoking initiation significantly predicted initiation among schizophrenia cases ($P_T < 0.01$, $\beta=0.087$, 95%CI:[0.049, 0.126], $P=9.57 \times 10^{-6}$, $n=5255$) accounting for 0.6% of the variance and results remained robust when SNPs from the 15q25 locus were removed from scores ($P_T < 0.01$, $\beta=0.083$, 95%CI:[0.042, 0.125], $P=8.12 \times 10^{-5}$, Nagelkerke's pseudo- $R^2=0.0046$). The scores based on TAG results for cigarettes-smoked-per-day also significantly predicted cigarettes-smoked-per-day among schizophrenia cases ($P_T < 0.01$, $\beta=0.005$, 95%CI:[0.002, 0.008], $P=8.57 \times 10^{-4}$, $n=3370$) accounting for 0.35% of the variance and results remained robust when SNPs from the 15q25 locus were removed from scores ($P_T < 0.01$, $\beta=0.006$, 95%CI:[0.003,

0.009], $P=4.42 \times 10^{-4}$, $R^2=0.0039$). For both smoking behaviors, the direction of the observed effect in schizophrenia was the same as that observed in the general population.

Next, we asked whether aggregate genetic risk of schizophrenia, as indexed by PGC2-based polygenic scores, was significantly associated with smoking behavior among schizophrenia cases (Figure S3). Neither smoking behaviors (smoking initiation, cigarettes-smoked-per-day) among schizophrenia-cases showed association with schizophrenia risk scores ($P>0.1$). Complete results for polygenic scoring analyses are reported in Tables S2-S7.

Finally, we estimated SNP- h^2 from GWAS summary statistics for schizophrenia stratified by smoking status. Among schizophrenia smokers (3832 schizophrenia-cases, 8518 controls), the heritability of schizophrenia was estimated as 0.237 (95% CI:[0.169,0.304]) and among schizophrenia non-smokers (1423 schizophrenia-cases, 8518 controls) was 0.133 (95% CI:[0.017,0.249]). The estimated genetic correlation between these groups (schizophrenia smokers and schizophrenia non-smokers) was 0.860 (95% CI: [0.497,1.224]) and was significantly different from 0 ($P=3.52 \times 10^{-6}$) but not from 1. These results suggest that the genetic risk for schizophrenia is largely overlapping between smoking and non-smoking schizophrenia patients.

III.4 Genome-wide association of smoking behaviors among schizophrenia cases

Genomic inflation factors (λ) were 1.017 and 1.005 for smoking initiation ($n=5255$) and cigarettes-smoked-per-day ($n=3370$), respectively. The discovery GWAS did not yield SNP associations significant at established genome-wide criteria (5.0×10^{-8}). The strongest evidence of SNP-based association was observed for cigarettes-smoked-per-day, upstream of the *CBWD2* gene at chromosome 2q13 (rs1900325; $P=1.01 \times 10^{-7}$). Subsequent follow-up of suggestively associated SNPs ($P < 10^{-6}$) in an independent European-ancestry cohort ($n=1802$) yielded a

significant finding between cigarettes-smoked-per-day and rs148253479 upstream of the *TMEM106B* gene at 7p21.3 (Table 4, discovery $P=1\times 10^{-6}$, replication $P=0.011$, combined $P=3.18\times 10^{-8}$). Regional association and forest plots for top associations are provided in the accompanying supplemental information (Figures S7-8). Notably, none of the previously identified smoking behavior-associated SNPs were detected at genome-wide significant thresholds in our GWAS of smoking behaviors within schizophrenia cases, likely due in part to the limited power to detect small SNP effects in our modest sample size (Table S12).

IV. Phenotypic and polygenic associations between smoking behavior and schizophrenia symptom dimensions

We considered whether smoking patterns among schizophrenia cases and genetic risk factors for smoking were related to the clinical presentation of schizophrenia. For sex, age, and each clinical variable considered, Table 5 gives the estimated effect and significance from logistic or linear regression. Age-of-onset of schizophrenia was found to have a nominal association with smoking initiation ($P=0.018$) indicating higher rates of initiation in cases with earlier onset. The positive symptom factor score showed a positive association with smoking initiation ($P=3.21\times 10^{-5}$) and cigarettes-smoked-per-day ($P=0.015$). Depressive symptoms were also nominally associated with cigarettes-smoked-per-day ($P=0.014$) indicating that those with higher depression scores endorsed smoking more cigarettes. No significant phenotypic associations were found between smoking behaviors and the negative and mania factor scores.

We followed-up phenotypic associations by examining the relationship between symptom dimensions and TAG-based polygenic scores for smoking behaviors (Tables S8-S11, Figure S4). Both the TAG-smoking initiation and TAG-cigarettes-smoked-per-day scores were

associated with positive symptoms at nominal levels of significance ($P=0.023$, $P=0.006$ respectively).

Discussion

Despite conspicuous epidemiological and molecular genetic evidence supporting a link between smoking behavior and schizophrenia, the biological basis of this relationship is not well understood. Given the availability of subject-level clinical data from the PGC study of schizophrenia, we were able to characterize smoking patterns among >5000 schizophrenia cases. Using polygenic risk score methodology and genome-wide summary statistics, we not only provide confirmatory evidence of aggregate genetic effects contributing to both smoking initiation and risk of schizophrenia, but demonstrate also that risk factors influencing smoking initiation and quantity are at least partially shared between schizophrenia patients and the general population.

Of particular importance, we have successfully demonstrated shared genetic liabilities to schizophrenia and smoking behaviors in European populations. While polygenic scores based on TAG results for smoking initiation and cigarettes-smoked-per-day were both strongly associated with increased risk of schizophrenia, the association with cigarettes-smoked-per-day was largely driven by the 15q25 locus. Although the results support a polygenic overlap between smoking behavior in the general population and schizophrenia risk, we cannot definitively rule out the possibility that some identified schizophrenia genetic variants may be in fact indexing liability to smoking behavior (rather than having a pleiotropic effect on both traits) because of the high prevalence of nicotine use among affected persons. Future research is needed to disentangle this

confounded relationship by collecting smoking behavior information for both schizophrenia cases *and* control subjects.

Polygenic scores for smoking initiation also significantly predicted initiation among schizophrenia cases. Taken together with an estimated genetic correlation of ~ 0.624 , this suggests that genetic factors influencing smoking behavior are at least partially shared between schizophrenia and non-schizophrenia populations. We could rule out the possibility that they are entirely independent, but better powered studies are needed to more precisely estimate the degree of overlap. Similarly, polygenic scores for smoking quantity were also significantly predictive of smoking quantity among schizophrenia patients, albeit to a lesser degree of statistical significance.

By contrast, polygenic scores based on PGC results for schizophrenia were not predictive of smoking behavior among schizophrenia patients. Our results suggest a shared genetic liability to smoking behavior and schizophrenia, and that genetic liability to smoking is shared between the general population and schizophrenia patients, while liability to schizophrenia is *not* associated with smoking behavior among schizophrenia-affected individuals. The latter could be partially due to power and the restricted range of the smoking liability distribution among the selected schizophrenia population, as recent studies have found schizophrenia-PRS to be associated with smoking behavior in substance use enriched samples (Chen et al., 2016; Hartz et al., 2017).

Exploratory GWAS of smoking initiation and cigarettes-smoked-per-day among schizophrenia cases did not yield genome-wide significant evidence of association in the discovery stages. The top association was observed for cigarettes-smoked-per-day (rs1900325; $P=1.01 \times 10^{-7}$) was upstream of *CBWD2*, which has been previously implicated in sleep and

metabolic traits(Doherty et al., 2018; Hammerschlag et al., 2017). In the replication phase, a single genome-wide significant association was observed between cigarettes-smoked-per-day and SNPs upstream of the *TMEM106B* gene, a much studied risk locus for frontotemporal lobar degeneration (FTLD)(Van Deerlin et al., 2010) that encodes a trans-membrane protein involved in lysosomal trafficking and dendritic branching(Brady et al., 2013; Schwenk et al., 2014). In addition to FTLD, the *TMEM106B* gene has demonstrated associations with the clinical presentation of Alzheimer disease(Rutherford et al., 2012), the volume of left-sided temporal lobe and interhemispheric structures(Adams et al., 2014), and amphetamine response(Hart et al., 2012). Although not genome-wide significant, our results support an association between the *CHRNA3/CHRNA5* locus (rs16969968) and cigarettes-smoked-per-day among schizophrenia cases (replication $P=0.0001$, Table S12). Interestingly, analysis of schizophrenia stratified by case smoking status revealed elevated odds ratio (and higher allele frequencies) for this SNP among schizophrenia cases that have ever smoked (Figure S9). Also notable is the lack of genome-wide significant associations between TAG-associated variants and smoking behaviors among schizophrenia cases, which could reflect our limited power to detect individual SNP effects in our current sample size (<35%; Table S12).

Consistent with the literature, schizophrenia cases had elevated smoking rates and smoked more cigarettes per day than smokers in the general population (>30 cigarettes: 16.5 versus 6.9% respectively)(de Leon and Diaz, 2005; Jamal et al., 2015). Earlier age of schizophrenia onset was associated with higher rates of smoking initiation. When examining clinical features of schizophrenia, the positive symptom factor score was associated with smoking behavior indicating that those endorsing hallucinations and delusions were more likely to initiate smoking and smoke more cigarettes. This is broadly consistent with the self-

medication hypothesis, by itself does not tell against other etiological hypotheses, as it might represent a process by which symptoms might be reduced, irrespective of etiology.

Pharmacological upregulation of nicotinic acetylcholinergic transmission using either acetylcholinesterase inhibitors or positive allosteric modulators (PAMs) have been shown to ameliorate symptoms of schizophrenia (Wallace and Bertrand, 2015). The clinical dimensions for which the literature most strongly supports a role for such treatments are negative and cognitive symptoms (Singh et al., 2012). However, animal models also support a potential role in positive symptom-like features of ketamine-induced psychosis as well (Nikiforuk et al., 2016).

Additionally, a nominal association was found between cigarettes-smoked-per-day and the depressive symptom dimension adding support for the role of nicotine on mood in schizophrenia patients. This might be consistent with an improvement in mood concomitant with an amelioration of symptoms of the illness overall. It might also represent an inherent antidepressant effect of agonizing nicotinic transmission, as has been suggested by studies using the forced swim test in rodents (Marcus et al., 2016; Onajole et al., 2016; Shang et al., 2016; Zhang et al., 2016).

The major limitation of this study was the number of available schizophrenia cases with detailed clinical and smoking-related data. Despite attaining a modest sample size for smoking analyses within schizophrenia cases, our power was limited to detect single-SNP associations with smoking initiation or cigarettes-smoked-per-day (Table S12). Another limitation was the use of self-report data to index smoking behavior among schizophrenia cases. Future research should incorporate objective nicotine metabolite biomarkers such as cotinine levels. The age of schizophrenia onset was determined retrospectively in some cases and since the duration of untreated psychosis can vary, the precision of the true onset is unknown. Also, because smoking

data on control subjects was not available for the majority of participating studies, we were limited in our ability to relate findings from the smoking analyses within schizophrenia cases to variation in the general population. Recently, a large-scale GWAS from the GSCAN consortium reported 55, 378, and 99 associated genetic variants with cigarettes-smoked-per-day, smoking initiation, and alcohol drinks per week respectively(Liu et al., 2019). Forthcoming research needs to examine shared genetic risk of schizophrenia across substances as well as gender and diverse populations. As available sample sizes and phenotypic data grow(Pardiñas et al., 2018; Schizophrenia Working Group of the Psychiatric Genomics Consortium et al., 2020) it will be possible to apply alternative methods such as Mendelian Randomization to assess causal processes between these phenotypes. Nonetheless, findings suggest a portion of the schizophrenia-smoking association is due to shared genetic etiology, as we were able to demonstrate partial overlap between genetic liability to smoking behavior in the general population and (1) schizophrenia risk, and (2) smoking behavior among schizophrenia patients. In addition to supporting genome-wide pleiotropic effects, our smoking GWAS within schizophrenia cases highlighted a schizophrenia-specific genetic liability for smoking quantity. Future research needs to address mechanisms underlying associations between these traits (e.g., Mendelian randomization, pharmacogenetics) to aid both schizophrenia and smoking treatment and prevention efforts.

Acknowledgements

Core funding for the Psychiatric Genomics Consortium is from the US National Institute of Mental Health (U01 MH094421). The work specific to this report was funded by the United States Department of Veterans Affairs Merit Review Program (5I01CX000278) to Ayman H. Fanous. Roseann E. Peterson is supported by National Institutes of Health (NIH) K01 grant MH113848 and The Brain & Behavior Research Foundation NARSAD grant 28632 P&S Fund. This project has received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement 279227 to Marcella Rietschel, and Research Council of Norway (grant #223273). Statistical analyses were carried out on the Genetic Cluster Computer (<http://www.geneticcluster.org>) which is financially supported by the Netherlands Scientific Organization (NWO 480-05-003) along with a supplement from the Dutch Brain Foundation and the VU University Amsterdam.

Disclosures

Q.S.L. is an employee of Janssen Research and Development, LLC. The other authors declare no conflict of interest.

Contributions

Authors R. Peterson, T. Bigdeli, and A. Fanous designed the study and wrote the first draft of the manuscript. R. Peterson, T. Bigdeli, and S. Ripke undertook the statistical analysis. All authors contributed to and have approved the final manuscript.

Tables

Table 1. Conceptual overview of analyses of schizophrenia and smoking behaviors.


Research Question	Cohorts & Sample Sizes	Analysis
Question 1: Are there genetic correlations between schizophrenia and smoking behaviors?	<i>Primary</i> - PGC-Schizophrenia European ancestry: 35,476 cases, 46,839 controls; TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181, cessation 41,278, age of smoking onset 24,114	genetic correlation (LD score regression)
	<i>Replication</i> - PGC-Schizophrenia East-Asian ancestry: 1,836 cases, 3,383 controls	trans-ethnic genetic correlation (popcorn)
Question 2: Do polygenic risk scores for smoking behaviors also predict schizophrenia case status?	<i>Training Set</i> - TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181, cessation 41,278, age of smoking onset 24,114	polygenic risk scores (cross-trait association)
	<i>Testing Set</i> - PGC-Schizophrenia: 35,476 cases, 46,839 controls	
Question 3: What is the genetic architecture of smoking behavior among schizophrenia patients?		
3.1 What is the SNP-based heritability of smoking behaviors among schizophrenia cases?	PGC-Schizophrenia Phenotype Working Group - 10 study sites: smoking initiation 5,255, cigarettes-per-day 3,370	SNP-based heritability (LD score regression)
3.2 Are genetic factors for smoking behaviors shared between populations with and without schizophrenia?	PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370; TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181	genetic correlation (LD score regression)
3.3 Do polygenic risk scores for smoking behaviors also predict these behaviors in schizophrenia patients?	<i>Training Set</i> - TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181	polygenic risk scores (within-trait across-cohort association)
	<i>Testing Set</i> - PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370	
3.4 Are there schizophrenia-specific genetic risk variants for smoking behaviors?	<i>Primary</i> - PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370	genome-wide association study meta-analysis of smoking behaviors among schizophrenia cases
	<i>Replication</i> - Janssen Pharmaceuticals: smoking initiation 1802, cigarettes-per-day 1802	
Question 4: Are there associations between smoking behaviors and clinical features of schizophrenia?		
4.1 Are smoking behaviors among schizophrenia cases associated with clinical presentation of schizophrenia?	PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370, age-schizophrenia-onset 4,658; symptom dimension factor scores: positive 3,846, negative 3,845, manic 3,740, depression 3,740	phenotypic associations (linear/logistic regression)
4.2 Do polygenic risk scores for smoking behaviors predict schizophrenia symptom dimensions?	<i>Training Set</i> - TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181	polygenic risk scores (across-trait across-cohort association)
	<i>Testing Set</i> - PGC-Schizophrenia: age-schizophrenia-onset 11,600; symptom dimension factor scores: positive 8,330, negative 8,427, manic 6,965, depression 6,964	

PGC is Psychiatric Genomics Consortium, TAG is Tobacco and Genetics consortium, LD is linkage disequilibrium, SNP is single nucleotide polymorphism.

*Table 2. Sample characteristics for each PGC2-schizophrenia cohort.**Table 3. Genetic correlations between TAG and PGC-schizophrenia phenotypes.**Table 4. Association results for top SNP associations.**Table 5. Association of smoking variables with clinical features in schizophrenia.*

Figure legends

Figure 1. Association of TAG-based polygenic scores with schizophrenia risk and smoking behaviors. For polygenic scores based on analyses of smoking behaviors described by TAG, the variance explained for selected outcomes in PGC-schizophrenia is shown on the y-axis, in terms of Nagelkerke's pseudo- R^2 (schizophrenia and smoking initiation) or R^2 (cigarettes-smoked-per-day); scores based on varying SNP P -value inclusion thresholds are displayed as colored bars. *logOnset* is log-transformed age-of-onset (see Methods).



References

- 1000 Genomes Project Consortium, Abecasis, G.R., Auton, A., Brooks, L.D., DePristo, M.A., Durbin, R.M., Handsaker, R.E., Kang, H.M., Marth, G.T., McVean, G.A., 2012. An integrated map of genetic variation from 1,092 human genomes. *Nature* 491, 56–65.
- Adams, H.H.H., Verhaaren, B.F.J., Vrooman, H.A., Uitterlinden, A.G., Hofman, A., van Duijn, C.M., van der Lugt, A., Niessen, W.J., Vernooij, M.W., Ikram, M.A., 2014. TMEM106B influences volume of left-sided temporal lobe and interhemispheric structures in the general population. *Biol. Psychiatry* 76, 503–508.
- Brady, O.A., Zheng, Y., Murphy, K., Huang, M., Hu, F., 2013. The frontotemporal lobar degeneration risk factor, TMEM106B, regulates lysosomal morphology and function. *Hum. Mol. Genet.* 22, 685–695.
- Brainstorm Consortium, Anttila, V., Bulik-Sullivan, B., Finucane, H.K., Walters, R.K., Bras, J., Duncan, L., Escott-Price, V., Falcone, G.J., Gormley, P., Malik, R., Patsopoulos, N.A., Ripke, S., Wei, Z., Yu, D., Lee, P.H., Turley, P., Grenier-Boley, B., Chouraki, V., Kamatani, Y., Berr, C., Letenneur, L., Hannequin, D., Amouyel, P., Boland, A., Deleuze, J.-F., Duron, E., Vardarajan, B.N., Reitz, C., Goate, A.M., Huentelman, M.J., Kamboh, M.I., Larson, E.B., Rogaeva, E., St George-Hyslop, P., Hakonarson, H., Kukull, W.A., Farrer, L.A., Barnes, L.L., Beach, T.G., Demirci, F.Y., Head, E., Hulette, C.M., Jicha, G.A., Kauwe, J.S.K., Kaye, J.A., Leverenz, J.B., Levey, A.I., Lieberman, A.P., Pankratz, V.S., Poon, W.W., Quinn, J.F., Saykin, A.J., Schneider, L.S., Smith, A.G., Sonnen, J.A., Stern, R.A., Van Deerlin, V.M., Van Eldik, L.J., Harold, D., Russo, G., Rubinsztein, D.C., Bayer, A., Tsolaki, M., Proitsi, P., Fox, N.C., Hampel, H., Owen, M.J., Mead, S., Passmore, P., Morgan, K., Nöthen, M.M., Rossor, M., Lupton, M.K., Hoffmann, P., Kornhuber, J., Lawlor, B., McQuillin, A., Al-Chalabi, A., Bis, J.C., Ruiz, A., Boada, M., Seshadri, S., Beiser, A., Rice, K., van der Lee, S.J., De Jager, P.L., Geschwind, D.H., Riemenschneider, M., Riedel-Heller, S., Rotter, J.I., Ransmayr, G., Hyman, B.T., Cruchaga, C., Alegret, M., Winsvold, B., Palta, P., Farh, K.-H., Cuenca-Leon, E., Furlotte, N., Kurth, T., Ligthart, L., Terwindt, G.M., Freilinger, T., Ran, C., Gordon, S.D., Borck, G., Adams, H.H.H., Lehtimäki, T., Wedenoja, J., Buring, J.E., Schürks, M., Hrafnisdottir, M., Hottenga, J.-J., Penninx, B., Artto, V., Kaunisto, M., Vepsäläinen, S., Martin, N.G., Montgomery, G.W., Kurki, M.I., Hämäläinen, E., Huang, H., Huang, J., Sandor, C., Webber, C., Muller-Myhsok, B., Schreiber, S., Salomaa, V., Loehrer, E., Göbel, H., Macaya, A., Pozo-Rosich, P., Hansen, T., Werge, T., Kaprio, J., Metspalu, A., Kubisch, C., Ferrari, M.D., Belin, A.C., van den Maagdenberg, A.M.J.M., Zwart, J.-A., Boomsma, D., Eriksson, N., Olesen, J., Chasman, D.I., Nyholt, D.R., Avbersek, A., Baum, L., Berkovic, S., Bradfield, J., Buono, R.J., Catarino, C.B., Cossette, P., De Jonghe, P., Depondt, C., Dlugos, D., Ferraro, T.N., French, J., Hjalgrim, H., Jamnadas-Khoda, J., Kälviäinen, R., Kunz, W.S., Lerche, H., Leu, C., Lindhout, D., Lo, W., Lowenstein, D., McCormack, M., Møller, R.S., Molloy, A., Ng, P.-W., Oliver, K., Privitera, M., Radtke, R., Ruppert, A.-K., Sander, T., Schachter, S., Schankin, C., Scheffer, I., Schoch, S., Sisodiya, S.M., Smith, P., Sperling, M., Striano, P., Surges, R., Thomas, G.N., Visscher, F., Whelan, C.D., Zara, F., Heinzen, E.L., Marson, A., Becker, F., Stroink, H., Zimprich, F., Gasser, T., Gibbs, R., Heutink, P., Martinez, M., Morris, H.R., Sharma, M., Ryten, M., Mok, K.Y., Pulit, S., Bevan, S., Holliday, E., Attia, J., Battey, T., Boncoraglio, G., Thijs, V., Chen, W.-M., Mitchell, B., Rothwell, P., Sharma, P.,

Sudlow, C., Vicente, A., Markus, H., Kourkoulis, C., Pera, J., Raffeld, M., Silliman, S., Boraska Perica, V., Thornton, L.M., Huckins, L.M., William Rayner, N., Lewis, C.M., Gratacos, M., Rybakowski, F., Keski-Rahkonen, A., Raevuori, A., Hudson, J.I., Reichborn-Kjennerud, T., Monteleone, P., Karwautz, A., Mannik, K., Baker, J.H., O'Toole, J.K., Trace, S.E., Davis, O.S.P., Helder, S.G., Ehrlich, S., Herpertz-Dahlmann, B., Danner, U.N., van Elburg, A.A., Clementi, M., Forzan, M., Docampo, E., Lissowska, J., Hauser, J., Tortorella, A., Maj, M., Gonidakis, F., Tziouvas, K., Papezova, H., Yilmaz, Z., Wagner, G., Cohen-Woods, S., Herms, S., Julià, A., Rabionet, R., Dick, D.M., Ripatti, S., Andreassen, O.A., Espeseth, T., Lundervold, A.J., Steen, V.M., Pinto, D., Scherer, S.W., Aschauer, H., Schosser, A., Alfredsson, L., Padyukov, L., Halmi, K.A., Mitchell, J., Strober, M., Bergen, A.W., Kaye, W., Szatkiewicz, J.P., Cormand, B., Ramos-Quiroga, J.A., Sánchez-Mora, C., Ribasés, M., Casas, M., Hervas, A., Arranz, M.J., Haavik, J., Zayats, T., Johansson, S., Williams, N., Dempfle, A., Rothenberger, A., Kuntsi, J., Oades, R.D., Banaschewski, T., Franke, B., Buitelaar, J.K., Arias Vasquez, A., Doyle, A.E., Reif, A., Lesch, K.-P., Freitag, C., Rivero, O., Palmason, H., Romanos, M., Langley, K., Rietschel, M., Witt, S.H., Dalsgaard, S., Børglum, A.D., Waldman, I., Wilmot, B., Molly, N., Bau, C.H.D., Crosbie, J., Schachar, R., Loo, S.K., McGough, J.J., Grevet, E.H., Medland, S.E., Robinson, E., Weiss, L.A., Bacchelli, E., Bailey, A., Bal, V., Battaglia, A., Betancur, C., Bolton, P., Cantor, R., Celestino-Soper, P., Dawson, G., De Rubeis, S., Duque, F., Green, A., Klauck, S.M., Leboyer, M., Levitt, P., Maestrini, E., Mane, S., De-Luca, D.M., Parr, J., Regan, R., Reichenberg, A., Sandin, S., Vorstman, J., Wassink, T., Wijsman, E., Cook, E., Santangelo, S., Delorme, R., Rogé, B., Magalhaes, T., Arking, D., Schulze, T.G., Thompson, R.C., Strohmaier, J., Matthews, K., Melle, I., Morris, D., Blackwood, D., McIntosh, A., Bergen, S.E., Schalling, M., Jamain, S., Maaser, A., Fischer, S.B., Reinbold, C.S., Fullerton, J.M., Guzman-Parra, J., Mayoral, F., Schofield, P.R., Cichon, S., Mühleisen, T.W., Degenhardt, F., Schumacher, J., Bauer, M., Mitchell, P.B., Gershon, E.S., Rice, J., Potash, J.B., Zandi, P.P., Craddock, N., Ferrier, I.N., Alda, M., Rouleau, G.A., Turecki, G., Ophoff, R., Pato, C., Anjorin, A., Stahl, E., Leber, M., Czerski, P.M., Cruceanu, C., Jones, I.R., Posthuma, D., Andlauer, T.F.M., Forstner, A.J., Streit, F., Baune, B.T., Air, T., Sinnamon, G., Wray, N.R., MacIntyre, D.J., Porteous, D., Homuth, G., Rivera, M., Grove, J., Middeldorp, C.M., Hickie, I., Pergadia, M., Mehta, D., Smit, J.H., Jansen, R., de Geus, E., Dunn, E., Li, Q.S., Nauck, M., Schoevers, R.A., Beekman, A.T., Knowles, J.A., Viktorin, A., Arnold, P., Barr, C.L., Bedoya-Berrio, G., Bienvenu, O.J., Brentani, H., Burton, C., Camarena, B., Cappi, C., Cath, D., Cavallini, M., Cusi, D., Darrow, S., Denys, D., Derks, E.M., Dietrich, A., Fernandez, T., Figuee, M., Freimer, N., Gerber, G., Grados, M., Greenberg, E., Hanna, G.L., Hartmann, A., Hirschtritt, M.E., Hoekstra, P.J., Huang, A., Huyser, C., Illmann, C., Jenike, M., Kuperman, S., Leventhal, B., Lochner, C., Lyon, G.J., Macciardi, F., Madruga-Garrido, M., Malaty, I.A., Maras, A., McGrath, L., Miguel, E.C., Mir, P., Nestadt, G., Nicolini, H., Okun, M.S., Pakstis, A., Paschou, P., Piacentini, J., Pittenger, C., Plessen, K., Ramensky, V., Ramos, E.M., Reus, V., Richter, M.A., Riddle, M.A., Robertson, M.M., Roessner, V., Rosário, M., Samuels, J.F., Sandor, P., Stein, D.J., Tsetsos, F., Van Nieuwerburgh, F., Weatherall, S., Wendland, J.R., Wolanczyk, T., Worbe, Y., Zai, G., Goes, F.S., McLaughlin, N., Nestadt, P.S., Grabe, H.-J., Depienne, C., Konkashbaev, A., Lanzagorta, N., Valencia-Duarte, A., Bramon, E., Buccola, N., Cahn, W., Cairns, M., Chong, S.A., Cohen, D., Crespo-Facorro, B., Crowley, J., Davidson, M., DeLisi, L., Dinan, T., Donohoe,

- G., Drapeau, E., Duan, J., Haan, L., Hougaard, D., Karachanak-Yankova, S., Khrunin, A., Klovins, J., Kučinskis, V., Lee Chee Keong, J., Limborska, S., Loughland, C., Lönnqvist, J., Maher, B., Mattheisen, M., McDonald, C., Murphy, K.C., Nenadic, I., van Os, J., Pantelis, C., Pato, M., Petryshen, T., Quested, D., Roussos, P., Sanders, A.R., Schall, U., Schwab, S.G., Sim, K., So, H.-C., Stögmänn, E., Subramaniam, M., Toncheva, D., Waddington, J., Walters, J., Weiser, M., Cheng, W., Cloninger, R., Curtis, D., Gejman, P.V., Henskens, F., Mattingsdal, M., Oh, S.-Y., Scott, R., Webb, B., Breen, G., Churchhouse, C., Bulik, C.M., Daly, M., Dichgans, M., Faraone, S.V., Guerreiro, R., Holmans, P., Kendler, K.S., Koeleman, B., Mathews, C.A., Price, A., Scharf, J., Sklar, P., Williams, J., Wood, N.W., Cotsapas, C., Palotie, A., Smoller, J.W., Sullivan, P., Rosand, J., Corvin, A., Neale, B.M., Schott, J.M., Anney, R., Elia, J., Grigoriu-Serbanescu, M., Edenberg, H.J., Murray, R., 2018. Analysis of shared heritability in common disorders of the brain. *Science* 360. <https://doi.org/10.1126/science.aap8757>
- Brown, B.C., Asian Genetic Epidemiology Network Type 2 Diabetes Consortium, Ye, C.J., Price, A.L., Zaitlen, N., 2016. Transethnic Genetic-Correlation Estimates from Summary Statistics. *Am. J. Hum. Genet.* 99, 76–88.
- Bulik-Sullivan, B., Finucane, H.K., Anttila, V., Gusev, A., Day, F.R., Loh, P.-R., ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, Duncan, L., Perry, J.R.B., Patterson, N., Robinson, E.B., Daly, M.J., Price, A.L., Neale, B.M., 2015. An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* 47, 1236–1241.
- Bulik-Sullivan, B.K., Loh, P.-R., Finucane, H.K., Ripke, S., Yang, J., Schizophrenia Working Group of the Psychiatric Genomics Consortium, Patterson, N., Daly, M.J., Price, A.L., Neale, B.M., 2015. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* 47, 291–295.
- Centers for Disease Control and Prevention (CDC), 2013. Vital signs: current cigarette smoking among adults aged ≥ 18 years with mental illness - United States, 2009-2011. *MMWR Morb. Mortal. Wkly. Rep.* 62, 81–87.
- Chen, J., Bacanu, S.-A., Yu, H., Zhao, Z., Jia, P., Kendler, K.S., Kranzler, H.R., Gelernter, J., Farrer, L., Minica, C., Pool, R., Milaneschi, Y., Boomsma, D.I., Penninx, B.W.J.H., Tyndale, R.F., Ware, J.J., Vink, J.M., Kaprio, J., Munafò, M., Chen, X., Cotinine meta-analysis group, FTND meta-analysis group, 2016. Genetic Relationship between Schizophrenia and Nicotine Dependence. *Sci. Rep.* 6, 25671.
- de Leon, J., Diaz, F.J., 2005. A meta-analysis of worldwide studies demonstrates an association between schizophrenia and tobacco smoking behaviors. *Schizophr. Res.* 76, 135–157.
- Doherty, A., Smith-Byrne, K., Ferreira, T., Holmes, M.V., Holmes, C., Pulit, S.L., Lindgren, C.M., 2018. GWAS identifies 14 loci for device-measured physical activity and sleep duration. *Nat. Commun.* 9, 5257.
- Freedman, R., 2014. $\alpha 7$ -nicotinic acetylcholine receptor agonists for cognitive enhancement in schizophrenia. *Annu. Rev. Med.* 65, 245–261.
- Gage, S.H., Jones, H.J., Taylor, A.E., Burgess, S., Zammit, S., Munafò, M.R., 2017. Investigating causality in associations between smoking initiation and schizophrenia using Mendelian randomization. *Sci. Rep.* 7, 40653.
- Gage, S.H., Munafò, M.R., 2015. Rethinking the association between smoking and schizophrenia. *Lancet Psychiatry* 2, 118–119.

- Guillozet-Bongaarts, A.L., Hyde, T.M., Dalley, R.A., Hawrylycz, M.J., Henry, A., Hof, P.R., Hohmann, J., Jones, A.R., Kuan, C.L., Royall, J., Shen, E., Swanson, B., Zeng, H., Kleinman, J.E., 2014. Altered gene expression in the dorsolateral prefrontal cortex of individuals with schizophrenia. *Mol. Psychiatry* 19, 478–485.
- Gurillo, P., Jauhar, S., Murray, R.M., MacCabe, J.H., 2015. Does tobacco use cause psychosis? Systematic review and meta-analysis. *Lancet Psychiatry* 2, 718–725.
- Hammerschlag, A.R., Stringer, S., de Leeuw, C.A., Sniekers, S., Taskesen, E., Watanabe, K., Blanken, T.F., Dekker, K., Te Lindert, B.H.W., Wassing, R., Jonsdottir, I., Thorleifsson, G., Stefansson, H., Gislason, T., Berger, K., Schormair, B., Wellmann, J., Winkelmann, J., Stefansson, K., Oexle, K., Van Someren, E.J.W., Posthuma, D., 2017. Genome-wide association analysis of insomnia complaints identifies risk genes and genetic overlap with psychiatric and metabolic traits. *Nat. Genet.* 49, 1584–1592.
- Hart, A.B., Engelhardt, B.E., Wardle, M.C., Sokoloff, G., Stephens, M., de Wit, H., Palmer, A.A., 2012. Genome-wide association study of d-amphetamine response in healthy volunteers identifies putative associations, including cadherin 13 (CDH13). *PLoS One* 7, e42646.
- Hartz, S.M., Horton, A.C., Hancock, D.B., Baker, T.B., Caporaso, N.E., Chen, L.-S., Hokanson, J.E., Lutz, S.M., Marazita, M.L., McNeil, D.W., Pato, C.N., Pato, M.T., Johnson, E.O., Bierut, L.J., 2018. Genetic correlation between smoking behaviors and schizophrenia. *Schizophr. Res.* 194, 86–90.
- Hartz, S.M., Horton, A.C., Oehlert, M., Carey, C.E., Agrawal, A., Bogdan, R., Chen, L.-S., Hancock, D.B., Johnson, E.O., Pato, C.N., Pato, M.T., Rice, J.P., Bierut, L.J., 2017. Association Between Substance Use Disorder and Polygenic Liability to Schizophrenia. *Biol. Psychiatry* 82, 709–715.
- Hartz, S.M., Pato, C.N., Medeiros, H., Cavazos-Rehg, P., Sobell, J.L., Knowles, J.A., Bierut, L.J., Pato, M.T., Genomic Psychiatry Cohort Consortium, 2014. Comorbidity of severe psychotic disorders with measures of substance use. *JAMA Psychiatry* 71, 248–254.
- Howes, O.D., Kapur, S., 2009. The dopamine hypothesis of schizophrenia: version III--the final common pathway. *Schizophr. Bull.* 35, 549–562.
- Howie, B., Marchini, J., Stephens, M., 2011. Genotype imputation with thousands of genomes. *G3* 1, 457–470.
- Howie, B.N., Donnelly, P., Marchini, J., 2009. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet.* 5, e1000529.
- International Schizophrenia Consortium, Purcell, S.M., Wray, N.R., Stone, J.L., Visscher, P.M., O'Donovan, M.C., Sullivan, P.F., Sklar, P., 2009. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460, 748–752.
- Jamal, A., Homa, D.M., O'Connor, E., Babb, S.D., Caraballo, R.S., Singh, T., Hu, S.S., King, B.A., 2015. Current cigarette smoking among adults - United States, 2005-2014. *MMWR Morb. Mortal. Wkly. Rep.* 64, 1233–1240.
- Kendler, K.S., Lönn, S.L., Sundquist, J., Sundquist, K., 2015. Smoking and schizophrenia in population cohorts of Swedish women and men: a prospective co-relative control study. *Am. J. Psychiatry* 172, 1092–1100.
- Kumari, V., Postma, P., 2005. Nicotine use in schizophrenia: the self medication hypotheses. *Neurosci. Biobehav. Rev.* 29, 1021–1034.

- Li, Q., Wineinger, N.E., Fu, D.-J., Libiger, O., Alphas, L., Savitz, A., Gopal, S., Cohen, N., Schork, N.J., 2017. Genome-wide association study of paliperidone efficacy. *Pharmacogenet. Genomics* 27, 7–18.
- Liu, M., Jiang, Y., Wedow, R., Li, Y., Brazel, D.M., Chen, F., Datta, G., Davila-Velderrain, J., McGuire, D., Tian, C., Zhan, X., 23andMe Research Team, HUNT All-In Psychiatry, Choquet, H., Docherty, A.R., Faul, J.D., Foerster, J.R., Fritsche, L.G., Gabrielsen, M.E., Gordon, S.D., Haessler, J., Hottenga, J.-J., Huang, H., Jang, S.-K., Jansen, P.R., Ling, Y., Mägi, R., Matoba, N., McMahon, G., Mulas, A., Orrù, V., Palviainen, T., Pandit, A., Reginsson, G.W., Skogholt, A.H., Smith, J.A., Taylor, A.E., Turman, C., Willemsen, G., Young, H., Young, K.A., Zajac, G.J.M., Zhao, W., Zhou, W., Bjornsdottir, G., Boardman, J.D., Boehnke, M., Boomsma, D.I., Chen, C., Cucca, F., Davies, G.E., Eaton, C.B., Ehringer, M.A., Esko, T., Fiorillo, E., Gillespie, N.A., Gudbjartsson, D.F., Haller, T., Harris, K.M., Heath, A.C., Hewitt, J.K., Hickie, I.B., Hokanson, J.E., Hopfer, C.J., Hunter, D.J., Iacono, W.G., Johnson, E.O., Kamatani, Y., Kardia, S.L.R., Keller, M.C., Kellis, M., Kooperberg, C., Kraft, P., Krauter, K.S., Laakso, M., Lind, P.A., Loukola, A., Lutz, S.M., Madden, P.A.F., Martin, N.G., McGue, M., McQueen, M.B., Medland, S.E., Metspalu, A., Mohlke, K.L., Nielsen, J.B., Okada, Y., Peters, U., Polderman, T.J.C., Posthuma, D., Reiner, A.P., Rice, J.P., Rimm, E., Rose, R.J., Runarsdottir, V., Stallings, M.C., Stančáková, A., Stefansson, H., Thai, K.K., Tindle, H.A., Tyrfinngsson, T., Wall, T.L., Weir, D.R., Weisner, C., Whitfield, J.B., Winsvold, B.S., Yin, J., Zuccolo, L., Bierut, L.J., Hveem, K., Lee, J.J., Munafò, M.R., Saccone, N.L., Willer, C.J., Cornelis, M.C., David, S.P., Hinds, D.A., Jorgenson, E., Kaprio, J., Stitzel, J.A., Stefansson, K., Thorgeirsson, T.E., Abecasis, G., Liu, D.J., Vrieze, S., 2019. Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. *Nat. Genet.* 51, 237–244.
- Maes, H.H., Sullivan, P.F., Bulik, C.M., Neale, M.C., Prescott, C.A., Eaves, L.J., Kendler, K.S., 2004. A twin study of genetic and environmental influences on tobacco initiation, regular tobacco use and nicotine dependence. *Psychol. Med.* 34, 1251–1261.
- Marcus, M.M., Björkholm, C., Malmerfelt, A., Möller, A., Pahlsson, N., Konradsson-Geuken, Å., Feltmann, K., Jardemark, K., Schilström, B., Svensson, T.H., 2016. Alpha7 nicotinic acetylcholine receptor agonists and PAMs as adjunctive treatment in schizophrenia. An experimental study. *Eur. Neuropsychopharmacol.* 26, 1401–1411.
- McGrath, J., Saha, S., Chant, D., Welham, J., 2008. Schizophrenia: a concise overview of incidence, prevalence, and mortality. *Epidemiol. Rev.* 30, 67–76.
- Metspalu, A., Köhler, F., Laschinski, G., Ganten, D., Roots, I., 2004. [The Estonian Genome Project in the context of European genome research]. *Dtsch. Med. Wochenschr.* 129 Suppl 1, S25–8.
- Nikiforuk, A., Potasiewicz, A., Kos, T., Popik, P., 2016. The combination of memantine and galantamine improves cognition in rats: The synergistic role of the $\alpha 7$ nicotinic acetylcholine and NMDA receptors. *Behav. Brain Res.* 313, 214–218.
- Olfson, M., Gerhard, T., Huang, C., Crystal, S., Stroup, T.S., 2015. Premature Mortality Among Adults With Schizophrenia in the United States. *JAMA Psychiatry* 72, 1172–1181.
- Onajole, O.K., Vallerini, G.P., Eaton, J.B., Lukas, R.J., Brunner, D., Caldarone, B.J., Kozikowski, A.P., 2016. Synthesis and Behavioral Studies of Chiral Cyclopropanes as Selective $\alpha 4\beta 2$ -Nicotinic Acetylcholine Receptor Partial Agonists Exhibiting an

- Antidepressant Profile. Part III. ACS Chem. Neurosci. 7, 811–822.
- Pardiñas, A.F., Holmans, P., Pocklington, A.J., Escott-Price, V., Ripke, S., Carrera, N., Legge, S.E., Bishop, S., Cameron, D., Hamshere, M.L., Han, J., Hubbard, L., Lynham, A., Mantripragada, K., Rees, E., MacCabe, J.H., McCarroll, S.A., Baune, B.T., Breen, G., Byrne, E.M., Dannlowski, U., Eley, T.C., Hayward, C., Martin, N.G., McIntosh, A.M., Plomin, R., Porteous, D.J., Wray, N.R., Caballero, A., Geschwind, D.H., Huckins, L.M., Ruderfer, D.M., Santiago, E., Sklar, P., Stahl, E.A., Won, H., Agerbo, E., Als, T.D., Andreassen, O.A., Bækvad-Hansen, M., Mortensen, P.B., Pedersen, C.B., Børglum, A.D., Bybjerg-Grauholm, J., Djurovic, S., Durmishi, N., Pedersen, M.G., Golimbet, V., Grove, J., Hougaard, D.M., Mattheisen, M., Molden, E., Mors, O., Nordentoft, M., Pejovic-Milovancevic, M., Sigurdsson, E., Silagadze, T., Hansen, C.S., Stefansson, K., Stefansson, H., Steinberg, S., Tosato, S., Werge, T., GERAD1 Consortium, CRESTAR Consortium, Collier, D.A., Rujescu, D., Kirov, G., Owen, M.J., O'Donovan, M.C., Walters, J.T.R., 2018. Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nat. Genet.* 50, 381–389.
- Parikh, V., Kutlu, M.G., Gould, T.J., 2016. nAChR dysfunction as a common substrate for schizophrenia and comorbid nicotine addiction: Current trends and perspectives. *Schizophr. Res.* 171, 1–15.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J., Sham, P.C., 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81, 559–575.
- Ripke, S., O'Dushlaine, C., Chambert, K., Moran, J.L., Kähler, A.K., Akterin, S., Bergen, S.E., Collins, A.L., Crowley, J.J., Fromer, M., Kim, Y., Lee, S.H., Magnusson, P.K.E., Sanchez, N., Stahl, E.A., Williams, S., Wray, N.R., Xia, K., Bettella, F., Børglum, A.D., Bulik-Sullivan, B.K., Cormican, P., Craddock, N., de Leeuw, C., Durmishi, N., Gill, M., Golimbet, V., Hamshere, M.L., Holmans, P., Hougaard, D.M., Kendler, K.S., Lin, K., Morris, D.W., Mors, O., Mortensen, P.B., Neale, B.M., O'Neill, F.A., Owen, M.J., Milovancevic, M.P., Posthuma, D., Powell, J., Richards, A.L., Riley, B.P., Ruderfer, D., Rujescu, D., Sigurdsson, E., Silagadze, T., Smit, A.B., Stefansson, H., Steinberg, S., Suvisaari, J., Tosato, S., Verhage, M., Walters, J.T., Multicenter Genetic Studies of Schizophrenia Consortium, Levinson, D.F., Gejman, P.V., Kendler, K.S., Laurent, C., Mowry, B.J., O'Donovan, M.C., Owen, M.J., Pulver, A.E., Riley, B.P., Schwab, S.G., Wildenauer, D.B., Dudbridge, F., Holmans, P., Shi, J., Albus, M., Alexander, M., Campion, D., Cohen, D., Dikeos, D., Duan, J., Eichhammer, P., Godard, S., Hansen, M., Lerer, F.B., Liang, K.-Y., Maier, W., Mallet, J., Nertney, D.A., Nestadt, G., Norton, N., O'Neill, F.A., Papadimitriou, G.N., Ribble, R., Sanders, A.R., Silverman, J.M., Walsh, D., Williams, N.M., Wormley, B., Psychosis Endophenotypes International Consortium, Arranz, M.J., Bakker, S., Bender, S., Bramon, E., Collier, D., Crespo-Facorro, B., Hall, J., Iyegbe, C., Jablensky, A., Kahn, R.S., Kalaydjieva, L., Lawrie, S., Lewis, C.M., Lin, K., Linszen, D.H., Mata, I., McIntosh, A., Murray, R.M., Ophoff, R.A., Powell, J., Rujescu, D., Van Os, J., Walshe, M., Weisbrod, M., Wiersma, D., Wellcome Trust Case Control Consortium 2, Donnelly, P., Barroso, I., Blackwell, J.M., Bramon, E., Brown, M.A., Casas, J.P., Corvin, A.P., Deloukas, P., Duncanson, A., Jankowski, J., Markus, H.S., Mathew, C.G., Palmer, C.N.A., Plomin, R., Rautanen, A., Sawcer, S.J., Trembath, R.C., Viswanathan, A.C., Wood,

- N.W., Spencer, C.C.A., Band, G., Bellenguez, C., Freeman, C., Hellenthal, G., Giannoulatou, E., Pirinen, M., Pearson, R.D., Strange, A., Su, Z., Vukcevic, D., Donnelly, P., Langford, C., Hunt, S.E., Edkins, S., Gwilliam, R., Blackburn, H., Bumpstead, S.J., Dronov, S., Gillman, M., Gray, E., Hammond, N., Jayakumar, A., McCann, O.T., Liddle, J., Potter, S.C., Ravindrarajah, R., Ricketts, M., Tashakkori-Ghanbaria, A., Waller, M.J., Weston, P., Widaa, S., Whittaker, P., Barroso, I., Deloukas, P., Mathew, C.G., Blackwell, J.M., Brown, M.A., Corvin, A.P., McCarthy, M.I., Spencer, C.C.A., Bramon, E., Corvin, A.P., O'Donovan, M.C., Stefansson, K., Scolnick, E., Purcell, S., McCarroll, S.A., Sklar, P., Hultman, C.M., Sullivan, P.F., 2013. Genome-wide association analysis identifies 13 new risk loci for schizophrenia. *Nat. Genet.* 45, 1150–1159.
- Ruderfer, D.M., Fanous, A.H., Ripke, S., McQuillin, A., Amdur, R.L., Schizophrenia Working Group of the Psychiatric Genomics Consortium, Bipolar Disorder Working Group of the Psychiatric Genomics Consortium, Cross-Disorder Working Group of the Psychiatric Genomics Consortium, Gejman, P.V., O'Donovan, M.C., Andreassen, O.A., Djurovic, S., Hultman, C.M., Kelsoe, J.R., Jamain, S., Landén, M., Leboyer, M., Nimgaonkar, V., Nurnberger, J., Smoller, J.W., Craddock, N., Corvin, A., Sullivan, P.F., Holmans, P., Sklar, P., Kendler, K.S., 2014. Polygenic dissection of diagnosis and clinical dimensions of bipolar disorder and schizophrenia. *Mol. Psychiatry* 19, 1017–1024.
- Rutherford, N.J., Carrasquillo, M.M., Li, M., Biscoglio, G., Menke, J., Josephs, K.A., Parisi, J.E., Petersen, R.C., Graff-Radford, N.R., Younkin, S.G., Dickson, D.W., Rademakers, R., 2012. TMEM106B risk variant is implicated in the pathologic presentation of Alzheimer disease. *Neurology* 79, 717–718.
- Sachidanandam, R., Craddock, N., Manolio, T.A., S. Nejentsev, N. Walker, D. Riches, M. Egholm, J.A. Todd, J.C. Cohen, E. Boerwinkle, T.H. Mosley, H.H. Hobbs, Levy, S., Wheeler, D.A., Bentley, D.R., Wang, J., Li, H., Lam, H.Y., Conrad, D.F., Irwin, J.A., Balaesque, P., MC. Wendl, R.K.W., Xing, J., Stranger, B.E., J. Marchini, B.H., Dixon, A.L., Genovese, G., MW. Nachman, S.L.C., Kondrashov, A.S., Roach, J.C., B Charlesworth M T Morgan, J. Maynard Smith, J.H., JJ. Cai, JM. Macpherson, G. Sella, DA. Petrov, BF. Voight, S. Kudaravalli, X. Wen, JK. Pritchard, LB. Barreiro, G. Laval, H. Quach, E. Patin, L. Quintana-Murci, Lamason, R.L., Van Kim, C.T.Y.C.J.P.C.C.L., Myers, S., S. Myers, C. Freeman, A. Auton, P. Donnelly, G. McVean, Baudat, F., E D Parvanov P M Petkov, Liu, J.Z., Sanna, S., AD. Ewing, H.H.K., Mills, R.E., Liti, G., Y. Li, C. Willer, S. Sanna, G. Abecasis, 2010. A map of human genome variation from population-scale sequencing. *Nature* 467, 1061–1073.
- Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 511, 421–427.
- Schizophrenia Working Group of the Psychiatric Genomics Consortium, Ripke, S., Walters, J.T.R., O'Donovan, M.C., 2020. Mapping genomic loci prioritises genes and implicates synaptic biology in schizophrenia. *medRxiv* 2020.09.12.20192922.
- Schwenk, B.M., Lang, C.M., Hogl, S., Tahirovic, S., Orozco, D., Rentzsch, K., Lichtenthaler, S.F., Hoogenraad, C.C., Capell, A., Haass, C., Edbauer, D., 2014. The FTLTD risk factor TMEM106B and MAP6 control dendritic trafficking of lysosomes. *EMBO J.* 33, 450–467.
- Severance, E.G., Yolken, R.H., 2008. Novel alpha7 nicotinic receptor isoforms and deficient cholinergic transcription in schizophrenia. *Genes Brain Behav.* 7, 37–45.
- Shang, J., Yamashita, T., Zhai, Y., Nakano, Y., Morihara, R., Fukui, Y., Hishikawa, N., Ohta,

- Y., Abe, K., 2016. Strong Impact of Chronic Cerebral Hypoperfusion on Neurovascular Unit, Cerebrovascular Remodeling, and Neurovascular Trophic Coupling in Alzheimer's Disease Model Mouse. *J. Alzheimers. Dis.* 52, 113–126.
- Simeone, J.C., Ward, A.J., Rotella, P., Collins, J., Windisch, R., 2015. An evaluation of variation in published estimates of schizophrenia prevalence from 1990–2013: a systematic literature review. *BMC Psychiatry* 15, 193.
- Singh, J., Kour, K., Jayaram, M.B., 2012. Acetylcholinesterase inhibitors for schizophrenia. *Cochrane Database Syst. Rev.* 1, CD007967.
- Strand, J.-E., Nybäck, H., 2005. Tobacco use in schizophrenia: a study of cotinine concentrations in the saliva of patients and controls. *Eur. Psychiatry* 20, 50–54.
- Sullivan, P.F., Kendler, K.S., Neale, M.C., 2003. Schizophrenia as a complex trait: evidence from a meta-analysis of twin studies. *Arch. Gen. Psychiatry* 60, 1187–1192.
- Thorgeirsson, T.E., Gudbjartsson, D.F., Surakka, I., Vink, J.M., Amin, N., Geller, F., Sulem, P., Rafnar, T., Esko, T., Walter, S., Gieger, C., Rawal, R., Mangino, M., Prokopenko, I., Mägi, R., Keskitalo, K., Gudjonsdottir, I.H., Gretarsdottir, S., Stefansson, H., Thompson, J.R., Aulchenko, Y.S., Nelis, M., Aben, K.K., den Heijer, M., Dirksen, A., Ashraf, H., Soranzo, N., Valdes, A.M., Steves, C., Uitterlinden, A.G., Hofman, A., Tönjes, A., Kovacs, P., Hottenga, J.J., Willemsen, G., Vogelzangs, N., Döring, A., Dahmen, N., Nitz, B., Pergadia, M.L., Saez, B., De Diego, V., Lezcano, V., Garcia-Prats, M.D., Ripatti, S., Perola, M., Kettunen, J., Hartikainen, A.-L., Pouta, A., Laitinen, J., Isohanni, M., Huei-Yi, S., Allen, M., Krestyaninova, M., Hall, A.S., Jones, G.T., van Rij, A.M., Mueller, T., Dieplinger, B., Haltmayer, M., Jonsson, S., Matthiasson, S.E., Oskarsson, H., Tyrfingsson, T., Kiemeny, L.A., Mayordomo, J.I., Lindholt, J.S., Pedersen, J.H., Franklin, W.A., Wolf, H., Montgomery, G.W., Heath, A.C., Martin, N.G., Madden, P.A.F., Giegling, I., Rujescu, D., Järvelin, M.-R., Salomaa, V., Stumvoll, M., Spector, T.D., Wichmann, H.-E., Metspalu, A., Samani, N.J., Penninx, B.W., Oostra, B.A., Boomsma, D.I., Tiemeier, H., van Duijn, C.M., Kaprio, J., Gulcher, J.R., ENGAGE Consortium, McCarthy, M.I., Peltonen, L., Thorsteinsdottir, U., Stefansson, K., 2010. Sequence variants at CHRNA3-CHRNA6 and CYP2A6 affect smoking behavior. *Nat. Genet.* 42, 448–453.
- Tidey, J.W., Colby, S.M., Xavier, E.M.H., 2014. Effects of smoking abstinence on cigarette craving, nicotine withdrawal, and nicotine reinforcement in smokers with and without schizophrenia. *Nicotine Tob. Res.* 16, 326–334.
- Tobacco and Genetics Consortium, 2010. Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nat. Genet.* 42, 441–447.
- Van Deerlin, V.M., Sleiman, P.M.A., Martinez-Lage, M., Chen-Plotkin, A., Wang, L.-S., Graff-Radford, N.R., Dickson, D.W., Rademakers, R., Boeve, B.F., Grossman, M., Arnold, S.E., Mann, D.M.A., Pickering-Brown, S.M., Seelaar, H., Heutink, P., van Swieten, J.C., Murrell, J.R., Ghetti, B., Spina, S., Grafman, J., Hodges, J., Spillantini, M.G., Gilman, S., Lieberman, A.P., Kaye, J.A., Woltjer, R.L., Bigio, E.H., Mesulam, M., Al-Sarraj, S., Troakes, C., Rosenberg, R.N., White, C.L., 3rd, Ferrer, I., Lladó, A., Neumann, M., Kretschmar, H.A., Hulette, C.M., Welsh-Bohmer, K.A., Miller, B.L., Alzualde, A., Lopez de Munain, A., McKee, A.C., Gearing, M., Levey, A.I., Lah, J.J., Hardy, J., Rohrer, J.D., Lashley, T., Mackenzie, I.R.A., Feldman, H.H., Hamilton, R.L., Dekosky, S.T., van der Zee, J., Kumar-Singh, S., Van Broeckhoven, C., Mayeux, R., Vonsattel, J.P.G., Troncoso, J.C., Kril, J.J., Kwok, J.B.J., Halliday, G.M., Bird, T.D., Ince, P.G., Shaw, P.J., Cairns, N.J.,

- Morris, J.C., McLean, C.A., DeCarli, C., Ellis, W.G., Freeman, S.H., Frosch, M.P., Growdon, J.H., Perl, D.P., Sano, M., Bennett, D.A., Schneider, J.A., Beach, T.G., Reiman, E.M., Woodruff, B.K., Cummings, J., Vinters, H.V., Miller, C.A., Chui, H.C., Alafuzoff, I., Hartikainen, P., Seilhean, D., Galasko, D., Masliah, E., Cotman, C.W., Tuñón, M.T., Martínez, M.C.C., Munoz, D.G., Carroll, S.L., Marson, D., Riederer, P.F., Bogdanovic, N., Schellenberg, G.D., Hakonarson, H., Trojanowski, J.Q., Lee, V.M.-Y., 2010. Common variants at 7p21 are associated with frontotemporal lobar degeneration with TDP-43 inclusions. *Nat. Genet.* 42, 234–239.
- Vink, J.M., Willemsen, G., Boomsma, D.I., 2005. Heritability of smoking initiation and nicotine dependence. *Behav. Genet.* 35, 397–406.
- Volkow, N.D., 2009. Substance use disorders in schizophrenia--clinical implications of comorbidity. *Schizophr. Bull.* 35, 469–472.
- Wallace, T.L., Bertrand, D., 2015. Neuronal $\alpha 7$ Nicotinic Receptors as a Target for the Treatment of Schizophrenia. *Int. Rev. Neurobiol.* 124, 79–111.
- Willer, C.J., Li, Y., Abecasis, G.R., 2010. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26, 2190–2191.
- Zhang, H.-K., Eaton, J.B., Fedolak, A., Gunosewoyo, H., Onajole, O.K., Brunner, D., Lukas, R.J., Yu, L.-F., Kozikowski, A.P., 2016. Synthesis and biological evaluation of novel hybrids of highly potent and selective $\alpha 4\beta 2$ -Nicotinic acetylcholine receptor (nAChR) partial agonists. *Eur. J. Med. Chem.* 124, 689–697.
- Zheng, J., Erzurumluoglu, A.M., Elsworth, B.L., Kemp, J.P., Howe, L., Haycock, P.C., Hemani, G., Tansey, K., Laurin, C., Early Genetics and Lifecourse Epidemiology (EAGLE) Eczema Consortium, Pourcain, B.S., Warrington, N.M., Finucane, H.K., Price, A.L., Bulik-Sullivan, B.K., Anttila, V., Paternoster, L., Gaunt, T.R., Evans, D.M., Neale, B.M., 2017. LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics* 33, 272–279.

Genome-wide analyses of smoking behaviors in schizophrenia: findings from the Psychiatric Genomics Consortium

Roseann E Peterson, PhD^{1,‡}; Tim B Bigdeli, PhD^{1,2,‡}; Stephan Ripke, MD, PhD^{3,4}; Silviu-Alin Bacanu, PhD¹; Pablo V Gejman, MD^{5,6}; Douglas F Levinson, MD⁷; Qingqin S Li, PhD⁸; Dai Wang, PhD⁸; Dan Rujescu, MD, PhD^{10,9}; Marcella Rietschel, MD, PhD¹¹; Daniel R Weinberger, MD^{12,13}; Richard E Straub, PhD¹²; James TR Walters, MD, PhD¹⁴; Bryan J Mowry, MD, FRANZCP^{15,16}; Roel A Ophoff, PhD^{17,18,19}; Ole A Andreassen, MD, PhD^{20,21}; Tõnu Esko, PhD^{22,23,24,25}; Tracey L Petryshen, PhD^{26,27,28}; Kenneth S Kendler, MD¹; Schizophrenia Working Group of the Psychiatric Genomics Consortium^a; Ayman H Fanous, MD^{1,2,29*}

¹Virginia Institute for Psychiatric and Behavioral Genetics, Department of Psychiatry, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

²Department of Psychiatry and Behavioral Sciences, State University of New York Downstate Medical Center, Brooklyn, NY, USA

³Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA

⁴Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA

⁵Department of Psychiatry and Behavioral Sciences, NorthShore University HealthSystem, Evanston, IL, USA

⁶Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL, USA

⁷Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA

⁸Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Raritan, NJ, USA

⁹Department of Psychiatry, University of Halle, Halle, Germany

¹⁰Department of Psychiatry, University of Munich, Munich, Germany

¹¹Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, University of Heidelberg, Heidelberg, Germany

¹²Lieber Institute for Brain Development, Baltimore, MD, USA

¹³Departments of Psychiatry, Neurology, Neuroscience and Institute of Genetic Medicine, Johns Hopkins School of Medicine, Baltimore, MD

¹⁴MRC Centre for Neuropsychiatric Genetics and Genomics, Institute of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK

¹⁵Queensland Brain Institute, The University of Queensland, Brisbane, Queensland, Australia

¹⁶Queensland Centre for Mental Health Research, University of Queensland, Brisbane, Queensland, Australia

¹⁷Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, CA, USA

¹⁸Department of Human Genetics, David Geffen School of Medicine, University of California, Los Angeles, CA, USA

¹⁹Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht, The Netherlands

²⁰NORMENT, KG Jebsen Centre for Psychosis Research, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

²¹Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway

²²Estonian Genome Center, University of Tartu, Tartu, Estonia

²³Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA, USA

²⁴Department of Genetics, Harvard Medical School, Boston, MA, USA

²⁵Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge, MA, USA

²⁶Center for Human Genetic Research and Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA

²⁷Department of Psychiatry, Harvard Medical School, Boston, MA, USA

²⁸The Broad Institute of MIT and Harvard, Boston MA

²⁹Mental Health Service Line, Washington VA Medical Center, Washington DC, USA

*Corresponding author

Department of Psychiatry and Behavioral Sciences
SUNY Downstate Medical Center
450 Clarkson Ave
Brooklyn, NY 11203

[‡]These authors contributed equally to this work

^aFull list of consortium members appears at the end of this Supplement.

Study descriptions

For participating studies, ascertainment and diagnosis has been described previously (1), and are reproduced here in brief. For each study, the first line gives the name of the principal investigator(s) name, PubMed ID for citations describing the sample in detail, the country of origin or study name, and the study identifier used herein.

Discovery studies (SCZ smoking behaviors)

Rietschel/Rujescu | 19571808 | Bonn/Mannheim, Germany | boco

Rujescu, D | 19571808 | Munich, Germany | munc

These German samples were collected by separate groups from Bonn/Mannheim and the University of Munich. For the PGC analyses, the samples were combined by chip and ancestry.

In Bonn/Mannheim, cases were ascertained as previously described (2). Controls were drawn from three population-based epidemiological studies (PopGen) (3), the Cooperative Health Research in the Region of Augsburg (KORA) study (4), and the Heinz Nixdorf Recall (HNR) study (5). All participants gave written informed consent and the local ethics committees approved the human subjects protocols.

Cases were ascertained from the Munich area of Germany, as described previously (2). The controls were unrelated volunteers randomly selected from the general population of Munich. All were screened to exclude a history of psychosis/central neurological disease either personally or in a first-degree relative. All participants gave written informed consent and the local ethics committees approved the human subjects protocols.

Petryshen, T | Not Published | Boston, US (CIDAR) | cims

Cases were recruited from inpatient and outpatient settings in the Boston area by clinician referral, through review of medical records, or through advertisements in local media. Cases were diagnosed with DSM-IV schizophrenia through a structured clinical interview (SCID) by trained interviewers with review of medical records and a best estimate diagnostic procedure including reliability trials across interviewers. A psychiatrist or a PhD-level mental health professional made the final diagnostic determination. Controls were ascertained through local advertisements from the same geographical area. Ethical approval was provided by local ethics committees and all participants gave written informed consent.

Walters, J | 21850710 | Cardiff, UK (CogUK) | cou3

Cases were recruited from community mental health teams in Wales and England on the basis of a clinical diagnosis of schizophrenia or schizoaffective disorder (depressed sub-type) as described previously (6). Diagnosis was confirmed following a SCAN (7) interview and review of case notes followed by consensus diagnosis according to DSM-IV (8) criteria. The samples were genotyped at the Broad Institute. The UK Multicentre Research Ethics Committee (MREC) approved the study and all participants provided valid informed consent.

Esko, T | 15133739 | Estonia (EGCUT) | egcu

The Estonian cohort comes from the population-based biobank of the Estonian Genome Project of University of Tartu (EGCUT) (9). The project was conducted according to the Estonian Gene Research Act and all participants provided informed consent (www.biobank.ee). In total, 52,000 individuals aged 18 years or older participated in this cohort (33% men, 67% women). The population distributions of the cohort reflect those of the Estonian population (83% Estonians, 14% Russians and 3% other). General practitioners (GP) and physicians in the hospitals randomly recruited the participants. A Computer-

Assisted Personal interview was conducted over 1-2 ours at doctors' offices. Data on demographics, genealogy, educational and occupational history, lifestyle and anthropometric and physiological data were assessed. Schizophrenia was diagnosed prior to the recruitment by a psychiatrist according to ICD-10 criteria and identified from the Estonian Biobank phenotype database. Controls were drawn from a larger pool of genotyped biobank samples by matching on gender, age and genetic ancestry. All the controls were population-based and have not been sampled for any specific disease.

Weinberger, D | 11381111 | NIMH CBDB | lie2, lie5

Subjects were recruited from the Clinical Brain Disorders Branch of the NIMH 'Sibling Study' as previously described (10). In brief, cases and controls gave informed consent and only participants of European ancestry were included in the current analysis. Cases completed a structured clinical interview and were diagnosed with schizophrenia-spectrum disorders. Samples were genotyped at the NIMH.

Gejman, P | 19571809 | US, Australia (MGS) | mgs2

European ancestry case samples were collected by the Molecular Genetics of Schizophrenia (MGS) collaboration across multiple sites in the USA and Australia as described in detail elsewhere (11). Cases gave written informed consent, and IRBs at each collecting site approved the human subjects protocol. A survey company (Knowledge Networks, under MGS guidance) collected the European ancestry control sample and ascertainment is described in detail elsewhere (12). DNA samples were genotyped at the Broad Institute.

Andreassen, O | 19571808 | Norway (TOP) | top8

In the TOP study (Tematisk område psykoser), cases of European ancestry, born in Norway, were recruited from psychiatric hospitals in the Oslo region. Patients were diagnosed according to SCID and further ascertainment details have been reported (13). Healthy control subjects were randomly selected from statistical records of persons from the same catchment area as the patient groups. All participants provided written informed consent and the human subjects protocol was approved by the Norwegian Scientific-Ethical Committee and the Norwegian Data Protection Agency.

Ophoff, R | 19571808 | Netherlands | ucla

The case sample consisted of inpatients and outpatients recruited through psychiatric hospitals and institutions throughout the Netherlands. Cases with DSM-IV schizophrenia were included in the analysis. Further details on ascertainment are provided elsewhere (2). Controls came from the University Medical Centre Utrecht and were volunteers with no psychiatric history. Ethical approval was provided by local ethics committees and all participants gave written informed consent.

*Replication studies (SCZ case-control)***Iwata, N | 20832056 | Japan | scz_jpn1_asn**

Case and control participants were recruited from the Tokai area of mainland Japan and self-identified as Japanese. Full details on sample ascertainment and ethical approval have been reported previously (14).

Liu, J | NP | Singapore (STCRP) | scz_tcr1_asn

The Singapore Translational and Clinical Research in Psychosis (STCRP) Study sample consisted of schizophrenia patients and healthy controls of Chinese ancestry. All patients were recruited from the Institute of Mental Health in Singapore from 2005-2008 and were aged 18-83 years. Diagnosis of schizophrenia was made using the Structured Clinical Interview for Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Research Version, Patient Edition (SCID-I/P) by trained raters. The controls were from the Singapore Prospective Study Program(15) and were randomly sampled from the Singapore population and approximately matched for age and sex to cases.

Sham, P | 24043878 | China | scz_hok2_asn

The case sample include patients with DSM-IV schizophrenia recruited in Hong Kong and Sichuan China described previously (16). The control group was a convenience sample gathered from several sources - subjects from a GWAS on bone mineral density (17), the control subjects from a GWAS on hypertension (18), and on liver cancer (19), and healthy control subjects recruited from Sichuan and Taiwan. All cases and controls gave written consent to participate. The was approved by the Institutional Review Boards of the University of Hong Kong and the West China Hospital at Sichuan University. Genotyping was performed at deCODE Genetics.

*Replication studies (SCZ smoking behaviors)***Esko, T; Wang, D | 15133739 | J&J cases, EGCUT controls | jr3a, jr3b, jri6, jrsa**

Cases were collected by Johnson and Johnson (J&J) as part of clinical collaborations with hospitals and outpatient centers in Eastern Europe. Cases were diagnosed according to DSM-IV criteria, following a structured clinical interview, with medical record review by a trained psychiatrist. There were reliability trials across centers. Most of the cases were from Estonian and Russia (>100) with intermediate numbers from Austria, the Czech Republic, Latvia, Lithuania, and Spain (50-100). There were smaller collections from Bulgaria, Hungary, and Poland (<50). Most of the Eastern European controls were from the Estonian Biobank project (EGCUT) (9) and were ancestrally matched with cases from the J&J sample.

Supplemental Tables

Supplemental Table S1. SNP-based heritability estimates for SI and CPD among PGC-SCZ cases. For each trait, LD-score regression summary output is displayed.

Study	Trait	K_{pop}	K_{sample}	β_0 (SE)	λ_{GC}	$h^2_{intercept}$ (SE)	$h^2_{constrain}$ (SE)
TAG	SI	0.350	0.562	1.003 (0.007)	1.093	0.116 (0.011)	0.120 (0.007)
	CPD	·	·	1.010 (0.007)	1.056	0.054 (0.016)	0.067 (0.012)
PGC	SI _{SCZ}	0.725	0.725	0.987 (0.006)	1.020	0.464 (0.158)	0.219 (0.112)
	CPD _{SCZ}	·	·	0.999 (0.006)	1.005	0.107 (0.138)	0.092 (0.096)

K_{pop} and K_{sample} are the assumed population and calculated sample prevalence, respectively. For each trait, λ_{GC} and β_0 , are the estimated genomic control factor (λ) and regression intercept, respectively. Estimates of heritability, h^2 , are reported on the liability scale for SI and the observed scale for CPD, for analyses in which the intercept was unconstrained ($h^2_{intercept}$) or constrained to 1 ($h^2_{constrain}$).

Supplemental Table S2. Association of TAG-SI polygenic scores with SCZ. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained in terms of Nagelkerke's R^2 ; β and SE are the beta regression coefficient and standard error from logistic regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	Z	P-value	P_{het}
1.00E-05	2.52E-05	0.104	0.099	1.049	0.294	0.629
1.00E-04	0.000128	0.091	0.038	2.364	0.018	0.740
0.001	0.000191	0.036	0.012	2.889	0.004	0.079
0.01	0.000602	0.027	0.005	5.122	3.03E-07	0.078
0.1	0.00122	0.016	0.002	7.280	3.38E-13	0.013
1	0.00133	0.012	0.002	7.608	2.83E-14	0.016

Supplemental Table S3. Association of TAG-CPD polygenic scores with SCZ. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained in terms of Nagelkerke's R^2 ; β and SE are the beta regression coefficient and standard error from logistic regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	Z	P -value	P_{het}
1.00E-05	0.000391	0.0260	0.0063	4.1334	3.58E-05	0.125
1.00E-04	6.62E-05	0.0077	0.0045	1.6997	0.089	0.126
0.001	5.01E-05	0.0027	0.0018	1.4794	0.139	0.228
0.01	5.97E-06	0.0004	0.0008	0.5104	0.610	0.831
0.1	1.72E-06	0.0001	0.0003	0.2736	0.784	0.533
1	1.61E-07	0.0000	0.0002	0.0838	0.933	0.435

Supplemental Table S4. Association of TAG-SI polygenic scores with SCZ-SI. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained in terms of Nagelkerke's R^2 ; β and SE are the beta regression coefficient and standard error from logistic regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	Z	P -value	P_{het}
1.00E-05	0.0004	0.4182	0.3797	1.1014	2.71E-01	0.442
1.00E-04	0.0004	0.1718	0.1460	1.1766	2.39E-01	0.794
0.001	0.0039	0.1717	0.0475	3.6168	3.01E-04	0.952
0.01	0.0058	0.0873	0.0197	4.4314	9.57E-06	0.463
0.1	0.0050	0.0345	0.0085	4.0873	4.44E-05	0.452
1	0.0040	0.0219	0.0060	3.6658	2.49E-04	0.629

Table S5. Association of TAG-CPD polygenic scores with SCZ-CPD. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained; β and SE are the beta regression coefficient and standard error from linear regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	Z	P -value	P_{het}
1.00E-05	0.0012	0.0241	0.0127	1.9026	5.72E-02	0.838
1.00E-04	0.0016	0.0202	0.0091	2.2170	2.67E-02	0.648
0.001	0.0008	0.0058	0.0037	1.5811	1.14E-01	0.378
0.01	0.0035	0.0052	0.0016	3.3370	8.57E-04	0.747
0.1	0.0031	0.0022	0.0007	3.1336	1.74E-03	0.541
1	0.0013	0.0009	0.0005	2.0533	4.01E-02	0.088

Table S6. Association of PGC-SCZ polygenic scores with SCZ-SI. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained in terms of Nagelkerke's R^2 ; β and SE are the beta regression coefficient and standard error from logistic regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	Z	P -value	P_{het}
1.00E-04	9.59E-05	-0.0288	0.0506	-0.5698	0.569	0.495
0.001	3.11E-05	-0.0108	0.0332	-0.3245	0.746	0.634
0.01	1.49E-05	0.0047	0.0209	0.2247	0.822	0.394
0.1	0.0002	0.0122	0.0133	0.9170	0.359	0.412
1	0.0004	0.0128	0.0109	1.1752	0.240	0.662

Table S7. Association of PGC-SCZ polygenic scores with SCZ-CPD. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained; β and SE are the beta regression coefficient and standard error from linear regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	Z	P -value	P_{het}
1.00E-04	0.0007	0.0389	0.0269	1.4419	0.149	0.362
0.001	0.0005	0.0226	0.0176	1.2879	0.198	0.168
0.01	0.0005	0.0137	0.0112	1.2231	0.221	0.149
0.1	0.0004	0.0079	0.0072	1.1025	0.270	0.155
1	0.0001	0.0036	0.0059	0.6110	0.541	0.437

Table S8. Association of TAG-SI polygenic scores with age-of-onset. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained; β and SE are the beta regression coefficient and standard error from linear regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	t -statistic	P -value	P_{het}
1.00E-05	1.03E-05	-0.034	0.14	-0.247	0.805	0.589
1.00E-04	5.93E-06	0.01	0.054	0.187	0.852	0.388
0.001	2.73E-05	-0.007	0.018	-0.4	0.689	0.391
0.01	2.68E-07	0	0.007	0.04	0.968	0.529
0.1	2.68E-06	0	0.003	0.126	0.9	0.0308
1	9.58E-06	-0.001	0.002	-0.237	0.812	0.0014

Table S9. Association of TAG-SI polygenic scores with positive symptoms. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained; β and SE are the beta regression coefficient and standard error from linear regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	t -statistic	P -value	P_{het}
1.00E-05	6.00E-04	-0.27	0.158	-1.707	0.0878	0.982
1.00E-04	0.00107	-0.141	0.062	-2.279	0.0227	0.643
0.001	2.13E-05	0.006	0.02	0.322	0.748	0.508
0.01	1.59E-05	0.002	0.008	0.278	0.781	0.852
0.1	1.85E-05	0.001	0.003	0.3	0.764	0.97
1	0.000136	0.002	0.002	0.813	0.416	0.907

Table S10. Association of TAG-CPD polygenic scores with positive symptoms. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained; β and SE are the beta regression coefficient and standard error from linear regression; P_{het} is the significance of Cochran's test for heterogeneity.

P_T	R^2	β	SE	t -statistic	P -value	P_{het}
1.00E-05	2.15E-05	0.003	0.01	0.323	0.747	0.159
1.00E-04	3.40E-05	0.003	0.007	0.406	0.685	0.457
0.001	0.00044	0.004	0.003	1.461	0.144	0.85
0.01	0.000569	0.002	0.001	1.662	0.0965	0.792
0.1	0.00115	0.001	0.001	2.368	0.0179	0.828
1	0.00155	0.001	0	2.748	0.00602	0.986

Table S11. Association of TAG-CPD polygenic scores with depression symptoms. For scores based on varying P -value thresholds (P_T), R^2 is the variance explained; β and SE are the beta regression coefficient and standard error from linear regression; P_{het} is the significance of Cochran's test for heterogeneity.


P_T	R^2	β	SE	t -statistic	P -value	P_{het}
1.00E-05	0.000244	0.011	0.011	1.066	0.286	0.449
1.00E-04	2.75E-05	0.003	0.008	0.358	0.721	0.512
0.001	4.49E-06	0	0.003	0.145	0.885	0.532
0.01	3.98E-07	0	0.001	0.043	0.966	0.00209
0.1	0.000115	0	0.001	0.732	0.464	0.0242
1	0.000132	0	0	0.786	0.432	0.0452

Table S12. *Post-hoc* power analyses for top associated SNPs from TAG. For genomewide-significant SNPs in the TAG study of smoking behaviors (20) the statistical power to detect these associations in the PGC-SCZ study is shown, Chr is chromosome, SNP is single-nucleotide polymorphism, Frq is effect allele frequency, β and SE are the beta regression coefficient and standard error from linear/logistic regression, P is P-value, N is sample size, CPD is cigarettes-per-day, SI is smoking initiation, Cessation is smoking cessation. The “GeneticsDesign” package in Bioconductor was used to estimate the power to detect SNP effects for quantitative traits (CPD) (<https://www.bioconductor.org/packages/devel/bioc/manuals/GeneticsDesign/>) and CaTS Power Calculator (http://csg.sph.umich.edu/abecasis/CaTS/gas_power_calculator/index.html) (21) was used for binary traits (SI, Cessation).

Marker		TAG Meta-Analysis				PGC-SCZ Results			
Chr	SNP Gene	A1/A2 A1 Frq.	Trait N	β SE	P	N	β SE	P	Power
15	rs16969968 <i>CHRNA3</i>	G/A 0.65	CPD 73853	-1.00 0.06	2.8x10 ⁻⁷²	3344	-0.09 0.03	8.0x10 ⁻⁴	0.376
10	rs1329650 <i>LOC100188947</i>	T/G 0.28	CPD 73853	-0.37 0.06	5.7x10 ⁻¹⁰	3344	-0.02 0.03	0.421	0.087
9	rs3733829 <i>CYP2A6</i>	G/A 0.36	CPD 73853	0.33 0.06	1.0x10 ⁻⁸	3344	-0.004 0.03	0.891	0.083
11	rs6265 <i>BDNF</i>	T/C 0.21	SI 143023	-0.06 0.01	1.8x10 ⁻⁸	4991	-0.02 0.06	0.761	0.050
9	rs3025343 <i>DBH</i>	G/A 0.84	Cessation 64924	0.12 0.02	3.6x10 ⁻⁸	2742	NA	NA	0.135

Supplemental Figures

Figure S1. Proportion of smoking initiation by PGC2 schizophrenia cohort. The percent of each PGC2-SCZ cohort endorsing ever smoked a cigarette, SI is smoking initiation.



Cohort	SI Definition
boco	Smoking ever, 100 cigarettes, CPD $\geq$ 1
cims	100 cigarettes, CPD $\geq$ 1, Nicotine Dependence
cou3	Ever regular smoker, current smoker, age started smoking, years smoking, CPD $\geq$ 1
egcu	Never/current/former smoker, CPD $\geq$ 1
lie2	Current smoker, past smoker, packs per day $>$ 0
lie5	Current smoker, past smoker, packs per day $>$ 0
mgs2	Ever smoked on a daily basis, packs per day $>$ 0
munc	Ever smoked, CPD $\geq$ 1
top8	Lifetime smoker, smoking years $>$ 0
ucla	Smoking ever, 100 cigarettes, CPD $\geq$ 1

Figure S2. Smoking quantity by PGC2 schizophrenia cohort. The proportion of each cohort endorsing each smoking quantity category.

Smoking Quantity by PGC2 Schizophrenia Cohort




Figure S3. Association of PGC-SCZ with smoking behaviors among cases. For polygenic scores based on PGC-SCZ results, the variance explained for smoking behaviors is shown on the y-axis, in terms of Nagelkerke's pseudo- R^2 (SI) or R^2 (CPD); scores based on varying SNP P -value inclusion thresholds are displayed as colored bars.




Figure S4. Association of TAG-based polygene scores with SCZ clinical features. For polygenic scores based on analyses of smoking behaviors described by TAG, the variance explained for selected clinical features in PGC-SCZ is shown on the y-axis, in terms of R^2 ; scores based on varying SNP P -value inclusion thresholds are displayed as colored bars.




Figure S5. Manhattan plots for random-effects meta-analyses of CPD and SI in SCZ. Red and blue lines indicate thresholds for genome-wide significance ($P < 5 \times 10^{-8}$) and replication follow-up ($P < 10^{-6}$). For regions significant at the latter, the most significant “independent” SNP within a 500kb region is displayed as a blue diamond; nearby SNPs in linkage disequilibrium ($r^2 > 0.1$) are highlighted.






Figure S7. Regional association and forest plots for SI associations. (*left*) Regional association plot created using LocusZoom (22). LD of each SNP with the “index” SNP, displayed as a large purple diamond, is indicated by its color. (*right*) Study abbreviation “repli” is the replication sample; all other study indicators are as given above; *HetP* is the *P*-value for Cochran’s test of heterogeneity.






rs754168 **SI** **chr13:112983288:A/C** **hetPval:0.5736**

study	freq	info	or	se	pval	n
boco	0.351	0.706	0.61	0.183	0.007	308
cims	0.333	0.98	0.956	0.744	0.952	33
cou3	0.34	0.882	0.677	0.165	0.018	501
egcu	0.347	0.874	1.186	0.219	0.435	234
lie2	0.377	0.848	1.002	0.297	0.994	128
lie5	0.344	0.746	0.92	0.168	0.616	479
mgs2	0.365	0.75	0.751	0.088	0.0011	2194
munc	0.334	0.746	0.84	0.204	0.392	406
top8	0.332	0.699	0.74	0.208	0.148	319
ucla	0.334	0.753	0.628	0.194	0.0163	389
meta	0.352	0.767	0.769	0.054	1.21e-06	4991
repli			0.85	0.104	0.115	1277
joint			0.785	0.048	4.58e-07	6268

Figure S8. Regional association and forest plots for CPD associations. (left) Regional association plot created using LocusZoom(22). LD of each SNP with the “index” SNP, displayed as a large purple diamond, is indicated by its color. (right) Study abbreviation “repli” is the replication sample; all other study indicators are as given above; *HetP* is the *P*-value for Cochran’s test of heterogeneity.










Figure S9. Association of rs16969968 with SCZ stratified by smoking status (SI). For analyses of SCZ in the 10 studies contributing smoking data on case subjects, the forest plot displays frequencies and association summary statistics for comparisons of cases versus controls, SI cases that smoke versus controls, and non-SI cases versus controls, overall and further stratified by sex.



^aFull list of PGC contributors

Stephan Ripke ^{1,2}, Benjamin M Neale ^{1,2,3,4}, Aiden Corvin ⁵, James TR Walters ⁶, Kai-How Farh ¹, Peter A Holmans ^{6,7}, Phil Lee ^{1,2,4}, Brendan Bulik-Sullivan ^{1,2}, David A Collier ^{8,9}, Hailiang Huang ^{1,3}, Tune H Pers ^{3,10,11}, Ingrid Agartz ^{12,13,14}, Esben Agerbo ^{15,16,17}, Margot Albus ¹⁸, Madeline Alexander ¹⁹, Farooq Amin ^{20,21}, Silviu A Bacanu ²², Martin Begemann ²³, Richard A Belliveau Jr ², Judit Bene ^{24,25}, Sarah E Bergen ^{2,26}, Elizabeth Bevilacqua ², Tim B Bigdeli ²², Donald W Black ²⁷, Richard Bruggeman ²⁸, Nancy G Buccola ²⁹, Randy L Buckner ^{30,31,32}, William Byerley ³³, Wiepke Cahn ³⁴, Guiqing Cai ^{35,36}, Dominique Campion ³⁷, Rita M Cantor ³⁸, Vaughan J Carr ^{39,40}, Noa Carrera ⁶, Stanley V Catts ^{39,41}, Kimberley D Chambert ², Raymond CK Chan ⁴², Ronald YL Chan ⁴³, Eric YH Chen ⁴⁴, Wei Cheng ⁴⁵, Eric FC Cheung ⁴⁶, Siow Ann Chong ⁴⁷, C Robert Cloninger ⁴⁸, David Cohen ⁴⁹, Nadine Cohen ⁵⁰, Paul Cormican ⁵, Nick Craddock ^{6,7}, James J Crowley ⁵¹, David Curtis ^{52,53}, Michael Davidson ⁵⁴, Kenneth L Davis ³⁶, Franziska Degenhardt ^{55,56}, Jurgen Del Favero ⁵⁷, Ditte Demontis ^{17,58,59}, Dimitris Dikeos ⁶⁰, Timothy Dinan ⁶¹, Srdjan Djurovic ^{14,62}, Gary Donohoe ^{5,63}, Elodie Drapeau ³⁶, Jubao Duan ^{64,65}, Frank Dudbridge ⁶⁶, Naser Durmishi ⁶⁷, Peter Eichhammer ⁶⁸, Johan Eriksson ^{69,70,71}, Valentina Escott-Price ⁶, Laurent Essioux ⁷², Ayman H Fanous ^{73,74,75,76}, Martilias S Farrell ⁵¹, Josef Frank ⁷⁷, Lude Franke ⁷⁸, Robert Freedman ⁷⁹, Nelson B Freimer ⁸⁰, Marion Friedl ⁸¹, Joseph I Friedman ³⁶, Menachem Fromer ^{1,2,4,82}, Giulio Genovese ², Lyudmila Georgieva ⁶, Ina Giegling ^{81,83}, Paola Giusti-Rodríguez ⁵¹, Stephanie Godard ⁸⁴, Jacqueline I Goldstein ^{1,3}, Vera Golimbet ⁸⁵, Srihari Gopal ⁸⁶, Jacob Gratten ⁸⁷, Lieuwe de Haan ⁸⁸, Christian Hammer ²³, Marian L Hamshire ⁶, Mark Hansen ⁸⁹, Thomas Hansen ^{17,90}, Vahram Haroutunian ^{36,91,92}, Annette M Hartmann ⁸¹, Frans A Henskens ^{39,93,94}, Stefan Herms ^{55,56,95}, Joel N Hirschhorn ^{3,11,96}, Per Hoffmann ^{55,56,95}, Andrea Hofman ^{55,56}, Mads V Hollegaard ⁹⁷, Masashi Ikeda ⁹⁸, Inge Joa ⁹⁹, Antonio Julià ¹⁰⁰, René S Kahn ¹⁰¹, Luba Kalaydjieva ^{102,103}, Sena Karachanak-Yankova ¹⁰⁴, Juha Karjalainen ⁷⁸, David Kavanagh ⁶, Matthew C Keller ¹⁰⁵, James L Kennedy ^{106,107,108}, Andrey Khrunin ¹⁰⁹, Yunjung Kim ⁵¹, Janis Klovins ¹¹⁰, James A Knowles ¹¹¹, Bettina Konte ⁸¹, Vaidutis Kucinskas ¹¹², Zita Ausrele Kucinskiene ¹¹², Hana Kuzelova-Ptackova ^{113,114}, Anna K Kähler ²⁶, Claudine Laurent ^{19,115}, Jimmy Lee ^{47,116}, S Hong Lee ⁸⁷, Sophie E Legge ⁶, Bernard Lerer ¹¹⁷, Miaoxin Li ¹¹⁸, Tao Li ¹¹⁹, Kung-Yee Liang ¹²⁰, Jeffrey Lieberman ¹²¹, Svetlana Limborska ¹⁰⁹, Carmel M Loughland ^{39,122}, Jan Lubinski ¹²³, Jouko Lönnqvist ¹²⁴, Milan Macek ^{113,114}, Patrik

KE Magnusson ²⁶, Brion S Maher ¹²⁵, Wolfgang Maier ¹²⁶, Jacques Mallet ¹²⁷, Sara Marsal ¹⁰⁰, Manuel Mattheisen ^{17,58,59,128}, Morten Mattingsdal ^{14,129}, Robert W McCarley ^{130,131}, Colm McDonald ¹³², Andrew M McIntosh ^{133,134}, Sandra Meier ⁷⁷, Carin J Meijer ⁸⁸, Bela Melegh ^{24,25}, Ingrid Melle ^{14,135}, Raquelle I Mesholam-Gately ^{130,136}, Andres Metspalu ¹³⁷, Patricia T Michie ^{39,138}, Lili Milani ¹³⁷, Vihra Milanova ¹³⁹, Younes Mokrab ⁸, Derek W Morris ^{5,63}, Ole Mors ^{17,58,140}, Kieran C Murphy ¹⁴¹, Robin M Murray ¹⁴², Inez Myin-Germeys ¹⁴³, Bertram Müller-Myhsok ^{144,145,146}, Mari Nelis ¹³⁷, Igor Nenadic ¹⁴⁷, Deborah A Nertney ¹⁴⁸, Gerald Nestadt ¹⁴⁹, Kristin K Nicodemus ¹⁵⁰, Liene Nikitina-Zake ¹¹⁰, Laura Nisenbaum ¹⁵¹, Annelie Nordin ¹⁵², Eadbhard O'Callaghan ¹⁵³, Colm O'Dushlaine ², F Anthony O'Neill ¹⁵⁴, Sang-Yun Oh ¹⁵⁵, Ann Olincy ⁷⁹, Line Olsen ^{17,90}, Jim Van Os ^{143,156}, Christos Pantelis ^{39,157}, George N Papadimitriou ⁶⁰, Sergi Papiol ²³, Elena Parkhomenko ³⁶, Michele T Pato ¹¹¹, Tiina Paunio ^{158,159}, Milica Pejovic-Milovancevic ¹⁶⁰, Diana O Perkins ¹⁶¹, Olli Pietiläinen ^{159,162}, Jonathan Pimm ⁵³, Andrew J Pocklington ⁶, Danielle Posthuma ^{163,164,165}, John Powell ¹⁴², Alkes Price ¹⁶⁶, Ann E Pulver ¹⁴⁹, Shaun Purcell ⁸², Digby Quested ¹⁶⁷, Henrik B Rasmussen ^{17,90}, Abraham Reichenberg ³⁶, Mark A Reimers ¹⁶⁸, Alexander L Richards ^{6,7}, Joshua L Roffman ^{30,32}, Panos Roussos ^{82,169}, Douglas M Ruderfer ⁸², Veikko Salomaa ⁷¹, Alan R Sanders ^{64,65}, Ulrich Schall ^{39,122}, Christian R Schubert ¹⁷⁰, Thomas G Schulze ^{77,171}, Sibylle G Schwab ¹⁷², Edward M Scolnick ², Rodney J Scott ^{39,173,174}, Larry J Seidman ^{130,136}, Jianxin Shi ¹⁷⁵, Engilbert Sigurdsson ¹⁷⁶, Teimuraz Silagadze ¹⁷⁷, Jeremy M Silverman ^{36,178}, Kang Sim ⁴⁷, Petr Slominsky ¹⁰⁹, Jordan W Smoller ^{2,4}, Hon-Cheong So ⁴³, Chris C A Spencer ¹⁷⁹, Eli A Stahl ^{3,82}, Hreinn Stefansson ¹⁸⁰, Stacy Steinberg ¹⁸⁰, Elisabeth Stogmann ¹⁸¹, Richard E Straub ¹⁸², Eric Strengman ^{183,184}, Jana Strohmaier ⁷⁷, T Scott Stroup ¹²¹, Mythily Subramaniam ⁴⁷, Jaana Suvisaari ¹²⁴, Dragan M Svrakic ⁴⁸, Jin P Szatkiewicz ⁵¹, Erik Söderman ¹², Srinivas Thirumalai ¹⁸⁵, Draga Toncheva ¹⁰⁴, Sarah Tosato ¹⁸⁶, Juha Veijola ¹⁸⁷, Peter M Visscher ⁸⁷, John Waddington ¹⁸⁸, Dermot Walsh ¹⁸⁹, Dai Wang ⁸⁶, Qiang Wang ¹¹⁹, Bradley T Webb ²², Mark Weiser ⁵⁴, Durk Wiersma ¹⁹⁰, Dieter B Wildenauer ¹⁹¹, Nigel M Williams ¹⁹², Stephanie Williams ⁵¹, Stephanie H Witt ⁷⁷, Aaron R Wolen ¹⁶⁸, Emily HM Wong ⁴³, Brandon K Wormley ²², Hualin Simon Xi ¹⁹³, Clement C Zai ^{106,107}, Xuebin Zheng ¹⁹⁴, Fritz Zimprich ¹⁸¹, Naomi R Wray ⁸⁷, Kari Stefansson ¹⁸⁰, Wellcome Trust Case-Control Consortium 2 ¹⁹⁵, Rolf Adolfsson ¹⁵², Ole A Andreassen ^{14,135}, Douglas HR Blackwood ¹³⁴, Elvira Bramon ¹⁹⁶, Joseph D Buxbaum ^{35,36,91,197}, Anders D Børghlum ^{17,58,59,140}, Sven Cichon ^{55,56,95,198}, Ariel Darvasi ¹⁹⁹, Enrico Domenici ²⁰⁰, Hannelore Ehrenreich ²³, Tõnu Esko ^{3,11,96,137},

Pablo V Gejman ^{64,65}, Michael Gill ⁵, Hugh Gurling ⁵³, Christina M Hultman ²⁶, Nakao Iwata ⁹⁸, Assen V Jablensky ^{39,201,202,203}, Erik G Jönsson ¹², Kenneth S Kendler ²⁰⁴, George Kirov ⁶, Jo Knight ^{106,107,108}, Todd Lencz ^{205,206,207}, Douglas F Levinson ¹⁹, Qingqin S Li ⁸⁶, Jianjun Liu ^{194,208}, Anil K Malhotra ^{205,206,207}, Steven A McCarroll ^{2,96}, Andrew McQuillin ⁵³, Jennifer L Moran ², Preben B Mortensen ^{15,16,17}, Bryan J Mowry ^{87,209}, Markus M Nöthen ^{55,56}, Roel A Ophoff ^{38,80,210}, Michael J Owen ^{6,7}, Psychosis Endophenotypes International Consortium ²¹¹, Aarno Palotie ^{4,162,212}, Carlos N Pato ¹¹¹, Tracey L Petryshen ^{130,212,213}, Marcella Rietschel ⁷⁷, Brien P Riley ²⁰⁴, Dan Rujescu ^{81,83}, Pak C Sham ²¹⁴, Pamela Sklar ^{82,91,169}, David St Clair ²¹⁵, Daniel R Weinberger ^{182,216}, Jens R Wendland ¹⁷⁰, Thomas Werge ^{17,90,217}, Mark J Daly ¹, Patrick F Sullivan ^{26,51,161}, Michael C O'Donovan ^{6,7}.

¹ Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA. ² Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA. ³ Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge, MA, USA. ⁴ Psychiatric and Neurodevelopmental Genetics Unit, Massachusetts General Hospital, Boston, MA, USA. ⁵ Neuropsychiatric Genetics Research Group, Department of Psychiatry, Trinity College Dublin, Ireland. ⁶ MRC Centre for Neuropsychiatric Genetics and Genomics, Institute of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK. ⁷ National Centre for Mental Health, Cardiff University, Cardiff, Wales. ⁸ Eli Lilly and Company Limited, Erl Wood Manor, Sunninghill Road, Windlesham, Surrey, UK. ⁹ Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, King's College London, London, UK. ¹⁰ Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Lyngby, Denmark. ¹¹ Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA, USA. ¹² Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden. ¹³ Department of Psychiatry, Diakonhjemmet Hospital, Oslo, Norway. ¹⁴ NORMENT, KG Jebsen Centre for Psychosis Research, Institute of Clinical Medicine, University of Oslo, Oslo, Norway. ¹⁵ Centre for Integrative Register-based Research, CIRRAU, Aarhus University, Aarhus, Denmark. ¹⁶ National Centre for Register-based Research, Aarhus University, Aarhus, Denmark. ¹⁷ The Lundbeck Foundation Initiative for Integrative Psychiatric Research, IPSYCH, Denmark. ¹⁸ State Mental Hospital, Haar, Germany. ¹⁹

Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA.²⁰

Department of Psychiatry and Behavioral Sciences, Atlanta Veterans Affairs Medical Center, Atlanta, GA, USA.²¹ Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, GA, USA.²²

Virginia Institute for Psychiatric and Behavioral Genetics, Department of Psychiatry, Virginia Commonwealth University, Richmond, VA, USA.²³ Clinical Neuroscience, Max Planck Institute of Experimental Medicine, Göttingen, Germany.²⁴ Department of Medical Genetics, University of Pécs, Pécs, Hungary.²⁵ Szentagothai Research Center, University of Pécs, Pécs, Hungary.²⁶ Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.²⁷ Department of Psychiatry, University of Iowa Carver College of Medicine, Iowa City, IA, USA.²⁸ University Medical Center Groningen, Department of Psychiatry, University of Groningen, The Netherlands.²⁹ School of Nursing, Louisiana State University Health Sciences Center, New Orleans, LA, USA.³⁰ Athinoula A. Martinos Center, Massachusetts General Hospital, Boston, MA, USA.³¹ Center for Brain Science, Harvard University, Cambridge MA, USA.³² Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA.³³ Department of Psychiatry, University of California at San Francisco, San Francisco, CA, USA.³⁴ University Medical Center Utrecht, Department of Psychiatry, Rudolf Magnus Institute of Neuroscience, The Netherlands.³⁵ Department of Human Genetics, Icahn School of Medicine at Mount Sinai, New York, NY, USA.³⁶ Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA.³⁷ Centre Hospitalier du Rouvray and INSERM U1079 Faculty of Medicine, Rouen, France.³⁸ Department of Human Genetics, David Geffen School of Medicine, University of California, Los Angeles, CA, USA.³⁹ Schizophrenia Research Institute, Sydney, Australia.⁴⁰ School of Psychiatry, University of New South Wales, Sydney, Australia.⁴¹ Royal Brisbane and Women's Hospital, University of Queensland, Brisbane, Australia.⁴² Institute of Psychology, Chinese Academy of Science, Beijing, PR China.⁴³ Department of Psychiatry, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, PR China.⁴⁴ Department of Psychiatry and State Key Laboratory for Brain and Cognitive Sciences, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, PR China.⁴⁵ Department of Computer Science, University of North Carolina, Chapel Hill, NC, USA.⁴⁶ Castle Peak Hospital, Hong Kong SAR, PR China.⁴⁷ Institute of Mental Health, Singapore.⁴⁸ Department of Psychiatry, Washington University, St. Louis, MO, USA.⁴⁹ Department of Child and Adolescent

Psychiatry, Pierre and Marie Curie Faculty of Medicine and Brain and Spinal Cord Institute (ICM), Paris, France. ⁵⁰ Formerly of Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Raritan, NJ, USA. ⁵¹ Department of Genetics, University of North Carolina, Chapel Hill, NC, USA. ⁵² Department of Psychological Medicine, Queen Mary University of London, UK. ⁵³ Molecular Psychiatry Laboratory, Division of Psychiatry, University College London, UK. ⁵⁴ Sheba Medical Center, Tel Hashomer, Israel. ⁵⁵ Department of Genomics, Life and Brain Center, Bonn, Germany. ⁵⁶ Institute of Human Genetics, University of Bonn, Bonn, Germany. ⁵⁷ Applied Molecular Genomics Unit, VIB Department of Molecular Genetics, University of Antwerp, Antwerp, Belgium. ⁵⁸ Centre for Integrative Sequencing, iSEQ, Aarhus University, Aarhus, Denmark. ⁵⁹ Department of Biomedicine, Aarhus University, Aarhus, Denmark. ⁶⁰ First Department of Psychiatry, University of Athens Medical School, Athens, Greece. ⁶¹ Department of Psychiatry, University College Cork, Ireland. ⁶² Department of Medical Genetics, Oslo University Hospital, Oslo, Norway. ⁶³ Cognitive Genetics and Therapy Group, School of Psychology and Discipline of Biochemistry, National University of Ireland Galway, Ireland. ⁶⁴ Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL, USA. ⁶⁵ Department of Psychiatry and Behavioral Sciences, NorthShore University HealthSystem, Evanston, IL, USA. ⁶⁶ Department of Non-Communicable Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK. ⁶⁷ Department of Child and Adolescent Psychiatry, University Clinic of Psychiatry, Skopje, Republic of Macedonia. ⁶⁸ Department of Psychiatry, University of Regensburg, Regensburg, Germany. ⁶⁹ Department of General Practice, Helsinki University Central Hospital, Helsinki, Finland. ⁷⁰ Folkhälsan Research Center, Helsinki, Finland. ⁷¹ National Institute for Health and Welfare, Helsinki, Finland. ⁷² Translational Technologies and Bioinformatics, Pharma Research and Early Development, F.Hoffman-La Roche, Basel, Switzerland. ⁷³ Department of Psychiatry, Georgetown University School of Medicine, Washington DC, USA. ⁷⁴ Department of Psychiatry, Keck School of Medicine of the University of Southern California, Los Angeles, CA, USA. ⁷⁵ Department of Psychiatry, Virginia Commonwealth University School of Medicine, Richmond, VA, USA. ⁷⁶ Mental Health Service Line, Washington VA Medical Center, Washington DC, USA. ⁷⁷ Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, University of Heidelberg, Heidelberg, Germany. ⁷⁸ Department of Genetics, University of Groningen, University Medical Centre Groningen, The Netherlands.

⁷⁹ Department of Psychiatry, University of Colorado Denver, Aurora, CO, USA. ⁸⁰ Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, CA, USA. ⁸¹ Department of Psychiatry, University of Halle, Halle, Germany. ⁸² Division of Psychiatric Genomics, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ⁸³ Department of Psychiatry, University of Munich, Munich, Germany. ⁸⁴ Departments of Psychiatry and Human and Molecular Genetics, INSERM, Institut de Myologie, Hôpital de la Pitié-Salpêtrière, Paris, France. ⁸⁵ Mental Health Research Centre, Russian Academy of Medical Sciences, Moscow, Russia. ⁸⁶ Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Raritan, NJ, USA. ⁸⁷ Queensland Brain Institute, The University of Queensland, Brisbane, Queensland, Australia. ⁸⁸ Academic Medical Centre University of Amsterdam, Department of Psychiatry, Amsterdam, The Netherlands. ⁸⁹ Illumina, Inc., La Jolla, CA, USA. ⁹⁰ Institute of Biological Psychiatry, MHC Sct. Hans, Mental Health Services Copenhagen, Denmark. ⁹¹ Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ⁹² JJ Peters VA Medical Center, Bronx, NY, USA. ⁹³ Priority Research Centre for Health Behaviour, University of Newcastle, Newcastle, Australia. ⁹⁴ School of Electrical Engineering and Computer Science, University of Newcastle, Newcastle, Australia. ⁹⁵ Division of Medical Genetics, Department of Biomedicine, University of Basel, Basel, Switzerland. ⁹⁶ Department of Genetics, Harvard Medical School, Boston, MA, USA. ⁹⁷ Section of Neonatal Screening and Hormones, Department of Clinical Biochemistry, Immunology and Genetics, Statens Serum Institut, Copenhagen, Denmark. ⁹⁸ Department of Psychiatry, Fujita Health University School of Medicine, Toyoake, Aichi, Japan. ⁹⁹ Regional Centre for Clinical Research in Psychosis, Department of Psychiatry, Stavanger University Hospital, Stavanger, Norway. ¹⁰⁰ Rheumatology Research Group, Vall d'Hebron Research Institute, Barcelona, Spain. ¹⁰¹ Department of Psychiatry, Rudolf Magnus Institute of Neuroscience, University Medical Center Utrecht, Utrecht, The Netherlands. ¹⁰² Centre for Medical Research, The University of Western Australia, Perth, Western Australia, Australia. ¹⁰³ Perkins Institute for Medical Research, The University of Western Australia, Perth, Western Australia, Australia. ¹⁰⁴ Department of Medical Genetics, Medical University, Sofia, Bulgaria. ¹⁰⁵ Department of Psychology, University of Colorado Boulder, Boulder, CO, USA. ¹⁰⁶ Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, Ontario, Canada. ¹⁰⁷ Department of Psychiatry, University of Toronto, Toronto, Ontario, Canada. ¹⁰⁸

Institute of Medical Science, University of Toronto, Toronto, Ontario, Canada. ¹⁰⁹ Institute of Molecular Genetics, Russian Academy of Sciences, Moscow, Russia. ¹¹⁰ Latvian Biomedical Research and Study Centre, Riga, Latvia. ¹¹¹ Department of Psychiatry and Zilkha Neurogenetics Institute, Keck School of Medicine at University of Southern California, Los Angeles, CA, USA. ¹¹² Faculty of Medicine, Vilnius University, Vilnius, Lithuania. ¹¹³ 2nd Faculty of Medicine and University Hospital Motol, Prague, Czech Republic. ¹¹⁴ Department of Biology and Medical Genetics, Charles University Prague, Prague, Czech Republic. ¹¹⁵ Pierre and Marie Curie Faculty of Medicine, Paris, France. ¹¹⁶ Duke-NUS Graduate Medical School, Singapore. ¹¹⁷ Department of Psychiatry, Hadassah-Hebrew University Medical Center, Jerusalem, Israel. ¹¹⁸ Centre for Genomic Sciences and Department of Psychiatry, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, PR China. ¹¹⁹ Mental Health Centre and Psychiatric Laboratory, West China Hospital, Sichuan University, Chendu, Sichuan, PR China. ¹²⁰ Department of Biostatistics, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA. ¹²¹ Department of Psychiatry, Columbia University, New York, NY, USA. ¹²² Priority Centre for Translational Neuroscience and Mental Health, University of Newcastle, Newcastle, Australia. ¹²³ Department of Genetics and Pathology, International Hereditary Cancer Center, Pomeranian Medical University in Szczecin, Szczecin, Poland. ¹²⁴ Department of Mental Health and Substance Abuse Services; National Institute for Health and Welfare, Helsinki, Finland. ¹²⁵ Department of Mental Health, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA. ¹²⁶ Department of Psychiatry, University of Bonn, Bonn, Germany. ¹²⁷ Centre National de la Recherche Scientifique, Laboratoire de Génétique Moléculaire de la Neurotransmission et des Processus Neurodégénératifs, Hôpital de la Pitié Salpêtrière, Paris, France. ¹²⁸ Department of Genomics Mathematics, University of Bonn, Bonn, Germany. ¹²⁹ Research Unit, Sørlandet Hospital, Kristiansand, Norway. ¹³⁰ Department of Psychiatry, Harvard Medical School, Boston, MA, USA. ¹³¹ VA Boston Health Care System, Brockton, MA, USA. ¹³² Department of Psychiatry, National University of Ireland Galway, Ireland. ¹³³ Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, UK. ¹³⁴ Division of Psychiatry, University of Edinburgh, Edinburgh, UK. ¹³⁵ Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway. ¹³⁶ Massachusetts Mental Health Center Public Psychiatry Division of the Beth Israel Deaconess Medical Center, Boston, MA, USA. ¹³⁷ Estonian Genome Center, University of Tartu, Tartu, Estonia. ¹³⁸

School of Psychology, University of Newcastle, Newcastle, Australia. ¹³⁹ First Psychiatric Clinic, Medical University, Sofia, Bulgaria. ¹⁴⁰ Department P, Aarhus University Hospital, Risskov, Denmark. ¹⁴¹ Department of Psychiatry, Royal College of Surgeons in Ireland, Ireland. ¹⁴² King's College London, UK. ¹⁴³ Maastricht University Medical Centre, South Limburg Mental Health Research and Teaching Network, EURON, Maastricht, The Netherlands. ¹⁴⁴ Institute of Translational Medicine, University Liverpool, UK. ¹⁴⁵ Max Planck Institute of Psychiatry, Munich, Germany. ¹⁴⁶ Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ¹⁴⁷ Department of Psychiatry and Psychotherapy, Jena University Hospital, Jena, Germany. ¹⁴⁸ Department of Psychiatry, Queensland Brain Institute and Queensland Centre for Mental Health Research, University of Queensland, Brisbane, Queensland, Australia. ¹⁴⁹ Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA. ¹⁵⁰ Department of Psychiatry, Trinity College Dublin, Ireland. ¹⁵¹ Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN, USA. ¹⁵² Department of Clinical Sciences, Psychiatry, Umeå University, Umeå, Sweden. ¹⁵³ DETECT Early Intervention Service for Psychosis, Blackrock, Dublin, Ireland. ¹⁵⁴ Centre for Public Health, Institute of Clinical Sciences, Queens University Belfast, Belfast, UK. ¹⁵⁵ Lawrence Berkeley National Laboratory, University of California at Berkeley, Berkeley, CA, USA. ¹⁵⁶ Institute of Psychiatry at King's College London, London, UK. ¹⁵⁷ Melbourne Neuropsychiatry Centre, University of Melbourne & Melbourne Health, Melbourne, Australia. ¹⁵⁸ Department of Psychiatry, University of Helsinki, Finland. ¹⁵⁹ Public Health Genomics Unit, National Institute for Health and Welfare, Helsinki, Finland. ¹⁶⁰ Medical Faculty, University of Belgrade, Belgrade, Serbia. ¹⁶¹ Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA. ¹⁶² Institute for Molecular Medicine Finland, FIMM, Helsinki, Finland. ¹⁶³ Department of Child and Adolescent Psychiatry, Erasmus University Medical Centre, Rotterdam, The Netherlands. ¹⁶⁴ Department of Complex Trait Genetics, Neuroscience Campus Amsterdam, VU University Medical Center Amsterdam, Amsterdam, The Netherlands. ¹⁶⁵ Department of Functional Genomics, Center for Neurogenomics and Cognitive Research, Neuroscience Campus Amsterdam, VU University, Amsterdam, The Netherlands. ¹⁶⁶ Department of Epidemiology, Harvard University, Boston, MA, USA. ¹⁶⁷ Department of Psychiatry, University of Oxford, Oxford, UK. ¹⁶⁸ Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, Richmond, VA, USA. ¹⁶⁹ Institute for Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, NY,

USA. ¹⁷⁰ PharmaTherapeutics Clinical Research, Pfizer Worldwide Research and Development, Cambridge, MA, USA. ¹⁷¹ Department of Psychiatry and Psychotherapy, University of Gottingen, Göttingen, Germany. ¹⁷² Psychiatry and Psychotherapy Clinic, University of Erlangen, Germany. ¹⁷³ Hunter New England Health Service, Newcastle, Australia. ¹⁷⁴ School of Biomedical Sciences, University of Newcastle, Newcastle, Australia. ¹⁷⁵ Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA. ¹⁷⁶ University of Iceland, Landspítali, National University Hospital, Reykjavik, Iceland. ¹⁷⁷ Department of Psychiatry and Drug Addiction, Tbilisi State Medical University (TSMU), Tbilisi, Georgia. ¹⁷⁸ Research and Development, Bronx Veterans Affairs Medical Center, New York, NY, USA. ¹⁷⁹ Wellcome Trust Centre for Human Genetics, Oxford, UK. ¹⁸⁰ deCODE Genetics, Reykjavik, Iceland. ¹⁸¹ Department of Clinical Neurology, Medical University of Vienna, Austria. ¹⁸² Lieber Institute for Brain Development, Baltimore, MD, USA. ¹⁸³ Department of Medical Genetics, University Medical Centre, Utrecht, The Netherlands. ¹⁸⁴ Rudolf Magnus Institute of Neuroscience, University Medical Centre Utrecht, The Netherlands. ¹⁸⁵ Berkshire Healthcare NHS Foundation Trust, Bracknell, UK. ¹⁸⁶ Section of Psychiatry, University of Verona, Verona, Italy. ¹⁸⁷ Department of Psychiatry, University of Oulu, Finland. ¹⁸⁸ Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, Dublin, Ireland. ¹⁸⁹ Health Research Board, Dublin, Ireland. ¹⁹⁰ Department of Psychiatry, University Medical Center Groningen, University of Groningen, The Netherlands. ¹⁹¹ Department of Psychiatry and Clinical Neurosciences, School of Psychiatry and Clinical Neurosciences, Queen Elizabeth II Medical Centre, Perth, Western Australia, Australia. ¹⁹² Department of Psychological Medicine and Neurology, MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Cardiff, Wales, UK. ¹⁹³ Computational Sciences CoE, Pfizer Worldwide Research and Development, Cambridge, MA, USA. ¹⁹⁴ Human Genetics, Genome Institute of Singapore, A*STAR, Singapore. ¹⁹⁵ WTCCC2. ¹⁹⁶ University College London, UK. ¹⁹⁷ Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ¹⁹⁸ Institute of Neuroscience and Medicine (INM-1), Research Center Juelich, Juelich, Germany. ¹⁹⁹ Department of Genetics, The Hebrew University of Jerusalem, Jerusalem, Israel. ²⁰⁰ Neuroscience Discovery and Translational Area, Pharma Research and Early Development, F.Hoffman-La Roche, Basel, Switzerland. ²⁰¹ School of Psychiatry and Clinical Neurosciences, The University of Western Australia, Perth, Australia. ²⁰² The Perkins Institute of Medical Research, Perth, Australia. ²⁰³

UWA Centre for Clinical Research in Neuropsychiatry.²⁰⁴ Virginia Institute for Psychiatric and Behavioral Genetics, Departments of Psychiatry and Human and Molecular Genetics, Virginia Commonwealth University, Richmond, VA, USA.²⁰⁵ The Feinstein Institute for Medical Research, Manhasset, NY, USA.²⁰⁶ The Hofstra NS-LIJ School of Medicine, Hempstead, NY, USA.²⁰⁷ The Zucker Hillside Hospital, Glen Oaks, NY, USA.²⁰⁸ Saw Swee Hock School of Public Health, National University of Singapore, Singapore.²⁰⁹ Queensland Centre for Mental Health Research, University of Queensland, Brisbane, Queensland, Australia.²¹⁰ Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht, The Netherlands.²¹¹ PEIC.²¹² The Broad Institute of MIT and Harvard, Cambridge, MA, USA.²¹³ Center for Human Genetic Research and Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA.²¹⁴ Centre for Genomic Sciences, State Key Laboratory for Brain and Cognitive Sciences, and Department of Psychiatry, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, PR China.²¹⁵ University of Aberdeen, Institute of Medical Sciences, Aberdeen, Scotland, UK.²¹⁶ Departments of Psychiatry, Neurology, Neuroscience and Institute of Genetic Medicine, Johns Hopkins School of Medicine, Baltimore, MD, USA.²¹⁷ Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.

Wellcome Trust Case-Control Consortium 2

Management Committee: Peter Donnelly^{179,218}, Ines Barroso²¹⁹, Jenefer M Blackwell^{220,221}, Elvira Bramon¹⁹⁶, Matthew A Brown²²², Juan P Casas^{223,224}, Aiden Corvin⁵, Panos Deloukas²¹⁹, Audrey Duncanson²²⁵, Janusz Jankowski²²⁶, Hugh S Markus²²⁷, Christopher G Mathew²²⁸, Colin N A Palmer²²⁹, Robert Plomin⁹, Anna Rautanen¹⁷⁹, Stephen J Sawcer²³⁰, Richard C Trembath²²⁸, Ananth C Viswanathan^{231,232}, Nicholas W Wood²³³.

Data and Analysis Group: Chris C A Spencer¹⁷⁹, Gavin Band¹⁷⁹, Céline Bellenguez¹⁷⁹, Peter Donnelly^{179,218}, Colin Freeman¹⁷⁹, Eleni Giannoulitou¹⁷⁹, Garrett Hellenthal¹⁷⁹, Richard Pearson¹⁷⁹, Matti Pirinen¹⁷⁹, Amy Strange¹⁷⁹, Zhan Su¹⁷⁹, Damjan Vukcevic¹⁷⁹.

DNA, Genotyping, Data QC, and Informatics: Cordelia Langford²¹⁹, Ines Barroso²¹⁹, Hannah Blackburn

²¹⁹, Suzannah J Bumpstead ²¹⁹, Panos Deloukas ²¹⁹, Serge Dronov ²¹⁹, Sarah Edkins ²¹⁹, Matthew Gillman ²¹⁹, Emma Gray ²¹⁹, Rhian Gwilliam ²¹⁹, Naomi Hammond ²¹⁹, Sarah E Hunt ²¹⁹, Alagurevathi Jayakumar ²¹⁹, Jennifer Liddle ²¹⁹, Owen T McCann ²¹⁹, Simon C Potter ²¹⁹, Radhi Ravindrarajah ²¹⁹, Michelle Ricketts ²¹⁹, Avazeh Tashakkori-Ghanbaria ²¹⁹, Matthew Waller ²¹⁹, Paul Weston ²¹⁹, Pamela Whittaker ²¹⁹, Sara Widaa ²¹⁹.

Publications Committee: Christopher G Mathew ²²⁸, Jenefer M Blackwell ^{220,221}, Matthew A Brown ²²², Aiden Corvin ⁵, Mark I McCarthy ²³⁴, Chris C A Spencer ¹⁷⁹.

Psychosis Endophenotype International Consortium

Maria J Arranz ^{156,235}, Steven Bakker ¹⁰¹, Stephan Bender ^{236,237}, Elvira Bramon ^{156,238,239}, David A Collier ^{8,9}, Benedicto Crespo-Facorro ^{240,241}, Jeremy Hall ¹³⁴, Conrad Iyegbe ¹⁵⁶, Assen V Jablensky ²⁴², René S Kahn ¹⁰¹, Luba Kalaydjieva ^{102,243}, Stephen Lawrie ¹³⁴, Cathryn M Lewis ¹⁵⁶, Kuang Lin ¹⁵⁶, Don H Linszen ²⁴⁴, Ignacio Mata ^{240,241}, Andrew M McIntosh ¹³⁴, Robin M Murray ¹⁴², Roel A Ophoff ⁸⁰, Jim Van Os ^{143,156}, John Powell ¹⁵⁶, Dan Rujescu ^{81,83}, Muriel Walshe ¹⁵⁶, Matthias Weisbrod ²³⁷, Durk Wiersma ¹⁹⁰.

²¹⁸ Department of Statistics, University of Oxford, Oxford, UK. ²¹⁹ Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK. ²²⁰ Cambridge Institute for Medical Research, University of Cambridge School of Clinical Medicine, Cambridge, UK. ²²¹ Telethon Institute for Child Health Research, Centre for Child Health Research, University of Western Australia, Subiaco, Western Australia, Australia. ²²² Diamantina Institute of Cancer, Immunology and Metabolic Medicine, Princess Alexandra Hospital, University of Queensland, Brisbane, Queensland, Australia. ²²³ Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK. ²²⁴ Department of Epidemiology and Public Health, University College London, London, UK. ²²⁵ Molecular and Physiological Sciences, The Wellcome Trust, London, UK. ²²⁶ Peninsula School of Medicine and Dentistry, Plymouth University, Plymouth, UK. ²²⁷ Clinical Neurosciences, St George's University of London, London, UK. ²²⁸ Department of Medical and Molecular Genetics, School of Medicine, King's College London, Guy's Hospital, London, UK. ²²⁹ Biomedical Research Centre, Ninewells Hospital and

Medical School, Dundee, UK. ²³⁰ Department of Clinical Neurosciences, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK. ²³¹ Institute of Ophthalmology, University College London, London, UK. ²³² National Institute for Health Research, Biomedical Research Centre at Moorfields Eye Hospital, National Health Service Foundation Trust, London, UK. ²³³ Department of Molecular Neuroscience, Institute of Neurology, London, UK. ²³⁴ Oxford Centre for Diabetes, Endocrinology and Metabolism, Churchill Hospital, Oxford, UK. ²³⁵ Fundació de Docència i Recerca Mútua de Terrassa, Universitat de Barcelona, Spain. ²³⁶ Child and Adolescent Psychiatry, University of Technology Dresden, Dresden, Germany. ²³⁷ Section for Experimental Psychopathology, General Psychiatry, Heidelberg, Germany. ²³⁸ Institute of Cognitive Neuroscience, University College London, London, UK. ²³⁹ Mental Health Sciences Unit, University College London, London, UK. ²⁴⁰ Centro Investigación Biomédica en Red Salud Mental, Madrid, Spain. ²⁴¹ University Hospital Marqués de Valdecilla, Instituto de Formación e Investigación Marqués de Valdecilla, University of Cantabria, Santander, Spain. ²⁴² Centre for Clinical Research in Neuropsychiatry, The University of Western Australia, Perth, Western Australia, Australia. ²⁴³ Western Australian Institute for Medical Research, The University of Western Australia, Perth, Western Australia, Australia. ²⁴⁴ Department of Psychiatry, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

References

1. Schizophrenia Working Group of the Psychiatric Genomics Consortium: Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 2014; 511:421–427
2. Stefansson H, Ophoff RA, Steinberg S, Andreassen OA, Cichon S, Rujescu D, Werge T, Pietiläinen OPH, Mors O, Mortensen PB, Sigurdsson E, Gustafsson O, Nyegaard M, Tuulio-Henriksson A, Ingason A, Hansen T, Suvisaari J, Lonnqvist J, Paunio T, Børghlum AD, Hartmann A, Fink-Jensen A, Nordentoft M, Hougaard D, Norgaard-Pedersen B, Böttcher Y, Olesen J, Breuer R, Möller H-J, Giegling I, Rasmussen HB, Timm S, Mattheisen M, Bitter I, Réthelyi JM, Magnusdottir BB, Sigmundsson T, Olason P, Masson G, Gulcher JR, Haraldsson M, Fossdal R, Thorgeirsson TE, Thorsteinsdottir U, Ruggeri M, Tosato S, Franke B, Strengman E, Kiemenev LA, Genetic Risk and Outcome in Psychosis (GROUP), Melle I, Djurovic S, Abramova L, Kaleda V, Sanjuan J, de Frutos R, Bramon E, Vassos E, Fraser G, Ettinger U, Picchioni M, Walker N, Touloupoulou T, Need AC, Ge D, Yoon JL, Shianna KV, Freimer NB, Cantor RM, Murray R, Kong A, Golimbet V, Carracedo

- A, Arango C, Costas J, Jönsson EG, Terenius L, Agartz I, Petursson H, Nöthen MM, Rietschel M, Matthews PM, Muglia P, Peltonen L, St Clair D, Goldstein DB, Stefansson K, Collier DA: Common variants conferring risk of schizophrenia. *Nature* 2009; 460:744–747
3. Krawczak M, Nikolaus S, von Eberstein H, Croucher PJP, El Mokhtari NE, Schreiber S: PopGen: population-based recruitment of patients and controls for the analysis of complex genotype-phenotype relationships. *Community Genet.* 2006; 9:55–61
 4. Wichmann H-E, Gieger C, Illig T, for the MONICA/KORA Study Group: KORA-gen - Resource for Population Genetics, Controls and a Broad Spectrum of Disease Phenotypes. *Das Gesundheitswesen* 2005; 67:26–30[cited 2016 Oct 31]
 5. Schmermund A, Möhlenkamp S, Stang A, Grönemeyer D, Seibel R, Hirche H, Mann K, Siffert W, Lauterbach K, Siegrist J, Jöckel K-H, Erbel R: Assessment of clinically silent atherosclerotic disease and established and novel risk factors for predicting myocardial infarction and cardiac death in healthy middle-aged subjects: rationale and design of the Heinz Nixdorf RECALL Study. *Risk Factors, Evaluation of Coronary Calcium and Lifestyle. Am. Heart J.* 2002; 144:212–218
 6. Carroll LS, Williams HJ, Walters J, Kirov G, O'Donovan MC, Owen MJ: Mutation screening of the 3q29 microdeletion syndrome candidate genes DLG1 and PAK2 in schizophrenia. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 2011; 156B:844–849
 7. Wing JK, Babor T, Brugha T, Burke J, Cooper JE, Giel R, Jablenski A, Regier D, Sartorius N: SCAN: Schedules for Clinical Assessment in Neuropsychiatry. *Arch. Gen. Psychiatry* 1990; 47:589–593
 8. Association, American Psychiatric: Diagnostic and statistical manual of mental disorders. 4th ed. Washington, D.C.: American Psychiatric Association; 1994.
 9. Metspalu A, Köhler F, Laschinski G, Ganten D, Roots I: [The Estonian Genome Project in the context of European genome research]. *Dtsch. Med. Wochenschr.* 2004; 129 Suppl 1:S25–8
 10. Egan MF, Goldberg TE, Kolachana BS, Callicott JH, Mazzanti CM, Straub RE, Goldman D, Weinberger DR: Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia. *Proc. Natl. Acad. Sci. U. S. A.* 2001; 98:6917–6922
 11. Shi J, Levinson DF, Duan J, Sanders AR, Zheng Y, Pe'er I, Dudbridge F, Holmans PA, Whittemore AS, Mowry BJ, Olincy A, Amin F, Cloninger CR, Silverman JM, Buccola NG, Byerley WF, Black DW, Crowe RR, Oksenberg JR, Mirel DB, Kendler KS, Freedman R, Gejman PV: Common variants on chromosome 6p22.1 are associated with schizophrenia. *Nature* 2009; 460:753–757
 12. Sanders AR, Levinson DF, Duan J, Dennis JM, Li R, Kendler KS, Rice JP, Shi J, Mowry BJ, Amin F, Silverman JM, Buccola NG, Byerley WF, Black DW, Freedman R, Cloninger CR, Gejman PV: The Internet-based MGS2 control sample: self report of mental illness. *Am. J. Psychiatry* 2010; 167:854–865
 13. Athanasiu L, Mattingsdal M, Kähler AK, Brown A, Gustafsson O, Agartz I, Giegling I, Muglia P, Cichon S, Rietschel M, Pietiläinen OPH, Peltonen L, Bramon E, Collier D, Clair DS, Sigurdsson E, Petursson H, Rujescu D, Melle I, Steen VM, Djurovic S, Andreassen OA:

Gene variants associated with schizophrenia in a Norwegian genome-wide study are replicated in a large European cohort. *J. Psychiatr. Res.* 2010; 44:748–753

14. Ikeda M, Aleksic B, Kinoshita Y, Okochi T, Kawashima K, Kushima I, Ito Y, Nakamura Y, Kishi T, Okumura T, Fukuo Y, Williams HJ, Hamshere ML, Ivanov D, Inada T, Suzuki M, Hashimoto R, Ujike H, Takeda M, Craddock N, Kaibuchi K, Owen MJ, Ozaki N, O'Donovan MC, Iwata N: Genome-wide association study of schizophrenia in a Japanese population. *Biol. Psychiatry* 2011; 69:472–478
15. Glessner JT, Reilly MP, Kim CE, Takahashi N, Albano A, Hou C, Bradfield JP, Zhang H, Sleiman PMA, Flory JH, Imielinski M, Frackelton EC, Chiavacci R, Thomas KA, Garriss M, Otieno FG, Davidson M, Weiser M, Reichenberg A, Davis KL, Friedman JI, Cappola TP, Margulies KB, Rader DJ, Grant SFA, Buxbaum JD, Gur RE, Hakonarson H: Strong synaptic transmission impact by copy number variations in schizophrenia. *Proc. Natl. Acad. Sci. U. S. A.* 2010; 107:10584–10589
16. Wong EHM, So H-C, Li M, Wang Q, Butler AW, Paul B, Wu H-M, Hui TCK, Choi S-C, So M-T, Garcia-Barcelo M-M, McAlonan GM, Chen EYH, Cheung EFC, Chan RCK, Purcell SM, Cherny SS, Chen RRL, Li T, Sham P-C: Common variants on Xq28 conferring risk of schizophrenia in Han Chinese. *Schizophr. Bull.* 2014; 40:777–786
17. Kung AWC, Xiao S-M, Cherny S, Li GHY, Gao Y, Tso G, Lau KS, Luk KDK, Liu J-M, Cui B, Zhang M-J, Zhang Z-L, He J-W, Yue H, Xia W-B, Luo L-M, He S-L, Kiel DP, Karasik D, Hsu Y-H, Cupples LA, Demissie S, Styrkarsdottir U, Halldorsson BV, Sigurdsson G, Thorsteinsdottir U, Stefansson K, Richards JB, Zhai G, Soranzo N, Valdes A, Spector TD, Sham PC: Association of JAG1 with bone mineral density and osteoporotic fractures: a genome-wide association study and follow-up replication studies. *Am. J. Hum. Genet.* 2010; 86:229–239
18. Guo Y, Tomlinson B, Chu T, Fang YJ, Gui H, Tang CS, Yip BH, Cherny SS, Hur Y-M, Sham PC, Lam TH, Thomas NG: A genome-wide linkage and association scan reveals novel loci for hypertension and blood pressure traits. *PLoS One* 2012; 7:e31489
19. Chan KY-K, Wong C-M, Kwan JS-H, Lee JM-F, Cheung KW, Yuen MF, Lai CL, Poon RT-P, Sham PC, Ng IO-L: Genome-wide association study of hepatocellular carcinoma in Southern Chinese patients with chronic hepatitis B virus infection. *PLoS One* 2011; 6:e28798
20. Tobacco and Genetics Consortium: Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nat. Genet.* 2010; 42:441–447
21. Skol AD, Scott LJ, Abecasis GR, Boehnke M: Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat. Genet.* 2006; 38:209–213
22. Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, Boehnke M, Abecasis GR, Willer CJ: LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* 2010; 26:2336–2337

Table 1. Conceptual overview of analyses of schizophrenia and smoking behaviors. PGC is Psychiatric Genomics Consortium, TAG is Tobacco and Genetics consortium, LD is linkage disequilibrium, SNP is single nucleotide polymorphism.

Research Question	Cohorts & Sample Sizes	Analysis
Question 1: Are there genetic correlations between schizophrenia and smoking behaviors?	<i>Primary</i> - PGC-Schizophrenia European ancestry: 35,476 cases, 46,839 controls; TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181, cessation 41,278, age of smoking onset 24,114	genetic correlation (LD score regression)
	<i>Replication</i> - PGC-Schizophrenia East-Asian ancestry: 1,836 cases, 3,383 controls	trans-ethnic genetic correlation (popcorn)
Question 2: Do polygenic risk scores for smoking behaviors also predict schizophrenia case status?	<i>Training Set</i> - TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181, cessation 41,278, age of smoking onset 24,114	polygenic risk scores (cross-trait association)
	<i>Testing Set</i> - PGC-Schizophrenia: 35,476 cases, 46,839 controls	
Question 3: What is the genetic architecture of smoking behavior among schizophrenia patients?		
3.1 What is the SNP-based heritability of smoking behaviors among schizophrenia cases?	PGC-Schizophrenia Phenotype Working Group - 10 study sites: smoking initiation 5,255, cigarettes-per-day 3,370	SNP-based heritability (LD score regression)
3.2 Are genetic factors for smoking behaviors shared between populations with and without schizophrenia?	PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370; TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181	genetic correlation (LD score regression)
3.3 Do polygenic risk scores for smoking behaviors also predict these behaviors in schizophrenia patients?	<i>Training Set</i> - TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181	polygenic risk scores (within-trait across-cohort association)
	<i>Testing Set</i> - PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370	
3.4 Are there schizophrenia-specific genetic risk variants for smoking behaviors?	<i>Primary</i> - PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370	genome-wide association study meta-analysis of smoking behaviors among schizophrenia cases
	Replication - Janssen Pharmaceuticals: smoking initiation 1802, cigarettes-per-day 1802	

Question 4: Are there associations between smoking behaviors and clinical features of schizophrenia?		
4.1 Are smoking behaviors among schizophrenia cases associated with clinical presentation of schizophrenia?	PGC-Schizophrenia: smoking initiation 5,255, cigarettes-per-day 3,370, age-schizophrenia-onset 4,658; symptom dimension factor scores: positive 3,846, negative 3,845, manic 3,740, depression 3,740	phenotypic associations (linear/logistic regression)
4.2 Do polygenic risk scores for smoking behaviors predict schizophrenia symptom dimensions?	<i>Training Set</i> - TAG-Smoking behaviors: smoking initiation 74,035, cigarettes-per-day 38,181 <i>Testing Set</i> - PGC-Schizophrenia: age-schizophrenia-onset 11,600; symptom dimension factor scores: positive 8,330, negative 8,427, manic 6,965, depression 6,964	polygenic risk scores (across-trait across-cohort association)

Tables

Table 2. Sample characteristics for each PGC2-SCZ cohort.

Cohort	<i>N</i> SCZ-cases	<i>Sex</i> % <i>female</i>	<i>Age</i> (mean)	<i>N</i> % SI	<i>N</i> CPD
boco	756	43.5%	36.7	538 77.3%	185
cims	37	16.2%	33.3	33 66.7%	23
cou3	513	39.2%	44.2	501 75.0%	278
egcu	234	26.9%	46.5	234 52.6%	123
lie2	133	28.6%	36.9	128 60.2%	69
lie5	497	25.7%	36.6	479 63.3%	285
mgs2	2348	30.5%	43.5	2227 78.8%	1674
munc	420	36.4%	37.9	406 74.4%	301
top8	344	43.6%	33.0	320 57.5%	176
ucla	450	23.8%	34.7	389 70.7%	256
TOTAL	6183	6183	6132	5255	3370

Note: PGC = Psychiatric Genomics Consortium, SCZ - schizophrenia, Age = age at assessment, SI = smoking initiation, CPD = cigarettes per day.

PGC GWAS of smoking behaviors in schizophrenia 2

Table 3. Genetic correlations between TAG and PGC-SCZ phenotypes.

Trait 1	Discovery			Replication		
	Trait 2	r_g (se)	P	Trait 2	r_g (se)	P'
SI _{TAG}	SCZ _{EUR}	0.159 (0.026)	5.05×10^{-10}	SCZ _{EAS}	0.040 (0.124)	0.744
CPD _{TAG}	SCZ _{EUR}	0.094 (0.034)	0.006	SCZ _{EAS}	-0.080 (0.161)	0.619
logOnset _{TAG}	SCZ _{EUR}	0.100 (0.038)	0.009	SCZ _{EAS}	0.463 (0.251)	0.064
Cessation _{TAG}	SCZ _{EUR}	-0.076 (0.035)	0.032	SCZ _{EAS}	-0.051 (0.193)	0.793
SI _{TAG}	SI _{SCZ, EUR}	0.624 (0.202)	0.002	.	.	.
CPD _{TAG}	CPD _{SCZ, EUR}	0.689 (0.443)	0.120	.	.	.

For each pair of traits, r_g is the estimated genetic correlation; P is the significance of $r_g \neq 0$; P' is a 1-sided test of whether $r_g > 0$ or $r_g < 0$ in the replication sample. For comparisons of TAG phenotypes to SCZ risk, *EUR* and *EAS* denote European and East-Asian cohorts.

PGC GWAS of smoking behaviors in schizophrenia 3

Table 4. Association results for top SNP associations.

Trait	Chr	SNP	A1/ A2	Discovery			Replication Phase		Combined P	Gene (+/-Kb)
				Frq Info	Z	P n	Z	P n		
SI	1	rs58215884	T/G	0.162 0.975	4.53	5.9×10 ⁻⁶ 4991	2.09	0.037 1776	6.8×10 ⁻⁷	FCRL2 (+7.2)
	5	rs1592907	A/G	0.454 0.983	-4.44	9.2×10 ⁻⁶ 4991	-2.19	0.029 1760	8.5×10 ⁻⁷	FBXL17 (+91.2)
	9	rs117381175	T/C	0.031 0.792	-4.55	5.3×10 ⁻⁶ 4991	-2.29	0.022 1746	6.9×10 ⁻⁷	intergenic
	13	rs754168	A/C	0.352 0.767	-4.85	1.2×10 ⁻⁶ 4991	-1.57	0.116 1277	4.6×10 ⁻⁷	LINC01044 (0)
CPD	1	rs3896119	A/G	0.018 0.832	4.77	1.9×10 ⁻⁶ 3321	1.45	0.148 1078	5.0×10 ⁻⁷	intergenic
	1	rs1210	T/C	0.028 0.745	5.06	4.2×10 ⁻⁷ 3321	2.11	0.034 1052	5.2×10 ⁻⁸	RGS8 (+4.7)
	2	rs1900325	T/C	0.479 0.924	5.32	1.1×10 ⁻⁷ 3344	0.18	0.854 1011	2.0×10 ⁻⁶	CBWD2 (-39.4)
	3	rs833663	A/G	0.038 0.982	4.58	4.7×10 ⁻⁶ 3344	2.07	0.038 1097	4.7×10 ⁻⁷	CADPS (0)
	7	rs148253479	A/C	0.988 0.694	-4.89	1.0×10 ⁻⁶ 2442	-2.54	0.011 1078	3.2×10 ⁻⁸	TMEM106B (-22.6)

SI is smoking initiation, CPD is cigarettes-per day, SNP and Chr information for build hg19; INFO is the statistical imputation information; Frq is the frequency of the reference (first listed) allele, and Z is its estimated standardized effect; P is the P-value for association; n is the sample size. The nearest gene within 100Kb is shown; its position relative to a gene is given parenthetically and with respect to direction of transcription (negative and positive kb values indicate up- and downstream positions).

Table 5. Association of smoking variables with clinical features in SCZ.

	Smoking initiation			Cigarettes per day		
	<i>N</i>	β (SE)	<i>P</i>	<i>N</i>	β (SE)	<i>P</i>
Sex	4991	-0.507 (0.069)	1.38×10^{-13}	3344	-0.210 (0.073)	0.004
Age	4991	-0.005 (0.003)	0.061	3344	0.018 (0.003)	5.53×10^{-10}
Age-of-onset	4658	-0.098 (0.038)	0.009	3168	-0.022 (0.021)	0.289
Positive	3846	0.157 (0.038)	3.23×10^{-5}	2796	0.071 (0.029)	0.015
Negative	3845	0.046 (0.038)	0.225	2794	-0.003 (0.029)	0.929
Mania	3740	0.034 (0.038)	0.367	2736	0.019 (0.019)	0.305
Depression	3740	-0.013 (0.038)	0.728	2735	0.047 (0.019)	0.014

For SCZ Age-of-onset and symptom factor scores, *N* is the number of subjects with non-missing data for both traits; β and SE are the beta regression coefficient and standard error from logistic or linear regression; *P* is the significance of the association between a given pair of traits.