

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/141037/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Gilbert, A., Drinkwater, K., McParland, L., Adams, R. , Glynn-Jones, R., Harrison, M., Hawkins, M. A., Sebag-Montefiore, D., Gilbert, D. C. and Muirhead, R. 2020. UK national cohort of anal cancer treated with intensity-modulated radiotherapy: one-year oncological and patient-reported outcomes. *European Journal of Cancer* 128 , pp. 7-16. 10.1016/j.ejca.2019.12.022

Publishers page: <http://dx.doi.org/10.1016/j.ejca.2019.12.022>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Supplementary data




Figure 1: Flowchart clarifying data included in publication

CORMAC Core Outcome Set outcomes by Early and Advanced risk groups




Figure 2: Overall QOL and function by Early vs Advanced cancer grouping




Figure 3: Cancer symptoms by Early vs Advanced cancer grouping




Figure 4: Bowel Symptoms by Early vs Advanced cancer grouping




Figure 5: Sexual symptoms by Early vs Advanced cancer grouping




Figure 6: Libido by Early vs Advanced cancer grouping