

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/141395/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Williams, O., Williams, C., Turner, D., Bull, M., Watkins, J. and Hurt, L. 2022. An epidemiological investigation of COVID-19 outbreaks in a group of care homes in Wales, UK: a retrospective cohort study. *Journal of Public Health* 44 (3) , pp. 606-613. [10.1093/pubmed/fdab150](https://doi.org/10.1093/pubmed/fdab150)

Publishers page: <https://doi.org/10.1093/pubmed/fdab150>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Title Page

Title of paper:

An epidemiological investigation of COVID-19 outbreaks in a group of care homes in Wales, UK: a retrospective cohort study

Contributing authors:

Mr Williams, O. (Public Health Wales, Cardiff, Wales, CF10 4BZ)

Dr Williams, C. (Public Health Wales, Cardiff, Wales, CF10 4BZ)

Mr Turner, D. (Public Health Wales, Cardiff, Wales, CF10 4BZ)

Mr Bull, M. (Public Health Wales, Cardiff, Wales, CF10 4BZ)

Ms Watkins, J. (Public Health Wales, Cardiff, Wales, CF10 4BZ)

Dr Hurt, L. (Cardiff University, Cardiff, Wales CF14 4YS)

Corresponding author:

Oliver Williams, 116 Westward Rise, Barry, CF62 6PQ. Oli_williams86@hotmail.com

Declarations:

None.

Acknowledgements:

We would like to acknowledge the time and information provided by the care homes whilst collecting data for this study. The fantastic staff and management provided all the information requested whilst being extremely busy in a difficult time. Thank you.

Abstract

Background

This study describes the epidemiology of COVID-19 outbreaks in four care homes in terms of spread, severity, presentation, and interventions.

Methods

Participants were 100 residents and 102 staff from 4 co-located care homes in Wales. Data were collected from the homes and Public Health Wales, including demographics, presentations, test status and results, hospital admissions and deaths. Genomic sequencing of confirmed case samples was completed, where possible. Epi-curves, crude attack rates, a Kaplan Meier survival curve and adjusted hazard ratios were calculated using R.

Results

14 confirmed and 43 possible resident cases, 23 confirmed and 47 possible staff cases occurred. Crude attack rates of possible and confirmed cases were 57% (residents) and 69% (staff). Genomic sequencing for 10 confirmed case PCR samples identified at least 5 different UK lineages of COVID-19. 42 (42%) residents died, 23 (55%) with COVID-19 or suspected COVID-19 recorded on the death certificate. The hazard ratio for death amongst resident possible and confirmed cases compared to null cases, adjusting for age and sex, was 13.26 (95% CI 5.61 to 31.34).

Conclusions

There were extensive outbreaks of COVID-19 in these homes with high crude attack rates and deaths. Universal testing and early isolation of residents are recommended.

Introduction

Background

Care homes provide accommodation to individuals needing significant help with personal care [1], with approximately 416,000 people living in care homes in the UK [2]. Care home residents live in close proximity to each other, which along with the typical older ages and common chronic underlying conditions of residents, means these facilities are at high risk for severe outbreaks of COVID-19 [3,4].

The pandemic has disproportionately impacted care home residents globally, with 19-72% of total COVID-19 deaths being reported in these settings [5]. Care homes in Wales have become hotspots for outbreaks, with significantly increased risks of death amongst residents compared to previous years [6].

Once outbreaks occur in care homes they can spread rapidly through the resident population and staff [4,7,8], leading to high mortality [7,9,10,11,12].

A particular challenge is transmission by asymptomatic individuals, who may or may not go on to be symptomatic [7,9,10,11,12,13]. Ladhani et al (2020) found that 43.8% of confirmed resident cases and 49.1% of confirmed staff cases were asymptomatic [14].

It is important to have more studies in UK settings, featuring both residents and staff and covering the full time-frame of an outbreak, to further understanding of COVID-19 outbreaks in care homes. Limited data are available on staff demographics and presentation, an understanding of which may assist in identification of possible cases for testing. The care homes involved in this study approached Public Health Wales to assist in an epidemiological investigation following high numbers of confirmed cases of COVID-19 and deaths. This study set out to describe the epidemiology of these outbreaks in terms of spread, severity, presenting symptoms and interventions implemented to reduce spread.

Methods

This was a retrospective cohort study, consisting of data for all staff and residents in 4 co-located care homes (Home A, Home B, Home C and Home D) between February and May 2020. Note that Home D is situated within Home C as a separate sealed-off unit.

The study time-frame covered 18 weeks, from 2 weeks prior to symptom onset in the first possible case in the homes (ensuring coverage of incubation periods) until 2 weeks after symptom onset in the final confirmed case.

Participant eligibility criteria:

- Any resident living in one of the homes at any time during the study time-frame.
- Any staff member working in one of the homes at any time during the study time-frame.

Definitions used within the study are as follows:

Possible case: any staff member or resident without a positive viral throat swab test, with onset of new symptoms (fever, cough, shortness of breath or anosmia) within the study time-frame.

Confirmed case: any resident or staff member with a positive viral throat swab test.

Null case: Any resident or staff member not meeting the possible or confirmed case definitions.

Symptom onset date: Date of onset was taken as the earliest date of new symptom onset. Asymptomatic confirmed cases were excluded from epi-curve and cumulative incidence graphs, as it was not possible to determine true onset date.

Data were collected from the Care Home managers, Public Health Wales and the local health board, and included the following:

- ***Test data:*** Prior to May 2020, testing was triggered for those who were symptomatic with symptoms of a cough, fever, shortness of breath or anosmia. From May, all staff and residents were tested as part of regular screening. Staff and residents were tested with viral throat swabs, which were analysed at the University Hospital of Wales' virology laboratory for Coronavirus SARS-CoV-2 RNA, by PCR with a test considered positive if Coronavirus SARS-CoV-2 RNA was found to be present.
- ***Care home data:***
 - Resident and staff demographics: counts, gender, and age (age of residents was taken as their age on 01/04/2020).
 - Presenting symptoms and onset dates, as recorded by the care home managers in their line lists. Residents were monitored daily for symptoms.
 - Location of confirmed and possible resident cases within each home.
 - Hospital admissions: Dates and reasons of hospital admissions were obtained from care home managers and local health board data via Clinical Portal.
 - Deaths: Dates and causes of death, based on death certificate recordings.
 - Staff areas of work: Individuals identified as working in more than one of the study homes were assigned to both for the purpose of descriptive analysis on that particular home, but were not double counted for analysis involving all homes.
 - Onset dates of COVID-19 response interventions: enhanced cleaning, personal protective equipment use and training, isolation of residents, restrictions on shared space and visiting cessation.
- ***Genomic data:*** Genomic sequencing data from available throat swab samples were obtained from Public Health Wales Microbiology department and the Pathogen Genomics Unit (PenGU). Samples were reverse-transcribed and then amplified using the ARTIC v3 primers and protocol [15]. Resulting amplicons were prepared for sequencing using Illumina Nextera XT library preparation kit

and sequencing was performed on the Illumina NextSeq using the NextSeq Mid-output v2.5 sequencing kit (300 cycles).

For each sample, all sequences were quality trimmed (Trim-Galore v0.6.5) and then aligned to the SARS-CoV-2 reference sequence (MN908947.3) using bwa v0.7.17. Once aligned, primer sequences were coordinate-trimmed and consensus fasta sequences generated using iVar (v1.2.2). This process is provided as a Nextflow workflow (<https://github.com/connor-lab/ncov2019-artic-nf>). Consensus fasta sequences were submitted to the COVID-19 Genomics UK Consortium (COG-UK) analysis environment hosted on MRC CLIMB, and each sequence was assigned a global lineage [16]. Alongside this, to facilitate analyses of local outbreaks, a more granular “UK lineage” and “phylotype” were assigned to each sample using a bespoke phylogenetic analysis pipeline (<https://github.com/COG-UK/grapevine/>).

Data Analysis

Data were analysed in R using tidyverse packages, including ggplot2 [17,18,19], to summarise demographic data, produce epidemic-curves, calculate hazard ratios (adjusted for age and sex) for death amongst possible and confirmed resident cases compared to null resident cases, and to generate a Kaplan Meier survival curve. We calculated crude attack rates as the total confirmed and possible cases divided by the relevant population at risk.

Results

Population at risk

The study included 100 residents (41% male and 59% female, age range 66 to 100 years, mean age 82.4 years) and 102 staff members (7% male and 93% female, age range 16 to 69 years, mean age 41 years) (*Table 1*). During the study time-frame 70 (69%) staff and 57 (57%) residents presented with a new symptom.

Testing coverage and results

Testing for COVID-19 occurred for 57 (57%) of residents and 75 (74%) of staff. There were 42 individuals (26 residents and 16 staff) presenting with a typical COVID-19 symptom who were not tested.

Of the 132 individuals with sample results, 46 had both recorded onset dates of symptoms and recorded sample dates. For these, there was a mean time of 20 days from symptom onset date to test date, with a range of -5 to 65 days.

There were 14 (14%) confirmed cases amongst the resident population and 23 (23%) amongst the staff population. Of the 14 residents testing positive, 8 (57%) were male and 6 (43%) were female, with an age range of 66 to 99 years and a mean age of 79.4 years. Of the 23 staff members testing positive, 2 (9%) were male and 21 (91%) were female, with an age range of 19 to 69 years and a mean age of 42.3 years.

Crude attack rates

The crude attack rates were 69% amongst staff and 57% amongst residents across the combined homes.

Table 1 includes a breakdown of the crude attack rates amongst staff and residents for each home.

Presenting symptoms

Recorded presenting symptoms amongst possible and confirmed cases are shown in *Figure 1*. At least one typical COVID-19 symptom (cough, fever, shortness of breath or anosmia) was recorded in 19 (82%) of the 23 confirmed staff cases and 10 (71%) of the 14 confirmed resident cases.

Spread of COVID-19 in the homes

Epidemic curves of all possible and confirmed cases amongst residents and staff by date of onset of initial symptoms for each home are presented in *Figure 2*.

Home A appears to have an initial point source pattern with a single peak in cases on the 13/03/2020, followed by an intermittent source pattern from then until the last confirmed case.

Home B had a couple of isolated cases in March before a sharp rise and peak at the end of March, peaking on the 02/04/2020 when there was an onset of 6 new suspected or confirmed cases in a single day. A continuous source pattern is then seen, suggesting person-to-person spread over the next 2-3 weeks.

Homes C and D have a propagated pattern of transmission with small intervals between small peaks.

Spot maps of possible and confirmed resident cases within each home did not appear to show any spatial clustering over time, with cases being spread throughout the homes rather than localised to specific floors, wings or areas.

Genomic sequencing

Identifiable lineages via genomic sequencing were obtainable from 10 PCR samples (27% of confirmed cases) taken amongst staff and residents at the homes (2 from residents and 8 from staff), with five different COVID-19 lineages identified.

Severity of COVID-19 in the homes

Hospital admissions were required for 4 residents, 3 of which were confirmed cases (21% of confirmed resident cases), and 4 staff members, who were all confirmed cases (17% of all confirmed staff cases).

There were 42 resident deaths across the 4 homes; 42% of the total resident population at risk, based on all-cause mortality. Of these, 91% died in the homes, and 9% died in hospital. In April, 30 residents died; more than twice the number of deaths in a single month than in any other month in the preceding 27 months, and 6 times higher than the monthly average for January 2018 to March 2020.

Home A had the lowest all-cause case fatality proportion at 27% (N=33), Home B had the highest case fatality proportion at 56% (N=34), whilst for Home C it was 42% (N=26) and in Home D it was 43% (N=7). There were no deaths amongst staff members.

Of the 42 resident deaths, 4 (10%) had COVID-19 recorded as the cause of death, 19 (45%) had suspected COVID-19 as the cause of death, and 19 (45%) had other causes of death recorded. Therefore 23 (55%) of the deaths during the study timeframe were given COVID-19 or suspected COVID-19 as cause of death on the death certificates. Of the 42 deaths, 30 (71%) displayed symptoms of COVID-19 prior to death.

Based on all-cause mortality, 29% of the confirmed and 61% of the possible resident cases died, compared to 28% of the null resident cases. Time to event analysis found a hazard ratio of 13.26 (95% CI 5.61 to 31.34) for being a possible or confirmed case amongst residents, after adjusting for age and sex. No significant differences in mortality were found between male and female residents, or between those aged under 80 and 80 plus.

A Kaplan Meier survival curve (**Figure 3**), for all-cause mortality amongst the resident population, shows a sharp drop in survival probability between approximately 30 days and 65 days after the start of the study time-frame, at the peaks of the outbreaks. The probability of survival at 30 days from the start of the study period was 96% (95% CI 92% to 100%), whereas at day 65 the probability of survival had reduced to 62% (95% CI 53% to 72%). Survival probability plateaued after day 73 at 58% (95% CI 49% to 69%), after which no further deaths occurred during the study period.

COVID-19 Interventions

All staff members testing positive stopped working with immediate effect and isolated for two weeks. All residents returning to a home from hospital were isolated in their rooms for 2 weeks.

Interventions included:

- Cessation of visiting: implemented from the 12/03/2020 at all homes.
- Isolation of all residents: Implemented first by Home A (12/03/2020), followed by Home D (25/03/2020), Home C (29/03/2020) and Home B (31/03/2020).
- PPE use: Implemented first by Home A (12/03/2020), then Homes B and D (25/03/2020) and Home C (29/03/2020).

Discussion

Main finding of this study

COVID-19 spread rapidly amongst staff and residents in the care homes following multiple introductions and likely subsequent person-to-person spread.

What is already known?

The significant outbreaks of COVID-19 observed at the 4 homes is in line with outbreaks observed in other epidemiological studies in the USA and UK [4,7,8,9,10,11,12,13,20].

Genomic sequencing of viable samples showed that there were at least 5 different UK lineages of COVID-19 present in the homes, demonstrating multiple introductions of COVID-19 to the homes, which has also been observed elsewhere [9,20]. However, limited conclusions can be drawn here due to the small number of sequenced samples.

Amongst confirmed cases of COVID-19 the most common presenting symptoms were fever and cough, which is in keeping with the general accepted primary symptoms of COVID-19 [23]. Anosmia was not reported as a symptom amongst any confirmed resident cases, which echoes the findings of other studies [7,9,11,12]. This could be due to anosmia being a difficult symptom for residents with cognitive or communication problems to express.

What this study adds

Morbidity and mortality was high, particularly amongst residents, with the majority (91%) of resident deaths with a cause of COVID-19 or suspected COVID-19 on their death certificates dying at the homes rather than in hospital. This is not data that has been reported in other studies, so it is not clear whether this is typical or not.

The majority of confirmed cases amongst both staff and residents presented with at least one typical symptom, along with symptoms of headaches, fatigue and myalgia in around a third to a half of confirmed staff cases, highlighting the importance of considering universal staff testing rather than symptom based screening.

Outbreak peaks within each home occurred in April, at a time when only limited symptom-based testing was taking place. Universal testing did not begin until May. Overall, almost half of the residents and a quarter of the staff in this study were not tested during the outbreaks, and for those that were there was an average delay from symptom onset to test date of 20 days. RT-PCR throat swab tests are not guaranteed to produce a positive result for SARS-CoV-2 infection, with a false-negative result becoming more likely the longer the time since symptom onset [21], and found to occur in up to 68% of pharyngeal swab tests [22]. Therefore, opportunities to confirm cases of COVID-19 may have been missed, leading to potential underestimations of disease prevalence and attack rates, with delays in implementation interventions to contain the outbreaks.

In all but one of the homes (Home A) crude attack rates were higher amongst staff than residents, which is opposite to findings of other studies [7,9,10,13]. This may be due to staff having higher risks of exposure to the virus in the community, where COVID-19 was circulating at the time.

The differences in dates of implementation of resident isolation between the homes may be an explanation for the differences in attack rates between the homes, and account for why Home A had the lowest attack

rates and Home B the highest. In homes A, B and C the peak of the outbreaks occur in the days following the implementation of isolation of residents and restriction of shared spaces as an intervention measure against spread of the disease. This suggests that the majority of transmissions occurred in the days prior to this intervention, with it taking around 2 weeks to take effect and reduce the outbreak.

The delayed and lack of testing early in the outbreaks and delays in isolating residents before they became symptomatic are both likely contributing factors to the extensive transmission of COVID-19 in these homes.

Limitations of this study

There are several limitations with this study. Firstly, it is based on a single group of care homes in Wales, so whilst representative of typical care homes in the UK, it may not be representative of other care homes that specialise in different areas of care or in other countries.

Not all staff members and residents were tested during the outbreak, so the true incidence and attack rates of COVID-19 within these study populations or a true timeline of events cannot be certain. The numbers presented for confirmed cases are almost certainly an underestimate, which is why both confirmed and possible cases were included in this study.

Genomic sequencing was only available for a portion of those residents and staff who were tested, so only limited conclusions can be drawn. Further research in this area is required to understand the true spread and transmission of COVID-19 in care homes.

Data was collected retrospectively on symptom onset dates, so may be open to recall bias. There may also be reporter bias amongst staff if they have been symptomatic as it could impact on work attendance.

Investigations into the causes of excess deaths in care homes to identify if this is due directly to COVID-19 alone, or also secondary indirect consequences of the pandemic would also be valuable.

Conclusion

These study results show that COVID-19 can spread rapidly and extensively through care homes amongst both residents and staff. All 4 co-sited homes had significant outbreaks, but it seems likely that the virus was introduced to each home on different occasions from multiple sources, rather than necessarily being spread between the homes. The high numbers of resident deaths highlight the vulnerability of this population to COVID-19.

Care homes should be enabled to take proactive steps to prevent introduction and transmission of COVID-19, including restricting visitors, universal testing and isolation of residents as required. Waiting for identification of the first case before taking action does not appear to be a sufficient strategy for preventing an outbreak.

References

- 1 Challis, D., Mozley, C.G., Sutcliffe, C. et al. 2000. Dependency in older people recently admitted to care homes. *Age and Ageing* 29, pp. 255-260.
- 2 Competition and Markets Authority. 2017. *Care homes market study: summary of final report* [Online]. Available at <https://www.gov.uk/government/publications/care-homes-market-study-summary-of-final-report/care-homes-market-study-summary-of-final-report#fn:3> . Accessed 1pm 29/07/2020.
- 3 Jordan, R.E., Adab, P. and Cheng, K.K. 2020. COVID-19: Risk factors for severe disease and death. *British Medical Journal* 368, pp. 1-2.
- 4 McMichael, T.M., Currie, D.W., Clark, S. et al. 2020. Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *The New England Journal of Medicine* 382, pp. 2005-2011.
- 5 Comas-Herrera, A., Zalakain, J., Lemmon, E. et al. 2020. *Mortality associated with COVID-19 outbreaks in care homes: early international evidence*. Article in LTCcovid.org, International Long-Term Care policy Network, CPEC-LSE.
- 6 Hollinghurst, J., Lyons, J., Fry, R. et al. 2021. The impact of COVID-19 on adjusted mortality risk in care homes for older adults in Wales, UK: a retrospective population-based cohort study for mortality in 2016-2020. *Age and Ageing* 50, pp. 25-31.
- 7 Arons, M.M., Hatfield, K.M., Reddy, S.C. et al. 2020. Presymptomatic SARS-CoV-2 infections and transmission in a skilled nursing facility. *The New England Journal of Medicine* 382, pp. 2081-2090.
- 8 Blackman, C., Farber, S., Feifer, R.A., Mor, V., White, E.M. 2020. An illustration of SARS-CoV-2 dissemination within a skilled nursing facility using heat maps. *Journal of the American Geriatrics Society* 00, pp. 1-5.
- 9 Graham, N.S.N., Junghans, C., Downes, R. et al. 2020. SARS-CoV-2 infection, clinical features and outcome of COVID-19 in United Kingdom nursing homes. *Journal of Infection* 38, pp. 1-9.

- 10 Goldberg, S.A., Lennerz, J., Klompas, M. et al. 2020. Presymptomatic transmission of SARS-CoV-2 amongst residents and staff at a skilled nursing facility: Results of real-time PCR and serologic testing. *Clinical Infectious Diseases*, pp. 1-4.
- 11 Patel, M.C., Chaisson, L.H., Borgetti, S. et al. 2020. Asymptomatic SARS-CoV-2 infection and COVID-19 mortality during an outbreak investigation in a skilled nursing facility. Oxford University Press for the Infectious Diseases Society of America.
- 12 Kimball, A., Hatfield, K.M., Arons, M. et al. 2020. Asymptomatic and presymptomatic SARS-CoV-2 infections in residents of a long-term care skilled nursing facility – King County, Washington, March 2020. *Morbidity Mortality Weekly Report* 69 (13), pp. 377-381.
- 13 Dora, A.V., Winnett, A., Jatt, L.P. et al. 2020. Universal and serial laboratory testing for SARS-CoV-2 at a long-term care skilled nursing facility for veterans – Los Angeles, California, 2020. *Morbidity and Mortality Weekly Report* 69 (21), pp. 651-655.
- 14 Ladhani, S.N., Yimmy Chow, J., Janarthanan, R. et al. 2020. Investigation of SARS-CoV-2 outbreaks in six care homes in London, April 2020. *The Lancet* 395, pp. 1-8.
- 15 Quick, J., Grubaugh, N.D., Loman, N.J. et al. 2017. Multiplex PCR method for MinION and Illumina sequencing of Zika and other virus genomes directly from clinical samples. *Nature Protocols* 12, pp. 1261-1276.
- 16 Rambaut, A., Holmes, E.C., O’Toole, A. et al. 2020. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nature Microbiology*.
- 17 Kassambara, A. Kosinski, M. and Przemyslaw, B. 2020. survminer: Drawing Survival Curves using 'ggplot2'. R package version 0.4.8. [Online]. Available at: <https://CRAN.R-project.org/package=survminer> Accessed 3pm 17.08.2020.
- 18 Therneau, T. 2020. A Package for Survival Analysis in R. R package version 3.1-12 [Online]. Available at: <https://CRAN.R-project.org/package=survival>. Accessed 3pm 17/08/2020.
- 19 Therneau, T.M. and Patricia M. Grambsch, P.M. 2000. Modelling Survival Data: Extending the Cox Model. Springer, New York. ISBN 0-387-98784-3.
- 20 Gallichotte, E.N., Quicke, K.M., Sexton, N.R. et al. 2020. Longitudinal surveillance for SARS-CoV-2 RNA among asymptomatic staff in five Colorado skilled nursing

facilities: epidemiologic, virologic and sequence analysis. medRxiv doi:

<https://dx.doi.org/10.1101/2020.06.08.20125989>. Preprint.

21 Wikramaratna, P.S., Paton, R.S., Ghafari, M. Lourenco, J. 2020. Estimating false-negative detection rate of SARS-CoV-2 by RT-PCR. *Eurosurveillance* 25 (50).

22 Wang, W., Xu, Y., Gao, R. et al. 2020. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA* 323 (18), pp. 1843-1844.

23 Public Health England. 2020. *Guidance COVID-19: epidemiology, virology and clinical features* [Online]. Available at

<https://www.gov.uk/government/publications/wuhan-novel-coronavirus-background-information/wuhan-novel-coronavirus-epidemiology-virology-and-clinical-features> .

Accessed 2pm 13/05/2020

FIGURE 1




Figure 1: Horizontal bar chart of presenting symptoms of staff and residents in Homes A, B, C and D, as a percentage of total possible and confirmed cases.

FIGURE 2




Figure 2: Epi-curves of all possible and confirmed cases amongst residents and staff by date of onset of initial symptoms in Home A (n=24), Home B (n=51), Home C (n=32) and Home D (n=16). Dashed black line

indicates date from which residents were isolated. Note – 8 confirmed cases had no date of onset and were therefore not included in the epi-curves.

FIGURE 3




Figure 3: Kaplan Meier survival curve illustrating resident survival probability by days since start of study period (16/02/2020) for all homes combined, based on all-cause mortality. Shaded area designates 95% confidence intervals. Final date is 31/05/2020 (end of the study period).

TABLE 1

Home	Residents				Staff			
	n	Possible cases ¹	Confirmed cases ¹	Crude attack rate ²	n	Possible cases ³	Confirmed cases ³	Crude attack rate ⁴
A	33	11 (33%)	2 (6%)	39%	36	9 (25%)	3 (8%)	33%
B	34	18 (53%)	6 (18%)	71%	30	13 (43%)	16 (53%)	97%
C	26	11 (42%)	5 (19%)	62%	27	16 (59%)	4 (15%)	74%
D	7	3 (43%)	1 (14%)	57%	13	12 (92%)	1 (8%)	100%
Total	100	43 (43%)	14 (14%)	57%	102 ⁵	47 ⁵ (46%)	23 ⁵ (23%)	69%

Table 1: Counts and percentages of resident and staff populations at risk, possible cases, confirmed cases and crude attack rates, by home and as a total. Note: 1-percentage is of resident n; 2-possible and confirmed

resident cases combined; 3-percentage is of staff n; 4-possible and confirmed staff cases combined; 5-some staff members worked in both Home C and Home D, so they have not been counted twice for the total staff value.