

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/141465/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Kopczynska, Maja, Sharif, Ben, Pugh, Richard, Otahal, Igor, Havalda, Peter, Groblewski, Wojciech, Lynch, Ceri, George, David, Sutherland, Jayne, Pandey, Manish, Jones, Phillippa, Murdoch, Maxene, Hatalyak, Adam, Jones, Rhidian, Kacmarek, Robert, Villar, Jesús, Szakmany, Tamas ORCID: <https://orcid.org/0000-0003-3632-8844> and on behalf of the PANDORA-WALES Investigators 2020. Prevalence and outcomes of acute hypoxaemic respiratory failure in Wales: The PANDORA-WALES Study. *Journal of Clinical Medicine* 9 (11), 3521. 10.3390/jcm9113521 file

Publishers page: <http://dx.doi.org/10.3390/jcm9113521>
<<http://dx.doi.org/10.3390/jcm9113521>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





Article

Prevalence and Outcomes of Acute Hypoxaemic Respiratory Failure in Wales: The PANDORA-WALES Study

Maja Kopczynska ^{1,2}, Ben Sharif ^{1,3}, Richard Pugh ⁴, Igor Otahal ⁵, Peter Havalda ⁵, Wojciech Groblewski ⁶, Ceri Lynch ³, David George ⁷, Jayne Sutherland ⁸, Manish Pandey ⁹, Phillippa Jones ¹⁰, Maxene Murdoch ¹⁰, Adam Hatalyak ¹¹, Rhidian Jones ¹², Robert M. Kacmarek ^{13,14}, Jesús Villar ^{15,16}, Tamas Szakmany ^{1,10,*}
and on behalf of the PANDORA-WALES Investigators [†]

¹ Department of Anaesthesia, Intensive Care and Pain Medicine, Division of Population Medicine, Heath Park Campus, Cardiff University, Cardiff CF14 4XN, UK; maya.kopczynska@gmail.com (M.K.); SharifBS@cardiff.ac.uk (B.S.)

² Salford Royal NHS Trust, Stott Lane, Manchester M6 8HD, UK

³ Anaesthetic Department, Royal Glamorgan Hospital, Cwm Taf Morgannwg University Health Board, Llantrisant CF72 8XR, UK; Ceri.Lynch5@wales.nhs.uk

⁴ Anaesthetic Department, Glan Clwyd Hospital, Betsi Cadwaladr University Health Board, Bodelwyddan, Rhyl LL18 5UJ, UK; Richard.Pugh@wales.nhs.uk

⁵ Anaesthetic Department, Glangwili Hospital, Hywel Dda University Health Board, Carmarthen SA31 2AF, UK; Igor.Otahal@wales.nhs.uk (I.O.); Peter.Havalda@wales.nhs.uk (P.H.)

⁶ Anaesthetic Department, Withybush Hospital, Hywel Dda University Health Board, Haverfordwest SA61 2PZ, UK; Wojciech.Groblewski@wales.nhs.uk

⁷ Anaesthetic Department, Wrexham Maelor Hospital, Betsi Cadwaladr University Health Board, Wrexham LL13 7TD, UK; David.George@doctors.org.uk

⁸ Ed Major Critical Care Unit, Morriston Hospital, Swansea Bay, University Health Board, Swansea SA6 6NL, UK; Jayne.Sutherland@doctors.org.uk

⁹ Critical Care Department, University Hospital Wales, Cardiff and Vale University Health Board, Cardiff CF14 4XW, UK; manish55pandey@gmail.com

¹⁰ Critical Care Directorate, Royal Gwent Hospital, Aneurin Bevan University Health Board, Newport, Gwent NP20 2UB, UK; phillippajones@gmail.com (P.J.); Maxene.Murdoch@doctors.org.uk (M.M.)

¹¹ Critical Care Directorate, Nevill Hall Hospital, Aneurin Bevan University Health Board, Abergavenny NP7 7EG, UK; adam.hatalyak@gmail.com

¹² Anaesthetic Department, Princess of Wales Hospital, Cwm Taf Morgannwg University Health Board, Bridgend CF31 1RQ, UK; Rhidianjones@hotmail.com

¹³ Department of Respiratory Care, Massachusetts General Hospital, Boston, MA 02114, USA; rkacmarek@partners.org

¹⁴ Department of Anesthesia, Harvard University, Boston, MA 02115, USA

¹⁵ CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, 28029 Madrid, Spain; jesus.villar54@gmail.com

¹⁶ Research Unit, Hospital Universitario Dr. Negrín, 35010 Las Palmas de Gran Canaria, Spain

* Correspondence: szakmany1@cardiff.ac.uk; Tel.: +44-2920723112

† Membership of the PANDORA-WALES Investigators is provided in the Acknowledgments.

Received: 9 September 2020; Accepted: 29 October 2020; Published: 31 October 2020



Abstract: Background: We aimed to identify the prevalence of acute hypoxaemic respiratory failure (AHRF) in the intensive care unit (ICU) and its associated mortality. The secondary aim was to describe ventilatory management as well as the use of rescue therapies. Methods: Multi-centre prospective study in nine hospitals in Wales, UK, over 2-month periods. All patients admitted to an ICU were screened for AHRF and followed-up until discharge from the ICU. Data were collected from patient charts on patient demographics, clinical characteristics, management and outcomes. Results: Out of

2215 critical care admissions, 886 patients received mechanical ventilation. A total of 197 patients met inclusion criteria and were recruited. Seventy (35.5%) were non-survivors. Non-survivors were significantly older, had higher SOFA scores and received more vasopressor support than survivors. Twenty-five (12.7%) patients who fulfilled the Berlin definition of acute respiratory distress syndrome (ARDS) during the ICU stay without impact on overall survival. Rescue therapies were rarely used. Analysis of ventilation showed that median V_t was 7.1 mL/kg PBW (IQR 5.9–9.1) and 21.3% of patients had optimal ventilation during their ICU stay. Conclusions: One in four mechanically ventilated patients have AHRF. Despite advances of care and better, but not optimal, utilisation of low tidal volume ventilation, mortality remains high.

Keywords: respiratory failure; ventilation; rescue therapies; survival

1. Introduction

Worldwide over 100 million patients are ventilated annually, mostly in operating theatres, with about 10 million ventilated in intensive care units (ICU). It is estimated that approximately one million patients develop acute hypoxemic respiratory failure (AHRF), although data on the exact prevalence and outcomes of this condition are sparse [1,2]. Acute respiratory distress syndrome (ARDS) is an acute inflammatory lung injury associated with AHRF [3]. Currently, the diagnosis of ARDS is challenging and relies on patient characteristics, such as a history of a predisposing illness, an acute onset, as well as radiological and physiological measurements [4]. Despite the introduction of a new ARDS definition by the Berlin criteria, it is still associated with mortality greater than 50% [3,5]. Although there are some previously published observational studies examining the incidence and mortality of patients with AHRF and ARDS, there are no studies specifically assessing the epidemiological characteristics, patterns of ventilation and clinical outcomes in patients with acute hypoxemic respiratory failure in the current era of lung protective ventilation [2,6,7]. Understanding the factors associated with AHRF outcomes could lead to development of effective interventions and improvements in care.

The primary aim of our “Prevalence and Outcome of Acute Hypoxemic Respiratory Failure in Wales (PANDORA-WALES)” study was to identify the prevalence of AHRF in ICU in Wales as well as its associated mortality. The secondary aim of the study was to investigate the values of parameters of ventilatory management as well as the use of rescue therapies for hypoxemia, such as prone positioning, recruitment manoeuvres and extracorporeal assist.

2. Experimental Section

2.1. Study Design and Participants

This was a multi-centre prospective observational study carried out in nine hospitals in Wales, United Kingdom. Patients were enrolled during two periods of two consecutive months (1 October 2017 to 30 November 2017 and 1 February 2018 to 31 March 2018). Data collection was performed using a secure digital data collecting platform [8].

During the study, consecutive patients admitted to ICU were screened daily and recruited to the study if they fulfilled all inclusion criteria: age ≥ 18 -years old; endotracheal intubation and mechanical ventilation; $PaO_2/FiO_2 \leq 300$ mmHg on invasive mechanical ventilation with a PEEP of 5 cm H_2O or more, and with a FiO_2 of 0.3 or more. Data were collected on patient demographics: reason for ICU admission, methods for diagnosis of the cause of AHRF, physiological and laboratory results, treating team management and complications during the ICU stay. Patients were followed-up until discharge from the ICU.

Data management, monitoring and reporting of the study were performed in accordance with the Good Clinical Practice Guidelines. The study was registered with each participating University Health Board and internal risk reviews classed it as a service evaluation, with no formal ethical approval required. The study was prospectively registered at an international registry (NCT03358043).

2.2. Statistical Analysis

Categorical variables are described as proportions and are compared using a Chi-square test. Continuous variables are described as median and inter-quartile range and compared using a Mann-Whitney U test between the groups. We compared patient groups ventilated with different modalities and survivors vs. non-survivors. We used univariate regression analysis to determine risk factors of mortality. A two-tailed p -value < 0.05 was considered statistically significant. All statistical tests were calculated using SPSS 23.0 (SPSS Inc., Chicago, IL, USA).

3. Results

3.1. Patient Characteristics

There were 2215 critical care admissions in the participating hospitals during the study period, with 886 patients receiving mechanical ventilation. A total of 197 (22.2%) patients met all of the inclusion criteria of AHRF and were recruited to the study (Figure 1). Median patient age was 60 years (IQR 49–70) and 59.4% (117/197) of patients were males. The summary of patient demographics and clinical characteristics is summarized in Table 1.




Figure 1. Prevalence and Outcome of Acute Hypoxemic Respiratory Failure in Wales (PANDORA-WALES) recruitment flow chart and eventual study sample. Mechanically ventilated patients who fulfilled study criteria of $\text{PaO}_2/\text{FiO}_2 \leq 300$ mmHg on invasive mechanical ventilation with a Positive End-Expiratory Pressure (PEEP) of 5 cm H₂O or more, and with a FiO_2 of 0.3 or more were included in the study. All other patients were ineligible.

Table 1. Summary of patient demographics and clinical characteristics.

Patient Characteristics	Number of Patients, Percentage (n = 197)
<i>Demographics</i>	
Age (years)	60 (49–70)
Male sex	117 (59.4%)
<i>Comorbidities</i>	
HTN	81 (41.1%)
HF	16 (8.1%)
Diabetes	27 (13.7%)
Obesity	26 (13.2%)
Liver cirrhosis	11 (5.6%)
Chronic kidney failure	17 (8.6%)
Immunosuppression	12 (6.1%)
Neuromuscular disease	1 (0.5%)
Malignancy	11 (5.6%)
Use of NIV at home	3 (1.5%)
Smoking	46 (23.4%)
Alcohol excess	32 (16.2%)
<i>Hospital type</i>	
Tertiary hospital	44 (22.3%)
District general hospital	153 (77.7%)
<i>Reason for ICU admission</i>	
Clinical condition	174 (88.3%)
Emergency surgery	32 (16.2%)
Planned surgery	2 (1.0%)
<i>Physiology at the time of recruitment</i>	
Baseline PaO ₂ /FiO ₂ –median mmHg (IQR)	150 (99–209)
V _T –median mL/kg PBW (IQR)	6.9 (6.1–8.2)
Respiratory rate–median (IQR)	19 (16–23)
Plateau pressure–median cm H ₂ O (IQR)	22 (17–25)
Peak pressure–median cm H ₂ O (IQR)	23 (19–29)
PEEP–median cm H ₂ O (IQR)	7 (5–10)
pH–median (IQR)	7.28 (7.17–7.37)
PaCO ₂ –median kPa (IQR)	6.2 (5.2–7.9)

Data are median (IQR) or n (%). HTN, hypertension; HF, heart failure; NIV, non-invasive ventilation; VT, tidal volume; PBW, predicted body weight; PEEP, positive end-expiratory pressure.

The diagnosis of the cause of respiratory failure was performed using chest X-rays in the majority of the patients (86.3%, 170/197). Other modalities of diagnosis included CT scan (24.4%, 48/197); blood culture (19.8%, 39/197); sputum culture (18.8%, 37/197); echocardiography (14.2%, 28/197); bronchoscopy (8.6%, 17/197) and bronchoalveolar lavage (4.1%, 8/197). Lung biopsy and autopsy post-mortem were not performed in any of the cases.

The causes of AHRF were as follows: pneumonia (29.9%, 59/197), sepsis (20.8%, 41/197), cardiogenic shock (12.7%, 25/197), central nervous system disorders (8.1%, 16/197), chronic obstructive pulmonary

disease exacerbation (5.6%, 11/197), hypovolaemic shock (4.6%, 9/197), drug overdose (3.6%, 7/197), asthma exacerbation (2.5%, 5/197), chest trauma (2.5%, 5/197), pancreatitis (2%, 4/197), pulmonary oedema (2%, 4/197), malignancy (1%, 2/197), pneumothorax (1%, 2/197), lung empyema (1%, 2/197), pulmonary embolism (1%, 2/197) and other (1.5%, 3/197).

3.2. Patient Ventilation

Analysis of all the measurements during the whole ICU stay showed that median V_T was 7.1 mL/kg PBW (IQR 5.9–9.1) (Figure 2A). PEEP of <6 cm H₂O was used in 43.9% patients, 6–10 cm H₂O in 45.8%, 11–15 cm H₂O in 8.8% and over 15 cm H₂O in 1.5% (Figure 2B).




Figure 2. (A) V_T measurements during the whole ICU stay. (B) PEEP measurements during the whole ICU stay. Absolute numbers are shown above the bars.

We also investigated what proportion of patients received an optimal V_T (less than 6 and between 6 and 7 mL/kg PBW) by analysing daily V_{Tmax} and V_{Tmin} during the whole ICU stay. A total of 21.1% of all V_{Tmax} measurements were within the optimal range and 75.5% of V_{Tmin} measurement were optimal. For V_{Tmax} , 77.9% of measurements were above 7 mL/kg PBW and 7.8% were below 6 mL/kg PBW. For V_{Tmin} , 23.7% of measurements were above 7 mL/kg PBW and 47.5% were below 6 mL/kg PBW. Forty-two patients (21.3%) had an optimal ventilation at some point during the ICU stay.

The majority of patients were treated with volume controlled (64%, 126/197) and pressure controlled (28.9%, 57/197) ventilation. Five patients were treated with other ventilation types: one with assisted spontaneous breathing and four with continuous positive airway pressure. We have investigated

an impact on ventilation type on delivered V_T values (Table 2). Volume controlled ventilation was associated with higher proportions of optimal V_T ($p = 0.049$) and lower proportions of V_T above 7mL/kg PBW ($p = 0.038$).

Table 2. Impact of ventilation type on V_T classification.

V_T	Ventilation Type		<i>p</i> Value
	Volume Controlled (n = 126)	Pressure Controlled (n = 57)	
V_T in optimal range	76 (60.3%)	25 (43.9%)	0.049
V_T above 7 mL/kg PBW	50 (39.7%)	32 (56.1%)	0.038

Ventilation types other than volume or pressure controlled were removed from the analysis due to small values. Optimal ventilation was described as less than 6 and between 6 and 7 mL/kg PBW. Data were missing for nine patients. *p*-value <0.05 is bold and underlined. For statistical analysis a Chi-square test was used.

3.3. Patient Management

Prior to enrolment in the study and meeting respiratory failure criteria, 9.6% patients (19/197) were treated with non-invasive ventilation and 19.8% (39/197) received high flow nasal oxygen therapy. On the day of recruitment and throughout the ICU stay, patients received other interventions and monitoring modalities presented in Table 3. On the day of recruitment, one in five patients were treated with continuous neuromuscular blockade and one in six had recruitment manoeuvres performed. Prone positioning was rarely used either as an initial treatment option or as a rescue therapy.

Table 3. Intervention and monitoring modalities provided to patients throughout ICU stay.

Intervention and Monitoring	Number of Patients, Percentage (n = 197)
<i>During the ICU stay</i>	
Oesophageal pressure monitoring	1 (0.5%)
Pulmonary artery catheter	1 (0.5%)
Nitric oxide	3 (1.5%)
Vasopressor use	80 (40.6%)
Corticosteroids	56 (28.4%)
Systemic vasodilators	6 (3%)
Blood transfusion	12 (6.1%)
Renal replacement therapy	26 (13.2%)
High frequency ventilation	2 (1%)
ECMO	1 (0.5%)
ECCO2R	1 (0.5%)
NAVA	1 (0.5%)
PiCCO	4 (2%)
NIV after being extubated	1 (0.5%)
<i>On the day of recruitment</i>	
Neuromuscular blockade	42 (21.3%)
Recruitment manoeuvres	35 (17.8%)
<i>Prone by protocol</i>	
First 48 h	2 (1%)
During the evolution	1 (0.5%)
As a rescue	1 (0.5%)

Data are presented as n (%). ECMO, Extracorporeal membrane oxygenation; ECCO2R, Extracorporeal carbon dioxide removal; NAVA, Neurally Adjusted Ventilatory Assist; PiCCO, Pulse index Contour Cardiac Output; NIV, non-invasive ventilation.

3.4. Complications and Survival Analysis

A significant proportion of patients suffered from complications, both respiratory and systemic such as shock or sepsis as presented in Table 4.

Table 4. Complications developed by enrolled patients. Data are presented as n (%).

Complication	Number of Patients, Percentage (n = 197)
Pneumothorax	2 (1%)
Pneumonia	36 (18.3%)
Pleural effusion	11 (5.5%)
Atelectasis	4 (2%)
New lung infiltrates	21 (10.7%)
Pulmonary oedema	5 (2.5%)
Shock	27 (13.7%)
Sepsis	38 (19.3%)
Major surgical operation	16 (8.1%)

Out of 197 recruited patients, 70 (35.5%) were non-survivors, 125 (63.5%) were survivors and data were missing for 2 (1.0%). The univariate analysis of factors associated with mortality is presented in Table 5.

Table 5. Univariate analysis of factors associated with patient mortality. Data are median (IQR) or n (%). *p*-value < 0.05 is bold and underlined.

Risk Factor	Survivor (n = 125)	Non-Survivor (n = 70)	<i>p</i> Value
Age	57 (47–68)	65 (52–72)	0.034
SOFA score	8 (6–9)	9 (7–11)	0.019
Alcohol excess	15 (12.0%)	17 (24.3%)	0.026
Inotropic support	73 (58.4%)	51 (72.9%)	0.044
Recruitment manoeuvres	17 (13.6%)	17 (24.3%)	0.059

Twenty-five (12.7%) patients fulfilled the Berlin definition of ARDS at any time during the ICU stay. Median number of days between ICU admission and ARDS development was 0 (IQR 0–1). Out of all patients with ARDS, 5 (36%) were non-survivors. Development of ARDS did not have an impact on overall survival in our patient cohort (*p* = 0.37).

4. Discussion

This is the first study prospectively investigating the prevalence, management processes and outcomes of patients developing AHRF in ICUs in Wales. A total of 22.2% of patients admitted for mechanical ventilation developed AHRF, which was associated with a significant mortality rate of 35.5%, similar to other European studies [1,9].

AHRF appears to be common in the ICUs [10,11], affecting almost a quarter of those needing mechanical ventilation. However, most large-scale prevalence studies concentrate on either patients undergoing mechanical ventilation or those meeting defined criteria for acute lung injury/ARDS [9,12]. In the multi-national LUNG-SAFE study, 34% of ventilated patients fulfilled AHRF criteria, but with no data showing if this differs across countries [13]. It must be noted that the LUNG-SAFE study had a significant bias towards tertiary centres, whilst our study was mainly conducted in district

general hospitals, which could lead to a different case-mix and could explain our lower prevalence [14]. We found that the incidence of ARDS using the Berlin definition [3] was almost five times lower than in LUNG-SAFE study conducted three years earlier—12.7% vs. 67.2% of patients with AHRF, respectively [13]. This observed discrepancy highlights the difficulties in adopting the Berlin definition as inclusion criteria for clinical trials [15]. Previous studies indicated that the PaO₂/FiO₂ ratio-based severity grading is open for interpretation and that measurements taken at a standardised ventilator setting may change the severity grading of more than half of ARDS patients [4]. Importantly, in the LUNG-SAFE study approximately 40% of patients diagnosed with ARDS, were not deemed to have the condition by the treating clinician [13]. Based on our data it is difficult to ascertain whether ARDS was “underdiagnosed” as postulated by the LUNG-SAFE study, or whether the clinicians interpreted the chest infiltrates appropriately in our study and the incidence of ARDS in Wales is within the previously described 10–15% category [9,13]. ARDS was prevalent in ICU admission in our cohort and less likely to develop as a complication of mechanical ventilation, in line with recent large international studies [10,16].

The outcome of AHRF in our study was similar to what was reported in the same period in several cohorts of mechanically ventilated hypoxaemic patients [10,17,18]. However, the observed mortality was almost 10% higher compared to cohorts in Italy and in Canada with similar demographics and similar degrees of hypoxaemia [19,20]. There are several possible explanations for these discrepancies. There is a well-documented disparity of critical care beds in the UK and in particular in Wales, compared with other developed countries [21,22]. This, in general, leads to higher acuity admissions to the ICU, with higher underlying risk of death, which has been described previously [23]. Indeed, our patient population was older, was more likely to have multi-organ failure based on the SOFA score and had higher prevalence of underlying medical problems compared to other studies [9]. Unsurprisingly, these previously well-described risk factors were associated with mortality in our cohort [9,24]. However, there is also a possibility that care processes and adherence to best practice guidance in mechanical ventilation was better in the studies conducted in Italy and Canada, accounting for better mortality [19,20]. In our cohort of patients with AHRF only one in five received optimal ventilation during their ICU stay, recommended as <7 mL/kg PBW [25]. Four out of five patients had V_{Tmax} values above 7 mL/kg PBW and almost a quarter of the measurements of V_{Tmin} were outside the recommended range. This finding is consistent with other studies in which patients received larger tidal volumes than expected [16]. We opted to use a strict criterion for optimal V_T as it has been known for decades that the physiological V_T is 6 mL/kg PBW in humans [26]. It has been shown that ventilation with use of large tidal volumes can cause hypoxaemia and release of inflammatory mediators, leading to increase in lung inflammation and injury to other organs [27]. Several retrospective analyses of large observational studies suggested that every 1mL/kg PBW increase of V_T is also associated with higher mortality [28,29]. Although in the recent PRoVENT multi-centre prospective study of mechanically ventilated patients at risk of ARDS, Neto et al. reported that V_T was not different between survivors and non-survivors, the patient population was significantly different compared to ours, as their patients had a much higher PaO₂/FiO₂ ratio and were more likely to be admitted following surgical procedures [12]. Importantly, the median V_T in our observations was 7 compared to 8 mL/kg PBW in the PRoVENT study, continuing the trend of V_T reduction over time in various trials [9,11–13]. Whilst a multi-centre randomized controlled trial (RCT) did not find difference in ventilator free days or mortality when comparing low to intermediate V_T strategies in patients without ARDS, the PReVENT trial had inadequate separation between the two arms after day 1, which means this study was likely underpowered to detect any true difference [30]. Our findings are in-line with the outcomes of the low V_T arm of the PReVENT trial, which has targeted 7 mL/kg PBW [30].

Rescue therapies were not used in the vast majority of the patients. Neuromuscular blocking agents were used only for 21.3% and recruitment manoeuvres for 17.8% of patients on the day of diagnosis of AHRF, whereas prone positioning and more expensive and invasive technologies such

as ECMO were used in less than 1% of patients, which is significantly lower than in the study by Bellani et al. [13]. At the time of the study, both neuromuscular blockade and prone positioning was recommended for severe ARDS based on previously published RCT, whilst uncertainty existed around the use of recruitment manoeuvres [5,31,32]. However, as the majority of our patients did not meet ARDS criteria, it is possible that the relatively low use of adjunctive measures reflects ongoing uncertainty about the quality of evidence supporting these interventions. Indeed, recent RCTs put some doubt about the efficacy of these interventions in the moderate to severe hypoxaemia category, with individualised ventilation strategies, including prone positioning, shown to be harmful compared to standard care with a low PEEP strategy [33]. Importantly, the harm measured in the experimental group was mostly attributed to the misclassification of focal or diffuse changes on chest imaging, further emphasising the difficulty of ARDS diagnosis using these modalities [33]. At the beginning of the study, the ART trial was published, reporting increased harm from a recruitment manoeuvre strategy in moderate to severe ARDS, where patients had similar levels of physiological derangement to our population [34]. This publication, which has been widely discussed in conventional and social media, might have had an impact on the use of recruitment manoeuvres during our data collection period. More recently, the use of early neuromuscular blockade has been shown to be inferior to the early mobilisation, awakening and spontaneous breathing trial approach in a large, multi-centre study recruiting patients at the severe end of the hypoxaemic respiratory failure spectrum [35]. These results cast significant doubt on the efficacy and safety of deep muscle paralysis in AHRF and our findings regarding neuromuscular blockade use are in line with the standard care arm of the ROSE-PETAL study [35].

The strengths of the study include wide participation of centres in Wales including both academic centres and general district hospitals. Capturing data over a four-month period was implemented to ensure representativeness. Information was collected prospectively enabling the analysis of real-life use of ventilation and rescue therapies. A similar study, using the same dataset, has been conducted in Spain (NCT03145974). This shows that this type of assessment of ICU performance could be standardized and used internationally.

There are significant limitations to our study: although data collection was digitised, it was still labour intensive, requiring everyday screening, meaning there is the potential that some patients were missed and not recruited in the study, however, retrospective reconciliation with ICU admissions did not reveal any. We have concentrated our data collection on the winter months, when AHRF incidence could be higher, skewing the result of the prevalence of the condition. However, the timing of data sampling has enabled us to make direct comparisons with the previously published LUNG-SAFE study, which used the same period [13]. As we have concentrated on the prevalence and initial treatment of AHRF, we have not collected detailed data on important treatment modalities, which might have affected outcomes such as tracheostomy, use of spontaneous breathing trials, rate of re-intubation, fluid management, daily interruption of sedation, use of thromboprophylaxis etc. Due to the nature of the study we also did not collect data on length of stay or longer-term outcomes. As the study sample was relatively small, we were only able to perform an univariate analysis to avoid potential overfitting of the model associated with a multivariate analysis. Nevertheless, our study has also proved that large-scale collaboration is possible in our hospitals and paved the way for other national studies [36,37]. As our study was concentrating on clinical variables, we did not collect detailed biomarker profiles for our patients and computer tomography was only performed in a quarter of the patients. Hence, detailed phenotyping of patients using these modalities, which has proved useful in the COVID-19 pandemic, could not be completed [38,39].

5. Conclusions

AHRF is associated with high mortality despite increased awareness and recognition of this condition. Despite advances in care, mechanical ventilation and use of adjunctive measures are still not optimal for the vast majority of patients.

Author Contributions: Conceptualization, R.M.K., J.V. and T.S.; formal analysis, M.K.; investigation, M.K., B.S., R.P., I.O., P.H., W.G., C.L., D.G., J.S., M.P., P.J., M.M., A.H., R.J. and T.S.; methodology, M.K., B.S., R.M.K., J.V. and T.S.; project administration, R.M.K., J.V. and T.S.; resources, T.S.; supervision, T.S.; visualization, M.K.; writing—original draft, M.K. and T.S.; writing—review & editing, all authors. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: PANDORA-WALES Investigators: Svetlana Kulikouskaya, Rachel Keeling, Anthony Hanks, Jessica Smith, Chunhei Li, Priscilla Chan, Emily Hubbard, Leanne Sparrow, Zak Deere, Michael Martin, Lauren McCarthy, Subhamay Ghosh, Rhidian Gareth Hughes, Jessica Smith, Ciaran Lisboa, Orsolya Minik, Nia Williams, Charlotte Cowling, Olivia Don, Iain Mclure, Nick Roberts, Andy Campbell, Venkat Sundaram, Antash Daryanani, Piroska Toth-Tarsoly, Zsolt Toth-Tarsoly, Maja Kopczynska Ben Sharif, Richard Pugh, Igor Otahal, Peter Havalda, Wojciech Groblewski, Ceri Lynch, David George, Jayne Sutherland, Manish Pandey, Phillippa Jones, Maxene Murdoch, Adam Hatalyak, Rhidian Jones, Robert M. Kacmarek, Jesús Villar and Tamas Szakmany.

Conflicts of Interest: R.M.K. serves as a consultant for Medtronic and Orange Med Inc., and received research grants from Medtronic and Orange Med Inc and provided a web seminar on High Flow Nasal Cannulas and Ventilator Management of the Obese Patient for Nihon Kohden. J.V. received a research grant from Maquet.

References

- Villar, J.; Blanco, J.; Añón, J.M.; Santos-Bouza, A.; Blanch, L.; Ambrós, A.; Gandía, F.; Carriedo, D.; Mosteiro, F.; Basaldúa, S.; et al. The ALIEN study: Incidence and outcome of acute respiratory distress syndrome in the era of lung protective ventilation. *Intensive Care Med.* **2011**, *37*, 1932–1941. [[CrossRef](#)] [[PubMed](#)]
- Luhr, O.R.; Antonsen, K.; Karlsson, M.; Aardal, S.; Thorsteinsson, A.; Frostell, C.; Bonde, J.; the ARF Study Group. Incidence and Mortality after Acute Respiratory Failure and Acute Respiratory Distress Syndrome in Sweden, Denmark, and Iceland. *Am. J. Respir. Crit. Care Med.* **1999**, *159*, 1849–1861. [[CrossRef](#)] [[PubMed](#)]
- The ARDS Definition Task Force. Acute Respiratory Distress Syndrome: The Berlin Definition of ARDS. *JAMA* **2012**, *307*, 2526–2533. [[CrossRef](#)]
- Villar, J.; Epidemiology, S.S.I.F.; Pérez-Méndez, L.; Blanco, J.; Anon, J.M.; Blanch, L.; Belda, J.; Santos-Bouza, A.; Fernández, R.L.; Kacmarek, R.M. A universal definition of ARDS: The PaO₂/FiO₂ ratio under a standard ventilatory setting—A prospective, multicenter validation study. *Intensive Care Med.* **2013**, *39*, 583–592. [[CrossRef](#)] [[PubMed](#)]
- Papazian, L.; Forel, J.-M.; Gacouin, A.; Penot-Ragon, C.; Perrin, G.; Loundou, A.; Jaber, S.; Arnal, J.-M.; Perez, D.; Seghboyan, J.-M.; et al. Neuromuscular Blockers in Early Acute Respiratory Distress Syndrome. *N. Engl. J. Med.* **2010**, *363*, 1107–1116. [[CrossRef](#)] [[PubMed](#)]
- Valta, P.; Uusaro, A.; Nunes, S.; Ruokonen, E.; Takala, J. Acute respiratory distress syndrome: Frequency, clinical course, and costs of care. *Crit. Care Med.* **1999**, *27*, 2367–2374. [[CrossRef](#)]
- Azevedo, L.C.P.; Park, M.; Salluh, J.I.F.; Rea-Neto, A.; Souza-Dantas, V.C.; Varaschin, P.; Oliveira, M.C.; Tierno, P.F.; Dal-Pizzol, F.; Silva, U.V.A.; et al. Clinical outcomes of patients requiring ventilatory support in Brazilian intensive care units: A multicenter, prospective, cohort study. *Crit. Care* **2013**, *17*, R63. [[CrossRef](#)]
- Sharif, B.; Lundin, R.M.; Morgan, P.; Hall, J.; Dhadda, A.; Mann, C.; Donoghue, D.; Brownlow, E.; Hill, F.; Carr, G.; et al. Developing a digital data collection platform to measure the prevalence of sepsis in Wales. *J. Am. Med. Inform. Assoc.* **2016**, *23*, 1185–1189. [[CrossRef](#)]
- Bertolini, G.; Lewandowski, K.; Bion, J.; Romand, J.-A.; Villar, J.S.; Thorsteinsson, A.; Damas, P.; Armaganidis, A.; Lemaire, F.; Brun-Buisson, C.; et al. Epidemiology and outcome of acute lung injury in European intensive care units. *Intensive Care Med.* **2004**, *30*, 51–61. [[CrossRef](#)]
- Grimaldi, D.; Hraiech, S.; Boutin, E.; Lacherade, J.C.; Boissier, F.; Pham, T.; Richard, J.C.; Thille, A.W.; Ehrmann, S.; Lascarrou, J.B.; et al. Hypoxemia in the ICU: Prevalence, treatment, and outcome. *Ann. Intensive Care* **2018**, *8*, 82. [[CrossRef](#)]
- Linko, R.; Okkonen, M.; Pettilä, V.; Perttilä, J.; Parviainen, I.; Ruokonen, E.; Tenhunen, J.; Ala-Kokko, T.; Varpula, T.; The FINNALI-study group. Acute respiratory failure in intensive care units. FINNALI: A prospective cohort study. *Intensive Care Med.* **2009**, *35*, 1352–1361. [[CrossRef](#)] [[PubMed](#)]

12. Neto, A.S.; Barbas, C.S.V.; Simonis, F.D.; Artigas-Raventos, A.; Canet, J.; Determann, R.M.; Anstey, J.; Hedenstierna, G.; Hemmes, S.N.T.; Hermans, G.; et al. Epidemiological characteristics, practice of ventilation, and clinical outcome in patients at risk of acute respiratory distress syndrome in intensive care units from 16 countries (PRoVENT): An international, multicentre, prospective study. *Lancet Respir. Med.* **2016**, *4*, 882–893. [[CrossRef](#)]
13. Bellani, G.; Laffey, J.G.; Pham, T.; Fan, E.; Brochard, L.; Esteban, A.; Gattinoni, L.; Van Haren, F.; Larsson, A.; McAuley, D.F.; et al. Epidemiology, Patterns of Care, and Mortality for Patients With Acute Respiratory Distress Syndrome in Intensive Care Units in 50 Countries. *JAMA* **2016**, *315*, 788–800. [[CrossRef](#)]
14. Laffey, J.G.; Madotto, F.; Pham, T.; Bajwa, E.K.; Lorente, J.A.; Jiménez, J.I.S.; McAuley, D.F.; Slutsky, A.S.; Laffey, J.G.; Bellani, G.; et al. Geo-economic variations in epidemiology, patterns of care, and outcomes in patients with acute respiratory distress syndrome: Insights from the LUNG SAFE prospective cohort study. *Lancet Respir. Med.* **2017**, *5*, 627–638. [[CrossRef](#)]
15. Villar, J.; Pérez-Méndez, L.; Kacmarek, R.M. The Berlin definition met our needs: No. *Intensive Care Med.* **2016**, *42*, 648–650. [[CrossRef](#)]
16. Pham, T.; Neto, A.S.; Pelosi, P.; Laffey, J.G.; De Haro, C.; Lorente, M.D.C.; Bellani, G.; Fan, E.; Brochard, L.J.; Pesenti, A.; et al. Outcomes of Patients Presenting with Mild Acute Respiratory Distress Syndrome. *Anesthesiology* **2019**, *130*, 263–283. [[CrossRef](#)]
17. Hsu, C.-H.; Hung, Y.; Chu, K.-A.; Chen, C.-F.; Yin, C.-H.; Lee, C.-C. Prognostic nomogram for elderly patients with acute respiratory failure receiving invasive mechanical ventilation: A nationwide population-based cohort study in Taiwan. *Sci. Rep.* **2020**, *10*, 1–10. [[CrossRef](#)]
18. Schmidt, M.F.S.; Amaral, A.C.K.B.; Fan, E.; Rubenfeld, G. Driving Pressure and Hospital Mortality in Patients without ARDS: A Cohort Study. *Chest* **2017**, *153*, 46–54. [[CrossRef](#)]
19. Urner, M.; Jüni, P.; Hansen, B.; Wettstein, M.S.; Ferguson, N.D.; Fan, E. Time-varying intensity of mechanical ventilation and mortality in patients with acute respiratory failure: A registry-based, prospective cohort study. *Lancet Respir. Med.* **2020**, *8*, 905–913. [[CrossRef](#)]
20. Rezoagli, E.; Gatti, S.; Villa, S.; Villa, G.; Muttini, S.; Rossi, F.; Faraldi, L.; Fumagalli, R.; Grasselli, G.; Foti, G.; et al. ABO blood types and major outcomes in patients with acute hypoxaemic respiratory failure: A multicenter retrospective cohort study. *PLoS ONE* **2018**, *13*, e0206403. [[CrossRef](#)]
21. Szakmany, T.; Walters, A.M.; Pugh, R.; Battle, C.; Berridge, D.M.; Lyons, R.A. Risk Factors for 1-Year Mortality and Hospital Utilization Patterns in Critical Care Survivors. *Crit. Care Med.* **2019**, *47*, 15–22. [[CrossRef](#)] [[PubMed](#)]
22. Sakr, Y.; Moreira, C.L.; Rhodes, A.; Ferguson, N.D.; Kleinpell, R.; Pickkers, P.; Kuiper, M.A.; Lipman, J.; Vincent, J.-L. The Impact of Hospital and ICU Organizational Factors on Outcome in Critically Ill Patients. *Crit. Care Med.* **2015**, *43*, 519–526. [[CrossRef](#)] [[PubMed](#)]
23. Capuzzo, M.; Volta, C.A.; Tassinati, T.; Moreno, R.P.; Valentin, A.; Guidet, B.; Iapichino, G.; Martin, C.; Perneger, T.; Combescure, C.; et al. Hospital mortality of adults admitted to Intensive Care Units in hospitals with and without Intermediate Care Units: A multicentre European cohort study. *Crit. Care* **2014**, *18*, 1–15. [[CrossRef](#)]
24. McNicholas, B.A.; Madotto, F.; Pham, T.; Rezoagli, E.; Masterson, C.H.; Horie, S.; Bellani, G.; Brochard, L.; Laffey, J.G. Demographics, management and outcome of women and men with Acute Respiratory Distress Syndrome in the LUNG SAFE prospective cohort study. *Eur. Respir. J.* **2019**, *54*, 1900609. [[CrossRef](#)]
25. Acute Respiratory Distress Syndrome Network; Brower, R.G.; Matthay, M.A.; Morris, A.; Schoenfeld, D.; Thompson, B.T.; Wheeler, A. Ventilation with Lower Tidal Volumes as Compared with Traditional Tidal Volumes for Acute Lung Injury and the Acute Respiratory Distress Syndrome. *N. Engl. J. Med.* **2000**, *342*, 1301–1308. [[CrossRef](#)]
26. Tenney, S.M.; Remmers, J.E. Comparative Quantitative Morphology of the Mammalian Lung: Diffusing Area. *Nat. Cell Biol.* **1963**, *197*, 54–56. [[CrossRef](#)] [[PubMed](#)]
27. Calfee, C.S.; Delucchi, K.L.; Sinha, P.; Matthay, M.A.; Hackett, J.; Shankar-Hari, M.; McDowell, C.; Laffey, J.G.; O’Kane, C.M.; McAuley, D.F.; et al. Acute respiratory distress syndrome subphenotypes and differential response to simvastatin: Secondary analysis of a randomised controlled trial. *Lancet Respir. Med.* **2018**, *6*, 691–698. [[CrossRef](#)]
28. Davies, J.D.; Senussi, M.H.; Mireles-Cabodevila, E. Should a Tidal Volume of 6 mL/kg Be Used in All Patients? *Respir. Care* **2016**, *61*, 774–790. [[CrossRef](#)]

29. Lanspa, M.J.; Peltan, I.D.; Jacobs, J.; Sorensen, J.S.; Carpenter, L.; Ferraro, J.P.; Brown, S.M.; Berry, J.G.; Srivastava, R.; Grissom, C.K. Driving pressure is not associated with mortality in mechanically ventilated patients without ARDS. *Crit. Care* **2019**, *23*, 424–428. [[CrossRef](#)]
30. Simonis, F.D.; Serpa Neto, A.; Binnekade, J.M.; Braber, A.; Bruin, K.C.M.; Determann, R.M.; Goekoop, G.-J.; Heidt, J.; Horn, J.; Innemee, G.; et al. Effect of a Low vs Intermediate Tidal Volume Strategy on Ventilator-Free Days in Intensive Care Unit Patients Without ARDS A Randomized Clinical Trial. *JAMA* **2018**, *320*, 1872–1880. [[CrossRef](#)]
31. Guérin, C.; Reignier, J.; Richard, J.-C.; Beuret, P.; Gacouin, A.; Boulain, T.; Mercier, E.; Badet, M.; Mercat, A.; Baudin, O.; et al. Prone Positioning in Severe Acute Respiratory Distress Syndrome. *N. Engl. J. Med.* **2013**, *368*, 2159–2168. [[CrossRef](#)] [[PubMed](#)]
32. Hodgson, C.L.; Goligher, E.C.; Young, M.E.; Keating, J.L.; Holland, A.E.; Romero, L.; Bradley, S.J.; Tuxen, D. Recruitment manoeuvres for adults with acute respiratory distress syndrome receiving mechanical ventilation. *Cochrane Database Syst. Rev.* **2016**, *11*, CD006667. [[CrossRef](#)] [[PubMed](#)]
33. Constantin, J.-M.; Jabaudon, M.; Lefrant, J.-Y.; Jaber, S.; Quenot, J.-P.; Langeron, O.; Ferrandière, M.; Grelon, F.; Seguin, P.; Ichai, C.; et al. Personalised mechanical ventilation tailored to lung morphology versus low positive end-expiratory pressure for patients with acute respiratory distress syndrome in France (the LIVE study): A multicentre, single-blind, randomised controlled trial. *Lancet Respir. Med.* **2019**, *7*, 870–880. [[CrossRef](#)]
34. Cavalcanti, A.B.; Suzumura, É.A.; Laranjeira, L.N.; Paisani, D.D.M.; Damiani, L.P.; Guimarães, H.P.; Romano, E.R.; Regenga, M.D.M.; Taniguchi, L.N.T.; de Oliveira, R.P.; et al. Effect of Lung Recruitment and Titrated Positive End-Expiratory Pressure (PEEP) vs Low PEEP on Mortality in Patients with Acute Respiratory Distress Syndrome. *JAMA* **2017**, *318*, 1335–1345. [[CrossRef](#)] [[PubMed](#)]
35. National Heart, Lung, and Blood Institute PETAL Clinical Trials Network; Moss, M.; Huang, D.T.; Brower, R.G.; Ferguson, N.D.; Ginde, A.A.; Gong, M.N.; Grissom, C.K.; Gundel, S.; Hayden, D.; et al. Early Neuromuscular Blockade in the Acute Respiratory Distress Syndrome. *N. Engl. J. Med.* **2019**, *380*, 1997–2008. [[CrossRef](#)]
36. Szakmany, T.; Pugh, R.; Kopczynska, M.; Lundin, R.M.; Sharif, B.; Morgan, P.; Ellis, G.; Abreu, J.; Kulikouskaya, S.; Bashir, K.; et al. Defining sepsis on the wards: Results of a multi-centre point-prevalence study comparing two sepsis definitions. *Anaesthesia* **2017**, *73*, 195–204. [[CrossRef](#)]
37. Kopczynska, M.; Sharif, B.; Unwin, H.; Lynch, J.; Forrester, A.; Zeicu, C.; Cleaver, S.; Kulikouskaya, S.; Chandy, T.; Ang, E.; et al. Real World Patterns of Antimicrobial Use and Microbiology Investigations in Patients with Sepsis outside the Critical Care Unit: Secondary Analysis of Three Nation-Wide Point Prevalence Studies. *J. Clin. Med.* **2019**, *8*, 1337. [[CrossRef](#)]
38. Sinha, P.; Calfee, C.S.; Cherian, S.; Brealey, D.; Cutler, S.; King, C.; Killick, C.; Richards, O.; Cheema, Y.; Bailey, C.; et al. Prevalence of phenotypes of acute respiratory distress syndrome in critically ill patients with COVID-19: A prospective observational study. *Lancet Respir. Med.* **2020**. [[CrossRef](#)]
39. Bos, L.D.J.; Paulus, F.; Vlaar, A.P.J.; Beenen, L.F.M.; Schultz, M.J. Subphenotyping Acute Respiratory Distress Syndrome in Patients with COVID-19: Consequences for Ventilator Management. *Ann. Am. Thorac. Soc.* **2020**, *17*, 1161–1163. [[CrossRef](#)]

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).