

CLINICAL PRACTICE GUIDELINE ENDORSED by the EUROPEAN SOCIETY OF ENDOCRINOLOGY (ESE)**THE 2021 EUROPEAN GROUP ON GRAVES' ORBITOPATHY (EUGOGO) CLINICAL PRACTICE GUIDELINES FOR THE MEDICAL MANAGEMENT OF GRAVES' ORBITOPATHY**

***Bartalena L¹, *Kahaly GJ², Baldeschi L³, Dayan CM⁴, Eckstein A⁵, Marcocci C⁶, Marinò M⁶, Vaidya B⁷, Wiersinga WM⁸ on behalf of EUGOGO**

*Luigi Bartalena and *George J. Kahaly equally contributed to this paper and share first authorship

¹Department of Medicine and Surgery, University of Insubria, Varese, Italy, ²Department of Medicine I, Johannes Gutenberg-University (JGU) Medical Center, Mainz, Germany, ³Department of Ophthalmology, Cliniques Universitaires Saint Luc, Catholic University of Louvain, Brussels, Belgium, ⁴Thyroid Research Group, Cardiff University School of Medicine, Cardiff, UK, ⁵Clinic for Ophthalmology, University Clinic, Essen, Germany, ⁶Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy, ⁷Department of Endocrinology, Royal Devon & Exeter Hospital and University of Exeter Medical School, Exeter, UK, ⁸Amsterdam University Medical Center, Amsterdam, the Netherlands

Short Title: Guidelines for Medical Management of GO

Key Words: Guidelines, Graves' Orbitopathy, Intravenous Glucocorticoids, TSH-Receptor Antibody, Smoking

Number of words: Abstract 285, Text: 7403, two tables, four Figures, 191 references

Correspondence: Prof. George J. Kahaly, Dept. of Medicine I, Johannes Gutenberg-University (JGU) Medical Center, Mainz 55101, Germany. Email gkahaly@uni-mainz.de

ABSTRACT

Graves' orbitopathy (GO) is the main extrathyroidal manifestation of Graves' disease (GD). Choice of treatment should be based on assessment of clinical activity and severity of GO. Early referral to specialized centers is fundamental for most patients with GO. Risk factors include smoking, thyroid dysfunction, high serum level of thyrotropin receptor antibodies, radioactive iodine (RAI) treatment, and hypercholesterolemia. In mild and active GO, control of risk factors, local treatments and selenium (selenium-deficient areas) are usually sufficient; if RAI treatment is selected to manage GD, low-dose oral prednisone prophylaxis is needed, especially if risk factors coexist. For both active moderate-to-severe and sight threatening GO, antithyroid drugs are preferred when managing Graves' hyperthyroidism. In moderate-to-severe and active GO intravenous (iv) glucocorticoids are more effective and better tolerated than oral glucocorticoids. Based on current evidence and efficacy/safety profile, costs and reimbursement, drug availability, long-term effectiveness and patient choice after extensive counselling, a combination of iv methylprednisolone and mycophenolate sodium is recommended as first-line treatment. A cumulative dose of 4.5 grams (g) of iv methylprednisolone in 12 weekly infusions is the optimal regimen. Alternatively, higher cumulative doses not exceeding 8 g can be used as monotherapy in most severe cases and constant/inconstant diplopia. Second-line treatments for moderate-to-severe and active GO include: a) a second course of iv methylprednisolone (7.5 g) subsequent to careful ophthalmic and biochemical evaluation, b) oral prednisone/prednisolone combined with either cyclosporine or azathioprine; c) orbital radiotherapy combined with oral or iv glucocorticoids, d) teprotumumab; e) rituximab and f) tocilizumab. Sight threatening GO is treated with several high single doses of iv methylprednisolone per week and, if unresponsive, with urgent orbital decompression. Rehabilitative surgery (orbital decompression, squint and eyelid surgery) is indicated for inactive residual GO manifestations.

INTRODUCTION

Graves' orbitopathy (GO), also called thyroid eye disease or thyroid-associated orbitopathy, is the major extrathyroidal manifestation of Graves' disease (GD), although it may less frequently occur in patients with chronic autoimmune thyroiditis (1). GO is relatively rare (estimated incidence: 0.54-0.9 cases/100,000/year in men, 2.67-3.3 cases/100,000/year in women) with more commonly mild and non-progressive cases and moderate-to-severe forms accounting for 5-6% of cases only (2, 3). GO has an impact on quality of life (QoL), even within mild disease (4-6), and poses a significant public health burden, in terms of direct and indirect costs (7). GO is a major therapeutic challenge in its moderate-to-severe forms, often incompletely responsive to available medical treatments. After publication of the 2016 European Thyroid Association (ETA)/European Group on GO (EUGOGO) guidelines for the management of GO (8), several relevant studies have been published, particularly randomized clinical trials (RCTs) of newer biological agents for the treatment of moderate-to-severe and active GO (1, 9-12). This prompted the EUGOGO Executive Committee to appoint an *ad hoc* task force committed to updating the guidelines, focusing on the medical management of GO. Synopsis of recommendations is presented in **Table 1**. Emergency and rehabilitative surgery of GO is briefly discussed, as it will be separately addressed by the ophthalmologist members of EUGOGO.

METHODS

Literature Search. Data acquisition was based on PubMed search strategies, with particular regard to papers published subsequent to the 2016 ETA/EUGOGO guidelines (8). In addition, the list of references of relevant citations and chapters of major textbooks were evaluated for any additional appropriate citation.

Grading. The GRADE system was used to make recommendations and express the quality of the evidence (13). The task force used the following coding system: (1) indicates a strong

recommendation and is associated with the phrase “we recommend”; (2) denotes a weak recommendation and is associated with the phrase “we suggest”. Evidence grading: $\emptyset\emptyset\emptyset\emptyset$ denotes very low quality evidence; $\emptyset\emptyset\emptyset\emptyset$, low quality; $\emptyset\emptyset\emptyset\emptyset$, moderate quality; $\emptyset\emptyset\emptyset\emptyset$, high quality. The draft was submitted to all members of EUGOGO and commented by e-mail. All members of the task force unanimously approved the final version and the 32 recommendations. Each EUGOGO site identified 1-2 members to be acknowledged as contributors. Among 48 potential contributors, written permission for being acknowledged was granted by 48 (100%).

CLASSIFICATION OF GO

Treatment decisions are based on clinical activity, severity and duration of GO (8, 14) as anti-inflammatory/immunosuppressive treatment is significantly less effective after 18 months of disease duration (15). **[Recommendation #1]**

Activity: The clinical activity score (CAS) is the best validated scoring system, although it has limitations, such as its binary (yes/no) feature, and should be therefore used for assessing activity (16). CAS is composed of seven items: GO is defined as active if CAS is $\geq 3/7$ (Table 2). A 10-item CAS, including increase in exophthalmos ≥ 2 mm, a decrease of eye movements in any direction of gaze $\geq 8^\circ$, and a decrease of visual acuity ≥ 1 line on the Snellen chart during a period of 1-3 months, is useful to evaluate recent progression and therefore activity of GO (16). A picture atlas to ensure consistent CAS assessment has been published (17). Other inflammation scores, such as the VISA score (18) may be useful, but an adequate validation is not available. Specific magnetic resonance imaging (MRI) sequences may be useful in quantifying disease activity and predicting response to anti-inflammatory treatment and outcome of GO (19-21); however costs and availability substantially limit MRI application in daily practice.

Severity: The EUGOGO classification into mild, moderate-to-severe, and sight threatening GO (Table 3) has been validated in clinical and research studies (8, 22) and should therefore be used. Other

scoring systems i.e. VISA (18), NOSPECS (23, 24), total eye score (25) allow quantification. MRI and CT scans provide information on the amount and distribution of orbital tissue expansion (muscle thickening, fat volume increase, apical crowding). Overall, orbital MRI is indicated in patients with unilateral or strongly asymmetric exophthalmos, suspected optic neuropathy, and euthyroidism with normal thyroid serology, while orbital CT is indicated prior to decompression surgery (19-21).

EARLY REFERRAL TO SPECIALIZED CENTERS

In the last 30 years a reduction in the incidence of GO in GD patients, as well as of its severity when present, has been reported (2, 3, 26) and recently confirmed by meta-analyses and meta-regression of published studies (27). This secular trend is multifactorial in origin (e.g., decrease in smoking habits, earlier diagnosis and better control of thyroid dysfunction). Improved interaction between endocrinologists and ophthalmologists lead to early diagnosis and treatment. In addition, mild GO can progress to more severe disease requiring expert advice and guidance for a general management plan (28). Therefore, it is fundamental to refer patients with overt GO and those at risk for deterioration of GO (mild and active GO, smokers, severe/unstable hyperthyroidism, high serum level of thyrotropin receptor antibodies [TSHR-Ab]) (8, 29) to thyroid-eye clinics, namely specialized centers providing combined endocrine and ophthalmic expertise (30, 31), as this will provide accurate and timely diagnosis to improve prognosis and QoL. Indeed, GO patient satisfaction was greater in those who attended such clinics (32). Primary care physicians, general practitioners, internists, endocrinologists or general ophthalmologists can manage the mildest cases without risk factors, unless progression occurs. **[Recommendation #2]**

ASSESSMENT OF TREATMENT OUTCOMES

Evaluation of treatment outcome should be standardized by using both a subjective primary outcome (*patient-reported outcome, PRO*) and an objective primary outcome (*clinician-reported*

outcome, CRO), assessed at a fixed time interval after the end of intervention. The preferred PRO is the validated disease-specific GO-QoL questionnaire (33, 34). The most appropriate CRO depends on the type of intervention. For moderate-to-severe and active GO a recently revised composite index is suggested (30). It is composed of entirely objective measures: ≥ 2 -mm reduction of lid aperture, ≥ 1 point reduction in 5-item CAS (excluding subjective, patient-reported spontaneous or gaze-evoked pain), ≥ 2 mm reduction in exophthalmos, $\geq 8^\circ$ increase of eye muscle duction (34). Improvement in ≥ 2 features in one eye without deterioration in the other eye might be considered a positive response to treatment (34). Other individual ocular, serological and imaging features can be included as secondary outcomes, including exophthalmos, eyelid aperture, ocular motility, visual acuity, CAS, intraocular pressure, orbital volume assessment, MRI, and TSHR-Ab measurement. Optimally, outcome of treatment should be assessed three months after the last therapeutic intervention (34), but, in addition, changes after six months can also be considered.

GENERAL MEASURES FOR ALL PATIENTS [Recommendations #3-6]

Control of risk factors.

When a patient with GD is seen, regardless of the clinical phenotype, every effort should be made to remove risk factors in order to prevent *de novo* occurrence and/or progression of GO. Adequate **control of thyroid dysfunction** is of paramount importance. Both hyper- and hypothyroidism negatively impact GO (35-37). In line with the expression of the TSHR as autoantigen on orbital target cells (38, 39) in patients with GD/GO, high serum concentrations of **TSHR-Ab** (>five-fold increase) are associated with the presence of GO both in children and adults with GD and Hashimoto's thyroiditis (40-43). Although neither regularly done in commercial laboratories nor routinely available to the clinician, dilution analysis of serum TSHR-Ab is both predictive for the occurrence of GO (positive and negative predictive values of 100% with a cut-off dilution titer >4)

(29), as well as for the response to antithyroid treatment of associated Graves' hyperthyroidism (42, 44).

All patients with GD, irrespective of the presence of GO, should be urged to **quit smoking**. The association between GO and smoking is evidence-based (3, 45, 46). Smoking increases the risk of GO in patients with GD (3); smokers have more severe GO (3); development or progression of GO after radioactive iodine (RAI) treatment is more frequent in smokers (47, 48); smokers have a delayed or worse outcome of immunosuppressive treatments (47, 49, 50); and smoking cessation is possibly associated with a better outcome of GO (51).

RAI bears a consistent risk of causing progression and/or *de novo* occurrence of GO (48, 52-55). In a large RCT, progression of GO occurred in 23 of 150 patients given RAI (15%), being persistent in eight (5%), hence requiring immunosuppressive treatment for GO (54). Both *de novo* occurrence and progression of GO are more likely in smokers (47, 48), in patients with duration of GD <5 years (56), and less likely in patients with long-standing and inactive GO (37). RAI-associated progression of GO can be prevented by a concomitant short-term course of oral prednisone (52, 54, 56, 57). The original regimen used a starting daily dose of 0.3-0.5 mg/kg/bodyweight, gradually tapered and withdrawn after three months [47]. Lower doses of oral prednisone (0.1-0.2 mg/kg/bodyweight as starting dose, gradually tapered and withdrawn after six weeks) (58) showed similar beneficial effects. As previously recommended [7], the 0.3-0.5mg dose should be used in patients who are at risk for progression and/or *de novo* development of GO (smokers, high TSHR-Ab levels, severe hyperthyroidism, preexisting GO). Prophylaxis using low-dose intravenous (iv) glucocorticoids has been proposed, but this requires one day of hospitalization per week for four weeks (56). Glucocorticoid prophylaxis is not only effective, but also safe (59).

High cholesterol is an emerging and potential risk factor for GO (60). The use of statins was associated with a reduced risk of GO occurrence in a large cohort retrospective study (61) and in a retrospective registry-based study (62). Association of high total and LDL-cholesterol with the

presence of GO was reported in one cross-sectional and one retrospective study (63, 64). Finally, in a retrospective study, the outcome of GO following iv glucocorticoid treatment was worse in patients with high LDL-cholesterol (65). These findings may reflect a pro-inflammatory action of cholesterol. Alternatively, they might be related to an anti-inflammatory effect of statins, irrespective of cholesterol levels. RCTs are lacking, but control of hypercholesterolemia by statins may be considered in patients with GO.

Local treatments.

Ocular surface inflammation and dry eye are frequent in GD patients; regardless of the presence of overt GO (66-68). In GO patients, several factors contribute to drying of the eye, i.e., increased width of the palpebral fissure, exophthalmos, blinking rate, lid lag, lagophthalmos, poor Bell's phenomenon due to restrictive elevation deficit, and altered tear film osmolality (69-71). Treatment with artificial tears during the day and ophthalmic gels/ointments with possible taping of the lids or using swimming goggles at nighttime, when severe lagophthalmos is present in the absence of an adequate Bell's phenomenon are recommended to GO patients since first observation and to patients with GD without overt GO but with dry-eye symptoms (72). Botulinum toxin injection in the levator muscle may reduce palpebral aperture (73).

MANAGEMENT OF MILD GO

Most patients with mild GO experience spontaneous resolution of eye manifestations. Therefore, a watchful strategy and local treatments are sufficient (8) (figure 1). Patients living in selenium-deficient areas can benefit from oral selenium supplementation. A randomized, double-blind, placebo-controlled trial of patients with mild GO, performed in Europe, reported a higher rate of improvement in both GO-QoL and overall ophthalmic outcome and a lower rate of progression to more severe GO in patients receiving sodium selenite (200 µg [91.2 µg selenium] daily for six months), compared to the placebo group (74). The benefit of selenium was maintained six months

after treatment withdrawal. According to country availability, sodium selenite can be replaced by seleniomethionine (100 µg daily). Whether selenium administration is of benefit in selenium-replete areas has to be confirmed. An European survey has shown that selenium supplementation is recommended by the majority of clinicians, both in patients with mild and moderate-to-severe GO (75), but there is no evidence of a beneficial adjuvant effect of selenium in patients with moderate-to-severe and active GO.

While a wait-and-see strategy is feasible in the majority of patients with mild GO, a very few patients may experience or develop a profound impact on QoL: in these exceptional patients, low-dose immunomodulation may be proposed if GO is active or rehabilitative surgery if GO is inactive, subsequent to extensive counselling and shared-decision (8). **[Recommendations #7-8]**

MANAGEMENT OF MODERATE TO SEVERE AND ACTIVE GO

After an initial phase in which inflammation and its manifestations are predominant (active phase), GO stabilizes (plateau phase) and then slowly remits leaving typical residual signs and symptoms (inactive phase); the whole process (natural history) is believed to last 18-24 months in untreated patients (3). In patients with moderate-to-severe and active GO the initial goal is to shorten the active phase of the disease and improve subjective and objective eye manifestations. Results of treatment are usually better if GO is treated early, within one year from its onset. Efficacy of immunosuppressive therapy varies between 50 and 80% according to the published trials (76), but rarely leads to a complete satisfactory response. Residual inactive disease benefits from rehabilitative surgery. Non-responders may require a second course of immunosuppressive therapy using different drugs/treatments, alone or in combination. Few patients still remain unresponsive or partially responsive and will need a surgical approach.

Extensive counselling is required when discussing the treatment plan. The patient should be informed that he/she would be engaged in a complex journey, clearly explaining benefits and risks

of the various therapeutic options. Response to initial therapy cannot be predicted. The patient should be aware that additional medical and/or surgical treatments might be needed, particularly rehabilitative surgery, even in responders, in an attempt to restore the pre-disease eye appearance and function. Selection of treatment follows several considerations, e.g. evidence-based effectiveness, short term/long term safety, evaluation of costs, reimbursement by the health system, drug-availability, infrastructure facilities for delivering highly specialized treatments and personal choice of the informed patient within a shared decision-making process.

[Recommendation #9]

Systemic and locally injected glucocorticoids [Recommendations #10-14]

High-dose systemic glucocorticoids have potent anti-inflammatory and immunosuppressive effects (77, 78) that have been applied successfully for the management of moderate-to-severe and active GO. Intravenous glucocorticoids have been indicated as the first-line treatment in moderate-to-severe and active GO (22, 79). A proof-of-concept RCT showed a significant improvement of GO outcome in patients treated with iv methylprednisolone compared to placebo (response rate 83% vs 11%) (80). Although oral glucocorticoids are effective, glucocorticoids are preferentially administered iv as the iv route has been shown in RCTs to be more effective (77-88% vs 51-63%) and better tolerated (81, 82). The most common protocol employs a cumulative dose of 4.5 g methylprednisolone, given in 12 weekly infusions (six infusions of 0.5 g, followed by six infusions of 0.25 g) (82). This 4.5 g regimen is very-well tolerated (83) and significantly improves QoL (84). While this regimen is appropriate for most patients, a higher cumulative dose of 7.5 g (starting with 0.75 g as a single iv dose) is reserved for more severe cases within the spectrum of moderate-to-severe and active GO, as the higher dose bears a higher risk of drug-induced adverse events (AEs) (85). Safety data suggest that, with the exception of sight threatening GO, single iv doses should not exceed 0.75 g, cumulative doses should be less than 8.0 g per cycle, and consecutive-day therapy

should be avoided, because these schedules are associated with a significantly higher rate and clinically relevant glucocorticoid-induced AEs, including liver toxicity and serious cardiovascular AEs (86-90).

Infusions should be performed slowly (1-2 hours) under strict surveillance. Therefore, prior to starting treatment and after ruling out infections (white blood cell count), cardiovascular risk, liver enzymes and markers of viral hepatitis are evaluated, in order to assess risks and contraindications (8). In addition, liver enzymes are closely monitored during treatment (89). Recent viral hepatitis, significant hepatic dysfunction, severe cardiovascular morbidity, or psychiatric disorders represent absolute contraindications to iv glucocorticoid treatment (78, 79), while diabetes and hypertension should be well controlled before starting treatment (8). Bone protection is recommended, and proton pump inhibitors used as appropriate (8). Response to iv glucocorticoids usually occurs early, but it may be delayed to the second half of treatment course (91). This is why partial responders to iv glucocorticoids should be offered to complete the 12-week regimen. In contrast, a clinical deterioration of clinical ophthalmic signs and symptoms requests a shift to second-line treatments (8, 91).

Intravenous glucocorticoid treatment needs to be carried out in specialized centers and facilities that may not be easily available in all countries. This partly explains why the oral route is still widely used either alone (92) or after initiating a treatment course with a few iv infusions (93) to reduce hospital admissions. In the case of oral glucocorticoids, treatment should start, as suggested by several RCTs (77, 81, 82), either with a fixed dose of 100 mg prednisone/prednisolone or, preferably, 1 mg/kg bodyweight, and be gradually tapered down by 5-10 mg/week until withdrawal (4-6 months). Combination with other treatments, including orbital radiotherapy or non-steroidal immunosuppressive drugs, (i.e. mycophenolate or cyclosporine) may work as a steroid-sparing procedure and increase the effectiveness of oral glucocorticoids.

Local (subconjunctival or parabolbar) glucocorticoid administration has been used in a few patients. In a RCT, retrobulbar injections of methylprednisolone acetate were less effective than systemic glucocorticoids when combined with orbital cobalt radiotherapy (94). In a prospective, single blind, placebo-controlled RCT, orbital injections of triamcinolone acetate into the inferolateral quadrant (4 weekly injections of 40 mg) reduced diplopia and extraocular muscle size (95). Furthermore, in a small RCT, subconjunctival upper eyelid injections of triamcinolone (1-3 injections of 20 mg) were reported to be effective for treatment of upper eyelid retraction in patients with short duration of GO (96). However, local glucocorticoid treatment has a significant risk of intraocular pressure elevation, may be associated with increased orbital lipomatosis, and bears a small but significant risk of retrobulbar hemorrhage, especially in patients with dual platelet inhibition (97, 98). Hence, local glucocorticoids may be considered in patients with contraindications to systemic administration of glucocorticoids only.

Mycophenolate [Recommendation #15]

Mycophenolate competitively and reversibly inhibits inosine monophosphate dehydrogenase, resulting in decreased antibody production by B cells and dual antiproliferative effect on both B- and T-cells (99). Mycophenolate induces apoptosis of activated T-cells, inhibits expression of adhesion molecules and recruitment of immune cells (100). In addition, mycophenolate inhibits fibroblast proliferation and functions (101-104). The drug is available worldwide as mycophenolate mofetil and/or enteric-coated mycophenolate sodium (105). Fractionated doses per day taken with the meals improve gastro-intestinal tolerance. A systematic review of the gastro-intestinal side effects between the two formulations did not demonstrate significant differences between gastro-intestinal-related QoL for patients using either form as maintenance immunosuppression (106). In a single-center trial, 174 euthyroid patients with moderate-to-severe and active GO were randomized to either an unusual combination of three infusions of iv glucocorticoids followed by

oral glucocorticoids, or mycophenolate mofetil (1g daily), both for 24 weeks (107). Mycophenolate mofetil demonstrated superior overall response rate (79%/91% at week 12/24 vs 51%/68% in the glucocorticoid group). Disease inactivation was observed in 94% of mycophenolate mofetil group (vs 69% in the combined glucocorticoid group) at week 24. Mycophenolate mofetil also performed well in proptosis and diplopia. Six percent of glucocorticoid patients developed disease reactivation, while none in the mycophenolate mofetil group relapsed.

In the EUGOGO's observer-masked multicenter trial (108, 109), 164 euthyroid patients with moderate-to-severe and active GO were randomized to weekly iv methylprednisolone for 12 weeks or a combination of iv methylprednisolone for 12 weeks and mycophenolate sodium 0.72 g daily (which is equivalent to 1g mycophenolate mofetil/day) for 24 weeks. In the intention-to-treat population at 12 weeks, responses were observed in 36 (49%) of 73 patients in the monotherapy group and 48 (63%) of 76 patients in the combination group, giving an odds ratio (OR) of 1.76 (95% CI 0.92–3.39, $p=0.089$). At week 24, 38 of 72 patients remaining in the monotherapy group and 53 of 75 patients remaining in the combination therapy group had responded to treatment (OR 2.16, 1.09–4.25, $p=0.026$). At week 36, 31 of 68 patients in the monotherapy group and 49 of 73 patients in the combination group had a sustained response (OR 2.44, 1.23–4.82, $p=0.011$). Thus, the combination group displayed statistically significant superior response rate at week 24 (71% vs 53%) and sustained response rate at week 36 (67% vs 45.5%). Overall, combination treatment demonstrated more significant improvements in CAS, swelling of eyelids and caruncle, orbital pain, chemosis, downgaze duction and elevation, as well as GO-QoL visual functioning score.

Evaluating both randomized trials, mycophenolate-treated groups demonstrated superior response rates at 12 weeks (107), 24 weeks (107, 108) and 36 weeks (108) when compared to their respective glucocorticoid monotherapy groups. Approximately 70% (versus 90% in the mycophenolate group) and 30% (versus 60-70% in the mycophenolate group) of patients achieved endpoints in most individual visual parameters of activity and severity, respectively. In addition, the mycophenolate

sodium + glucocorticoid group of the EUGOGO trial performed better than mycophenolate alone (107) in terms of improvement of pain and eye movement. However, longer-term follow-up and subsequent rehabilitative surgery data are currently not available and may be regarded as limitations of both RCTs.

Higher mean age, more prevalent smoking habit, longer disease duration, and a greater proportion of TSHR-Ab-positive patients may explain the lower response rates in the EUGOGO trial. Neither trial reported any serious infection or treatment-related mortality (110, 111). The combination treatment did not increase the risk of infection and hepatotoxicity when compared to iv methylprednisolone monotherapy. Furthermore, “real-world” efficacy and safety of mycophenolate mofetil in patients with active moderate-to-sight-threatening GO was demonstrated over a 4-year observation period (112). Therefore, the risk-benefit ratio of low dose mycophenolate, either as monotherapy or in combination with iv glucocorticoids treatment in active moderate-to-severe GO, is highly favorable given its reassuring safety profile and promising efficacy (110, 111). Hence, the combination of low dose mycophenolate sodium and iv methylprednisolone was both safe and affordable in view of its superior efficacy compared to current standard of care.

Orbital Radiotherapy [Recommendation #16]

Orbital radiotherapy for GO has been shown by several RCTs to be more effective than sham irradiation in improving diplopia and ductions (113, 114), although its efficacy was questioned by two additional RCTs (115, 116). In another RCT, orbital radiotherapy was found to be as effective as oral prednisone (117), and other RCTs have shown that orbital radiotherapy synergistically potentiates the effects of oral glucocorticoids (118, 119). RCTs showing that this synergistic effect holds true using iv glucocorticoids are missing. However, two retrospective studies showed that combination of orbital radiotherapy and iv glucocorticoids was more effective than iv glucocorticoids alone in improving eye motility and reducing GO severity (120, 121). Usually, a 20

Gray (Gy) cumulative dose per orbit fractionated in 10 daily doses over 2-weeks is given (122). However, a regimen of one Gy per week over 20-weeks was shown to be equally effective and better tolerated (123). Mild and transient exacerbation of ocular symptoms may occur during orbital radiotherapy, which is controlled by concomitant administration of low-dose oral prednisone. Although orbital radiotherapy is safe (124-127), it should be avoided in patients with hypertensive or diabetic retinopathy, or, in view of a remote carcinogenetic risk, in patients younger than 35 years (8). In a 17-year long-term follow-up study comparing a single dose of one Gy vs. two Gy, radiation-induced retinopathy was observed in 5% of patients with GO, diabetes mellitus and hypertension approximately 10 years after orbital radiotherapy, however, in none of those irradiated with the lower dose (128). In summary, orbital radiotherapy is effective, particularly on ocular motility, and safe, being devoid of major adverse events even after a long-term follow-up (122).

Cyclosporine [Recommendation #17]

Cyclosporine is a potent immunosuppressive agent that inhibits calcineurin pathway reducing T-cell proliferation and IL-2 secretion. Two small, early RCTs have assessed its efficacy in patients with moderate to severe GO. The combination of cyclosporine (initial dose: 5-7.5mg/kg bodyweight/day) and oral prednisolone (initial dose: 50-100mg/day) had a better ophthalmic outcome and a lower relapse rate than oral prednisolone monotherapy (129). In another study (130), significantly fewer patients responded to cyclosporine (7.5mg/kg bodyweight/day) as compared to oral prednisolone (initial dose 60mg/daily) (22% vs 61%). However, more than half of the non-responders to either drug alone showed subsequent improvement with prednisolone-cyclosporine combination confirming the potential beneficial effect of a combined cyclosporine-glucocorticoid treatment and its superiority versus either cyclosporine or oral glucocorticoid monotherapy. Of note, there are no RCTs comparing cyclosporine with iv glucocorticoids for the treatment of GO.

Azathioprine [Recommendation #18]

Azathioprine is an antiproliferative agent with a similar mode of action to mycophenolate, frequently used as a “steroid-sparing agent” in autoimmune/inflammatory conditions. Azathioprine was ineffective in GO as a single agent (131), but observational studies suggested benefit in combination with low-dose glucocorticoids (132). In the randomized and blinded CIRTED study (n=126), azathioprine over 12 months was studied in combination with high-dose oral glucocorticoids and orbital radiotherapy in a factorial designed trial (116). Although the majority of patients returned for primary end-point review (82%), 66% of participants allocated to azathioprine and 45% of those allocated to placebo did not complete the full 48 weeks of treatment. Withdrawals from the azathioprine group were for known AEs of azathioprine. Despite low adherence rates, in intention-to-treat analyses the point estimate for the OR for improvement in patients treated with azathioprine was substantial (2.56, 95%CI 0.98–6.66; $P=0.054$) and a sensitivity analysis in which patients who withdrew during the trial were recoded to unfavorable outcomes regardless of their status at 48 weeks, the effect of azathioprine treatment was enhanced (3.65, 95% CI 1.34–9.86; $P=0.011$). In addition, in a post-hoc per protocol analysis of patients who completed their allocated therapy the OR for improvement on azathioprine was large (6.83, 1.66–28.1; $P=0.008$). The major benefit was a reduction in the relapse rate after glucocorticoid withdrawal. Hence, azathioprine may be a valuable steroid-sparing agent when continued after an oral glucocorticoid taper, although it is frequently not well tolerated. Benefits in combination with iv glucocorticoids are unknown.

Teprotumumab [Recommendation #19]

The insulin-like growth factor-1 (IGF-1) receptor is over-expressed in GO orbital fibroblasts and lymphocytes (133, 134). It forms a functional complex with the TSHR, and mediates TSHR

downstream signaling (135). Teprotumumab is a fully humanized immunoglobulin (Ig) G1 monoclonal inhibiting antibody, which binds to the extracellular portion of IGF-1R and blocks its activation and signaling by endogenous ligands. Binding of teprotumumab also leads to internalization and degradation of IGF-1R resulting in up to 95% reduction of accessible receptor protein on the cell surface (135).

The safety and efficacy of teprotumumab was evaluated sequentially in two RCTs, which comprised 170 patients with moderate-to-severe and active GO (136, 137). Both trials had similar designs and patients were randomly assigned to teprotumumab (83 patients; once every three weeks iv for eight doses over 24 weeks) or placebo (87 patients). Seventy-three percent in teprotumumab groups (vs 14% in placebo groups) were overall responders with both CAS and proptosis improvement. Individually, CAS of 0-1 (62% vs 22%) and proptosis response (77% vs 15%) were much more common in teprotumumab groups. Proptosis response occurred early at week six in most patients. The mean reduction in proptosis by week 24 ranged from 2.9 to 3.3 mm. Teprotumumab treatment was also associated with a significant improvement in GO-QoL score. Recently, systematic analyses and off-treatment follow-up results from the two RCTs were published (138). One year after final dose, integrated proptosis, diplopia, and composite responses were 67%, 69%, and 83%, respectively. In the teprotumumab group, orbital decompression surgery was required in a few patients during the follow-up observation period and dysthyroid optic neuropathy (DON) occurred in one patient four months after the last teprotumumab infusion. The most common AEs reported with teprotumumab included muscle spasms (25%), nausea (17%), alopecia (13%), diarrhea (13%), fatigue (10%), hearing impairment (10%), and hyperglycemia (8%). Also, Teprotumumab is contraindicated for those with inflammatory bowel disease and in pregnancy. In the systematic analysis, most teprotumumab AEs were mild-moderate during treatment, with three related serious AEs (diarrhea, infusion reaction and Hashimoto's encephalopathy/confusion) leading to study discontinuation. In line with this, a case of serious teprotumumab-induced amyloid encephalopathy

was recently reported, which was unresponsive to high-dose glucocorticoids or immunoglobulin G therapy, but remitted after plasmapheresis (139). The current dosing regimen of teprotumumab has proven effective for GO, however dose ranging studies including variable concentrations, infusion frequencies, and durations of teprotumumab therapy in the setting of GO have not been performed (140). Therefore, although teprotumumab has become the first drug approved by the US Food and Drug Administration for the treatment of adult GO, its incorporation into routine clinical practice is currently limited by the lack of comprehensive long-term efficacy and safety data, absence of head-to-head comparison with iv glucocorticoids, restricted geographical availability, reimbursement (outside the US), and costs.

Rituximab [Recommendation #20]

Rituximab is a chimeric human and mouse monoclonal antibody against CD20 surface antigen expressed on B cells that causes immunosuppression through B-cell depletion. After several retrospective case-series (141-143) suggested a potential benefit of rituximab for the treatment of GO, two double blind, but low-powered single-center RCTs have evaluated rituximab in patients with moderate-to-severe and active GO with conflicting results. The US study (144) randomized 25 patients to receive two infusions of either rituximab (1000 mg each) or placebo (saline) two weeks apart: no additional advantage of rituximab over placebo was found in reducing CAS or severity of GO at 24 or 52 weeks. In contrast, the Italian study (145) demonstrated better ophthalmic and QoL outcomes with rituximab as compared to iv glucocorticoids: 32 patients were randomly assigned to receive either rituximab (two doses of 1000 mg two weeks apart or a single dose of 500 mg) or iv glucocorticoids (cumulative dose, 7.5 g). At 24 weeks, all patients treated with rituximab showed inactivation of GO as compared to 69% in the iv glucocorticoid group. At 52 weeks, none in the rituximab group and 31% in the iv glucocorticoid group had reactivation of GO (145). As compared

to the US study (144), participants in the Italian trial had a shorter average duration of GO (4.5 vs 12.2 months), which may explain the discrepant results.

In the above studies, DON developed in two patients and vasculitis in one rituximab-treated patient (144). A severe cytokine release syndrome presenting with marked periorbital edema and decrease of vision, then controlled with glucocorticoids, occurred in two patients (145). A recent non-randomized prospective study of 17 patients (8 steroid naive, 9 unresponsive to iv glucocorticoids) has suggested an efficacy of a low dose (100-mg single infusion) of rituximab in moderate-to-severe and active GO (146). Over 90% of these patients showed disease inactivation by 12 weeks, and no patient had reactivation after a 76-week follow-up. Mild infusion related AEs frequently seen with the higher dose of rituximab were rare; nevertheless, one patient developed a cytokine release syndrome (146).

Tocilizumab [Recommendation #21]

Tocilizumab is a humanized monoclonal antibody against the interleukin (IL)-6 receptor approved for use in rheumatoid arthritis. In addition to playing a role on T and B cell activation as pro-inflammatory cytokine, IL-6 also acts directly on orbital pre-adipocytes to promote volume expansion (147). In an RCT of GO patients considered to have failed initial glucocorticoid therapy (n=32), subjects treated with iv tocilizumab monotherapy on week 0, 4, 8, 12 showed greater reductions in CAS (86% achieving CAS <3 vs 35% in the placebo group, p<0.005) at week 16 (148). Tocilizumab was generally well tolerated, but there was a higher rate of infections and headache in the tocilizumab group; benefit was predominantly on soft tissue signs (148). In an uncontrolled observational study of 48 patients, resistant to established therapies (mostly iv glucocorticoids, 90%), tocilizumab given monthly iv or weekly subcutaneously was well tolerated and the majority of patients showed improvement (92%) (149). Likewise, a small study of eight glucocorticoid-resistant patients with moderate-to-severe and active GO, showed a beneficial effect of tocilizumab

on CAS and exophthalmos (150). Similar findings were recently reported in a single-center retrospective observation over nine years with 54 patients analyzed (151), nevertheless larger RCTs including tocilizumab in naive patients with GO of short duration are warranted. Currently, data suggests that tocilizumab may cause rapid resolution of inflammatory signs in glucocorticoid-resistant moderate-to-severe and active GO.

Other Immunomodulators

Iv immunoglobulin treatment resulted in decreased specific autoantibody titers and clinical improvement in several autoimmune diseases (152). The random iv administration of anti-idiotypic immunoglobulins (1g/kg/bodyweight) in patients with moderate-to-severe and active GO was as effective as (62% response rate) and better tolerated than oral prednisolone (153). High costs, the need for iv administration, and its small potential risk for transmitting infectious agents limit the routine use of iv immunoglobulins in the treatment of GO. In comparison, when randomly tested in moderate-to-severe and active GO, ciamexone did not show any beneficial effect vs placebo (154). Also negative were four RCTs evaluating somatostatin-analogs (either octreotide or lanreotide) in GO (155-158). The level of tumor necrosis factor-alpha (TNF α) is elevated in GO patients compared to controls (159, 160). However, selected anti-TNF α agents tested in uncontrolled, small studies of GO patients had limited efficacy (161-163). Adding methotrexate to iv glucocorticoids in an uncontrolled small study in patients with active GO was safe and allowed reduced administration of glucocorticoids without compromising their efficacy (164).

First-line and second-line treatments for moderate-to-severe and active GO

When reviewing all currently published RCTs and evaluating pros and cons, efficacy and safety, as well as comparing the present evidence-base reports with those available six years ago, the

following first-line treatments (figure 2) and alternative second-line approaches (Figure 3) are recommended:

First-line treatments: As demonstrated by two large RCTs (107, 108) including more than 300 patients with moderate-to-severe and active GO and involving mycophenolate and glucocorticoids, this combination therapy shows a beneficial efficacy/safety profile (110) with a statistically significant and clinically relevant higher benefit than iv glucocorticoid monotherapy. Hence, and as shown in Figure 2, the combination of iv methylprednisolone (moderate cumulative dose of 4.5 g over 12 weeks) + mycophenolate sodium 0.72 g per day for 24 weeks is recommended as first-line treatment for most patients with moderate-to-severe and active GO. If the enteric-coated mycophenolate sodium is not available, mycophenolate mofetil is administered (1 g of mofetil is equivalent to 0.72 g of the sodium formulation).

In the most severe forms (including constant/inconstant diplopia, severe soft-tissue signs) within the spectrum of moderate-to-severe and active GO, a higher cumulative dose of iv methylprednisolone (7.5 g) as monotherapy is also recommended as alternative first-line approach. RCTs comparing the higher dose (7.5 g) in combination with mycophenolate to iv glucocorticoid monotherapy are not available.

Second-line treatments: As already recommended in the 2016 ETA/EUGOGO guidelines (8) and subsequent to a careful ophthalmic and biochemical (liver enzymes) evaluation, after 3-4 weeks a second course of iv methylprednisolone monotherapy administering the higher cumulative dose of 7.5 g and starting with single doses of 0.75 g for six weeks is a further acknowledged and valid second-line therapy. Of note, a cumulative dose of 8 g iv methylprednisolone per cycle is allowed. Alternatively, the combination of oral prednisone/prednisolone and cyclosporine are recommended, as two RCTs demonstrated the benefits of this combination (8, 129, 130). In addition, azathioprine can be used together with oral glucocorticoids because of its steroid-sparing action in one RCT (Figure 3).

The largest clinical experience as of today for an alternative evidence-based therapy is with the combination of oral prednisone/prednisolone and orbital radiotherapy (118, 119). However, in view of the evidence that iv glucocorticoids are more effective and better tolerated than oral glucocorticoids [1], and that iv glucocorticoids combined with orbital radiotherapy are more effective than oral glucocorticoids combined with orbital radiotherapy [83], we suggest that orbital radiotherapy combined with iv glucocorticoids can be considered a second-line therapy (expert opinion), particularly in patients with eye muscle dysfunction, although a prospective RCT directly comparing iv glucocorticoid monotherapy vs. iv glucocorticoids combined with orbital radiotherapy is not available and an evidence-based recommendation for a single glucocorticoid dosage is missing.

Rituximab can be recommended as second-line treatment at the dosage reported in one RCT (one 500 mg shot) (145) or at lower dose (one 100 mg shot) [145], however not in patients with a potential risk for DON (144, 145). Rituximab is the only drug with a head to head comparison of iv glucocorticoids (145). Tocilizumab may also be considered in glucocorticoid-resistant patients, though robust data concerning efficacy and safety are still missing. Finally, Teprotumumab, as a promising and effective drug for GO (136, 137), is currently available in the US only. Subsequent to its clearance by the European Medicine Agency, publication of long-term efficacy and safety data, information on need for post-teprotumumab rehabilitative surgery, and, optimally, subsequent to head-to-head comparison with iv glucocorticoids within a future RCT, teprotumumab will likely play a relevant role in the management of patients with moderate-to severe and active GO provided it is available and affordable to each patient [**Recommendations #22-25**].

Overall and based on the features of GO that they are most likely to be effective for, intravenous glucocorticoids, mycophenolate, tocilizumab, rituximab, and cyclosporine substantially decreased inflammatory ophthalmic signs while orbital radiotherapy (preferably in combination with

glucocorticoids) significantly improved eye muscle motility and/or diplopia. In comparison, Teprotumumab showed the strongest effect on exophthalmos.

MANAGEMENT OF SIGHT THREATENING GO

Sight threatening GO is an emergency that is treated immediately. Impairment or loss of vision can be due to DON, severe corneal exposure breakdown, and, in rare cases, eyeball subluxation causing acute optic neuropathy due to stretching of the optic nerve, increase in the intraocular pressure, and/or corneal breakdown. A small RCT (165) showed that in DON patients immediate decompression did not result in a better outcome compared to iv glucocorticoids given as first-line treatment. In a retrospective study of 24 DON patients (40 eyes), more than 40% of patients showed permanent restoration of normal visual function after high-dose iv methylprednisolone therapy (88). Based on these results, the first line treatment of DON is high-dose (single doses of 500 to 1000 mg) iv methylprednisolone for three consecutive days or more preferably (for safety reasons) (86, 90) on every second day (alternate days) during the first week, which can be repeated for another week (figure 4). When the response is absent or poor with a deterioration in visual acuity or visual fields, urgent orbital decompression surgery is mandatory. Blepharorrhaphy, tarsorrhaphy, lid lengthening, extraocular muscle recession, gluing, antibiotics, and transplantation are used to protect the cornea when severe corneal exposure (ulceration) or corneal breakdown occurs. Subluxation of the eyeball is treated with orbital decompression [**Recommendations #26-27**].

ORBITAL/OPHTHALMIC SURGERY IN THE TREATMENT OF GO

In the active phase of GO, decompression surgery is indicated in patients with severe exposure keratopathy and, as second-line treatment, in patients with DON not responding to iv

glucocorticoids. Local treatment (tarsorrhaphies, corneal patches or gluing) can be used in the same phase as temporary measures to shield the cornea for superficial damages or to correct extreme corneal thinning, thus decreasing risks of spontaneous eyeball perforation or perforation in the course of subsequent decompression surgery. In the post-inflammatory, inactive phase, residual disfigurements (exophthalmos, lid retractions, eyelid and periorbital puffiness, strabismus and correlated symptoms such eye grittiness, retro/peri-ocular tension, and diplopia) can be treated by a combination of decompression, ophthalmic plastic and strabismus surgery (166).

TREATMENT OF HYPERTHYROIDISM IN PATIENTS WITH GO [Recommendations #28-32]

Graves' hyperthyroidism can be managed by thionamide antithyroid drugs (ATDs), RAI treatment, or, less frequently, total thyroidectomy (Tx) (42, 167). ATDs and Tx *per se* do not modify the natural history of mild GO (53, 54, 168), although RCTs on moderate-to-severe GO are lacking. Long-term ATD treatment is beneficial for GO due to normalization of thyroid function and associated decline of TSHR-Ab serum levels (169), which are a biomarker for GO (43, 170). Avoiding iatrogenic hypothyroidism in treating patients with GD/GO (Figures 1-4) is an important principle of medical management. With this respect, within a large EUGOGO prospective multicentre observational cohort study of 344 patients with Graves' hyperthyroidism, the prevalence of biochemical euthyroidism during treatment with ATDs was higher during the ATD titration regimen compared to the „block and replace“ regimen. *De novo* development of GO did not differ significantly between the two regimens (171).

A small RCT (172) and a retrospective case study (173) showed that early Tx may be associated with a better outcome of immunosuppressive treatment for moderate-to-severe and active GO. At variance, RAI treatment is associated with a definite risk for GO, which can be prevented, in patients at risk, by oral low-dose and short-term prednisone prophylaxis, given concomitantly with RAI treatment, and prompt correction of post-RAI hypothyroidism. Total thyroid ablation (i.e., Tx

followed by RAI ablation of thyroid remnants) has been proposed, and two RCTs have shown a beneficial effect of this procedure following iv glucocorticoid treatment in patients with moderate-to-severe and active GO, in the short, but not in the long term (174-176).

The optimal treatment for hyperthyroidism in patients with GO is an unsolved dilemma, and there is no current evidence as to the superiority of the conservative approach (ATDs) with respect to the ablative approach (Tx, RAI, total thyroid ablation) or *vice versa* (177). If GO, either mild or moderate-to-severe is stably and longstanding inactive, any treatment for hyperthyroidism can be selected, based on standardized criteria and patient preference after extensive counselling (167), as it is unlikely to cause recurrence or progression of GO (37, 178). In patients with residual, but longstanding inactive moderate-to-severe GO, oral glucocorticoid prophylaxis can be considered if RAI treatment is selected and risk factors are still present. If GO is active and mild, ATD treatment (or Tx) is mostly preferred. RAI treatment can be used in combination with oral prednisone /prednisolone prophylaxis (167). When GO is moderate-to-severe and active, management of GO should be prioritized, because delayed treatment is associated with a lower response rate (8). While GO is being treated, hyperthyroidism is controlled by ATDs, possibly given longer than the usual 18-24 months (178, 179). Large thyroid glands (>50 ml) and/or nodular goiters can be surgically treated if mechanical signs of tracheal compression or suspicion of thyroid cancer are present. When GO is sight threatening, its emergency treatment either medical and/or surgical, is an absolute priority: hyperthyroidism is stabilized by ATDs until treatment of GO is completed (8).

MANAGEMENT OF GO DURING VIRAL PANDEMIC

Patients with mild GO should receive the usual local treatment (Figure 1) and should be urged to quit smoking to prevent progression of GO (180). As to moderate-to-severe and active GO, glucocorticoids and other immunosuppressive agents could make patients more susceptible to infections (1). Discontinuation of long-term glucocorticoid treatment may be associated with

adrenal failure, which, in turn, increases the risk of developing infections and related mortality, likely including COVID-19 (181). Nevertheless, unless immunised or having had COVID, all of us have no immunity to COVID, so adding immunosuppression will not increase that risk. Furthermore, dexamethasone or methylprednisolone or tocilizumab have now become the standard of care for COVID (182, 183). Also, it is now clear that very high dose steroids used in COVID does not cause adrenal suppression and normal adrenal function was observed in patients who survived COVID-19 Infection (184). In line with this, withdrawal of iv glucocorticoid treatment in GO patients is not associated with adrenal failure (185, 186).

No studies are available on the use of iv glucocorticoids or other immunosuppressive agents for GO during the current pandemic. With ongoing vaccination of the population, the risk related to immunosuppression will gradually decrease. Although the impact of immunomodulatory /immunosuppressive therapies on the efficacy of vaccination against COVID is not known, and steroids are known to decrease the efficacy of other vaccines, it seems reasonable to propose that patients already under treatment continue iv glucocorticoids or other immunosuppressive treatments under careful monitoring (180). Oral treatments can be continued at home, strictly following rules of social distancing, shielding and hygiene. Sight-threatening GO is an emergency and should be treated as such, irrespective of viral pandemic.

CONCLUSIONS AND PERSPECTIVES

It is reasonable to recommend the combination of iv methylprednisolone and mycophenolate (sodium) as the updated standard of care in moderate-to-severe and active GO, in view of its practicability and superior efficacy to weekly iv glucocorticoid monotherapy. Biologicals, especially teprotumumab and, to a lesser degree, tocilizumab or rituximab, hold great promise in future management of GO and can be useful if patients are intolerant or resistant to standard immunosuppressive treatment. However, they were not rigorously tested in large RCTs against

current standard of care, namely, iv glucocorticoids. The fact that they may not be widely available or affordable, as well as the lack of information, say, on the need of subsequent rehabilitative surgery, further add to their current limitations. As multiple pathogenic pathways are implicated in GO, several targeted therapies are worth exploring in clinical trials, e.g., monoclonal antibodies and/or small molecules targeting the TSHR (187, 188) or the CD40 molecule expressed in both thyrocytes and orbital fibroblasts (189), or anti-IL-23/anti-IL-17 for the IL-23/IL-17 axis and sirolimus for the mTOR pathway (190). Worthwhile is also a modulating impact on the microbiome in patients with GO (191). Overall, any novel therapeutic strategy in GO must be examined in RCTs, hopefully adopting the same assessment of treatment primary outcomes, before any conclusion regarding efficacy (i.e., proptosis and diplopia) and safety can be drawn.

ACKNOWLEDGMENTS

The following is the list of 48 EUGOGO members listed as collaborators / contributors to this guideline manuscript

Ankara (Turkey): Goksun Ayvaz, Onur Konuk; Belgrade (Serbia): Jasmina Ciric, Bijliana Beleslin; Brussels (Belgium): Antonella Boschi, Maria Cristina Burlacu; Cardiff (UK): Dan Morris; Catania (Italy): Rosario Le Moli, Antonio Marino; Edinburgh (UK): Justin McKee, Nicola Zammit; Essen (Germany): Dagmar Führer; Exeter (UK): Ioana Pereni; Göttingen: Michael Schittkowski, Dirk Raddatz; London (UK): Vickie Lee, Karim Meeran; Lyon (France): Juliette Abeillon, Thia Soui Thcong; Mainz: Katharina Ponto; Milan (Italy): Ilaria Muller, Nicola Currò; München (Germany): Christoph Hintschich, Roland Gärtner; Newcastle (UK): Simon Pearce, Lucy Clarke; Odense (Denmark): Thomas Brix, Dorte Bechtold; Olten (Switzerland): Gottfried Rudovfsky, Nicole Fichter; Paris (France): Laurence Du Pasquier, Julia Meney; Pisa (Italy): Francesca Menconi, Giulia Lanzolla; Singapore: Gangadhara

Sundar, Samantha Peiling Yung; Thessaloniki (Greece): Kostas Boboridis, Anagnostis Panagiotis; Valencia (Spain): Marta Pérez Lopez, Carlos Javier Sanchez; Varese (Italy): Maria Laura Tanda, Simone Donati; Vienna (Austria): Andrea Papp, Shuren Li; Warsaw (Poland): Anna Jablonska, Piotr Miskiewicz; Zagreb (Croatia): Jelena Juri Mandic, Maja Baretic.

Lara Frommer, MSc. and Jan Längericht (pharmacist), both JGU Thyroid Lab, Mainz, Germany are acknowledged for their editorial assistance.

DISCLOSURES

Luigi Bartalena: Nothing to disclose

George J. Kahaly: The Johannes Gutenberg University (JGU) Medical Center, Mainz, Germany has received research-associated funding from the JGU Medical Faculty, AdvanceCor (Germany), Apitope (Belgium), Berlin-Chemie (Germany), Byondis (The Netherlands), Horizon (USA), Immunovant (USA), ISAR (Germany), Mediomics (USA), Merck (Germany), Novartis (USA), Quidel (USA), River Vision (USA), and Roche (Switzerland). GJK consults for GlycoEra (Switzerland), Immunovant, Mediomics, Merck, Novartis, Quidel, & VasaraGen (USA).

Lelio Baldeschi: Nothing to disclose

Colin Dayan: Payment for lectures from Berlin-Chemie and Sanofi-Aventis

Anja Eckstein: Took part in multicenter studies from Horizon, Immunovant and Novartis. Lecture honoraria from Sanofi Aventis and Novo Nordisk

Claudio Marcocci: Consultant for Horizon Pharma, His institution received research funding for performing Teprotumumab study 2 (NCT03298867)

Michele Marinò: His institution received research funding from Horizon Pharma for performing Teprotumumab studies OPTIC and OPTIC-X (EudraCT 2017-002763-18 / EudraCT 2017-002713-58)

Bijay Vaidya: Lecture Honoraria from Berlin-Chemie

Wilmar Wiersinga: Consultant for Riverside, Horizon and Immunovant.

LEGENDS

Figure 1: Algorithm for the management of mild Graves' orbitopathy

TSHR-Ab: Thyrotropin receptor antibody, GO: Graves' orbitopathy, GD: Graves' disease

Figure 2: Algorithm for the first-line management of moderate-to-severe and active Graves' orbitopathy

The combination of a moderate cumulative dose of intravenous methylprednisolone + a moderate daily dose of oral enteric-coated mycophenolate sodium (first pathway) is the EUGOGO recommended first-line treatment for patients with moderate-to-severe and active GO (with or without diplopia). If mycophenolate sodium is not available, the other formulation mycophenolate mofetil is administered. Of note, 0.72 gram of mycophenolate sodium are equivalent to 1 gram of mycophenolate mofetil.

An alternative first-line treatment is the administration of high single doses of intravenous methylprednisolone starting with 0.75 gram per day and week for six consecutive weeks. This regimen is recommended for patients with constant/inconstant diplopia, severe proptosis and severe inflammatory soft-tissue changes.

GO: Graves' orbitopathy, GD: Graves' disease

Figure 3: Algorithm for the second-line management of moderate-to-severe and active Graves' orbitopathy

There are currently six alternative second-line treatments ("six pathways") for persistent moderate-to-severe and active GO as a non-response to a first-line treatment: a) a second cycle of iv methylprednisolone (starting with 0.75 g per infusion per week, allowed is a cumulative dose of 8 g per cycle) and subsequent to careful ophthalmic and biochemical (liver enzymes) evaluation; b) oral

glucocorticoids with either cyclosporine or azathioprine; c) orbital radiotherapy with either oral or intravenous glucocorticoids; d) teprotumumab (availability and affordability pending); e) rituximab (not in patients at risk for optic neuropathy); f) tocilizumab (considered in glucocorticoid-resistant patients). Head to head comparison data are available for rituximab against iv methylprednisolone (145). Overall and based on the features of GO that they are most likely to be effective for, intravenous glucocorticoids, mycophenolate, tocilizumab, rituximab, and cyclosporine substantially decreased inflammatory ophthalmic signs while orbital radiotherapy (preferably in combination with glucocorticoids) significantly improved eye muscle motility and/or diplopia. In comparison, Teprotumumab showed the strongest effect on exophthalmos. Of note, with the exception of Teprotumumab (FDA cleared for the treatment of active and moderately-severe GO in January 2020), all drugs stated in Figures 2 and/or 3 can be given as off-label treatment

GO: Graves' orbitopathy, GD: Graves' disease

Figure 4 Algorithm for the management of sight threatening Graves' orbitopathy

The first line treatment for optic neuropathy is high-dose iv methylprednisolone (single doses of 500 to 1000 mg) for three consecutive days or most preferably and for safety reasons on every second day (alternate days) during the first week, which can be repeated for another week. When the response is absent or poor with a deterioration in visual acuity or visual fields, urgent orbital decompression surgery is mandatory.

GO: Graves' orbitopathy, GD: Graves' disease

Table 1: Evidence-based recommendations for the management of Graves' orbitopathy (GO)

Table 2: Classification (clinical activity and severity) of Graves' orbitopathy (GO)

REFERENCES

1. Kahaly GJ. Management of Graves Thyroidal and Extrathyroidal Disease: An Update. *J Clin Endocrinol Metab.* 2020;105(12).
2. Perros P, Zarkovic M, Azzolini C, Ayvaz G, Baldeschi L, Bartalena L, Boschi A, Bournaud C, Brix TH, Covelli D, *et al.* PREGO (presentation of Graves' orbitopathy) study: changes in referral patterns to European Group On Graves' Orbitopathy (EUGOGO) centres over the period from 2000 to 2012. *Br J Ophthalmol.* 2015;99(11):1531-5.
3. Bartalena L, Piantanida E, Gallo D, Lai A, Tanda ML. Epidemiology, Natural History, Risk Factors, and Prevention of Graves' Orbitopathy. *Front Endocrinol (Lausanne).* 2020;11:615993.
4. Kahaly GJ, Petrak F, Hardt J, Pitz S, Egle UT. Psychosocial morbidity of Graves' orbitopathy. *Clin Endocrinol (Oxf).* 2005;63(4):395-402.
5. Wiersinga WM. Quality of life in Graves' ophthalmopathy. *Best Pract Res Clin Endocrinol Metab.* 2012;26(3):359-70.
6. Ponto KA, Hommel G, Pitz S, Elflein H, Pfeiffer N, Kahaly GJ. Quality of life in a german graves orbitopathy population. *Am J Ophthalmol.* 2011;152(3):483-90 e1.
7. Ponto KA, Merkesdal S, Hommel G, Pitz S, Pfeiffer N, Kahaly GJ. Public health relevance of Graves' orbitopathy. *J Clin Endocrinol Metab.* 2013;98(1):145-52.
8. Bartalena L, Baldeschi L, Boboridis K, Eckstein A, Kahaly GJ, Marcocci C, Perros P, Salvi M, Wiersinga WM, European Group on Graves O. The 2016 European Thyroid Association/European Group on Graves' Orbitopathy Guidelines for the Management of Graves' Orbitopathy. *Eur Thyroid J.* 2016;5(1):9-26.
9. Wiersinga WM. Advances in treatment of active, moderate-to-severe Graves' ophthalmopathy. *Lancet Diabetes Endocrinol.* 2017;5(2):134-42.
10. Smith TJ, Bartalena L. Will biological agents supplant systemic glucocorticoids as the first-line treatment for thyroid-associated ophthalmopathy? *Eur J Endocrinol.* 2019;181(5):D27-D43.
11. Davies TF, Andersen S, Latif R, Nagayama Y, Barbesino G, Brito M, Eckstein AK, Stagnaro-Green A, Kahaly GJ. Graves' disease. *Nat Rev Dis Primers.* 2020;6(1):52.
12. Taylor PN, Zhang L, Lee RWJ, Muller I, Ezra DG, Dayan CM, Kahaly GJ, Ludgate M. New insights into the pathogenesis and nonsurgical management of Graves orbitopathy. *Nat Rev Endocrinol.* 2020;16(2):104-16.
13. Swiglo BA, Murad MH, Schunemann HJ, Kunz R, Vigersky RA, Guyatt GH, Montori VM. A case for clarity, consistency, and helpfulness: state-of-the-art clinical practice guidelines in endocrinology using the grading of recommendations, assessment, development, and evaluation system. *J Clin Endocrinol Metab.* 2008;93(3):666-73.
14. European Group on Graves O, Wiersinga WM, Perros P, Kahaly GJ, Mourits MP, Baldeschi L, Boboridis K, Boschi A, Dickinson AJ, Kendall-Taylor P, *et al.* Clinical assessment of patients with Graves' orbitopathy: the European Group on Graves' Orbitopathy recommendations to generalists, specialists and clinical researchers. *Eur J Endocrinol.* 2006;155(3):387-9.
15. Terwee CB, Prummel MF, Gerding MN, Kahaly GJ, Dekker FW, Wiersinga WM. Measuring disease activity to predict therapeutic outcome in Graves' ophthalmopathy. *Clin Endocrinol (Oxf).* 2005;62(2):145-55.

16. Mourits MP, Koornneef L, Wiersinga WM, Prummel MF, Berghout A, van der Gaag R. Clinical criteria for the assessment of disease activity in Graves' ophthalmopathy: a novel approach. *Br J Ophthalmol*. 1989;73(8):639-44.
17. Dickinson AJ, Perros P. Controversies in the clinical evaluation of active thyroid-associated orbitopathy: use of a detailed protocol with comparative photographs for objective assessment. *Clin Endocrinol (Oxf)*. 2001;55(3):283-303.
18. Dolman PJ, Rootman J. VISA Classification for Graves orbitopathy. *Ophthalmic Plast Reconstr Surg*. 2006;22(5):319-24.
19. Kahaly GJ. Imaging in thyroid-associated orbitopathy. *Eur J Endocrinol*. 2001;145(2):107-18.
20. Muller-Forell W, Kahaly GJ. Neuroimaging of Graves' orbitopathy. *Best Pract Res Clin Endocrinol Metab*. 2012;26(3):259-71.
21. Jain AP, Gellada N, Ugradar S, Kumar A, Kahaly G, Douglas R. Teprotumumab reduces extraocular muscle and orbital fat volume in thyroid eye disease. *Br J Ophthalmol*. 2020.
22. Bartalena L, Baldeschi L, Dickinson A, Eckstein A, Kendall-Taylor P, Marcocci C, Mourits M, Perros P, Boboridis K, Boschi A, *et al*. Consensus statement of the European Group on Graves' orbitopathy (EUGOGO) on management of GO. *Eur J Endocrinol*. 2008;158(3):273-85.
23. Werner SC. Modification of the classification of the eye changes of Graves' disease. *Am J Ophthalmol*. 1977;83(5):725-7.
24. Eckstein AK, Plicht M, Lax H, Hirche H, Quadbeck B, Mann K, Steuhl KP, Esser J, Morgenthaler NG. Clinical results of anti-inflammatory therapy in Graves' ophthalmopathy and association with thyroidal autoantibodies. *Clin Endocrinol (Oxf)*. 2004;61(5):612-8.
25. Wiersinga WM, Prummel MF, Mourits MP, Koornneef L, Buller HR. Classification of the eye changes of Graves' disease. *Thyroid*. 1991;1(4):357-60.
26. Tanda ML, Piantanida E, Liparulo L, Veronesi G, Lai A, Sassi L, Pariani N, Gallo D, Azzolini C, Ferrario M, *et al*. Prevalence and natural history of Graves' orbitopathy in a large series of patients with newly diagnosed graves' hyperthyroidism seen at a single center. *J Clin Endocrinol Metab*. 2013;98(4):1443-9.
27. Ippolito S, Cusini C, Lasalvia P, Gianfagna F, Veronesi G, Gallo D, Masiello E, Premoli P, Sabatino J, Mercuriali A, *et al*. Change in newly diagnosed Graves' disease phenotype between the twentieth and the twenty-first centuries: meta-analysis and meta-regression. *J Endocrinol Invest*. 2020.
28. Wiersinga W, Zarkovic M, Bartalena L, Donati S, Perros P, Okosieme O, Morris D, Fichter N, Lareida J, von Arx G, *et al*. Predictive score for the development or progression of Graves' orbitopathy in patients with newly diagnosed Graves' hyperthyroidism. *Eur J Endocrinol*. 2018;178(6):635-43.
29. Kahaly GJ, Wuster C, Olivo PD, Diana T. High Titers of Thyrotropin Receptor Antibodies Are Associated With Orbitopathy in Patients With Graves Disease. *J Clin Endocrinol Metab*. 2019;104(7):2561-8.
30. Perros P, Dayan CM, Dickinson AJ, Ezra D, Estcourt S, Foley P, Hickey J, Lazarus JH, MacEwen CJ, McLaren J, *et al*. Management of patients with Graves' orbitopathy: initial assessment, management outside specialised centres and referral pathways. *Clin Med (Lond)*. 2015;15(2):173-8.

31. Baretic M, Juri MJ, Kusacic KS, Gudelj L, Radic M, Zlatar M. Awareness of EUGOGO guidelines and attitudes regarding smoking in Graves' orbitopathy in Croatia. *Taiwan J Ophthalmol*. 2021;DOI: 10.4103/tjo.tjo_2_21.
32. Estcourt S, Hickey J, Perros P, Dayan C, Vaidya B. The patient experience of services for thyroid eye disease in the United Kingdom: results of a nationwide survey. *Eur J Endocrinol*. 2009;161(3):483-7.
33. Terwee CB, Dekker FW, Mourits MP, Gerding MN, Baldeschi L, Kalmann R, Prummel MF, Wiersinga WM. Interpretation and validity of changes in scores on the Graves' ophthalmopathy quality of life questionnaire (GO-QOL) after different treatments. *Clin Endocrinol (Oxf)*. 2001;54(3):391-8.
34. Bartalena L, Wiersinga WM. Proposal for Standardization of Primary and Secondary Outcomes in Patients with Active, Moderate-to-Severe Graves' Orbitopathy. *Eur Thyroid J*. 2020;9(Suppl 1):3-16.
35. Prummel MF, Wiersinga W, Mourits M, Koornneef L, Berghout A, van der Gaag R. Amelioration of eye changes of Graves' ophthalmopathy by achieving euthyroidism. *Acta Endocrinol (Copenh)*. 1989;121:185-9.
36. Prummel MF, Wiersinga WM, Mourits MP, Koornneef L, Berghout A, van der Gaag R. Effect of abnormal thyroid function on the severity of Graves' ophthalmopathy. *Arch Intern Med*. 1990;150(5):1098-101.
37. Perros P, Kendall-Taylor P, Neoh C, Frewin S, Dickinson J. A prospective study of the effects of radioiodine therapy for hyperthyroidism in patients with minimally active graves' ophthalmopathy. *J Clin Endocrinol Metab*. 2005;90(9):5321-3.
38. Diana T, Ungerer M, Wuster C, Fassbender J, Li Z, Reimann A, Holthoff HP, Kanitz M, Kahaly GJ. A cyclic peptide significantly improves thyroid function, thyrotropin-receptor antibodies and orbital mucine /collagen content in a long-term Graves' disease mouse model. *J Autoimmun*. 2021;122:102666.
39. Hai YP, Lee ACH, Frommer L, Diana T, Kahaly GJ. Immunohistochemical analysis of human orbital tissue in Graves' orbitopathy. *J Endocrinol Invest*. 2020;43(2):123-37.
40. Diana T, Brown RS, Bossowski A, Segni M, Niedziela M, Konig J, Bossowska A, Ziora K, Hale A, Smith J, *et al*. Clinical relevance of thyroid-stimulating autoantibodies in pediatric graves' disease-a multicenter study. *J Clin Endocrinol Metab*. 2014;99(5):1648-55.
41. Kahaly GJ, Diana T, Glang J, Kanitz M, Pitz S, Konig J. Thyroid Stimulating Antibodies Are Highly Prevalent in Hashimoto's Thyroiditis and Associated Orbitopathy. *J Clin Endocrinol Metab*. 2016;101(5):1998-2004.
42. Kahaly GJ, Diana T, Kanitz M, Frommer L, Olivo PD. Prospective Trial of Functional Thyrotropin Receptor Antibodies in Graves Disease. *J Clin Endocrinol Metab*. 2020;105(4).
43. George A, Diana T, Langericht J, Kahaly GJ. Stimulatory Thyrotropin Receptor Antibodies Are a Biomarker for Graves' Orbitopathy. *Front Endocrinol (Lausanne)*. 2020;11:629925.
44. Leschik JJ, Diana T, Olivo PD, Konig J, Krahn U, Li Y, Kanitz M, Kahaly GJ. Analytical performance and clinical utility of a bioassay for thyroid-stimulating immunoglobulins. *Am J Clin Pathol*. 2013;139(2):192-200.
45. Wiersinga WM. Smoking and thyroid. *Clin Endocrinol (Oxf)*. 2013;79(2):145-51.

46. Khong JJ, Finch S, De Silva C, Rylander S, Craig JE, Selva D, Ebeling PR. Risk Factors for Graves' Orbitopathy; the Australian Thyroid-Associated Orbitopathy Research (ATOR) Study. *J Clin Endocrinol Metab.* 2016;101(7):2711-20.
47. Bartalena L, Marcocci C, Tanda ML, Manetti L, Dell'Unto E, Bartolomei MP, Nardi M, Martino E, Pinchera A. Cigarette smoking and treatment outcomes in Graves ophthalmopathy. *Ann Intern Med.* 1998;129(8):632-5.
48. Traisk F, Tallstedt L, Abraham-Nordling M, Andersson T, Berg G, Calissendorff J, Hallengren B, Hedner P, Lantz M, Nystrom E, *et al.* Thyroid-associated ophthalmopathy after treatment for Graves' hyperthyroidism with antithyroid drugs or iodine-131. *J Clin Endocrinol Metab.* 2009;94(10):3700-7.
49. Eckstein A, Quadbeck B, Mueller G, Rettenmeier AW, Hoermann R, Mann K, Steuhl P, Esser J. Impact of smoking on the response to treatment of thyroid associated ophthalmopathy. *Br J Ophthalmol.* 2003;87(6):773-6.
50. Matheis N, Lantz M, Grus FH, Ponto KA, Wolters D, Brorson H, Planck T, Shahida B, Pitz S, Pfeiffer N, *et al.* Proteomics of Orbital Tissue in Thyroid-Associated Orbitopathy. *J Clin Endocrinol Metab.* 2015;100(12):E1523-30.
51. Pfeilschifter J, Ziegler R. Smoking and endocrine ophthalmopathy: impact of smoking severity and current vs lifetime cigarette consumption. *Clin Endocrinol (Oxf).* 1996;45(4):477-81.
52. Bartalena L, Marcocci C, Bogazzi F, Panicucci M, Lepri A, Pinchera A. Use of corticosteroids to prevent progression of Graves' ophthalmopathy after radioiodine therapy for hyperthyroidism. *N Engl J Med.* 1989;321(20):1349-52.
53. Tallstedt L, Lundell G, Torring O, Wallin G, Ljunggren JG, Blomgren H, Taube A. Occurrence of ophthalmopathy after treatment for Graves' hyperthyroidism. The Thyroid Study Group. *N Engl J Med.* 1992;326(26):1733-8.
54. Bartalena L, Marcocci C, Bogazzi F, Manetti L, Tanda ML, Dell'Unto E, Bruno-Bossio G, Nardi M, Bartolomei MP, Lepri A, *et al.* Relation between therapy for hyperthyroidism and the course of Graves' ophthalmopathy. *N Engl J Med.* 1998;338(2):73-8.
55. Ponto KA, Zang S, Kahaly GJ. The tale of radioiodine and Graves' orbitopathy. *Thyroid.* 2010;20(7):785-93.
56. Vannucchi G, Covelli D, Campi I, Curro N, Dazzi D, Rodari M, Pepe G, Chiti A, Guastella C, Lazzaroni E, *et al.* Prevention of Orbitopathy by Oral or Intravenous Steroid Prophylaxis in Short Duration Graves' Disease Patients Undergoing Radioiodine Ablation: A Prospective Randomized Control Trial Study. *Thyroid.* 2019;29(12):1828-33.
57. Shiber S, Stiebel-Kalish H, Shimon I, Grossman A, Robenshtok E. Glucocorticoid regimens for prevention of Graves' ophthalmopathy progression following radioiodine treatment: systematic review and meta-analysis. *Thyroid.* 2014;24(10):1515-23.
58. Lai A, Sassi L, Compri E, Marino F, Sivelli P, Piantanida E, Tanda ML, Bartalena L. Lower dose prednisone prevents radioiodine-associated exacerbation of initially mild or absent graves' orbitopathy: a retrospective cohort study. *J Clin Endocrinol Metab.* 2010;95(3):1333-7.
59. Rosetti S, Tanda ML, Veronesi G, Masiello E, Premoli P, Gallo D, Cusini C, Donati S, Sabatino J, Ippolito S, *et al.* Oral steroid prophylaxis for Graves' orbitopathy after

radioactive iodine treatment for Graves' disease is not only effective, but also safe. *J Endocrinol Invest.* 2020;43(3):381-3.

60. Lanzolla G, Vannucchi G, Ionni I, Campi I, Sileo F, Lazzaroni E, Marino M. Cholesterol Serum Levels and Use of Statins in Graves' Orbitopathy: A New Starting Point for the Therapy. *Front Endocrinol (Lausanne).* 2019;10:933.
61. Stein JD, Childers D, Gupta S, Talwar N, Nan B, Lee BJ, Smith TJ, Douglas R. Risk factors for developing thyroid-associated ophthalmopathy among individuals with Graves disease. *JAMA Ophthalmol.* 2015;133(3):290-6.
62. Nilsson A, Tsoumani K, Planck T. Statins Decrease the Risk of Orbitopathy in Newly Diagnosed Patients with Graves' disease. *J Clin Endocrinol Metab.* 2021.
63. Sabini E, Mazzi B, Profilo MA, Mautone T, Casini G, Rocchi R, Ionni I, Menconi F, Leo M, Nardi M, *et al.* High Serum Cholesterol Is a Novel Risk Factor for Graves' Orbitopathy: Results of a Cross-Sectional Study. *Thyroid.* 2018;28(3):386-94.
64. Lanzolla G, Sabini E, Profilo MA, Mazzi B, Sframeli A, Rocchi R, Menconi F, Leo M, Nardi M, Vitti P, *et al.* Relationship between serum cholesterol and Graves' orbitopathy (GO): a confirmatory study. *J Endocrinol Invest.* 2018;41(12):1417-23.
65. Naselli A, Moretti D, Regalbuto C, Arpi ML, Lo Giudice F, Frasca F, Belfiore A, Le Moli R. Evidence That Baseline Levels of Low-Density Lipoproteins Cholesterol Affect the Clinical Response of Graves' Ophthalmopathy to Parenteral Corticosteroids. *Front Endocrinol (Lausanne).* 2020;11:609895.
66. Gurdal C, Sarac O, Genc I, Kirimlioglu H, Takmaz T, Can I. Ocular surface and dry eye in Graves' disease. *Curr Eye Res.* 2011;36(1):8-13.
67. Gupta A, Sadeghi PB, Akpek EK. Occult thyroid eye disease in patients presenting with dry eye symptoms. *Am J Ophthalmol.* 2009;147(5):919-23.
68. Matheis N, Grus FH, Breitenfeld M, Knych I, Funke S, Pitz S, Ponto KA, Pfeiffer N, Kahaly GJ. Proteomics Differentiate Between Thyroid-Associated Orbitopathy and Dry Eye Syndrome. *Invest Ophthalmol Vis Sci.* 2015;56(4):2649-56.
69. Selter JH, Gire AI, Sikder S. The relationship between Graves' ophthalmopathy and dry eye syndrome. *Clin Ophthalmol.* 2015;9:57-62.
70. Ismailova DS, Fedorov AA, Grusha YO. Ocular surface changes in thyroid eye disease. *Orbit.* 2013;32(2):87-90.
71. Eckstein AK, Finkenrath A, Heiligenhaus A, Renzing-Kohler K, Esser J, Kruger C, Quadbeck B, Steuhl KP, Gieseler RK. Dry eye syndrome in thyroid-associated ophthalmopathy: lacrimal expression of TSH receptor suggests involvement of TSHR-specific autoantibodies. *Acta Ophthalmol Scand.* 2004;82(3 Pt 1):291-7.
72. Vogel R, Crockett RS, Oden N, Laliberte TW, Molina L, Sodium Hyaluronate Ophthalmic Solution Study G. Demonstration of efficacy in the treatment of dry eye disease with 0.18% sodium hyaluronate ophthalmic solution (vismed, rejena). *Am J Ophthalmol.* 2010;149(4):594-601.
73. Uddin JM, Davies PD. Treatment of upper eyelid retraction associated with thyroid eye disease with subconjunctival botulinum toxin injection. *Ophthalmology.* 2002;109(6):1183-7.
74. Marcocci C, Kahaly GJ, Krassas GE, Bartalena L, Prummel M, Stahl M, Altea MA, Nardi M, Pitz S, Boboridis K, *et al.* Selenium and the course of mild Graves' orbitopathy. *N Engl J Med.* 2011;364(20):1920-31.

75. Negro R, Hegedus L, Attanasio R, Papini E, Winther KH. A 2018 European Thyroid Association Survey on the Use of Selenium Supplementation in Graves' Hyperthyroidism and Graves' Orbitopathy. *Eur Thyroid J*. 2019;8(1):7-15.
76. Kahaly GJ. Management of moderate-to-severe-GO. In: Wiersinga WM, Kahaly GJ, editors. *Graves' Orbitopathy*; Karger; 2017. p. 140-71.
77. Zang S, Kahaly GJ. Steroids and the immune response in Graves' orbitopathy *Immune Endoc Metab Agents Med Chem*. 2011 11(2):90 - 8.
78. Langericht J, Kramer I, Kahaly GJ. Glucocorticoids in Graves' orbitopathy: mechanisms of action and clinical application. *Ther Adv Endocrinol Metab*. 2020;11:2042018820958335.
79. Zang S, Ponto KA, Kahaly GJ. Clinical review: Intravenous glucocorticoids for Graves' orbitopathy: efficacy and morbidity. *J Clin Endocrinol Metab*. 2011;96(2):320-32.
80. van Geest RJ, Sasim IV, Koppeschaar HP, Kalmann R, Stravers SN, Bijlsma WR, Mourits MP. Methylprednisolone pulse therapy for patients with moderately severe Graves' orbitopathy: a prospective, randomized, placebo-controlled study. *Eur J Endocrinol*. 2008;158(2):229-37.
81. Marcocci C, Bartalena L, Tanda ML, Manetti L, Dell'Unto E, Rocchi R, Barbesino G, Mazzi B, Bartolomei MP, Lepri P, *et al*. Comparison of the effectiveness and tolerability of intravenous or oral glucocorticoids associated with orbital radiotherapy in the management of severe Graves' ophthalmopathy: results of a prospective, single-blind, randomized study. *J Clin Endocrinol Metab*. 2001;86(8):3562-7.
82. Kahaly GJ, Pitz S, Hommel G, Dittmar M. Randomized, single blind trial of intravenous versus oral steroid monotherapy in Graves' orbitopathy. *J Clin Endocrinol Metab*. 2005;90(9):5234-40.
83. Riedl M, Kolbe E, Kampmann E, Kramer I, Kahaly GJ. Prospectively recorded and MedDRA-coded safety data of intravenous methylprednisolone therapy in Graves' orbitopathy. *J Endocrinol Invest*. 2015;38(2):177-82.
84. Hoppe E, Lee ACH, Hoppe D, Kahaly GJ. Predictive Factors for Changes in Quality of Life after Steroid Treatment for Active Moderate-to-Severe Graves' Orbitopathy: A Prospective Trial. *Eur Thyroid J*. 2021;9(6):313-20.
85. Bartalena L, Krassas GE, Wiersinga W, Marcocci C, Salvi M, Daumerie C, Bournaud C, Stahl M, Sassi L, Veronesi G, *et al*. Efficacy and safety of three different cumulative doses of intravenous methylprednisolone for moderate to severe and active Graves' orbitopathy. *J Clin Endocrinol Metab*. 2012;97(12):4454-63.
86. Zang S, Ponto KA, Pitz S, Kahaly GJ. Dose of intravenous steroids and therapy outcome in Graves' orbitopathy. *J Endocrinol Invest*. 2011;34(11):876-80.
87. Marcocci C, Watt T, Altea MA, Rasmussen AK, Feldt-Rasmussen U, Orgiazzi J, Bartalena L, European Group of Graves O. Fatal and non-fatal adverse events of glucocorticoid therapy for Graves' orbitopathy: a questionnaire survey among members of the European Thyroid Association. *Eur J Endocrinol*. 2012;166(2):247-53.
88. Curro N, Covelli D, Vannucchi G, Campi I, Pirola G, Simonetta S, Dazzi D, Guastella C, Pignataro L, Beck-Peccoz P, *et al*. Therapeutic outcomes of high-dose intravenous steroids in the treatment of dysthyroid optic neuropathy. *Thyroid*. 2014;24(5):897-905.
89. Sisti E, Coco B, Menconi F, Leo M, Rocchi R, Latrofa F, Profilo MA, Mazzi B, Vitti P, Marcocci C, *et al*. Age and Dose Are Major Risk Factors for Liver Damage Associated

with Intravenous Glucocorticoid Pulse Therapy for Graves' Orbitopathy. *Thyroid*. 2015;25(7):846-50.

90. Miskiewicz P, Jankowska A, Brodzinska K, Milczarek-Banach J, Ambroziak U. Influence of Methylprednisolone Pulse Therapy on Liver Function in Patients with Graves' Orbitopathy. *Int J Endocrinol*. 2018;2018:1978590.
91. Bartalena L, Veronesi G, Krassas GE, Wiersinga WM, Marcocci C, Marino M, Salvi M, Daumerie C, Bournaud C, Stahl M, *et al*. Does early response to intravenous glucocorticoids predict the final outcome in patients with moderate-to-severe and active Graves' orbitopathy? *J Endocrinol Invest*. 2017;40(5):547-53.
92. Wang Y, Sharma A, Padnick-Silver L, Francis-Sedlak M, Holt RJ, Foley C, Massry G, Douglas RS. Trends in Treatment of Active, Moderate-to-Severe Thyroid Eye Disease in the United States. *J Endocr Soc*. 2020;4(12):bvaa140.
93. Nedeljkovic Beleslin B, Ciric J, Stojkovic M, Savic S, Lalic T, Stojanovic M, Miletic M, Knezevic M, Stankovic B, Zarkovic M. Comparison of efficacy and safety of parenteral versus parenteral and oral glucocorticoid therapy in Graves' orbitopathy. *Int J Clin Pract*. 2020;74(11):e13608.
94. Marcocci C, Bartalena L, Panicucci M, Marconcini C, Cartei F, Cavallacci G, Laddaga M, Campobasso G, Baschieri L, Pinchera A. Orbital cobalt irradiation combined with retrobulbar or systemic corticosteroids for Graves' ophthalmopathy: a comparative study. *Clin Endocrinol (Oxf)*. 1987;27(1):33-42.
95. Ebner R, Devoto MH, Weil D, Bordaberry M, Mir C, Martinez H, Bonelli L, Niepomniszcz H. Treatment of thyroid associated ophthalmopathy with periocular injections of triamcinolone. *Br J Ophthalmol*. 2004;88(11):1380-6.
96. Lee SJ, Rim TH, Jang SY, Kim CY, Shin DY, Lee EJ, Lee SY, Yoon JS. Treatment of upper eyelid retraction related to thyroid-associated ophthalmopathy using subconjunctival triamcinolone injections. *Graefes Arch Clin Exp Ophthalmol*. 2013;251(1):261-70.
97. Gupta OP, Boynton JR, Sabini P, Markowitch W, Jr., Quatela VC. Proptosis after retrobulbar corticosteroid injections. *Ophthalmology*. 2003;110(2):443-7.
98. Huebert I, Heinicke N, Kook D, Boost KA, Miller CV, Mayer WJ, Haritoglou C, Kampik A, Gandorfer A, Hintschich C, *et al*. Dual platelet inhibition in cases of severe retrobulbar hemorrhage following retrobulbar and peribulbar anesthesia. *J Cataract Refract Surg*. 2015;41(10):2092-101.
99. Allison AC, Eugui EM. Mycophenolate mofetil and its mechanisms of action. *Immunopharmacology*. 2000;47(2-3):85-118.
100. Staatz CE, Tett SE. Pharmacology and toxicology of mycophenolate in organ transplant recipients: an update. *Arch Toxicol*. 2014;88(7):1351-89.
101. Azzola A, Havryk A, Chhajed P, Hostettler K, Black J, Johnson P, Roth M, Glanville A, Tamm M. Everolimus and mycophenolate mofetil are potent inhibitors of fibroblast proliferation after lung transplantation. *Transplantation*. 2004;77(2):275-80.
102. Roos N, Poulalhon N, Farge D, Madelaine I, Mauviel A, Verrecchia F. In vitro evidence for a direct antifibrotic role of the immunosuppressive drug mycophenolate mofetil. *J Pharmacol Exp Ther*. 2007;321(2):583-9.
103. Petrova DT, Brandhorst G, Brehmer F, Gross O, Oellerich M, Armstrong VW. Mycophenolic acid displays IMPDH-dependent and IMPDH-independent effects on renal fibroblast proliferation and function. *Ther Drug Monit*. 2010;32(4):405-12.

104. Zhang L, Grennan-Jones F, Draman MS, Lane C, Morris D, Dayan CM, Tee AR, Ludgate M. Possible targets for nonimmunosuppressive therapy of Graves' orbitopathy. *J Clin Endocrinol Metab.* 2014;99(7):E1183-90.
105. Gabardi S, Tran JL, Clarkson MR. Enteric-coated mycophenolate sodium. *Ann Pharmacother.* 2003;37(11):1685-93.
106. Gardiner KM, Tett SE, Staatz CE. Is Conversion from Mycophenolate Mofetil to Enteric-Coated Mycophenolate Sodium Justifiable for Gastrointestinal Quality of Life? *Drugs R D.* 2018;18(4):271-82.
107. Ye X, Bo X, Hu X, Cui H, Lu B, Shao J, Wang J. Efficacy and safety of mycophenolate mofetil in patients with active moderate-to-severe Graves' orbitopathy. *Clin Endocrinol (Oxf).* 2017;86(2):247-55.
108. Kahaly GJ, Riedl M, Konig J, Pitz S, Ponto K, Diana T, Kampmann E, Kolbe E, Eckstein A, Moeller LC, *et al.* Mycophenolate plus methylprednisolone versus methylprednisolone alone in active, moderate-to-severe Graves' orbitopathy (MINGO): a randomised, observer-masked, multicentre trial. *Lancet Diabetes Endocrinol.* 2018;6(4):287-98.
109. Orgiazzi J. Adding the Immunosuppressant Mycophenolate Mofetil to Medium-Dose Infusions of Methylprednisolone Improves the Treatment of Graves' Orbitopathy. *Clin Thyroidol.* 2018;30(1):10-4.
110. Lee ACH, Riedl M, Frommer L, Diana T, Kahaly GJ. Systemic safety analysis of mycophenolate in Graves' orbitopathy. *J Endocrinol Invest.* 2020;43(6):767-77.
111. Riedl M, Kuhn A, Kramer I, Kolbe E, Kahaly GJ. Prospective, systematically recorded mycophenolate safety data in Graves' orbitopathy. *J Endocrinol Invest.* 2016;39(6):687-94.
112. Quah Qin Xian N, Alnahrawy A, Akshikar R, Lee V. Real-World Efficacy and Safety of Mycophenolate Mofetil in Active Moderate-to-Sight-Threatening Thyroid Eye Disease. *Clin Ophthalmol.* 2021;15:1921-32.
113. Mourits MP, van Kempen-Harteveld ML, Garcia MB, Koppeschaar HP, Tick L, Terwee CB. Radiotherapy for Graves' orbitopathy: randomised placebo-controlled study. *Lancet.* 2000;355(9214):1505-9.
114. Prummel MF, Terwee CB, Gerding MN, Baldeschi L, Mourits MP, Blank L, Dekker FW, Wiersinga WM. A randomized controlled trial of orbital radiotherapy versus sham irradiation in patients with mild Graves' ophthalmopathy. *J Clin Endocrinol Metab.* 2004;89(1):15-20.
115. Gorman CA, Garrity JA, Fatourech V, Bahn RS, Petersen IA, Stafford SL, Earle JD, Forbes GS, Kline RW, Bergstralh EJ, *et al.* A prospective, randomized, double-blind, placebo-controlled study of orbital radiotherapy for Graves' ophthalmopathy. *Ophthalmology.* 2001;108(9):1523-34.
116. Rajendram R, Taylor PN, Wilson VJ, Harris N, Morris OC, Tomlinson M, Yarrow S, Garrett H, Herbert HM, Dick AD, *et al.* Combined immunosuppression and radiotherapy in thyroid eye disease (CIRTED): a multicentre, 2 x 2 factorial, double-blind, randomised controlled trial. *Lancet Diabetes Endocrinol.* 2018;6(4):299-309.
117. Prummel MF, Mourits MP, Blank L, Berghout A, Koornneef L, Wiersinga WM. Randomized double-blind trial of prednisone versus radiotherapy in Graves' ophthalmopathy. *Lancet.* 1993;342(8877):949-54.
118. Marcocci C, Bartalena L, Bogazzi F, Bruno-Bossio G, Lepri A, Pinchera A. Orbital radiotherapy combined with high dose systemic glucocorticoids for Graves'

- ophthalmopathy is more effective than radiotherapy alone: results of a prospective randomized study. *J Endocrinol Invest*. 1991;14(10):853-60.
119. Bartalena L, Marcocci C, Chiovato L, Laddaga M, Lepri G, Andreani D, Cavallacci G, Baschieri L, Pinchera A. Orbital cobalt irradiation combined with systemic corticosteroids for Graves' ophthalmopathy: comparison with systemic corticosteroids alone. *J Clin Endocrinol Metab*. 1983;56(6):1139-44.
120. Kim JW, Han SH, Son BJ, Rim TH, Keum KC, Yoon JS. Efficacy of combined orbital radiation and systemic steroids in the management of Graves' orbitopathy. *Graefes Arch Clin Exp Ophthalmol*. 2016;254(5):991-8.
121. Oeverhaus M, Witteler T, Lax H, Esser J, Fuhrer D, Eckstein A. Combination Therapy of Intravenous Steroids and Orbital Irradiation is More Effective Than Intravenous Steroids Alone in Patients with Graves' Orbitopathy. *Horm Metab Res*. 2017;49(10):739-47.
122. Tanda ML, Bartalena L. Efficacy and safety of orbital radiotherapy for graves' orbitopathy. *J Clin Endocrinol Metab*. 2012;97(11):3857-65.
123. Kahaly GJ, Rosler HP, Pitz S, Hommel G. Low- versus high-dose radiotherapy for Graves' ophthalmopathy: a randomized, single blind trial. *J Clin Endocrinol Metab*. 2000;85(1):102-8.
124. Marquez SD, Lum BL, McDougall IR, Katkuri S, Levin PS, MacManus M, Donaldson SS. Long-term results of irradiation for patients with progressive Graves' ophthalmopathy. *Int J Radiat Oncol Biol Phys*. 2001;51(3):766-74.
125. Marcocci C, Bartalena L, Rocchi R, Marino M, Menconi F, Morabito E, Mazzi B, Mazzeo S, Sartini MS, Nardi M, *et al*. Long-term safety of orbital radiotherapy for Graves' ophthalmopathy. *J Clin Endocrinol Metab*. 2003;88(8):3561-6.
126. Wakelkamp IM, Tan H, Saeed P, Schlingemann RO, Verbraak FD, Blank LE, Prummel MF, Wiersinga WM. Orbital irradiation for Graves' ophthalmopathy: Is it safe? A long-term follow-up study. *Ophthalmology*. 2004;111(8):1557-62.
127. Godfrey KJ, Kazim M. Radiotherapy for Active Thyroid Eye Disease. *Ophthalmic Plast Reconstr Surg*. 2018;34(4S Suppl 1):S98-S104.
128. Kahaly GJ, Rosler HP, Pitz S, Krummenauer F, Hommel G. Radiotherapy for Graves' ophthalmopathy. *J Clin Endocrinol Metab* 2001;86:2327-8 (Letter).
129. Kahaly G, Schrezenmeir J, Krause U, Schweikert B, Meuer S, Muller W, Dennebaum R, Beyer J. Cyclosporin and prednisone v. prednisone in treatment of Graves' ophthalmopathy: a controlled, randomized and prospective study. *Eur J Clin Invest*. 1986;16(5):415-22.
130. Prummel MF, Mourits MP, Berghout A, Krenning EP, van der Gaag R, Koornneef L, Wiersinga WM. Prednisone and cyclosporine in the treatment of severe Graves' ophthalmopathy. *N Engl J Med*. 1989;321(20):1353-9.
131. Perros P, Weightman DR, Crombie AL, Kendall-Taylor P. Azathioprine in the treatment of thyroid-associated ophthalmopathy. *Acta Endocrinol (Copenh)*. 1990;122(1):8-12.
132. Chalvatzis NT, Tzamalidis AK, Kalantzis GK, El-Hindy N, Dimitrakos SA, Potts MJ. Safety and efficacy of combined immunosuppression and orbital radiotherapy in thyroid-related restrictive myopathy: two-center experience. *Eur J Ophthalmol*. 2014;24(6):953-9.
133. Tsui S, Naik V, Hoa N, Hwang CJ, Afifiyan NF, Sinha Hikim A, Gianoukakis AG, Douglas RS, Smith TJ. Evidence for an association between thyroid-stimulating

- hormone and insulin-like growth factor 1 receptors: a tale of two antigens implicated in Graves' disease. *J Immunol.* 2008;181(6):4397-405.
134. Douglas RS, Naik V, Hwang CJ, Afifiyan NF, Gianoukakis AG, Sand D, Kamat S, Smith TJ. B cells from patients with Graves' disease aberrantly express the IGF-1 receptor: implications for disease pathogenesis. *J Immunol.* 2008;181(8):5768-74.
135. Smith TJ, Janssen J. Insulin-like Growth Factor-I Receptor and Thyroid-Associated Ophthalmopathy. *Endocr Rev.* 2019;40(1):236-67.
136. Smith TJ, Kahaly GJ, Ezra DG, Fleming JC, Dailey RA, Tang RA, Harris GJ, Antonelli A, Salvi M, Goldberg RA, *et al.* Teprotumumab for Thyroid-Associated Ophthalmopathy. *N Engl J Med.* 2017;376(18):1748-61.
137. Douglas RS, Kahaly GJ, Patel A, Sile S, Thompson EH, Perdok R, Fleming JC, Fowler BT, Marcocci C, Marino M, *et al.* Teprotumumab for the Treatment of Active Thyroid Eye Disease. *N Engl J Med.* 2020;382(4):341-52.
138. Kahaly GJ, Douglas RS, Holt RJ, Sile S, Smith TJ. Teprotumumab for patients with active thyroid eye disease: a pooled data analysis, subgroup analyses, and off-treatment follow-up results from two randomised, double-masked, placebo-controlled, multicentre trials. *Lancet Diabetes Endocrinol.* 2021;9(6):360-72.
139. Hoang TD, Nguyen NT, Chou E, Shakir MK. Rapidly progressive cognitive decline associated with teprotumumab in thyroid eye disease. *BMJ Case Rep.* 2021;14(5).
140. Winn BJ, Kersten RC. Teprotumumab: Interpreting the Clinical Trials in the Context of Thyroid Eye Disease Pathogenesis and Current Therapies. *Ophthalmology.* 2021.
141. Salvi M, Vannucchi G, Campi I, Curro N, Dazzi D, Simonetta S, Bonara P, Rossi S, Sina C, Guastella C, *et al.* Treatment of Graves' disease and associated ophthalmopathy with the anti-CD20 monoclonal antibody rituximab: an open study. *Eur J Endocrinol.* 2007;156(1):33-40.
142. Mitchell AL, Gan EH, Morris M, Johnson K, Neoh C, Dickinson AJ, Perros P, Pearce SH. The effect of B cell depletion therapy on anti-TSH receptor antibodies and clinical outcome in glucocorticoid-refractory Graves' orbitopathy. *Clin Endocrinol (Oxf).* 2013;79(3):437-42.
143. Khanna D, Chong KK, Afifiyan NF, Hwang CJ, Lee DK, Garneau HC, Goldberg RA, Darwin CH, Smith TJ, Douglas RS. Rituximab treatment of patients with severe, corticosteroid-resistant thyroid-associated ophthalmopathy. *Ophthalmology.* 2010;117(1):133-9 e2.
144. Stan MN, Garrity JA, Carranza Leon BG, Prabin T, Bradley EA, Bahn RS. Randomized controlled trial of rituximab in patients with Graves' orbitopathy. *J Clin Endocrinol Metab.* 2015;100(2):432-41.
145. Salvi M, Vannucchi G, Curro N, Campi I, Covelli D, Dazzi D, Simonetta S, Guastella C, Pignataro L, Avignone S, *et al.* Efficacy of B-cell targeted therapy with rituximab in patients with active moderate to severe Graves' orbitopathy: a randomized controlled study. *J Clin Endocrinol Metab.* 2015;100(2):422-31.
146. Vannucchi G, Campi I, Covelli D, Curro N, Lazzaroni E, Palomba A, Soranna D, Zambon A, Fugazzola L, Muller I, *et al.* Efficacy Profile and Safety Implications of Very Low Dose Rituximab in Patients with Graves' Orbitopathy. *Thyroid.* 2020.
147. Hamed Azzam S, Kang S, Salvi M, Ezra DG. Tocilizumab for thyroid eye disease. *Cochrane Database Syst Rev.* 2018;11:CD012984.

148. Perez-Moreiras JV, Gomez-Reino JJ, Maneiro JR, Perez-Pampin E, Romo Lopez A, Rodriguez Alvarez FM, Castillo Laguarda JM, Del Estad Cabello A, Gessa Sorroche M, Espana Gregori E, *et al.* Efficacy of Tocilizumab in Patients With Moderate-to-Severe Corticosteroid-Resistant Graves Orbitopathy: A Randomized Clinical Trial. *Am J Ophthalmol.* 2018;195:181-90.
149. Sanchez-Bilbao L, Martinez-Lopez D, Revenga M, Lopez-Vazquez A, Valls-Pascual E, Atienza-Mateo B, Valls-Espinosa B, Maiz-Alonso O, Blanco A, Torre-Salaberri I, *et al.* Anti-IL-6 Receptor Tocilizumab in Refractory Graves' Orbitopathy: National Multicenter Observational Study of 48 Patients. *J Clin Med.* 2020;9(9).
150. Ceballos-Macias Jose J, Rivera-Moscoso R, Flores-Real Jorge A, Vargas-Sanchez J, Ortega-Gutierrez G, Madriz-Prado R, Velasco-Ramos PC, Munoz-Monroy Omar E, Meneses-Perez Anna C, Fernandez-Morales Irma N, *et al.* Tocilizumab in glucocorticoid-resistant graves orbitopathy. A case series report of a mexican population. *Ann Endocrinol (Paris).* 2020;81(2-3):78-82.
151. Perez-Moreiras JV, Varela-Agra M, Prada-Sanchez MC, Prada-Ramallal G. Steroid-Resistant Graves' Orbitopathy Treated with Tocilizumab in Real-World Clinical Practice: A 9-Year Single-Center Experience. *J Clin Med.* 2021;10(4).
152. Kazatchkine MD, Kaveri SV. Immunomodulation of autoimmune and inflammatory diseases with intravenous immune globulin. *N Engl J Med.* 2001;345(10):747-55.
153. Kahaly G, Pitz S, Muller-Forell W, Hommel G. Randomized trial of intravenous immunoglobulins versus prednisolone in Graves' ophthalmopathy. *Clin Exp Immunol.* 1996;106(2):197-202.
154. Kahaly G, Lieb W, Muller-Forell W, Mainberger M, Beyer J, Vollmar J, Staiger C. Ciamezone in endocrine orbitopathy. A randomized double-blind, placebo-controlled study. *Acta Endocrinol (Copenh).* 1990;122(1):13-21.
155. Dickinson AJ, Vaidya B, Miller M, Coulthard A, Perros P, Baister E, Andrews CD, Hesse L, Heverhagen JT, Heufelder AE, *et al.* Double-blind, placebo-controlled trial of octreotide long-acting repeatable (LAR) in thyroid-associated ophthalmopathy. *J Clin Endocrinol Metab.* 2004;89(12):5910-5.
156. Wemeau JL, Caron P, Beckers A, Rohmer V, Orgiazzi J, Borson-Chazot F, Nocaudie M, Perimenis P, Bisot-Locard S, Bourdeix I, *et al.* Octreotide (long-acting release formulation) treatment in patients with graves' orbitopathy: clinical results of a four-month, randomized, placebo-controlled, double-blind study. *J Clin Endocrinol Metab.* 2005;90(2):841-8.
157. Stan MN, Garrity JA, Bradley EA, Woog JJ, Bahn MM, Brennan MD, Bryant SC, Achenbach SJ, Bahn RS. Randomized, double-blind, placebo-controlled trial of long-acting release octreotide for treatment of Graves' ophthalmopathy. *J Clin Endocrinol Metab.* 2006;91(12):4817-24.
158. Chang TC, Liao SL. Slow-release lanreotide in Graves' ophthalmopathy: A double-blind randomized, placebo-controlled clinical trial. *J Endocrinol Invest.* 2006;29(5):413-22.
159. Wakelkamp IM, Gerding MN, Van Der Meer JW, Prummel MF, Wiersinga WM. Both Th1- and Th2-derived cytokines in serum are elevated in Graves' ophthalmopathy. *Clin Exp Immunol.* 2000;121(3):453-7.

160. Kishazi E, Dor M, Eperon S, Oberic A, Turck N, Hamedani M. Differential profiling of lacrimal cytokines in patients suffering from thyroid-associated orbitopathy. *Sci Rep.* 2018;8(1):10792.
161. Durrani OM, Reuser TQ, Murray PI. Infliximab: a novel treatment for sight-threatening thyroid associated ophthalmopathy. *Orbit.* 2005;24(2):117-9.
162. Paridaens D, van den Bosch WA, van der Loos TL, Krenning EP, van Hagen PM. The effect of etanercept on Graves' ophthalmopathy: a pilot study. *Eye (Lond).* 2005;19(12):1286-9.
163. Ayabe R, Rootman DB, Hwang CJ, Ben-Artzi A, Goldberg R. Adalimumab as steroid-sparing treatment of inflammatory-stage thyroid eye disease. *Ophthalmic Plast Reconstr Surg.* 2014;30(5):415-9.
164. Sipkova Z, Insull EA, David J, Turner HE, Keren S, Norris JH. Early use of steroid-sparing agents in the inactivation of moderate-to-severe active thyroid eye disease: a step-down approach. *Clin Endocrinol (Oxf).* 2018;89(6):834-9.
165. Wakelkamp IM, Baldeschi L, Saeed P, Mourits MP, Prummel MF, Wiersinga WM. Surgical or medical decompression as a first-line treatment of optic neuropathy in Graves' ophthalmopathy? A randomized controlled trial. *Clin Endocrinol (Oxf).* 2005;63(3):323-8.
166. Baldeschi L. Rehabilitative surgery. In: Wiersinga WM, Kahaly GJ, editors. *Graves' Orbitopathy A Multidisciplinary Approach - Questions and Answers.* 3rd ed. Basel, Switzerland: Karger; 2017.
167. Kahaly GJ, Bartalena L, Hegedus L, Leenhardt L, Poppe K, Pearce SH. 2018 European Thyroid Association Guideline for the Management of Graves' Hyperthyroidism. *Eur Thyroid J.* 2018;7(4):167-86.
168. Maccocci C, Bruno-Bossio G, Manetti L, Tanda ML, Miccoli P, Iacconi P, Bartolomei MP, Nardi M, Pinchera A, Bartalena L. The course of Graves' ophthalmopathy is not influenced by near total thyroidectomy: a case-control study. *Clin Endocrinol (Oxf).* 1999;51(4):503-8.
169. Laurberg P, Wallin G, Tallstedt L, Abraham-Nordling M, Lundell G, Topping O. TSH-receptor autoimmunity in Graves' disease after therapy with anti-thyroid drugs, surgery, or radioiodine: a 5-year prospective randomized study. *Eur J Endocrinol.* 2008;158(1):69-75.
170. Diana T, Ponto KA, Kahaly GJ. Thyrotropin receptor antibodies and Graves' orbitopathy. *J Endocrinol Invest.* 2021;44(4):703-12.
171. Zarkovic M, Wiersinga W, Perros P, Bartalena L, Donati S, Okosieme O, Morris D, Fichter N, Lareida J, Daumerie C, *et al.* Antithyroid drugs in Graves' hyperthyroidism: differences between "block and replace" and "titration" regimes in frequency of euthyroidism and Graves' orbitopathy during treatment. *J Endocrinol Invest.* 2021;44(2):371-8.
172. Erdogan MF, Demir O, Ersoy RU, Gul K, Aydogan BI, Uc ZA, Mete T, Ertek S, Unluturk U, Cakir B, *et al.* Comparison of Early Total Thyroidectomy with Antithyroid Treatment in Patients with Moderate-Severe Graves' Orbitopathy: A Randomized Prospective Trial. *Eur Thyroid J.* 2016;5(2):106-11.
173. Meyer Zu Horste M, Pateronis K, Walz MK, Alesina P, Mann K, Schott M, Esser J, Eckstein AK. The Effect of Early Thyroidectomy on the Course of Active Graves' Orbitopathy (GO): A Retrospective Case Study. *Horm Metab Res.* 2016;48(7):433-9.

174. Menconi F, Marino M, Pinchera A, Rocchi R, Mazzi B, Nardi M, Bartalena L, Marcocci C. Effects of total thyroid ablation versus near-total thyroidectomy alone on mild to moderate Graves' orbitopathy treated with intravenous glucocorticoids. *J Clin Endocrinol Metab.* 2007;92(5):1653-8.
175. Moleti M, Violi MA, Montanini D, Trombetta C, Di Bella B, Sturniolo G, Presti S, Alibrandi A, Campenni A, Baldari S, *et al.* Radioiodine ablation of postsurgical thyroid remnants after treatment with recombinant human TSH (rhTSH) in patients with moderate-to-severe graves' orbitopathy (GO): a prospective, randomized, single-blind clinical trial. *J Clin Endocrinol Metab.* 2014;99(5):1783-9.
176. Leo M, Marcocci C, Pinchera A, Nardi M, Megna L, Rocchi R, Latrofa F, Altea MA, Mazzi B, Sisti E, *et al.* Outcome of Graves' orbitopathy after total thyroid ablation and glucocorticoid treatment: follow-up of a randomized clinical trial. *J Clin Endocrinol Metab.* 2012;97(1):E44-8.
177. Bartalena L, Macchia PE, Marcocci C, Salvi M, Vermiglio F. Effects of treatment modalities for Graves' hyperthyroidism on Graves' orbitopathy: a 2015 Italian Society of Endocrinology Consensus Statement. *J Endocrinol Invest.* 2015;38(4):481-7.
178. Elbers L, Mourits M, Wiersinga W. Outcome of very long-term treatment with antithyroid drugs in Graves' hyperthyroidism associated with Graves' orbitopathy. *Thyroid.* 2011;21(3):279-83.
179. Laurberg P, Berman DC, Andersen S, Bulow Pedersen I. Sustained control of Graves' hyperthyroidism during long-term low-dose antithyroid drug therapy of patients with severe Graves' orbitopathy. *Thyroid.* 2011;21(9):951-6.
180. Bartalena L, Chiovato L, Marcocci C, Vitti P, Piantanida E, Tanda ML. Management of Graves' hyperthyroidism and orbitopathy in time of COVID-19 pandemic. *J Endocrinol Invest.* 2020;doi: 10.1007/s40618-020-01293-7.
181. Puig-Domingo M, Marazuela M, Giustina A. COVID-19 and endocrine diseases. A statement from the European Society of Endocrinology. *Endocrine.* 2020;68(1):2-5.
182. Group WHOREAfC-TW, Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, Angus DC, Annane D, Azevedo LCP, Berwanger O, *et al.* Association Between Administration of Systemic Corticosteroids and Mortality Among Critically Ill Patients With COVID-19: A Meta-analysis. *JAMA.* 2020;324(13):1330-41.
183. Alexaki VI, Henneicke H. The Role of Glucocorticoids in the Management of COVID-19. *Horm Metab Res.* 2021;53(1):9-15.
184. Clarke SA, Phylactou M, Patel B, Mills EG, Muzi B, Izzi-Engbeaya C, Choudhury S, Khoo B, Meeran K, Comninos AN, *et al.* Normal adrenal and thyroid function in patients who survive COVID-19 infection. *J Clin Endocrinol Metab.* 2021.
185. Jespersen S, Nygaard B, Kristensen LO. Methylprednisolone Pulse Treatment of Graves' Ophthalmopathy Is Not Associated with Secondary Adrenocortical Insufficiency. *Eur Thyroid J.* 2015;4(4):222-5.
186. Ambroziak U, Bluszcz G, Bednarczyk T, Miskiewicz P. The influence of Graves' orbitopathy treatment with intravenous glucocorticoids on adrenal function. *Endokrynol Pol.* 2017;68(4):430-3.
187. Neumann S, Krieger CC, Gershengorn MC. Targeting TSH and IGF-1 Receptors to Treat Thyroid Eye Disease. *Eur Thyroid J.* 2020;9(Suppl 1):59-65.
188. Krause G, Eckstein A, Schulein R. Modulating TSH Receptor Signaling for Therapeutic Benefit. *Eur Thyroid J.* 2020;9(Suppl 1):66-77.

189. Lee ACH, Kahaly GJ. Novel Approaches for Immunosuppression in Graves' Hyperthyroidism and Associated Orbitopathy. *Eur Thyroid J.* 2020;9(Suppl 1):17-30.
190. Roos JCP, Murthy R. Sirolimus (rapamycin) for the targeted treatment of the fibrotic sequelae of Graves' orbitopathy. *Eye (Lond).* 2019;33(4):679-82.
191. Masetti G, Ludgate M. Microbiome and Graves' Orbitopathy. *Eur Thyroid J.* 2020;9(Suppl 1):78-85.

General recommendations

- Refrain from smoking
- Treat thyroid dysfunction (preferably with antithyroid drugs, especially if risk factors for deterioration/progression of GO are present (see below))
- Avoid iatrogenic hypothyroidism in treating patients with GD/GO
- Referral to thyroid-eye clinics if risk factors present (active GO, smoker, high TSHR-Ab, unstable / severe hyperthyroidism)
- Search for dry eye syndrome

Management

Local treatment

- Artificial tears, especially when dry eye present
- Ophthalmic gels (cornea protection during the night)

Systemic adjunct therapy for active GO

- Selenium supplementation for six months (fasting intake)


Quality of life markedly impaired



Discuss low dose immunomodulatory (active GO) or rehabilitative surgery (inactive GO) following extensive counselling and shared decision

Box 2**MODERATE-TO-SEVERE AND ACTIVE GO
FIRST – LINE TREATMENT****General
Recommendations**

- Referral to thyroid-eye clinic for counselling and treatment plan shared with patient
- Stop smoking
- Treat thyroid dysfunction with antithyroid drugs
- Avoid iatrogenic hypothyroidism in treating patients with GD/GO

**INACTIVE GO**


Rehabilitative surgery as needed or required by the patient

MODERATE-TO-SEVERE AND ACTIVE GO

SECOND – LINE TREATMENTS

General Recommendations

- Referral to thyroid-eye clinic for counselling and treatment plan shared with patient
- Stop smoking
- Treat thyroid dysfunction with antithyroid drugs
- Avoid iatrogenic hypothyroidism in treating patients with GD/GO



Rehabilitative surgery (orbital decompression, squint / lid surgery) as needed or required by the patient

Box 4

SIGHT - THREATENING GO (Optic Neuropathy)**General recommendations**

- Immediate referral to thyroid-eye clinic
- Stop smoking
- Avoid radioactive iodine treatment
- Stabilize thyroid dysfunction with antithyroid drugs
- Avoid iatrogenic hypothyroidism in treating patients with GD/GO

Specific Management

Intravenous methylprednisolone (0.5 – 1 gram, as single dose repeated on three consecutive or alternate days)



Daily monitoring of ophthalmic parameters



After one week, evaluation if therapy can be continued



Yes



Further with intravenous methylprednisolone as in week one



Response



0.5 g intravenous methylprednisolone 1x / wk. (cumulative dose <8 g / cycle)



Partial response



Urgent orbital decompression surgery (Imaging recommended)



No response and/or deterioration of ophthalmic signs



Number		RECOMMENDATIONS	Strength of recommendation and level of evidence
1)	Assessment	Clinical activity and severity of Graves' orbitopathy (GO) should be assessed according to standardized criteria and GO be categorized as active or inactive, and mild, moderate-to-severe, or sight threatening, and should include evaluation of quality of life (QoL) by the EUGOGO disease-specific GO-QoL questionnaire	(1, $\emptyset\emptyset\emptyset\emptyset$)
2)	Specialized centres and risk factors	We recommend that primary-care physicians, general practitioners, general internists and specialists should refer patients with overt GO and mild cases at risk to deteriorate (clinically active GO, smokers, severe/unstable hyperthyroidism, high serum thyrotropin receptor antibody [TSHR-Ab] titres), to combined thyroid-eye clinics or specialized centres providing both endocrine and ophthalmic expertise, as this will provide accurate and timely diagnosis to improve prognosis and QoL	(1, $\emptyset\emptyset\emptyset\emptyset$)
3)	Quit smoking	Physicians should urge all patients with Graves' hyperthyroidism, irrespective of the presence/absence of GO, to refrain from smoking	(1, $\emptyset\emptyset\emptyset\emptyset$)
4)	Thyroid dysfunction	Euthyroidism should be promptly restored and stably maintained in all patients with GO	(1, $\emptyset\emptyset\emptyset\emptyset$)
5)	Glucocorticoid prophylaxis	Oral prednisone/prednisolone prophylaxis should be given to radioactive iodine (RAI)-treated patients at risk of progression or <i>de novo</i> development of GO (smokers, severe/unstable hyperthyroidism, high serum TSHR-Ab). Regimen: high risk: 0.3-0.5 mg/kg/bodyweight as starting dose, tapered and withdrawn after three months; low risk: 0.1-0.2 mg/kg/bodyweight, tapered and withdrawn after six weeks. Patients with longstanding and stably inactive GO can receive RAI without prednisone/prednisolone cover if risk factors for GO progression, particularly smoking and high serum TSHR-Ab titers are absent. Uncontrolled post-RAI hypothyroidism should be avoided.	(1, $\emptyset\emptyset\emptyset\emptyset$)
6)	Local treatments	All patients with GO should be extensively treated locally with artificial tears at all times in the course of their disease unless corneal exposure requires higher protection than ophthalmic gels or ointment, especially at nighttime	(1, $\emptyset\emptyset\emptyset\emptyset$)
7)	Mild GO	Mild GO should be treated with local treatments and general measures to control risk factors; a 6-month selenium supplementation should be given to patients with mild and active GO of recent onset, because it improves eye manifestations and QoL, and usually prevents GO progression to more severe forms	(1, $\emptyset\emptyset\emptyset\emptyset$)

8)	Mild GO	If the impact of the disease on the QoL outweighs risks, then low-dose immunomodulatory therapy (active GO) or rehabilitative surgery (inactive GO) are proposed subsequent to extensive counselling and shared decision	(2, 0000)
9)	Counselling and selection of treatment for moderate-to-severe and active GO	Extensive counselling is warranted to explain aims and expectations, benefits and risks of different therapies. Selection of treatment relies on evidence-based effectiveness, safety, evaluation of costs, reimbursement by the health system, drug availability, facilities for delivering highly specialized treatments, and personal choice of the informed patient within a shared decision-making process.	(1, 0000)
10)	Cumulative dose of intravenous glucocorticoids	The cumulative dose of intravenous (iv) glucocorticoids should not exceed 8.0 g for each cycle; GO patients with evidence of recent viral hepatitis, significant hepatic dysfunction, severe cardiovascular morbidity, uncontrolled hypertension, should not be administered iv glucocorticoids; diabetes should be well controlled before starting treatment. We strongly recommend that such a treatment be applied in experienced centres only that manage potentially serious adverse events	(1, 0000)
11)	Cumulative and single dose of intravenous glucocorticoids	An intermediate dose of iv glucocorticoids, i.e., a starting dose of 0.5 g iv methylprednisolone once weekly for six weeks, followed by 0.25 g once weekly for 6 weeks, cumulative dose 4.5 g, should be used in most cases of moderate-to-severe and active GO.	(1, 0000)
12)	Cumulative and single dose of intravenous glucocorticoids	High-dose regimen, i.e., a starting dose of 0.75 g iv methylprednisolone once weekly for six weeks, followed by 0.5 g once weekly for six weeks, cumulative dose 7.5 g, should be reserved for the more severe cases (constant/inconstant diplopia, severe proptosis, severe soft-tissue pathology or involvement) within the moderate-to-severe and active GO spectrum	(1, 0000)
13)	Glucocorticoid withdrawal	Clinicians should monitor each individual patient receiving glucocorticoid therapy for response to treatment and adverse events. When drug-induced side effects outweigh benefits, clinicians should consider withdrawing glucocorticoid treatment in favor of another modality, or watchful monitoring	(2, 0000)
14)	Local injections of triamcinolone	Local subconjunctival/periocular injections of triamcinolone acetate may be considered when systemic glucocorticoids are absolutely contraindicated	(2, 0000)
15)	Mycophenolate	Mycophenolate has a positive efficacy / safety profile in patients with moderate-to-severe and active GO, both as monotherapy and in combination with iv glucocorticoids	1, 0000
16)	Orbital radiotherapy	Orbital radiotherapy is considered an effective second-line treatment for moderate-to-severe and	(1, 0000)

		active GO, in combination with glucocorticoids, particularly in the presence of diplopia and/or restriction of extraocular motility	
17)	Cyclosporine	The combination of cyclosporine and oral glucocorticoids is a valid second-line treatment for moderate-to-severe and active GO	(1, ∅∅∅∅)
18)	Azathioprine	Consideration can be given to azathioprine as a second-line and glucocorticoid-sparing agent in combination with oral glucocorticoids	(1, ∅∅∅∅)
19)	Teprotumumab	Very promising drug with strong reduction of exophthalmos, diplopia and improvement of QoL. Currently second-line option as longer term data, availability, affordability, costs, and need for subsequent rehabilitative surgery are pending	(1, ∅∅∅∅)
20)	Rituximab	Rituximab can be considered a second-line treatment for patients with moderate-to-severe and active GO of recent onset (<12 months) if refractory to iv glucocorticoids, as long as dysthyroid optic neuropathy (DON) is excluded. We strongly recommend that such a treatment be applied in experienced centers only that manage potentially serious adverse events	(1, ∅∅∅∅)
21)	Tocilizumab	Tocilizumab may be given consideration as a second-line treatment for moderate-to-severe and active glucocorticoid-resistant GO	(1, ∅∅∅∅)
22)	First-line treatment for moderate-to-severe and active GO	Intravenous methylprednisolone in combination with oral mycophenolate sodium (or mofetil) represent the first-line treatment for moderate-to-severe and active GO	(1, ∅∅∅∅)
23)	First-line treatment for moderate-to-severe and active GO	In the more severe forms of moderate-to-severe and active GO, including constant/inconstant diplopia, severe inflammatory signs and exophthalmos >25mm, intravenous methyl-prednisolone at the highest cumulative dose (7.5 g per cycle) as monotherapy represents an additional valid first-line treatment	(1, ∅∅∅∅)
24)	Second-line treatments for moderate-to-severe and active GO	If response to primary treatment is poor and GO is still moderate-to-severe and active, subsequent to careful ophthalmic and biochemical (liver enzymes) evaluation, the following second-line treatments should be considered: <ul style="list-style-type: none"> • Second course of iv methylprednisolone monotherapy, starting with high single doses (0.75 g) and a maximal cumulative dose of 8 g per cycle • Oral prednisone/prednisolone combined with either cyclosporine or azathioprine • Orbital radiotherapy combined with oral or intravenous glucocorticoids • Teprotumumab • Rituximab • Tocilizumab 	(1, ∅∅∅∅)

25)	Combination of orbital radiotherapy and iv glucocorticoids	Based on expert opinion only (as randomized trials are not available), the task force suggests combination of orbital radiotherapy and iv methylprednisolone as a potential second-line treatment for moderate-to-severe and active GO	(2, ∅∅∅∅)
26)	Treatment of sight threatening GO	Optic neuropathy should be treated immediately with high single doses of iv methylprednisolone (0.5-1 g of methylprednisolone daily for either three consecutive days or more preferably on every second day), and urgent orbital decompression should be performed if response is absent or poor within 1-2 weeks. Recent eyeball subluxation should undergo orbital decompression as soon as possible	(1, ∅∅∅∅)
27)	Treatment of sight threatening GO	Severe corneal exposure should be urgently treated medically or by means of progressively more invasive surgeries in order to avoid progression to corneal breakdown; the latter should be immediately surgically addressed	(2, ∅∅∅∅)
28)	Thyroid treatment in patients with GO	Mild and inactive GO: any treatment for hyperthyroidism can be used based on standardized criteria and patient choice	(1, ∅∅∅∅)
29)	Thyroid treatment in patients with GO	Mild and active GO: antithyroid drugs (ATDs) or thyroidectomy are preferred and prednisone/prednisolone prophylaxis should be used if RAI treatment is selected	(1, ∅∅∅∅)
30)	Thyroid treatment in patients with GO	Moderate-to-severe, longstanding and inactive GO: as for mild and inactive GO, but consideration should be given to prednisone/prednisolone prophylaxis if RAI treatment is selected and risk factors (smoking, high TSHR-Ab) are present	(1, ∅∅∅∅)
31)	Thyroid treatment in patients with GO	Moderate-to-severe and active GO: hyperthyroidism should be treated with ATDs until treatment of GO is completed	(1, ∅∅∅∅)
32)	Thyroid treatment in patients with GO	Sight threatening GO: in this emergency condition, treatment of GO is an absolute priority; hyperthyroidism should be treated with ATDs until treatment of GO is completed	(1, ∅∅∅∅)

TABLE 1: THE EUGOGO EVIDENCE-BASED RECOMMENDATIONS FOR THE MANAGEMENT OF GRAVES' ORBITOPATHY (GO)

TABLE 2.
ASSESSMENT OF ACTIVITY BY THE CLINICAL ACTIVITY SCORE (CAS)*

ASSESSMENT OF ACTIVITY

1. Spontaneous retrobulbar pain
2. Pain on attempted upward or downward gaze
3. Redness of eyelids
4. Redness of conjunctiva
5. Swelling of caruncle or plica
6. Swelling of eyelids
7. Swelling of conjunctiva (chemosis)

CAS <3 = Inactive GO; CAS \geq 3 = Active GO

A 10-item CAS, including an increase in exophthalmos of \geq 2 mm, a decrease in eye motility of \geq 8°, or a decrease in visual acuity in the last 1-3 months, is useful to assess progression of GO after the first visit.

*Modified according to Wiersinga et al. (14) and reproduced with the permission

TABLE 3 CLASSIFICATION OF SEVERITY OF GRAVES' ORBITOPATHY (GO)

Classification	Features
Mild GO	<p>Patients whose features of GO have only a minor impact on daily life insufficient to justify immunomodulation or surgical treatment. They usually have one or more of the following:</p> <ul style="list-style-type: none"> • minor lid retraction (<2 mm) • mild soft-tissue involvement • exophthalmos • <3 mm above normal for race and gender • no or intermittent diplopia and corneal exposure responsive to lubricants
Moderate-to-Severe GO	<p>Patients without sight-threatening GO whose eye disease has sufficient impact on daily life to justify the risks of immunosuppression (if active) or surgical intervention (if inactive). They usually have two or more of the following:</p> <ul style="list-style-type: none"> • lid retraction ≥ 2 mm • moderate or severe soft-tissue involvement • exophthalmos ≥ 3 mm above normal for race and gender • inconstant or constant diplopia
Sight threatening (very severe) GO	<p>Patients with dysthyroid optic neuropathy and/or corneal breakdown</p>