

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/145969/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Enticott, Gareth , Earl, Lynsey and Gates, M. Carolyn 2022. A systematic review of social research data collection methods used to investigate voluntary animal disease reporting behaviour. *Transboundary and Emerging Diseases* 69 (8) , pp. 2573-2587. 10.1111/tbed.14407

Publishers page: <https://doi.org/10.1111/tbed.14407>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Title

A Systematic Review of Social Research Data Collection Methods Used to Investigate Voluntary Animal Disease Reporting Behaviour

Running Title

Review of Methodologies to Assess Disease Reporting Behaviour

Accepted in *Transboundary and Emerging Diseases*, November 2021

Authors

Gareth Enticott^{§†}, Lynsey Earl[‡], and M. Carolyn Gates*

[†]Cardiff School of Geography and Planning, Cardiff University, King Edward VII Avenue, Cardiff CF10 3WA, United Kingdom

[‡] Diagnostic and Surveillance Services, Biosecurity New Zealand - Tiakitanga Pūtaiao Aotearoa, Ministry for Primary Industries - Manatū Ahu Matua, PO Box 2526 Wellington 6140 New Zealand

*School of Veterinary Science, Massey University, Private Bag 11-222, Palmerston North 4442, New Zealand

§Correspondence: enticottg@cardiff.ac.uk

Acknowledgements

We are grateful to the Ministry for Primary Industries (New Zealand) for supporting this work.

Ethics Statement

The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to. No ethical approval was required as this is a review article with no original research data.

Conflict of Interest Statement

The authors declare no conflicts of interest.

Summary

Voluntary detection of emerging disease outbreaks is considered essential for limiting their potential impacts on livestock industries. However, many of the strategies employed by animal health authorities to capture data on potential emerging disease threats rely on farmers and veterinarians identifying situations of concern and then voluntarily taking appropriate actions to notify animal health authorities. To improve the performance of these systems, it is important to understand the range of socio-cultural factors influencing the willingness of individuals to engage with disease reporting such as trust in government, perceived economic impacts, social stigma, and perceptions of 'good farming'. The objectives of this systematic review were to assess how different social research methodologies have been employed to understand the role these socio-cultural dimensions play in voluntary disease reporting and to discuss limitations to address in future research. The review uncovered 39 relevant publications that employed a range of quantitative and qualitative methodologies including surveys, interviews, focus groups, scenarios, observations, mixed-methods, interventions, and secondary data analysis. While these studies provided valuable insights, one significant challenge remains eliciting accurate statements of behaviour and intentions rather than those that reflect desirable social norms. There is scope to develop methodological innovations to study the decision to report animal disease to help overcome the gap between what people say they do and their observable behaviour. A notable absence is studies exploring specific interventions designed to encourage disease reporting. Greater clarity in specifying the disease contexts, behavioural mechanisms and outcomes, and the relationships between them would provide a more theoretically informed and policy relevant understanding of how disease reporting works, for which farmers, and in which disease contexts.

Keywords

Biosecurity; Disease Reporting; Farmer Behaviour; Veterinary Behaviour; Infectious Disease

1. Introduction

Voluntary detection of emerging disease outbreaks is considered to be an integral component of emergency preparedness and ‘essential for the timely detection, reporting and communication of occurrence, incursion or emergence of diseases’ ([World Organisation for Animal Health \(OIE\), 2021](#)). Governments may facilitate voluntary detection through various strategies such as developing rapid diagnostic tests, establishing surveillance networks, providing information to farmers and veterinarians, and creating dedicated reporting methods such as telephone or internet-based systems ([Hoinville et al., 2013](#)). As many of these strategies rely on individuals correctly identifying the clinical signs of disease and then voluntarily taking actions to inform the appropriate authorities, these voluntary animal disease reporting systems need to be culturally appropriate to be effective. Like other aspects of disease control, a range of socio-cultural factors such as trust in government, perceived economic impacts, social stigma, and perceptions of ‘good farming’ will influence when, who and how voluntary disease reporting occurs ([Gates, Earl, & Enticott, 2021](#)). Creating effective disease reporting systems therefore requires knowledge of how different socio-cultural factors influence farmers’ and vets’ decisions to report disease. However, understanding these factors requires overcoming significant methodological barriers. For example, retrospective accounts of disease reporting can be compromised by inaccurate recall ([Coughlin, 1990](#); [Gilbert, Häslar, & Rushton, 2014](#)) or social desirability bias ([Burton, 2004](#); [Krumpal, 2013](#)). Similarly, prospective accounts face the challenge of the value-action gap: the difference between how people describe their behaviour and their observable behaviour ([Kollmuss & Agyeman, 2002](#)). Accordingly, this paper has two aims. The first aim is to provide a systematic review of research methodologies used to investigate the social and behavioural challenges facing voluntary disease reporting systems. In doing so, the second aim is to promote discussion on effective methodological strategies to understand the effectiveness of voluntary disease reporting and benefit future research into voluntary disease reporting.

2. Methodology

2.1 Defining Voluntary Disease Reporting

In referring to the disease reporting practices of farmers and animal keepers, terminology refers to active and passive surveillance ([Hoinville et al., 2013](#)). In relying on farmers and animal keepers to report disease, surveillance systems therefore depend on voluntary behaviour. These voluntary behaviours occur along a chain of activities beginning with observation, discussion, and research through to the reporting of suspicious or clinical signs of disease to an official channel or organisation ([Gates et al., 2021](#)). The outcome of these behaviours is the early detection of disease by the responsible veterinary authority. We therefore define voluntary disease reporting as a process encompassing the autonomous actions of different stakeholders following a disease incursion that results in regulatory authorities being made aware of a potential disease threat. Our particular focus for this review is on the methodologies that can be used to explore what makes farmers who notice clinical signs in affected animals decide to contact their veterinarian or directly notify animal health authorities

Research on voluntary disease reporting is defined by attention to three different contexts (see figure 1). Firstly, the actor context refers to the behaviour of specific individuals in the chain of activities. For example, a farmer may report suspicions to his/her vet, other farmers, or farm advisors who may subsequently discuss it informally for advice with other colleagues or refer it to other veterinary authorities as part of a disease reporting system. Secondly, the disease context refers to the temporal nature of disease reporting research: actors' behaviour may be discussed in relation to current disease threats, their behaviour in relation to past events, or anticipatory actions for future disease incursions. Finally, the agricultural context refers to how responses to disease reporting systems may vary across agricultural sectors and policy regimes.




Figure 1: Contexts of animal disease reporting research

2.2 Review Methodology

Figure 2 describes the methodology used for this review. Firstly, a literature search was conducted to identify relevant published academic studies relating to voluntary disease reporting and detection. The focus was on reporting of animal disease and the outcome of voluntary reporting behaviours, i.e. the early detection of disease. Where relevant, the review followed the PRISMA guidelines for systematic reviews (Page et al., 2021) and was undertaken using Scopus (www.scopus.com) using the following search terms :

- disease AND "early detection" AND animal AND farm* AND decision*
- disease AND reporting OR detection OR surveillance AND animal OR crop AND farm* AND decision* AND (LIMIT-TO (SUBJAREA , "SOCI"))

The two searches generated 710 and 742 documents respectively. Each document in each search was analysed for relevance and removed from the sample where appropriate. As the review focussed on social science methodologies, initial study selection was made by (author 1) to reflect their expertise. Subsequently, all authors assessed whether studies fitted the definition of voluntary disease reporting and were cross-checked with a parallel review undertaken by author 2 reported elsewhere (anonymised

reference). This led to the exclusion of studies where they focused on generic biosecurity preventive behaviours. Studies were included where they conformed to our definition of disease reporting systems and any of the disease contexts described in Figure 1.




Figure 2: Review Methodology

Studies were not limited to farmers or past disease outbreaks. Literature searches focused on the social scientific literature, but this did not preclude social scientific studies published in mainstream veterinary

journals (e.g. Preventive Veterinary Medicine). Thirteen additional documents were added following a process of ‘snowball reviewing’ ([Greenhalgh & Peacock, 2005](#)): using Scopus, citations of the most highly cited and relevant documents identified in the initial reviews were analysed and included where relevant. In addition, one relevant unpublished Ph.D. thesis was included in the list of documents. Grey literature was not included in the review: Scopus does not contain unpublished policy reports and snowball reviewing did not identify any additional unpublished studies to include.

In total, 39 publications were included in the review which either used social research methods (e.g. interviews, surveys) to analyse farmers’ behaviour, or analysed secondary data to explore the impacts of policy changes to farmers’ decisions (results of the review are shown in appendix 1). Thirty-two exclusively focused on the subject of reporting disease incidence. A further six papers were included that explicitly mentioned disease surveillance and disease reporting in the paper’s abstract, but also provided information on other biosecurity activities. Finally, one paper was included that referred to an on-farm disease alerting system. Six papers were identified employing theoretical or conceptual modelling to explore the effectiveness of financial rewards and penalties. As these are conceptual in nature and do not rely on social research methods, they are excluded from the final analysis. Five papers dealt with crop or plant diseases. These were retained because they conformed to the definition of Voluntary disease reporting, their methods were relevant to animal disease, and the underlying behavioural mechanisms are similar.

The quality of studies was not adjudicated prior to inclusion. The majority of studies provided a high-level of detail when describing the social research methods used in the study. All studies provided details on the number of research participants: the range varied considerably (from 5 – 1140) reflecting the varied demands of different research approaches. Methodological details tended to be more specific when identifiable conceptual frameworks were used. For many other qualitative studies, however, methodological description was limited to broader descriptions such as pointing to the topics discussed in focus groups or interviews, rather than the precise wording of the questions.

3. Results

3.1 Disease Reporting Contexts

For the actor context, most papers (29) focused on farmers. Of these, 20 were exclusively focused on farmers with two (2) focussed on smallholders. Other papers examined farmers' views alongside other food chain actors including: veterinarians (4), food processors (1) and farm advisors (2). Five papers focused exclusively on vets and disease reporting and a further one examined vets and policy actors' views.

For the agricultural context, nine publications examined disease reporting in all forms of livestock. Eight studies analysed disease reporting in the cattle sector. Six studies focused on pigs. Studies of disease reporting in sheep were less common: two papers exclusively focussed on sheep and another examined sheep and cattle disease reporting together. Poultry accounted for three papers. The review found two in the aquatic sector, one study relating to equine disease reporting, one for bees and two for crops. In relation to the disease context, most papers (26) did not distinguish between different diseases, of which nine did not distinguish between livestock sectors. Most papers (29) examined actors' responses to endemic disease outbreaks: no studies examined disease reporting systems that had contributed to disease eradication. Ten papers were explicitly oriented towards finding out actors' intentions towards future disease threats, whether these were diseases where there had been previous outbreaks (such as Foot-and-Mouth Disease) or none recorded (such as varroosis).

3.2 Descriptive Summary of Methodologies

Most of the publications included in the review (26) used only one research method with the remainder employing a 'mixed methods' ([Ivankova, Creswell, & Stick, 2006](#)) research design. The most frequently used method was interviews (20) followed by surveys (18) and focus groups (9). Eleven studies exclusively used a survey (whether it was online, postal, using the telephone, or in person). In addition, six other studies used a survey alongside other methods. Methods are equally distributed across different subjects and sectors. Studies that focused on all diseases across all sectors did, however,

exclusively use survey methods. The following sections provide further details on these methodologies and how they have been employed to understand disease reporting.

3.2.1 Surveys

Studies relying on survey methodologies employ a range of different approaches. Some rely on open-ended or generic survey questions collecting information on practices and knowledge, whilst others rely on more advanced conceptual frameworks that collect quantitative data using Likert scale questions for advanced statistical modelling. Examples of the former include two papers by ([Hammond, Hardie, Hauser, & Reid, 2016a, 2016b](#)) investigating whether a high priority pest would be reported to the Australian crop disease surveillance programme. These surveys point towards gaps in knowledge in surveillance programmes and respondents' confidence in their ability to correctly identify plant pests. Similarly, [Guinat, Wall, Dixon, and Pfeiffer \(2016\)](#) study on disease reporting in pigs in the UK uses a range of questions relating to respondents' knowledges and perceptions, showing that farmers with low levels of knowledge, concern and awareness of African Swine Fever are less likely to report it ([see also Hernández-Jover, Higgins, Bryant, Rast, & McShane, 2016](#)).

Other studies have attempted to operationalise frameworks such as the Theory of Planned Behaviour or Health Belief Model using structured surveys with Likert-type response scales, followed by quantitative analysis and modelling. For example, [Palmer, Sully, and Fozdar \(2009\)](#) study of disease reporting is informed by aspects of the Health Belief Model (HBM) which emphasise the role of perceived risks, barriers to reporting, trust, control beliefs and self-efficacy, position within the community and attitudes towards surveillance. [Wright, Jorgensen, and Smith \(2018\)](#) rely on the Theory of Planned Behaviour to analyse intentions to report disease amongst Australian livestock producers to either the Government or private vets. Similarly, [Pramuwidyatama, Hogeveen, and Saatkamp \(2020\)](#) survey instrument is based on the theory of planned behaviour in order to quantitatively analyse motivations of broiler farmers to prevent Avian Influenza. The study shows that farmers responding to the survey tended not to be “aware of the clinical signs of HPAI; hence, only severe HPAI outbreaks will be reported. As a result, information about HPAI is less obvious and “trustful” (p.7).

Structured questionnaires have also been used to compare reporting behaviours between farmers according to their exposure to disease. [Gilbert et al. \(2014\)](#) use a survey of 69 cattle farmers to assess factors associated with farmers' decisions to submit biological samples as a means of identifying early stages of infection. The survey posed two different kinds of questions to farmers: firstly, 'prospective assessment' in which farmers were asked how often they performed specific actions, and secondly 'retrospective assessment' in which farmers were asked to recall specific events over the last 6 months. Analysis revealed that farmers prospective assessments generally over-estimate actual disease reporting. In addition, farmers were asked questions about their attitudes and motivations towards veterinarians and disease surveillance. Results also indicated that farmers who were members of herd health schemes would be more likely to submit samples. Likewise, [Randrianantoandro, Kubota, and Kono \(2018\)](#) use a questionnaire survey of 201 pig farmers in Madagascar about African Swine Fever (ASF) in order to analyse their responses to financial compensation. Their results show that as compensation increases, more farmers are willing to report ASF cases but this also depends on farm-related characteristics, farmers' knowledge about ASF, administration of the classical swine fever vaccine, and previous experiences with ASF.

Other studies combined structured and open-ended questions within a survey to explore and explain differences in farmers' disease reporting practices. [Lupo, Osta Amigo, Mandard, Peroz, and Renault \(2014\)](#) adopted a retrospective case-control design in which 27 non-reporting and 89 reporting farmers were surveyed by telephone using this approach. Analysis of farmers' reasons for not reporting disease, such as the absence of compensation, is illustrated both quantitatively and qualitatively using quotations from open-ended survey questions. Separate quantitative modelling is conducted to show how different factors (such as farm size, prior receipt of financial compensation) are associated with farmers who have reported disease and those who have not.

3.2.2 Interviews

Although interviews are a popular method to explore disease reporting, the studies reveal a diversity in approaches. Face to face interviews can be used to collect objective information, but as this information could be collected by a questionnaire, the value of interviews lies in using qualitative semi-structured approaches to explore in-depth phenomena such as behaviour and decision-making. For some studies in the review, detailed methodological descriptions are provided, but in others interview topic guides, questions and approach are not provided ([Garza, Ågren, & Lindberg, 2020](#); [Palmer, Fozdar, & Sully, 2009](#)). Where studies looked at reporting for all diseases across a sector, interviews took a broad approach, often not directly referring to disease reporting. For instance, the open-ended interview schedule in Fox et al's ([2020](#)) study of shellfish in Northern Ireland covers a range of biosecurity related issues and enables the authors to report generic attitudes and behaviours pertaining to disease reporting. From this, [Fox et al. \(2020\)](#) report how farmers' perceptions of stress, predation, and luck mean that dead oysters are not reported and thrown away despite requirements to report them. Similarly, two studies by [Sawford, Vollman, and Stephen \(2012\)](#); ([2013](#)) examined veterinarians' decision making about diagnostic submissions in Canada and Sri Lanka. Both studies employed the same open-ended interview schedule that revolved around pre-established key themes, such as decision making around laboratory submissions and participation in disease monitoring.

Interview methods were more specific where they examined specific diseases or practices. For example, [Bronner, Hénaux, Fortané, Hendriks, and Calavas \(2014\)](#)'s study analyses why farmers do not report abortions in relation to brucellosis control. Firstly, their research design seeks to differentiate respondents between areas of recent and historical outbreak histories. Secondly, their questions are specifically directed to understanding abortion as an indicator of disease, and farmers' relationships with the surveillance system. Thus, interviews focused on farmers' own definitions of abortion and their knowledge of surveillance systems.

The approach adopted in [Burns, Ribble, McLaws, Kelton, and Stephen \(2013\)](#)'s study of smallholders' attitudes to avian influenza in Canada is both generic and specific. The interview schedule combines generic questions about biosecurity, context, and motivations for keeping poultry with more specific

follow-up questions in relation to the health problems farmers mention in the interview. Although studies like [Burns et al. \(2013\)](#) seek to gather some contextual background, there are specific interview techniques that aim to situate behaviour in the historical context of each farm. A good example is the Biographical Narrative Interpretive Method (BNIM) used by [McFarland et al. \(2020\)](#) in their study of factors influencing sample submission to animal health surveillance services in Ireland. The BNIM relies on a single question for inducing narrative (referred to as the SQUIN) which gives the opportunity to recount the history of a particular incident or event and what the respondent considers to be the most significant factors and changes that influenced that event. In this case, the SQUIN was:

"As you know, I'm interested in how farmers make decisions to send or not send carcasses or fetuses to the lab. Can you tell me the story of when it happens that an animal dies on the farm?"

In McFarland et al.'s study, participants with different levels of engagement with the surveillance system were selected: some were regular submitters, others occasional, and the remainder who had never submitted any samples. [McFarland et al. \(2020\)](#)'s study sample was relatively small (five BNIM interviews were conducted), but it can be used with more respondents if necessary (McFarland et al.'s study also used focus groups). Other studies of tree health by [Porth, Dandy, and Marzano \(2015\)](#) provide a similar historical narrative to understanding disease reporting. These methods allow the authors to trace the network of actors involved in the reporting and management of an outbreak of Asian Longhorn Beetle (ALB) in Kent, England. In tracing this history, the study is able to provide details on the reasons why the outbreak was discovered. Although detailed interview methods like the BNIM are not referred to in the paper, this detailed retrospective social network approach provides a potentially more useful research design compared with interviews with people who have not encountered disease.

3.2.3 Focus Groups

Three studies in the review exclusively used focus groups to analyse disease reporting. Firstly, [Robinson and Epperson \(2013\)](#); [Robinson et al. \(2012\)](#) use focus groups to explore disease reporting practices amongst private vets in Mississippi (US) and Northern Ireland (UK). Both studies broadly

adopted the same approach in which general questions about the decision and process of sample submission was discussed. The subsequent discussion elicited a range of factors that influenced sample submission, which were both social and technical. For example, veterinarians reported that farmers had become keener to submit samples but this depended on the social characteristics of farmers. Veterinarians also pointed to technical challenges of sample submission, such as health and safety issues, inadequately packaged samples and biosecurity issues associated with transportation. Importantly, the focus groups also highlighted how the relationship between veterinarians and pathologists was an important part of the reporting process. Pathologists were not included in this study, nor do they appear to be part of other studies.

3.2.4 Using Scenarios

For both qualitative interviews and quantitative survey methodologies, scenarios were used as a means of eliciting participants' intentions towards disease outbreaks. Scenarios represent a helpful methodological device to stimulate thinking or challenge preconceptions and to examine the causes and consequences of what may happen in future ([Quine et al., 2011](#)). They are likely to work best when they are plausible ([Oreszczyn & Carr, 2008](#)), developed consensually ([Oreszczyn, Lane, & Carr, 2010](#)) and apply to a specific purpose ([van der Heijden, 1996](#)). [Bradfield, Wright, Burt, Cairns, and Van Der Heijden \(2005\)](#) suggest scenario methodologies work best in relation to four purposes (making sense of puzzling situations; developing strategy; anticipation; and adaptive organisational learning) all of which apply to voluntary disease reporting. In health psychology, scenarios are referred to as vignettes, which may take the form of text, pictures or video ([Hughes & Huby, 2002](#)) and used to elicit attitudes and beliefs about complex and potentially sensitive situations ([De Bruyn et al., 2020](#)). Nevertheless, where they are poorly designed and/or executed ([see Gould, 1996](#)), vignettes may fail to capture the reality of people's lives, limiting their generalizability ([Hughes & Huby, 2002](#)).

Examples of this approach included [Pham et al. \(2017\)](#) who used a scenario based discrete choice experiment to investigate willingness to report swine diseases in Vietnam. The survey presented a series

of choice-sets from which farmers would choose from two reporting options and a non-reporting option under three different policy scenarios. The exemplar scenario given to respondents was:

“Imagine that more than 40% of your pigs are sick with clinical signs as suspected cases from one of the three notified diseases (PRRS, FMD, and CSF). Your sick pigs did not respond after 3–5 days of treatment and the live weight of fattening pigs ranged from 20 to 50 kg. Choose one of the following options related to disease reporting to report swine disease in your holding to veterinary or local authorities”.

[Pham et al. \(2017\)](#) conclude that the most important factors in reporting are the type of culling implemented and the likelihood (as opposed to the amount) of receiving compensation. Importantly, the choice-sets relied on a scenario to generate responses. [Delgado et al. \(2014\)](#) also use scenarios in their analysis of intentions to report foot-and-mouth disease amongst cattle producers in the USA. Their study used a two-stage research design in which two surveys were returned by the same respondents over a 6-month period with a different scenario in each. This approach allows [Delgado et al. \(2014\)](#) to show how disease reporting intentions and perceptions are linked to different disease scenarios.

In qualitative research, [Pfeiffer \(2018\)](#) utilises a scenario-based approach to explore disease reporting. Three hypothetical scenarios were presented to farmers which were preceded by a general discussion about biosecurity issues and use of veterinary services on farms. Scenarios were presented alongside a picture of a sheep farm, although the picture was not relevant to the scenario itself. Following the discussion, focus group participants were asked to rank the barriers and incentives in order of importance. Results from the focus groups revealed the importance of context, which was defined as farmers’ sense of self as a sheep farmer. This meant that self-reliance was a key strategy in resolving any problems rather than contacting a veterinarian which would happen when they had reached the limit of their knowledge and experience. Other qualitative studies using scenario methods include [Hamilton-Webb, Naylor, Little, and Maye \(2016\)](#) who use three scenarios to elicit participants’ views on disease reporting under different contexts and policy environments. In their case, the scenarios were co-

developed with the UK government department responsible for disease management, although such co-production of realistic scenarios could involve a wider range of relevant participants. The scenarios used reflected different compensation options, and participants were asked opinions of the likely reactions to the scenario.

3.2.5 Ethnographic Observation

One study used ethnographic methods to examine farmers' disease reporting practices. Whilst ethnography uses a range of methods such as interviewing ([Hammersley & Atkinson, 2019](#)), it is often associated with different forms of observation. For example, in [Phillips \(2020\)](#) study of beekeeping and potential infestation with varroa mite, interviews with beekeepers and policy officers was accompanied by participant observation at meetings, events, field-days, club beekeeping, and training. Research activities were also spread over a period of seven years. The longitudinal and observation focus allows Phillips to show how a sense of anticipation and 'good beekeeping' is created as a means to encourage early reporting of signs of varroa mite infestation.

3.2.6 Mixed Methods

Mixed methods refers to the use of a combination of qualitative and quantitative methods to capitalise on their respective strengths ([Creswell & Clark, 2017](#)). The approach allows small scale research (such as semi-structured interviews) to sequentially inform methodologies (such as surveys) used on a larger scale ([Ivankova et al., 2006](#)). Alternatively, different forms of data can be triangulated to provide 'convergent validation' ([Fielding, 2012](#)). A third of the studies (13) used a combination of methods, such as semi-structured interviews and large-scale quantitative surveys. Examples of sequential mixed methods designs included [Delgado, Norby, Dean, McIntosh, and Scott \(2012\)](#); ([Delgado et al., 2014](#)) who used semi-structured interviews to design a structured survey based around the Theory of Planned Behaviour. In this study, the aim was to examine farmers' likely reactions to an outbreak of foot-and-mouth disease in Texas (US). Telephone interviews were initially used to understand respondents' knowledge of FMD, before being asked to comment on the feasibility of an outbreak scenario. As such, participants were drawn from a range of public and private organisations that would be involved in

outbreak management. Subsequently, a survey instrument was developed based on the Theory of Planned Behaviour and the results from the telephone survey, which was subsequently evaluated and modified during a 2-day workshop. In doing so, it highlights the importance of mixed methods when researching disease reporting.

Other studies that used this sequential mixed methods approach included two studies by Elbers et al ([2010](#); [2010](#)) that analysed farmers' disease reporting behaviours for classical swine fever and avian influenza. The first study on avian influenza analysed poultry farmers' and veterinarians' reporting practices employing focus groups, semi-structured interviews and lastly an electronic survey. The focus groups and interviews involved farmers and veterinarians and highlighted a range of different themes relating to the failure to report disease and identified potential solutions to disease reporting barriers. Finally, a structured survey focused on the conditions required to report a suspicious case, barriers to reporting and the respondents' feelings and economic consequences of reporting a suspicious incident. Other mixed studies began with quantitative methods. For example, [Hernández-Jover et al. \(2016\)](#) used results from a postal survey of 919 Australian farmers to select interview participants by stratifying farmers according to demographic and husbandry practices. [Limon et al. \(2014\)](#) used a survey containing projective questions such as asking what would 'most people in the community do if almost all their animals die within one week?'. The results were subsequently discussed at focus groups and interviews.

3.2.7 Intervention Studies

Four studies sought to analyse disease reporting practices by analysing a specific intervention or policy change designed to help farmers and other stakeholders submit samples or report suspected disease. [Rao and Zhang \(2020\)](#) use a natural experiment of different insurance regimes in the pig industry in China. Using a structured in-person survey, results showed that insurance was associated with an 18.2% increase in likelihood of reporting diseased animals, but that insurance also acted as moral hazard increasing pig mortality (see [Zhang, Zhu, and Turvey \(2016\)](#)). [Struchen et al. \(2016\)](#) provide an example of an intervention study examining the use of a disease reporting tool developed to be used by

mobile devices. The application was designed to collect disease information on equine diseases and can be used only by registered ‘sentinel practitioners’, i.e. equine veterinarians. The study analysed disease reporting using the application and found that disease reporting continued to increase compared with previous years. Telephone interviews with participating veterinarians explored motivations for using the application, reasons for low levels of reporting, and the use of reporting devices. Low reporting was connected to disease absence, technical difficulties, and lack of benefits. In contrast, [Beyene et al. \(2018\)](#)’s compared disease reporting amongst 12 final year veterinary students, six of which were provided with a smartphone application. Comparing the two groups revealed that the smartphone group reported disease faster than those without it, although delays were encountered due to poor cell-phone reception.

In the management of tree-health, mobile applications and citizen-science approaches have proved a popular way of enhancing disease surveillance. In the UK, the ‘Observatree’ initiative has been analysed by [Crow et al. \(2020\)](#) and [Hall, Raum, Morris, and O’Brien \(2017\)](#). The Observatree initiative is a programme of activities designed to give people interested in tree health the skills and resources to identify priority tree pests and diseases, thereby enabling them to report them. For this project, tree health data was submitted online which are then shared with relevant authorities such as government pathologists and entomologists. Other initiatives in other countries follow a similar procedure. For example, the LIFE ARTEMIS project in Slovenia used a mobile application called “Invazivke” to allow diseases to be easily reported ([Crow et al., 2020](#)). Evaluations of these projects have focused on the aims, motivations, educational outcomes, and reporting practices of those volunteers involved to understand their motivations and tracking the social outcomes ([Hall et al., 2017](#)).

3.2.8 Secondary Data Analysis

One study used secondary data analysis to analyse the influence of veterinarians and other local stakeholders on farmers’ disease reporting habits. [Bronner, Morignat, and Calavas \(2015\)](#)’s study relies on observable behaviour as a way of avoiding the bias common to farmers’ stated preferences/behaviours found in other studies ([for example Gilbert et al., 2014](#)). Such analyses are of

course limited by the availability of data, and the data itself: attitudinal, motivational, and other social factors are not routinely collected so these analyses cannot incorporate them. Bronner et al's study compares the number of veterinarians in specific regions of France to levels of disease reporting by farmers. Results show that whilst disease reporting significantly varies between agricultural sectors and herd sizes, so does the size of veterinary practices in each area. Specifically, in areas where veterinary practices had more than one veterinarian, farmers were more likely to report disease. The absence of other contextual data poses limitations to this analysis. However, where these data exist, they may be utilised to explore the relationships between structural constraints to disease reporting and the role of the veterinary profession.

4. Discussion

This review has highlighted the range of social science methods used to investigate behavioural factors associated with the voluntary reporting of animal disease. The range of methods used is unsurprising given that disease reporting involves a range of activities involving different actors, and which allows research questions to be posed in a variety of ways. Some methods may be better suited to answer some of these questions than others. For example, understanding how farmers make sense of what counts as a suspicious disease and the thresholds for reporting may be best derived from in-depth qualitative methods such as focus groups, interviews, and observation. Where studies are attempting to estimate the effect of different behavioural interventions or capture information at scale about farmers' behaviour, quantitative methods are more likely to be relevant. At the same time, however, the familiarity of these methodological approaches also points towards specific improvements and innovations that could be used to improve understandings of voluntary disease reporting. These are discussed in turn below.

Firstly, a significant challenge facing studies of voluntary disease reporting is using methods that elicit accurate statements of behaviour ([Gilbert et al., 2014](#)), rather than those that reflect desirable social norms ([Burton, 2004](#)). In fact, beyond comparing subjective accounts with objective reports (see Gilbert

et al, 2014), no studies in the review seek to compare these objective and subjective reports, or provide significant new insights into how to overcome the gap between values and actions. The use of prospective questions that are framed in relation to the action of others (such as good or bad farmers, neighbours, or specific contexts) can potentially help to overcome issues of social desirability bias. However, many of the studies in the review lack detailed information on their precise question wording making it difficult to ascertain whether this approach has been used. Other approaches such as the biographical narrative approach used by [McFarland et al. \(2020\)](#) may also be useful given their reliance on more general and open-ended questions which allow the respondent to tell their own story, situate it within their own context and validate their own experiences.

The use of scenarios is another way in which to explore farmers' intentions in non-judgmental ways ([Soleri & Cleveland, 2005](#)). Whilst the review found several studies that used these approaches, there is potential to adapt how they are used. In other studies, participatory and interactive scenarios have been used, allowing participants to dynamically and recursively map key issues in discussing their own and others' responses to a given situation ([Oreszczyn & Carr, 2008](#)). For example, [Khan et al. \(2020\)](#) developed a card game to explore support for different regulatory approaches to anti-microbial resistance. This game involved a simple card-sorting exercise in which participants were asked to associate actors with specific characteristics. Similarly, [Maye, Enticott, and Naylor \(2017\)](#) use an influence mapping exercise in which participants were asked to identify a range of biosecurity 'influencers' and place them on concentric circles to indicate their importance to them in different biosecurity scenarios. These interactive methods combined with scenarios can potentially make a greater methodological contribution to understanding disease reporting.

Other innovative methods have been developed in order to address the methodological challenges of the values-action gap, particularly where the actions are controversial or test social expectations. For example, the randomised response technique (RRT) is widely used in research relating to behaviours that may be illegal or on the boundary of illegality and appropriate conduct ([Lensvelt-Mulders, Hox, van der Heijden, & Maas, 2005](#)). When using the RRT, the question-answer process is subject to a

randomising device (usually a device) which indicates whether the respondent should answer the question truthfully, or whether pre-determined answers (such as yes or no) are recorded. As the randomising device is not visible to researchers, the process provides the respondent with a sense of security when providing answers that may be incriminating. In agricultural studies, this method has been used to assess the level of illegal wildlife culling ([Cross, St. John, Khan, & Petroczi, 2013](#)). Other more complex games may also be used to simulate disease reporting behaviours. For example, [Merrill, Koliba, et al. \(2019\)](#); ([Merrill, Moegenburg, et al., 2019](#)) developed a biosecurity computer game to examine how rewards and penalties influenced whether biosecurity actions were taken (such as showering before entering a biosecure pig unit). Whilst these games simulated behavioural responses in disease management, arguably they can help distance the participant from more formal research settings which appear as ‘accountability settings’, which are more likely to elicit socially desirable behaviours.

Secondly, another way of dealing with the values-action gap is through different research designs, such as the comparative approach used by [Lupo et al. \(2014\)](#). However, the fact that the review finds only one study explicitly employing a case-control research design suggests limited methodological innovation in approaching this challenge. Whilst other studies, for example [Bronner et al. \(2014\)](#), [Kuchler and Hamm \(2000\)](#) and [Rao and Zhang \(2020\)](#), try to account for disease incidence by conducting research in different areas or take advantage of changes to long-term policies, other methodologies and research designs should be employed to assess farmers’ behaviour. The use of mixed methods may also help in this respect, either by allowing triangulation of different data sources, or using one method to construct culturally relevant and meaningful survey questions, and/or identifying and selecting appropriate research participants. Whilst some studies used this approach ([Delgado et al., 2014](#); [Elbers, Gorgievski-Duijvesteijn, Zarafshani, et al., 2010](#)), there is potential for greater use of a mixed methods design in future studies of disease reporting.

Another way of managing the challenge of the values-action gap is through studies of interventions designed to improve voluntary reporting of disease. However, it was surprising to find few studies of

specific interventions seeking to improve disease reporting. Some studies such as [Struchen et al. \(2016\)](#) examined the role of mobile apps, but there were no examples of controlled trials of different interventions. Elsewhere, other studies have analysed the use of on-farm sensors used to alert farmers of potential health and welfare problems in livestock ([Bruijnjs, Hogeveen, Garforth, & Stassen, 2013](#); [Eckelkamp & Bewley, 2020](#); [Mazrier, Tal, Aizinbud, & Bargai, 2006](#)). As the potential of sensors and other wearable technologies increases, such interventions may become more relevant to unexpected and exotic diseases. However, the promise of technological fixes suggested by the emergence of smart agricultural technologies may be compromised by social and economic dimensions of farming affecting their adoption ([Rotz et al., 2019](#)). Unless there are unpublished studies of interventions designed to improve disease reporting, the review indicates a sizeable research gap that can be filled by future intervention studies.

Finally, the review also suggests the need for greater theorisation of how disease reporting ‘works’ and can be improved, particularly given the potential for intervention studies. Approaches to policy evaluation stress the need for studies to be organised around a ‘theory of change’ which conceptualise how policies should work. For Pawson and Tilley ([Pawson, Greenhalgh, Harvey, & Walshe, 2005](#); [Pawson & Tilley, 2004](#)), a theory of change should encompass the relationships between contexts (C), mechanisms (M) and outcomes (O). Understanding the CMO relationship, they argue, is crucial to understanding what works for whom and how. Adopting this approach to analysing disease reporting could offer significant advantages, not least because existing studies often elide different contexts and ways disease reporting works, and multiple outcomes. For example, the review found that in many studies, disease contexts, and agricultural contexts are combined meaning the specificity of contextual influences is lost. Indeed, the majority of papers had no specific focus on any particular disease. Even where studies have analysed a range of diseases, there remains a need to understand the interactions between different diseases and control efforts: are farmers’ attitudes towards disease control for one disease the same as others he/she faces, and is so why does this occur? This question is not adequately answered in the studies in this review, which instead seem to conceptualise farmers as having universal

responses to disease rather than diseases. In this sense, this review suggests the need for further contextually specific studies both in terms of the diseases analysed and the agricultural sectors involved.

Alongside context, mechanisms refer to specific behavioural responses that are in some way triggered by an event, intervention, or policy. Mechanisms therefore reflect an identifiable pattern of action that can be analysed in some way. In Pawson and Tilley's terms, mechanisms are therefore not the policy itself, but in terms of disease reporting reflect farmers' responses and causal motivations. Pawson and Tilley suggest that not only should policies be based on these likely causal mechanisms, but that also policy evaluation should be based upon them. Where they are not explicit, researchers should estimate how policies are seeking to have a behavioural influence. [Maye, Enticott, and Naylor \(2020\)](#) provide an example of how mechanisms operate in animal health, identifying 'seeing is believing' as a mechanism to promote vaccine acceptability and confidence. By involving farmers in the process of wildlife vaccination, allowing to see it being conducted and its uncertainties managed, this mechanism should lead to outcomes of increased confidence in vaccines.

Whilst this approach can help understand how and for whom animal disease policies and initiatives work, this review finds a very narrow focus on specific behavioural mechanisms, if at all. For example, [Bronner et al. \(2015\)](#) connect larger veterinary practices with increased disease reporting, but the precise causal mechanisms for this is unclear: does contact with a greater variety of vets promote greater social responsibility, and if so why? The extent to which these behavioural mechanisms are evaluated is not present in this or other studies. One exception is Hall et al's (2017) analysis of the Observatree project whose analysis implicitly considers how developing social capital and a shared sense of responsibility can contribute to disease reporting. [Phillips \(2020\)](#) study of potential infestation with varroa mite also highlights how a 'sense of anticipation' is crucial to disease reporting and can be created by various organisations and activities.

Where studies do focus on behavioural interventions – specifically the role of compensation – their evaluation causal mechanisms are nevertheless limited. These studies tend to focus on farmers as

economic maximisers, that is the provision of compensation will promote behavioural responses that are consistent with maximising profit. Studies of the effects of changes to compensation regimes such as [Kuchler and Hamm \(2000\)](#) make no attempt to unpack the varied motivations and mechanisms by which farmers respond to financial incentives. As studies in the review show, other social factors are also likely to shape these responses. However, studies in the review that explicitly analyse these factors – whether qualitative attempts to examine the role of self-identity or in surveys employing the theory of good farming – are largely focused on farmers’ intentions, rather than their actual behaviour.

Contexts and mechanisms when correctly aligned should promote specific outcomes. Where papers in the review refer to outcomes, they relate to whether a disease is reported or not, and/or speed of reporting (for example [Beyene et al., 2018](#); [Struchen et al., 2016](#)). However, other studies in the review show, there are a range of social factors that influence the decision to report disease or not (for example [Bronner et al., 2014](#); [McFarland et al., 2020](#); [Palmer, Fozdar, et al., 2009](#)). It is therefore likely that attempts to influence disease reporting will also need to have other social outcomes in order to function efficiently. This is clearly shown in the analyses of citizen-science approaches to disease reporting whose success is dependent on the social capital generated by the activities they encompass ([Hall et al., 2017](#)). Methodologies used to evaluate disease reporting initiatives – as well as the interventions themselves – therefore need to be sensitive to these broader social outcomes to understand how and whether disease reporting works.

5. Conclusion

This paper presents the results of a systematic review of the methodologies used to investigate behaviours towards voluntary animal disease reporting. Whilst early detection of disease is a key element of a successful disease surveillance and control programme, the review finds relatively few (39) published academic studies, some of which cover a range of different diseases and livestock sectors. The review finds a broad range of methodological approaches, involving both qualitative and quantitative methodologies. Methodological quality appears to be high, but a key recommendation for

future studies is to provide a greater level of methodological detail, such as the precise question wording used in personal interviews and focus groups. Moreover, there is scope to further methodological innovations to study the decision to report animal disease to help overcome the gap between what people say they do and their observable behaviour. A notable absence is studies exploring specific interventions designed to encourage disease reporting. Greater clarity in specifying the disease contexts, behavioural mechanisms, and outcomes, and the relationships between them would likely provide a more theoretically informed and policy relevant understanding of how disease reporting works, for which farmers and in which disease contexts.

1 Appendix

2

Table A1: Details of papers included in the systematic review

Publication	Primary Method	Other Methods	Disease Context	No. Participants	Methodological Details	Diseases	Sector	Participants
Fox M, Christley R, Lupo C, et al. (2020) Preventing and mitigating farmed bivalve disease: a Northern Ireland case study. <i>Aquaculture International</i> 28(6): 2397-2417.	Interviews		Retrospective	16	Interview schedule is provided	All	Aquatic	Farmers and processors
Lupo C, Osta Amigo A, Mandard YV, et al. (2014) Improving early detection of exotic or emergent oyster diseases in France: Identifying factors associated with shellfish farmer reporting behaviour of oyster mortality. <i>Preventive Veterinary Medicine</i> 116(1-2): 168-182.	Interviews		Retrospective	116	Case-control study design - interviews with farmers who have reported in the past and those that are currently reporting. Details of the questions are provided.	All	Aquatic	Farmers
Phillips C (2020) The force of Varroa: Anticipatory experiences in beekeeping biosecurity. <i>Journal of Rural Studies</i> 76: 58-66.	Observation	Interviews	Anticipatory	42 interviews	some details are provided	Varroa	Bees	Beekeepers
Beyene TJ, Asfaw F, Getachew Y, et al. (2018) A Smartphone-Based Application Improves the Accuracy, Completeness, and Timeliness of Cattle Disease Reporting and Surveillance in Ethiopia. <i>Frontiers in Veterinary Science</i> 5: 2.	Intervention Study		Contemporary	n/a	Analysis of the effectiveness of a smartphone disease recording system in Ethiopia	All	Cattle	farmers and vets
Sawford K, Vollman AR and Stephen C (2013) A Focused Ethnographic Study of Alberta Cattle Veterinarians' Decision Making about Diagnostic Laboratory Submissions and Perceptions of Surveillance Programs. <i>PLOS ONE</i> 8(5).	Interviews		Retrospective	10	A list of questions / topic guide is provided.	All	Cattle	Vets

Bronner A, Hénaux V, Fortané N, et al. (2014) Why do farmers and veterinarians not report all bovine abortions, as requested by the clinical brucellosis surveillance system in France? <i>BMC Veterinary Research</i> 10.	Interviews		Retrospective	20 interviews (12 farmers, 8 vets)	A topic guide is provided	Brucellosis	Cattle	Farmers and vets
Bronner A, Morignat E and Calavas D (2015) Respective influence of veterinarians and local institutional stakeholders on the event-driven surveillance system for bovine brucellosis in France. <i>BMC Veterinary Research</i> 11(1).	Secondary data analysis		Retrospective	n/a	The paper uses data from the surveillance system	Brucellosis	Cattle	farmers and vets
McFarland L, Macken-Walsh A, Claydon G, et al. (2020) Irish dairy farmers' engagement with animal health surveillance services: Factors influencing sample submission. <i>Journal of Dairy Science</i> 103(11): 10614-10627.	Interviews	Focus Groups	Retrospective	5	Interviews used the Biographical Narrative Method	All	Cattle	Farmers
Hernández-Jover M, Higgins V, Bryant M, et al. (2016) Biosecurity and the management of emergency animal disease among commercial beef producers in New South Wales and Queensland (Australia). <i>Preventive Veterinary Medicine</i> 134: 92-102.	Interviews	Survey	Anticipatory	182 surveys, 34 interviews	Interview topic guide is provided. Survey variables are provided.	All	Cattle	Farmers
Delgado AH, Norby B, Scott HM, et al. (2014) Distribution of cow-calf producers' beliefs about reporting cattle with clinical signs of foot-and-mouth disease to a veterinarian before or during a hypothetical outbreak. <i>Preventive Veterinary Medicine</i> 117(3-4): 505-517.	Survey	Use of scenarios	Anticipatory	524	scenario based survey, drawing on Theory of Planned Behaviour. Survey questions are provided.	FMD	Cattle	Farmers

Delgado AH, Norby B, Dean WR, et al. (2012) Utilizing qualitative methods in survey design: Examining Texas cattle producers' intent to participate in foot-and-mouth disease detection and control. <i>Preventive Veterinary Medicine</i> 103(2): 120-135.	Workshop	Interviews, use of scenarios	Anticipatory	40 interviews, unspecified number in workshops	Aim is to develop a survey instrument using qualitative methods. Survey is based on scenarios which are provided.	FMD	Cattle	Farmers
Hammond, N. E. B., D. Hardie, C. E. Hauser and S. A. Reid (2016). "Can general surveillance detect high priority pests in the Western Australian Grains Industry?" <i>Crop Protection</i> 79: 8-14.	Survey		Anticipatory	101	Survey variables not provided	All	Crops	Farmers and advisors
Hammond NEB, Hardie D, Hauser CE, et al. (2016) How would high priority pests be reported in the Western Australian grains industry? <i>Crop Protection</i> 79: 26-33.	Survey		Anticipatory	84 (28% response rate from 300)	Survey variables are provided.	All	Crops	Farmers and advisors
Struchen R, Hadorn D, Wohlfender F, et al. (2016) Experiences with a voluntary surveillance system for early detection of equine diseases in Switzerland. <i>Epidemiology and Infection</i> 144(9): 1830-1836.	Intervention Study	Interviews, survey	Contemporary	6	Research examines the use of a reporting intervention - a mobile reporting device. Survey variables are not provided.	All	Equine	vets
Robinson PA, Epperson WB, Huston CL, et al. (2012) Factors influencing diagnostic sample submission by food animal veterinarians in Mississippi. <i>Veterinaria Italiana</i> 48(1): 31-39.	Focus Groups		Retrospective	12	No detail on the form of the focus groups is given.	All	Livestock	Vets
Robinson PA and Epperson WB (2013) Farm animal practitioners' views on their use and expectations of veterinary diagnostic Laboratories. <i>Veterinary Record</i> 172(19): 503.	Focus Groups		Retrospective	11	Focus Group questions are provided	All	Livestock	Vets
Garza M, Ågren ECC and Lindberg A (2020) Nudging in Animal Disease Control and Surveillance: A Qualitative Approach to Identify Strategies Used to Improve Compliance With Animal Health Policies. <i>Frontiers in Veterinary Science</i> 7.	Interviews		Contemporary	24	Interview schedule is not provided.	All	Livestock	Vets / Policy

Sawford K, Vollman AR and Stephen C (2012) A Focused Ethnographic Study of Sri Lankan Government Field Veterinarians' Decision Making about Diagnostic Laboratory Submissions and Perceptions of Surveillance. PLOS ONE 7(10).	Interviews		Retrospective	40	A list of questions / topic guide is provided.	All	Livestock	Vets
Gilbert WH, Häsler BN and Rushton J (2014) Influences of farmer and veterinarian behaviour on emerging disease surveillance in England and Wales. Epidemiology and Infection 142(1): 172-186.	Survey		Contemporary	69	Surveys are used to calculate the statistical probability of reporting samples to a laboratory. Survey is provided.	All	Livestock	Farmers
Hernández-Jover M, Hayes L, Woodgate R, et al. (2019) Animal health management practices among smallholder livestock producers in Australia and their contribution to the surveillance system. <i>Frontiers in Veterinary Science</i> 6(JUN).	Survey		Retrospective	1140	survey variables are provided.	All	Livestock	Smallholders
Wright BK, Jorgensen BS and Smith LDG (2018) Understanding the biosecurity monitoring and reporting intentions of livestock producers: Identifying opportunities for behaviour change. <i>Preventive Veterinary Medicine</i> 157: 142-151.	Survey		Anticipatory	200	Survey based on Theory of Reasoned Action. Survey variables are provided.	All	Livestock	Farmers
Hamilton-Webb, A., R. Naylor, R. Little and D. Maye (2016). "Compensation and exotic livestock disease management: The views of animal keepers and veterinarians in England." <i>Veterinary Record</i> 179(20): 513.	Interviews	Focus Groups, use of scenarios	Retrospective	82 interviews, 9 focus groups (8-12 in each)	scenarios are provided, broad topics are provided for the interviews	All	Livestock	Farmers and vets
Limon G, Lewis EG, Chang YM, et al. (2014) Using mixed methods to investigate factors influencing reporting of livestock diseases: A case study among smallholders in Bolivia. <i>Preventive Veterinary Medicine</i> 113(2): 185-196.	Interviews	Focus Groups, survey	Retrospective	240	Topic list for the interviews is provided.	All	Livestock	Farmers
Guinat C, Wall B, Dixon L, et al. (2016) English pig farmers' knowledge and behaviour towards African swine fever suspicion and reporting. PLOS ONE 11(9).	Survey		Retrospective	109	Survey variables are provided.	African Swine Fever	Pigs	Farmers
Randrianantoandro, T. N., S. Kubota and H. Kono (2018). "Farmers' knowledge and incentive in reporting infectious animal disease: The case of african swine fever in madagascar." <i>Journal of Animal and Plant Sciences</i> 28(1): 270-279.	Survey		Contemporary	201	Contingent valuation	African Swine Fever	Pigs	Farmers
Elbers ARW, Gorgievski-Duijvesteijn MJ, van der Velden PG, et al. (2010) A socio-psychological investigation into limitations and incentives concerning reporting a clinically suspect situation aimed at improving early detection of classical swine fever outbreaks. <i>Veterinary Microbiology</i> 142(1): 108-118.	Interviews	Focus Groups, survey	Retrospective	15 focus group participants, 17 interviews, 409 survey responses	Survey questions are provided. No details on interview or focus group questions.	Classical Swine Fever	Pigs	Farmers

Zhang, Y., X. Zhu and C. G. Turvey (2016). "On the Impact of Agricultural Livestock Microinsurance on Death-Loss, Production and Vaccine Use: Observations from a Quasi-Natural Experiment in China." <u>The Geneva Papers on Risk and Insurance - Issues and Practice</u> 41(2): 225-243.	Observational data	Survey	Contemporary	405 surveys	Survey provides background demographic data. Uses data from a longitudinal insurance/compensation scheme	All	Pigs	Farmers
Rao, X. and Y. Zhang (2020). "Livestock insurance, moral hazard, and farmers' decisions: a field experiment among hog farms in China." <u>Geneva Papers on Risk and Insurance: Issues and Practice</u> 45(1): 134-156.	Survey	Observational data	Contemporary	135 longitudinal survey participants	Survey provides background demographic data. Uses data from a longitudinal insurance/compensation scheme	All	Pigs	Farmers
Pham HTT, Peyre M, Trinh TQ, et al. (2017) Application of discrete choice experiment to assess farmers' willingness to report swine diseases in the Red River Delta region, Vietnam. <i>Preventive Veterinary Medicine</i> 138: 28-36.	Survey	Use of scenarios	Anticipatory	196	statistical analysis of responses to determine farmers reporting behaviours, using scenarios in a discrete choice experiment. Scenarios are detailed in the paper	All	Pigs	Farmers
Burns TE, Ribble C, McLaws M, et al. (2013) Perspectives of an underrepresented stakeholder group, backyard flock owners, on poultry health and avian influenza control. <i>Journal of Risk Research</i> 16(2): 245-260.	Interviews		Retrospective	18	Interview questions are provided	Avian Influenza	Poultry	Smallholders
Pramuwidyatama MG, Hogeveen H and Saatkamp HW (2020) Understanding the Motivation of Western Java Smallholder Broiler Farmers to Uptake Measures Against Highly Pathogenic Avian Influenza (HPAI). <i>Frontiers in Veterinary Science</i> 7.	Survey		Anticipatory	203	Based on theory of planned behaviour. Some questions provided.	Avian Influenza	Poultry	Farmers
Elbers ARW, Gorgievski-Duijvesteijn MJ, Zarafshani K, et al. (2010) To report or not to report: A psychosocial investigation aimed at improving early detection of avian influenza outbreaks. <i>OIE Revue Scientifique et Technique</i> 29(3): 435-449.	Interviews	Focus Groups, Survey	Retrospective	33 farmers & 334 vet survey responses, 9 interviews, 14 participants in 4 focus groups	Survey questions are provided. No details on interview or focus group questions.	Avian Influenza	Poultry	Farmers
Pfeiffer CN (2018) Improving disease surveillance in Australia's sheep industries: investigations of syndromic surveillance, farmer behaviour and sheep trade networks. University of Melbourne. Unpublished PhD Thesis	Focus Groups	Use of scenarios	Anticipatory	33 (3xfocus groups)	Focus groups used scenarios for discussion which are provided	All	Sheep	Farmers

Kuchler, F. and S. Hamm (2000). "Animal disease incidence and indemnity eradication programs*." <i>Agricultural Economics</i> 22 (3): 299-308.	Observational data		Contemporary	n/a	Uses data from a longitudinal insurance/compensation scheme	Scrapie	Sheep	Farmers
Palmer, S., F. Fozdar and M. Sully (2009). "The Effect of Trust on West Australian Farmers' Responses to Infectious Livestock Diseases." <i>Sociologia Ruralis</i> 49 (4): 360-374.	Interviews		Retrospective	37 interviews, 455 surveys	Interviews, topic guide used but not provided	All	Sheep and Cattle	Farmers
Palmer S, Sully M and Fozdar F (2009) Farmers, Animal Disease Reporting and The Effect of Trust: A Study of West Australian Sheep and Cattle Farmers. <i>Rural Society</i> 19(1): 32-48.	Survey		Retrospective	37 interviews, 455 surveys	Survey variables are provided.	All	Sheep and Cattle	Farmers
Hall, C., S. Raum, J. Morris and L. O'Brien (2017). Observatree: Key lessons Qualitative study of the 'Observatree' citizen science project. https://www.forestresearch.gov.uk/documents/7252/Observatree_evaluation_report.pdf .	Intervention Study	Interviews, Focus Group	Contemporary	16 interviews	analysis of citizen science initiatives	All	Trees	All
Porth, E. F., N. Dandy and M. Marzano (2015). ""My garden is the one with no trees:" Residential Lived Experiences of the 2012 Asian Longhorn Beetle Eradication Programme in Kent, England." <i>Human Ecology</i> 43 (5): 669-679.	Interviews		Retrospective	11	no topic guide. Interviews trace actors in the outbreak.	ALB	Trees	All
Crow, P., A. Perez-Sierra, A. Kavčič, K. Lewthwaite, M. Kolšek, N. Ogris, B. Piškur, J. K. Veenvliet, S. Zidar, S. Sancisi-Frey and M. de Groot (2020). "Using Citizen science to monitor the spread of tree pests and diseases: Outcomes of two projects in slovenia and the UK." <i>Management of Biological Invasions</i> 11 (4): 703-719.	Intervention Study	Interviews	Contemporary	900/500 survey responses	analysis of citizen science initiatives	All	Trees	All

3

4

5

6 Bibliography

7

- 8 Beyene, T. J., Asfaw, F., Getachew, Y., Tufa, T. B., Collins, I., Beyi, A. F., & Revie, C. W. (2018). A
9 smartphone-based application improves the accuracy, completeness, and timeliness of cattle
10 disease reporting and surveillance in Ethiopia. *Frontiers in Veterinary Science*, 5(JAN).
11 doi:10.3389/fvets.2018.00002
- 12 Bradfield, R., Wright, G., Burt, G., Cairns, G., & Van Der Heijden, K. (2005). The origins and
13 evolution of scenario techniques in long range business planning. *Futures*, 37(8), 795-812.
14 doi:<https://doi.org/10.1016/j.futures.2005.01.003>
- 15 Bronner, A., Hénaux, V., Fortané, N., Hendriks, P., & Calavas, D. (2014). Why do farmers and
16 veterinarians not report all bovine abortions, as requested by the clinical brucellosis
17 surveillance system in France? *BMC Vet Res*, 10(1), 93. doi:10.1186/1746-6148-10-93
- 18 Bronner, A., Morignat, E., & Calavas, D. (2015). Respective influence of veterinarians and local
19 institutional stakeholders on the event-driven surveillance system for bovine brucellosis in
20 France. *BMC Vet Res*, 11(1). doi:10.1186/s12917-015-0499-1
- 21 Bruijnjs, M., Hogeveen, H., Garforth, C., & Stassen, E. (2013). Dairy farmers' attitudes and intentions
22 towards improving dairy cow foot health. *Livestock Science*, 155(1), 103-113.
23 doi:<http://dx.doi.org/10.1016/j.livsci.2013.04.005>
- 24 Burns, T. E., Ribble, C., McLaws, M., Kelton, D., & Stephen, C. (2013). Perspectives of an
25 underrepresented stakeholder group, backyard flock owners, on poultry health and avian
26 influenza control. *Journal of Risk Research*, 16(2), 245-260.
27 doi:10.1080/13669877.2012.726244
- 28 Burton, R. J. F. (2004). Reconceptualising the 'behavioural approach' in agricultural studies: a socio-
29 psychological perspective. *Journal of Rural Studies*, 20(3), 359-371.
30 doi:<http://dx.doi.org/10.1016/j.jrurstud.2003.12.001>
- 31 Coughlin, S. S. (1990). Recall bias in epidemiologic studies. *Journal of Clinical Epidemiology*, 43(1),
32 87-91. doi:[https://doi.org/10.1016/0895-4356\(90\)90060-3](https://doi.org/10.1016/0895-4356(90)90060-3)
- 33 Creswell, J. W., & Clark, V. L. P. (2017). *Designing and conducting mixed methods research*: Sage
34 publications.
- 35 Cross, P., St. John, F. A. V., Khan, S., & Petroczi, A. (2013). Innovative Techniques for Estimating
36 Illegal Activities in a Human-Wildlife-Management Conflict. *PLoS ONE*, 8(1), e53681.
37 doi:10.1371/journal.pone.0053681
- 38 Crow, P., Perez-Sierra, A., Kavčič, A., Lewthwaite, K., Kolšek, M., Ogris, N., . . . de Groot, M.
39 (2020). Using Citizen science to monitor the spread of tree pests and diseases: Outcomes of
40 two projects in slovenia and the UK. *Management of Biological Invasions*, 11(4), 703-719.
41 doi:10.3391/mbi.2020.11.4.06
- 42 De Bruyn, S., Wouters, E., Ponnet, K., Tholen, R., Masquillier, C., Remmen, R., & Van Hal, G.
43 (2020). Off-label prescribing of stimulant medication to students: a qualitative study on the
44 general practitioner perspective. *Sociology of Health & Illness*, 42(7), 1657-1672.
45 doi:<https://doi.org/10.1111/1467-9566.13166>
- 46 Delgado, A. H., Norby, B., Dean, W. R., McIntosh, W. A., & Scott, H. M. (2012). Utilizing
47 qualitative methods in survey design: Examining Texas cattle producers' intent to participate
48 in foot-and-mouth disease detection and control. *Preventive Veterinary Medicine*, 103(2-3),
49 120-135. doi:<http://dx.doi.org/10.1016/j.prevetmed.2011.09.012>
- 50 Delgado, A. H., Norby, B., Scott, H. M., Dean, W., McIntosh, W. A., & Bush, E. (2014). Distribution
51 of cow-calf producers' beliefs about reporting cattle with clinical signs of foot-and-mouth
52 disease to a veterinarian before or during a hypothetical outbreak. *Preventive Veterinary
53 Medicine*, 117(3-4), 505-517. doi:<http://dx.doi.org/10.1016/j.prevetmed.2014.09.011>
- 54 Eckelkamp, E. A., & Bewley, J. M. (2020). On-farm use of disease alerts generated by precision dairy
55 technology. *Journal of Dairy Science*, 103(2), 1566-1582. doi:10.3168/jds.2019-16888
- 56 Elbers, A. R. W., Gorgievski-Duijvesteijn, M. J., van der Velden, P. G., Loeffen, W. L. A., &
57 Zarafshani, K. (2010). A socio-psychological investigation into limitations and incentives
58 concerning reporting a clinically suspect situation aimed at improving early detection of

- 59 classical swine fever outbreaks. *Vet Microbiol*, 142(1), 108-118.
60 doi:<https://doi.org/10.1016/j.vetmic.2009.09.051>
- 61 Elbers, A. R. W., Gorgievski-Duijvesteijn, M. J., Zarafshani, K., & Koch, G. (2010). To report or not
62 to report: A psychosocial investigation aimed at improving early detection of avian influenza
63 outbreaks. *OIE Revue Scientifique et Technique*, 29(3), 435-449.
- 64 Fielding, N. G. (2012). Triangulation and Mixed Methods Designs: Data Integration With New
65 Research Technologies. *Journal of Mixed Methods Research*, 6(2), 124-136.
66 doi:10.1177/1558689812437101
- 67 Fox, M., Christley, R., Lupo, C., Moore, H., Service, M., & Campbell, K. (2020). Preventing and
68 mitigating farmed bivalve disease: a Northern Ireland case study. *Aquaculture International*,
69 28(6), 2397-2417. doi:10.1007/s10499-020-00597-y
- 70 Garza, M., Ågren, E. C. C., & Lindberg, A. (2020). Nudging in Animal Disease Control and
71 Surveillance: A Qualitative Approach to Identify Strategies Used to Improve Compliance
72 With Animal Health Policies. *Frontiers in Veterinary Science*, 7, 383.
- 73 Gates, M. C., Earl, L., & Enticott, G. (2021). Factors influencing the performance of voluntary farmer
74 disease reporting in passive surveillance systems: a scoping review. *Preventive Veterinary
75 Medicine*, 105487. doi:<https://doi.org/10.1016/j.prevetmed.2021.105487>
- 76 Gilbert, W. H., Häsler, B. N., & Rushton, J. (2014). Influences of farmer and veterinarian behaviour
77 on emerging disease surveillance in England and Wales. *Epidemiology and Infection*, 142(1),
78 172-186. doi:10.1017/S0950268813000484
- 79 Gould, D. (1996). Using vignettes to collect data for nursing research studies: how valid are the
80 findings? *Journal of Clinical Nursing*, 5(4), 207-212. doi:[https://doi.org/10.1111/j.1365-
81 2702.1996.tb00253.x](https://doi.org/10.1111/j.1365-2702.1996.tb00253.x)
- 82 Greenhalgh, T., & Peacock, R. (2005). Effectiveness and efficiency of search methods in systematic
83 reviews of complex evidence: audit of primary sources. *BMJ (Clinical research ed.)*,
84 331(7524), 1064-1065. doi:10.1136/bmj.38636.593461.68
- 85 Guinat, C., Wall, B., Dixon, L., & Pfeiffer, D. U. (2016). English pig farmers' knowledge and
86 behaviour towards African swine fever suspicion and reporting. *PLoS ONE*, 11(9).
87 doi:10.1371/journal.pone.0161431
- 88 Hall, C., Raum, S., Morris, J., & O'Brien, L. (2017). *Observatree: Key lessons Qualitative study of the
89 'Observatree' citizen science project.*
90 https://www.forestresearch.gov.uk/documents/7252/Observatree_evaluation_report.pdf.
- 91 Hamilton-Webb, A., Naylor, R., Little, R., & Maye, D. (2016). Compensation and exotic livestock
92 disease management: The views of animal keepers and veterinarians in England. *Veterinary
93 Record*, 179(20), 513. doi:10.1136/vr.103571
- 94 Hammersley, M., & Atkinson, P. (2019). *Ethnography : principles in practice* (Fourth edition. ed.).
95 London ;; Routledge.
- 96 Hammond, N. E. B., Hardie, D., Hauser, C. E., & Reid, S. A. (2016a). Can general surveillance detect
97 high priority pests in the Western Australian Grains Industry? *Crop Protection*, 79, 8-14.
98 doi:<https://doi.org/10.1016/j.cropro.2015.10.004>
- 99 Hammond, N. E. B., Hardie, D., Hauser, C. E., & Reid, S. A. (2016b). How would high priority pests
100 be reported in the Western Australian grains industry? *Crop Protection*, 79, 26-33.
101 doi:10.1016/j.cropro.2015.10.005
- 102 Hernández-Jover, M., Higgins, V., Bryant, M., Rast, L., & McShane, C. (2016). Biosecurity and the
103 management of emergency animal disease among commercial beef producers in New South
104 Wales and Queensland (Australia). *Preventive Veterinary Medicine*, 134, 92-102.
105 doi:10.1016/j.prevetmed.2016.10.005
- 106 Hoinville, L. J., Alban, L., Drewe, J. A., Gibbens, J. C., Gustafson, L., Häsler, B., . . . Stärk, K. D. C.
107 (2013). Proposed terms and concepts for describing and evaluating animal-health surveillance
108 systems. *Preventive Veterinary Medicine*, 112(1), 1-12.
109 doi:<https://doi.org/10.1016/j.prevetmed.2013.06.006>
- 110 Hughes, R., & Huby, M. (2002). The application of vignettes in social and nursing research. *Journal
111 of Advanced Nursing*, 37(4), 382-386. doi:<https://doi.org/10.1046/j.1365-2648.2002.02100.x>

- 112 Ivankova, N. V., Creswell, J. W., & Stick, S. L. (2006). Using Mixed-Methods Sequential
113 Explanatory Design: From Theory to Practice. *Field Methods*, 18(1), 3-20.
114 doi:10.1177/1525822X05282260
- 115 Khan, M. S., Durrance-Bagale, A., Mateus, A., Sultana, Z., Hasan, R., & Hanefeld, J. (2020). What
116 are the barriers to implementing national antimicrobial resistance action plans? A novel
117 mixed-methods policy analysis in Pakistan. *Health Policy and Planning*.
118 doi:10.1093/heapol/czaa065
- 119 Kollmuss, A., & Agyeman, J. (2002). Mind the Gap: Why do people act environmentally and what
120 are the barriers to pro-environmental behavior? *Environmental Education Research*, 8(3),
121 239-260. doi:10.1080/13504620220145401
- 122 Krumpal, I. (2013). Determinants of social desirability bias in sensitive surveys: a literature review.
123 *Quality & Quantity*, 47(4), 2025-2047. doi:10.1007/s11135-011-9640-9
- 124 Kuchler, F., & Hamm, S. (2000). Animal disease incidence and indemnity eradication programs.
125 *Agricultural Economics*, 22(3), 299-308. doi:[https://doi.org/10.1111/j.1574-](https://doi.org/10.1111/j.1574-0862.2000.tb00076.x)
126 [0862.2000.tb00076.x](https://doi.org/10.1111/j.1574-0862.2000.tb00076.x)
- 127 Lensvelt-Mulders, G. J. L. M., Hox, J. J., van der Heijden, P. G. M., & Maas, C. J. M. (2005). Meta-
128 Analysis of Randomized Response Research: Thirty-Five Years of Validation. *Sociological*
129 *Methods & Research*, 33(3), 319-348. doi:10.1177/0049124104268664
- 130 Limon, G., Lewis, E. G., Chang, Y. M., Ruiz, H., Balanza, M. E., & Guitian, J. (2014). Using mixed
131 methods to investigate factors influencing reporting of livestock diseases: A case study
132 among smallholders in Bolivia. *Preventive Veterinary Medicine*, 113(2), 185-196.
133 doi:10.1016/j.prevetmed.2013.11.004
- 134 Lupo, C., Osta Amigo, A., Mandard, Y. V., Peroz, C., & Renault, T. (2014). Improving early
135 detection of exotic or emergent oyster diseases in France: Identifying factors associated with
136 shellfish farmer reporting behaviour of oyster mortality. *Preventive Veterinary Medicine*,
137 116(1-2), 168-182. doi:10.1016/j.prevetmed.2014.05.002
- 138 Maye, D., Enticott, G., & Naylor, R. (2017). Using scenario-based influence mapping to examine
139 farmers' biosecurity behaviour. *Land Use Policy*, 66, 265-277.
140 doi:<https://doi.org/10.1016/j.landusepol.2017.04.026>
- 141 Maye, D., Enticott, G., & Naylor, R. (2020). Theories of Change in Rural Policy Evaluation.
142 *Sociologia Ruralis*, 60(1), 198-221. doi:10.1111/soru.12269
- 143 Mazrier, H., Tal, S., Aizinbud, E., & Bargai, U. (2006). A field investigation of the use of the
144 pedometer for the early detection of lameness in cattle. *The Canadian veterinary journal = La*
145 *revue veterinaire canadienne*, 47(9), 883-886.
- 146 McFarland, L., Macken-Walsh, Á., Claydon, G., Casey, M., Douglass, A., McGrath, G., & McAloon,
147 C. G. (2020). Irish dairy farmers' engagement with animal health surveillance services:
148 Factors influencing sample submission. *Journal of Dairy Science*, 103(11), 10614-10627.
149 doi:10.3168/jds.2019-17889
- 150 Merrill, S. C., Koliba, C. J., Moegenburg, S. M., Zia, A., Parker, J., Sellnow, T., . . . Smith, J. M.
151 (2019). Decision-making in livestock biosecurity practices amidst environmental and social
152 uncertainty: Evidence from an experimental game. *PLoS ONE*, 14(4), e0214500.
153 doi:10.1371/journal.pone.0214500
- 154 Merrill, S. C., Moegenburg, S., Koliba, C. J., Zia, A., Trinity, L., Clark, E., . . . Smith, J. M. (2019).
155 Willingness to Comply With Biosecurity in Livestock Facilities: Evidence From
156 Experimental Simulations. *Frontiers in Veterinary Science*, 6, 156-156.
157 doi:10.3389/fvets.2019.00156
- 158 Oreszczyn, S., & Carr, S. (2008). Improving the link between policy research and practice: using a
159 scenario workshop as a qualitative research tool in the case of genetically modified crops.
160 *Qualitative Research*, 8(4), 473-497.
- 161 Oreszczyn, S., Lane, A., & Carr, S. (2010). The role of networks of practice and webs of influencers
162 on farmers' engagement with and learning about agricultural innovations. *Journal of Rural*
163 *Studies*, 26(4), 404-417. doi:<http://dx.doi.org/10.1016/j.jrurstud.2010.03.003>
- 164 Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., . . .
165 Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting
166 systematic reviews. *BMJ*, 372, n71. doi:10.1136/bmj.n71

- 167 Palmer, S., Fozdar, F., & Sully, M. (2009). The Effect of Trust on West Australian Farmers'
168 Responses to Infectious Livestock Diseases. *Sociologia Ruralis*, 49(4), 360-374.
169 doi:<https://doi.org/10.1111/j.1467-9523.2009.00495.x>
- 170 Palmer, S., Sully, M., & Fozdar, F. (2009). Farmers, Animal Disease Reporting and The Effect of
171 Trust: A Study of West Australian Sheep and Cattle Farmers. *Rural Society*, 19(1), 32-48.
172 doi:10.5172/rsj.351.19.1.32
- 173 Pawson, R., Greenhalgh, T., Harvey, G., & Walshe, K. (2005). Realist review - a new method of
174 systematic review designed for complex policy interventions. *Journal of health services*
175 *research & policy*, 10(1_suppl), 21-34. doi:10.1258/1355819054308530
- 176 Pawson, R., & Tilley, N. (2004). *Realist Evaluation*. London: Cabinet Office.
- 177 Pfeiffer, C. N. (2018). *Improving disease surveillance in Australia's sheep industries: investigations*
178 *of syndromic surveillance, farmer behaviour and sheep trade networks*. (PhD), University of
179 Melbourne. Retrieved from <http://hdl.handle.net/11343/219385>
- 180 Pham, H. T. T., Peyre, M., Trinh, T. Q., Nguyen, O. C., Vu, T. D., Rukkamsuk, T., & Antoine-
181 Moussiaux, N. (2017). Application of discrete choice experiment to assess farmers'
182 willingness to report swine diseases in the Red River Delta region, Vietnam. *Preventive*
183 *Veterinary Medicine*, 138, 28-36. doi:10.1016/j.prevetmed.2017.01.002
- 184 Phillips, C. (2020). The force of Varroa: Anticipatory experiences in beekeeping biosecurity. *Journal*
185 *of Rural Studies*, 76, 58-66. doi:10.1016/j.jrurstud.2020.04.002
- 186 Porth, E. F., Dandy, N., & Marzano, M. (2015). "My garden is the one with no trees:" Residential
187 Lived Experiences of the 2012 Asian Longhorn Beetle Eradication Programme in Kent,
188 England. *Human Ecology*, 43(5), 669-679. doi:10.1007/s10745-015-9788-3
- 189 Pramuwidyatama, M. G., Hogeveen, H., & Saatkamp, H. W. (2020). Understanding the Motivation of
190 Western Java Smallholder Broiler Farmers to Uptake Measures Against Highly Pathogenic
191 Avian Influenza (HPAI). *Frontiers in Veterinary Science*, 7. doi:10.3389/fvets.2020.00362
- 192 Quine, C. P., Barnett, J., Dobson, A. D., Marcu, A., Marzano, M., Moseley, D., . . . Uzzell, D. (2011).
193 Frameworks for risk communication and disease management: the case of Lyme disease and
194 countryside users. *Philosophical transactions of the Royal Society of London. Series B,*
195 *Biological sciences*, 366(1573), 2010-2022. doi:10.1098/rstb.2010.0397
- 196 Randrianantoandro, T. N., Kubota, S., & Kono, H. (2018). Farmers' knowledge and incentive in
197 reporting infectious animal disease: The case of african swine fever in madagascar. *Journal of*
198 *Animal and Plant Sciences*, 28(1), 270-279.
- 199 Rao, X., & Zhang, Y. (2020). Livestock insurance, moral hazard, and farmers' decisions: a field
200 experiment among hog farms in China. *Geneva Papers on Risk and Insurance: Issues and*
201 *Practice*, 45(1), 134-156. doi:10.1057/s41288-019-00151-9
- 202 Robinson, P. A., & Epperson, W. B. (2013). Farm animal practitioners' views on their use and
203 expectations of veterinary diagnostic laboratories. *Veterinary Record*, 172(19), 503-503.
204 doi:10.1136/vr.101366
- 205 Robinson, P. A., Epperson, W. B., Huston, C. L., Pace, L. W., Wills, R. W., & Cosby, A. G. (2012).
206 Factors influencing diagnostic sample submission by food animal veterinarians in
207 Mississippi. *Veterinaria Italiana*, 48(1), 31-39.
- 208 Rotz, S., Gravely, E., Mosby, I., Duncan, E., Finnis, E., Horgan, M., . . . Fraser, E. (2019). Automated
209 pastures and the digital divide: How agricultural technologies are shaping labour and rural
210 communities. *Journal of Rural Studies*, 68, 112-122.
211 doi:<https://doi.org/10.1016/j.jrurstud.2019.01.023>
- 212 Sawford, K., Vollman, A. R., & Stephen, C. (2012). A Focused Ethnographic Study of Sri Lankan
213 Government Field Veterinarians' Decision Making about Diagnostic Laboratory Submissions
214 and Perceptions of Surveillance. *PLoS ONE*, 7(10). doi:10.1371/journal.pone.0048035
- 215 Sawford, K., Vollman, A. R., & Stephen, C. (2013). A Focused Ethnographic Study of Alberta Cattle
216 Veterinarians' Decision Making about Diagnostic Laboratory Submissions and Perceptions of
217 Surveillance Programs. *PLoS ONE*, 8(5). doi:10.1371/journal.pone.0064811
- 218 Soleri, D., & Cleveland, D. A. (2005). Scenarios as a tool for eliciting and understanding farmers'
219 biological knowledge. *Field Methods*, 17(3), 283-301.
- 220 Struchen, R., Hadorn, D., Wohlfender, F., Balmer, S., Süptitz, S., Zinsstag, J., & Vial, F. (2016).
221 Experiences with a voluntary surveillance system for early detection of equine diseases in

222 Switzerland. *Epidemiology and Infection*, 144(9), 1830-1836.
223 doi:10.1017/S0950268816000091
224 van der Heijden, K. (1996). *Scenarios: The Art of Strategic Conversation*. Chichester: John Wiley.
225 World Organisation for Animal Health (OIE). (2021). Terrestrial Animal Health Code, 29th Ed. OIE,
226 Paris. Available at: [www.oie.int/en/international-standard-setting/terrestrial-code/access-](http://www.oie.int/en/international-standard-setting/terrestrial-code/access-online/)
227 [online/](http://www.oie.int/en/international-standard-setting/terrestrial-code/access-online/) (accessed on 06 September 2021).
228 Wright, B. K., Jorgensen, B. S., & Smith, L. D. G. (2018). Understanding the biosecurity monitoring
229 and reporting intentions of livestock producers: Identifying opportunities for behaviour
230 change. *Preventive Veterinary Medicine*, 157, 142-151. doi:10.1016/j.prevetmed.2018.07.007
231 Zhang, Y., Zhu, X., & Turvey, C. G. (2016). On the Impact of Agricultural Livestock Microinsurance
232 on Death-Loss, Production and Vaccine Use: Observations from a Quasi-Natural Experiment
233 in China. *The Geneva Papers on Risk and Insurance - Issues and Practice*, 41(2), 225-243.
234 doi:10.1057/gpp.2016.3
235