

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/146413/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Guerrero-Sanchez, Sergio, Burger, Richard, Shah Fitri Bin Rosli, Mohd, Bin Moh Natsyir, Nazrul, Bin Rahman, Roslee, Saiman, Silvester, Orozco Ter Wengel, Pablo and Goossens, Benoit 2021. Homing: a case-study on the spatial memory of the Asian water monitor lizard *Varanus salvator* in the Kinabatangan floodplain. *Herpetological Bulletin* 158 (4) 10.33256/hb158.3739

Publishers page: <http://dx.doi.org/10.33256/hb158.3739>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



**Homing: a case-study on the spatial memory of the Asian water monitor lizard *Varanus salvator* in the Kinabatangan floodplain**

**SERGIO GUERRERO-SANCHEZ<sup>1,2,\*</sup>, RICHARD BURGER<sup>1,2</sup>, MOHD SHAH FITRI BIN ROSLI<sup>2</sup>, NAZRUL BIN MOH NATSYIR<sup>2</sup>, ROSLEE BIN RAHMAN<sup>2</sup>, SILVESTER SAIMIN<sup>3</sup>, PABLO OROZCO-TERWENGEL<sup>1</sup>, BENOIT GOOSSENS<sup>1,2,3,4</sup>**

<sup>1</sup>Organisms and Environment Division, School of Biosciences, Cardiff University, Sir Martin Evans Building, Museum Avenue, Cardiff CF10 3AX, UK

<sup>2</sup>Danau Girang Field Centre, c/o Sabah Wildlife Department, Wisma Muis, 88100 Kota Kinabalu, Sabah, Malaysia

<sup>3</sup>Sabah Wildlife Department, Wisma Muis, 88100 Kota Kinabalu, Sabah, Malaysia


<sup>4</sup>Sustainable Places Research Institute, Cardiff University, 33 Park Place, Cardiff CF10 3BA, UK

\*Corresponding author e-mail: [ekio0474@gmail.com](mailto:ekio0474@gmail.com)

Translocation may be used to improve the biological health of animal populations (Wolf et al., 1996) or to mitigate the impact of human-wildlife conflicts (Fisher & Lindenmayer, 2000). However, wildlife species may respond differently to translocations; they can show a tendency to either travel long distances in an attempt to return to their original location (homing), or to exhibit larger home ranges compared to resident individuals (Bradley, 2005; Wolf, et al., 2009). Homing behaviour is negatively correlated with translocation distance (Bowman et al., 2002; Villaseñor et al., 2013), and it is associated with several factors such as the identification of landscape landmarks and resource availability in the original home range (Powell and Mitchel, 2012). Thus, understanding the spatial memory and navigation skills of an organism can be fundamental to predict the success of management actions, such as translocations. This report describes the response of an Asian water monitor (*Varanus salvator*) translocated within the Kinabatangan floodplain in Sabah (Malaysian Borneo) and its return journey to its home territory.

In February 2018, a message was received that a monitor lizard, GPS-tagged as part of a long-term telemetry study, had been feeding on poultry in an oil palm plantation estate (Hillco, Felda Global Ventures Sdn. Bhd.; N5° 25'02" N, E118° 01'46" E). The 17 kg individual (presumably male) was subsequently translocated to a forested area (Lot 6 of the Lower Kinabatangan Wildlife Sanctuary; 5° 24'05" N, 118° 04'27" E), 5.27 km away from its original home range, which had been previously estimated using 2472 locations over 299


tracking days with a fix success rate of 75% (Guerrero-Sanchez et al., unpublished data). A new GPS tracker (Advanced Telemetry Systems Inc., North Isanti, MN, USA) was deployed in order to monitor its adaptations to the new environment (Fig. 1).



**Figure 1.** The Asian water monitor with GPS tracker navigating the Kinabatangan river, Sabah, Malaysian Borneo

The new tracker was set to record one GPS location every 90 minutes during daytime; night time was not recorded as water monitors are not active nocturnally. The lizard was tracked for 11 weeks post-translocation, collecting a total of 621 GPS locations. The data show that the lizard took about seven weeks to return to its original home range, but instead of traveling in a straight line or following the river, it travelled through the forest by way of three different plantation “spots” (Fig 2). The last two of those spots were on the same side of the river as the home range, with which there was contiguous plantation habitat, but to reach the home range the monitor instead went through the forest and made further river crossings. This route may have been chosen as unpublished data suggest that forested areas offer more protection to the monitor lizards than oil palm plantations and have prey in equal abundance.





**Figure 2.** Weekly occupied areas of the translocated monitor lizard in the Kinabatangan floodplain. Polygons represent a 95% home range (kernel density estimate; KDE). Release spot is within the “week 1” polygon. Arrows show the flow of the lizard’s movements and the orange star marks the location of the chicken house. KDE was fitted using the package Animove HR for R.

Homing behaviour is not rare in reptiles (Read et al., 2007; Pittman et al., 2014); it has been reported that the saltwater crocodile (*Crocodylus porosus*) can travel up to 400 km back to its original home range after being translocated (Read et al., 2007). Burmese pythons (*Python bivittatus*) possess a well-developed bearing ability that allow translocated individuals to head back home without the need to follow straight lines (Pittman et al., 2014). The natal habitat preference induction theory suggests that when translocated individuals of certain species are looking for a new home they search for habitat attributes similar to those encountered early in life (Davis and Stamps, 2004). Furthermore, the length of time a released individual spends at a release site can be informative about its acceptance or rejection of a new home, while the overall distance travelled during its return can indicate the degree of preference for the special features of its original habitat (Hayward et al., 2007). The time taken by the lizard in this study to return to its original home range, as well as the time

spent in certain key areas, (i.e. a different location of an oil palm plantation), suggest that this particular individual was willing to look for a suitable ‘new home’ with similar features to its original one, but ended up rejecting these areas, possibly due to the presence of other individuals, or unsuitable environmental features (i.e. prey and shelter availability, intense human activity). This report suggests two main drivers influencing the lizard’s behaviour: (1) the well-identified habitat of its original home range as a source of predictable food resources and safety, and (2) the discontinuous distribution of these features within the landscape, forcing this individual to avoid these areas and keep moving towards its original home range. We cannot discard the role of the navigational ability and spatial memory that might help the lizard to locate itself within the landscape and find the safest route to his original range (Pittman et al., 2014).

Although it is unclear whether monitors exhibit strictly territorial behaviour (Pascoe et al., 2019), antagonism is likely to occur between males, not only as territorial defence but also as competition for both food and reproductive females (Pascoe et al., 2019). Interestingly, after the return of the lizard to its original home range it was tracked for four more weeks and the data show that it remained within the boundaries of its home range. This behaviour suggests that the lizard not only recognized its home but also that probably no other large individual occupied it during its absence. Hence, what we witnessed could be part of a territorial behaviour, which should be taken into consideration in further studies of human-monitor lizard conflict mitigation.

The water monitor’s knowledge of the most relevant elements in its original home range, such as absence of other lizards and the features associated with food and cover, might work as a stimulus for its return to its original home. All these findings suggest that the species may have a well-developed spatial memory, as well as a strong attachment to the well-known features of its home range. These characteristics should be considered in areas where there are human-lizard conflicts and whenever translocated lizards are moved to areas already abundant in monitor lizards. The presence of large monitors in these selected areas can have a counterproductive effect if they result in translocated individuals fleeing and returning to their original homes. In order to get a better understanding of territoriality and habitat preferences of monitor lizards, we recommend carrying out long-term experiments on translocations, using GPS telemetry and considering treatments with varying translocation

distance and varying habitat similarities, especially for areas where human-lizard interactions are a burden.

## ACKNOWLEDGEMENTS

This study was financially supported by Danau Girang Field Centre and Cardiff University. SGS was supported by the National Council for Science and Technology (CONACYT; Mexico; scholarship No. 235294). Safety and animal welfare protocols were reviewed and approved by Sabah Wildlife Department and the Sabah Biodiversity Centre (permit JKM/MBS.1000-2/2 JLD.3-7). The authors would like to thank students and staff members of the Danau Girang Field Centre for their support with the fieldwork, as well as the Director of Sabah Wildlife Department for his support on this project, and Felda Global Holdings Sdn. Bhd. for granting permission to conduct part of the fieldwork within their plantation estates. We would also like to thank our colleagues Rudi Delvaux, for the two photos in Figure 1, and Liesbeth Frias, for her comments on early versions of this manuscript and proofreading.

## REFERENCES

- Bowman, J., Jaeger, J.A.G. & Fahrig, L. (2002). Dispersal distance of mammals is proportional to home range size. *Ecology* 83(7): 2049-2055. [doi.org/10.1890/0012-9658\(2002\)083\[2049:DDOMIP\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[2049:DDOMIP]2.0.CO;2).
- Bradley, E.H., Pletscher, D.H., Bangs, E.E., Kunkel, K.E., Smith, D.W., Mack, C.M., Meier, T.J., Fontaine, J.A., Niemeyer, C.C. & Jimenez, M.D. (2005). Evaluating wolf translocation as a nonlethal method to reduce livestock conflicts in the northwestern United States. *Conservation Biology* 19(5): 1498-1508. [doi.org/10.1111/j.1523-1739.2005.00102.x](https://doi.org/10.1111/j.1523-1739.2005.00102.x).
- Davis, J.M. & Stamps, J.A. (2004). The effect of natal experience on habitat preferences. *Trends in Ecology and Evolution*. 19(8): 411-416. [doi.org/10.1016/j.tree.2004.04.006](https://doi.org/10.1016/j.tree.2004.04.006).
- Fischer, J. & Lindenmayer, D.B. (2000). An assessment of the published results of animal relocations. *Biological Conservation* 96(1): 1-11. [doi:10.1016/S0006-3207\(00\)00048-3](https://doi.org/10.1016/S0006-3207(00)00048-3).
- Hayward, M.W., Kerley, G.I.H., Adendorff, J., Moolman, L., O'Brien, J., Sholto-Douglas, A., Bisset, C., Bean, P., Fogarty, A., Howarth, D. & Slater, R. (2007). The reintroduction

127 of large carnivores to the Eastern Cape, South Africa: an assessment. *Oryx* 41(2): 205-  
 128 214. [doi.org/10.1017/S0030605307001767](https://doi.org/10.1017/S0030605307001767).

129 Pascoe, J.H., Flesh, J.S., Duncan, M.G., Pla, M.L. & Mulley, R.C. (2019). Territoriality and  
 130 seasonality in the home range of adult male free-ranging Lace monitor (*Varanus varius*)  
 131 in South-eastern Australia. *Herpetological Conservation and Biology* 14(1): 97-104.  
 132 [www.herpconbio.org/Volume\\_14/Issue\\_1/Pascoe\\_etal\\_2019.pdf](http://www.herpconbio.org/Volume_14/Issue_1/Pascoe_etal_2019.pdf).

133 Pitman, S.E., Hart, K.M., Cherkiss, M.S., Snow, R.W., Fujisaki, I., Smith, B., Mazzotti, F.J.,  
 134 & Dorcas, M.E. (2014). Homing of invasive Burmese pythons in South Florida: evidence  
 135 for map and compass senses in snakes. *Biology Letters* 10: 20140040.  
 136 [doi.org/10.1098/rsbl.2014.0040](https://doi.org/10.1098/rsbl.2014.0040).

137 Powell, R.A. & Mitchel, I.M.S. (2012). What is a home range? *Journal of Mammalogy* 93(4):  
 138 948-958. [doi.org/10.1644/11-MAMM-S-177.1](https://doi.org/10.1644/11-MAMM-S-177.1).

139 Read, M.A., Grigg, G.C., Irwin, S.R., Shanahan D. & Franklin, C.E. (2007). Satellite tracking  
 140 reveals long distance coastal travel and homing by translocated estuarine crocodiles,  
 141 *Crocodylus porosus*. *PLoS ONE* 2(9): e949. [doi:10.1371/journal.pone.0000949](https://doi.org/10.1371/journal.pone.0000949).

142 Stamps, J.A. & Swaisgood, R.R. (2007). Someplace like home: experience, habitat selection  
 143 and conservation biology. *Applied Animal Behaviour Science* 102(3-4): 392-409.  
 144 [doi.org/10.1016/j.applanim.2006.05.038](https://doi.org/10.1016/j.applanim.2006.05.038).

145 Villaseñor, N.R., Escobar, M.A.H. & Estades, C.F. (2013). There is no place like home: High  
 146 homing rate and increased mortality after translocation of a small mammal. *European*  
 147 *Journal of Wildlife Research* 59: 749-760. [doi.org/10.1007/s10344-013-0730-y](https://doi.org/10.1007/s10344-013-0730-y).

148 Wolf, C.M., Griffith, B., Reed, C. & Temple, S.A. (1996). Avian and mammalian  
 149 translocations: Update and reanalysis of 1987 survey data. *Conservation Biology* 10(4):  
 150 1142-1154. [doi.org/10.1046/j.1523-1739.1996.10041142.x](https://doi.org/10.1046/j.1523-1739.1996.10041142.x).

151 Wolf, M., Frair, J., Merrill, E. & Turchin, P. (2009). The attraction of the known: the  
 152 importance of spatial familiarity in habitat selection in wapiti *Cervus elaphus*. *Ecography*  
 153 32: 401-410. [doi.org/10.1111/j.1600-0587.2008.05626.x](https://doi.org/10.1111/j.1600-0587.2008.05626.x).