

Paternal Postnatal Depression Among Fathers of Newborn in Saudi Arabia

American Journal of Men's Health
January-February 2019: 1–12
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1557988319831219
journals.sagepub.com/home/jmh



Naila A. Shaheen, MD^{1,2,3} , **Yousra AlAtiq, MPhil^{2,3,4},**
Abin Thomas, MSc^{1,2,3}, **Hussam Ali Alanazi, MD⁵,**
Ziad E. AlZahrani, MHA^{2,3,6}, **Samah Abdul Ra'ouf Younis, BDS^{2,3,6},**
and Mohamed A. Hussein, PhD^{1,2,3}

Abstract

Paternal postnatal depression (PPND) is not a commonly recognized phenomenon. The aim of the study was to identify the Edinburgh Postnatal Depression Scale (EPDS) cutoff for Saudi fathers, to estimate PPND prevalence and to determine the risk factors of PPND among fathers of newborn in Saudi Arabia. A cross-sectional study of fathers with babies born up to 6 months prior to the survey was conducted. Fathers were screened using EPDS and demographic questionnaire. The fathers were selected using systematic random sampling from visitors to the birth registration office. A subsample of participants from the postnatal wards in a tertiary care was invited for additional evaluation by a psychologist using the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)* criteria for major depressive disorders. Receiver operating characteristic curve was utilized to identify fathers with depression; a cutoff of 8/9 was optimal to achieve sensitivity 77.8% and specificity 81.3%. Adjusted prevalence of PPND was reported with corresponding Wilson 95% confidence interval. Two hundred and ninety fathers completed the EPDS and demographic questionnaire. Of 72 invited participants, 57 (79.16%) attended the diagnostic interview. The average age of fathers was 34.97 ± 8.56 years, the average maternal age was 29.18 ± 7.41 years, average age of the newborn was 43.13 ± 35.88 days. PPND adjusted prevalence was 16.6% (95% CI [8.5, 25.6]). Paternal mental health needs equal attention during and postdelivery of newborn. Fathers should receive perinatal and postnatal mental health assessment to prevent behavioral problems in their children and disruption of relationship with their spouse.

Keywords

paternal depression, Edinburgh Postnatal Depression Scale, postpartum depression, paternal postnatal depression, fathers, validation

Received June 5, 2018; revised January 6, 2019; accepted January 14, 2019

Depression is a leading contributor to global disease burden: over 300 million people worldwide were estimated to suffer from the condition in 2015 (World Health Organization, 2017). The prevalence of depression in the United States has doubled between 2005 and 2015 (Weinberger et al., 2018). Despite a plethora of available treatments, depression continues to be a burden because of the low detection rate. Detection is even more challenging in unsuspected groups, such as men, due to the stigma attached to the condition by society. Men suppress their feelings and are less likely to acknowledge depression. Men can exhibit symptoms that are less characteristic of depression, including irritability, anger, and aggression (O'Hara & McCabe, 2013). Research on

parental mental health after childbirth focuses mainly on the adjustment of women during the transition to parenthood (Edward, Castle, Mills, Davis, & Casey, 2015).

Postpartum depression (PPD) is defined as “a non-psychotic depressive episode of mild to moderate severity, beginning in or extending into the first postnatal year” (Miller, 2002). PPD among women has been thoroughly investigated by researchers for over 50 years (Brockington, 2004). Conversely, depression among the fathers of newborn, termed paternal postnatal depression (PPND) or “sad dads,” is a phenomenon recognized infrequently, is underscreened and underdiagnosed (Brockington, 2004; Musser, Ahmed, Foli, & Coddington, 2013). The concept of PPND came to the fore in 1990 (Brockington, 2004),



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

and has gained significant attention in recent years (Condon, Corkindale, & Boyce, 2004).

To date, two reviews had reported the estimated rate of depression among fathers of newborn. A meta-analysis in 2010 has reported PPND estimated rate as 10% (95% CI [8.5%, 12.7%]) with a mix of studies reporting depression rates at single versus two or more time points during the antenatal/postpartum period (Paulson & Bazemore, 2010). A more recent meta-analysis in 2016 reported the estimated rate of PPND 8.4% (95% CI [7.2%, 9.6%]; Cameron, Sedov, & Tomfohr-Madsen, 2016). Studies have shown that PPND estimated prevalence tends to decline with time during the postpartum period (Goodman, 2004). Prevalence of PPD among both partners has been reported as 19.6% at 4 weeks postpartum, which decreased to 4.7% at 8 weeks postpartum (Paulson & Bazemore, 2010).

In spite of a series of studies conducted throughout Europe and the East between 2001 and 2015 to validate a screening tool for fathers of newborn (Edmondson, Psychogiou, Vlachos, Netsi, & Ramchandani, 2010; Lai, Tang, Lee, Yip, & Chung, 2010; Loscalzo, Giannini, Contena, Gori, & Benvenuti, 2015; Massoudi, Hwang, & Wickberg, 2013; Matthey, Barnett, Kavanagh, & Howie, 2001; Tran, Tran, & Fisher, 2012), a recent insight considered only the mothers of newborn at high risk of developing depression (Reynolds & Patel, 2017).

Several risk factors have been associated with PPND, such as unemployment, unplanned pregnancy, poor marital relationship, poor social support, lack of family support, previous history of depression, presence of depression during antenatal period, maternal depression, joint family versus nuclear family (Condon et al., 2004; Nishimura & Ohashi, 2010; Paulson & Bazemore, 2010). Studies have reported an association between PPND and the development of emotional and behavioral psychiatric disorders among their children (Pinheiro et al., 2006; Ramchandani et al., 2008). PPND needs to be recognized as a serious public health issue. It is deemed necessary to identify fathers with PPND in order to prevent developing childhood emotional and behavioral problems (Ramchandani, Stein, Evans, & O'Connor, 2005).

The reported percent of total years lived with disability (YLD) in Saudi Arabia is 9.5% due to depression (World Health Organization, 2017). The prevalence of maternal postpartum depression (MPPD) among Saudi women is 14% (Al-Modayfer, Alatiq, Khair, & Abdelkawi, 2015) compared to internationally reported figures of 4.5%–20% (O'Hara & McCabe, 2013). However, PPND among fathers of a newborn baby is relatively a new concept in Middle East and Saudi Arabia. A study conducted in an Arab country; Egypt, had reported 31.8% prevalence of depression in expecting fathers (Moussa et al., 2012) compared to reported 13.3% estimated rate in Canada among first-time expecting fathers (Da Costa et al., 2017). To our knowledge, there is no current estimate for the prevalence of PPND in Saudi Arabia. Furthermore, estimates provided from other published studies might have limited generalizability to the Saudi population due to the difference in cultural and social values.

The shifting of home responsibilities, with an increase in working women and greater involvement of fathers in childcare, puts men at increased risk of developing depression. Fathers of a newborn baby also experience changes in life, like mothers (Kim & Swain, 2007). Simultaneously, men are approximately 50% less likely to seek mental health treatment than women (Gonzalez, Alegria, & Prihoda, 2005). The cost on services utilized by fathers diagnosed with PPND was (£1,103.51), higher than those without depression (£945.03; Edeka, Petrou, & Ramchandani, 2011), that emphasizes the need for early detection and treatment of PPND. The current study addresses the limitations of the literature by identifying the Edinburgh Postnatal Depression Scale (EPDS) cutoff for Saudi fathers, to estimate the prevalence of depression among fathers of newborn and to determine the risk factors for PPND.

Methods

Study Design and Participants

This is a cross-sectional study conducted among Saudi fathers with babies born up to 6 months before survey administration during 2016. The study sample included 290

¹Department of Biostatistics and Bioinformatics, King Abdullah International Medical Research Center, Riyadh, Kingdom of Saudi Arabia

²King Saud bin Abdulaziz University for Health Sciences, Riyadh, Kingdom of Saudi Arabia

³Ministry of National Guard Health Affairs, Riyadh, Kingdom of Saudi Arabia

⁴Department of Psychiatry, King Abdulaziz Medical City, Riyadh, Kingdom of Saudi Arabia

⁵University of Hail, Kingdom of Saudi Arabia

⁶Research Office/Research Quality Management Section, King Abdullah International Medical Research Center, Riyadh, Kingdom of Saudi Arabia

Corresponding Author:

Naila A. Shaheen, MD, Department of Biostatistics and Bioinformatics, King Abdullah International Medical Research Center, King Saud bin Abdulaziz University for Health Sciences, Ministry of National Guard Health Affairs, P.O. Box 22490, Riyadh 11426, Kingdom of Saudi Arabia. Emails: drnaila@hotmail.com/ashrafna@ngha.med.sa

fathers who could read and write. Arabic was the native language of all participants. All participants had filled a self-filled survey that included a validated screening tool (EPDS) for identifying depression and set of questions composed of: demographics (six items), employment-related factors (five items), family-related factors (six items), wife-related factors (four items), pregnancy-related factors (seven items), psychological well-being factors (three items), and social support factors (two items). Out of 290, a subsample of 72 (25%) fathers was invited in a tertiary care center's postnatal ward for diagnostic interviews by a psychologist. The information obtained from the subsample was used to determine the EPDS optimal score for the Saudi population as well as used to correct for the apparent prevalence obtained from the screening tool.

Participants were selected using systematic sampling based on the rate of visitation from those attending postnatal wards at a tertiary care hospital or the birth registration office in Riyadh, Saudi Arabia (where all fathers are required to visit within 30 days of childbirth to obtain their child's birth certificate). Participants filling only the demographic and EPDS questionnaire were verbally consented at the introduction of the survey. The questionnaire was distributed by the study team. The completed questionnaires were collected immediately in an envelope to maintain the confidentiality of the responses. Ethical approval was obtained from the King Abdullah International Medical Research Center Institutional Review Board with the approval number (RC13/206/R).

Instrument

The EPDS is a 10-item self-rating scale designed to screen for depression among mothers (Cox, Holden, & Sagovsky, 1987). The EPDS has been demonstrated to be a valid tool to identify depression among fathers (Edmondson et al., 2010; Lai et al., 2010; Loscalzo et al., 2015; Massoudi et al., 2013; Matthey et al., 2001; Tran et al., 2012). The fathers of newborn were screened using an Arabic version of the EPDS translated by Ghubash, Abou-Saleh, and Daradkeh (1997). The scale identifies symptoms over the preceding 7 days. Each item has four possible responses, which are scored from 0 to 3; the total score ranges from 0 to 30. Fathers were excluded if they had a history of psychiatric illness, received medication for the psychiatric illness, if their newborn were older than 6 months, or if they missed out any of the EPDS responses.

Diagnostic Interview

Out of 290 participants, 72 (24.8%) participants who completed the demographic and EPDS questionnaire were immediately invited to participate in the diagnostic interview and undergo evaluation by a psychologist using

the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)* criteria for major depressive disorders. Fathers who agreed to participate in the diagnostic interview were enrolled after providing written consent. The psychologist was blinded to their EPDS responses. The participants who were identified with major depression by the psychologist were advised to consult their primary care physician for further evaluation and assessment. The *DSM-5* scoring system was considered the gold standard.

The potential risk factors of depression among fathers of newborn were identified as demographics, employment related, family support, family set-up (joint vs. nuclear family), social support, pregnancy related, family history of depression, wife having postpartum depression, and psychological well-being (feeling isolated from the partner, whether had received overspeeding ticket postdelivery) correlates.

Statistical Analysis

Quantitative variables, including parents' age, child's age, and the number of children living in the house, were summarized and reported in terms of means, standard deviation, median, and interquartile range (IQR). Categorical variables, such as educational level, occupation, income level, family history of psychiatric illness, and information related to pregnancy, spouse, and family, were reported in terms of frequency tables and percentages. All variables were compared between fathers with and without depression using the Chi-square test of independence and Wilcoxon rank sum test. Statistical tests were declared significant at an α -level of less than .05. The cutoff value for the EPDS score was determined using a receiver operating characteristic (ROC) curve. Participants with a total EPDS score of ≥ 9 were considered to have depression. The prevalence of depression among fathers was estimated by dividing the number of fathers identified as depressed based on an EPDS cutoff of 9 by the total number of fathers in the study. The proportion was then corrected for the estimated sensitivity and specificity of the EPDS tool in the Saudi population. The apparent and adjusted prevalence was reported in terms of point estimate and corresponding Wilson 95% confidence intervals. Logistic regression was used to explore the risk factors for depression. The results were reported as odds ratios, 95% confidence intervals, and p values. Analyses were performed using SAS version 9.4 (SAS Institute, Cary, NC, USA).

Results

Study Sample

A total of 347 fathers were invited to participate in the study. Of them, 57 (16.42%) were excluded because they had missed some EPDS responses or had a history of

psychiatric illness. Thus, 290 fathers completed the EPDS survey. A convenient sample of 72 (24.8%) fathers were invited to undergo a structured interview with a psychologist, of whom 57 (79.16%) consented and agreed. The 15 (20.83%) who did not agree to participate in the diagnostic interview were not statistically different (in terms of demographics and other factors) from the 57 (79.17%) who had participated.

Sociodemographic Characteristics

Of the 290 fathers enrolled in the study, the mean reported paternal age was 34.97 ($SD = 8.56$) years, maternal age was 29.18 ($SD = 7.41$) years, and the mean reported age of the newborn was 43.13 ($SD = 35.88$) days (Table 1). Out of 290 reported, 67 (23.04%) were first-time fathers, whereas 184 (63.23%) were experienced fathers. Two hundred and seventy one (93.12%) fathers had one newborn. Sixty-seven (23.02%) fathers were in their twenties, the majority 147 (50.51%) were in their thirties, 44 (15.12%) were in their forties, and 13 (4.46%) were in their fifties or older.

One hundred and forty-one (50.9%) fathers reported that they had received higher education. Of 259 fathers who had an occupation, 92 (35.52%) were employed in the armed forces, 53 (20.46%) were professionals (doctors/engineers), 36 (13.9%) were managers, and 36 (13.9%) were clerical support workers, 21 (8.11%) were associate professionals, and 21 (8.11%) were sales professionals. With regard to total income, 74 (27.41%) fathers reported a total monthly income of Saudi-Arabian Riyal (SAR) $\geq 15,000$. The majority of the fathers 263 (94.27%) were employed; 218 (83.2%) had full-time employment, and 187 (70.57%) had a permanent job. Seventy-eight (28.36%) fathers owned their own home. The median number of children living in each household was 3 (IQR = 3). Demographic/employment-related characteristics were not significantly different between fathers with and without depression. However, the employment status, permanent versus temporary, was significantly different between fathers with and without depression ($\chi^2 = 5.103, p = .024$; Table 1).

Prevalence of Depression Among Fathers of Newborn

Of the 290 participants, 98 (27.9%; 95% CI [23.1, 33.4]) appeared to be depressed according to the EPDS cutoff score of 8/9. The estimated prevalence was corrected for tool sensitivity (77.8%) and specificity (81.3%), and the final estimated prevalence was reduced to 16.6%, with a 95% confidence limit (Wilson score) of 8.5%–25.6%.

Figure 1 summarizes the unadjusted prevalence at different cutoffs reported in the literature. The prevalence increases or decreases by using the cutoffs determined for other samples.

Marital, Wife, Family Support, and Psychological Well being Related Correlates of Depression

Several factors related to marital relationship, family, and social support were examined among the fathers. Most of the fathers, 276 (98.92%), were married and living with their wives. Almost all fathers, 274 (98.56%), reported a good marital relationship, and the majority, 261 (92.55%), had just one wife. Only 25 (9.03%) reported that their wives were depressed. One hundred and fifty-five (55.96%) fathers reported that their wife went to her mother's house for delivery, whereas 102 (52.31%) fathers reported that their wife was joined by her mother at their home. The family setup was reported as nuclear by 228 (81.14%) of the fathers. Almost all fathers, 280 (99.64%), reported a good relationship with their parents, and 244 (86.83%) received the support of friends. The marital-, wife-, and family-related characteristics were not different between fathers with and without depression.

Twenty-four (8.66%) fathers reported that they were spending more time at work to disconnect from home and family; however, this was not significantly different between fathers with and without depression ($\chi^2 = 0.025, p = .875$). Speeding ticket was received by 47 (16.97%) of the fathers after their newborn's birth. Thirty-six (13%) fathers felt isolated and disconnected from their partner. The number of fathers who reported feeling isolated and disconnected from their partners was significantly different between fathers with and without depression ($\chi^2 = 6.322, p = .012$; Table 2).

Pregnancy-Related Correlates of Depression

Almost half of the fathers, 159 (56.9%), reported that the pregnancy was planned. Only 12 (4.3%) had conceived using in vitro fertilization (IVF). Almost all fathers, 262 (93.91%), had attended antenatal checkups with their wives, and 224 (80.58%) had attended their newborn's delivery. The mode of delivery was reported as vaginal by 191 (67.97%) fathers, and cesarean section by 91 (34.73%) fathers. Thirty-one (11.07%) fathers reported complications during childbirth. Forty-four (15.83%) fathers had lost a child before. Twenty-two (8.27%) fathers reported that their newborn had been diagnosed with congenital birth defects/syndromes. Variables related to pregnancy were not significantly different between fathers with and without depression (Table 2).

ROC Curve Analysis

The results of the ROC curve analysis are summarized in Table 3 and Figure 2. The area under the curve was 0.81. A score of 8/9 was identified as the optimal cutoff for

Table 1. Demographic/Employment-Related Characteristics of Fathers With/Without Depression.

Variables <i>n</i> = response rate	Overall <i>n</i> = 290	Depressed <i>n</i> = 98	Not depressed <i>n</i> = 192	Statistic	
	Mean ± SD	Mean ± SD	Mean ± SD	Z- NPar ^{^^}	<i>p</i> value
Newborn's age in days (<i>n</i> = 216)	43.13 ± 35.88	45.95 ± 33.92	42.1 ± 36.62	0.955	.341 [^]
Father's age in years (<i>n</i> = 271)	34.97 ± 8.56	34.28 ± 8.4	35.22 ± 8.62	0.454	.455 [^]
Mother's age in years (<i>n</i> = 269)	29.18 ± 7.41	28.96 ± 8.7	29.26 ± 6.91	0.438	.439 [^]
	Median (IQR)	Median (IQR)	Median (IQR)		
Number of children in household (<i>n</i> = 251)	3 (3)	3 (3)	3 (3)	0.602	.602 ^{**}
Father's educational level (<i>n</i> = 277)	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)	Chi-square/exact	.842 [*]
Primary	5 (1.81)	1 (1.28)	4 (2.01)		
Secondary	103 (37.18)	31 (39.74)	72 (36.18)		
Diploma	28 (10.11)	9 (11.54)	19 (9.55)		
Higher education	141 (50.9)	37 (47.44)	104 (52.26)		
Number of newborn babies (<i>n</i> = 279)					
One newborn	271 (97.13)	73 (94.81)	198 (98.02)	-	.223 [*]
Twins	8 (2.87)	4 (5.19)	4 (1.98)		
Father's employment information					
Are you currently employed? (<i>n</i> = 279)					
No	16 (5.73)	2 (2.63)	14 (6.9)	-	.25 ^{**}
Yes	263 (94.27)	74 (97.37)	189 (93.1)		
Employment status if employed (<i>n</i> = 265)					
Permanent	187 (70.57)	59 (80.82)	128 (66.67)	5.103	.024^{**}
Temporary	78 (29.43)	14 (19.18)	64 (33.33)		
Job commitment (<i>n</i> = 262)					
Full-time	218 (83.21)	62 (86.11)	156 (82.11)	0.6	.439 ^{**}
Part-time	44 (16.79)	10 (13.89)	34 (17.89)		
Profession (<i>n</i> = 259)					
Armed forces occupations	92 (35.52)	25 (36.76)	67 (35.08)		
Professional (doctors/engineers)	53 (20.46)	10 (14.71)	43 (22.51)		
Legislator, senior official, or manager	36 (13.9)	9 (13.24)	27 (14.14)	4.579	.469 ^{**}
Clerical support worker	36 (13.9)	12 (17.65)	24 (12.57)		
Technician or associate professional	21 (8.11)	8 (11.76)	13 (6.81)		
Service, sales, or others	21 (8.11)	4 (5.88)	17 (8.9)		
Total household income (SAR; <i>n</i> = 270)					
<3,000	21 (7.78)	2 (2.74)	19 (9.64)	4.842	.304 ^{**}
3,000–4,999	76 (28.15)	22 (30.14)	54 (27.41)		
5,000–8,999	34 (12.59)	12 (16.44)	22 (11.17)		
9,000–14,999	65 (24.07)	16 (21.92)	49 (24.87)		
>15,000	74 (27.41)	21 (28.77)	53 (26.9)		
Do you own your home? (<i>n</i> = 275)					
No	197 (71.64)	53 (71.62)	144 (71.64)	0.000	.997 ^{**}
Yes	78 (28.36)	21 (28.38)	57 (28.36)		

Note. **p* value is based on Fisher's exact test. ***p* value is based on the Chi-square test. [^]*p* value is based on the Wilcoxon rank-sum test (^{^^}non-parametric Z value). SAR = Saudi-Arabian Riyal; SD = standard deviation; IQR = interquartile range. Bold values are statistically significant (*p*-value less than 0.05).

screening for depression among fathers. A cutoff of 8/9 yielded a sensitivity of 77.8% and a specificity of 81.3%. At a cutoff of 8/9, 7 of 9 depressed fathers and 39 of 48

nondepressed fathers were classified correctly, giving an overall accuracy of 80.7%. The gold standard test (*DSM-5*) showed that nine (15.79%) fathers were depressed.




Figure 1. Apparent prevalence of depression in the study sample at different Edinburgh Postnatal Depression Scale cutoffs based on the literature.

Risk Factors of Depression

The potential risk factors of PPNP among fathers of a newborn during postnatal period are summarized in Table 4. Multiple logistic regression using maximum likelihood estimation was used to obtain the odds ratio. Variable selection was based on the response rate, theoretical relevance, pattern of correlation with outcome, and computational feasibility. Initial model included demographic, employment-related, social support of friends/family, family history of depression, marital relationship, pregnancy-related, wife having postpartum depression, and psychological well-being correlates, with less than 30% missing values. Variables with least relevance and that showed weak relationship with outcome variable were eliminated one by one to achieve the final model. The final results included close relative with depression, family setup, social support of friends/family, if the wife is reported as depressed by father, marital relationship, father spending more time at work, and father feeling isolated and disconnected from partner. None of the correlates were identified as statistically significant, except that depressed fathers were more likely to feel isolated or disconnected from their partners compared with nondepressed fathers, with a significant odds ratio of 2.74 (95% CI [1.23, 6.1]) (Table 4).

Discussion

The EPDS has been validated in fathers of newborn across Europe and Far East countries, with a variation in

reported cutoffs. This is the first study to establish the validity of the EPDS among Arabic-speaking fathers of newborn in the Middle East and Saudi Arabia. The established EPDS cutoff in the current study is 8/9, higher than those established for depression (major or minor) in Vietnam (4/5) (Tran et al., 2012), but more or less similar to the one reported for (major or minor) depression (10/11) in Australia, UK, and China (Edmondson et al., 2010; Lai et al., 2010; Matthey et al., 2001). Italian and Swedish studies reported higher cutoffs, 12/13 compared with 8/9 (Loscalzo et al., 2015; Massoudi et al., 2013). The use of different cutoffs had been reported in a series of Australian studies, cutoff 5/6 for depression or anxiety disorder, and 10 or more for reporting minor or major depression (Matthey et al., 2001). However, using the EPDS-3A (three items; 3, 4, and 5), a score of 4 or more can be used for screening anxiety disorder among fathers (Matthey, 2008; Matthey & Agostini, 2017). The sensitivity, 77.8%, and specificity, 81.3%, in the current study are low compared with those reported in Chinese (91% sensitivity, 97% specificity) and Italian (90% sensitivity and 90% specificity) studies (Lai et al., 2010; Loscalzo et al., 2015), but higher than that reported in the Vietnamese study (68% sensitivity, 77% specificity; Tran et al., 2012).

The current study results depict an EPDS cutoff closer to that of the UK/China, but substantially different from that reported by Italy/Australia. This wide variation in cutoff is caused by the geographic location of the study setting as well as cultural factors. One reason for the variation is the use of heterogeneous samples: in the Italian

Table 2. Information Related to the Family, Wife, and Pregnancy for Fathers With/Without Depression.

Correlates of depression <i>n</i> = response rate	Overall <i>n</i> = 290 <i>n</i> (%)	Depressed <i>n</i> = 98 <i>n</i> (%)	Not depressed <i>n</i> = 192 <i>n</i> (%)	Statistics (chi-square/ exact)	<i>p</i> value
Information related to newborn					
Has your newborn child been diagnosed with any of these conditions? (<i>n</i> = 266)					
Congenital birth defects	9 (3.38)	3 (4.05)	6 (3.13)	-	.507**
Down syndrome	6 (2.26)	3 (4.05)	3 (1.56)		
Any other syndrome	7 (2.63)	1 (1.35)	6 (3.13)		
No	244 (91.73)	67 (90.54)	177 (92.19)		
Information related to family					
What is your family setup? (<i>n</i> = 281)					
Nuclear family	228 (81.14)	59 (76.62)	169 (82.84)	1.413	.235**
Joint family	53 (18.86)	18 (23.38)	35 (17.16)		
Do you have a good relationship with your parents? (<i>n</i> = 281)					
No	1 (0.36)	0 (0)	1 (0.49)	-	1*
Yes	280 (99.64)	77 (100)	203 (99.51)		
Do you have the support of friends? (<i>n</i> = 281)					
No	37 (13.17)	10 (12.99)	27 (13.24)	0.003	.956**
Yes	244 (86.83)	67 (87.01)	177 (86.76)		
Do you have the support of family? (<i>n</i> = 281)					
No	16 (5.71)	4 (5.19)	12 (5.91)	-	1**
Yes	264 (94.29)	73 (94.81)	191 (94.09)		
Have any of your close relatives been diagnosed with depression (e.g., mother, father, brother, or sister)? (<i>n</i> = 265)					
No	244 (92.08)	62 (87.32)	182 (93.81)	3.001	.083**
Yes	21 (7.92)	9 (12.68)	12 (6.19)		
Information related to wife					
Number of wives (<i>n</i> = 282)					
One	261 (92.55)	72 (93.51)	189 (92.2)	0.14	.709**
More than one	21 (7.45)	5 (6.49)	16 (7.8)		
Marital status (<i>n</i> = 279)					
Married: living together	276 (98.92)	75 (98.68)	201 (99.01)	-	1*
Married: living separately	3 (1.08)	1 (1.32)	2 (0.99)		
Is your marital relationship good? (<i>n</i> = 278)					
No	4 (1.44)	1 (1.33)	3 (1.48)	-	1*
Yes	274 (98.56)	74 (98.67)	200 (98.52)		
Do you think that your wife is depressed? (<i>n</i> = 277)					
No	252 (90.97)	66 (86.84)	186 (92.54)	2.179	.14**
Yes	25 (9.03)	10 (13.16)	15 (7.46)		
Did your wife go to her mother's house postdelivery? (<i>n</i> = 277)					
No	122 (44.04)	29 (38.67)	93 (46.04)	1.206	.272**
Yes	155 (55.96)	46 (61.33)	109 (53.96)		
If not, did anyone join her at home? (<i>n</i> = 195)					
No	93 (47.69)	26 (44.07)	67 (49.26)	0.446	.505**
Yes	102 (52.31)	33 (55.93)	69 (50.74)		
Information related to pregnancy					
Was this pregnancy planned? (<i>n</i> = 279)					
No	120 (43.01)	30 (39.47)	90 (44.33)	0.533	.465**
Yes	159 (56.99)	46 (60.53)	113 (55.67)		
Was the pregnancy conceived through IVF? (<i>n</i> = 279)					
No	266 (95.34)	70 (93.33)	196 (96.55)	-	.316*
Yes	12 (4.3)	5 (6.67)	7 (3.45)		

(continued)

Table 2. (continued)

Correlates of depression <i>n</i> = response rate	Overall <i>n</i> = 290 <i>n</i> (%)	Depressed <i>n</i> = 98 <i>n</i> (%)	Not depressed <i>n</i> = 192 <i>n</i> (%)	Statistics (chi-square/ exact)	<i>p</i> value
Did you attend antenatal checkups with your wife? (<i>n</i> = 279)					
No	17 (6.09)	6 (7.89)	11 (5.42)	-	.414*
Yes	262 (93.91)	70 (92.11)	192 (94.58)		
Did you attend the delivery of your child? (<i>n</i> = 278)					
No	54 (19.42)	14 (18.42)	40 (19.8)	0.067	.795**
Yes	224 (80.58)	62 (81.58)	162 (80.2)		
Mode of delivery (<i>n</i> = 281)					
Vaginal delivery	191 (67.97)	56 (73.68)	135 (65.85)	1.562	.211**
Cesarean section	90 (32.03)	20 (26.32)	70 (34.15)		
Did your wife experience complications during childbirth? (<i>n</i> = 280)					
No	249 (88.93)	63 (82.89)	186 (91.18)	3.857	.05**
Yes	31 (11.07)	13 (17.11)	18 (8.82)		
Had you ever lost a child before the birth of this child? (<i>n</i> = 278)					
No	234 (84.17)	61 (81.33)	173 (85.22)	0.622	.431**
Yes	44 (15.83)	14 (18.67)	30 (14.78)		
Information related to father's emotional status					
Do you spend more time at work to disconnect from home and family? (<i>n</i> = 277)					
No	253 (91.34)	67 (91.78)	186 (91.18)	0.025	.875**
Yes	24 (8.66)	6 (8.22)	18 (8.82)		
Do you feel isolated and disconnected from your partner? (<i>n</i> = 277)					
No	241 (87)	59 (78.67)	182 (90.1)	6.322	.012**
Yes	36 (13)	16 (21.33)	20 (9.9)		
Have you ever received a speeding ticket after your child's birth? (<i>n</i> = 277)					
No	230 (83.03)	61 (81.33)	169 (83.66)	0.211	.646**
Yes	47 (16.97)	14 (18.67)	33 (16.34)		

Note. **p* value based on Fisher's exact test. ***p* value based on the chi-square test. IVF = in vitro fertilization. Bold values are statistically significant (*p*-value less than 0.05).

Table 3. Receiver Operating Characteristic Curve Analysis.

EPDS cutoff	1 - Specificity	Sensitivity
≥ 5	0.56	0.89
≥ 6	0.44	0.89
≥ 7	0.33	0.78
≥ 8	0.19	0.78
≥ 9*	0.19	0.78
≥ 10	0.15	0.67
≥ 11	0.04	0.56
≥ 12	0.02	0.22
≥ 13	0.02	0.22
≥ 14	0.02	0.11

Note. EPDS = Edinburgh Postnatal Depression Scale. *The cutoff used to identify depression.

study, the sample of fathers was very diverse in terms of inclusion criteria, and the majority of fathers had grown-up children (Loscalzo et al., 2015). The time of screening depression has also varied across studies; antenatal




Figure 2. Sensitivity and false-negative rate of Edinburgh Postnatal Depression Scale scores in the study sample.

Table 4. Risk Factors of Depression Among Fathers of Newborn.

Risk factors of depression	Estimate (SE)	Odds ratio (95% CI)	p value
Intercept	-2.06 (1.47)	–	.162
Close relatives with depression (yes vs. no)	0.87 (0.5)	2.4 [0.89, 6.44]	.083
Family setup (joint vs. nuclear)	0.12 (0.18)	1.29 [0.63, 2.64]	.494
Social support of friends (yes vs. no)	0.06 (0.5)	1.06 [0.4, 2.83]	.907
Social support of family (yes vs. no)	0.85 (0.95)	2.34 [0.37, 15.02]	.369
Wife depressed (yes vs. no)	0.03 (0.57)	1.03 [0.34, 3.13]	.954
Marital relationship (good vs. bad)	-0.41 (1.45)	0.67 [0.04, 11.4]	.779
Spend more time at work (yes vs. no)	-0.27 (0.57)	0.76 [0.25, 2.34]	.638
Feeling isolated and disconnected from partner (yes vs. no)	1.01 (0.41)	2.74 [1.23, 6.1]	.013*

Note. Probabilistic model is based on the probability of having depression. CI = confidence interval; SE = standard error. The italics are reference groups. *Statistically significant p-value.

versus postnatal. Another reason for the variation is that the symptoms of depression vary between men and women; men suffer with anger, irritability, emotional rigidity, and sleep disturbance (Edward et al., 2015). Men are more likely to have difficulty focusing on work and abuse alcohol/drugs. The symptoms also vary within populations. The prevalence of PPND varies depending on the cutoff used. The use of nonvalidated cutoff leads to a different clinical interpretation of the rates of perinatal depression (Matthey, Henshaw, Elliott, & Barnett, 2006).

The advantages of using the EPDS are that it is open access, it takes only a short time to administer, it is easy to understand, and it is a reasonable tool for screening fathers of newborn for PPND provided that recalibration is performed based on the specific study setting, population, and culture. Studies have reported use of the Gotland Male Depression Scale (GMDS) instrument apart from the EPDS. The Swedish study reported combining the EPDS and GMDS as the Edinburgh–Gotland Depression Scale (EGDS) which showed higher sensitivity than the EPDS alone (Madsen & Juhl, 2007; Psouni, Agebjorn, & Linder, 2017).

The result of this study indicates that 16.6% fathers of newborn have PPND. This is the first study to report the prevalence of postnatal depression among fathers of newborn in Saudi Arabia. Prior published results have reported PPND ranging between 2% and 31% (Bergstrom, 2013; Edoka et al., 2011; Goodman, 2008; Matthey et al., 2001; Moussa et al., 2012; O'Hara & McCabe, 2013; Zhang et al., 2016). In the Egyptian study, the reported depression rate (31.4%) among expecting fathers is higher compared to 16.6%; however the results are based on the non-validated cutoff 10 or higher during antenatal period (Moussa et al., 2012). The results are more or less similar to the one reported in Sweden, 10.3% at 3 months postpartum (Bergstrom, 2013); in the UK, 12.7% at 7 weeks postpartum (Edoka

et al., 2011); in the United States, 13% at 2 months postpartum (Goodman, 2008); and in China, 20.4% at 2 weeks postpartum (Zhang et al., 2016). The prevalence is higher than that reported in Turkey, 1.8% at 2–6 months postpartum (O'Hara & McCabe, 2013), and in Australia, 3.3% at 6–9 weeks postpartum (Matthey et al., 2001). The comparison with prior published studies is difficult since different methodological approach and diagnostic criteria of depression was used. These studies were selected for comparison because the EPDS was used as a screening tool. However, the time of PPND assessment varied across the studies.

The prevalence of PPND among Saudi fathers, 16.6%, reported in this study is very similar to that of MPPD, 14%, reported among Saudi mothers (Al-Modayfer et al., 2015). The estimated prevalence of PPND can be as high as 50% if the mother is also depressed (Goodman, 2004). Depression among fathers tends to start later than among mothers, and the rate of depression among fathers increases at 6 and 9 months postchildbirth (Goodman, 2004). The estimated prevalence of PPND tends to decline over time during the postnatal period (Goodman, 2004).

There are several confounding factors in PPND research, the most important of which is the lack of standard criteria to diagnose PPND. Other factors that can lead to divergent prevalence results include different methodologic approaches, different assessment tools for screening, clinical heterogeneity, the timing of depression assessment (ante-partum/early postpartum vs. late postpartum), study setting, and social and cultural restraints (Paulson & Bazemore, 2010). Such differences make the generalization of results very difficult.

Several risk factors are associated with PPND, such as unemployment, unplanned pregnancy, poor marital relationship, lack of family support, previous history of depression, the presence of depression, the presence of

maternal depression, joint families, sex of the baby, and antenatal paternal depression (Moussa et al., 2012). Among them, MPPD has been identified as the most significant risk factor for the development of PPND, and fathers are 8.4 times more likely to get depressed if their partner is depressed (Pinheiro et al., 2006). Researchers have recommended that fathers should be screened for depression, especially if their partners are depressed (Pinheiro et al., 2006). A study conducted in Japan reported no association between maternal and paternal depression, but reported that PPND was associated with employment status, past history of psychiatric treatment, and unplanned pregnancy (Nishimura & Ohashi, 2010).

The relationship of several factors with the risk of depression among fathers was examined in this study. The risk factors were identified based on prior literature, and cultural and social values. Interestingly, no significant association was identified between PPND and suspected risk factors such as employment status, social support of family/friends, family history of depression, family setup, maternal postpartum depression (as reported by fathers), marital relationship, and spending more time at work. The only risk factor significantly associated with PPND was feeling isolated and disconnected from one's partner. These results are inconsistent with prior studies that reported associations between similar risk factors and paternal depression (Moussa et al., 2012; Nishimura & Ohashi, 2010). One possible reason of insignificance is sample size to assess risk factors. The second possible reason is use of unstandardized methods for risk factors measurement. One of the compelling predictors of PPND reported in the literature is maternal depression (Goodman, 2004). Past history of depression is also associated with PPND (Spry et al., 2018). In the current study, fathers with a history of depression were excluded, which indicates that PPND can appear as first-time depression among men. Fathers who reported social support during pregnancy reported less depression (Castle, Slade, Barranco-Wadlow, & Rogers, 2008).

The symptoms of depression vary between men and women; the most commonly reported depressive symptoms among fathers of newborn are exhaustion, irritability, despair, anger, emotional rigidity, self-criticism, sleep disturbance, risk-taking behavior, and alcohol and/or drug abuse (Brownhill, Wilhelm, Barclay, & Schmied, 2005; Goodman, 2004; Martin, Neighbors, & Griffith, 2013). The causes of PPND among fathers are a sense of responsibility, the inability to cope with changes due to the arrival of a new baby, changes in lifestyle, changes in the marital relationship, lack of sleep, increased expenses, and increased workload (Goodman, 2004).

The study fills a gap in the literature regarding PPND among Saudi fathers. The findings provide direction for perinatal mental health assessments during the transition

to parenthood. The results were strengthened by EPDS validation in the study sample. The results of the field survey are applicable to the general population because a random sample of fathers was selected from all fathers living in the central region and visiting the birth registration office. The generated results are unbiased because the psychologist was blinded to the participants' EPDS scores. Unlike other studies, fathers with a prior history of depression were excluded. The sample was homogeneous in terms of the age of the baby. The prevalence is reported after adjustment for sensitivity and specificity.

Limitations

The current study has some limitations. The tool was less sensitive, and requires validation keeping in mind the social, cultural, and religious beliefs of society. A focus group discussion is needed to develop a tool specifically intended to be used among fathers.

Because of the study setting, assessment of the newborn mothers for depression was impractical. Consequently, the ability to understand the impact of maternal depression on paternal depression was impaired. To mitigate the issue, a simple, self-reported question about the mother's depression status was included in the questionnaire. Because of the lack of a control group of normal individuals, it was impossible to assess whether the observed depression rates differed from that reported in the general population. This study has relied on a single psychologist to assess the ground of truth of depression. This approach might have an implication on the reproducibility of the estimated prevalence. Finally, the sample size is not very large which might impact the generalizability of the results; however the sample size is close to the average sample size (313) reported across studies.

The EPDS is a reasonably sensitive tool for screening fathers of newborn for PND provided that recalibration is performed based on specific populations and cultures. However, the current diagnostic criteria of PND are biased toward women's response to depression. Diagnostic instruments to measure symptoms specific to depression in men may result in an increase in detection of the condition (Martin et al., 2013). There is a need to develop an instrument focused on the mental health of men during the perinatal period.

Conclusion

Depression in men is a serious problem, and most men with the condition suffer without acknowledging it because of the stigma attached to depression by society. PPND must be recognized as a serious public health issue. Both men and women experience depression, but

the willingness of men to share their feelings after a newborn's birth differs from that of women (Brownhill et al., 2005). It is necessary to identify fathers with PPND to prevent the development of behavioral problems in their children and disruption of their relationship with their spouse. Perinatal mental health assessment should focus on the family as a unit. Most laypeople and health-care workers are unaware of PPND. Therefore, clinicians working in obstetrics and pediatrics must be educated about PPND. There is a need to develop an instrument focused on the mental health of men during the perinatal period. Future long-term cohort studies at different time points (antenatal, postnatal, and during the years of child growth) are recommended to determine the implications of PPND.

Acknowledgments

The authors would like to thank all the fathers who shared their experiences. The authors would also like to thank the KAIMRC for providing the grant and the postnatal ward nurses for their support.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by the King Abdullah International Medical Research Center (KAIMRC) with the grant number RC13/206/R. The funder had no further role in the study design, data collection, analysis or interpretation of the data, report writing, or decision to submit the article for publication. The views expressed in paper are solely of authors.

ORCID iD

Naila A. Shaheen  <https://orcid.org/0000-0001-8756-3591>

References

- Al-Modayfer, O., Alatiq, Y., Khair, O., & Abdelkawi, S. (2015). Postpartum depression and related risk factors among Saudi females. *International Journal of Culture and Mental Health, 8*(3), 316–324.
- Bergstrom, M. (2013). Depressive symptoms in new first-time fathers: Associations with age, sociodemographic characteristics, and antenatal psychological well-being. *Birth, 40*(1), 32–38.
- Brockington, I. (2004). Postpartum psychiatric disorders. *The Lancet, 363*(9405), 303–310.
- Brownhill, S., Wilhelm, K., Barclay, L., & Schmied, V. (2005). "Big build": Hidden depression in men. *Australian & New Zealand Journal of Psychiatry, 39*(10), 921–931.
- Cameron, E. E., Sedov, I. D., & Tomfohr-Madsen, L. M. (2016). Prevalence of paternal depression in pregnancy and the postpartum: An updated meta-analysis. *Journal of Affective Disorders, 206*, 189–203.
- Castle, H., Slade, P., Barranco-Wadlow, M., & Rogers, M. (2008). Attitudes to emotional expression, social support and postnatal adjustment in new parents. *Journal of Reproductive and Infant Psychology, 26*(3), 180–194.
- Condon, J. T., Boyce, P., & Corkindale, C. J. (2004). The first-time fathers study: A prospective study of the mental health and wellbeing of men during the transition to parenthood. *Australian & New Zealand Journal of Psychiatry, 38*(1–2), 56–64.
- Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry: The Journal of Mental Science, 150*, 782–786.
- Da Costa, D., Zekowitz, P., Dasgupta, K., Sewitch, M., Lowensteyn, I., Cruz, R., . . . Khalife, S. (2017). Dads get sad too: Depressive symptoms and associated factors in expectant first-time fathers. *American Journal of Men's Health, 11*(5), 1376–1384.
- Edmondson, O. J. H., Psychogiou, L., Vlachos, H., Netsi, E., & Ramchandani, P. G. (2010). Depression in fathers in the postnatal period: Assessment of the Edinburgh Postnatal Depression Scale as a screening measure. *Journal of Affective Disorders, 125*(1–3), 365–368.
- Edoka, I. P., Petrou, S., & Ramchandani, P. G. (2011). Healthcare costs of paternal depression in the postnatal period. *Journal of Affective Disorders, 133*(1–2), 356–360.
- Edward, K., Castle, D., Mills, C., Davis, L., & Casey, J. (2015). An integrative review of paternal depression. *American Journal of Men's Health, 9*(1), 26–34.
- Ghubash, R., Abou-Saleh, M. T., & Daradkeh, T. K. (1997). The validity of the Arabic Edinburgh Postnatal Depression Scale. *Social Psychiatry and Psychiatric Epidemiology, 32*(8), 474–476.
- Gonzalez, J. M., Alegria, M., & Prihoda, T. J. (2005). How do attitudes toward mental health treatment vary by age, gender, and ethnicity/race in young adults? *Journal of Community Psychology, 33*(5), 611–629.
- Goodman, J. H. (2004). Paternal postpartum depression, its relationship to maternal postpartum depression, and implications for family health. *Journal of Advanced Nursing, 45*(1), 26–35.
- Goodman, J. H. (2008). Influences of maternal postpartum depression on fathers and on father–infant interaction. *Infant Mental Health Journal, 29*(6), 624–643.
- Kim, P., & Swain, J. E. (2007). Sad dads: Paternal postpartum depression. *Psychiatry (Edmont), 4*(2), 35–47.
- Lai, B. P. Y., Tang, A. K. L., Lee, D. T. S., Yip, A. S. K., & Chung, T. K. H. (2010). Detecting postnatal depression in Chinese men: A comparison of three instruments. *Psychiatry Research, 180*(2–3), 80–85. doi:10.1016/j.psychres.2009.07.015
- Loscalzo, Y., Giannini, M., Contena, B., Gori, A., & Benvenuti, P. (2015). The Edinburgh Postnatal Depression Scale for

- fathers: A contribution to the validation for an Italian sample. *General Hospital Psychiatry*, 37(3), 251–256.
- Madsen, S. A., & Juhl, T. (2007). Paternal depression in the postnatal period assessed with traditional and male depression scales. *The Journal of Men's Health & Gender*, 4(1), 26–31. doi:10.1016/j.jmhg.2006.10.017
- Martin, L. A., Neighbors, H. W., & Griffith, D. M. (2013). The experience of symptoms of depression in men vs women: Analysis of the national comorbidity survey replication. *JAMA Psychiatry*, 70(10), 1100–1106.
- Massoudi, P., Hwang, C. P., & Wickberg, B. (2013). How well does the Edinburgh Postnatal Depression Scale identify depression and anxiety in fathers? A validation study in a population based Swedish sample. *Journal of Affective Disorders*, 149(1–3), 67–74.
- Matthey, S. (2008). Using the Edinburgh Postnatal Depression Scale to screen for anxiety disorders. *Depression and Anxiety*, 25(11), 926–931.
- Matthey, S., & Agostini, F. (2017). Using the Edinburgh Postnatal Depression Scale for women and men—some cautionary thoughts. *Archives of Women's Mental Health*, 20(2), 345–354.
- Matthey, S., Barnett, B., Kavanagh, D. J., & Howie, P. (2001). Validation of the Edinburgh Postnatal Depression Scale for men, and comparison of item endorsement with their partners. *Journal of Affective Disorders*, 64(2–3), 175–184.
- Matthey, S., Henshaw, C., Elliott, S., & Barnett, B. (2006). Variability in use of cut-off scores and formats on the Edinburgh Postnatal Depression Scale: Implications for clinical and research practice. *Archives of Women's Mental Health*, 9(6), 309–315. doi:10.1007/s00737-006-0152-x
- Miller, J. (2002). The Scottish Intercollegiate Guidelines Network (SIGN). *The British Journal of Diabetes & Vascular Disease*, 2(1), 47–49.
- Moussa, S., Emad, M., Khoweiled, A., Amer, D., Refaat, O., & Goueli, T. (2012). Antenatal depression in expectant fathers (an Egyptian study). *Egyptian Journal of Psychiatry*, 33(2), 90–96.
- Musser, A. K., Ahmed, A. H., Foli, K. J., & Coddington, J. A. (2013). Paternal postpartum depression: What health care providers should know. *Journal of Pediatric Health Care: Official Publication of National Association of Pediatric Nurse Associates & Practitioners*, 27(6), 479–485.
- Nishimura, A., & Ohashi, K. (2010). Risk factors of paternal depression in the early postnatal period in Japan. *Nursing & Health Sciences*, 12(2), 170–176.
- O'Hara, M. W., & McCabe, J. E. (2013). Postpartum depression: Current status and future directions. *Annual Review of Clinical Psychology*, 9, 379–407.
- Paulson, J. F., & Bazemore, S. D. (2010). Prenatal and postpartum depression in fathers and its association with maternal depression: A meta-analysis. *JAMA*, 303(19), 1961–1969.
- Pinheiro, R. T., Magalhaes, P. V. S., Horta, B. L., Pinheiro, K. A. T., Silva, R. A., & Pinto, R. H. (2006). Is paternal postpartum depression associated with maternal postpartum depression? Population-based study in Brazil. *Acta Psychiatrica Scandinavica*, 113(3), 230–232.
- Psouni, E., Agebjorn, J., & Linder, H. (2017). Symptoms of depression in Swedish fathers in the postnatal period and development of a screening tool. *Scandinavian Journal of Psychology*, 58(6), 485–496.
- Ramchandani, P., Stein, A., Evans, J., & O'Connor, T. G. (2005). Paternal depression in the postnatal period and child development: A prospective population study. *The Lancet*, 365(9478), 2201–2205. doi:10.1016/S0140-6736(05)66778-5
- Ramchandani, P. G., Stein, A., O'Connor, T. G., Heron, J., Murray, L., & Evans, J. (2008). Depression in men in the postnatal period and later child psychopathology: A population cohort study. *Journal of the American Academy of Child & Adolescent Psychiatry*, 47(4), 390–398. doi:10.1097/CHI.0b013e31816429c2
- Reynolds, C. F., & Patel, V. (2017). Screening for depression: The global mental health context. *World Psychiatry*, 16(3), 316–317.
- Spry, E., Giallo, R., Moreno-Betancur, M., Macdonald, J., Becker, D., Borschmann, R., ... Olsson, C. A. (2018). Preconception prediction of expectant fathers' mental health: 20-year cohort study from adolescence. *BJPsych Open*, 4, 58–60.
- Tran, T. D., Tran, T., & Fisher, J. (2012). Validation of three psychometric instruments for screening for perinatal common mental disorders in men in the north of Vietnam. *Journal of Affective Disorders*, 136(1–2), 104–109.
- Weinberger, A. H., Gbedemah, M., Martinez, A. M., Nash, D., Galea, S., & Goodwin, R. D. (2018). Trends in depression prevalence in the USA from 2005 to 2015: Widening disparities in vulnerable groups. *Psychological Medicine*, 48(8), 1308–1315.
- World Health Organization. (2017). *Depression and other common mental disorders: Global health estimates*. Geneva: World Health Organization.
- Zhang, Y.-P., Zhang, L.-L., Wei, H.-H., Zhang, Y., Zhang, C.-L., & Porr, C. (2016). Post partum depression and the psychosocial predictors in first-time fathers from northwestern China. *Midwifery*, 35, 47–52.