

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/148263/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Deli, John, González-Beiras, Camila, Guldan, Georgia S., Moses, Rachael L., Dally, Jordanna, Moseley, Ryan, Lundy, Fionnuala T., Corbacho-Monne, Marc, Walker, Stephen L., Ubals Cazorla, Maria, Ouchi, Dan, Fang, Rui, Briggs, Marie, Kiapranis, Robert, Yahimbu, Martha, Mitjà, Oriol and Prescott, Thomas A.K. 2022. Ficus septica exudate, a traditional medicine used in Papua New Guinea for treating infected cutaneous ulcers: in vitro evaluation and clinical efficacy assessment by cluster randomised trial. *Phytomedicine* 99, 154026. 10.1016/j.phymed.2022.154026

Publishers page: <http://doi.org/10.1016/j.phymed.2022.154026>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1
2 **Ficus septica exudate, a traditional medicine used in Papua New**
3 **Guinea for treating infected cutaneous ulcers: in vitro evaluation**
4 **and clinical efficacy assessment by cluster randomised trial**
5

6 John Deli^{1*}, Camila González-Beiras^{2*}, Georgia S Guldan¹, Rachael L. Moses³, Jordanna Dally³,
7 Ryan Moseley,³ Fionnuala T. Lundy⁴, Marc Corbacho-Monne², Stephen L Walker⁵, Maria Ubals
8 Cazorla², Dan Ouchi², Rui Fang⁶, Marie Briggs⁶, Robert Kiapranis⁷, Martha Yahimbu¹, Oriol Mitjà^{8*},
9 Thomas A.K. Prescott^{6*†}

10
11 ¹*Division of Public Health, School of Medicine and Health Sciences, University of Papua New Guinea*

12 ²*Fight AIDS and Infectious Diseases Foundation, Hospital Universitari Germans Trias i Pujol, Universitat Autònoma de Barcelona, Spain*

13 ³*Oral and Biomedical Sciences, School of Dentistry, Cardiff Institute Tissue Engineering and Repair, Cardiff University, Cardiff, UK*

14 ⁴*Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry and Biomedical Sciences, Queen's University*
15 *Belfast, BT9 7BL, UK*

16 ⁵*London School of Hygiene and Tropical Medicine, London, United Kingdom.*

17 ⁶*Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, UK*

18 ⁷*Papua New Guinea Forest Research Institute, Lae, Papua New Guinea*

19 ⁸*Fight AIDS and Infectious Diseases Foundation, Hospital Universitari Germans Trias i Pujol, Universitat Autònoma de Barcelona, Spain*
20 *and Lihir Medical Center–International SOS, Lihir Island, Papua New Guinea*

21
22 *Co-first authors and co-last authors

23
24 † Corresponding author. Tel.: +44 208 332 5393; fax: +44 208 332 5310.
25 *E-mail address:* t.prescott@kew.org (T.A.K Prescott).
26
27
28

29 Keywords: Ficuseptine; *Ficus septica*; Papua New Guinea; clinical trial; ulcer.

30
31 Abbreviations: DPP (dual path platform, syphilis screen and confirm); ELISA (enzyme-linked immunosorbent
32 assay); FCS (foetal calf serum); GM-CSF (granulocyte-macrophage colony-stimulating factor); IL-6 (interleukin-
33 6), LPS (lipopolysaccharide); MMP (matrix metalloproteinase); NMR (nuclear magnetic resonance); PBS
34 (phosphate buffered saline); TNF- α (tumour necrosis factor- α); Tx (treatment).

35 **ABSTRACT**

36 **Background and objectives:** Infected cutaneous ulcers are a major health problem for children living
37 in rural areas of Papua New Guinea. The inaccessibility of affected populations and lack of access to
38 basic healthcare, makes a local plant-based therapy an attractive treatment option. We assessed *Ficus*
39 *septica* exudate in biological assays relevant to wound healing. We then carried out a clinical trial to
40 determine the exudate's efficacy in healing small cutaneous ulcers compared with chlorhexidine
41 cetrimide cream, and soap and water washing.

42 **Methods:** Pre-clinical *in vitro* assessment of the exudate was carried out using assays to monitor the
43 pro-inflammatory responses of M1 macrophages and neutrophils, antibacterial assays using known
44 ulcer pathogens, an Ames test for mutagenicity and LC-MS chemical analysis of the exudate. An open
45 label cluster-randomised clinical trial was performed, enrolling participants from three different clusters
46 with skin lesions less than 1cm in diameter. Each cluster comprising 50 participants was randomly
47 assigned to one of three treatment arms namely topical exudate, topical chlorhexidine with cetrimide
48 cream, and standard care (soap and water treatment), all administered daily for 2 days. The primary
49 outcome was clinical healing/improvement measured at days 7 and 14, assessed by three dermatologists
50 using blinded photographs. Primary analysis was non-inferiority of *F. septica* treatment based on the
51 risk difference for healing/improvement.

52 **Results:** *In vitro*, the exudate which is rich in the alkaloid ficuseptine, was found to be non-mutagenic
53 whilst also inhibiting pro-inflammatory responses and exhibiting antibacterial activity. When
54 administered to participants enrolled in the clinical trial, no significant differences were observed
55 between the healing efficacy of *F. septica* exudate and the two comparators (chlorhexidine cream and
56 soap/water treatment). At day 14, but not at day 7, the efficacy of *F. septica* exudate for
57 healing/improving the ulcers was non-inferior to chlorhexidine or water/soap treatment.

58 **Conclusions:** *F. septica* exudate is non-mutagenic and has both bactericidal and anti-inflammatory
59 properties. When applied topically to small cutaneous ulcers, the exudate has a healing effect that is
60 non-inferior to chlorhexidine and standard treatment with soap and water at day 14. Our findings, which
61 should be confirmed in larger clinical trials, have important public health implications.

62

63

64

65

66 Introduction

67 Infected cutaneous ulcers are a painful condition that are common in amongst children in Papua New
68 Guinea. The ulcers present as single or multiple non-purulent skin lesions showing epidermal loss, and
69 are chronic, generally lasting more than 2 weeks (Noguera-Julian et al. 2019). They are believed to
70 develop from a small scratch or trauma, which subsequently becomes infected. In rural tropical regions,
71 cutaneous ulcers are most prevalent in children, with infection rates that may range from 5 to 15%
72 (Noguera-Julian et al. 2019). Although accurate figures for Papua New Guinea are lacking, our own
73 observations suggest that most children living in rural villages either have active ulcer infections or
74 show scars from previous ulcers. As an estimated 6.4 million people (80% of Papua New Guinea's
75 population) live in rural areas (Schmidt Mueller et al., 2020), cutaneous ulcers represent a significant
76 area of unmet clinical need. Furthermore, the unusual circumstances in Papua New Guinea, in which
77 relatively large and scattered rural populations live in highly inaccessible rainforest villages has
78 prevented the development of an effective aid post network, and as a result, in these areas, conventional
79 treatment options for cutaneous ulcers are extremely limited. For these populations, early treatment of
80 small, infected lesions with a locally available plant antiseptic is preferable to late treatment with
81 antibiotics in a hard-to-reach clinic and may, therefore, represent a valuable treatment option.
82 Acknowledging the limited access to medicines in rural areas, the Papua New Guinea standard
83 treatment guidelines for children recommend the use of soap and water to prevent skin disease and
84 ulcers (Paediatric Society of Papua New Guinea, 2016); however, evidence for this approach from
85 clinical trials is currently lacking.

86 The microbiology of cutaneous ulcers in Papua New Guinea is a developing research area; the aetiology
87 of cutaneous ulcers may differ, with *Treponema pallidum* subsp *pertenue* (the causative agent of yaws
88 disease) and *Haemophilus ducreyi* (the causative agent of chancroid) suggested to be causative agents
89 (Mitja et al. 2014; Noguera-Julian et al. 2019). Earlier work dating back to the 1980s that used culture-
90 based identification methods had identified *Fusobacterium ulcerans* as a potential pathogen (Aadrians
91 and Shah 1988; Lupi et al. 2006) whereas another study carried out in the highlands of Papua New
92 Guinea identified β -haemolytic *Streptococci*, *Staphylococcus aureus*, *Corynebacterium diphtheriae* and
93 *Corynebacterium haemolyticum* as potential wound pathogens (Montgomery 1985). More recent
94 metagenomics-based studies carried out in the Solomon Islands, identified *H. ducreyi* as a prominent
95 pathogen (Marks et al., 2014), and further application of metagenomics to yaws-like ulcers in Lihir
96 Island (Papua New Guinea) revealed a range of bacterial species including *H. ducreyi*, *Streptococcus*
97 *pyogenes* and *Streptococcus dysgalactiae* (Griesenauer et al. 2021; Noguera-Julian et al. 2019).

98 An additional aspect to be considered is the role of excessive inflammation in these ulcers. Although
99 little is known regarding the role of inflammation in small cutaneous ulcers in Papua New Guinea,
100 extensive research carried out on venous ulcers, diabetic ulcers, and pressure sores in high income

101 countries, has revealed a correlation between the chronic nature of the ulcers and persistent or
102 dysregulated inflammation (Zhao et al. 2016). In these types of ulcers, pro-inflammatory neutrophils
103 and M1 macrophages, which predominate during early dermal healing, persist in the ulcers, resulting
104 in wounds perpetually remaining in the early inflammatory phase (Eming et al. 2007). The neutrophils
105 and macrophages are responsible for elevated levels of pro-inflammatory cytokines, such as IL-1 and
106 IL-6, and TNF- α . These in turn increase the expression of matrix metalloproteinases (MMPs), which
107 degrade extracellular matrix components within these wounds (Barrientos et al. 2008; Moseley et al.
108 2004). Additionally, fibroblasts present in nonhealing wounds display impaired proliferative and
109 migratory responses as a consequence of enhanced cellular senescence within chronic wound
110 environments (Wall et al. 2008). Thus, topical exudate treatments that eradicate wound infections,
111 reduce the subsequent inflammatory responses, or stimulate fibroblast proliferation and keratinocyte
112 healing responses may reduce the chronic non-healing nature of cutaneous ulcers (Moses et al. 2020b).

113 *F. septica* Burm.f. (Moraceae), is a tree or shrub with yellowish exudate, and has a distribution that
114 includes Taiwan, Malesia (including New Guinea and the Bismark Archipelago), Australia (Northern
115 Queensland) through to Vanuatu (Berg Corner, E.J.H. 2005). *F. septica* exudate has traditionally been
116 used as a topical treatment for cutaneous ulcers in Papua New Guinea and our own ethnobotanical
117 observations in the Kaulong speaking area of New Britain reveal the exudate is still used as a topical
118 treatment for cutaneous ulcers (Baumgartner et al. 1990). The exudate is obtained by breaking off the
119 figs and allowing the exudate to drip from the broken peduncle onto the surface of infected cutaneous
120 ulcers in the same manner as a topical antiseptic. The tree is commonly found in secondary forest near
121 villages and is therefore a highly accessible treatment option. Extensive phytochemical isolation from
122 stems, roots, and leaves of *F. septica* has yielded a wide variety of phenanthroindolizidine alkaloids,
123 ficuseptine and ficuseptines A-N (Damu et al. 2005, 2009; Kubo et al. 2016; Ueda et al. 2009; Wu et
124 al., 2002). In addition, vanillic acid, (5-acetyl-2-hydroxyphenyl)- β -D-glucopyranoside, the coumarins
125 umbelliferone and esculin, the flavonoids genistin and kaempferitrin, squalene, and uracil have been
126 isolated from the leaves (Wu et al., 2002). *F. septica* has antimicrobial activity which has been attributed
127 to the alkaloids ficuseptine and antofine (Baumgartner et al. 1990; Prescott et al. 2015). Furthermore,
128 two alkaloids reported from *F. septica*, tylophorine and ficuseptine-A, both inhibit nitric oxide
129 production in lipopolysaccharide (LPS)/interferon- γ -stimulated macrophages, without causing
130 significant cytotoxicity (Yang et al. 2006).

131 In the present study, we investigated the suitability of *F. septica* exudate as a low-cost treatment option
132 for small cutaneous ulcers in rural areas of Papua New Guinea. *F. septica* is particularly suitable as a
133 first-line treatment because it is commonly found growing near areas of human habitation. Our study
134 includes both an *in vitro* assessment of the exudate's suitability as a topical ulcer treatment, as well as
135 a clinical trial to determine its efficacy.

136 **Materials and methods**

137 *Collection and preparation of plant material*

138 Permission for collection and issuing of export licences for botanical specimens was provided by the
139 National Herbarium of Papua New Guinea. Two exudate samples were collected along with
140 corresponding voucher specimens; one sample for *in vitro* pre-clinical testing and the other sample for
141 the clinical trial. The exudate was collected by precisely replicating the traditional method which was
142 observed first-hand during previous ethnobotanical fieldwork (Prescott et al., 2012). The exudate was
143 harvested by picking the figs and allowing a drop of exudate to form on the broken peduncle (stalk);
144 the exudate droplet was then touched onto the inside edge of a 2ml cryovial and the process repeated
145 until the vial was filled with exudate. The sample for *in vitro* testing was collected in Umbi Village,
146 West New Britain in March 2017. A voucher specimen (T.A.K.P. 164) from the same tree is lodged at
147 the herbarium, Royal Botanic Gardens, Kew. To prepare this exudate sample for *in vitro* work, the
148 exudate was centrifuged at 13,400 g, and then filtered through a 0.02 µm PTFE syringe filter under
149 sterile conditions and stored at -20 °C. Subsequent lyophilisation and weighing of the centrifuged and
150 filtered exudate sample used for the *in vitro* pre-clinical experiments revealed it to have a concentration
151 of 147.43 mg exudate metabolites per ml of exudate.

152 The sample for the clinical trial was collected directly into plastic vials using the same method but
153 without centrifugation and filtration. The exudate was collected in November 2019 from two separate
154 tree specimens in Lihir Island and Kokopo, East New Britain, vouchers for each respectively, (T.A.K.P.
155 176) and (T.A.K.P. 177) are lodged at Kew and the Forest Research Institute, Lae, Papua New Guinea.
156 The two exudate samples were pooled together to make a single sample and stored briefly at 5 °C before
157 beginning the clinical trial. A portion of the clinical trial exudate sample was centrifuged and filtered
158 in the same way as the *in vitro* pre-clinical sample and, after lyophilisation and weighing, was found to
159 have a concentration of 158.48 mg/ml. For all samples, botanical identifications were carried out by
160 comparison with authentic herbarium reference material.

161 *Liquid chromatography–mass spectrometry (LC–MS) analysis of exudate sample*

162 Liquid chromatography-mass spectrometry (LC-MS) was carried out with a ThermoFisher Scientific
163 ‘Vanquish’ UHPLC system coupled to a Thermo Scientific ‘LTQ-Orbitrap XL’ hybrid linear ion trap-
164 orbitrap mass analyser fitted with an electrospray ionisation (ESI) source. Samples (5 µL) were injected
165 into a Phenomenex Luna C18 column (150 x 3 mm internal diameter, 3 µm particle size) and eluted at
166 0.4 ml/min and 30 °C using a linear gradient of MeOH, H₂O, and MeCN with 1% formic acid (0:90:10–
167 90:0:10 v/v over 60 min). MS¹ spectra at 30,000 resolution were recorded by Fourier transform mass
168 spectrometry (FTMS) in both positive and negative modes in the *m/z* range 125–1800. Simultaneously,
169 the linear ion-trap (ITMS) recorded low resolution MS¹, MS², and MS³ spectra (*m/z* 125-1800).

170 For quantification of ficuseptine in exudate samples, a sevenfold serial dilution of a pure ficuseptine
171 standard was prepared using a sample obtained previously (Prescott et al. 2015). A seven point dilution
172 series from 500µg/ml to 7µg/ml ficuseptine was prepared and then analysed by LC-MS along with a
173 five hundredfold dilution of the exudate samples. For the clinical trial exudate sample, a dilution was
174 prepared from the pure unfiltered unrefined sample. Quantification was achieved by measuring UV
175 peak area at 285nm and then interpolating the readings from exudate samples to the linear range of the
176 ficuseptine standard curve ($r^2=0.999$).

177 *Ames tests with Salmonella typhimurium TA98 and TA100 strains*

178 The 24-well plate bacterial reverse mutation test was performed following standard methods (Escobar
179 et al., 2013; Maron and Ames, 1983; Mortelmans and Zeiger, 2000). Briefly, 5 µL of solvent, positive
180 control or test item were added to wells of a 24-well plate containing minimal glucose agar. A mixture
181 of 5 µL of overnight bacterial culture, 25 µL of sodium phosphate buffer or S9 mix (Molecular
182 Toxicology Inc, USA), and 100 µl of molten 0.6% agar were maintained at approximately 50 °C and
183 supplemented immediately with 0.05 mM biotin and histidine. Once the agar had solidified, plates were
184 inverted and incubated for 3 days at 37 °C. After incubation, wells of all plates were examined both
185 macroscopically and microscopically for evidence of cytotoxicity or precipitates and any other effect
186 relevant to the interpretation of the test. Microscopic examination was used to check the condition of
187 the histidine-requiring micro-colonies that make up the ‘background lawns’. Revertant (his+) colony
188 numbers were scored manually.

189 *Antibacterial assays*

190 Bacterial strains: *S. aureus* (NCTC 6571), *S. dysgalactiae* (NCTC 13762), *S. pyogenes* (NCTC 8198),
191 *C. diphtheria* (NCTC 11397) and *F. ulcerans* (NCTC 12112) were grown at 37 °C. Logarithmic phase
192 cultures of *S. aureus*, *S. dysgalactiae* and *S. pyogenes* were grown aerobically in brain heart infusion
193 (BHI) broth (Sigma-Aldrich, Poole, UK). *C. diphtheria* was grown in BHI under microaerophilic
194 conditions (candle jar) and *F. ulcerans* was grown anaerobically in fastidious anaerobic broth (Lab M,
195 Heywood, UK).

196 The disc diffusion assays were carried out as described previously (Moses et al. 2020a, b). Positive
197 control discs contained 15 µL 10 % w/v povidone iodine (Vetasept) or 0.2 % chlorhexidine. The
198 minimum inhibitory concentration (MIC; lowest concentration of the test compounds to inhibit visible
199 microorganism growth after overnight incubation) and minimum bactericidal concentration (MBC;
200 lowest concentration of the test compounds to prevent microorganism growth after subculturing) were
201 performed as follows. Sterile filtered exudate samples were diluted in broth to obtain desired exudate
202 concentrations alongside serial dilutions of 0.2% w/v chlorhexidine, the starting dilution being 5% (100
203 µg/ml). The bacterial inoculum was adjusted for each organism, to yield a cell concentration of 5×10^4

204 colony forming units (CFU)/ml, and 95 μ L of inoculum was added to microtitre plate wells (Nunc,
205 Thermo Fisher Scientific, Loughborough, UK). A total volume of 5 μ L exudate solution was then added
206 to wells and incubated overnight. After overnight incubation, plates were examined for visible
207 microorganism growth and the lowest concentration of the test compound at which no visible bacterial
208 growth occurred was recorded as the MIC. Samples were then removed from each well, serially diluted
209 and plated out on Columbia blood agar plates for bacterial viability counting of *S. aureus*, *S.*
210 *dysgalactiae*, *S. pyogenes*, *C. diphtheria* or plated on fastidious anaerobic agar plates for *F. ulcerans*
211 enumeration (Southern Group Laboratory, Corby, UK). The MBC was recorded as the lowest
212 concentration of the test compound that prevented bacterial growth on subculturing.

213 *Isolation of neutrophils and macrophages from whole human blood*

214 Neutrophils and non-polarised macrophages were isolated from whole blood, based on the original
215 method of Moseley et al. (2003), as described recently (Moses et al. 2020a, b). Whole blood (in 20 ml
216 aliquots) was collected from healthy, human volunteers (age range 20-40 years).

217 *Pro-inflammatory cytokine release by human neutrophils and M1 macrophages*

218 Pro-inflammatory cytokine quantification was carried out, as described previously (Moses et al. 2020a,
219 b). Briefly, isolated neutrophil or macrophage cultures (5×10^5 cells/ml) were established in the absence
220 and presence of non-cytotoxic dilutions of *F. septica* exudate. Il-6 and TNF- α levels in cell culture
221 supernatants were determined using standard ELISA procedures. Experiments were performed on n=3
222 independent occasions, with data expressed as pg/ml.

223 *Dermal fibroblast and keratinocyte proliferation*

224 Primary dermal fibroblasts derived from normal skin were purchased from ATCC (Teddington, UK),
225 (HaCaTs) were obtained from the German Cancer Research Centre (Heidelberg, Germany). Cell
226 proliferation assays were carried out, as described previously with experiments performed on n=3
227 independent occasions, and data expressed as a % versus untreated controls (Moses et al., 2020).

228 *Clinical trial participants and study setting*

229 Following the *in vitro* assessment, we conducted a prospective, open label, cluster-randomized
230 controlled trial at three neighbouring schools with 150 students aged 9 to 18. The trial was carried out
231 in East New Britain, Papua New Guinea, between Nov 1, and Dec 15, 2019. Each school which
232 corresponded to a single cluster was randomly assigned to one of the three treatment arms after
233 screening by a physician.

234 Investigators subsequently assessed all children within the 9 to 18 age range in each school, starting
235 from the lowest grades. All children with skin lesions of less than 1 cm in diameter were assessed for
236 eligibility. Exclusion criteria were the need for antibiotic therapy or a known history of hypersensitivity

237 or allergy to the investigated product. Eligible children were subsequently offered the opportunity to
238 enrol in the study until the 50 designated spaces per treatment arm had been filled.

239 Children who agreed to enrol in the study provided verbal assent, and their parents or legal guardians
240 gave signed written consent for participation, according to standard protocols implemented and
241 described previously (Gonzalez-Beiras et al. 2017). The protocol was approved by the Research and
242 Ethics Committee of the School of Medicine and Health Sciences, University of Papua New Guinea
243 and is registered at ClinicalTrials.gov with Identifier: NCT04453124. The study followed the guidelines
244 of the Declaration of Helsinki and Tokyo for humans.

245 *Randomization and intervention*

246 Schools were randomly assigned to receive either *F. septica* exudate, chlorhexidine with cetrimide
247 cream (Savlon) antiseptic cream (Novartis Consumer Health, UK) or recommended standard care
248 (washing the ulcer with soap and water). All interventions were administered twice, at baseline (day 0)
249 and again at day 1, by the research team.

250 To apply the exudate, 50 μ L exudate was pipetted onto the tip of the index finger of a volunteer nurse
251 wearing surgical gloves. The exudate was then applied directly to the ulcer surface taking care to ensure
252 the exudate was evenly distributed on the ulcer. New gloves were used for each patient. The Savlon
253 cream (cetrimide 0.5%w/w and chlorhexidine gluconate 0.1% w/w) was applied in the same way, with
254 200mg of the cream applied to the glove. The standard care, soap and water wash was carried out by a
255 nurse according to the Papua New Guinea health guidelines using locally available sodium palmate
256 based “WasWas” brand of soap.

257 *Trial procedures*

258 Before the intervention, a physician carried out a structured assessment of the skin, arms, face, and legs
259 of each participant (baseline assessment). Information was recorded from each enrolled participant on
260 a case report form (CRF) specifically designed for the study. A *Treponema* DPP syphilis screen and
261 confirm assay (Chembio Diagnostic Systems) was carried out at baseline (pre-treatment) according to
262 manufacturer’s instructions as described previously (Ayove et al. 2014), along with a lesional swab,
263 taken using a sterile cotton swab and stored in DNA/RNA shield (Cambridge Biosciences). Patients
264 testing positive with the DPP assay and showing no clinical resolution after two weeks were treated
265 with azithromycin, as described previously (Mitja et al. 2014). The physician selected one skin lesion
266 that met the eligibility criteria and recorded its location in the CRF for further assessments. A
267 photograph of each lesion was taken at baseline (before treatment) and at days 7 and 14 post-treatment.
268 Adhesive labels printed with QR codes to identify each patient were applied next to the ulcers when
269 pictures were taken.

270 Photographs of each ulcer were blinded by digitally removing the identification code present in the
271 photograph. The photos were sent to three dermatologists, who graded the ulcers according to level of
272 improvement at 7 and 14 days relative to baseline into the following categories: worse, unchanged,
273 improved, and healed. Discordant grading was resolved between the three dermatologists using expert
274 opinion from co-author S.L.W, (London School of Hygiene and Tropical Medicine, UK).

275 *PCR of ulcer swabs*

276 DNA was extracted using the DNA Advance Magnetic Bead Extraction Kit (Beckman Coulter, UK)
277 following a modification of the manufacturer's protocol. Briefly, 200 μ L of extraction buffer comprising
278 173 μ L lysis buffer with 5 μ L of lysozyme 20 mg/ml, 7 μ L proteinase K (40 mg/ml) and 5 μ L 1 M
279 DTT, were added to each sample and incubated for 1 h at 56 °C. Samples were mixed by pipetting with
280 100 μ L of Bind1 Buffer, followed by 170 μ L of Bind2 Buffer. A magnet was applied to bind the beads
281 and the solution removed. Three washes of 340 μ L 80% ethanol were applied, each time removing the
282 tubes from the magnet and reapplying them to remove liquid. Samples were eluted into 100 μ L of
283 Elution Buffer. Samples were quantified by fluorimetry using the Qubit dsDNA High Sensitivity Kit
284 (Invitrogen, Carlsbad, USA) and also quantified on a Nanodrop spectrophotometer (GE Healthcare,
285 Amersham, UK) to provide an indication of DNA purity. A260/A280 ratios were >1.70. Primers were
286 designed and wet-lab validated for specificity, sensitivity, linearity over 7 log, and efficiency >95% by
287 qStandard (UK). For each target, a standard of 107 copies was prepared from purified PCR products
288 and serially diluted 10-fold to 10 copies. Primer sequences for the single copy Pol A gene of both
289 bacterial species were as follows; *T. pallidum* F: aatggtgtggtcgcgtttgat R: gaagagaaactccggaacaataag,
290 *H. ducrey*ii F: cgaatatcgcgcataaattgctga R: ttagtctgaggtaacgagtggaaca. Primers for human (YWHAZ)
291 gDNA were: F: acttttggtacattgtggcttca R: ccgccaggacaaaccagtat. For qPCR 1 ng DNA was used per
292 reactions and qPCR was performed as described previously (Lowe et al., 2019).

293 *Statistical analysis*

294 Results of the *in vitro* analyses were summarized as mean \pm standard error of the mean (SEM) of n=3
295 independent experiments. Statistical significance was determined by one-way analysis of variance
296 (ANOVA) with post-Dunnett's test for multiple comparisons. The significant threshold was set at an
297 alpha-error of 0.05.

298 Clinical data were described using frequencies and percentages of each category of lesion progression:
299 worse, unchanged, improved, and healed. The primary endpoint of the clinical assessment was the
300 percentage of participants with an improved/healed lesion within each study arm. We assessed the non-
301 inferiority of *F. septica* exudate vs. chlorhexidine and *F. septica* exudate vs. standard care (soap and
302 water) based the absolute risk difference between arms using the standard normal approximation to
303 obtain the confidence interval (CI) of the risk difference (Tang et al., 2014); non-inferiority was


304 considered when the lower boundary of the 95% CI of the risk difference was greater than -10
305 percentage units. All analyses were performed in R 3.6.3 and package dani (Quartagno 2020).

306 Results and discussion

307 LC-MS analysis reveals the alkaloid ficuseptine is a major component of the exudate

308 Previous phytochemical work on the leaves, roots, and stems of *F. septica* has yielded a variety of
309 phenanthroindolizidine alkaloids (Damu et al. 2005). Our previous work on the related species *Ficus*
310 *botryocarpa* identified the alkaloid ficuseptine as the principle active antibacterial compound
311 (Prescott et al. 2015). Our LC-MS analysis of *F. septica* exudate also showed ficuseptine as a major
312 component (Figure 1.). We also detected ficuseptine C, seco-dehydroantofine, and ficuseptine D
313 along with a variety of caffeoylgalactaric acid and sinapoylgalactaric acid isomers. Their
314 identifications were based on comparison with mass spectrometric and UV spectra data reported in
315 the literature (Baumgartner et al. 1990; Damu et al. 2005; Kubo et al. 2016; Ueda et al. 2009). Both
316 exudate samples presented the same major peaks and, in both cases, the antibacterial alkaloid
317 ficuseptine was the major peak. As ficuseptine has already been demonstrated to be an active
318 antibacterial principle we decided to quantify concentrations of the compound in both the clinical trial
319 exudate sample and the sample used in the pre-clinical investigations. Quantitation using a ficuseptine
320 standard revealed the sample for *in vitro* pre-clinical testing contained 53.9mg/ml ficuseptine whereas
321 the clinical trial sample (un-processed as used in the trial) contained 23.3mg/ml ficuseptine,
322 ($r^2=0.999$).

323



| Peak | Identification |
|------|-------------------------------------|
| 1-3 | caffeoylgalactaric acid or isomer |
| 4 | ficuseptamine C |
| 5 | caffeoylgalactaric acid or isomer |
| 6-9 | sinapoylgalactaric acid and isomers |
| 10 | seco-dehydroantofine |
| 11 | ficuseptine D |
| 12 | ficuseptine |

325

326 **Fig. 1.** Photometric Diode Array (PDA) chromatogram of *F. septica* exudate sample used in the
 327 clinical trial. Identity of peaks were confirmed based on their mass spectrometry fragments and UV
 328 spectra.

329 *F. septica* exudate is antibacterial towards wound pathogens known to colonise
 330 cutaneous ulcers in Papua New Guinea

331 Application of a topical antiseptic to the ulcer surface can reduce bacterial bio-burden and concomitant
 332 bacterial toxin production allowing wound progression and healing. We carried out disc diffusion assays
 333 and broth micro-dilution assays with non-fastidious bacterial pathogens that have been previously
 334 isolated or detected from cutaneous ulcers in Papua New Guinea (Montgomery 1985; Aadrians and
 335 Shah 1988; Noguera-Julian et al. 2019). The topical wound antiseptics chlorhexidine and povidone
 336 iodine were included as positive control agents. A disc diffusion assay was used as it measures the
 337 ability of a test substance to diffuse across an aqueous surface. This is conceptually similar to the *in*
 338 *vivo* application of the *F. septica* exudate to the ulcer surface, where diffusion across the aqueous wound
 339 milieu is important to ensure elimination of bacterial bio-burden. In the disc diffusion assay (Table 1.)
 340 *S. aureus* was the most sensitive organism. The exudate shows greater activity than a positive control
 341 comprising 0.2 % w/v chlorhexidine which is the concentration of chlorhexidine used in antiseptic
 342 formulations. The exudate is slightly less active than the 10 % w/v povidone iodine sample. In the broth
 343 microdilution assay the exudate was most active against *S. aureus* and *C. diphtheriae* and least active
 344 with *S. pyogenes* where full inhibition of growth was not observed. The two different exudate samples
 345 showed similar levels of activity in their MIC values but differed more in the MBC values (Table 2).
 346 For both exudate samples, the MIC and MBC values were well within the concentrations of plant
 347 exudate that would be present on the ulcer surface where the volume of exudate is generally greater
 348 than the volume of wound exudate present. This suggests the exudate could exert a bactericidal activity
 349 at the ulcer surface.

350

| | <i>S. aureus</i> | <i>S. pyogenes</i> | <i>F. ulcerans</i> |
|---------------------------|------------------|--------------------|--------------------|
| <i>F. septica</i> exudate | 5.9 | 5.5 | 2.2 |
| Chlorhexidine | 3.3 | 4.5 | 2.0 |
| Povidone iodine | 6.9 | 6.0 | 1.8 |

351

352 **Table 1.** Zones of inhibition (radius, in mm) with *S. aureus*, *S. pyogenes* and *F. ulcerans* in a disc
 353 diffusion assay. Each 6 mm paper disc was treated with 15 µL *F. septica* exudate (*in vitro* sample), 15
 354 µL 10 % w/v povidone iodine or 15 µL 0.2 % w/v chlorhexidine, before applying to agar plates seeded
 355 with *S. aureus* (NCTC 6571), *S. pyogenes* (NCTC 8198) and *F. ulcerans* (NCTC 12112). Povidone and
 356 chlorhexidine results were obtained as part of a multiple sample experiment carried out in parallel and
 357 reported previously (Moses et al. 2020a, b).

| | Exudate sample 1 MIC | Exudate sample 1 MBC | Exudate sample 2 MIC | Exudate sample 2 MBC | CHX MIC | CHX MBC |
|------------------------|-------------------------|-------------------------|-------------------------|-------------------------|------------|-----------|
| <i>C. diphtheriae</i> | 461 (0.31) | 921 (0.63) | 495 (0.31) | 991 (0.63) | 1 (0.04) | 3 (0.16) |
| <i>F. ulcerans</i> | 921 (0.63) | 7372 (5) | 991 (0.63) | 3962 (2.5) | 2 (0.08) | 6 (0.31) |
| <i>S. pyogenes</i> | - | - | - | - | 0.4 (0.02) | 0.4(0.02) |
| <i>S. aureus</i> | 230 (0.16) | 7372 (5) | 247 (0.16) | 991 (0.63) | 2 (0.08) | 13 (0.63) |
| <i>S. dysgalactiae</i> | 1843 (1.25) | 3686 (2.5) | 991 (0.63) | 7924 (5) | 2 (0.08) | 6 (0.31) |

358


359 **Table 2.** Minimum inhibitory concentrations and minimum bactericidal concentrations for two *F.*
360 *septica* exudate samples and the positive control chlorhexidine. Values outside brackets indicate MIC
361 and MBC values expressed as µg/ml. Values in brackets indicate MIC and MBCs as % v/v dilutions
362 where the undiluted exudate is 100% and the chlorhexidine (CHX) dilution is % v/v of a 0.2% w/v
363 stock solution. Exudate sample 1 is the sample used for pre-clinical *in vitro* testing, exudate sample 2
364 is the clinical trial sample. For *S. pyogenes* both exudate samples gave partial but incomplete
365 inhibition at the test concentrations used.

366

367 *The exudate is not mutagenic in Ames tests with Salmonella typhimurium TA98 and*
368 *TA100*

369 The exudate was tested for mutagenic activity in a bacterial reverse mutation assay. *S. typhimurium*
370 LT2 bacteria strains TA98 and TA100 were used as indicator organisms to detect frame shift and base
371 substitutions respectively. The assay uses standardised protocols described previously (Maron and
372 Ames 1983; Mortelmans and Zeiger 2000; Escobar et al. 2013). Rat liver S9 homogenate was used to
373 check for metabolic activation of the exudate, and the known mutagens sodium azide and 2-
374 aminoanthraene were used as positive controls. In accordance with standard procedures, a positive
375 response was considered when the minimum mean number of revertants per well increased by
376 threefold over the corresponding solvent control. The results (Figure 2.) show that while the known
377 mutagenic positive controls show increases above threefold for each strains as expected, with the
378 exudate there were no consistent increases in numbers of revertant colonies for either strain including
379 in the presence of S9 metabolic activation.

380




381 **Fig. 2.** The exudate does not induce mutations in the TA98 Salmonella strain (A) or the TA100 strain
 382 (B) when tested at concentrations up to 0.56mg/ml (3.703% v/v). Shaded bars indicate S9 metabolic
 383 activation, unshaded without S9 activation. Figures in mg/ml indicate concentration of *F. septica*
 384 exudate. 2AA (0.37µg/ml 2-aminoanthracene), 2NF (0.37µg/ml nitrofluorene), NaN₃ (0.19µg/ml
 385 sodium azide). Error bars indicate the standard error of the mean (SEM).

386 *The exudate stimulates proliferation of keratinocytes but not, dermal fibroblasts*

387 Both fibroblasts and keratinocytes play an important role in wound healing, with both cell types required
 388 for deposition of extracellular matrix components such as collagens, proteoglycans and hayaluronin
 389 (Moseley et al. 2004). However, fibroblasts present in chronic wounds exhibit reduced proliferative and
 390 migratory capabilities, due to the increased onset of cellular senescence within chronic wound
 391 environments, while keratinocytes also show reduced migratory responses (Wall et al. 2008; Pastar et
 392 al. 2014). We examined the effect of the exudate on fibroblasts and keratinocytes. Figures. 3A and 3B
 393 show that although the fibroblasts showed no significant increases in proliferation in response to the
 394 exudate (all $p > 0.05$), the keratinocytes exhibited significantly increased proliferation at 24 h post-
 395 treatment ($p < 0.05$ at 15ng/ml exudate).

396




397

398 **Fig. 3.** Exudate effects on dermal fibroblast proliferation (A) and keratinocyte proliferation (B) at 24 h
 399 (shaded) and 72 h (unshaded), relative to treatment with media only. Error bars indicate standard error
 400 of the mean (SEM) values, based on data analysis from n=3 independent experiments. *P* values
 401 * $p < 0.05$.

402 *The exudate completely down regulates cytokine expression from M1 macrophages*
 403 *and neutrophils*

404 We next looked at the effect of the exudate on the pro-inflammatory responses of M1 macrophages and
 405 neutrophils that are known to reside within chronic wounds. Excessive inflammatory responses from
 406 these cell types in chronic wounds is known to result in elevated levels of IL-1, IL-6, and TNF- α , which
 407 in turn results in upregulation of MMP expression and degradation of extracellular matrix (Barrientos et
 408 al. 2008; Moseley et al. 2004). Primary cells obtained from whole human blood were used as they

409 closely resemble the status of these cells *in vivo*. For the neutrophils, cells treated with and without LPS
 410 were included as controls to show the extent of the inflammatory response. For macrophages, control
 411 cells treated with or without GM-CSF were included to show polarisation occurred as expected. Figure
 412 4A-4D shows that for both neutrophils and M1 macrophages, the exudate potently downregulates IL-6
 413 or TNF- α secretion. The neutrophils are more sensitive to the exudate than the macrophages, with
 414 complete knock down of IL-6 or TNF- α levels occurring at 1.5 μ g/ml exudate (0.001% v/v) compared
 415 with 15 μ g/ml for IL-6 from macrophages. Interestingly, very low amounts of the exudate 0.15 μ g/ml
 416 caused a significant increase in IL-6 expression from the macrophages, suggesting the exudate may
 417 elicit both pro-inflammatory and anti-inflammatory effects depending on the concentrations used.




418
 419 **Fig. 4.** Top two graphs show the effect of the exudate on IL-6 (A) and TNF- α (B) secretion by M1
 420 macrophages. All conditions included stimulation with GM-CSF, apart from the -GM-CSF controls as
 421 indicated. Controls -GM-CSF and +GM-CSF show responses of pre-macrophages and polarised M1
 422 macrophages respectively, each in the absence of exudate. Concentrations indicated in μ g/ml are GM-
 423 CSF-stimulated cells treated with *F. septica* exudate at the specified concentrations. The bottom two
 424 graphs show the effect of the exudate on IL-6 (C) and TNF- α (D) secretion by neutrophils. All
 425 conditions included stimulation with LPS, apart from the -LPS control as indicated. Controls -LPS and
 426 +LPS indicate cell responses with and without LPS, in the absence of exudate. Bars with concentrations
 427 indicated in μ g/ml are LPS-stimulated cells with *F. septica* exudate at specified concentrations. For all
 428 graphs, error bars indicate standard error of the mean (SEM) values, based on data analysis from n=3
 429 independent experiments. *P* values **p*<0.05, ***p*<0.01, ****p*<0.005.

430 *A clinical trial of the exudate shows similar levels of healing to chlorhexidine and*
 431 *cetrimide cream*

432 The clinical assessment included 150 children aged 9 to 18 randomized at school level to treat skin
 433 ulcers with a standard care consisting of soap and water (control), *F. septica* exudate, or chlorhexidine
 434 with cetrimide cream (Savlon) antiseptic cream. This age range was chosen as the ulcers are particularly
 435 prevalent in children. All participants received two treatment rounds; 134 attended at least one follow-
 436 up visit and were therefore included in the non-inferiority analysis (Figure 5).

437




438

439 **Fig. 5.** A consort flow diagram of the clinical trial. The three treatment arms were as follows:
 440 recommended standard care (washing with soap and water), chlorhexidine and cetrimide cream and *F.*
 441 *septica* exudate. The clinical assessment was carried out on blinded photographs of ulcers. ENB: East
 442 New Britain

443


444 Figure 6 shows the result of the clinical assessment of skin lesions at days 7 and 14. The assessment
 445 was performed by expert dermatologists using blinded ulcer photographs. The dermatologists
 446 categorized the ulcers into one of four healing states, namely: worse, unchanged, improved, and healed.
 447 On both day 7 and 14, most participants had either improved or healed skin lesions, irrespective of the
 448 type of treatment (Figure 6). Accordingly, the risk difference estimate revealed no significant
 449 differences between treatments in the likelihood of improvement/healing at day 7 or 14 (Figure 7).

450 According to the pre-set noninferiority margin, *F. septica* exudate was non-inferior to chlorhexidine
 451 cream or standard care with water and soap at day 14; the noninferiority criterion was not met at day 7.
 452



453
 454 **Fig. 6.** Proportion of participants within each of the categories of lesion progression (i.e., worse,
 455 unchanged, improved, and healed) at each assessment. Results are presented for each treatment arm and
 456 assessments at days 7 and 14.

457
 458
 459



460

461 **Fig. 7.** Absolute risk difference for improvement/healing between treatment with *F. septica* exudate
 462 and the two comparators: chlorhexidine with cetrimide cream (Savlon) antiseptic cream and
 463 recommended standard care (washing the ulcer with soap and water). Dots and horizontal lines represent
 464 the absolute risk difference and the 95% confidence interval (CI). The dashed line shows the non-
 465 inferiority margin for the lower boundary of the 95% CI (-10 percentage points).

466

467 *H. ducreyi* and *T. pallidum* subspecies *pertenue*, are two bacterial species that have been detected in
 468 previous work on larger yaws-like ulcers in the South Pacific (Noguera-Julian et al. 2019). PCR
 469 diagnostics were carried out on wound swabs taken from each patient to establish if these bacterial
 470 species were also present in the smaller, less than 1cm diameter ulcers under investigation in this study.
 471 Participants in the three study arms showed a high proportion of *H. ducreyi* and a low proportion of *T.*
 472 *pallidum* subspecies *pertenue*, which is the causative agent of yaws disease (Table 4). We also used a
 473 DPP finger prick DPP syphilis screen and confirm assay, to detect active yaws cases, similarly revealing
 474 a low proportion in each treatment arm.

475

| | <i>F. septica</i> exudate | CHX/cetrimide | Soap and water |
|--------------------------|---------------------------|---------------|----------------|
| <i>H. ducreyi</i> (PCR) | 92% | 98% | 100% |
| <i>T. pallidum</i> (PCR) | 0% | 4% | 4% |
| <i>T. pallidum</i> DPP | 0% | 4% | 6% |

476

477 **Table 4.** Proportion of patients in each treatment arm with PCR positive ulcers for *H. ducreyi*, *T.*
 478 *pallidum* subspecies *pertenue* as well as proportions of patients with active yaws infection determined
 479 with a DPP syphilis screen and confirm assay.

480 **Conclusions**

481 This is the first study to investigate the potential of *F. septica* as a first-line treatment for cutaneous
 482 ulcers and is the first clinical trial study of a plant medicine for infected cutaneous ulcers of the type
 483 commonly found in the South Pacific. The *in vitro* results presented here demonstrate that *F. septica*
 484 exudate exhibits bactericidal activity against both Gram-positive and Gram-negative bacterial
 485 pathogens that are known to infect ulcers in Papua New Guinea. The exudate also showed strong
 486 inhibition of pro-inflammatory responses from neutrophils and M1 macrophages, and it was non-
 487 mutagenic when tested in Ames assays looking for frameshift and base substitution mutations. Chemical
 488 analysis of the exudate revealed the presence of a variety of different alkaloids, but with the known
 489 antibacterial alkaloid ficuseptine the dominant major component. Thus, from the *in vitro* data, *F. septica*
 490 exudate exhibits potentially useful traits for a topical treatment for infected cutaneous ulcers of the type
 491 commonly encountered in Papua New Guinea. Furthermore, *F. septica* is a common plant species and

492 is often found growing in close proximity to villages in Papua New Guinea. The physical characteristics
493 of the exudate are well suited to its traditional use as a topical treatment for cutaneous ulcers. Its
494 sufficiently fluid to transfer from the tree to the ulcer but still viscous enough to adhere to the ulcers.
495 Highly potent stimulation of keratinocytes was also observed but caution should be applied when
496 considering this effect in an *in vivo* context as the topical application of the plant exudate would result
497 in much higher concentrations of plant exudate at the ulcer surface and the *in vitro* results do not show
498 stimulation of keratinocytes at higher concentrations.

499 Next, we conducted a cluster-randomized trial to investigate the healing effect of *F. septica* exudate on
500 small skin lesions. To assess the microbiology of the ulcers, PCR diagnostics were carried out on lesion
501 swabs, revealing the presence of both *H. ducreyi* and *T. pallidum* subspecies *pertenue*. Of the two
502 species, *H. ducreyi* is by far the most prevalent, with detection rates ranging from 92% to 100% in the
503 three treatment arms. Although previous work has applied metagenomics and PCR to large, yaws-like
504 ulcers (Mitja et al. 2014), this is the first study to specifically examine small ulcers, less than 1cm
505 diameter. The results provide further confirmation of the association of *H. ducreyi* with infected
506 cutaneous ulcers in the South Pacific and suggest that *H. ducreyi* infection occurs early on in small
507 ulcers rather than being acquired later on as a secondary infection of larger ulcers. The results also
508 demonstrate the potential for small cutaneous ulcers to act as a gateway for yaws disease by providing
509 a route into the body to establish a systemic infection.

510 The clinical trial compared healing outcomes of the exudate with chlorhexidine antiseptic cream and
511 the standard treatment (soap and water) which is recommended in Papua New Guinea treatment
512 guidelines. The analysis showed no significant differences between treatments. More interestingly, *F.*
513 *septica* exudate was non-inferior to both chlorhexidine cream or soap and water at day 14. Our findings,
514 which should be confirmed in a larger clinical trial, have important public health implications in
515 preventing complications of skin ulcers. Although chlorhexidine cream is available in pharmacies in
516 Papua New Guinea, a substantial section of the Papua New Guinean population lack ready access to
517 pharmacies, aid posts or clinics. Furthermore, even soap and water treatment using clean water is not
518 readily available to patients living in remoter areas of the country. Therefore, educational programs for
519 self-care of small skin lesions using *F. septica* exudate, might form part of a low-cost health intervention
520 for treating ulcers in remote areas of Papua New Guinea. Larger clinical trials with an adequate sample
521 size are warranted to permit the inclusion of *F. septica* treatment into clinical guidelines and public
522 health strategies.

523 **Author contributions**

524 John Deli (Investigation), Camila González-Beiras (Project administration; Investigation), Georgia S.
525 Guldan (Supervision), Ryan Moseley (Supervision), Rachael L. Moses (Investigation), Jordanna

526 Dally (Investigation), Fionnuala T. Lundy (Formal analysis; Supervision), Marc Corbacho-Monne
527 (Formal analysis), Stephen L Walker (Formal analysis), Maria Ubals (Formal analysis), Dan Ouchi
528 (Formal analysis) Rui Fang (Formal analysis), Marie Briggs (Formal analysis) Robert Kiapranis
529 (Project administration), Martha Yahimbu (Project administration), Oriol Mitjà (Conceptualization;
530 Supervision), Thomas A.K. Prescott (Conceptualization; Investigation; Funding acquisition)

531 **Conflict of interest**

532 The authors declare no conflict of interest. The Savlon antiseptic cream used in this study was purchased
533 from a high street pharmacy and the manufacturer did not provide funding for this study. All data were
534 generated in-house, and no paper mill was used. All authors agree to be accountable for all aspects of
535 work ensuring integrity and accuracy

536 **Funding**

537 This work was funded by a grant from the Christensen Fund for Melanesia 2016-8040.

538 **Acknowledgements**

539 We like to acknowledge the help provided by members of the Umbi village in New Britain in providing
540 crucial ethnobotanical advice, and to local healthcare workers from Rabaul Hospital for assistance in
541 the clinical trial. We are also grateful to Lydia Sahami for help with the clinical trial, Dr Patricia de
542 Winter for assistance with qPCR, Catherine Fulton for *in vitro* antibacterial testing and Gerard Carot-
543 Sans for editing the manuscript.

544

545 **References**

- 546 Aadrians, B., Shah, H., 1988. *Fusobacterium ulcerans* sp. nov. from tropical ulcers. *Int. J. Syst.*
547 *Bacteriol.*, 38, 447–448.
- 548 Ayove, T., Houniei, W., Wangnapi, R., Bieb, S. V, Kazadi, W., Luke, L.N., Manineng, C., Moses, P.,
549 Paru, R., Esfandiari, J., Alonso, P.L., de Lazzari, E., Bassat, Q., Mabey, D., Mitja, O., 2014.
550 Sensitivity and specificity of a rapid point-of-care test for active yaws: a comparative study.
551 *Lancet Glob. Heal.* 2, e415-21. doi:10.1016/S2214-109X(14)70231-1
- 552 Barrientos, S., Stojadinovc, O., Golinko, M.S., Brem, H., Tomic-Canic, M., 2008. Growth factors and
553 cytokines in wound healing. *Wound. Repair. Regen.* 16, 585–601. doi:10.1111/j.1524-
554 475X.2008.00410.x
- 555 Baumgartner, B., Erdelmeier, C.A., Wright, A.D., Rali, T., Sticher, O., 1990. An antimicrobial alkaloid
556 from *Ficus septica*. *Phytochemistry* 29, 3327-3330.
- 557 Berg, C. C., Corner, E. J. H., Nooteboom, H.P., 2005. *Flora Malesiana. Series I, Seed plants.* Volume
558 17, Part 2: Moraceae (*Ficus*) 730 pp. Nationaal Herbarium Nederland (Leiden, Netherlands).
- 559 Damu, A.G., Kuo, P.C., Shi, L.S., Li, C.Y., Kuoh, C.S., Wu, P.L., Wu, T.S., 2005.
560 Phenanthroindolizidine alkaloids from the stems of *Ficus septica*. *J. Nat. Prod.* 68, 1071–1075.
561 doi:10.1021/np050095o
- 562 Damu, A.G., Kuo, P.C., Shi, L.S., Li, C.Y., Su, C.R., Wu, T.S., 2009. Cytotoxic phenanthroindolizidine
563 alkaloids from the roots of *Ficus septica*. *Planta Med.* 75, 1152–1156. doi:10.1055/s-0029-
564 1185483
- 565 Eming, S.A., Krieg, T., Davidson, J.M., 2007. Inflammation in wound repair: molecular and cellular
566 mechanisms. *J. Invest. Dermatol.* 127, 514–525. doi:10.1038/sj.jid.5700701
- 567 Escobar, P.A., Kemper, R.A., Tarca, J., Nicolette, J., Kenyon, M., Glowienke, S., Sawant, S.G.,
568 Christensen, J., Johnson, T.E., McKnight, C., Ward, G., Galloway, S.M., Custer, L., Gocke, E.,
569 O'Donovan, M.R., Braun, K., Snyder, R.D., Mahadevan, B., 2013. Bacterial mutagenicity
570 screening in the pharmaceutical industry. *Mutat. Res.* 752, 99–118.
571 doi:10.1016/j.mrrev.2012.12.002
- 572 Gonzalez-Beiras, C., Kapa, A., Vall-Mayans, M., Paru, R., Gavilan, S., Houine, W., Bieb, S., Sanz, S.,
573 Martins, R., Mitja, O., 2017. Single-dose azithromycin for the treatment of *Haemophilus ducreyi*
574 skin ulcers in Papua New Guinea. *Clin. Infect. Dis.* 65 (12): 2085-2090.
- 575 Griesenauer, B., González-Beiras, C., Fortney, K.R., Lin, H., Gao, X., Godornes, C., Nelson, D.E.,
576 Katz, B.P., Lukehart, S.A., Mitjà, O., Dong, Q., Spinola, S.M., 2021. *Streptococcus pyogenes* is

577 associated with idiopathic cutaneous ulcers in children on a yaws-endemic island. MBio. 12, 1–
578 16. doi:10.1128/mBio.03162-20

579 Paediatric Society of Papua New Guinea., 2016. Standard treatment for common illnesses of children
580 in Papua New Guinea: a manual for nurses, health extension officers and doctors. Port Moresby:
581 PNG Department of Health.

582 Kubo, M., Yatsuzuka, W., Matsushima, S., Harada, K., Inoue, Y., Miyamoto, H., Matsumoto, M.,
583 Fukuyama, Y., 2016. Antimalarial Phenanthroindolizine Alkaloids from *Ficus septica*. Chem.
584 Pharm. Bull.. 64, 957–960. doi:10.1248/cpb.c16-00181

585 Lupi, O., Madkan, V., Tying, S.K., 2006. Tropical dermatology: bacterial tropical diseases. J. Am.
586 Acad. Dermatol. 54, 559–580. doi:10.1016/j.jaad.2005.03.066

587 Maron, D.M., Ames, B.N., 1983. Revised methods for the Salmonella mutagenicity test. Mutat. Res.
588 113, 173–215. doi:10.1016/0165-1161(83)90010-9

589 Mitja, O., Lukehart, S.A., Pokowas, G., Moses, P., Kapa, A., Godornes, C., Robson, J., Cherian, S.,
590 Houinei, W., Kazadi, W., Siba, P., de Lazzari, E., Bassat, Q., 2014. *Haemophilus ducreyi* as a
591 cause of skin ulcers in children from a yaws-endemic area of Papua New Guinea: a prospective
592 cohort study. Lancet Glob. Heal. 2, e235-41. doi:10.1016/S2214-109X(14)70019-1

593 Montgomery, J., 1985. The aerobic bacteriology of infected skin lesions in children of the Eastern
594 Highlands Province. PNG Med. J. 28, 93–103.

595 Mortelmans, K., Zeiger, E., 2000. The Ames Salmonella/microsome mutagenicity assay. Mutat. Res.,
596 455, 29–60. doi:10.1016/s0027-5107(00)00064-6

597 Moseley, R., Stewart, J.E., Stephens, P., Waddington, R.J., Thomas, D.W., 2004. Extracellular matrix
598 metabolites as potential biomarkers of disease activity in wound fluid: lessons learned from other
599 inflammatory diseases? Br. J. Dermatol. 150, 401–413. doi:10.1111/j.1365-2133.2004.05845.x

600 Moseley, R., Walker, M., Waddington, R.J., Chen, W.Y., 2003. Comparison of the antioxidant
601 properties of wound dressing materials--carboxymethylcellulose, hyaluronan benzyl ester and
602 hyaluronan, towards polymorphonuclear leukocyte-derived reactive oxygen species.
603 Biomaterials. 24, 1549–1557. doi:10.1016/s0142-9612(02)00540-9

604 Moses, R.L., Dally, J., Lundy, F.T., Langat, M., Kiapranis, R., Tsolaki, A.G., Moseley, R., Prescott,
605 T.A.K., 2020a. *Lepiniopsis ternatensis* sap stimulates fibroblast proliferation and down regulates
606 macrophage TNF-alpha secretion. Fitoterapia. 141, 104478. doi:10.1016/j.fitote.2020.104478

607 Moses, R.L., Fang, R., Dally, J., Briggs, M., Lundy, F.T., Kiapranis, R., Moseley, R., Prescott, T.A.K.,
608 2020b. Evaluation of *Cypholophus macrocephalus* sap as a treatment for infected cutaneous ulcers

609 in Papua New Guinea. *Fitoterapia*. 143, 104554. doi:10.1016/j.fitote.2020.104554

610 Noguera-Julian, M., Gonzalez-Beiras, C., Parera, M., Ubals, M., Kapa, A., Paredes, R., Mitja, O., 2019.
611 Etiological characterization of the cutaneous ulcer syndrome in Papua New Guinea using shotgun
612 metagenomics. *Clin. Infect. Dis.* 68, 482–489. doi:10.1093/cid/ciy502

613 Pastar, I., Stojadinovic, O., Yin, N.C., Ramirez, H., Nusbaum, A.G., Sawaya, A., Patel, S.B., Khalid,
614 L., Isseroff, R.R., Tomic-Canic, M., 2014. Epithelialization in wound healing: A comprehensive
615 review. *Adv. Wound Care*. 3, 445–464. doi:10.1089/wound.2013.0473

616 Prescott, T.A., Briggs, M., Kiapranis, R., Simmonds, M.S., 2015. Medicinal plants of Papua New
617 Guinea’s Miu speaking population and a focus on their use of plant-slaked lime mixtures. *J.*
618 *Ethnopharmacol.* 174, 217–223. doi:10.1016/j.jep.2015.08.019

619 Prescott, T. A., Kiapranis, R., and Maciver, S.K., 2012. Comparative ethnobotany and in-the-field
620 antibacterial testing of medicinal plants used by the Bulu and inland Kaulong of Papua New
621 Guinea. *J. Ethnopharmacol.* 139 (2): 497-503.

622 Quartagno, M., 2020. Package ‘dani’: Design and Analysis of Non-Inferiority Trials. [https://cran.r-](https://cran.r-project.org/package=dani)
623 [project.org/package=dani](https://cran.r-project.org/package=dani) (accessed 4.2.21).

624 Schmidt Mueller, V., Rosenbach, G., E., 2020. Rural households in Papua New Guinea afford better
625 diets with income from small businesses. *Food Policy*. 97, 101964.

626 Ueda, J.Y., Takagi, M., Shin-ya, K., 2009. Aminocaprophenone- and pyrrolidine-type alkaloids from
627 the leaves of *Ficus septica*. *J. Nat. Prod.* 72, 2181–2183. doi:10.1021/np900580f

628 Wall, I.B., Moseley, R., Baird, D.M., Kipling, D., Giles, P., Laffafian, I., Price, P.E., Thomas, D.W.,
629 Stephens, P., 2008. Fibroblast dysfunction is a key factor in the non-healing of chronic venous leg
630 ulcers. *J. Invest. Dermatol.* 128, 2526–2540. doi:10.1038/jid.2008.114

631 Wu, P., Rao, K.V., Su, C., Kuoh, C., Wu, T., 2002. Phenanthroindolizidine alkaloids and their
632 cytotoxicity from the leaves of *Ficus septica*. *Heterocycles*. 57, 2401–2408.

633 Yang, C.W., Chen, W.L., Wu, P.L., Tseng, H.Y., Lee, S.J., 2006. Anti-inflammatory mechanisms of
634 phenanthroindolizidine alkaloids. *Mol. Pharmacol.* 69, 749–758. doi:10.1124/mol.105.017764

635 Zhao, R., Liang, H., Clarke, E., Jackson, C., Xue, M., 2016. Inflammation in chronic wounds. *Int. J.*
636 *Mol. Sci.* 17. doi:10.3390/ijms17122085

637

638