

Additionally, only 51% agreed that they felt able to refer a patient with diabetes to the most appropriate diabetes service based on type of diabetes and medication prescribed. This highlights an important issue, as cohesive shared-care and clear referral pathways are key when considering effective diabetes management. **Conclusion.** Psychiatric inpatient admission could be used opportunistically to improve the healthcare disparities for people with comorbid diabetes and SMI. This national survey highlights key areas that would need to be addressed to standardise and optimise diabetes care in this setting. This includes appropriate training, clear guidelines and cohesive shared-care pathways.

The Effectiveness of Mindfulness-Based Interventions for Anxiety Disorders in Adults: A Systematic Narrative Review

Ms Lisa Harrison*, Dr Katja Umla-Runge and Dr Athanasios Hassoulas

Cardiff University, Cardiff, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2022.197

Aims. In recent years there has been accelerated clinical interest in Mindfulness based interventions (MBI's) leading to an upswell in research due the impact of its wide clinical application. Mindfulness Based Cognitive Therapy (MBCT) and Acceptance and Commitment Therapy (ACT) have recently been investigated for the beneficial treatment of anxiety-based disorders in adults. The aim of the current review was to appraise and synthesise findings of studies published within the last decade, in determining the efficacy of MBCT and ACT in treating anxiety disorders in adults, given gaps identified in the existing literature.

Methods. Scoping searches were conducted using MEDLINE, PsycINFO, Emcare, and Cochrane databases. The Synthesis Without Meta-analysis protocol (SWiM) was adopted for this review, in evaluating the efficacy of MBCT and ACT for anxiety disorders in adults. The review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Standards.

Results. The results of this review suggest that MBCT and ACT are effective therapeutic modalities in improving anxiety in adult populations. The results are, however, tentative. Whilst both MBI's show promise in the treatment of anxiety disorders, with the paucity of existing systematic reviews and methodological flaws within overall primary study design, the results should be interpreted with caution.

Conclusion. The overall therapeutic effectiveness of MBI's has been assessed and the general data support its efficacy. However, a judicious approach is required as results continue to remain inconclusive grounded in the totality of the evidence.

The current review revealed the ongoing methodological concerns encountered in determining the comparative effectiveness of MBCT and ACT for anxiety disorders in adults. Due to the current limited number of comparative studies of mindfulness based with mindfulness informed interventions, it could be suggested that a lack of systematic research is slowly influencing a collective understanding of MBI's being a homogenous group of treatments. The lack of delineation can have an impact on research, clinical practice and policy making. Further high quality research is required to continue to bridge the science practice gap. Without depth of understandings associated with the mechanisms of change and the impact that contextual aspects have on the outcome effectiveness, there are significant implications for practice and patient care. It is of importance that the adaptation and

subsequent developments in clinical practice do not outpace the research base to fully understand the mechanisms that make each MBI effective, for which population and diagnoses.

A Narrative Review of Defensive Medical Practice in Psychiatry

Dr Khalil Hassanally*

CNWL, London, United Kingdom. University of Bristol, Bristol, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2022.198

Aims. Defensive medical practice has become an increasingly global phenomenon and encompasses all medical specialties. In the UK it was defined in the case of *Sidaway v Board of Governors of the Bethlem Royal Hospital* [1985] UKHL 1 (21 February 1985) as “the practice of doctors advising and undertaking the treatment which they think is legally safe even though they may believe that it is not the best for their patient”. This narrative review surveys the literature to establish the forms in which defensive practice may manifest itself within psychiatry.

Methods. In this narrative review, various terms pertaining to defensive medical practice in psychiatry were searched in both medical and legal databases.

Results. Though the literature in psychiatry compared to other medical specialties is more limited, some common themes occur across all jurisdictions surveyed. Defensive psychiatric practice included admitting the patient even though they may be managed within the community (as reported by 21% of psychiatrists surveyed in the North of England) and employing more coercive practice, either using the mental health legislation or implied or actual threats. Once hospitalised, defensive practice manifests itself by placing patients on higher levels of nursing observations than necessary.

Across inpatient and outpatient settings between one and two thirds of psychiatrists reported altering the way they document to attend to medicolegal considerations. Prescribing habits were also altered due to fears of litigation; an Israeli study found that almost half of psychiatrists surveyed reported they prescribed smaller doses of medication than what they felt was required to pregnant woman and ninety percent reported the same when it came to the treatment of elderly patients.

When looked at by seniority it was felt that junior doctors were more prone to admitting patients defensively than consultants. In this respect, psychiatry differs from most other medical specialties as, in general, the evidence suggests that increased seniority is more likely to lead to admission.

Conclusion. Defensive practice in psychiatry appears to be widespread and takes a number of different forms. However, the research in psychiatry is limited and does not explore key areas common to other medical specialties such as clinician avoidance of certain cases or increased use of diagnostic tests. Furthermore, there is little examination of how psychiatrists may utilise mental health legislation within their defensive practice.

Breast Feeding Experiences of NHS Staff Returning to Work From Maternity Leave: A National Study

Dr Hollie Hearfield^{1*}, Dr Jennie Collier² and Dr Fauzia Paize³

¹Ancora House, Chester, United Kingdom; ²Wirral CAMHS, Wirral, United Kingdom and ³Liverpool Women's Hospital, Liverpool, United Kingdom

*Presenting author.