CARDIFF UNIVERSITY PRIFYSGOL CAERDYD

ORCA – Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:https://orca.cardiff.ac.uk/id/eprint/150866/

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Bryan, Paul, Ivaki, Mohammad and Scheuer, Julian 2022. On the classification of ancient solutions to curvature flows on the sphere. Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 25 (1), pp. 53-76. 10.2422/2036-2145.202105_006

Publishers page: https://doi.org/10.2422/2036-2145.202105_006

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See http://orca.cf.ac.uk/policies.html for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



ON THE CLASSIFICATION OF ANCIENT SOLUTIONS TO CURVATURE FLOWS ON THE SPHERE

PAUL BRYAN, MOHAMMAD N. IVAKI, AND JULIAN SCHEUER

ABSTRACT. We consider the evolution of hypersurfaces on the unit sphere S^{n+1} by smooth functions of the Weingarten map. We introduce the notion of 'quasi-ancient' solutions for flows that do not admit non-trivial, convex, ancient solutions. Such solutions are somewhat analogous to ancient solutions for flows such as the mean curvature flow, or 1-homogeneous flows. The techniques presented here allow us to prove that any convex, quasi-ancient solution of a curvature flow which satisfies a backwards in time uniform bound on mean curvature must be stationary or a family of shrinking geodesic spheres. The main tools are geometric, employing the maximum principle, a rigidity result in the sphere and an Aleksandrov reflection argument. We emphasize that no homogeneity or convexity/concavity restrictions are placed on the speed, though we do also offer a short classification proof for several such restricted cases.

1. INTRODUCTION

We consider the evolution of a closed, convex hypersurfaces M^n by

(1.1)
$$\partial_t x = -F\nu, \ x \colon (-T,0) \times M^n \to \mathbb{S}^{n+1},$$

where \mathbb{S}^{n+1} is equipped with the round metric of constant curvature 1, ν is the outward unit normal to $M_t = x(t, M^n)$ and F is a function of the principal curvatures of M_t . Whenever we refer to (1.1), F will be understood to have the following properties without further mentioning.

1.1. Assumption. Let $\Gamma_+ = \{(\kappa_i) \in \mathbb{R}^n : \kappa_i > 0, i = 1, \dots, n\}$. Assume that $F \in C^{\infty}(\Gamma_+) \cap C^0(\overline{\Gamma}_+)$ is

- (i) symmetric,
- (ii) positive on Γ_+ with F(0) = 0 and
- (iii) increasing, i.e., for all $1 \le i \le n$ we have $\frac{\partial F}{\partial \kappa_i} \ge 0$.

We emphasize that no homogeneity, convexity or concavity restrictions are placed on the speed - all we require is that the speed is parabolic, positive and symmetric (equivalently, the speed is invariant under isometries). It *shrinks* strictly convex hypersurfaces. The assumption F(0) = 0 ensures that equators are static solutions and that the choice of an *outward* unit normal makes sense for all nontrivial solutions.

We consider quasi-ancient, convex solutions of (1.1). Here quasi-ancient refers to solutions existing on the same time interval (-T, 0) as the maximal flow of strictly convex geodesic spheres, compare Definition 3.3. For many common flows, such as

Date: June 29, 2022.

²⁰¹⁰ Mathematics Subject Classification. 53C44, 58J35, 35E10.

the mean curvature flow $F(\mathcal{W}) = \text{Trace}(\mathcal{W}) = H$, the maximal interval of a flow of geodesic spheres is the infinite time interval $(-\infty, 0)$. However, for flows $F = H^p$ with $p \in (0, 1)$, the maximal time interval is infact bounded; see Section 3.

The following theorem is the main result of the paper.

1.2. **Theorem.** Let $n \ge 1$ and let M_t be a smooth, quasi-ancient, convex solution of (1.1) where F satisfies Assumption 1.1 and such that

$$\limsup_{t \to -T} \max_{M_t} H < \infty.$$

Suppose that either

(i) M_t is strictly convex for all $t \in (-T, 0)$, or (ii) $F \in C^1(\overline{\Gamma}_+)$. Then M_t is a flow of geodesic spheres.

In certain situations, such as 1-homogeneous speeds that are either convex, or concave and comparable to H, the theorem is quite easy to prove, and we offer a short proof in these cases in Section 4 without assuming a bound on the curvature. For instance, this should be compared with the result in [17, Theorem 6.1] where a short proof in the case of the mean curvature flow in the sphere may be found. The main idea is to find the correct quantity to which the maximum principle may be applied. In the paper referred to it is the pinching quantity $(|A|^2 - nH^2)/H^2$ used so often for the mean curvature flow. Here we use W/F but also remark that the curvature ball estimates from [1] may also be used.

The main thrust of the theorem is that we have very strong rigidity in the sphere, and we can prove the result in great generality - *all we require is weak parabolicity* of the flow. At this level of generality, it is quite difficult to obtain suitable uniform regularity estimates, in particular, Evans-Krylov higher regularity estimates are not known for arbitrary non-linear equations, and hence less direct PDE methods are required. We obtain the theorem by first using the rigidity results of [21] to argue that quasi-ancient solutions with bounded H must limit to an equator backwards in time. Then the Aleksandrov reflection technique developed in [8, 10] applies to show the symmetry is preserved under the flow, completing the classification.

Let us remark that one may seek other methods of classification, using more PDE theoretic techniques, but they tend to suffer the drawback that they only apply (without modification) to speeds F that are non-singular and non-degenerate on the equator. The methods we describe here apply even in the case of *singular* or degenerate speeds. Very simple examples of such speeds are H^p , which are singular for 0 and degenerate for <math>p > 1 whenever H = 0, in particular along the equator. We find that the geometric methods are quite appealing and of interest in their own right, even in cases where PDE methods are applicable, these methods offering a powerful, complementary alternative to direct PDE techniques, and supplanting them in cases where such methods are unknown.

One limitation of our method is that at this stage, we assume a bound on the mean curvature H. Although there are certainly solutions emanating from convex polyhedra (and hence with unbounded H), it is not at all clear to us whether such solutions can be quasi-ancient. In Section 7, we give two examples (for non-geometric flows) showing what could go wrong. The first converges to a lune (intersection of hemispheres), with unbounded H and the second converges to an equator, but is not a family of shrinking spheres. Since these examples are not geometric flows,

they are not of the type considered here, but they do show that parabolic flows on the sphere may admit ancient solutions that are not a family of shrinking geodesics spheres.

It is also worth comparing the techniques and results here to the Euclidean case. The latter does not exhibit such strong rigidity, and other ancient, convex solutions may occur. Indeed, for the curve shortening flow in the plane, the classification [11] states that solutions are either shrinking circles or the Angenent oval (also known as the paper clip solution) [2]. In higher dimensions a characterization of when ancient, convex mean curvature flows are shrinking spheres have been given [14, 17], and one generally expects a greater variety of ancient, convex solutions to exist.

Convex, ancient solutions are of interest as they arise as rescalings of singularities, cf. [16], of mean convex, mean curvature flow, cf. [15, 26]. Singularity models for mean curvature flow in the sphere have been studied in [22], where mean convex singularities with an ambient curvature dependent pinching condition are classified according to a blow-up yielding ancient solutions in Euclidean space.

Finally, we want to mention that after the first version of this paper appeared on arXiv.org in 2016, many other results on ancient solutions to curvature flows appeared, see [5, 24] for a classification of ancient solutions using Aleksandrov reflection and [6, 7, 19, 20] for other related results.

This paper is laid out as follows: In Section 2, we establish some notation and basic equations to be used in this paper. In Section 3, we introduce the notion of quasi-ancient solutions and establish some facts about such solutions. In Section 4, we give a short classification result for 1-homogeneous concave or convex speeds and for the curve shortening flow. In Section 5, we use the rigidity results from [21] to show that the backwards limits of convex quasi-ancient solutions with bounded H are equators. In Section 6, we use an Aleksandrov reflection technique to complete the classification. In Section 7, we look at two examples of non-geometric flows for which there exist convex, ancient solutions that are not shrinking spheres, illustrating the necessary restriction to geometric flows and justifying our use of geometric methods.

Acknowledgment

The authors would like to thank Knut Smoczyk and the Institut für Differentialgeometrie at Leibniz Universität for hosting a research visit where part of this work took place. The first author was a Riemann Fellow at the Riemann Center for Geometry and Physics, Leibniz Universität and was also supported by the EPSRC on a Programme Grant entitled "Singularities of Geometric Partial Differential Equations" reference number EP/K00865X/1. The work of the second author was supported by Austrian Science Fund (FWF) Project M1716-N25 and the European Research Council (ERC) Project 306445. All three authors would like to thank Slack.com and Overleaf.com for providing wonderful platforms for collaboration. The figure in this paper was created using the wonderful SageMath Cloud (cloud.sagemath.com).

2. Preliminaries

It is well known, e.g. [13, Chapter 2], that a curvature function as in Assumption 1.1 also can be considered to depend on the Weingarten map,

$$F = F(\mathcal{W}) = F(h_i^i)$$

in case of which the derivatives of the speed F are written as

$$F_j^i = \frac{\partial F}{\partial h_i^j}.$$

It is also possible to consider F as a function of the metric and second fundamental form,

$$F(g,h) = F(g^{ik}h_{kj}),$$

where we write

$$F^{ij} = \frac{\partial F}{\partial h_{ij}}, \quad F^{ij,kl} = \frac{\partial^2 F}{\partial h_{kl} \partial h_{ij}}.$$

These derivatives are related by

$$F^{ij} = g^{ik} F^j_k.$$

Let us also define the operator

$$\Box = F^{ij} \nabla_{ij}^2,$$

where ∇ is the Levi-Civita connection of the induced metric g. Occasionally ∇ will denote the Levi-Civita connection of the round metric of \mathbb{S}^n , in which case we will explicitly say so.

2.1. Lemma. The following evolution equations hold.

(2.1)
$$\partial_t h_i^j = \Box h_i^j + F^{kl} (h^2)_{kl} h_i^j - (F^{kl} h_{kl} - F) (h^2)_i^j + F^{kl, rs} \nabla_i h_{kl} \nabla^j h_{rs} \\ + (F + F^{kl} h_{kl}) \delta_i^j - F^{kl} g_{kl} h_i^j,$$

(2.2)
$$\partial_t F = \Box F + F F^{ij} (h^2)_{ij} + F F^{ij} g_{ij}$$

Proof. The proofs are standard; see [13, Lemma 2.4.3, Lemma 2.3.4].

We will occasionally have to work with graphs u over an equator $E \subset \mathbb{S}^{n+1}$ around a point $e \in \mathbb{S}^{n+1}$. In geodesic polar coordinates around e the spherical metric takes the form

(2.3)
$$d\bar{s}^2 = d\rho^2 + \vartheta(\rho)^2 \sigma_{ij}(x) dx^i dx^j,$$

where ρ is the radial distance to $e, \vartheta(\rho) = \sin(\rho)$ and σ the round metric on the equator $E \simeq \mathbb{S}^n$. Hence, if a hypersurface M is given as a graph of a function u over E,

$$M = \{ (\rho, (x^i)) \in \mathbb{S}^{n+1} \colon \rho = u(x), \ x \in E \},\$$

a straightforward computation yields the following representation of the Weingarten map in terms of the function u, namely

(2.4)
$$h_j^i = \frac{\vartheta'}{\vartheta\vartheta} \delta_j^i + \frac{\vartheta'}{\vartheta^3\vartheta^3} \nabla^i u \nabla_j u - \frac{g^{ik}}{\vartheta} \nabla_{kj}^2 u,$$

where $v^{-1} = \partial_{\rho} \cdot \nu$. Covariant derivatives as well as index raising are performed with respect to σ_{ij} ; see, for example, [25, (3.82)].

A one-parameter family of graphs satisfying the flow (1.1) is a solution of

(2.5)
$$\frac{\partial}{\partial t}u = -F\left(\frac{\vartheta'}{v\vartheta}\delta_j^i + \frac{\vartheta'}{v^3\vartheta^3}\nabla^i u\nabla_j u - \frac{g^{ik}}{v}\nabla_{kj}^2 u\right)v = \Phi(x, u, \nabla u, \nabla^2 u);$$

see [13, p. 98-99].

3. Ancient and quasi-ancient Solutions

We are interested in solutions with maximal possible lifetime. To understand this maximal time, we define T_S to be the lifespan of the strictly convex spherical solution of (1.1). By the strictly convex spherical solution we mean a family of geodesic spheres shrinking under the flow (1.1) collapsing to a point at time t = 0and existing on the maximal interval $(-T_S, 0)$. T_S can be finite or infinite, as the following lemma shows.

3.1. Lemma. The following assertions hold:

- i: Let $p \in (0,1)$ and consider (1.1) with speed $F = f^p$ where f is strictly increasing, concave and 1-homogeneous. Then the strictly convex spherical solution exists only on a finite time interval $(-T_S, 0)$ with $0 < T_S < \infty$, collapses to a point at t = 0 and converges to an equator at $t = -T_S$.
- ii: Let F be 1-homogeneous, then the strictly convex spherical solution of (1.1) is ancient.

Proof. (i) We may assume that f(1, ..., 1) = n. Since F is constant on a geodesic sphere, the evolution equation (2.2) for a flow of geodesic spheres yields

$$\frac{d}{dt}f \ge \frac{1}{n}f^{p+2} + nf^p,$$

where we used [13, Lemma 2.2.19, Lemma 2.2.20]. Finite lifespan forward in time follows immediately. Integration over an interval $(a, b) \subset (-T_S, 0)$ yields

$$0 \le f^{1-p}(a) \le f^{1-p}(b) - n(1-p)(b-a).$$

Allowing $a \to -\infty$ gives the finite time existence backwards in time.

(ii) Assume F(1, ..., 1) = n and let r be the radius of the spherical solution. Then (1.1) yields

$$\dot{r} = -n \cot r.$$

The maximal spherical solution satisfies

$$r(0) = 0$$
 and $r(-T) = \frac{\pi}{2}$

and hence integration of this ODE yields $T = \infty$.

3.2. Lemma. Let x be a non-equatorial, convex solution of (1.1), defined on the open interval (-T, 0), where 0 is the collapsing time, then $T \leq T_S$.

Proof. Suppose $T > T_S + \epsilon$ for some $\epsilon > 0$. $M = M_{-T_S - \frac{\epsilon}{2}}$ bounds a convex body \hat{M} and is strictly contained in an open hemisphere due to the classical paper [12]. Then there exists a geodesic sphere S with $\hat{M} \subset \hat{S}$. By the avoidance principle, the flow with initial hypersurface M collapses before the spherical flow thus contradicting $T > T_S$.

In view of this lemma, the following definition is reasonable.

3.3. **Definition.** A convex solution of (1.1) defined on an interval (-T, 0) is called *quasi-ancient*, if $T = T_S$. For convenience we also call the static equatorial solution quasi-ancient.

By the definition, convex ancient solutions are also quasi-ancient. For convex speeds of homogeneity 1, the following proposition gives a bound on mean curvature backwards in time for ancient solutions, using a Harnack inequality. For genuine quasi-ancient solutions $(T_S < \infty)$, the Harnack inequality (whenever it holds) does not give such a bound since we cannot send $s \to -\infty$ as in the proof. For example, a Harnack inequality holds for the flows $F = H^p$, $0 ; see [9, Theorem 1] and from Lemma 3.1, <math>T_S < \infty$. One could envisage backwards limits as convex polyhedra and hence with unbounded H, but it is not clear that these solutions exist on the maximal time interval $(-T_S, 0)$, i.e., we do not know if they can be quasi-ancient solutions. For the classification of quasi-ancient solutions, we thus make the additional assumption that H is bounded, and defer the question of whether solutions with unbounded H can exist as an interesting study for a later date.

For a strictly convex hypersurface, define (b^{ij}) to be the inverse of (h_{ij}) .

3.4. **Proposition.** Suppose F is a strictly monotone, 1-homogeneous and convex curvature function. Then any strictly convex ancient solution of (1.1) satisfies

$$\partial_t F - b^{ij} \nabla_i F \nabla_j F \ge 0$$

Therefore, for any $t_0 > 0$ and all $t \leq -t_0$ we have $H(\cdot, t) \leq c(t_0)$, where $c(t_0) < \infty$ depends only on M_{-t_0} .

Proof. For any t > s, the Harnack estimate of [9, Theorem 1] implies that

$$\partial_t F - b^{ij} \nabla_i F \nabla_j F + \frac{1}{2} \frac{F}{t-s} > 0.$$

Allowing $s \to -\infty$ proves the first claim.

In particular, for strictly convex, ancient solutions, $b^{ij} > 0$ and hence $F(\cdot, t)$ is a non-decreasing function in t. For the second claim then, observe that for any 1-homogeneous, convex F we have

$$F \ge \frac{F(1,\ldots,1)}{n}H$$

see [13, Lemma 2.2.20]. Therefore, strictly convex, ancient solutions satisfy

$$H(\cdot,t) \le \frac{n}{F(1,\ldots,1)} F(\cdot,t) \le \frac{n}{F(1,\ldots,1)} F(\cdot,-t_0).$$

for all $t \leq -t_0$.

4. CONVEX, CONCAVE AND HOMOGENEOUS SPEEDS

In this section, we give a quick classification of ancient solutions in three special cases.

4.1. **Theorem.** Let $n \ge 2$ and let F be strictly monotone, 1-homogeneous, and either convex, or concave with $H \le cF$ for some c > 0. Then any strictly convex ancient solution of the flow (1.1) with speed F is a family of contracting geodesic spheres.

Proof. We may assume F is normalized so that $F(1, \ldots, 1) = n$. We will give the details for F convex. The concave case is similar, and we remark below where the proof deviates slightly.

Convexity and 1-homogeneity implies that $F \ge \frac{F(1,...,1)}{n}H$; see [13, Lemma 2.2.20]. The chosen normalization then implies,

(4.1)
$$\frac{\kappa_{\min}}{F} \le \frac{\kappa_{\min}}{H} \le \frac{1}{n}$$

Now we prove the reverse inequality, $\frac{\kappa_{\min}}{F} \geq \frac{1}{n}$. The tensor $w_i^j = \frac{h_i^j}{F}$ satisfies the evolution equation

$$\partial_t w_i^j = \Box w_i^j - 2F^{kl}g_{kl}w_i^j + 2\delta_i^j + \frac{1}{F}F^{kl,rs}\nabla_i h_{kl}\nabla^j h_{rs}$$
$$- 2F^{kl}\nabla_k h_i^j \nabla_l \left(\frac{1}{F}\right) - 2\frac{h_j^i}{F^3}F^{kl}\nabla_k F\nabla_l F.$$

At a critical point of w_i^i we have

$$0 = \nabla_k \left(\frac{h_i^i}{F}\right) = \frac{\nabla_k h_i^i}{F} + h_i^i \nabla_k \left(\frac{1}{F}\right),$$

and hence the second line above vanishes:

$$-2F^{kl}\nabla_k h_i^i \nabla_l \left(\frac{1}{F}\right) - 2\frac{h_i^i}{F^3}F^{kl}\nabla_k F \nabla_l F$$
$$= 2F^{kl} \left(Fh_i^i \nabla_k \left(\frac{1}{F}\right)\nabla_l \left(\frac{1}{F}\right) - \frac{h_i^i}{F^3}\nabla_k F \nabla_l F\right) = 0.$$

Homogeneity, convexity and the normalization F(1, ..., 1) = n imply that

$$F^{kl}g_{kl} \le n$$
, and $F^{kl,rs} \ge 0$,

see the proof of [13, Lemma 2.2.19]. Fixing an index i and working in normal coordinates centered on a local minimum we have

$$\frac{1}{F}F^{kl,rs}\nabla_i h_{kl}\nabla^i h_{rs} = \frac{1}{F}F^{kl,rs}\nabla_i h_{kl}\nabla_i h_{rs}$$

is non-negative. Thus at a spatial minimum,

$$\partial_t \left(w_i^i - \frac{1}{n} \right) \ge \Box \left(w_i^i - \frac{1}{n} \right) - 2n \left(w_i^i - \frac{1}{n} \right).$$

By the maximum principle, for any s < 0, we obtain the bound

$$w_i^i - \frac{1}{n} \ge c_s e^{-2n(t-s)}$$

for $t \geq s$, and where

$$c_s := \min \frac{\kappa_{\min}}{F}(\cdot, s) - \frac{1}{n}.$$

Moreover, we have the uniform bounds

$$-\frac{1}{n} \le c_s \le 0$$

from equation (4.1).

Taking the limit $s \to -\infty$ we then obtain for all t < 0,

(4.2)
$$\frac{\kappa_{\min}}{F} \ge \frac{1}{n}.$$

Combining equations (4.1) and (4.2) we obtain

$$\frac{\kappa_{\min}}{F} \le \frac{\kappa_{\min}}{H} \le \frac{1}{n} \le \frac{\kappa_{\min}}{F}.$$

Hence, $H \equiv n\kappa_{\min}$ and the flow is by totally umbilical, closed hypersurfaces, and hence by geodesic spheres.

In the concave case, we obtain

$$w_i^i - \frac{1}{n} \le d_s e^{-2n(t-s)},$$

where $d_s := \max \frac{\kappa_{\max}}{F}(\cdot, s) - \frac{1}{n} \ge 0$. Using the additional assumption $\frac{H}{F} \le c$, we obtain that

$$d_s \le \max \frac{H(\cdot, s)}{F} - \frac{1}{n} \le c - \frac{1}{n}$$

Now by letting $s \to -\infty$, we conclude that $H \equiv n\kappa_{\text{max}}$ and the claim follows. \Box

4.2. Remark. The bounds,

$$\frac{\kappa_{\max}}{F} \le \frac{1}{n}, \quad F \text{ concave}$$
$$\frac{\kappa_{\min}}{F} \ge \frac{1}{n}, \quad F \text{ convex}$$

may also be obtained from the curvature ball estimates of [1, Theorem 1.1], which imply that any strictly convex solution of the flow with a 1-homogeneous, concave speed F satisfies

$$\frac{\kappa_{\max}}{F}(x,t) \le \frac{1}{n} + d_s e^{-2n(t-s)},$$

with d_s as in the proof above. For convex speeds, the curvature ball estimates imply that

$$\frac{\kappa_{\min}}{F}(x,t) \ge \frac{1}{n} + c_s e^{-2n(t-s)},$$

where c_s is from the proof above. The proof applying the maximum principle to h_i^i/F above, though weaker than the curvature ball estimates, is significantly simple and suffices for our purposes.

4.3. Remark. In virtue of the evolution equations in Lemma 2.1, for strictly increasing, 1-homogeneous, convex speeds F with $F \in C^1(\bar{\Gamma}_+)$, any non-equatorial, convex, ancient solution of (1.1) is in fact strictly convex. To see this, apply the strong maximum principle to the evolution equation of F to conclude that F must be strictly positive for any time. On the other hand, the last term in the evolution equation of κ_{\min} is bounded below by

$$F + F - n\kappa_{\min} \ge F + H - n\kappa_{\min} \ge F > 0.$$

Therefore, the statement of Theorem 4.1 holds under the weaker assumption of convexity of ancient solutions rather than strict convexity.

We end this section by providing a short proof for the classification result of [10].

4.4. **Theorem.** [10] The only convex, ancient solutions of the curve shortening flow on the sphere are shrinking geodesic circles or equators.

8

Proof. Suppose γ_t is a convex, ancient solution to the curve shortening flow. Let s denote the arc-length parameter of γ_t . By Lemma 2.1 and $\frac{d}{dt}ds = -\kappa^2 ds$ we have

$$\frac{d}{dt}\left(\left(\int_{\gamma_t} \kappa ds\right)^2 + \left(\int_{\gamma_t} ds\right)^2\right) = 2\left(\left(\int_{\gamma_t} \kappa ds\right)^2 - \int_{\gamma_t} ds \int_{\gamma_t} \kappa^2 ds\right) \le 0$$

Therefore,

$$q(t) = \left(\int_{\gamma_t} \kappa ds\right)^2 + \left(\int_{\gamma_t} ds\right)^2$$

is a non-increasing function. By [10, Lemma 4.1], $\lim_{t \to \infty} \int_{\gamma_t} \kappa ds = 0$; therefore,

$$\lim_{t \to -\infty} q(t) = 4\pi^2$$

On the other hand, by the Gauss-Bonnet theorem and the isoperimetric inequality [23], $q(t) \ge 4\pi^2$ and hence

$$q(t) \equiv 4\pi^2.$$

By the characterization of the equality cases, γ_t are either shrinking geodesic circles or equators.

5. RIGIDITY AND BACKWARDS LIMIT

Now we turn to more general speeds F. The aim of this section is to prove that for a quasi-ancient solution of (1.1) with bounded mean curvature for $t \to -T_S$ the backwards limit of the flow hypersurfaces M_t is an equator. We will use the rigidity result of [21] to achieve this. For convenience, we state this result:

5.1. **Theorem.** [21, Theorem 1.1] Let $n \ge 1$ and $\hat{M} \subset \mathbb{S}^{n+1}$ be a weakly convex body in a hemisphere. Let $x_0 \in \mathbb{S}^{n+1}$ be such that \hat{M} is contained in the closed hemisphere $\mathcal{H}(x_0)$ with equator $\mathcal{S}(x_0)$. Suppose that \hat{M} satisfies an interior sphere condition at all points $p \in \hat{M} \cap \mathcal{S}(x_0)$. Then either \hat{M} is equal to $\mathcal{H}(x_0)$ or \hat{M} is contained in an open hemisphere.

Here \hat{M} is a weakly convex body in a hemisphere $\mathcal{H}(x_0)$, if it is a compact set with non-empty interior and for every two points $p, q \in \hat{M}$ there exists a minimizing geodesic connecting p with q, while being contained in \hat{M} . \hat{M} satisfies an interior sphere condition at $p \in \partial \hat{M}$ with radius R, if there exists a geodesic ball B_R with radius R, such that

$$\partial B_R \cap \partial M = \{p\} \text{ and } B_R \subset \operatorname{int}(M).$$

Note that the points p in Theorem 5.1 are automatically boundary points of \hat{M} .

In what follows, in addition to the notion of "weakly convex body", we will also need the notions of "weakly convex set", "convex set" and "convex body" for which we refer the reader to [21, Definition 3.2].

The next lemma is the first step in providing the assumptions of Theorem 5.1.

5.2. Lemma. Let x be a convex, quasi-ancient solution of (1.1) with backwards bounded mean curvature. Then there holds:

(1) For all $t_0 < 0$ there exists a uniform radius R > 0, such that the enclosed convex bodies \hat{M}_t , $-T_S < t \le t_0$, of the flow hypersurfaces M_t satisfy a uniform interior sphere condition with radius R.

(2) For every $y_0 \in \operatorname{int} M_{t_0}$ the hypersurfaces M_t , $-T_S < t \le t_0$, can be written as a graph in geodesic polar coordinates around y_0 and the corresponding graph functions satisfy uniform C^2 -estimates.

Proof. Fix an interior point $y_0 \in \inf \hat{M}_{t_0}$. Since for a contracting flow the enclosed convex bodies of the flow hypersurfaces are decreasing, they are increasing backwards in time. By the proof of [21, Lemma 3.9] there exists a closed hemisphere $\mathcal{H}(x_0)$, such that

$$\hat{M}_t \subset \mathcal{H}(x_0).$$

In our situation all hypersurfaces M_t , $-T_S < t \le t_0$, satisfy

$$B_{\epsilon}(y_0) \subset \operatorname{int} \hat{M}_{\epsilon}$$

and

$$B_{\epsilon}(\hat{y}_0) \subset M_t^c$$

with a uniform ϵ , where \hat{y}_0 denotes the antipodal point of y_0 . Now we prove the two claims.

- (1) Consider the stereographic projection with \hat{y}_0 corresponding to infinity. The image hypersurfaces are then strictly convex hypersurfaces in Euclidean space with uniformly bounded second fundamental form. Blaschke's rolling theorem (see [4]) gives the interior sphere condition.
- (2) Write the M_t as graphs in geodesic polar coordinates around y_0 ,

$$M_t = \{ (\rho, x^i) \colon \rho = u(t, x^i) \}.$$

Due to (2.3), on the set in which M_t range, the metrics g_{ij} , $\bar{g}_{ij} = \sin^2 \rho \sigma_{ij}$ and σ_{ij} are all equivalent. In view of [13, Theorem 2.7.10], for all convex hypersurfaces M_t the quantity

$$v^2 = 1 + \bar{g}^{ij} \nabla_i u \nabla_j u$$

is uniformly bounded by a constant which only depends on ϵ . Hence by the equivalence of norms, M_t are uniformly C^1 -bounded in the sense that the corresponding functions $u(t, \cdot)$ are uniformly $C^1(\mathbb{S}^n)$ -bounded. Recalling equation (2.4), due to the curvature estimates we obtain uniform $C^2(\mathbb{S}^n)$ -estimates for u.

5.3. Corollary. Let x be a convex and quasi-ancient solution of (1.1) with backwards bounded mean curvature. Then there exists a unique backwards limiting hypersurface M_{-T_S} and the flow hypersurfaces M_t converge to M_{-T_S} in $C^{1,\beta}$, $0 < \beta < 1$, in the sense that for a common graph representation as in Lemma 5.2 there holds

$$u(t,\cdot) \to u(-T_S,\cdot)$$

in the norm of $C^{1,\beta}(\mathbb{S}^n)$.

Proof. In view of the point-wise monotonicity of $u(t, \cdot)$ backwards in time, we obtain a point-wise limit. The $C^{1,\beta}$ -convergence follows from compactness.

5.4. **Theorem.** The hypersurface M_{-T_s} defined in Corollary 5.3 is an equator.

Proof. Since the convex bodies \hat{M}_t are increasing backwards in time and due to the uniform convergence of M_t to M_{-T_s} , the set

$$\hat{M}_{-T_S} := \overline{\bigcup_{t < 0} \hat{M}_t}$$

is a compact body with

$$\partial M_{-T_S} = M_{-T_S}$$

Since $\operatorname{int}(\hat{M}_{-T_S})$ is a convex set, it is especially a weakly convex set in a hemisphere. Thus \hat{M}_{-T_S} is a weakly convex body in a hemisphere. The proof of [21, Lemma 6.1] can literally be applied to show that \hat{M}_{-T_S} satisfies a uniform interior sphere condition as well. We can apply [21, Theorem 1.1] and obtain that \hat{M}_{-T_S} is either strictly contained in an open hemisphere or is equal to a closed hemisphere. The first alternative is not possible since the solution is quasi-ancient.¹ We conclude that $\partial \hat{M}_{-T_S} = M_{-T_S}$ is an equator of \mathbb{S}^{n+1} .

6. Aleksandrov Reflection and Classification

In this section, we use the result of Theorem 5.4 to classify convex and quasiancient solutions of contracting curvature flows on \mathbb{S}^{n+1} as either equators or shrinking geodesic spheres. The proof uses Aleksandrov reflection as in [8, 10] to show that the symmetry of the backwards limit is preserved along the flow. Here we give a very general version with minimal assumptions on the flow: all we require is that the flow limits to an equator at $-T_S$ in C^0 , with uniform C^2 bounds, and that the flow is geometric and parabolic, compare Assumption 1.1.

6.1. Notation. We consider \mathbb{S}^{n+1} to be embedded into \mathbb{R}^{n+2} by the standard inclusion. Let

$$e = e_{n+2} = (0, \dots, 0, 1) \in \mathbb{R}^{n+2}.$$

For a vector $V \in \mathbb{S}^{n+1}$ let $H_V^{\pm} = \{x \in \mathbb{R}^{n+2} : \pm \langle x, V \rangle > 0\}$, and for a set $S \in \mathbb{R}^{n+2}$ define $S_V^{\pm} = S \cap H_V^{\pm}$. We denote the equator in \mathbb{S}^{n+1} around e by $E = e^{\perp} \cap \mathbb{S}^{n+1}$. Furthermore, δ_V denotes the signed angle that V makes with the hyperplane e^{\perp} , $\delta_V = \arcsin \langle V, -e \rangle$. The reflection map across the hyperplane V^{\perp} is denoted by

$$R_V \colon \mathbb{R}^{n+2} \to \mathbb{R}^{n+2}$$
$$x \mapsto x - 2 \langle x, V \rangle V$$

which is an isometry of \mathbb{R}^{n+2} . Finally, for $x \in \mathbb{S}^{n+1} \setminus \{\pm e\}$, let γ_x denote the unique minimizing geodesic in \mathbb{S}^{n+1} from e to -e that passes through x and then define

$$\operatorname{pr}(x) = \begin{cases} y, & x \in \mathbb{S}^{n+1} \setminus \{\pm e\} \\ E, & x = e \text{ or } x = -e. \end{cases}$$

where $y \in E$ is the unique element on the image of γ_x lying in E.

The reader may find it useful to refer to Figure 1 for the arguments in this section.

¹On the contrary, suppose \hat{M}_{-T_S} is contained in an open hemisphere, then there exists an open geodesic ball B of radius $\rho \leq \pi/2 - \epsilon$, for some $\epsilon > 0$, enclosing \hat{M}_{-T_S} . Its boundary sphere ∂B has a maximal finite time existence $T < T_S - c_\epsilon$ for some constant c_ϵ . For all $t > -T_S$ we have $M_t \subset B$. Hence the flow starting from M_t exists only for $-t \leq T$. Letting $t \to -T_S$ yields a contradiction.



FIGURE 1. Reflection in the (e, V)-plane showing the reflected equator, M_t , $R_V(M_t)$ and some geodesics through the north pole (dotted lines).

6.2. Star-shaped hypersurfaces and one-sided reflections.

6.1. *Remark.* Recall a nonempty set $S \in \mathbb{S}^{n+1}$ to be star-shaped around e, if $\pm e \notin S$ and if every minimizing geodesic from e to -e hits S at most once, i.e.,

$$\forall x \in E \colon \# (\operatorname{Im} \gamma_x \cap S) \le 1.$$

In this case we have a well-defined graph function of S,

$$f_S \colon \operatorname{pr} S \to \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$
$$\sigma \mapsto \operatorname{arcsin} \left\langle x_{\sigma}, e \right\rangle,$$

where x_{σ} is the unique preimage of σ in S under the projection pr. We obtain a parametrization of the set S via the correspondence

$$x_{\sigma} = (\sigma, f_S(\sigma)).$$

6.2. **Definition.** Let S, T be two star-shaped sets around e with the corresponding functions f_S and f_T .

(1) We say that S lies above T, denoted by $S \ge T$, if

$$\forall \sigma \in \operatorname{pr} S \cap \operatorname{pr} T \colon f_S(\sigma) \ge f_T(\sigma).$$

(2) Let $V \in \mathbb{R}^{n+2}$ and suppose $R_V(S)$ and T are star-shaped around e. We say that S one-sided reflects above T, if

$$R_V(S_V^+) \ge T_V^-.$$

6.3. Lemma. Let $V \in \mathbb{S}^{n+1}$ and $0 \leq \delta_V \leq \delta_0 < \frac{\pi}{4}$. Then

- i: $R_V(E)$ is star-shaped around e.
- ii: There exists a constant $c = c(\delta_0) > 0$ with the following property. If a closed and convex hypersurface $M \subset \mathbb{S}^{n+1}$ is star-shaped around e and satisfies

$$|f_M| \leq c,$$

then the reflection $R_V(M)$ is star-shaped around e.

Proof. (i) Since $R_V(E)$ is also an equator, the only way not to be star-shaped around e would be $e \in R_V(E)$. But then, letting $e = R_V(x)$ with $x \in E$, and using the Cauchy-Schwarz inequality, we obtain

$$1 = \langle R_V(x), e \rangle = 2 \sin \delta_V \langle x, V \rangle \le 2 \sin \delta_V \cos \delta_V,$$

which is impossible in the given range of δ_V .

(ii) Let us put

$$2r = \min_{\{V: \ 0 \le \delta_V \le \delta_0\}} \operatorname{dist}(e, R_V(E)) > 0.$$

The minimum is precisely achieved for vectors V with $\delta_V = \delta_0$ and thus r depends on δ_0 . Suppose $S_r(\pm e)$ are the geodesic spheres of radii r with centers at $\pm e$. Note that $S_r(e)$ is contained in the interior of one of the open hemispheres formed by $R_V(E)$, and $S_r(-e)$ is contained in the interior of the opposite hemisphere.

Write $E_{\pm\varepsilon}$ for the geodesic spheres at heights $\pm\varepsilon$. By continuity, there exists $c(\delta_0)$ such that for all $0 < \varepsilon \leq c(\delta_0)$, $S_{\frac{r}{2}}(e)$ is contained in the interior of the smallest spherical cap cut by $R_V(E_{\varepsilon})$, and $S_{\frac{r}{2}}(-e)$ is contained in the interior of the smallest spherical cap cut by $R_V(E_{-\varepsilon})$.

Assume that $-c(\delta_0) \leq f_M \leq c(\delta_0)$. Then using $E_{\pm c(\delta_0)}$ as barriers we can see that one of the bodies enclosed by $R_V(M)$ contains only e in its interior and the other contains -e in its interior. Now convexity of $R_V(M)$ implies that it is star-shaped around e.

It is easily seen that the equator E one-sided reflects above itself with respect to V^{\perp} , if $0 < \delta_V < \frac{\pi}{4}$. Now we show that this is also true for any convex hypersurface M, which is C^1 -close to E.

6.4. **Proposition.** Suppose $V \in \mathbb{S}^{n+1}$ and $0 < \delta_1 \leq \delta_V \leq \delta_0 < \frac{\pi}{4}$. There exists $\alpha = \alpha(\delta_0, \delta_1) > 0$ with the following property. If M is a closed and convex hypersurface that is star-shaped around e and

$$0 \le f_M, \quad |f_M|_{C^1} \le \alpha,$$

then M one-sided reflects above itself.

Proof. Assume without loss of generality that $f_M \leq c(\delta_0)$, where $c(\delta_0)$ is given in the previous lemma. We will prove

$$R_V(M_V^+) \ge M_V^-.$$

Suppose there exist $x \in M_V^+$ and $y \in M_V^-$ with

$$\operatorname{pr}(R_V(x)) = \operatorname{pr}(y) \in \operatorname{pr}(R_V(M_V^+)) \cap \operatorname{pr}(M_V^-).$$

Consider the projection map

pr:
$$S_{\varepsilon} \subset \mathbb{R}^{n+2} \to E \subset \mathbb{R}^{n+1}$$

pr $(a) = \frac{a - \langle a, e \rangle e}{|a - \langle a, e \rangle e|},$

where S_{ε} is the compact set that is made by removing two opposite caps of height $0 < \varepsilon < c(\delta_0)$ from the unit sphere at e, -e. Since pr is smooth on S_{ε} , it is Lipschitz and we have the estimate

$$|\operatorname{pr}(a) - \operatorname{pr}(b)| \le l_{\varepsilon}|a - b|, \quad \forall a, b \in S_{\varepsilon}$$

for some constant $0 < l_{\varepsilon} < \infty$. In addition, we have

$$\langle x, e \rangle = \sin(f_M(\operatorname{pr}(x))), \quad \forall x \in M,$$

$$2 \operatorname{arcsin}\left(\frac{|\operatorname{pr}(a) - \operatorname{pr}(b)|}{2}\right) = d(\operatorname{pr}(a), \operatorname{pr}(b)),$$
$$\operatorname{arcsin}(z) \le \frac{\pi}{2}z, \quad \forall \ 0 \le z \le 1.$$

Therefore, we obtain

$$\begin{aligned} \langle x, e \rangle - \langle y, e \rangle &\geq -|\sin(f_M(\operatorname{pr}(x))) - \sin(f_M(\operatorname{pr}(y)))| \\ &\geq -d(\operatorname{pr}(x), \operatorname{pr}(y))|f_M|_{C^1(\mathbb{S}^n)} \\ &= -2 \operatorname{arcsin}(\frac{1}{2}|\operatorname{pr}(x) - \operatorname{pr}(y)|)|f_M|_{C^1} \\ &\geq -\frac{\pi}{2}|\operatorname{pr}(x) - \operatorname{pr}(y)||f_M|_{C^1}. \end{aligned}$$

Let us put $\alpha = \min\{\frac{2\sin\delta_1}{\pi l_{\varepsilon}}, c(\delta_0)\}$. To finish the proof, note that

$$\langle R_V(x), e \rangle - \langle y, e \rangle = \langle R_V(x), e \rangle - \langle x, e \rangle + \langle x, e \rangle - \langle y, e \rangle$$

$$\geq 2 \langle x, V \rangle \sin \delta_V - \frac{\pi}{2} |\operatorname{pr}(x) - \operatorname{pr}(y)| |f_M|_{C^1}$$

$$= |R_V(x) - x| \sin \delta_V - \frac{\pi}{2} |\operatorname{pr}(x) - \operatorname{pr}(R_V(x))| |f_M|_{C^1}$$

$$\geq \left(\sin \delta_1 - \frac{\pi}{2} l_{\varepsilon} \alpha \right) |R_V(x) - x| \geq 0$$

and hence,

$$\operatorname{arcsin}(\langle R_V(x), e \rangle) \ge \operatorname{arcsin}(\langle y, e \rangle)$$

as required in Definition 6.2.

6.3. Reflecting quasi-ancient curvature flows. The final aim of this section is to show that for a solution of a curvature flow defined on an interval $(-T_S, 0)$, which converges backwards in time to E in C^1 and has bounded curvature, the flow hypersurfaces are invariant under reflection about every hyperplane V^{\perp} with $V \in E$. Hence we somehow need to let δ_1 go to zero in the preceding proposition. In fact, until now we have only obtained that for a given $\delta_1 > 0$, there exists T_{δ_1} such that all flow hypersurfaces in an interval $(-T_S, T_{\delta_1})$ one-sided reflect above themselves. Thus we cannot yet deduce that any flow hypersurface has this property for $\delta_1 = 0$. To eliminate the dependence of the interval $(-T_S, T_{\delta_1})$ on δ_1 , we will use the parabolic maximum principle. 6.5. Lemma. Let the closed and convex hypersurfaces M_t , $-T_S < t < 0$, satisfy the parabolic curvature flow equation (1.1) under either assumption (i) or (ii) in Theorem 1.2 and suppose

$$M_t \to E, \quad t \to -T_S,$$

in the sense that e lies in the enclosed convex bodies by M_t for all sufficiently small times and the graph functions satisfy

$$0 \le f_{M_t} \quad and \quad \lim_{t \to -T_S} |f_{M_t}|_{C^1} = 0.$$

Then there exists $T > -T_S$, such that for all $V \in \mathbb{S}^{n+1}$ satisfying

$$0 < \delta_V \le \frac{\pi}{8},$$

 M_t one-sided reflects above itself for all $t \in (-T_S, T)$.

Proof. We may assume M_t is not the equator E and that $f_{M_t} \leq \pi/4$ for all $t \leq T_*$. By Lemma 6.3 and the C^1 -convergence of M_t we find a time $T \in (-T_S, T_*]$, such that M_t and $R_V(M_t)$ are both star-shaped around e and e also lies in the convex body of $R_V(M_t)$ for all $t \in (-T_S, T]$. We show that this T has the desired property.

By Proposition 6.4, we know that for a given $V \in \mathbb{S}^{n+1}$ with (suppressing the subscript V)

$$0 < \delta \le \frac{\pi}{8},$$

there exists $-T_S < \tilde{T}_{\delta} \leq T_*$, such that M_t one-sided reflects above itself for all $t \in (-T_S, \tilde{T}_{\delta})$. Pick a $-T_S < T_{\delta} < \tilde{T}_{\delta}$ and define the domain

$$\Omega = \bigcup_{t \in (T_{\delta}, T)} \left(\operatorname{pr}(M_t^-) \cap \operatorname{pr}(R(M_t^+)) \times \{t\} \subset E \times (-T_S, T). \right)$$

Recall from (2.5) that the time dependent radial graph functions of M_t satisfy

$$\frac{\partial}{\partial t}w = -F\left(\frac{\vartheta'}{v\vartheta}\delta^i_j + \frac{\vartheta'}{v^3\vartheta^3}\nabla^i w\nabla_j w - \frac{g^{ik}}{v}\nabla^2_{kj}w\right)v,$$

where $\vartheta = \vartheta(w) = \sin(w)$, $w(\cdot, t) = \pi/2 - f_{M_t}(\cdot)$, $v^2 = 1 + \vartheta^{-2} |\nabla w|^2$ and the connection and the norm $|\cdot|$ are those belonging to the spherical metric σ_{ij} . Since the Weingarten operator is invariant under ambient isometries, the reflected hypersurfaces $R_V(M_t)$ with the corresponding graph functions u satisfy the same curvature flow equation.

If the claim of the lemma were false, then there existed a time $t_* \in (T_{\delta}, T)$ and a point $x_* \in \operatorname{pr}(M_{t_*}^-) \cap \operatorname{pr}(R(M_{t_*}^+))$, such that

(6.1)
$$u(x_*, t_*) > w(x_*, t_*) \ge \frac{\pi}{4}.$$

Formally we would like to apply the parabolic maximum principle to conclude that u must remain below w, since this is the case at the initial time T_{δ} and also at the boundaries $\partial \left(\operatorname{pr}(M_t^-) \cap \operatorname{pr}(R(M_t^+)) \right) \subset \operatorname{pr}\left(V^{\perp} \cap M_t\right)$.²

²Note that for two sets A and B we have

 $\partial(A \cap B) \subset \partial A \cup \partial B.$

Since pr is a diffeomorphism from M_t to E and M_t is star-shaped,

$$\partial(\operatorname{pr}(M_t^-)) = \operatorname{pr}(\partial M_t^-) = \operatorname{pr}(V^{\perp} \cap M_t)$$

Similarly, $\partial(\operatorname{pr}(R(M_t^+))) = \operatorname{pr}(R(\partial M_t^+)) = \operatorname{pr}(R(V^{\perp} \cap M_t)) = \operatorname{pr}(V^{\perp} \cap M_t).$

However, in this situation the application of the standard comparison principles to equations of the form

$$\dot{w} = \Phi(x, w, \nabla w, \nabla^2 w)$$

is not straightforward; note to the restriction that F is generally only defined on non-negative definite endomorphisms and hence the standard proof of comparison principles for fully nonlinear operators, namely forming convex combinations of two solutions, might leave the domain of Φ . Nevertheless, assuming (6.1), we obtain a contradiction using the following argument.

A slight rewriting of the second fundamental form of $R(M_t)$ described by the graph function u yields, cf. (2.4),

$$\begin{split} h_{j}^{i} &= \frac{\vartheta'}{\sqrt{\vartheta^{2} + |\nabla u|^{2}}} \delta_{j}^{i} + \frac{\vartheta'}{(\vartheta^{2} + |\nabla u|^{2})^{\frac{3}{2}}} \nabla^{i} u \nabla_{j} u - \frac{\vartheta g^{ik}}{\sqrt{\vartheta^{2} + |\nabla u|^{2}}} \nabla_{kj}^{2} u \\ &= \frac{\vartheta g^{ik}}{\sqrt{\vartheta^{2} + |\nabla u|^{2}}} \left(\frac{\vartheta'}{\vartheta} g_{kj} + \frac{\vartheta'}{\vartheta (\vartheta^{2} + |\nabla u|^{2})} g_{km} \sigma^{mr} \nabla_{r} u \nabla_{j} u - \nabla_{kj}^{2} u \right) \\ &\equiv \frac{\vartheta(u) g^{ik}}{\sqrt{\vartheta^{2}(u) + |\nabla u|^{2}}} A_{kj}(x, u, u, \nabla u, \nabla^{2} u), \end{split}$$

where $\vartheta = \vartheta(u)$ and we set

$$A_{kj}(x,z,u,p,r) = \frac{\vartheta'(z)}{\vartheta(z)}g_{kj}(p,u) + \frac{\vartheta'(z)}{\vartheta(z)\left(\vartheta^2(z) + |p|^2\right)}g_{km}(p,u)\sigma^{mr}p_rp_j - r_{kj}.$$

Here the dependence on x is hidden in σ^{mr} . Moreover, $g_{kj}(p, u) = p_k p_j + \vartheta^2(u)\sigma_{kj}$ and we artificially introduced z in the argument in order to distinguish increasing behavior from decreasing behavior in u. Since $\cot(z)$ is decreasing in z, we have

(6.2)
$$A_{ij}\left(x, z_2, u, \nabla u, \nabla^2 u\right) X^i X^j < A_{ij}\left(x, z_1, u, \nabla u, \nabla^2 u\right) X^i X^j$$

for all $z_1, z_2 \in (0, u(x, t))$ with $z_2 > z_1$ and all non-zero vectors $X \in TM_t$. Also write

$$h_j^i(x,z,u,p,r) = \frac{\vartheta(z)g^{ik}(p,z)}{\sqrt{\vartheta^2(z) + |p|^2}} A_{kj}(x,z,u,p,r).$$

Note that if u graphs a convex hypersurface, (6.2) implies that

$$h^i_j(x,z,u(x,t),\nabla u(x,t),\nabla^2 u(x,t)) > 0, \quad \forall \ 0 < z < u(x,t)$$

as a self-adjoint endomorphism and thus h_i^i lies in the domain of F. Write

$$\Phi(x, z, u, p, r) = -F(h_j^i(x, z, u, p, r))\sqrt{1 + \vartheta^{-2}(z)|p|^2}, \quad \forall \ 0 < z < u(x, t).$$

Choose

(6.3)
$$\lambda > \sup_{(x,t)\in E\times[T_{\delta},T]} \sup_{z\in Q(x,t)} \frac{\partial\Phi}{\partial z}(x,z,u(x,t),\nabla u(x,t),\nabla^2 u(x,t)),$$

where for $(x,t) \in E \times [T_{\delta},T]$ we defined $Q(x,t) = \{z : \frac{\pi}{4} \leq z \leq u(x,t)\}$. Such a choice of λ is possible since when z is away from 0, Φ is smooth.

In view of (6.1), there exists $(x_0, t_0) \in \overline{\Omega}$ not lying on the parabolic boundary of Ω such that for

$$\tilde{u} = ue^{-\lambda t}$$
 and $\tilde{w} = we^{-\lambda t}$,

16

there holds

$$0 < \tilde{u}(x_0, t_0) - \tilde{w}(x_0, t_0) = \max_{\bar{\Omega}} (\tilde{u} - \tilde{w}).$$

Therefore,

(6.4)
$$[w(x_0, t_0), u(x_0, t_0)] \subset Q(x_0, t_0).$$

Note that at (x_0, t_0) we have $\dot{\tilde{u}} \geq \dot{\tilde{w}}, \nabla u = \nabla w$ and $\nabla^2 u \leq \nabla^2 w$. Hence using (6.4), (6.3) and our choice of λ , we obtain, at (x_0, t_0) , that

$$(\dot{u} - \Phi(x_0, u, u, \nabla u, \nabla^2 u)) e^{-\lambda t_0} = \dot{\tilde{u}} - \Phi(x_0, u, u, \nabla u, \nabla^2 u) e^{-\lambda t_0} + \lambda \tilde{u} > \dot{\tilde{u}} - \Phi(x_0, w, u, \nabla u, \nabla^2 u) e^{-\lambda t_0} + \lambda \tilde{w} \ge \dot{\tilde{w}} - \Phi(x_0, w, w, \nabla w, \nabla^2 w) e^{-\lambda t_0} + \lambda \tilde{w} = (\dot{w} - \Phi(x_0, w, w, \nabla w, \nabla^2 w) e^{-\lambda t_0},$$

a contradiction, since both sides are zero in view of the flow equation.

$$\square$$

Proof of Theorem 1.2. From Lemma 6.5 we have $R_V((M_t)_V^+) \ge (M_t)_V^-$ everywhere for all $t \in (-T_S, -T)$ and any $\delta \in (0, \delta_0)$. By continuity therefore, sending $\delta \to 0$ we have $R_{\rm V}((M_t)_{\rm V}^+) \ge (M_t)_{\rm V}^-$ for all $t \in (-T_S, -T)$ and any V satisfying $\langle {\rm V}, e \rangle = 0$. Now we need some properties of R_V following from the fact that $\langle V, e \rangle = 0$:

- $R_{\mathrm{V}}^2 = \mathrm{id},$ $S \ge T \Rightarrow R_{\mathrm{V}}(S) \ge R_{\mathrm{V}}(T),$ $R_{\mathrm{V}} = R_{-\mathrm{V}},$ and

•
$$R_{\mathrm{V}} = R_{-\mathrm{V}}, \epsilon$$

• $S_{\rm V}^{\pm} = S_{-{\rm V}}^{\mp}$.

Thus we obtain

$$(M_t)_{\mathcal{V}}^+ = R_{\mathcal{V}}(R_{\mathcal{V}}((M_t)_{\mathcal{V}}^+)) \ge R_{\mathcal{V}}((M_t)_{\mathcal{V}}^-)$$
$$= R_{-\mathcal{V}}((M_t)_{-\mathcal{V}}^+) \ge (M_t)_{-\mathcal{V}}^-$$
$$= (M_t)_{\mathcal{V}}^+.$$

So we must have equality all the way through and hence the middle line implies

$$R_{-V}((M_t)^+_{-V}) = (M_t)^-_{-V}$$

for any V.

Therefore, M_t is invariant under R_V for any V satisfying $\langle V, e \rangle = 0$, hence it is a geodesic sphere for every $t \in (-T_S, -T)$ and thus it is a geodesic sphere for every $t \in (-T_S, 0).$

7. Non-geometric counterexamples

The examples in this section highlight the necessity of geometric invariance, without which our methods can not be applied. First, we have a flow that is parabolic - but not geometric - admitting a convex, ancient solution for which H becomes unbounded. As mentioned earlier, we do not yet know whether such singular behavior can occur for isotropic, geometric flows. Certainly this is not possible whenever a differential Harnack inequality holds and the spherical solutions are ancient as in Proposition 3.4.

7.1. Example. Let $\bar{\gamma}_t$, evolve by the curve shortening flow in \mathbb{R}^2 , and let $\gamma_t \subset \mathbb{S}^2$ be the inverse gnomonic projection (see, e.g., [3]) of $\bar{\gamma}_t$. The spherical radial functions

$$\rho: \mathbb{S}^1 \times [0,T) \to \mathbb{R}$$

evolve by

(7.1)
$$\partial_t \rho(\cdot, t) = -\kappa \frac{\sqrt{\sin^2 \rho + \rho_\theta^2}}{\sin \rho} \frac{\sin^2 \rho + \rho_\theta^2}{\tan^2 \rho + (\tan \rho)_\theta^2} (\cdot, t)$$

where, $\kappa(\cdot, t)$ is the curvature of the curve γ_t with radial function $\rho(\cdot, t)$.

To obtain the evolution equation, in polar coordinates we can express the curvature as follows:

$$\kappa = \frac{-\rho_{\theta\theta} \sin \rho + 2\rho_{\theta}^2 \cos \rho + \cos \rho \sin^2 \rho}{(\sin^2 \rho + \rho_{\theta}^2)^{\frac{3}{2}}}$$

Write $\bar{\rho}(\cdot, t)$ for the radial function of $\bar{\gamma}_t$. We recall from [3, p. 8] that $\bar{\rho} = \tan \rho$. Using this formula and the expression of κ we can write the curvature of $\bar{\gamma}_t$, $\bar{\kappa}(\cdot, t)$, as follows:

$$\kappa = \left(\frac{\bar{\rho}^2 + \bar{\rho}_{\theta}^2}{(1 + \bar{\rho}^2)(\sin^2 \rho + \rho_{\theta}^2)}\right)^{\frac{3}{2}} \bar{\kappa} = \left(\frac{\bar{\rho}^2 + 1}{\bar{h}^2 + 1}\right)^{\frac{3}{2}} \bar{\kappa}$$

Here $\bar{h} = \frac{\bar{\rho}^2}{\sqrt{\bar{\rho}^2 + \bar{\rho}_{\theta}^2}}$ is the support function of $\bar{\gamma}$. Therefore,

$$\begin{aligned} \partial_t \bar{\rho} &= -\bar{\kappa} (1+\bar{\rho}^2) \left(\frac{\bar{\rho}^2 + \bar{\rho}_{\theta}^2}{(1+\bar{\rho}^2)(\sin^2\rho + \rho_{\theta}^2)} \right)^{\frac{3}{2}} \frac{\sqrt{\sin^2\rho + \rho_{\theta}^2}}{\sin\rho} \frac{\sin^2\rho + \rho_{\theta}^2}{\tan^2\rho + (\tan\rho)_{\theta}^2} \\ &= -\bar{\kappa} (1+\bar{\rho}^2)^{\frac{3}{2}} \left(\frac{\bar{\rho}^2 + \bar{\rho}_{\theta}^2}{(1+\bar{\rho}^2)(\sin^2\rho + \rho_{\theta}^2)} \right)^{\frac{3}{2}} \frac{\sqrt{\sin^2\rho + \rho_{\theta}^2}}{\bar{\rho}} \frac{\sin^2\rho + \rho_{\theta}^2}{\tan^2\rho + (\tan\rho)_{\theta}^2} \\ &= -\bar{\kappa} \frac{\sqrt{\bar{\rho}^2 + \bar{\rho}_{\theta}^2}}{\bar{\rho}}. \end{aligned}$$

The curve shortening flow in \mathbb{R}^2 has non-trivial ancient solutions, the Angenent ovals. Thus, there exists a non-spherical, convex, ancient solution to the flow (7.1). This ancient solution converges backwards in time to a lune (the intersection of two hemispheres) with a pair of antipodal points at which κ is unbounded.

The second example exhibits a (non-geometric) flow for which the symmetry of the backwards limit is not preserved. That is, the backwards limit is an equator, yet the flow is not by geodesic spheres. This example is not as troubling as the first, since the non-geometric nature of the flow prohibits the use of the Aleksandrov reflection technique in Section 6, but it does illustrate the necessity of geometric invariance for our methods.

7.2. *Example.* Using the notation of the previous example, let $\bar{\gamma}_t$ evolve by the affine normal flow in \mathbb{R}^2 :

$$\partial_t \bar{\rho} = -\bar{\kappa}^{\frac{1}{3}} \frac{\sqrt{\bar{\rho}^2 + \bar{\rho}_\theta^2}}{\bar{\rho}}.$$

Let γ_t be the inverse gnomonic projection. Then the spherical radial functions evolve by

$$\rho: \mathbb{S}^1 \times [0,T) \to \mathbb{R}$$

(7.2)
$$\partial_t \tan \rho(\cdot, t) = -\kappa^{\frac{1}{3}} \frac{\sqrt{\sin^2 \rho + \rho_{\theta}^2}}{\sin \rho} (\cdot, t).$$

Origin-centered ellipses are ancient solutions to the affine normal flow in \mathbb{R}^2 (see, for example, [18]). Thus, there exists a non-spherical, convex, ancient solution to the flow (7.2). This solution converges backwards in time to an equator.

References

- B. Andrews, X. Han, H. Li, and Y. Wei, Non-collapsing for hypersurface flows in the sphere and hyperbolic space, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) XIV (2015), no. 1, 331–338.
- [2] S. Angenent, Shrinking doughnuts, In: "Nonlinear diffusion equations and their equilibrium states" 3, N. G. Lloyd, W. M. Ni, L. A. Peletier, J. Serrin (eds.), Birkhäuser Boston, MA, 1992, 21–38.
- [3] F. Besau and E. Werner, The spherical convex floating body, Adv. Math. 301 (2016), 867–901.
- [4] W. Blaschke, "Kreis und kugel", 2. ed., De Gruyter, 1956.
- [5] T. Bourni and M. Langford, Classification of convex ancient free boundary mean curvature flows in the ball, arxiv:2204.06710, 2022.
- [6] T. Bourni, M. Langford, and A. Mramor, On the construction of closed nonconvex nonsoliton ancient mean curvature flows, Int. Math. Res. Not. IMRN 2021 (2021), no. 1, 757–768.
- [7] T. Bourni, M. Langford, and G. Tinaglia, Collapsing ancient solutions of mean curvature flow, J. Differential Geom. 119 (2021), no. 2, 187–219.
- [8] P. Bryan and M. N. Ivaki, Harnack estimate for mean curvature flow on the sphere, Asian J. Math. 24 (2020), no. 1,165–175.
- [9] P. Bryan, M. N. Ivaki, and J. Scheuer, Harnack inequalities for evolving hypersurfaces on the sphere, Comm. Anal. Geom. 26 (2018), no. 5, 1047–1077.
- [10] P. Bryan and J. Louie, Classification of convex ancient solutions to curve shortening flow on the sphere, J. Geom. Anal. 26 (2016), no. 2, 858–872.
- [11] P. Daskalopoulos, R. Hamilton, and N. Sesum, Classification of compact ancient solutions to the curve shortening flow, J. Differential Geom. 84 (2010), no. 3, 455–464.
- [12] M. Do Carmo and F. Warner, Rigidity and convexity of hypersurfaces in spheres, J. Differential Geom. 4 (1970), no. 2, 133–144.
- [13] C. Gerhardt, "Curvature problems", Series in Geometry and Topology, vol. 39, International Press of Boston Inc., Sommerville, 2006.
- [14] R. Hashhofer and O. Hershkovits, Ancient solutions of the mean curvature flow, Comm. Anal. Geom. 24 (2016), no. 3, 593–604.
- [15] G. Huisken and C. Sinestrari, Convexity estimates for mean curvature flow and singularities of mean convex surfaces, Acta Math. 183 (1999), no. 1, 45–70.
- [16] G. Huisken and C. Sinestrari, Mean curvature flow singularities for mean convex surfaces, Calc. Var. Partial Differential Equations 8 (1999), no. 1, 1–14.
- [17] G. Huisken and C. Sinestrari, Convex ancient solutions of the mean curvature flow, J. Differential Geom. 101 (2015), no. 2, 267–287.
- [18] M. N. Ivaki, Classification of compact convex ancient solutions of the planar affine normal flow, J. Geom. Anal. 26 (2016), no. 1, 663–671.
- [19] X. Liu and C. L. Terng, Ancient solutions to mean curvature flow for isoparametric submanifolds, Math. Ann. 378 (2020), no. 1–2, 289–315.
- [20] S. Lynch and H. T. Nguyen, Pinched ancient solutions to the high codimension mean curvature flow, Calc. Var. Partial Differential Equations 60 (2021), no. 1, art. 29.
- [21] M. Makowski and J. Scheuer, Rigidity results, inverse curvature flows and Alexandrov-Fenchel type inequalities in the sphere, Asian J. Math. 20 (2016), no. 5, 869–892.
- [22] H. T. Nguyen, Convexity and cylindrical estimates for mean curvature flow in the sphere, Trans. Amer. Math. Soc. 367 (2015), no. 7, 4517–4536.
- [23] T. Rado, The isoperimetric inequality on the sphere, Amer. J. Math. 57 (1935), no. 4, 765– 770.
- [24] S. Risa and C. Sinestrari, Strong spherical rigidity of ancient solutions of expansive curvature flows, Bull. Lond. Math. Soc. 52 (2020), no. 1, 94–99.

- [25] J. Scheuer, Non-scale-invariant inverse curvature flows in hyperbolic space, Calc. Var. Partial Differential Equations 53 (2015), no. 1, 91–123.
- [26] B. White, The nature of singularities in mean curvature flow of mean-convex sets, J. Amer. Math. Soc. 16 (2002), no. 1, 123–138.

DEPARTMENT OF MATHEMATICS, MACQUARIE UNIVERSITY, NSW 2109, AUSTRALIA, paul.bryan@mq.edu.au

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TECHNISCHE UNI-VERSITÄT WIEN, WIEDNER HAUPTSTR 8-10, 1040 WIEN, AUSTRIA, mohammad.ivaki@tuwien.ac.at

SCHOOL OF MATHEMATICS, CARDIFF UNIVERSITY, SENGHENNYDD ROAD, CARDIFF CF24 4AG, WALES, scheuerj@cardiff.ac.uk

20