

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/151472/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Gallimore, Awen 2022. Tumor-associated high endothelial venules: inroads enabling immune control of cancer progression. *Cancer Immunology Research* 10 (4) , 371. 10.1158/2326-6066.CIR-22-0112

Publishers page: <http://dx.doi.org/10.1158/2326-6066.CIR-22-0112>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



## Tumor-associated high endothelial venules: Inroads enabling immune control of cancer progression

Awen Gallimore

Systems Immunity University Research Institute, Cardiff University, UK

Email: [gallimoream@cardiff.ac.uk](mailto:gallimoream@cardiff.ac.uk)

### Abstract

*Infiltration of lymphocytes into solid tumors represents a significant bottleneck to successful control of tumor growth. The nature of tumor blood vessels is an important factor governing both quantitative and qualitative features of the immune infiltrate. In this issue, Sawada and colleagues identify a genetic signature for blood vessels, most notably high-endothelial venules (HEV), which are associated with tertiary lymphoid structures (TLS) and improved clinical outcome.*

*See related article by Sawada et al. XXX*

HEVs are specialized blood vessels that are normally found only in secondary lymph organs. They serve to recruit L-selectin–expressing lymphocytes into the lymph node (LN) parenchyma where they encounter cognate antigen. Antigen stimulation drives expansion of HEV networks through proliferation of endothelial cells and the co-operative activities of vascular, stromal and immune cells. These changes facilitate increased blood flow and lymphocyte trafficking to support development of adaptive immunity.

HEVs are present in human cancers where they are often associated with organized TLS and/or a greater degree of lymphocytic infiltration. Many studies show that TLS are associated with longer survival and/or are predictive of favorable responses to immunotherapy (1). In mouse models, intratumoural HEVs can be induced in several ways including depletion of regulatory T cells, adoptive T-cell transfer and delivery of anti–PD-L1 and anti-VEGFR2 (1, 2). Overall, these studies indicate that tumors successfully treated with immunotherapy contain HEVs (1, 2).

Key questions emerging from the above relate to the molecular identity and function of intratumoural HEVs. In this issue, Sawada and colleagues compare the transcriptomes of HEVs and non-HEVs in human breast cancer samples. HEVs and non-HEVs are distinguished using a monoclonal antibody, MECA-79, which binds to the sulfated, fucosylated glycoprotein, peripheral node addressin (PNAd); the principal marker of HEV. Laser-capture microdissection is used to isolate PNAd<sup>+</sup> and PNAd<sup>−</sup> endothelial cells from TLS-rich and TLS-free (no PNAd<sup>+</sup> cells) areas for gene expression analyses, which reveals a molecular signature identifying tumor HEVs. The authors focus on two upregulated genes, *TSPAN7* and *MEOX2*, which they find are expressed on almost all tumor HEVs and are associated with high T-cell and B-cell infiltrates and improved survival. *TSPAN7*, which encodes tetraspanin 7, is compatible with the tumor HEVs having a role in extravasation of lymphocytes whereas *MEOX2* encodes for a negative regulator of angiogenic activity. Previous work in mice indicates that vascular normalization using anti-VEGFR2 promotes HEV formation, thus suggesting that HEV development is dependent on repression of angiogenesis. Several other potentially interesting genes that could hold clues to understanding tumor HEV function as well as how they are formed and maintained are also uncovered in the study.

Early studies drew on the LN paradigm of HEV function to explain the role of tumor-associated HEVs. However, it is becoming increasingly clear that whilst tumor and LN HEV share

many features, there are disparities, which have been revealed by differential PNA<sup>+</sup> staining as well as gene-expression profiling of tumor and LN HEV. This is not surprising given a recent study which highlighted significant HEV heterogeneity even within LN (3), specifically indicating that HEV exist in different functional states depending on the activation status of the LN. An interesting finding of the study by Sawada et al is that there is an overlap between PNA<sup>+</sup> and PNA<sup>-</sup> vessels isolated from TLS-rich areas of the tumor, characterized by TSPAN<sup>hi</sup>PNA<sup>-</sup> vessels. It is possible that these vessels also contribute to the pool of tumor-infiltrating lymphocytes and a successful immune response.

Overall, this study provides a molecular signature of tumor vessels that may be useful for interrogating cancer datasets and for generating new hypotheses aimed at identifying the critical features of HEV or HEV-like vessels that can facilitate robust antitumor immune responses. Such information will help identify routes for inducing *useful* vessels to enhance the efficacy of current and novel immunotherapies.

## References

1. Schumacher TN, Thommen DS. Tertiary lymphoid structures in cancer. *Science*. 2022 Jan 7;375(6576):eabf9419. doi: 10.1126/science.abf9419. Epub 2022 Jan 7.
2. Colbeck EJ, Ager A, Gallimore A, Jones GW. Tertiary Lymphoid Structures in Cancer: Drivers of Antitumor Immunity, Immunosuppression, or Bystander Sentinels in Disease? *Front Immunol*. 2017 Dec 19;8:1830. doi: 10.3389/fimmu.2017.01830. eCollection 2017.
3. Veerman K, Tardiveau C, Martins F, Coudert J, Girard JP. Single-Cell Analysis Reveals Heterogeneity of High Endothelial Venules and Different Regulation of Genes Controlling Lymphocyte Entry to Lymph Nodes. *Cell Rep*. 2019 Mar 12;26(11):3116-3131.e5. doi: 10.1016/j.celrep.2019.02.042.
4. Sawada-J et al – THIS PAPER

## Acknowledgement

Awen Gallimore is supported by grants from Breast Cancer Now, The Wellcome Trust and Cancer Research UK