

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/154510/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Nagendrababu, Venkateshbabu, Vinothkumar, Thilla Sekar, El-Karim, Ikhlas, Rossi-Fedele, Giampiero, Dogramaci, Esma J., Dummer, Paul M. H. and Duncan, Henry F. 2023. Dental patient-reported outcomes in endodontics - a narrative review. *Journal of Evidence-Based Dental Practice* 23 (1) , 101805. 10.1016/j.jebdp.2022.101805

Publishers page: <https://doi.org/10.1016/j.jebdp.2022.101805>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



DENTAL PATIENT-REPORTED OUTCOMES IN ENDODONTICS - A NARRATIVE REVIEW

VENKATESHBABU NAGENDRABABU^a, THILLA SEKAR VINOTHKUMAR^{b,c}, IKHLAS EL-KARIM^d, GIAMPIERO ROSSI-FEDELE^e, ESMA J. DOĞRAMACI^e, PAUL M. H. DUMMER^f, HENRY F. DUNCAN^g

^a Department of Preventive and Restorative Dentistry, College of Dental Medicine, University of Sharjah, Sharjah, UAE

^b Department of Restorative Dental Sciences, College of Dentistry, Jazan University, Jazan, Saudi Arabia

^c Department of Conservative Dentistry and Endodontics, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences, Chennai, India

^d School of Medicine Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, UK

^e Adelaide Dental School, University of Adelaide, Adelaide, South Australia, Australia

^f School of Dentistry, College of Biomedical and Life Sciences, Cardiff University, Cardiff, UK

^g Division of Restorative Dentistry & Periodontology, Dublin Dental University Hospital, Trinity College Dublin, Lincoln Place, Dublin 2, Ireland

CORRESPONDING AUTHOR

Venkateshbabu Nagendrababu,
Department of Preventive and Restorative Dentistry,
College of Dental Medicine,
University of Sharjah,
Sharjah,
UAE.

Mail id: hivenkateshbabu@yahoo.com, vnagendrababu@sharjah.ac.ae

Highlights

The main dPROs following endodontic treatment include pain, tenderness, tooth function, need for further intervention, adverse effects and Oral Health–Related Quality of Life.

dPROs are essential for endodontic treatment as they enable dentists and patients to discuss and identify the most appropriate management options, whilst providing the opportunity for researchers to improve the methodology and design of future clinical trials to ensure the patient's interests are at the center of the study.

Clinicians and researchers working in Endodontology should focus on patient wellbeing and routinely use dPROs

ABSTRACT

Recently in oral healthcare settings, the focus of assessing treatment outcomes has shifted from the perspective of the clinician towards that of the patient. Endodontology is a specialty of dentistry concerned with the prevention and treatment of pulp and periapical diseases. Research in endodontology and its associated treatment outcomes have focused mainly on clinician-reported outcomes (CROs) and not patient-reported outcomes (PROs). As a result, there is a need to emphasize the importance and relevance of dPROs to researchers and clinicians. The aim of this review is to present an overview of dPROs and dPROMs within clinical endodontics in an attempt to create a better understanding of the patient experience, highlight the need to place the patient at the center of treatment, enhance patient care and encourage more research into dPROs. The key dPROs following endodontic treatment include pain, tenderness, tooth function, need for further intervention, adverse effects (exacerbation of symptoms, tooth discoloration) and Oral Health–Related Quality of Life. dPROs are important following endodontic treatment because they assist clinicians and patients when they discuss and select the most appropriate management options, help clinicians make decisions on pre-operative assessment, prevention and treatment, and improve the methodology and design of future clinical studies. Clinicians and researchers in endodontology should prioritize patient welfare and undertake routine analyses of dPROs using appropriate and robust measures. Due to the lack of agreement over the reporting and definition of endodontic treatment outcomes, a project to define a "Core Outcome Set for Endodontic Treatment Methods (COSET)" is currently ongoing. In the future, a new and exclusive assessment tool should be developed to reflect the viewpoints of patients receiving endodontic treatment more accurately.

KEYWORDS

Endodontics, Oral Health–Related Quality of Life, Pain, Patient reported outcomes, Patient-reported outcome measures

INTRODUCTION

PATIENT-REPORTED OUTCOMES and PATIENT-REPORTED OUTCOME MEASURES

Patient-reported outcomes (PROs) are reports of a patient's health status that come directly from the patient without interpretation by a clinician,¹ whereas clinician-reported outcomes (CROs) are a report of a patient's health status by a trained healthcare professional.² Patient-reported outcome measures (PROMs) are objective or subjective measurements used to evaluate the effectiveness of an intervention using tools or instruments, generally in the form of self-reported questionnaires.³ It should be noted that although by definition the patient's interpretations should be unfiltered by a clinician, a PRO in reality represents more, with the subjective views of the patient including personal weights, emotions and the consideration of both values and expectations being considered. The role of PROs is significant in enhancing clinical care because it strengthens the relationship between clinicians and patients⁴ and places the patient at the center of the process.

PROs include:

- Symptoms associated with a disease/condition or therapeutic side-effects such as pain, anxiety, or and the need for pain relief;
- Functional outcomes such as physical, emotional, or cognitive functioning; and
- Multi-dimensional categories such as Health-Related Quality of Life (HRQoL).^{5,6}

DENTAL PATIENT-REPORTED OUTCOMES and DENTAL PATIENT-REPORTED OUTCOME MEASURES

In dentistry, any report of a patient's oral health status (or wider general health if relevant to oral health) that comes directly from the patient, without interpretation by a clinician or anyone else, is referred to as a dental patient-reported outcome (dPRO).⁷ dPROs are being utilized increasingly in oral health research to explore patients' perceptions of the success and impact of dental treatments.⁸ A dental patient-reported outcome measure (dPROM) is an instrument, questionnaire, scale, or survey that measures or captures dPROs.⁷ The dPROM scores can be reported numerically or

graphically to the clinician at chairside,⁹ and variations between pre- and post-treatment dPROM ratings can be utilized to determine the outcome of treatment as well as assist patients in making more informed treatment choices.¹⁰ Notably, a methodological study by Tao et al., identified only 76 of 315 randomised controlled trials in orthodontics used dPROs as either primary or secondary outcomes.¹¹ dPROs are critical for the development of evidence-based dentistry in a pragmatic primary care setting,¹² as they help to improve research impact and ‘real world’ relevance.¹³ They are essential for value-based oral health care, which is concerned with improving oral health outcomes for patients in relation to the associated costs.¹⁴ The principle dPRO used to date is the Oral Health-Related Quality of Life (OHRQoL), which is defined as “people’s perspective on their oral health status including eating, sleeping and engaging in social interaction; their self-esteem; and their satisfaction with respect to their oral health”.¹⁵

SCOPE OF ENDODONTOLOGY

Endodontology is “concerned with the study of the form, function, and health of, injuries to and diseases of the dental pulp and periradicular region, their prevention and treatment; the principal disease being apical periodontitis, caused by infection”.¹⁶ The principal treatment interventions include vital pulp treatment, root canal treatment, root canal retreatment, surgical endodontics, and regenerative endodontic therapies.¹⁷ Endodontic treatment focuses on the prevention or resolution of pulp and apical disease with the overarching aim being to improve the patients’ quality of life by preserving their natural teeth. As a consequence, patients can function optimally, allowing them to speak, eat, and smile more naturally, as well as improve their dentofacial esthetics, self-confidence, and psychosocial wellbeing.^{17–20}

AIM OF THE REVIEW

Clinicians and researchers working within endodontology should be concerned about the welfare of patients and are well-placed to routinely assess PROs using relevant tools and measures. Hence, the aim of this review is to provide an overview of dPROs and dPROMs to allow better understanding of the patient experience, consider the need to place the patient at the center of the outcome to ultimately improve patient care, as well as to promote dPRO research in endodontology.

dPROs IN ENDODONTICS

Assessing the outcome of endodontic treatment has traditionally focused on CROs such as clinical and radiographic examinations, microbial culture and analysis, and pulp sensibility testing. On the other hand, PROs such as pain, and quality of life have been undervalued.^{16,21,22} For example, De-Deus & Canabarro²³ reported that CROs (*e.g.* radiographic healing) were the primary outcome of clinical research comparing single- versus multiple-visit root canal treatment. The range of clinical procedures in endodontics have been adopted not only for their efficacy and biological consequences but also for their ability to minimize patient suffering.²⁴ The results of a recent scoping review revealed that there are 300 CROs but only 114 PROs contained within studies on root canal treatment, retreatment, and apexification published during the last four decades (1980–2020).²⁵ Evidently, PROs are underutilized in endodontics when compared with CROs.²⁵ As a result, there is a paucity of evidence linking endodontic treatment outcomes to patient-reported factors, such as pain, discomfort, the need for medication, and the cost of the procedure.^{26,27}

In general, the use of PROs in clinical studies has risen in recent years, with 6168 (45.1%) of the 13,666 trials recorded in the Australian New Zealand Clinical Trials Registry incorporating a PRO.^{5,6} Regulatory bodies including the US Food Drug Administration and the European Medicines Agency both urge that data from PROs be considered in the assessment of clinical trial endpoints. Within the development of clinical guidelines using a GRADE-framework, there is an insistence that outcomes consider the patient and are patient-centered, as a minimum.²¹ In the last four decades, a decrease in the reporting of CROs such as radiographic healing and success of endodontic treatment, which includes root canal treatment, retreatment and apexification, while a small but steady increase in the reporting of PROs such as pain assessment, and quality of life has occurred.²⁵

PROs can be used at any time and provide the patient's perspective for a specific health status; however, they are most often used to evaluate the outcome of treatment and the overall quality of care, both of which are crucial to the well-being of patients. For example, when it comes to understanding whether root canal treatment is successful or not, the individual patient, who is generally unaware of the condition of the periapical tissues of their root-filled tooth, may not always agree with the criteria considered by their

clinician. If the tooth remains symptom-free following root canal treatment despite radiographic indications of ongoing periapical pathosis, the treatment may still be judged successful by the patient.^{28,29} On the other hand, the patient may be dissatisfied with the treatment even if no objective evidence of periapical pathosis is present, as pain or other problems may persist.³⁰ As a result, a holistic view for evaluating the outcome of endodontic treatment should ideally include a set of core criteria assessing the patient's perspective on the outcome of treatment.^{29,31,32}

IMPORTANCE of dPROs in ENDODONTICS

The importance of dPROs following endodontic treatment includes^{6,19,33,34}:

- Helping clinicians and patients select the most appropriate treatment by providing a more comprehensive view of treatment costs and benefits. For example, a recurring debate in clinical practice involves whether to save a natural tooth by initiating root canal treatment, or whether to extract followed by replacement. This is not a simple cost-benefit analysis; however, preserving the natural tooth offers other benefits with tooth loss being shown to have a negative influence on the overall OHRQoL.^{19,35}
- Providing guidance to aid clinical decision-making in the assessment, prevention, and management of pulpal and periapical diseases. For example, a study comparing patients' quality of life following surgical endodontic treatment using two different techniques (a traditional technique without an operating microscope and a technique using an operating microscope and minimal osteotomy). The results of this study will be useful for the clinician to select the most appropriate technique, from a patient's perspective, when performing surgical endodontics.³⁶
- Developing clinical practice guidelines in endodontology and other disciplines should always include essential PROs, ranked into those that are "most critical," "critical," or "important".^{22,27} The *European Society of Endodontology* (ESE) is in the process of developing clinical practice guidelines for pulpal and periapical diseases. PROs play an important role in the development of these guidelines as the patient is a key stakeholder in guideline development.²²
- Analyzing the perspective of the patient on the effectiveness of treatment are valuable sources of information that can be overlooked when the assessment and

opinion of the clinician is used to filter the patient's perspective of the clinical intervention. The effects of many intervention are not well understood by clinicians, so it is important to obtain the patient's perspective and view of the treatment itself and its outcome.¹

- Allowing an estimation of treatment benefits and risks separately from effectiveness measures used traditionally, allows PROs to be used as effective research endpoints for the development and evaluation of new drugs and treatments. For example, during vital pulp treatment the use of some hydraulic calcium silicate cements is associated with tooth discoloration, albeit with excellent healing outcomes.³⁷ An assessment of aesthetics as an important component of OHRQoL has led to shifts in the criteria for evaluating new endodontic materials and therapies.
- Helping to improve the quality of future clinical trial methodology and study design.

dPROs FOR ENDODONTIC TREATMENT – EUROPEAN SOCIETY OF ENDODONTOLOGY (ESE) INITIATIVE

The ESE is currently engaged in the process of developing new practice guidelines for the treatment of pulpitis, the non-surgical treatment of apical periodontitis, the surgical treatment of apical periodontitis and the regenerative treatment of apical periodontitis for the benefit of both clinicians and patients.²¹ To assist robust systematic reviews of the literature, online Delphi surveys and meetings were conducted to identify and rank (*critical*, and *important*) the relevant patient and clinician-reported outcome measures with follow-up intervals. As part of this project, recommendations were also offered regarding the acceptable minimum follow-up duration. In summary, the results of the Delphi survey and online meeting concluded that the “*critical*” outcomes were “pain and tenderness” whereas “*important*” outcomes were “tooth function, need for further intervention, adverse effects , and OHRQoL”.²²

dPROs USED IN ENDODONTICS

Pain

Pain arising from the pulp or periapical region before, during and after endodontic treatment is of prime concern for both patients and clinicians.²⁶ Pain is a subjective feeling that can occur spontaneously or in response to touch or biting (tenderness) and is the primary reason people seek dental care.³⁸ In dentistry, pain is an important dPRO in addition to its association with the Oral Health Impact Profile (OHIP) tool³⁹ and has a major impact on one of the four dimensions of OHRQoL (orofacial pain).⁴⁰ The outcome measurement tools commonly used for objective assessment of pain are clinical examination, numerical rating scale (NRS), visual analogue scale (VAS), verbal rating scale (VRS) and Likert-scale.⁴¹⁻⁴³ The VAS used to record patients' self-reported pain ranges from either 1 to 10 or 1 to 100.^{41,44} In the VRS, patients are requested to score the intensity of pain ranging from 1 to 4 as none, slight, moderate and severe.⁴³ Similarly, in an 11-point NRS ranging from 0 (no pain) to 10 (worst pain), patients are asked to mark a numbered line on a chart corresponding to their level of pain.⁴⁵ Pain levels can be further divided or dichotomized into: 0 = none, 1-3 = mild, 4-7 = moderate and 8-10 = severe.⁴⁶

The majority of endodontic outcome studies do not consider pain as a criterion for failure possibly because of its subjective nature.⁴⁷⁻⁵⁰ However, pain on tooth percussion can indicate failure of endodontic treatment as this sign is an indicator of underlying periapical inflammation that may not be captured by radiographic examination alone.⁵¹ Pain may also play an important role in a patients' decision-making process in selecting between root canal treatment (tooth retention) versus extraction followed by an implant-supported crown.⁵² Notably, the pain experienced by patients during their current treatment may influence their future treatment decisions, depending on their personal endurance limits.⁵³

Other subjective pain-related outcomes that do not have dedicated outcome measurement tools include: 'flare-up' and need for medication, which can be recorded through a careful clinical history and be considered as "*critical*" outcomes.²² A 'flare-up' is an unpleasant experience of severe pain and swelling following endodontic treatment that necessitates an unscheduled emergency visit to a clinician.⁵⁴ The reported incidence rate of a 'flare-up' after root canal treatment ranges from 2.3 to 3.2%^{55,56} as opposed to 10.4% in regenerative endodontic procedures.⁵⁷ It is usually relieved by prophylactic

antibiotics, steroidal and non-steroidal anti-inflammatory drugs, opioid and non-opioid analgesics, but in a true 'flare-up' the treating clinician is contacted.⁵⁸ Although there is no tool to measure the need for medication,⁴³ this can be recorded at subsequent clinical visits or by a patient in a diary.²²

Oral Health–Related Quality of Life (OHRQoL)

OHRQOL is the most significant dPRO and a major element of HRQoL, as it is directly related to the effects of oral disease and dental treatment on patients.⁵⁹ To measure OHRQoL, instruments (questionnaires) are employed to gather patient-perceived impact, and instrument scores are utilized to quantify patient suffering.⁵⁹ OHRQoL is a global outcome measure that encompasses other measures considered by the European Society of Endodontology.⁶⁰ Nonetheless, some instruments used in this field are often not sufficiently broad to capture all aspects of OHRQoL.⁶⁰ OHRQoL is highly beneficial in areas with limited oral health care resources because it can be used to direct limited resources to the patients who require them most,⁶¹ though only in conjunction with the relevant clinical outcome measures.³⁹

Overall, root canal treatment is associated with an improvement in OHRQoL.^{18,62–65} Similar OHRQoL scores have been reported where root canal treatment is compared with other treatment interventions in the primary care setting.⁶⁴ Such findings assist in discussing the negative perception of root canal treatment reported by patients compared with other dental treatment modalities. Similarly, negative beliefs regarding endodontic surgery have been confuted with regard to OHRQoL.⁶⁶ Recently published clinical endodontic studies highlight a lack of correlation between the healing of apical periodontitis, a commonly assessed outcome in endodontics, and OHRQoL;^{63,67} however, association with other PROs remains a possibility.⁶³

Inconsistent findings regarding OHRQoL have been demonstrated when endodontic treatment is compared with extractions. One study reported no significant differences,⁶⁴ whereas another, with a shorter recall period, highlighted that participants receiving root canal treatment had a significant improvement in OHRQoL compared with those having an extraction.¹⁹ It should be noted that in the above studies, teeth were also extracted for non-endodontic reasons. Therefore, direct comparison of the management of endodontic

diseases may not be appropriate. Subjects receiving rehabilitation of an edentulous space with a single implant-supported prosthesis versus root canal treatment reported comparable OHRQoL scores.³³ It appears that no study has directly compared endodontic treatment with monitoring of apical periodontitis, the latter being a highly prevalent disease globally.⁶⁸ Notably, a high prevalence of apical periodontitis associated with root-filled teeth has been reported,⁶⁸ therefore, the presence of endodontic diseases may influence OHRQoL scores.

As clinical outcomes can be influenced by pre-, intra- and post-operative factors;⁶⁹ the same should be expected for OHRQoL. Pre-operative factors may encompass both subject-level and tooth-level factors. Subject-level factors that have been associated with better OHRQoL include patient age and gender, with the elderly (those over 65 years of age) and female subjects reporting better quality of life;⁷⁰ however, there are partially contrasting findings in a second comparable study where men and subjects younger than 35 years of age had poorer health outcomes in some factors.²⁶ The latter study also reported greater OHRQoL scores amongst those of higher socio-occupational status.²⁶ Pre-operative tooth-level factors include the type of tooth, with anterior teeth having a more significant association with higher OHRQoL scores compared with molar teeth.^{26,70} The presence of pre-operative pain was also associated with negative outcomes,⁷⁰ with improvement of OHRQoL scores being more pronounced in teeth with vital pulps compared with those with necrotic pulps with a history of missing teeth.⁶² Intra-operative factors have focused mostly on instrumentation protocols, with limited differences amongst them.⁷¹⁻⁷⁴ Considering the multitude of potentially relevant tooth-related and/or intra-operative factors such as diagnosis, irrigant solution sequence and delivery, number of sessions, use of intra-canal medications, root canal filling material and techniques, it is difficult to reach a definitive conclusion on the role of a specific treatment step on OHRQoL. Finally, the role of the operator may also influence the OHRQoL. Some studies have reported no difference in outcomes of root canal treatment when it is carried out by either dental students, general dentists or specialists,^{62,65} whereas another study suggested more favorable responses in some domains when treatment was carried out by general dentists, which was justified by the likely increased complexity in the cases managed by specialists.⁷⁵

The following quality of life questionnaires have been used in Endodontics :²⁰ OHIP-14, Modified health-related quality of life index, OQOL measure 6-item and 12-item Versions, OHIP-17, Modified OHIP-49, Ad hoc post-operative QoL questionnaire (POQoL), American chronic pain association quality of life scale (QoLS) and Patient Perception questionnaire. An important concluding note is that endodontic research does not necessarily use previously validated OHRQoL instruments. Most studies use versions of OHIP, while other studies use instruments that appear to be created *ad hoc* for the purposes of the study. This, together with the high variability of clinical techniques adopted globally, makes reaching a definitive conclusion regarding the impact of endodontic treatment in OHRQoL difficult. Further understanding of the association between endodontic treatments and OHRQoL may be supported by the application of a disease-specific instrument for the measurement of this crucial outcome in the discipline. The development and validation of the latter should take into consideration the Food and Drug Administration Guidance on the development of PROs in medicine.⁷⁶

Function

Being able to bite and chew with an endodontically-treated tooth is considered an “important” outcome.²² This has been indirectly evaluated by recording the incidence of vertical fractures,⁷⁷ chewing ability⁶³ and patient comfort/discomfort.²⁶ Objective assessment of vertical root fractures of root canal treated teeth is difficult due to the fact that such teeth are often extracted and are unavailable for further analysis. The reported incidence rate for vertical root fractures (1.2%) has been evaluated during subsequent treatment procedures, including periapical surgery.⁷⁷ Chewing ability following root canal treatment has been evaluated using a subjective patient-centered questionnaire.⁶³ Similarly, patient comfort following any endodontic treatment could be assessed via a VAS ranging from 1 to 10²⁶ or an NRS-11 ranging from 0-10.⁴⁵

Adverse effects

Adverse effects associated with endodontic treatment, for managing teeth with pulpitis and apical periodontitis, are also considered “important” outcomes.²² Discoloration due to the leaching of root canal sealers or endodontic materials in the pulp chamber is one of the most common adverse effects of root canal treatment. Discoloration can be measured subjectively or objectively measured using a spectrophotometer.⁷⁸ That said

although discoloration can be used as a quantifiable outcome measured that does not make it a PRO, achieving that would involve questions such as to whether the discolored tooth made the patient feel emotional distress, lack confidence or have esthetic impairment. Many outcome assessment studies have recommended thorough debridement of the pulp chamber and internal bleaching to prevent sealer-induced discoloration.^{79–81} Other rare adverse effects include allergic reactions to certain chemical constituents of dental materials including root canal sealers containing formaldehyde or zinc oxide,^{82–84} intracanal medicaments such as tetracycline in Ledermix paste⁸⁵ and root canal irrigants such as sodium hypochlorite⁸⁶ and chlorhexidine.⁸⁷ Calcium hydroxide extrusion beyond the root apex can also result in damage to connective tissue and skin necrosis.⁸⁸ All such adverse effects are likely to impact on the OHRQoL of patients.⁸⁹

Patient satisfaction

Patient satisfaction is the feedback on the quality of treatment delivered adds value when assessing the patients' perception of a successful or unsuccessful treatment and is usually measured by means of a questionnaire.²⁸ Satisfaction is dependent on various factors, such as tooth retention, healing, functionality, absence of symptoms, and the patient's own perception of care.^{28,30,31} Therefore the tool should be comprehensive including all relevant aspects (outcome-related and process-related quality of care) of satisfaction.³¹ An example includes one that consists of eight items measuring the patients' perception of root canal treatment, present pain intensity, satisfaction, chewing ability and cost of treatment that was used in a cohort study conducted in the Swedish public dental health service.²⁹ In that study, 75% of patients were satisfied with the root canal treatment procedure and were willing to undergo the procedure again.²⁹

REPORTING OF ENDODONTIC OUTCOME STUDIES

An increase in the number and quality of studies reporting the outcomes of different endodontic treatment modalities has recently been reported.⁹⁰ These studies varied in design and included mostly comparative prospective and retrospective cohort studies, randomized control trials and case series. This variability in study design is a well-known cause of heterogeneity that negatively impacts the synthesis of evidence. This is compounded by evidence of great variability in the reporting outcomes included in these

studies,^{91,92} The ambiguity and variation in how outcomes are described across different studies is a universal problem in medical research⁹³ that results in inconsistencies and inefficient use of clinical research findings that greatly impact clinical care.

Attempts has been made to develop classifications and taxonomies for mapping outcomes into specific domains to facilitate consistent reporting. For instance, Fletcher and Fletcher's⁹⁴ 5Ds model mapped outcomes into 5 domains: death, disease, discomfort, disability, dissatisfaction. This model has adopted and modified with addition of a 6th domain "destitution" to map endodontic treatment outcomes into patient-centered outcomes.²⁵

Other systems/classifications such as that developed in the Patient-Reported Outcomes Measurement Information System (PROMIS),⁹⁵ provides a structure for classifying patient-reported measures only, whereas outcomes collected by health care providers, and those affecting wider society are not included. An ideal taxonomy/classification for outcomes must clearly differentiate between high-level outcome types and cover potential outcomes in a logical hierarchical structure of sufficient scope and granularity to be applied across different outcome studies.⁹⁶ In this regard Dodd et al.⁹⁶ developed a comprehensive 38-item taxonomy for reporting outcomes in health research. Using this taxonomy, outcomes are mapped to 5 main core areas: survival, clinical/physiological changes, life impact, resource use and adverse events. Within each core area disease and treatment specific outcome measure domains are created adding more granularity and scope. This taxonomy has been created to provide a high-level difference between outcome domains in order to improve uniformity of outcome classification in electronic databases and it has been adopted in the development of many core outcome sets (COS) projects, systematic reviews and clinical trials. The taxonomy has the added advantage of flexibility in mapping outcomes to specific outcome domains within each core area. For instance, within the clinical and physiological core area, many patients and clinician reported outcome measures such as pain, healing, success/failure of treatment can be mapped, while OHRQoL measures are covered in the functionality core area and health economics outcomes in use of resources core area. The main problem, however, with using this or other outcome classifications is the lack of consensus on the definition of endodontic treatment outcome itself. For example, the definition of success has been

modified over the years with earlier classifications using terms such as stringent or lenient where in the former, success is characterized by clinical and radiographic normalcy while the latter requires only an absence of clinical signs and symptoms.⁹⁷ Others have adopted strict and loose criteria for success, where strict is defined as absence of clinical signs and symptoms and radiographic evidence of complete healing, whereas success based on loose criteria is defined by absence of signs and symptoms and radiographic evidence of incomplete healing.^{98,99}

CORE OUTCOME SET (COS) IN ENDODONTICS

The lack of consensus in reporting and defining endodontic treatment outcomes has stimulated recent initiatives for development of COS for endodontic treatment. Using a scoping review methodology, Azarpazhooh et al.¹⁰⁰ developed a framework for standardized data collection and reporting of endodontic outcome studies. Although this provided valuable information and identified problems with outcome reporting, the final outcome was not a consensus-based COS development. COS is defined as an agreed, standardized set of outcomes that should be included, measured and reported as a minimum in all trials and outcome studies in a particular field.⁹³ The process involves identification of reported outcomes through literature searches, followed by a consensus Delphi process to agree on the most important outcomes for inclusion. The contribution of appropriate stakeholders is essential.⁹³ Therefore, the creation of a Core Outcome Set for Endodontic Treatment modalities (COSET) using a standardized and approved methodology in line with the Core Outcome Set-STAndards for Development (COS-STAD) and Core Outcome Set-STAndards for Reporting the (COS-STAR) recommendations^{101,102} is currently underway.

An *a priori* protocol for the development of the COSET project was published³² and registered in COMET.¹⁰³ This is an international collaborative project and will be carried out into two phases: (1) a structured scoping review to identify all outcomes reported for the range of endodontic treatment modalities including non-surgical root canal treatment, surgical root canal treatment, vital pulp treatment and revitalization. (2) a consensus process using the Delphi methodology and semi-structured interviews to agree upon the most important outcomes. This project is different to other COS studies in dentistry in that it puts the patients at the center of the process and considers their

perspective in the COS to be developed. The expected outcome will be the development of a consensus-based list of outcomes for each endodontic treatment modality, how these outcomes are measured and the optimal timing for their measurement.

FUTURE DIRECTIONS

- Several reports have confirmed a lack of clinical studies employing dPROs in endodontics.^{23,100} Future clinical studies should incorporate dPROs such as discomfort, and OHRQoL along with important evidence based CROs. dPROMs used in clinical trials must be reliable and valid. This would lead to a patient-centered, evidence-based practice of dentistry, thereby reducing research waste, and ultimately increasing the value of treatment and research in endodontology.
- Research has demonstrated that disease-specific assessment is important, for example, sarcoma and autosomal hereditary bleeding disorders in patients.^{104,105} Most of the PROMs employed in endodontics have been generic. Hence, a need arises to develop a validated dPROM that is specific to endodontology to standardize the outcome evaluation to guide further research and enhance patient care.
- Future studies should be conducted to identify the dPROMs that can be used in endodontics and map those outcome measures to the 4-dimensional framework (Oral function, Orofacial appearance, Orofacial pain, and Psychosocial impact). This will eventually result in the selection of the most appropriate outcome measures for patients undergoing endodontic treatment in the future.
- It is critical that future research adheres to a standardized approach for recording and reporting OHRQoL.²⁰ It has been noted in a previous systematic review analyzing OHRQoL in endodontology, that meta-analyses were not possible due to the lack of a standardized method for recording and reporting OHRQoL scores.²⁰ The standard approach should include OHRQoL scores obtained prior to the intervention and a sufficient recall period (e.g. 6 months). It is recommended to assess this dPRO using previously validated instruments and, if possible, an endodontic disease-specific tool.

CONCLUSION

With the development of dPROs and an awareness of what is important to patients, both practitioners and researchers now have the ability to improve endodontic treatment and research, ultimately benefiting the profession, individual patients and society. Appropriate use of dPROs is critical in endodontology to appropriately reflect the impact of pulpal or periapical diseases or endodontic treatment on patients. However, currently endodontic-specific PROMs are lacking. A new and specific measuring instrument should be developed to better reflect the perspectives of patients receiving endodontic treatment.

References

1. U.S. Department of Health and Human Services FDA Center for Drug Evaluation and Research, U.S. Department of Health and Human Services FDA Center for Biologics Evaluation and Research, U.S. Department of Health and Human Services FDA Center for Devices and Radiological Health. Guidance for industry: patient-reported outcome measures: use in medical product development to support labeling claims: draft guidance. *Health Qual Life Outcomes*. 2006;4:79.
2. Powers JH, Patrick DL, Walton MK, et al. Clinician-Reported Outcome Assessments of Treatment Benefit: Report of the ISPOR Clinical Outcome Assessment Emerging Good Practices Task Force. *Value Health*. 2017;20:2-14.
3. Johnston B, DL P, T D, et al. Patient-reported outcomes. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ WV, ed. *Cochrane Handbook for Systematic Reviews of Interventions Version 6.3 (Updated February 2022)*. Cochrane; 2022. www.training.cochrane.org/handbook.
4. Pantaleon L. Why measuring outcomes is important in health care. *J Vet Intern Med*. 2019;33:356-362.
5. Mercieca-Bebber R, Williams D, Tait MA, et al. Trials with patient-reported outcomes registered on the Australian New Zealand Clinical Trials Registry (ANZCTR). *Qual Life Res*. 2018;27:2581-2591.
6. Mercieca-Bebber R, King MT, Calvert MJ, Stockler MR, Friedlander M. The importance of patient-reported outcomes in clinical trials and strategies for future optimization. *Patient Relat Outcome Meas*. 2018;9:353-367.
7. John MT. Health Outcomes Reported by Dental Patients. *J Evid Based Dent Pract*. 2018;18:332-335.
8. Liu YC, Shih MC, Tu YK. Using dental patient-reported outcomes (dPROS) in meta-analyses: a scoping review and methodological investigation. *J Evid Based Dent Pract*. 2022;22:101658.
9. Liang M, Lian Q, Kotsakis GA, Michalowicz BS, John MT, Chu H. Bayesian Network Meta-analysis of Multiple Outcomes in Dental Research. *J Evid Based Dent Pract*. 2020;20:101403.
10. Paulson DR, Pattanaik S, Chanthavisouk P, John MT. Including the patient's oral health perspective in evidence-based decision-making. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2021;64:959-966.

11. Tao Z, Zhao T, Ngan P, Qin D, Hua F, He H. The use of Dental Patient-Reported Outcomes among randomized controlled trials in orthodontics: A methodological study. *J Evid Based Dent Pract.* 2022;In press.
12. Reissmann DR. Dental Patient-Reported Outcome Measures Are Essential for Evidence-Based Prosthetic Dentistry. *J Evid Based Dent Pract.* 2019;19:1-6.
13. Hua F. Increasing the Value of Orthodontic Research Through the Use of Dental Patient-Reported Outcomes. *J Evid Based Dent Pract.* 2019;19:99-105.
14. Listl S. Value-Based Oral Health Care: Moving Forward With Dental Patient-Reported Outcomes. *J Evid Based Dent Pract.* 2019;19:255-259.
15. U.S. Department of Health and Human Services. *Oral Health in America: A Report of the Surgeon General.*; 2000.
16. European Society of Endodontology. Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. *Int Endod J.* 2006;39:921-930.
17. Torabinejad M, Bahjri K. Essential elements of evidenced-based endodontics: steps involved in conducting clinical research. *J Endod.* 2005;31:563-569.
18. Liu P, McGrath C, Cheung GSP. Improvement in oral health-related quality of life after endodontic treatment: a prospective longitudinal study. *J Endod.* 2014;40:805-810.
19. Wigsten E, Kvist T, Jonasson P, EndoReCo, Davidson T. Comparing Quality of Life of Patients Undergoing Root Canal Treatment or Tooth Extraction. *J Endod.* 2020;46:19-28.e1.
20. Neelakantan P, Liu P, Dummer PMH, McGrath C. Oral health-related quality of life (OHRQoL) before and after endodontic treatment: a systematic review. *Clin Oral Investig.* 2020;24:25-36.
21. Duncan HF, Nagendrababu V, El-Karim IA, Dummer PMH. Outcome measures to assess the effectiveness of endodontic treatment for pulpitis and apical periodontitis for use in the development of European Society of Endodontology (ESE) S3 level clinical practice guidelines: a protocol. *Int Endod J.* 2021;54:646-654.
22. Duncan HF, Nagendrababu V, El-Karim I, Dummer PMH. Outcome measures to assess the effectiveness of endodontic treatment for pulpitis and apical periodontitis for use in the development of European Society of Endodontology

- S3-level clinical practice guidelines: A consensus-based development. *Int Endod J*. 2021;54:2184-2194.
23. De-Deus G, Canabarro A. Strength of recommendation for single-visit root canal treatment: grading the body of the evidence using a patient-centred approach. *Int Endod J*. 2017;50:251-259.
 24. Su Y, Wang C, Ye L. Healing rate and post-obturation pain of single- versus multiple-visit endodontic treatment for infected root canals: a systematic review. *J Endod*. 2011;37:125-132.
 25. Azarpazhooh A, Sgro A, Cardoso E, et al. A Scoping Review of 4 Decades of Outcomes in Nonsurgical Root Canal Treatment, Nonsurgical Retreatment, and Apexification Studies-Part 2: Outcome Measures. *J Endod*. 2022;48:29-39.
 26. Montero J, Lorenzo B, Barrios R, Albaladejo A, Mirón Canelo JA, López-Valverde A. Patient-centered Outcomes of Root Canal Treatment: A Cohort Follow-up Study. *J Endod*. 2015;41:1456-1461.
 27. Bergenholtz G, Kvist T. Evidence-based endodontics. *Endod Top*. 2014;31:3-18.
 28. Friedman S, Mor C. The success of endodontic therapy--healing and functionality. *J Calif Dent Assoc*. 2004;32:493-503.
 29. Wigsten E, Al Hajj A, Jonasson P, et al. Patient satisfaction with root canal treatment and outcomes in the Swedish public dental health service: A prospective cohort study. *Int Endod J*. 2021;54:1462-1472.
 30. Polycarpou N, Ng YL, Canavan D, Moles DR, Gulabivala K. Prevalence of persistent pain after endodontic treatment and factors affecting its occurrence in cases with complete radiographic healing. *Int Endod J*. 2005;38:169-178.
 31. Bergenholtz G, Spångberg L. Controversies in endodontics. *Crit Rev Oral Biol Med*. 2004;15:99-114.
 32. El-Karim IA, Duncan HF, Cushley S, et al. A protocol for the Development of Core Outcome Sets for Endodontic Treatment modalities (COSET): an international consensus process. *Trials*. 2021;22.
 33. Gatten DL, Riedy CA, Hong SK, Johnson JD, Cohenca N. Quality of life of endodontically treated versus implant treated patients: a University-based qualitative research study. *J Endod*. 2011;37:903-909.
 34. Coleman RL, Beck JT, Baranda JC, et al. The Use of Patient-Reported Outcome Measures in Phase I Oncology Clinical Trials. *Oncology*. 2021;99:444-453.

35. Gerritsen AE, Allen PF, Witter DJ, Bronkhorst EM, Creugers NHJ. Tooth loss and oral health-related quality of life: a systematic review and meta-analysis. *Health Qual Life Outcomes*. 2010;8:126.
36. Tsesis I, Shoshani Y, Givol N, Yahalom R, Fuss Z, Taicher S. Comparison of quality of life after surgical endodontic treatment using two techniques: a prospective study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2005;99:367-371.
37. Careddu R, Duncan HF. How does the pulpal response to Biodentine and ProRoot mineral trioxide aggregate compare in the laboratory and clinic? *Br Dent J*. Published online October 19, 2018.
38. Montero J, Albaladejo A, Zalba J. Influence of the usual motivation for dental attendance on dental status and oral health-related quality of life. *Med Oral Patol Oral y Cir Bucal*. Published online 2014:e225-e231.
39. Slade GD, Spencer AJ. Development and evaluation of the Oral Health Impact Profile. *Community Dent Health*. 1994;11:3-11.
40. John MT. Standardization of dental patient-reported outcomes measurement using ohip-5 – Validation of “Recommendations for use and scoring of oral health impact profile versions.” *J Evid Based Dent Pract*. 2022;22:101645.
41. Atav Ates A, Dumani A, Yoldas O, Unal I. Post-obturation pain following the use of carrier-based system with AH Plus or iRoot SP sealers: a randomized controlled clinical trial. *Clin Oral Investig*. 2019;23:3053-3061.
42. Tan HSG, Lim KC, Lui JN, Lai WMC, Yu VSH. Postobturation Pain Associated with Tricalcium Silicate and Resin-based Sealer Techniques: A Randomized Clinical Trial. *J Endod*. 2021;47:169-177.
43. Ferreira N de S, Gollo EKF, Boscato N, Arias A, Silva EJNI da. Postoperative pain after root canal filling with different endodontic sealers: a randomized clinical trial. *Braz Oral Res*. 2020;34.
44. Aslan T, Dönmez Özkan H. The effect of two calcium silicate-based and one epoxy resin-based root canal sealer on postoperative pain: a randomized controlled trial. *Int Endod J*. 2021;54:190-197.
45. Ballal NV, Duncan HF, Rai N, Jalan P, Zehnder M. Sodium Hypochlorite Reduces Postoperative Discomfort and Painful Early Failure after Carious Exposure and Direct Pulp Capping-Initial Findings of a Randomized Controlled Trial. *J Clin Med*. 2020;9.

46. Mostafa M, El-Shrief Y, Anous W, et al. Postoperative pain following endodontic irrigation using 1.3% versus 5.25% sodium hypochlorite in mandibular molars with necrotic pulps: a randomized double-blind clinical trial. *Int Endod J*. 2020;53:154-166.
47. Friedman S, Abitbol S, Lawrence HP. Treatment outcome in endodontics: the Toronto Study. Phase 1: initial treatment. *J Endod*. 2003;29:787-793.
48. Farzaneh M, Abitbol S, Lawrence HP, Friedman S, Toronto Study. Treatment outcome in endodontics-the Toronto Study. Phase II: initial treatment. *J Endod*. 2004;30:302-309.
49. Marquis VL, Dao T, Farzaneh M, Abitbol S, Friedman S. Treatment outcome in endodontics: the Toronto Study. Phase III: initial treatment. *J Endod*. 2006;32:299-306.
50. de Chevigny C, Dao TT, Basrani BR, et al. Treatment outcome in endodontics: the Toronto study--phase 4: initial treatment. *J Endod*. 2008;34:258-263.
51. Bernstein SD, Horowitz AJ, Man M, et al. Outcomes of endodontic therapy in general practice: a study by the Practitioners Engaged in Applied Research and Learning Network. *J Am Dent Assoc*. 2012;143:478-487.
52. Azarpazhooh A, Dao T, Ungar WJ, et al. Patients' Values Related to Treatment Options for Teeth with Apical Periodontitis. *J Endod*. 2016;42:365-370.
53. Andrade EB, Bianchini MA, Lucchiari N. Combine or Separate Future Pain? The Impact of Current Pain on Decisions about Future Dental Treatments. Milanese S, ed. *PLoS One*. 2013;8:e64057.
54. Bassam S, El-Ahmar R, Salloum S, Ayoub S. Endodontic postoperative flare-up: An update. *Saudi Dent J*. 2021;33:386-394.
55. Onay EO, Ungor M, Yazici AC. The evaluation of endodontic flare-ups and their relationship to various risk factors. *BMC Oral Health*. 2015;15:142.
56. Azim AA, Azim KA, Abbott P V. Prevalence of inter-appointment endodontic flare-ups and host-related factors. *Clin Oral Investig*. 2017;21:889-894.
57. Meschi N, EzEldeen M, Garcia AET, et al. Regenerative Endodontic Procedure of Immature Permanent Teeth with Leukocyte and Platelet-rich Fibrin: A Multicenter Controlled Clinical Trial. *J Endod*. 2021;47:1729-1750.
58. Santini M, Da Rosa RA, Ferreira MB, et al. Medications Used for Prevention and Treatment of Postoperative Endodontic Pain: A Systematic Review. *Eur Endod J*.

- 2021;6:15-24.
59. John MT. Foundations of oral health-related quality of life. *J Oral Rehabil*. 2021;48:355-359.
 60. Dođramacı E, Rossi-Fedele G. Patient-related outcomes and Oral Health-Related Quality of Life in endodontics. *Int Endod J*.:Unpublished results.
 61. Lawal F, Omara M. Applicability of dPROs in low resource settings - bridging the gap in oral diagnostics, monitoring and evaluation in the clinical and community settings. *J Evid Based Dent Pract*. 2022;In press.
 62. Hamasha AA, Hatiwsh A. Quality of life and satisfaction of patients after nonsurgical primary root canal treatment provided by undergraduate students, graduate students and endodontic specialists. *Int Endod J*. 2013;46:1131-1139.
 63. He J, White RK, White CA, Schweitzer JL, Woodmansey KF. Clinical and Patient-centered Outcomes of Nonsurgical Root Canal Retreatment in First Molars Using Contemporary Techniques. *J Endod*. 2017;43:231-237.
 64. Chew T, Brennan D, Rossi-Fedele G. Comparative Longitudinal Study on the Impact Root Canal Treatment and Other Dental Services Have on Oral Health-related Quality of Life Using Self-reported Health Measures (Oral Health Impact Profile-14 and Global Health Measures). *J Endod*. 2019;45:985-993.e1.
 65. Dugas NN, Lawrence HP, Teplitsky P, Friedman S. Quality of life and satisfaction outcomes of endodontic treatment. *J Endod*. 2002;28:819-827.
 66. Iqbal MK, Kratchman SI, Guess GM, Karabucak B, Kim S. Microscopic periradicular surgery: perioperative predictors for postoperative clinical outcomes and quality of life assessment. *J Endod*. 2007;33:239-244.
 67. Khoo ST, Ode W, Lopez V, Yu VSH, Lai C, Lui JN. Factors Influencing Quality of Life after Surgical and Nonsurgical Interventions of Persistent Endodontic Disease. *J Endod*. 2020;46:1832-1840.
 68. Tibúrcio-Machado CS, Michelon C, Zanatta FB, Gomes MS, Marin JA, Bier CA. The global prevalence of apical periodontitis: a systematic review and meta-analysis. *Int Endod J*. 2021;54:712-735.
 69. Ng YL. *Factors Affecting Outcome of Non-Surgical Root Canal Treatment*. University of London, University College London (United Kingdom); 2008.
 70. Zilinskaite-Petrauskiene I, Haug SR. A Comparison of Endodontic Treatment Factors, Operator Difficulties, and Perceived Oral Health-related Quality of Life

- between Elderly and Young Patients. *J Endod.* 2021;47:1844-1853.
71. Pasqualini D, Corbella S, Alovisei M, et al. Postoperative quality of life following single-visit root canal treatment performed by rotary or reciprocating instrumentation: a randomized clinical trial. *Int Endod J.* 2016;49:1030-1039.
 72. Oliveira PS, da Costa KNB, Carvalho CN, Ferreira MC. Impact of root canal preparation performed by ProTaper Next or Reciproc on the quality of life of patients: a randomized clinical trial. *Int Endod J.* 2019;52:139-148.
 73. Diniz-de-Figueiredo FE, Lima LF, Oliveira LS, Bernardino IM, Paiva SM, Faria-E-Silva AL. The impact of two root canal treatment protocols on the oral health-related quality of life: a randomized controlled pragmatic clinical trial. *Int Endod J.* 2020;53:1327-1338.
 74. Multari S, Alovisei M, Berutti E, et al. Postoperative Quality of Life after Single-Visit Root Canal Treatment Performed with Reciprocating Shaping Systems: An Observational Study. *Appl Sci.* 2021;11:273.
 75. Vena DA, Collie D, Wu H, et al. Prevalence of persistent pain 3 to 5 years post primary root canal therapy and its impact on oral health-related quality of life: PEARL Network findings. *J Endod.* 2014;40:1917-1921.
 76. U.S. Department of Health and Human, Administration SF and D, (CDER) C for DE and R, (CBER) C for BE and R, Center for Devices and Radiological Health (CDRH). *Guidance for Industry Patient-Reported Outcome Measures: Use in Medical Product Development to Support Labeling Claims.*; 2009.
 77. Maddaloni M, Gagliani M, Citterio CL, Karanxha L, Pellegatta A, Del Fabbro M. Prevalence of vertical root fractures in teeth planned for apical surgery. A retrospective cohort study. *Int Endod J.* 2018;51:969-974.
 78. Bosenbecker J, Barbon FJ, de Souza Ferreira N, Morgental RD, Boscato N. Tooth discoloration caused by endodontic treatment: A cross-sectional study. *J Esthet Restor Dent.* 2020;32:569-574.
 79. Ekici MA, Ekici A, Kaskatı T, Helvacioğlu Kıvanç B. Tooth crown discoloration induced by endodontic sealers: a 3-year ex vivo evaluation. *Clin Oral Investig.* 2019;23:2097-2102.
 80. Gürel MA, Kıvanç BH, Ekici A, Alaçam T. Evaluation of crown discoloration induced by endodontic sealers and colour change ratio determination after bleaching. *Aust Endod J.* 2016;42:119-123.

81. Ioannidis K, Beltes P, Lambrianidis T, Kapagiannidis D, Karagiannis V. Crown discoloration induced by endodontic sealers: spectrophotometric measurement of Commission International de l'Eclairage's L*, a*, b* chromatic parameters. *Oper Dent*. 2013;38:E1-12.
82. Karabucak B, Stoopler ET. Root canal treatment on a patient with zinc oxide allergy: a case report. *Int Endod J*. 2007;40:800-807.
83. Braun JJ, Zana H, Purohit A, et al. Anaphylactic reactions to formaldehyde in root canal sealant after endodontic treatment: four cases of anaphylactic shock and three of generalized urticaria. *Allergy*. 2003;58:1210-1215.
84. Haïkel Y, Braun JJ, Zana H, Boukari A, de Blay F, Pauli G. Anaphylactic shock during endodontic treatment due to allergy to formaldehyde in a root canal sealant. *J Endod*. 2000;26:529-531.
85. Kaufman AY, Solomonov M, Galieva D, Abbott P V. Allergic reaction to the tetracycline component of Ledermix paste: a case report. *Int Endod J*. 2014;47:1090-1097.
86. Çalışkan MK, Türkün M, Alper S. Allergy to sodium hypochlorite during root canal therapy: a case report. *Int Endod J*. 1994;27:163-167.
87. Pemberton MN, Gibson J. Chlorhexidine and hypersensitivity reactions in dentistry. *Br Dent J*. 2012;213:547-550.
88. Al-Sheeb F, Al Mannai G, Tharupeedikayil S. Nicolau Syndrome after Endodontic Treatment: A Case Report. *J Endod*. 2022;48:269-272.
89. John MT, Hujoel P, Miglioretti DL, LeResche L, Koepsell TD, Micheelis W. Dimensions of oral-health-related quality of life. *J Dent Res*. 2004;83:956-960.
90. Azarpazhooh A, Cardoso E, Sgro A, et al. A Scoping Review of 4 Decades of Outcomes in Nonsurgical Root Canal Treatment, Nonsurgical Retreatment, and Apexification Studies-Part 1: Process and General Results. *J Endod*. 2022;48:15-28.
91. Cushley S, Duncan HF, Lappin MJ, et al. Pulpotomy for mature carious teeth with symptoms of irreversible pulpitis: A systematic review. *J Dent*. 2019;88:103158.
92. Cushley S, Duncan HF, Lappin MJ, et al. Efficacy of direct pulp capping for management of cariously exposed pulps in permanent teeth: a systematic review and meta-analysis. *Int Endod J*. 2021;54:556-571.
93. Williamson PR, Altman DG, Blazeby JM, et al. Developing core outcome sets for

- clinical trials: Issues to consider. *Trials*. 2012;13.
94. Fletcher RH, Fletcher SW, Fletcher GS. *Clinical Epidemiology : The Essentials*. 5th ed. Wolters Kluwer Health/Lippincott Williams & Wilkins; 2014.
 95. National Institutes of Health. PROMIS: Clinical Outcomes Assessment. The Common Fund: Patient-Reported Outcomes Measurement Information System. Accessed March 5, 2022. <http://commonfund.nih.gov/promis/index>
 96. Dodd S, Clarke M, Becker L, Mavergames C, Fish R, Williamson PR. A taxonomy has been developed for outcomes in medical research to help improve knowledge discovery. *J Clin Epidemiol*. 2018;96:84-92.
 97. Friedman S. Prognosis of initial endodontic therapy. *Endod Top*. 2002;2:59-88.
 98. Ng YL, Mann V, Gulabivala K. A prospective study of the factors affecting outcomes of non-surgical root canal treatment: part 2: tooth survival. *Int Endod J*. 2011;44:610-625.
 99. Ng YL, Mann V, Rahbaran S, Lewsey J, Gulabivala K. Outcome of primary root canal treatment: systematic review of the literature - part 1. Effects of study characteristics on probability of success. *Int Endod J*. 2007;40:921-939.
 100. Azarpazhooh A, Khazaei S, Jafarzadeh H, et al. A Scoping Review of Four Decades of Outcomes in Nonsurgical Root Canal Treatment, Nonsurgical Retreatment, and Apexification Studies: Part 3-A Proposed Framework for Standardized Data Collection and Reporting of Endodontic Outcome Studies. *J Endod*. 2022;48:40-54.
 101. Kirkham JJ, Gorst S, Altman DG, et al. Core Outcome Set-STAndards for Reporting: The COS-STAR Statement. *PLoS Med*. 2016;13.
 102. Kirkham JJ, Davis K, Altman DG, et al. Core Outcome Set-STAndards for Development: The COS-STAD recommendations. *PLoS Med*. 2017;14.
 103. El-Karim I, Duncan HF, Cushley S, et al. A protocol for the development of Core Outcome Sets for Endodontic Treatment (COSET): an international consensus process. Accessed April 13, 2022. <https://comet-initiative.org/Studies/Details/1879>
 104. Van-Hoorn ES, Houwing ME, Al Arashi W, et al. Patient-reported outcomes in autosomal inherited bleeding disorders: A systematic literature review. *Haemophilia*. 2022;28:197-214.
 105. Almeida A, Martins T, Lima L. Patient-Reported Outcomes in Sarcoma: A scoping review. *Eur J Oncol Nurs*. 2021;50:101897.