


Figure 1. Recruitment flowchart



Footnote: *due to administrative error

Figure 2: Percentage of participants (and 95% CI) who meet criteria for PTSD, sub-threshold PTSD, total PTSD, and detected by their clinical teams (ie. a diagnosis was recorded in their notes)





Figure 3: Percentage of participants (and 95% CI) who score 1 or more symptoms as moderate or more severe, by symptom domain



Post-traumatic stress disorder (PTSD) in people with psychosis and people at increased risk of developing psychosis

Supplementary document

CTQ questions used in this study

1. "I knew that there was someone to take care of me and protect me" (physical neglect)
2. "There was someone in my family who helped me feel that I was important or special" (emotional neglect)
3. "I felt loved" (emotional neglect)
4. "People in family hit me so hard that it left me with bruises or marks" (physical abuse)
5. "People in my family said hurtful or insulting things to me" (emotional abuse)
6. "I got hit or beaten so badly that it was noticed by someone like a teacher, neighbour, or doctor" (physical abuse)
7. "I felt that someone in my family hated me" (emotional abuse)
8. "Someone tried to touch me in sexual way, or tried to make me touch them" (sexual abuse)
9. "There was someone to take me to the doctor if I needed it" (physical neglect)
10. "I believe that I was sexually abused" (sexual abuse)

Post-traumatic stress disorder (PTSD) in people with psychosis and people at increased risk of developing psychosis

Table 3. Number (%) of people with a trauma history who were interested in receiving TF-therapy, by trauma-type

	N	Percentage	95% CI
Emotional abuse	33	75%	0.60, 0.87
Physical abuse	18	75%	0.53, 0.90
Sexual abuse	17	81%	0.58, 0.95
Emotional neglect	25	74%	0.56, 0.87
Physical neglect	23	72%	0.53, 0.86