


Data supplement

1. eFigure 1. Flow diagram of participants included in the analytical sample
2. Table S1. Self-Reported Suicide Attempt, Self-Harm, and Psychological Distress by Gender Minority Status in a Sample with Complete Data (n = 7,829)



eFigure 1. Flow diagram of participants included in the analytical sample

Table S1. Self-Reported Suicide Attempt, Self-Harm, and Psychological Distress by Gender Minority Status in a Sample with Complete Data (n = 7 829)

Outcome	Risk ratio (95% confidence interval)			
	Model 1	Model 2	Model 3	Model 4
Suicide attempt	3.28 (2.05, 5.23)	1.68 (1.00, 2.81)	1.80 (1.07, 3.02)	1.15 (0.74, 1.79)
Self-harm				
Cut	5.29 (4.18, 6.69)	2.40 (1.83, 3.14)	2.50 (1.93, 3.23)	1.69 (1.30, 2.19)
Burned	5.06 (3.09, 8.29)	2.19 (1.23, 3.89)	2.39 (1.36, 4.20)	1.48 (0.86, 2.54)
Bruised or pinched	4.07 (3.32, 4.98)	1.97 (1.58, 2.45)	1.98 (1.59, 2.45)	1.50 (1.22, 1.85)
Overdose	5.57 (2.94, 10.54)	2.82 (1.37, 5.82)	3.14 (1.57, 6.28)	1.85 (0.95, 3.62)
Pull hair	3.99 (2.70, 5.91)	1.82 (1.18, 2.80)	1.89 (1.22, 2.90)	1.24 (0.82, 1.88)
Other way	6.86 (4.77, 9.89)	2.77 (1.81, 4.25)	2.72 (1.76, 4.22)	1.97 (1.30, 2.99)
Psychological distress	Coefficient (95% confidence interval)			
Kessler K6 screening scale total	5.90 (4.89, 6.91)	3.42 (2.40, 4.44)	3.49 (2.48, 4.51)	2.52 (1.57, 3.47)
Strengths and Difficulties total (child)	5.31 (4.15, 6.47)	2.51 (1.31, 3.70)	2.61 (1.42, 3.80)	1.50 (0.38, 2.62)
Strengths and Difficulties total (parent)	2.22 (1.03, 3.41)	0.42 (-0.84, 1.67)	0.58 (-0.66, 1.83)	0.05 (-1.18, 1.29)

Model 1: gender identity; Model 2: model 1 plus sexual identity; Model 3: model 2 plus substance use; Model 4: model 2 plus victimisation