

## SHORT REPORT

## Haematological Malignancy – Clinicals

# A randomised evaluation of low-dose cytosine arabinoside plus lenalidomide versus single-agent low-dose cytosine arabinoside in older patients with acute myeloid leukaemia: Results from the LI-1 trial

Mhairi Copland<sup>1</sup>   | Cono Ariti<sup>2,3</sup> | Ian F. Thomas<sup>2</sup> | Laura Upton<sup>2</sup> |  
 Mia Sydenham<sup>2</sup> | Priyanka Mehta<sup>4</sup>  | Shahid Islam<sup>5</sup> | Lars Kjeldsen<sup>6</sup> |  
 Alan K. Burnett<sup>1</sup> | Robert K. Hills<sup>7</sup>  | Nigel Russell<sup>8</sup>  | Mike Dennis<sup>9</sup>   |

on behalf of the UK NCRI AML Study Group

<sup>1</sup>Paul O’Gorman Leukaemia Research Centre, School of Cancer Sciences, University of Glasgow, Glasgow, UK

<sup>2</sup>Centre for Trials Research, Cardiff University, Cardiff, UK

<sup>3</sup>Oxon Epidemiology, Madrid, Spain

<sup>4</sup>University Hospitals of Bristol and Weston NHS Trust, Bristol, UK

<sup>5</sup>Department of Haematology, Waikato Hospital, Hamilton, New Zealand

<sup>6</sup>Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark

<sup>7</sup>Nuffield Department of Population Health, University of Oxford, Oxford, UK

<sup>8</sup>Guy’s and St Thomas’ NHS Trust, London, UK

<sup>9</sup>The Christie, Manchester, UK

## Correspondence

Mhairi Copland, Paul O’Gorman Leukaemia Research Centre, School of Cancer Sciences, University of Glasgow, Glasgow G12 0ZD, UK.  
 Email: [mhairi.copland@glasgow.ac.uk](mailto:mhairi.copland@glasgow.ac.uk)

## Funding information

Blood Cancer UK; Celgene

## Summary

Improving outcomes for older patients with acute myeloid leukaemia remains an unmet need. As part of the LI-1 trial, we evaluated lenalidomide (LEN) in combination with low-dose cytosine arabinoside (LDAC) in patients aged >60 years unfit for intensive therapy and compared this to LDAC alone. Two hundred and two patients, randomised 1:1, were evaluable. Overall response rate (CR+CRi) was higher for LDAC+LEN versus LDAC (26% and 13.7% respectively  $p=0.031$ ). However, there was no difference in overall survival between the arms (14% and 11.5% at 2 years for LDAC+LEN and LDAC respectively). The addition of LEN was associated with increased toxicity and supportive care requirements.

## KEY WORDS

acute myeloid leukaemia, clinical trial, elderly, lenalidomide, low-dose cytosine arabinoside

## INTRODUCTION

Many patients with acute myeloid leukaemia (AML) diagnosed after 60 years of age are not considered suitable for intensive remission-induction chemotherapy, either due to comorbidities or frailty associated with advanced age.<sup>1</sup> Despite treatment with either low-dose cytosine arabinoside

(LDAC) or a hypomethylating agent,<sup>2,3</sup> survival is usually poor, with 1-year overall survival (OS) after LDAC of 21%–32% in NCRI AML16 and historical arms of LI-1.<sup>3</sup> Combination therapy with a backbone of LDAC or a hypomethylating agent with additional agents represents an attractive option, with the potential to improve patient outcomes without substantially increasing toxicity.<sup>4–7</sup>

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *British Journal of Haematology* published by British Society for Haematology and John Wiley & Sons Ltd.

Recently, the BCL-2 inhibitor venetoclax has been approved and widely adopted for the treatment of older, less fit patients with AML in combination with either azacitidine or LDAC.<sup>4,5</sup> This has demonstrated superior complete remission (CR) rates compared to azacitidine or LDAC alone: 36.7% for azacitidine+venetoclax compared to 17.9% for azacitidine alone in VIALE-A,<sup>4</sup> and 48% for LDAC+venetoclax compared to 13% for LDAC+placebo in VIALE-C.<sup>5</sup> Improvements in survival are also seen, with a median OS of 14.7 months for azacitidine+venetoclax, 9.6 months for azacitidine alone in VIALE-A, 8.4 months for LDAC+venetoclax and 4.1 months for LDAC+placebo in VIALE-C.<sup>4,5</sup> These survival benefits are modest with most patients still dying of AML, and there remains a need to develop new well-tolerated, outpatient-based, effective therapeutic strategies for older, frail patients with AML to further improve outcomes.

Lenalidomide (LEN; Revlimid™), a derivative of thalidomide, is an immunomodulatory drug used to treat myeloma<sup>8</sup> and some cases of myelodysplastic syndrome (MDS),<sup>9</sup> and has potent anti-neoplastic, anti-angiogenic, anti-inflammatory and pro-erythropoietic properties.<sup>10</sup> Early-phase trials of LEN in AML have demonstrated clinical activity with acceptable toxicity.<sup>11,12</sup>

We assessed the efficacy and tolerability of LDAC+LEN versus LDAC alone in patients aged 60+ unsuitable for intensive therapy.

## METHODS

### Design and eligibility

The LI-1 trial (ISRCTN40571019) was an international multicentre, multiarm, randomised phase II/III trial developed to study the efficacy and tolerability of novel non-intensive therapies in AML using a 'pick-a-winner' design.<sup>6,13</sup> In LI-1, the comparator arm was LDAC, and there was no comparison of different experimental arms. Patients in the LDAC control arm were recruited and randomised 1:1, with the experimental arms available contemporaneously.

Patients aged  $\geq 60$  years, with de novo or secondary AML or high-risk MDS ( $>10\%$  marrow blasts), considered unfit for intensive therapy were eligible. Patients with a prior diagnosis of MDS with  $<10\%$  blasts who had failed a demethylating agent but subsequently developed AML were also eligible. Impaired renal or hepatic function (defined as serum creatinine  $>174 \mu\text{mol/L}$ , total bilirubin  $\geq 1.5$  times the upper limit of normal (ULN), aspartate aminotransferase or alanine aminotransferase  $\geq 2.5$  times ULN) were exclusion criteria. Patients with a history of myocardial infarction, unstable angina, cerebrovascular accident/transient ischaemic attack within 6 months were also excluded.

LDAC was given at 20 mg BD SC on days 1–10 of each course. LEN was administered orally once daily in a flat

10 mg dose for 21 days, where Day 1 was Day 1 of LDAC, with courses occurring at 5-week intervals for courses 1–4. Patients considered to be benefitting after four courses, that is in remission or stable disease, could continue to receive treatment until disease progression, either with LDAC+LEN at 6-week intervals or LEN only at 4-week intervals if the patient had experienced significant toxicity.

All patients provided written informed consent. The LI1 trial was sponsored by Cardiff University and approved by the Research Ethics Committee for Wales in compliance with the Declaration of Helsinki.

### End-points and toxicity

The aim within the experimental arms was doubling 2-year survival from 11% to 22% (HR 0.69), with planned interim assessments after 50 and 100 patients were recruited per arm. At the first interim assessment, overall response rate (ORR), was the primary end-point, defined as either CR or CR without evidence of adequate count recovery (CRi), and was required to be at least 2.5% higher in the experimental arm. CR was defined as normocellular marrow with  $<5\%$  leukaemic blasts, evidence of normal myeloid maturation, neutrophil and platelet recovery in the absence of platelet transfusions ( $>1 \times 10^9/\text{L}$  and  $>100 \times 10^9/\text{L}$  respectively). Patients in CR but failing to achieve neutrophils  $>1 \times 10^9/\text{L}$  and platelets  $>100 \times 10^9/\text{L}$  were designated as CRi. At the second interim assessment, the primary end-point was OS, with an HR of  $<0.85$  in the experimental arm for the trial to continue.

The co-primary objectives at the final analysis were OS, defined as time from trial randomisation to death from any cause or last follow-up; ORR (CR+CRi) and reasons for failure; duration of response; relapse rates; and deaths in the first CR. Relapse-free survival (RFS) was defined as time from remission (CR/CRi) to death or relapse, censored at last follow-up; relapse risk (RR) was defined as time from remission (CR/CRi) to relapse, censored at death in CR and last follow-up; death in CR (DCR) was defined as time from remission (CR/CRi) to death, censored at relapse and last follow-up. Secondary objectives were haematological recovery times, defined as time from end of course to recovery of platelets to  $>100 \times 10^9/\text{L}$  and neutrophils to  $>1 \times 10^9/\text{L}$ , censored at next course, or time last known not to have been recovered; adverse events and toxicity defined by the National Cancer Institute Common Terminal Criteria for Adverse Events (NCI CTCAE) version 4; and resource usage, including number of units of blood and platelets per course; number of days on intravenous antibiotics or in hospital per course.

### Statistical analysis

All analyses are by intention-to-treat. Categorical end-points (e.g. CR rates) were compared using logistic regression, giving odds ratios and confidence intervals.

Continuous/scale variables were analysed by non-parametric (Wilcoxon rank sum) tests. Time-to-event outcomes were analysed using the log-rank test and Kaplan–Meier survival curves. Hazard ratios and 95% confidence intervals were calculated using the statistics from the log-rank test.<sup>14</sup>

## RESULTS

### Patient characteristics

Between January 2017 and June 2019, 206 patients from Denmark (8%), New Zealand (16%) and the United Kingdom (76%) were randomised (Consort Diagram; [Figure S1](#)). Four patients were randomised in error and removed from subsequent analyses. Thus, 202 patients were evaluable. The median age was 78 years (range 62–89); 92.6% of patients were aged  $\geq 70$  years and 35.1%  $\geq 80$  years. Baseline patient demographics are shown in [Table 1](#). Overall, 153/202 (75.7%) patients had de novo AML, 40/202 (19.8%) secondary AML and 9/202 (4.5%) high-risk MDS. Cytogenetic data were available for 173/202 patients (85.6%). One patient had favourable cytogenetics, 133 normal/intermediate and 39 adverse karyotypes. Of interest, 13/202 (6.4%) patients had a del5q abnormality, but in 12/13, this was as part of a complex karyotype or in association with a *TP53* mutation. The one patient with isolated del5q was randomised to the control arm, did not achieve CR/CRi and died from sepsis after two cycles of LDAC.

The most prevalent baseline comorbidities were cardiovascular disease, infection, arrhythmias and diabetes, all affecting  $>10\%$  of the study population. Less frequent comorbidities are shown in [Figure S2A](#). The haematopoietic cell transplantation-comorbidity index (HCT-CI) was available for 198/202 patients and was 0 in 36.1%, 1–2 in 33.1% and  $\geq 3$  in 28.7%, indicating an overall frail population ([Figure S2B](#)).

One hundred and ninety-seven patients received their allocated therapy: 100 in the LDAC arm and 97 in the LDAC+LEN arm ([Figure S1](#)). A median of two courses (range 0–24; mean 3.28) was delivered in the LDAC arm and one course in the LDAC+LEN arm (range 0–25; mean 3.48; [Figure S3](#)).

### Response

Overall response (CR/CRi) was achieved in 40/202 patients (19.8%) ([Table 2](#)). There was a significant difference in ORR between the LDAC and LDAC+LEN arms (LDAC 13.7% and LDAC+LEN 26%, respectively, OR 0.45 [0.22, 0.93],  $p=0.031$ ).

Despite the difference in ORR, 1- and 2-year OS showed no significant difference between the LDAC and LDAC+LEN arms at the second interim analysis (22.9% and 11.5% in the LDAC arm and 29.7% and 14% in the LDAC+LEN

arm, respectively; HR 0.94 [0.69, 1.2],  $p=0.719$  at 2 years). Median OS was 4.6 months for LDAC versus 3.5 months for LDAC+LEN; HR 0.96 (0.71, 1.30),  $p=0.798$  ([Figure S4A](#)). One-year OS for patients that did not enter CR/CRi was 6.8% for LDAC+LEN versus 16.9% for LDAC ( $p=0.028$ ). The most common cause of death in both arms was a resistant/recurrent disease. There was no difference in survival after remission, RFS or survival after relapse between the two arms ([Figure S4B–D](#)). Note, however, that the study was powered to detect a difference in OS, so with the low response rate observed, it was unlikely differences in RFS could be detected. Analysis by AML or patient characteristics did not identify any subgroup in which LDAC+LEN had an OS benefit ([Figure S5](#)).

### Toxicity and resource usage

Most adverse events (AEs) were grade 1/2 in both arms ([Figure S6](#)). During cycle 1, there were 78 versus 51 grade 3/4 AEs in the LDAC+LEN and LDAC arms respectively ( $p=0.02$ ). This included five thrombotic events in the LDAC+LEN arm (4 grade 3 and 1 grade 4) and none in the LDAC arm. Thirty- and 60-day mortality were not significantly different between the arms (19.2% for LDAC vs. 19.4% for LEN+LDAC at 30 days [OR 1.02 {0.52, 1.92};  $p=0.96$ ] and 31.3% vs. 41.2% in the LDAC vs. LDAC+LEN arms at 60 days [OR 1.33 {0.84, 2.12};  $p=0.23$ ]; [Table 2](#)).

In course 1, supportive care requirements were higher in terms of both days of antibiotics (7 vs. 3;  $p=0.001$ ) and hospitalisation days (11 vs. 6.5;  $p=0.005$ ) for the LDAC+LEN arm ([Figure S7](#)). There was no difference in transfusion requirements.

## CONCLUSION

Despite improving the CR/CRi rate, the combination of LDAC+LEN did not improve OS, RFS or time in remission in elderly patients with AML. The addition of LEN to LDAC resulted in increased toxicity, including episodes of thrombosis, and increased supportive care requirements. Alternative strategies to improve survival for elderly patients with AML remain a significant clinical need.

### AUTHOR CONTRIBUTIONS

Mhairi Copland, Mike Dennis and the UK NCRI AML Study Group designed and implemented the trial. Mike Dennis and Mhairi Copland were co-chief investigators; they reviewed the data and wrote the manuscript. Alan K. Burnett designed the trial, wrote the protocol and was chief investigator until Q3 2014; Robert K. Hills designed the trial and wrote the protocol. Ian F. Thomas supervised the data collection and reviewed the data. Cono Ariti analysed the data. Laura Upton and Mia Sydenham supervised the data collection and reviewed the data. Priyanka Mehta, Shahid Islam and Lars Kjeldsen were the top recruiters in their

TABLE 1 Baseline clinical characteristics.

	Randomisation		Total N = 202
	LDAC + Lenalidomide N = 100	LDAC N = 102	
<b>Age at entry (years)</b>			
Mean (SD)	77.5 (5.3)	77.2 (5.5)	77.3 (5.4)
Median (IQR)	77.7 (73.5, 81.1)	77.5 (73.4, 81.2)	77.7 (73.4, 81.2)
Min, max	(62.2, 89.1)	(63.1, 92.2)	(62.2, 92.2)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>Age group (years)</b>			
60–64	2 (2.0)	1 (1.0)	3 (1.5)
65–69	4 (4.0)	8 (7.8)	12 (5.9)
70–74	29 (29.0)	27 (26.5)	56 (27.7)
75–79	28 (28.0)	32 (31.4)	60 (29.7)
80+	37 (37.0)	34 (33.3)	71 (35.1)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>Sex</b>			
Female	46 (46.0)	39 (38.2)	85 (42.1)
Male	54 (54.0)	63 (61.8)	117 (57.9)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>WHO performance status</b>			
0	15 (15.0)	15 (14.7)	30 (14.9)
1	58 (58.0)	59 (57.8)	117 (57.9)
2	22 (22.0)	23 (22.5)	45 (22.3)
3	5 (5.0)	5 (4.9)	10 (5.0)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>AML type</b>			
De novo	75 (75.0)	78 (76.5)	153 (75.7)
Secondary	21 (21.0)	19 (18.6)	40 (19.8)
High-risk MDS	4 (4.0)	5 (4.9)	9 (4.5)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>WBC categories (10<sup>9</sup>/L)</b>			
0.0–9.9	66 (66.0)	67 (65.7)	133 (65.8)
10.0–49.9	25 (25.0)	29 (28.4)	54 (26.7)
50–99.9	7 (7.0)	5 (4.9)	12 (5.9)
100+	2 (2.0)	1 (1.0)	3 (1.5)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>Cytogenetic status</b>			
Favourable	0 (0.0)	1 (1.0)	1 (0.5)
Normal/Intermediate	63 (63.0)	70 (68.6)	133 (65.8)
Adverse	17 (17.0)	22 (21.6)	39 (19.3)
Unknown	20 (20.0)	9 (8.8)	29 (14.4)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)
<b>Wheatley index</b>			
Good	3 (3.0)	2 (2.0)	5 (2.5)
Standard	47 (47.0)	42 (41.2)	89 (44.1)
Poor	50 (50.0)	58 (56.9)	108 (53.5)
Missing (%)	0 (0.0)	0 (0.0)	0 (0.0)

Abbreviations: AML, acute myeloid leukaemia; IQR, interquartile range; LDAC, low-dose cytosine arabinoside; MDS, myelodysplastic syndrome; SD, standard deviation; WBC, white blood cell count.

**TABLE 2** Response and survival outcomes.

	Randomisation		OR/HR (95% CI)	p value
	LDAC + Lenalidomide	LDAC		
	N = 100	N = 102		
Patient status, n (%)				
Resistant disease	55 (55.0)	69 (67.6)	1.71 (0.97, 3.03)	0.066
Induction death	19 (19.0)	19 (18.6)	0.98 (0.48, 1.98)	0.946
Achieved CR/Cri	26 (26.0)	14 (13.7)	0.45 (0.22, 0.93)	0.031
Response outcomes n (%)				
CR	24 (24.0)	12 (11.8)	0.42 (0.20, 0.90)	0.026
Cri	2 (2.0)	2 (2.0)	0.98 (0.14, 7.10)	0.984
ORR (CR + Cri)	26 (26.0)	14 (13.7)	0.45 (0.22, 0.93)	0.031
Survival end-points (months)				
Median survival	3.5	4.6		
30-day mortality	19.4	19.2	1.02 (0.54, 1.92)	0.957
60-day mortality	41.2	31.3	1.33 (0.84, 2.12)	0.225
1-year survival	29.7	23.3	0.94 (0.68, 1.31)	0.732
2-year survival	14.3	11.1	0.95 (0.70, 1.29)	0.730
1-year relapse-free survival	57.7	35.7	1.26 (0.52, 3.02)	0.611
2-year relapse-free survival	30.8	14.3	1.57 (0.77, 3.22)	0.220
2-year survival after remission	45.8	35.7	0.66 (0.27, 1.61)	0.322
1-year survival after relapse	22.2	20.0	1.53 (0.65, 3.63)	0.351
1-year survival no CR/Cri	6.8	17.3	1.47 (1.04, 2.09)	0.024

Note: Response end-points are reported as n (%) and the odds ratio comparing LDAC to LDAC + LEN. Survival end-points are reported as Kaplan–Meier estimates (%) and hazard ratios comparing LDAC + LEN to LDAC.

Abbreviations: CR, complete response; Cri, complete response with incomplete recovery of counts; HR, hazard ratio; LDAC, low-dose cytosine arabinoside; OR, odds ratio; ORR, overall response rate.

countries. Nigel Russell: designed the trial; reviewed the data. All authors reviewed the manuscript.

## ACKNOWLEDGEMENTS

We are grateful to the patients for contributing to this trial. Thank you to the NIHR research teams for supporting local trial delivery and handling of patient samples.

## FUNDING INFORMATION

We are grateful to the NIHR for supporting local trial delivery, Blood Cancer UK for research support, the Haematology Clinical Trials Unit and Centre for Trials Research, Cardiff University for managing the trial and Celgene for providing drug and additional support for this Investigator-Initiated Trial.

## CONFLICT OF INTEREST STATEMENT

M.C. has received research funding from Cyclacel and Incyte, is/has been an advisory board member for Novartis, Incyte, Jazz Pharmaceuticals, Pfizer and Servier, and has received honoraria from Astellas, Novartis, Incyte, Pfizer and Jazz Pharmaceuticals. P.M. has received advisory fees and speaker honoraria from Jazz Pharmaceuticals, Pfizer, Astellas, Servier and Celgene. L.K. has received honoraria from Celgene.

## DATA AVAILABILITY STATEMENT

Data from this trial will be made available upon request to the trial sponsor (Cardiff University).

## ETHICS STATEMENT

All patients provided written informed consent. The LI1 trial was sponsored by Cardiff University and approved by the Research Ethics Committee (REC) for Wales in compliance with the Declaration of Helsinki.

## ORCID

Mhairi Copland  <https://orcid.org/0000-0002-7655-016X>

Priyanka Mehta  <https://orcid.org/0000-0001-8561-1705>

Robert K. Hills  <https://orcid.org/0000-0003-0166-0062>

Nigel Russell  <https://orcid.org/0000-0003-1893-8155>

Mike Dennis  <https://orcid.org/0000-0002-2540-8673>

## TWITTER

Mhairi Copland  CoplandMhairi

Mike Dennis  mikewdennis

## REFERENCES

1. Dennis M, Copland M, Kaur H, Kell J, Nikolousis E, Mehta P, et al. Management of older patients with frailty and acute myeloid

- leukaemia: a British Society for Haematology good practice paper. *Br J Haematol.* 2022;199(2):205–21.
2. Saiz-Rodríguez M, Labrador J, Cuevas B, Martínez-Cuadrón D, Campuzano V, Alcaraz R, et al. Use of azacitidine or decitabine for the up-front setting in acute myeloid leukaemia: a systematic review and meta-analysis. *Cancers (Basel).* 2021;13(22):5677.
  3. Dennis M, Hills RK, Russell NR, Copland M, Thomas I, McMullin M-F, et al. An evaluation of 17 years of low dose cytarabine as therapy for AML patients not fit for intensive treatment, including patients with adverse cytogenetics, shows improving survival, potential underutilisation and highlights the need for new therapy. *Blood.* 2017;130(Suppl 1):3874.
  4. DiNardo CD, Jonas BA, Pullarkat V, Thirman MJ, Garcia JS, Wei AH, et al. Azacitidine and venetoclax in previously untreated acute myeloid leukemia. *N Engl J Med.* 2020;383(7):617–29.
  5. Wei AH, Montesinos P, Ivanov V, DiNardo CD, Novak J, Laribi K, et al. Venetoclax plus LDAC for newly diagnosed AML ineligible for intensive chemotherapy: a phase 3 randomized placebo-controlled trial. *Blood.* 2020;135(24):2137–45.
  6. Dennis M, Burnett A, Hills R, Thomas I, Ariti C, Severinsen MT, et al. A randomised evaluation of low-dose cytosine arabinoside (ara-C) plus tosedostat versus low-dose ara-C in older patients with acute myeloid leukaemia: results of the LI-1 trial. *Br J Haematol.* 2021;194(2):298–308.
  7. Dennis M, Thomas IF, Ariti C, Upton L, Burnett AK, Gilkes A, et al. Randomized evaluation of quizartinib and low-dose ara-C vs low-dose ara-C in older acute myeloid leukemia patients. *Blood Adv.* 2021;5(24):5621–5.
  8. McCaughan GJ, Gandolfi S, Moore JJ, Richardson PG. Lenalidomide, bortezomib and dexamethasone induction therapy for the treatment of newly diagnosed multiple myeloma: a practical review. *Br J Haematol.* 2022;199(2):190–204.
  9. Garcia-Manero G, Almeida A, Fenaux P, Gattermann N, Giagounidis A, Goldberg SL, et al. Clinical benefit-risk profile of lenalidomide in patients with lower-risk myelodysplastic syndromes without del(5q): results of a phase III trial. *Clin Lymphoma Myeloma Leuk.* 2019;19(4):213–219.e4.
  10. Martínez-Høyer S, Karsan A. Mechanisms of lenalidomide sensitivity and resistance. *Exp Hematol.* 2020;91:22–31.
  11. Pollyea DA, Kohrt HE, Gallegos L, Figueroa ME, Abdel-Wahab O, Zhang B, et al. Safety, efficacy and biological predictors of response to sequential azacitidine and lenalidomide for elderly patients with acute myeloid leukemia. *Leukemia.* 2012;26(5):893–901.
  12. Wei A, Tan P, Perruzza S, Govindaraj C, Fleming S, McManus J, et al. Maintenance lenalidomide in combination with 5-azacitidine as post-remission therapy for acute myeloid leukaemia. *Br J Haematol.* 2015;169(2):199–210.
  13. Hills RK, Burnett AK. Applicability of a "Pick a Winner" trial design to acute myeloid leukemia. *Blood.* 2011;118(9):2389–94.
  14. Peto R, Pike MC, Armitage P, Breslow NE, Cox DR, Howard SV, et al. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. II. Analysis and examples. *Br J Cancer.* 1977;35(1):1–39.

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Copland M, Ariti C, Thomas IF, Upton L, Sydenham M, Mehta P, et al. A randomised evaluation of low-dose cytosine arabinoside plus lenalidomide versus single-agent low-dose cytosine arabinoside in older patients with acute myeloid leukaemia: Results from the LI-1 trial. *Br J Haematol.* 2023;00:1–6. <https://doi.org/10.1111/bjh.19220>