



DATA NOTE

The genome sequence of the common earthworm, *Lumbricus terrestris* (Linnaeus, 1758) [version 1; peer review: 3 approved]

Mark L. Blaxter¹, David Spurgeon², Peter Kille³,
Wellcome Sanger Institute Tree of Life programme,
Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective,
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹Tree of Life, Wellcome Sanger Institute, Hinxton, England, UK

²UK Centre for Ecology & Hydrology, Wallingford, England, UK

³Cardiff School of Biosciences, Cardiff University, Cardiff, Wales, UK

V1 First published: 30 Oct 2023, 8:500
<https://doi.org/10.12688/wellcomeopenres.20178.1>
Latest published: 30 Oct 2023, 8:500
<https://doi.org/10.12688/wellcomeopenres.20178.1>

Abstract

We present a genome assembly from an individual *Lumbricus terrestris* (the common earthworm; Annelida; Clitellata; Haplotaxida; Lumbricidae). The genome sequence is 1,056.5 megabases in span. Most of the assembly is scaffolded into 18 chromosomal pseudomolecules. The mitochondrial genome has also been assembled and is 15.93 kilobases in length.

Keywords

Lumbricus terrestris, common earthworm, genome sequence, chromosomal, Haplotaxida



This article is included in the [Tree of Life gateway](#).

Open Peer Review

Approval Status

	1	2	3
version 1			
30 Oct 2023	view	view	view

1. **Yoshiaki Morino** , University of Tsukuba, Tsukuba, Japan
2. **Sudhakar Sivasubramaniam**, Manonmaniam Sundaranar University, Tirunelveli, India
3. **Arun Arumugaperumal** , Rajalakshmi Engineering College, Chennai, India
3. **José María Martín-Durán** , Queen Mary University of London, London, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: **Blaxter ML:** Investigation, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Spurgeon D:** Resources; **Kille P:** Resources;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (206194) and the Darwin Tree of Life Discretionary Award (218328).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2023 Blaxter ML *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Blaxter ML, Spurgeon D, Kille P *et al.* **The genome sequence of the common earthworm, *Lumbricus terrestris* (Linnaeus, 1758) [version 1; peer review: 3 approved]** Wellcome Open Research 2023, 8:500

<https://doi.org/10.12688/wellcomeopenres.20178.1>

First published: 30 Oct 2023, 8:500 <https://doi.org/10.12688/wellcomeopenres.20178.1>

Species taxonomy

Eukaryota; Metazoa; Eumetazoa; Bilateria; Protostomia; Spiralia; Lophotrochozoa; Annelida; Clitellata; Oligochaeta; Crassicitellata; Lumbricina; Lumbricidae; Lumbricinae; *Lumbricus*; *Lumbricus terrestris* complex; *Lumbricus terrestris* (Linnaeus, 1758) (NCBI:txid6398).

Background

The Darwin Tree of Life project aims to sequence to reference quality the genomes of species of Ireland and Britain (Blaxter *et al.*, 2022). While most species we target make their homes on these islands, some are ecosystem engineers that shape the environments in which they and others live. One such group is the lumbricid annelids, or earthworms, which collectively generate and maintain soils by processing and degrading mineral particles, cycling biological material into the soil, aerating and draining soil through their burrows and providing rich habitats for soil microbes in their casts (Kooch & Jalilvand, 2008). The importance of earthworms as not just inhabitants but prime generators of soil was noted by Charles Darwin in his last book “*The Formation of Vegetable Mould Through the Action of Worms, with Observations on their Habits*” (Darwin, 1881).

Lumbricus terrestris, the common earthworm (see (NBN Atlas Partnership, 2023)) was described by Linnaeus in his “*Systema Naturae*” in 1758 (genus #246, on pages 647–648 in a paraphyletic “*Vermes*”) (Linnaeus, 1758). Now placed in Lumbricidae within the oligochaete annelids (phylum Annelida, class Oligochaeta, order Crassicitellata), *L. terrestris* is the largest of the earthworms of Ireland and Britain, and has, since Darwin, been the focus of intensive study especially in the areas of ecotoxicology and ecotoxicogenomics (Brulle *et al.*, 2010; Sheppard, 1998; Sturzenbaum *et al.*, 2009). As *L. terrestris*, like other earthworms, carries out respiratory exchange through its skin, and consumes and processes soil matter through its gut, it is particularly exposed to geogenic and anthropogenic contamination of soils (Stürzenbaum *et al.*, 2004). Here we present a chromosomally-complete assembly of *L. terrestris*, collected from the Wellcome Sanger Institute campus, to support ongoing work on this keystone species.

Genome sequence report

The genome was sequenced from one *Lumbricus terrestris* (Figure 1) collected from Wellcome Genome Campus, Hinxton, UK (52.08, 0.18). We generated 33-fold coverage using Pacific Biosciences single-molecule HiFi long reads. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data. Manual assembly curation corrected 199 missing joins or mis-joins and removed 221 haplotypic duplications, reducing the assembly length by 11.9% and the scaffold number by 33.33%, and decreasing the scaffold N50 by 4.45%.

The final assembly has a total length of 1056.5 Mb in 361 sequence scaffolds with a scaffold N50 of 56.6 Mb (Table 1).



Figure 1. Photograph of *Lumbricus terrestris* by Michael Linnenbach.

The snailplot in Figure 2 provides a summary of the assembly statistics, while the distribution of assembly scaffolds on GC proportion and coverage is shown in Figure 3. The cumulative assembly plot in Figure 4 shows curves for subsets of scaffolds assigned to different phyla. Most (98.77%) of the assembly sequence was assigned to 18 chromosomal-level scaffolds. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size (Figure 5; Table 2). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.

The estimated Quality Value (QV) of the final assembly is 57.3 with *k*-mer completeness of 99.99%, and the assembly has a BUSCO v5.3.2 completeness of 91.4% (single = 88.1%, duplicated = 3.4%), using the metazoa_odb10 reference set (*n* = 954).

Metadata for specimens, spectral estimates, sequencing runs, contaminants and pre-curation assembly statistics can be found at <https://links.tol.sanger.ac.uk/species/6398>.

Methods

Sample acquisition and nucleic acid extraction

Lumbricus terrestris specimens were collected from Wellcome Genome Campus (latitude 52.08, longitude 0.18) on 2020-03-12. The specimen was taken from soil in woodland by turning leaf litter and subsoil. The collectors were Mark Blaxter (Wellcome Sanger Institute), Peter Kille (University of Cardiff) and David Spurgeon (UK Centre for Ecology & Hydrology) and David Spurgeon identified the specimen. The specimen was flash frozen in liquid nitrogen. One specimen

Table 1. Genome data for *Lumbricus terrestris*, wLumTerr1.1.

Project accession data		
Assembly identifier	wLumTerr1.1	
Species	<i>Lumbricus terrestris</i>	
Specimen	wLumTerr1	
NCBI taxonomy ID	6398	
BioProject	PRJEB59400	
BioSample ID	SAMEA7524126	
Isolate information	wLumTerr1: bodywall and intestine (DNA sequencing and Hi-C scaffolding) wLumTerr2: bodywall (RNA sequencing)	
Assembly metrics*		Benchmark
Consensus quality (QV)	57.3	≥ 50
<i>k</i> -mer completeness	99.99%	≥ 95%
BUSCO**	C:91.4%[S:88.1%,D:3.4%], F:3.6%,M:5.0%,n:954	C ≥ 95%
Percentage of assembly mapped to chromosomes	98.77%	≥ 95%
Sex chromosomes	-	localised homologous pairs
Organelles	Mitochondrial genome assembled	complete single alleles
Raw data accessions		
PacificBiosciences SEQUEL II	ERR10841333, ERR10841334, ERR10841335	
Hi-C Illumina	ERR10851548, ERR10851550	
PolyA RNA-Seq Illumina	ERR10851549	
Genome assembly		
Assembly accession	GCA_949752735.1	
Accession of alternate haplotype	GCA_949752785.1	
Span (Mb)	1056.5	
Number of contigs	1587	
Contig N50 length (Mb)	1.6	
Number of scaffolds	361	
Scaffold N50 length (Mb)	56.6	
Longest scaffold (Mb)	98.9	

* Assembly metric benchmarks are adapted from column VGP-2020 of “Table 1: Proposed standards and metrics for defining genome assembly quality” from (Rhie *et al.*, 2021).

** BUSCO scores based on the metazoa_odb10 BUSCO set using v5.3.2. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at <https://blobtoolkit.genomehubs.org/view/wLumTerr1.1/dataset/CATKHU01/busco>.

was used for DNA sequencing and Hi-C data (specimen ID SAN0001206, ToLID wLumTerr1) and another specimen (specimen ID SAN0001207, ToLID wLumTerr2) was used for RNA sequencing.

DNA was extracted at the Tree of Life laboratory, Wellcome Sanger Institute (WSI). The wLumTerr1 sample was weighed and dissected on dry ice with tissue set aside for Hi-C sequencing. Bodywall and intestinal tissue was disrupted




Figure 2. Genome assembly of *Lumbricus terrestris*, wLumTerr1.1: metrics. The BlobToolKit Snailplot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 1,056,547,181 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (98,895,420 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (56,588,617 and 43,887,615 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the metazoa_odb10 set is shown in the top right. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/wLumTerr1.1/dataset/CATKHU01/snail>.

using a Nippi Powermasher fitted with a BioMasher pestle. High molecular weight (HMW) DNA was extracted using the Qiagen MagAttract HMW DNA extraction kit. HMW DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system with speed setting 30. Sheared DNA was purified by solid-phase reversible immobilisation using AMPure PB beads with a 1.8X ratio of beads to sample to remove the shorter fragments and concentrate the DNA sample. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

RNA was extracted from tissue of wLumTerr2 in the Tree of Life Laboratory at the WSI using TRIzol, according to the manufacturer's instructions. RNA was then eluted in 50 μ l

RNAse-free water and its concentration assessed using a Nanodrop spectrophotometer and Qubit Fluorometer using the Qubit RNA Broad-Range (BR) Assay kit. Analysis of the integrity of the RNA was done using Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay.

Sequencing

Pacific Biosciences HiFi circular consensus DNA sequencing libraries were constructed according to the manufacturers' instructions. Poly(A) RNA-Seq libraries were constructed using the NEB Ultra II RNA Library Prep kit. DNA and RNA sequencing was performed by the Scientific Operations core at the WSI on Pacific Biosciences SEQUEL II (HiFi) and Illumina HiSeq 4000 (RNA-Seq) instruments. Hi-C data were also generated from body wall and intestinal tissue of wLumTerr1 using the Arima2 kit and sequenced on the HiSeq X Ten and Illumina NovaSeq 6000 instrument.




Figure 3. Genome assembly of *Lumbricus terrestris*, wLumTerr1.1: BlobToolKit GC-coverage plot. Scaffolds are coloured by phylum. Circles are sized in proportion to scaffold length. Histograms show the distribution of scaffold length sum along each axis. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/wLumTerr1.1/dataset/CATKHU01/blob>.

Genome assembly, curation and evaluation

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021) and haplotypic duplication was identified and removed with purge_dups (Guan *et al.*, 2020). The assembly was then scaffolded with Hi-C data (Rao *et al.*, 2014) using YaHS (Zhou *et al.*, 2023). The assembly was checked for contamination and corrected as described previously (Howe *et al.*, 2021). Manual curation was performed using HiGlass (Kerpedjiev *et al.*, 2018) and Pretext (Harry, 2022). The mitochondrial genome was assembled using MitoHiFi

(Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) or MITOS (Bernt *et al.*, 2013) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.

A Hi-C map for the final assembly was produced using bwa-mem2 (Vasimuddin *et al.*, 2019) in the Cooler file format (Abdennur & Mirny, 2020). To assess the assembly metrics, the *k*-mer completeness and QV consensus quality values were calculated in Merqury (Rhie *et al.*, 2020). This




Figure 4. Genome assembly of *Lumbricus terrestris*, wLumTerr1.1: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/wLumTerr1.1/dataset/CATKHU01/cumulative>.




Figure 5. Genome assembly of *Lumbricus terrestris*, wLumTerr1.1: Hi-C contact map of the wLumTerr1.1 assembly, visualised using HiGlass. Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/l/?d=OFTjARz6QZKy250DRKimLg>.

Table 2. Chromosomal pseudomolecules in the genome assembly of *Lumbricus terrestris*, wLumTerr1.

INSDC accession	Chromosome	Length (Mb)	GC%
OX457036.1	1	98.9	40.5
OX457037.1	2	84.43	40.5
OX457038.1	3	77.17	41.0
OX457039.1	4	72.47	40.5
OX457040.1	5	69.63	40.5
OX457041.1	6	63.79	41.0
OX457042.1	7	61.74	41.0
OX457043.1	8	56.59	41.0
OX457044.1	9	55.79	41.0
OX457045.1	10	49.13	41.0
OX457046.1	11	47.79	40.5
OX457047.1	12	46.9	41.0
OX457048.1	13	46.6	41.0
OX457049.1	14	46.28	40.5
OX457050.1	15	45.14	40.5
OX457051.1	16	43.89	41.0
OX457052.1	17	39.89	41.0
OX457053.1	18	36.07	41.5
OX457054.1	MT	0.02	36.0

work was done using Nextflow (Di Tommaso *et al.*, 2017) DSL2 pipelines “sanger-tol/readmapping” (Surana *et al.*, 2023a) and “sanger-tol/genomenote” (Surana *et al.*, 2023b). The genome was analysed within the BlobToolKit environment (Challis *et al.*, 2020) and BUSCO scores (Manni *et al.*, 2021; Simão *et al.*, 2015) were calculated.

Table 3 contains a list of relevant software tool versions and sources.

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material

Table 3. Software tools: versions and sources.

Software tool	Version	Source
BlobToolKit	4.1.7	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.3.2	https://gitlab.com/ezlab/busco
Hifiasm	0.16.1-r375	https://github.com/chhylp123/hifiasm
HiGlass	1.11.6	https://github.com/higlass/higlass
Mercury	MercuryFK	https://github.com/thegenemyers/MERQUERY.FK
MitoHiFi	2	https://github.com/marcelauliano/MitoHiFi
PretextView	0.2	https://github.com/wtsi-hpag/PretextView
purge_dups	1.2.3	https://github.com/dfguan/purge_dups
sanger-tol/genomenote	v1.0	https://github.com/sanger-tol/genomenote
sanger-tol/readmapping	1.1.0	https://github.com/sanger-tol/readmapping/tree/1.1.0
YaHS	1.2a	https://github.com/c-zhou/yahs

- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Lumbricus terrestris* (common earthworm). Accession number PRJEB59400; <https://identifiers.org/ena.embl/PRJEB59400>. (Wellcome Sanger Institute, 2023) The genome sequence is released openly for reuse. The *Lumbricus terrestris* genome sequencing initiative is part of the Darwin Tree of Life (DToL) project. All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the [Ensembl](#) pipeline at

the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#).

Author information

Members of the Wellcome Sanger Institute Tree of Life programme are listed here: <https://doi.org/10.5281/zenodo.4783585>.

Members of Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective are listed here: <https://doi.org/10.5281/zenodo.4790455>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.5013541>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

Acknowledgements

Peter Kille, David Spurgeon and Mark Blaxter would like to acknowledge the inspirational mentorship of A. John Morgan (1948–2016) in all things *Lumbricus*.

References

- Abdennur N, Mirny LA: **Cooler: Scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics.* 2020; **36**(1): 311–316. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, et al.: **MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bernt M, Donath A, Jühling F, et al.: **MITOS: Improved *de novo* metazoan mitochondrial genome annotation.** *Mol Phylogenet Evol.* 2013; **69**(2): 313–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Blaxter M, Mieszkowska N, Di Palma F, et al.: **Sequence locally, think globally: The Darwin Tree of Life Project.** *Proc Natl Acad Sci U S A.* 2022; **119**(4): e2115642118. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brulle F, Morgan AJ, Cocquerelle C, et al.: **Transcriptomic underpinning of toxicant-mediated physiological function alterations in three terrestrial invertebrate taxa: A review.** *Environ Pollut.* 2010; **158**(9): 2793–2808. [PubMed Abstract](#) | [Publisher Full Text](#)
- Challis R, Richards E, Rajan J, et al.: **BlobToolKit - interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, et al.: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Darwin C: **The Formation of Vegetable Mould through the Action of Worms, with Observations on Their Habits.** London, UK: John Murray, 1881. [Reference Source](#)
- Di Tommaso P, Chatzou M, Floden EW, et al.: **Nextflow enables reproducible computational workflows.** *Nat Biotechnol.* 2017; **35**(4): 316–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Guan D, McCarthy SA, Wood J, et al.: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired REad TEXTure Viewer): A desktop application for viewing pretext contact maps.** 2022; [Accessed 19 October 2022]. [Reference Source](#)
- Howe K, Chow W, Collins J, et al.: **Significantly improving the quality of genome assemblies through curation.** *GigaScience.* Oxford University Press, 2021; **10**(1): gjaa153. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, et al.: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kooch Y, Jalilvand H: **Earthworms as Ecosystem Engineers and the Most Important Detritivores in Forest Soils.** *Pak J Biol Sci.* 2008; **11**(6): 819–825. [PubMed Abstract](#) | [Publisher Full Text](#)
- Linnaeus C: **Systema Naturae.** 10th ed. Stockholm: Holmiae Salvius, 1758.
- Manni M, Berkeley MR, Seppely M, et al.: **BUSCO update: Novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- NBN Atlas Partnership: ***Lumbricus terrestris* Linnaeus, 1758 Common Earthworm.** *NBN Atlas.* 2023; [Accessed 9 September 2023]. [Reference Source](#)
- Rao SSP, Huntley MH, Durand NC, et al.: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, et al.: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, Walenz BP, Koren S, et al.: **Merqury: Reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol.* 2020; **21**(1): 245. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sheppard S: **Advances in Earthworm Ecotoxicology: Proceedings from the Second International Workshop on Earthworm Ecotoxicology, April 1997, Amsterdam, the Netherlands.** Setac Press, 1998. [Reference Source](#)
- Simão FA, Waterhouse RM, Ioannidis P, et al.: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.** *Bioinformatics.* 2015; **31**(19): 3210–3212. [PubMed Abstract](#) | [Publisher Full Text](#)
- Stürzenbaum SR, Andre J, Kille P, et al.: **Earthworm genomes, genes and proteins: the (re)discovery of Darwin's worms.** *Proc Biol Sci.* 2009; **276**(1658): 789–797. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stürzenbaum SR, Georgiev O, Morgan AJ, et al.: **Cadmium Detoxification in**

Earthworms: From Genes to Cells. *Environ Sci Technol.* 2004; **38**(23): 6283–6289.
[PubMed Abstract](#) | [Publisher Full Text](#)

Surana P, Muffato M, Qi G: **sanger-tol/readmapping: sanger-tol/readmapping v1.1.0 - Hebridean Black (1.1.0).** *Zenodo.* 2023a; [Accessed 21 July 2023].
[Publisher Full Text](#)

Surana P, Muffato M, Sadasivan Baby C: **sanger-tol/genomenote (v1.0.dev).** *Zenodo.* 2023b; [Accessed 21 July 2023].
[Publisher Full Text](#)

Uljiano-Silva M, Ferreira JGRN, Krashennikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity**

reads. *BMC Bioinformatics.* 2023; **24**(1): 288.

[Publisher Full Text](#)

Vasimuddin M, Misra S, Li H, *et al.*: **Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS).* IEEE, 2019; 314–324.
[Publisher Full Text](#)

Wellcome Sanger Institute: **The genome sequence of the common earthworm, *Lumbricus terrestris* (Linnaeus, 1758).** European Nucleotide Archive. [dataset], accession number PRJEB59400, 2023.

Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:   

Version 1

Reviewer Report 25 January 2024

<https://doi.org/10.21956/wellcomeopenres.22341.r71542>

© 2024 **Martín-Durán J.** This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

 **José María Martín-Durán** 

School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK

This genome note reports the assembly of the common earthworm *Lumbricus terrestris*. This is a broadly distributed segmented worm, very familiar to the public. The assembly meets the high standards of the Darwin Tree of Life initiative and will be an essential resource to a diverse range of disciplines, from developmental biology to ecology, ecotoxicology and conservation.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Evolutionary biology, comparative genomics, evo-devo, annelids

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 19 January 2024

<https://doi.org/10.21956/wellcomeopenres.22341.r71560>

© 2024 Sivasubramaniam S et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Sudhakar Sivasubramaniam**

Department of Biotechnology, Manonmaniam Sundaranar University, Tirunelveli, Tamil Nadu, India

Arun Arumugaperumal 

Department of Biotechnology, Rajalakshmi Engineering College, Chennai, Tamilnadu, India

The authors report the genome sequence of *Lumbricus terrestris* earthworm. The genome size was 1056.5 Mbp and the sequences were assembled in to 18 chromosomes. The genome completeness analysis showed that the assembly was 91.4% complete.

The authors have used best performing softwares for the bioinformatics work. They could have gone an extra mile to predict the non-coding genes present in the genome. It would have been an interesting data to look in to the exon-intron junctions and gene structure.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Regeneration and stem cell biology, Genomics and Transcriptomics

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 19 January 2024

<https://doi.org/10.21956/wellcomeopenres.22341.r71561>

© 2024 Morino Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Yoshiaki Morino 

Institute of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

The paper reports genome sequencing of the common earthworm, *Lumbricus terrestris*. The chromosome-level assembly was generated using standard methods and is sufficiently high quality. The significance of the species is well introduced. I have only two minor comments:

1. Concerning the absence of sex chromosomes, it would be beneficial to include a mention in the manuscript that earthworms, including this species, are generally hermaphrodites.
2. The manuscript should describe the source tissues used for the RNA-seq.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Marine invertebrate, Evo-Devo

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
