

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/167641/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Tseng, Chun-Chih, Woolley, Thomas , Jiang, Ting-Xin, Wu, Ping, Maini, Philip, Widelitz, Randall and Chuong, Cheng-Ming 2024. Inhibition of gap junctions stimulates Turing-type periodic feather pattern formation during chick skin development. PLoS Biology

Publishers page:

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



S1 Appendix

Technical details for the computational simulations

All codes and data can be found at <https://zenodo.org/doi/10.5281/zenodo.10865520>. Definitions of the parameters and active regions during each simulation stage. Note the threshold value defines the morphogen region that becomes feather placode.

Since we are dealing with nondimensional parameter values pattern sizes are simply relative to one another. Thus, parameter values were chosen to both produce patterns and so that the final relative sizes of the patterns were consistent with observations.

Figure 13B

| Stage | (D_{u_i}, D_{v_i}) | Ω_i | Threshold |
|-------|----------------------|---|-----------------|
| 1 | (1,40) | $[0,15] \times [0,25]$, | $u_{Bud} > 1.5$ |
| 2 | (3/4,30) | $u_{Bud} < 1.5$ | $u_1 > 2.6$ |
| 3 | (2/5,8) | $u_1 < 0.59$ | $u_2 > 2.2$ |
| 4 | (1/10,4) | $u_2 < 0.8$ | $u_3 > 2.4$ |
| 5 | (1/10,4) | $(u_{Bud} > 0.8)(u_1 > 0.9)(u_2 > 0.9)$ | $u_4 > 2.55$ |

Figure 13C

| Stage | (D_{u_i}, D_{v_i}) | Ω_i | Threshold |
|-------|----------------------|---|-----------------|
| 1 | (1,40) | $[0,15] \times [0,25]$, | $u_{Bud} > 1.5$ |
| 3 | (2/5,8) | $u_{Bud} < 1.5$ | $u_2 > 2.2$ |
| 4 | (1/10,4) | $u_3 < 0.8$ | $u_3 > 2.4$ |
| 5 | (1/10,4) | $(u_{Bud} > 0.8)(u_1 > 0.9)(u_2 > 0.9)$ | $u_4 > 2.55$ |

Figure 13D

| Stage | (D_{u_i}, D_{v_i}) | Ω_i | Threshold |
|---------|----------------------|---|-----------------|
| 1 | (1,40) | $[0,15] \times [0,25]$, | $u_{Bud} > 1.5$ |
| 2 | (3/4,30) | $u_{Bud} < 1.5$ | $u_1 > 2.6$ |
| 3, 4, 5 | (1/30,4/3) | $(u_1 > 0.8)(u_1 < 1.2)((y < 2)(y > -6) + (y < 15)(y > 6) + (y < -11))$ | $u_2 > 2.2$ |

Initial conditions

For the initial bud simulation the initial condition is a spot of activator in the centre,

$I_1(x, y) = \exp(-(x - 7.5)^2 - (y - 12.5)^2)$. Although a specific condition is used, it is a fairly weak requirement because the same pattern will form under any initial conditions due to the wavelengths chosen by the Turing pattern. The only difference will be that the pattern will be translated left, or right, because of

the periodic boundary conditions. Thus, we choose the centered Gaussian to centralize the pattern and, hence, provide simple visual results.

For every other stage the last condition is a source of activator at the back of the bud, $I(x, y) = \exp(-x^2 - (y + 10.5)^2) + \exp(-x^2 - (y - 22.5)^2) + \exp(-(x - 15)^2 - (y - 22.5)^2)$.