



In vitro acaricidal activity of several natural products against ibex-derived *Sarcoptes scabiei*

Jesús M. Pérez^{a,*}, Emiliano N. Jesser^{b,2}, Jorge O. Werdin^{b,3}, Colin Berry^{c,4}, Mohamed A. Gebely^{c,d,5}, Raquel Crespo-Ginés^{a,e,6}, José E. Granados^{f,7}, Antonio J. López-Montoya^{g,8}

^a Departamento de Biología Animal, Vegetal y Ecología, Universidad de Jaén, Campus Las Lagunillas, s.n., Jaén E-23071, Spain

^b Departamento de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur, San Juan 670, Bahía Blanca B 8000CPB, Argentina

^c School of Biosciences, Cardiff University, Museum Avenue, Cardiff CF10 3AX, UK

^d Department of Parasitology and Animal Diseases, Veterinary Research Institute, National Research Centre, Dokki, Giza 12622, Egypt

^e Instituto de Investigación en Recursos Cinegéticos (IREC-CSIC, UCLM, JCCM), Ronda de Toledo 12, Ciudad Real E-13071, Spain

^f Centro Administrativo Parque Nacional y Parque Natural Sierra Nevada, Carretera Antigua Sierra Nevada, Km 7, E-18071, Pinos Genil, Granada, Spain

^g Department of Statistics and Operational Research, Jaén University, Campus Las Lagunillas, s.n., Jaén E-23071, Spain

ARTICLE INFO

Keywords:

Bacillus thuringiensis

Essential plant oils

Nanoparticles

Sarcoptes scabiei

Survival

ABSTRACT

In this study we analysed the effect of the temperature, diverse strains of *Bacillus thuringiensis*, *Lysinibacillus sphaericus* and nanoformulations with essential plant oils (EONP) on the survival of *Sarcoptes scabiei* mites derived from naturally-infested Iberian ibex (*Capra pyrenaica*). In general, mites maintained at 12°C survived more than those maintained at 35°C (40.7 hr and 31.2 hr, respectively). Mites with no treatment survived 27.6 h on average. Mites treated with *B. thuringiensis* serovar. *konkukian* and geranium EONP showed significant reduction in their survival. Despite the fact that these agents seem to be promising candidates for controlling sarcoptic mange in the field, further research is still needed to get stable, efficient and eco-friendly acaricides.

1. Introduction

Sarcoptes scabiei is an astigmatid mite causing a dermal disease, namely sarcoptic mange, in domestic and wild mammalian hosts, including man, worldwide, reaching high morbidity and mortality rates (Bornstein et al., 2001; Pence and Ueckermann, 2002; Arlian and Morgan, 2017). Transmission of this mite between susceptible hosts may be direct, indirect, or a combination of both (Browne et al., 2022).

Control of this disease in wild populations is a challenging task. Although multiple doses of subcutaneous ivermectin (200–400 µg/kg) is the treatment most commonly used (Rowe et al., 2019), its implementation with free-ranging animals is very difficult from a logistic

viewpoint. Moreover, this approach may have undesirable impact on non-target organisms and favours the development of resistance by the mite (Walton et al., 2000), among other “secondary” effects (Moroni et al., 2020).

The development of resistance of *S. scabiei* against acaricidal compounds is increasing (Currie et al., 2004; Mounsey et al., 2010; Andriantsoanirina et al., 2014). Therefore, it is crucial to develop new drugs for treating scabies (Walton et al., 2004). *Bacillus thuringiensis* is a Gram-positive bacterium which produces one or several crystalline proteins referred to as δ-endotoxins (Hill and Pinnock, 1998). After being ingested by susceptible arthropods, the *B. thuringiensis* δ-endotoxin crystals are dissolved in the midgut with a consequent production of

* Corresponding author.

E-mail address: jperez@ujaen.es (J.M. Pérez).

¹ 0000-0001-9159-0365

² 0000-0002-4526-6449

³ 0000-0001-5152-3782

⁴ 0000-0002-9943-548X

⁵ 0000-0003-3127-2781

⁶ 0000-0001-7901-7670

⁷ 0000-0002-9787-9896

⁸ 0000-0002-0453-7978

activated toxic polypeptides commonly known as δ -endotoxin crystal proteins (Cry proteins) (Höfte and Whiteley, 1989), which may belong to a number of distinct structural families (Crickmore et al., 2021). These toxins seem to disrupt the selective permeability of the cell membrane, which ultimately causes the arthropod death from starvation and/or septicemia (Knowles & Dow, 1993). Like *B. thuringiensis*, *Lysinibacillus sphaericus* is another gram positive bacterium able to produce a range of insecticidal proteins and which exerts its effects in a similar manner (Berry, 2012).

Essential oils (EO) are mixtures of diverse volatile compounds synthesized by plants to protect themselves and are considered as new ecofriendly insecticides, since they may show good biological activity against a number of insect pests, low toxicity to humans and rapid degradation in the environment (Jesser et al., 2020a). However, EO show some disadvantages such as instability, volatility, and low solubility in water, which may limit their applications (Jesser et al., 2020b).

In recent years, the integration of nanotechnology in the field of biopesticides has garnered significant attention. An innovative method, the nanoformulation of EO, has emerged as a solution to shield active compounds from environmental conditions and prevent the gradual loss of EO (Kumar et al., 2020). Several materials, including proteins, synthetic emulsifiers, polysaccharides (such as starch and chitosan), and polyethers (like polyethylene glycol and poly- ϵ -caprolactone), have been explored for their effectiveness in nanoformulating EO or its constituents (Athanasios et al., 2018; De Luca et al., 2021). Polyethylene glycol 6000 (PEG 6000) has been extensively investigated over the past few decades for medical, food industry, and pest control applications. This material showed a broad range of solubility, lacks antigenicity and immunotoxicity, and is easily excreted from living organisms without toxicity concerns. PEG 6000 polymeric nanoparticles loaded with EO (EONP) are considered as one of the most important emerging trends in insect pest control (Campolo et al., 2017; Werdin-González et al., 2014; 2017).

Nanoformulations of *B. thuringiensis* Cry proteins and EO contribute to increase the activity period of such active compounds in the environment and allow reductions in the amount to be used (de Oliveira et al., 2014; De Oliveira et al., 2021).

The aim of this study was to test the potential acaricide effect of several treatments with *B. thuringiensis* and *Lysinibacillus sphaericus* Cry proteins and essential oils nanoparticles formulations.

2. Materials and methods

2.1. Preparation of *Bacillus thuringiensis* and *Lysinibacillus sphaericus*

Bacillus thuringiensis and *Lysinibacillus sphaericus* spore/crystal preparations were produced using a method previously described (Jones et al., 2008), by growing the strains in Embrapa medium (Monnerat et al., 2007) until approximately 95% sporulation (judged by phase contrast microscopy) after which spores and crystals were harvested by centrifugation and washed in distilled water before lyophilisation and storage at 4°C. Bacterial strains used were obtained as follows: *Lysinibacillus sphaericus* strain IAB59, *Bacillus thuringiensis* serovar. *higo* strain T44001, *Bacillus thuringiensis* serovar. *israelensis* strain 4Q7, *Bacillus thuringiensis* serovar. *kurstaki* strain HD1 from the *Bacillus* Genetic Stock Center: *Bacillus thuringiensis* GP 138, a kind gift from Prof Alejandra Bravo, UNAM, Mexico; and *Bacillus thuringiensis* serovar. *konkukian* strain 97–27 and *Lysinibacillus sphaericus* strain 2362 from the collection of the Pasteur institute, France.

2.2. Preparation of nanoparticles

The EONP was synthesized using the melt-dispersion method, a procedure previously outlined by Werdin-González et al. (2014). Initially, 20 g of PEG 6000 were heated to 65°C on a hotplate stirrer. Subsequently, 2 g of geranium (*Geranium maculatum*) or peppermint

(*Mentha piperita*) EO were added to the molten PEG 6000. Geranium and peppermint oils were purchased from Swiss-Just (Switzerland). PEG 6000 was acquired from Merck, Germany. Concurrently, the mixture of PEG 6000 and EO was stirred with a D-500 Handheld Homogenizer (D-lab instrument limited) for 15 minutes at 15,000 rpm. The EONP spontaneously formed when the mixture was cooled to -4°C . After 45 minutes at that temperature, the resulting mixture was thoroughly ground in a refrigerated mortar box at 0°C , and the product was sifted through a stainless-steel sieve with a mesh size of 230. The EONP were stored in airtight polyethylene pouches at $27 \pm 2^{\circ}\text{C}$ within desiccators containing calcium chloride for seven days before further experimentation.

2.3. EO and EOPN composition

According to Yeguerman et al. (2022), the chemical composition of both pre- and post-formulation essential oils (EOs) was analysed using gas chromatography-mass spectrometry (GC-MS) with an Agilent 7890B gas chromatograph coupled to an Agilent 5977 A mass spectrometer. A HP-5MS capillary column (30 m \times 0.25 mm i.d. \times 0.25 μm film thickness) was utilized, with helium serving as the carrier gas at a flow rate of 1.0 mL min $^{-1}$. The oven temperature was initially set at 50°C for 2 minutes, then ramped at $5^{\circ}\text{C min}^{-1}$ to 200°C and held for 15 minutes. Injection block temperature was maintained at 280°C , and 1 μL aliquots of samples were injected. Ionization energy of 70 eV was used for mass spectrometry, scanning from 35 to 550 m/z . Retention indices (RI) of components were determined using a series of n-alkanes (C8-C20). Further identification was accomplished using NIST 2.0 database. Relative percentages of individual components were calculated by averaging gas chromatography with flame ionization detection (GC-FID) peak areas obtained on a DB-5 column under similar conditions. Key components like α -pinene, limonene, menthol, pulegone, and geraniol were confirmed by comparing with their standard samples (Sigma-Aldrich) via co-injection. Essential oils were extracted from polymeric nanoparticles (EOPN) by dissolving 0.5 g of each sample in 5 mL of distilled water, heating at 65°C for 30 minutes with magnetic stirring. Upon melting of PEG 6000, 4 mL of petroleum ether was added, and the mixture was stirred for 2 hours. Afterward, the ether phase containing the extracted EOs was collected, diluted to a concentration of 0.001 mg mL $^{-1}$ (0.1% v/v), and subjected to GC-MS and GC analysis for component identification.

2.4. EONP size measurement

A Malvern Nano ZS90 instrument was used to determine the size of the EONP. The Polydispersity Index (PDI) was calculated as the square of the standard deviation divided by the square of the mean size, serving as an indicator of the homogeneity or heterogeneity in the size distribution of the particles, following the method described by Pascoli et al. (2018). Each sample, consisting of 0.2 g of EONP, was suspended in 10 mL of distilled water for 30 minutes. Subsequently, the dispersion was filtered using Whatman N $^{\circ}$ 1 filter paper and allowed to equilibrate for 2 hours. Data were statistically compared using one-way analysis of variance (ANOVA), followed by the LSD test (N = 4).

2.5. EONP encapsulation efficiency

As outlined by Werdin-González et al. (2014) the encapsulation efficiency was assessed using spectrophotometric methods. For this, 0.1 g of EONP were individually dissolved in 2 mL of an absolute ethanol-water solution (75:25). The resulting mixture was then centrifuged at 9000 rpm for 10 minutes. The supernatant was carefully collected and subjected to analysis via UV-vis spectrophotometry, employing a Shimadzu UV-1203 photometer with the Kinetics-2-Program Pack (P/N 206–62029–10; Shimadzu Corp., Kyoto, Japan) at a wavelength of 290 nm. This process was repeated for four

samples, and the quantity of EO was determined by referring to an appropriate calibration curve for free EO in ethanol.

Encapsulation efficiency (EE) was determined from:

$$EE(\%) = \frac{\text{weight of loaded EO}}{\text{weight of initial EO}} \times 100$$

One way analysis of variance (ANOVA) and LSD were used in order to compare the data (N = 4). The mean physical and chemical characteristics of the EONP formulations are included in Table 2 and 3, respectively.

2.6. Mite collection

Ibex (*Capra pyrenaica*) with severe mange in the consolidation and chronic stages (Espinosa et al., 2017), with lesions affecting ≥ 50% of the host skin surface were selected as mite donors (Fig. 1a). The ibex were chemically immobilized with a mixture of ketamine (3 mg/kg) and xylazine (3 mg/kg) (Casas-Díaz et al., 2011), and then euthanized with T-61 Intervet® (combination of embutramide and mebezonium iodide) at a dose of 1 mL/1.5 kg.

For mite extraction we painted glass Petri dishes black (14 cm diameter), except in a central circle at the bottom (5.5. cm diameter). Then, a 25 W lamp was placed 7–8 cm below the central circle and several skin pieces from the donor ibex were placed around this circle (Fig. 1b). In this way, a temperature gradient was created allowing the mites to concentrate in the central area of the dish (Andrews, 1981) after overnight exposure to the lamp (Fig. 1c). Once the mites left the host skin, the skin pieces were removed (Fig. 1d). The aim of this method was to obtain live mites without manipulating them, therefore, avoiding mechanical damage to mites which could affect their survival.

The first assay (including mite extraction and following counts) was carried out at 12°C and 70% relative humidity (RH). During the remaining assays the plates were maintained in an incubator at 35°C and 45% RH. The number of control plates and treatments of each assay are included in Table 1. After an initial count (including both live and dead mites), live mites (those showing some kind of movement) were counted twice a day until the death of all the mites.

The acaricidal activity of geranium EONP and peppermint EONP was

Table 1
Dates of assays for analyse mite survival and the treatments tested.

	Assay 1 (19 Nov 2018)	Assay 2 (19 Apr 2021)	Assay 3 (28 Apr 2021)	Assay 4 (12 May 2021)	Assay 5 (19 May 2021)	Assay 6 (21 Jun 2021)
CONTROL	1	2	2	2	2	2
Ls 2362 (0.0106 g)	1				2	
Ls IAB59 (0.0023 g)				4		
Bt <i>israelensis</i> 4Q2 (0.0234 g)	1					
Bt <i>higo</i> T44001 (0.033 g)	1	2				
Bt <i>kurstaki</i> HD1 (0.044 g)		2				
Bt GP 138 (0.072 g)			2			
Bt <i>konkukian</i> 97–27 (0.0156 g)			2			
Geranium EONP (35 µg/cm ²)			2		2	
Peppermint EONP (70 µg/ cm ²)						3

Table 2

Average Size (AS), in nanometers, polydispersity (PDI), and Encapsulation Efficiency (EE) of geranium and peppermint EONP, after 7 Days Post-formulation. Figures represent mean value ± standard error.

	AS	PDI	EE (%)
Geranium EONP	259 ± 12 a	0.228 ± 0.007 a	90.5 ± 2.32 a
Peppermint EONP	381 ± 29 b	0.532 ± 0.013 b	72.25 ± 1.6 b

^a Different letters within the same row indicate statistical differences (LSD; p < 0.05).

evaluated at 35 µg cm⁻² and 70 µg cm⁻², respectively. The decision to use the concentration of peppermint for this nanoparticle was based on research by Jesser et al. (2020a), which indicated that the bioactivity of geranium EONP was higher than that of peppermint EONP against *Plodia interpunctella* (Lepidoptera: Pyralidae). The nanoparticles were dispersed in the central circle at the bottom base of glass Petri dish.

2.7. Statistical analysis

Survival of mites subjected to the different treatments was analysed using the non-parametric Kaplan-Meier estimate via survival curves (Kaplan and Meier, 1958). The Log-rank test (Kleinbaum and Klein, 2012) with Bonferroni correction allowed for multiple pairwise comparisons between the survival curves of each treatment in order to determine significant differences between them.

All statistical analyses were performed using R version 4.3.1 (R Core Team, 2023). We used the survfit() function to conduct the Kaplan-Meier estimations with the survival package (Therneau, 2023). The survival curves were drawn using ggsurvplot() function of the survminer package (Kassambara et al., 2020). The log-rank test was carried out using survdiff() function of the survival package. Multiple pairwise comparisons were conducted with the pairwise_survdiff() function of the survminer package. The statistical significance level set in all statistical analyses was 0.05.

3. Results and discussion

First, the chemical analysis of EONP revealed that β-citronellol and geraniol were the predominant compounds in geranium EONP (Table 3). Moreover, components such as linalool, menthone, citronellyl formate, and geranyl formate, which, in the pre-formulation sample, were between 8% and 11%, had a significant reduction after formulation (<1.7%). Additionally, minor components present in the original sample (<3%) were undetectable after formulation. In contrast, menthol emerged as the primary compound in peppermint oil and its nanoparticles, as indicated in Table 3. After formulation, a slight decrease was noted in the concentrations of isomenthone, p-menthen-3-one, and menthol acetate. Furthermore, minor components present in the initial sample (<6%) were not detected after formulation.

Mites in control plates survived, on average, 27.6 h. Mean and median survival times are shown in Table 3. We must take into account that this time measurement started with mite extraction, but, on average, ibex death to laboratory mite extraction took around 10.9 ± 6.1 h. As expected, average survival of mites maintained at low temperature (12 °C) reached the highest values: 40.7 h, compared with that at 35°C: 31.2 hr.

Kaplan-Meier survival analysis is depicted in Figs. 2 and 3. The overall survival function can be seen in Fig. 2a. Log-rank test (Fig. 2b) showed statistically significant differences in the mean survival time for most of the treatments (χ² = 38309, p < 0.0001). In particular, *B. thuringiensis* serovar. *konkukian* 97–27 and geranium EONP reduced mite survival significantly. Conversely, mites treated with *B. thuringiensis* GP 138, *B. thuringiensis* *higo* T44001, Ls IAB59 and peppermint EONP survived more than those maintained in the control plates (Table 4; Fig. 3). The remaining treatments did not show

Table 3
Chemical analysis of pre / post-formulation of the oils from geranium and peppermint.

RT (min)	Compounds	Geranium EO		Peppermint EO	
		Preformulation	Postformulation	Preformulation	Postformulation
7.16	α - pinene	-	-	1.92	-
8.36	β - pinene	-	-	1.85	-
9.87	Limonene	-	-	3.36	-
9.93	1–8 cineol	-	-	5.88	-
13.06	Linalool	12.67	9.95	-	-
13.55	Isomenthone	-	-	16.90	6.95
13.85	Menthone	11.14	1.38	-	-
14.10	Menthol	-	-	52.51	81.37
14.35	p-menten-3-ona	-	-	10.43	7.57
16.14	β -citronellol	26.14	38.12	-	-
16.48	Geraniol	23.19	47.89	-	-
16.98	Citronellyl Formate	10.37	1.71	-	-
17.70	Geranyl Formate	7.94	0.95	-	-
18.04	Menthol acetate	-	-	7.15	4.11
20.85	Geranyl Acetate	2.01	-	-	-
20.86	Caryophyllene	2.58	-	-	-
23.70	Neryl Acetate	2.98	-	-	-




Fig. 1. A: Skin of the scabietic donor ibex. B: Several ibex skin pieces were placed into a painted glass Petri dish; note that the center of the plate remains transparent. C: the light applied to the bottom of the plate generated a temperature gradient into the plate. This gradient favoured mite migration from the skin to the centre of the plate. D: protective wear was needed for skin and plates manipulation.

significant differences in mite survival compared with controls.

4. Discussion

The *B. thuringiensis* serovar. *konkukian* and geranium EONP formulation gave promising results in our assays, and could be effective in reducing mite survival time. Different encapsulation strategies (for bacteria, Cry proteins and single spores) aimed to increase ingestion of *B. thuringiensis* Cry proteins, need to be tested. Moreover, host contact time and the effect of the temperature and UV radiation on the

persistence of *B. thuringiensis* in the field must be addressed before performing *in vivo* assays (de Oliveira et al., 2021). The MXPA patent 02008705 (Ramirez, 2004) is a nanoencapsulation technique of a mixture of *B. thuringiensis* Cry proteins and spores with high residual activity. On the other hand, Ureña-Saborío et al. (2017) performed chitosan/TPP nanoparticles containing bacterial metabolic infiltrates of the strain *B. thuringiensis* SER-217, and achieved their efficient release in an aqueous medium, together with increasing protection and stability of such compounds.

B. thuringiensis strain GP 138, which has previously shown activity




Fig. 2. Kaplan-Meier survival curves. Left graph (A) shows the overall survival curve without considering any treatments. Right graph (B) shows the survival curves by treatment, the p-value of the log-rank test and the pairwise multiple comparisons with Bonferroni correction. Significant differences were indicated by different lowercase letters ($p < 0.05$).




Fig. 3. Kaplan-Meier survival curves separated for better visualization.

against the tick *Rhipicephalus microplus* (Fernández-Ruvalcaba et al., 2010), did not show any reduction in *Sarcoptes scabiei* survival in this study. *B. thuringiensis konkukian* strain 97-27, however, did show activity. This is a genome-sequenced strain of *B. thuringiensis* (Han et al., 2006) that, in contrast to other *B. thuringiensis* strains tested, is not recorded as producing known delta endotoxins or invertebrate-active toxins produced during the vegetative stage of growth. The genome does encode toxins with reported roles in mammalian food poisoning such as CytK, and the tripartite toxins Hbl and Nhe (in common with

several strains of *B. thuringiensis*). The activity of these proteins against invertebrates has not been reported and it is possible that these or other, as yet uncharacterised, proteins or small molecule toxins are responsible for its activity against *S. scabiei* in this study. This possibility warrants further investigation to identify the agent responsible for the activity observed.

This study used geranium and peppermint essential oils (EO) to formulate PEG-6000 nanoparticles, due to their bioactivity against various tick species (An and Tak, 2022; Awad et al., 2022; Klafke et al.,

Table 4

Parameters estimated by treatment via Kaplan-Meier analysis. From left to right: type of treatment, number of mites subjected to this treatment, mean survival time, standard error of the mean and median survival time. Mean and median are measured in hours; se: standard error.

Treatment	n	mean	se	median
Bt GP 138 (0.072 g)	8829	48.8	0.1339	47
Bt <i>higo</i> T44001 (0.033 g)	1147	28.8	0.2713	26.2
Bt <i>israelensis</i> 4Q2 (0.0234 g)	488	26.4	0.1388	26.2
Bt <i>konkukian</i> 97–27 (0.0156 g)	2001	14.8	0.0690	13.8
Bt <i>kurstaki</i> HD1 (0.044 g)	91	34.8	0.8064	29
CONTROL	11211	27.6	0.1077	26.2
Ls 2362 (0.0106 g)	2564	26.3	0.0504	26.2
Ls IAB59 (0.0023 g)	278	33	0.4005	29.8
Geranium EONP (35 µg/cm ²)	8603	18.1	0.0826	13.8
Peppermint EONP (70 µg/cm ²)	3744	40.5	0.1285	47

2021; Voronova et al., 2022). In the case of geranium oil, citronellol and geraniol are the important constituents that responsible for the acaricidal activity in *Rhipicephalus annulatus* (Ibrahium et al., 2022). Similarly, the bioactivity of peppermint EO, attributed to menthol and isomenthone, has been observed against *Tetranychus cinnabarinus* and *Tetranychus urticae* (Abd-Allah et al., 2022). Enan (2001) suggested that the toxicity of constituents of essential oils against insect pests might be related to the octopaminergic nervous system of insects, while De Oliveira et al. (1997) proposed that certain monoterpenes inhibit cytochrome P450-dependent monooxygenases. Moreover, Ryan and Byrne (1988) identified a connection between the toxicity of monoterpenes, their capacity to inhibit acetylcholinesterase (AChE), and their effectiveness against insects or ticks.

Regarding the bioactivity of nanoparticles, it was observed that geranium EONP exhibited greater efficacy compared to peppermint EONP. This result could be attributed to the physicochemical characteristics of the nanoparticles. Peppermint EONP had size of 390 nm and were polydisperse these values are higher than geranium EONP. It is well-known that nanoparticle size plays a crucial role in the penetration of bioactive compounds through the cuticle. Hashem et al. (2018) demonstrated that EO nanoformulations enhance cuticle penetration, allowing products to penetrate insects more easily. Furthermore, the nanoscale size of EONP could extend the exposure time of bioactive compounds to insect pests, covering larger areas of the insect cuticle. Additionally, nanoparticles can alter the delivery pattern of EO active ingredients, thereby enhancing their efficacy (Iavicoli et al., 2017). Moreover, the encapsulation efficiency (EE) of peppermint EONP was 72%, which is lower than that of geranium EONP. It will probably be necessary to use a higher amount of peppermint EONP for an effective pest management program.

In our study, temperature and relative humidity (RH) were maintained during the different assays. When off the host, *Sarcoptes* mites are unable to use water vapor actively from unsaturated ambient air (Arlan and Veselica, 1979) and, therefore, their survival time is strongly affected by ambient RH (Arlan et al., 1984). Mellanby et al. (1942) found that *S. scabiei* (obtained from human scrapings) did not move when temperature was below 15–16°C, but did so rapidly above 20°C; and heating at 50°C for 10 minutes was enough to exterminate the mite. At cooler temperatures (e.g., 4 °C) black bear-derived mites survived over a week (Niedringhaus et al., 2019). Moreover, at low temperatures, survival of *S. scabiei* increases with relative humidity (Davis and Moon, 1987; Arlan et al., 1989). Thus, environmental conditions (mainly temperature and relative humidity) will strongly affect mite survival when off the host and, therefore, its ability to be transmitted indirectly, to spread, to establish and to persist (Castro et al., 2017; Montecino-Latorre et al., 2019; Loredó et al., 2020; Browne et al., 2021).

5. Conclusion

In conclusion, the acaricidal activity of NP formulations of *B. thuringiensis konkukian* strain 97–27, of other *B. thuringiensis* strains and of other plant essential oils at different concentrations deserve to be studied in more detail, before considering *in vivo* assays. In a complementary way, the effect of both temperature and RH on the survival of ibex-derived mites (when off the host) need to be analysed in depth by maintaining mites *in vitro* at a wider range of such conditions.

Ethics approval

Procedures carried out in this work were approved by the regional government (Junta de Andalucía): Project 15/12/2018/163, and also by the Ethics Committee of the Jaén University.

Funding

Research activities of JEG and JMP are partially supported by the Junta de Andalucía Government (RNM-118 group of the PAIDI), and by the Jaén University (Action 1b). The AUIP (Asociación Universitaria Iberoamericana de Postgrado) scholarship facilitated EJ's mobility between the National University of the South (Argentina) and the Jaén University (Spain) to conduct this research.

CRediT authorship contribution statement

Jesús M. Pérez: Writing – review & editing, Writing – original draft, Supervision, Investigation, Funding acquisition, Conceptualization. **Jorge O Werdin:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Investigation, Conceptualization. **Emiliano J Jesser:** Writing – review & editing, Investigation, Data curation, Conceptualization. **Colin Berry:** Writing – review & editing, Writing – original draft, Supervision, Resources, Investigation, Conceptualization. **Raquel Crespo-Ginés:** Writing – review & editing, Writing – original draft, Investigation. **Mohamed A. Gebely:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Investigation, Conceptualization. **Antonio J López-Montoya:** Writing – review & editing, Writing – original draft, Formal analysis, Data curation. **José E Granados:** Writing – review & editing, Writing – original draft, Resources, Investigation.

Declaration of Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships.

Acknowledgements

Authors are indebted to Antonio Rodríguez, José López and Isidro Puga for their help in obtaining biological samples for our study.

References

- Abd-Allah, G.E., Habashy, M.G., Shalaby, M.M., 2022. Efficacy of mint derivatives, *Mentha spicata* L., against two species of *Tetranychus* spp. (Acari: Tetranychidae) and the predator, *Neoseiulus* sp. Egypt. Acad. J. Biol. Sci. A Entomol. 15, 63–70. <https://doi.org/10.21608/EAJBSA.2022.224349>.
- An, H., Tak, J.H., 2022. Miticidal and repellent activity of thirty essential oils and their synergistic interaction with vanillin against *Tetranychus urticae* Koch (Acari: Tetranychidae). Ind. Crops Prod. 182, 114872 <https://doi.org/10.1016/j.indcrop.2022.114872>.
- Andrews, J.R., 1981. The extraction of *Sarcoptes scabiei* from mammalian hosts. J. Parasitol. 67, 753–754.
- Andriantsoanirina, V., Izri, A., Botterel, F., Foulet, F., Chosidow, O., Durand, R., 2014. Molecular survey of knockdown resistance to pyrethroids in human scabies mites. Clin. Microbiol. Infect. 20, O139–O141. <https://doi.org/10.1111/1469-0691.12334>.
- Arlan, L.G., Morgan, M.S., 2017. A review of *Sarcoptes scabiei*: past, present and future. Parasites Vectors 10, 297. <https://doi.org/10.1186/s13071-017-2234-1>.

- Arlian, L.G., Runyan, R.A., Achar, S., Estes, S.A., 1984. Survival and infestivity of *Sarcoptes scabiei* var. *canis* and var. *hominis*. *J. Am. Acad. Dermatol.* 11, 210–215. [https://doi.org/10.1016/S0190-9622\(84\)70151-4](https://doi.org/10.1016/S0190-9622(84)70151-4).
- Arlian, L.G., Veselica, M.M., 1979. Water balance in insects and mites. *Comp. Biochem. Physiol.* 64, 191–200. [https://doi.org/10.1016/0300-9629\(79\)90650-9](https://doi.org/10.1016/0300-9629(79)90650-9).
- Arlian, L.G., Vyzenski-Moher, D.L., Pole, M.J., 1989. Survival of adults and development stages of *Sarcoptes scabiei* var. *canis* when off the host. *Exp. Appl. Acarol.* 6, 181–187. <https://doi.org/10.1007/BF01193978>.
- Athassiou, C.G., Kavallieratos, N.G., Benelli, G., Losic, D., Usha Rani, P., Desneux, N., 2018. Nanoparticles for pest control: current status and future perspectives. *J. Pest Sci.* 91, 1–15. <https://doi.org/10.1007/s10340-017-0898-0>.
- Awad, S.E., Salah, K.B.H., Jghel, M.M., Alkhaibari, A.M., Shami, A.A., Alghamdi, R.A., Awad, A.E., 2022. Chemical characterization of clove, basil and peppermint essential oils; evaluating their toxicity on the development stages of two-spotted spider mites grown on cucumber leaves. *Life* 12, 1751. <https://doi.org/10.3390/life12111751>.
- Berry, C., 2012. The bacterium, *Lysinibacillus sphaericus*, as an insect pathogen. *J. Invert. Pathol.* 109, 1–10. <https://doi.org/10.1016/j.jip.2011.11.008>.
- Bornstein, S., Mörner, T., Samuel, W.M., 2001. *Sarcoptes scabiei* and sarcoptic mange. In: Samuel, W.M., Pybus, M.J., Kocan, A.A. (Eds.), *Parasitic diseases of wild mammals*, 2nd edn. Iowa State University Press, Ames, pp. 107–119.
- Browne, E., Diressen, M.M., Ross, R., Roach, M., Carver, S., 2021. Environmental suitability of bare-nosed wombat burrows for *Sarcoptes scabiei*. *Int. J. Parasitol.* Parasites Wildl. 16, 37–47. <https://doi.org/10.1016/j.ijppaw.2021.08.003>.
- Campolo, O., Cherif, A., Ricupero, M., Siscaro, G., Grissa-Lebdi, K., Russo, A., Cucci, L. M., Di Pietro, P., Satriano, C., Desneux, N., Biondi, A., Zappalà, L., Palmeri, V., 2017. Citrus peel essential oil nanoformulations to control the tomato borer, *Tuta absoluta*: chemical properties and biological activity. *Sci. Rep.* 7, 13036 <https://doi.org/10.1038/s41598-017-13413-0>.
- Casas-Díaz, E., Marco, I., López-Olvera, J.R., Mentaberre, G., Lavín, S., 2011. Comparison of xylazine-ketamine and medetomidine-ketamine anaesthesia in the Iberian ibex (*Capra pyrenaica*). *Eur. J. Wildl. Res.* 57, 887–893. <https://doi.org/10.1007/s10344-011-0500-7>.
- Castro, I., de la Fuente, A., Fandos, P., Cano-Manuel, F.J., Granados, J.E., Sorieguero, R.C., Alasaad, S., Pérez, J.M., 2017. On the population biology of *Sarcoptes scabiei* infesting Iberian ibex (*Capra pyrenaica*). *Int. J. Acarol.* 42, 7–11. <https://doi.org/10.1080/01647954.2015.1109710>.
- Crickmore, N., Berry, C., Panneerselvam, S., Mishra, R., Connor, T.R., Bonning, B.C., 2021. A structure-based nomenclature for *Bacillus thuringiensis* and other bacteria-derived pesticidal proteins. *J. Invertebr. Pathol.* 186, 107438 <https://doi.org/10.1016/j.jip.2020.107438>.
- Currie, B.J., Harumal, P., McKinnon, M., Walton, S.F., 2004. First documentation of in vivo and in vitro ivermectin resistance in *Sarcoptes scabiei*. *Clin. Infect. Dis.* 39, e8–e12. <https://doi.org/10.1086/421776>.
- Davis, D.P., Moon, R.D., 1987. Survival of *Sarcoptes scabiei* (De Geer) stored in three media at three temperatures. *J. Parasitol.* 73, 661–662.
- De Luca, I., Pedram, P., Moieni, A., Cerruti, P., Peluso, G., Di Salle, A., Germann, N., 2021. Nanotechnology development for formulating essential oils in wound dressing materials to promote the wound-healing process: a review. *Appl. Sci.* 11, 1713. <https://doi.org/10.3390/app11041713>.
- De Oliveira, A.C., Ribeiro-Pinto, L.F., Paumgarten, F.J., 1997. In vitro inhibition of CYP2B1 monooxygenase by β -myrcene and other monoterpene compounds. *Toxicol. Lett.* 92, 39–46. [https://doi.org/10.1016/S0378-4274\(97\)00034-9](https://doi.org/10.1016/S0378-4274(97)00034-9).
- de Oliveira, J.L., Campos, E.V.R., Bakshi, M., Abhilash, P.C., Fraceto, L.F., 2014. Application of nanotechnology for the encapsulation of botanical insecticides for sustainable agriculture: prospects and promises. *Biotechnol. Adv.* 32, 1550–1561. <https://doi.org/10.1016/j.biotechadv.2014.10.010>.
- de Oliveira, J.L., Fernandes Fraceto, L., Bravo, A., Polaczyk, R.A., 2021. Encapsulation strategies for *Bacillus thuringiensis*: from now to the future. *J. Agric. Food Chem.* 69, 4564–4577. <https://doi.org/10.1021/acs.jafc.0c07118>.
- Enan, E., 2001. Insecticidal activity of essential oils: octopaminergic sites of action. *Comp. Biochem. Physiol. C. Toxicol. Pharmacol.* 130, 325–337. [https://doi.org/10.1016/S1532-0456\(01\)00255-1](https://doi.org/10.1016/S1532-0456(01)00255-1).
- Espinosa, J., Ráez-Bravo, A., López-Olvera, J.R., Pérez, J.M., Lavín, S., Tvarijonavičute, A., Cano-Manuel, F.J., Fandos, P., Sorieguero, R.C., Granados, J.E., Romero, D., Velarde, R., 2017. Histopathology, microbiology and the inflammatory process associated with *Sarcoptes scabiei* infection in the Iberian ibex, *Capra pyrenaica*. *Parasites Vectors* 10, 596. <https://doi.org/10.1186/s13071-017-2542-5>.
- Fernández-Ruvalcaba, M., Peña-Chora, G., Romo-Martínez, A., Hernández-Velázquez, V., Bravo de la Parra, A., Pérez De La Rosa, D., 2010. Evaluation of *Bacillus thuringiensis* pathogenicity for a strain of the tick, *Rhipicephalus microplus*, resistant to chemical pesticides. *J. Insect Sci.* 10, 186. <https://doi.org/10.1673/031.010.14146>.
- Han, C.S., Xie, G., Challacombe, J.F., Altherr, M.R., Bhotika, S.S., Bruce, D., Campbell, C. S., Campbell, M.L., Chen, J., Chertkov, O., Cleland, C., Dimitrijevic, M., Doggett, N. A., Fawcett, J.J., Glavina, T., Goodwin, L.A., Hill, K.K., Hitchcock, P., Jackson, P.J., Keim, P., Kewalramani, A.R., Longmire, J., Malfatti, L.S., McMurry, K., Meincke, L.J., Misra, M., Moseman, B.L., Mundt, M., Munk, C., Okinaka, R.T., Parson-Quintana, B., Reilly, L.P., Richardson, P., Robinson, D.L., Rubin, E., Saunders, E., Tapia, R., Tesmer, J.G., Thayer, N., Thompson, L.S., Tice, H., Ticknor, L.O., Wills, P.L., Brettin, T.S., Gilna, P., 2006. Pathogenomic sequence analysis of *Bacillus cereus* and *Bacillus thuringiensis* isolates closely related to *Bacillus anthracis*. *J. Bacteriol.* 188, 3382–3390. <https://doi.org/10.1128/JB.188.9.3382-3390.2006>.
- Hashem, A.S., Awadalla, S.S., Zayed, G.M., Maggi, F., Benelli, G., 2018. *Pimpinella anisum* essential oil nanoemulsions against *Tribolium castaneum* - insecticidal activity and mode of action. *Environ. Sci. Pollut. Res.* 25, 18802–18812. <https://doi.org/10.1007/s11356-018-2068-1>.
- Hill, C.A., Pinnock, D.E., 1998. Histopathological effects of *Bacillus thuringiensis* on the alimentary canal of the sheep louse, *Bovicola ovis*. *J. Invert. Pathol.* 72, 9–20. <https://doi.org/10.1006/jipa.1998.4761>.
- Höfte, H., Whiteley, H.R., 1989. Insecticidal crystal proteins of *Bacillus thuringiensis*. *Microbiol. Rev.* 53, 242–255. <https://doi.org/10.1128/mr.53.2.242-255.1989>.
- Iavicoli, L., Leso, V., Beezhold, D.H., Shvedova, A.A., 2017. Nanotechnology in agriculture: opportunities, toxicological implications, and occupational risks. *Toxicol. Appl. Pharmacol.* 329, 96–111. <https://doi.org/10.1016/j.taap.2017.05.025>.
- Ibrahiam, S.M., Aboelhadid, S.M., Wahba, A.A., Farghali, A.A., Miller, R.J., Abdel-Baki, A.A.S., Al-Quraishy, S., 2022. Preparation of geranium oil formulations effective for control of phenotypic resistant cattle tick *Rhipicephalus annulatus*. *Sci. Rep.* 12, 11693 <https://doi.org/10.1038/s41598-022-14661-5>.
- Jesser, E., Yeguerman, C., Stefanazzi, N., Gomez, R., Murray, A.P., Ferrero, A.A., Werdin-Gonzalez, J.O., 2020a. Ecofriendly approach for the control of a common insect pest in the food industry, combining polymeric nanoparticles and post-application temperature. *J. Agric. Food Chem.* 68, 5951–5958. <https://doi.org/10.1021/acs.jafc.9b06604>.
- Jesser, E., Yeguerman, C., Gili, V., Santillan, G., Murray, A.P., Domini, C., Werdin-González, J.O., 2020b. Optimization and characterization of essential oil nanoemulsions using ultrasound for new ecofriendly insecticides. *ACS Sustain. Chem. Eng.* 8, 7981–7992. <https://doi.org/10.1021/acscuschemeng.0c02224>.
- Jones, G.W., Wirth, M.C., Monnerat, R.G., Berry, C., 2008. The Cry48Aa-Cry49Aa binary toxin from *Bacillus sphaericus* exhibits highly restricted target specificity. *Environ. Microbiol.* 10, 2418–2424. <https://doi.org/10.1111/j.1462-2920.2008.01667.x>.
- Kaplan, E.L., Meier, P., 1958. Nonparametric estimation from incomplete observations. *J. Am. Stat. Assoc.* 53, 457–481. <https://doi.org/10.2307/2281868>.
- Kassambara, A., Kosinski, M., Biecek, P., 2020. *Survminer: drawing survival curves using 'ggplot2'*. R package version 0.4.8. (<https://CRAN.R-project.org/package=survminer>).
- Klafke, G.M., Thomas, D.B., Miller, R.J., de León, A.A.P., 2021. Efficacy of a water-based botanical acaricide formulation applied in portable spray box against the southern cattle tick, *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae), infesting cattle. *Ticks Tick. Borne Dis.* 12, 101721 <https://doi.org/10.1016/j.ttbdis.2021.101721>.
- Kleinbaum, D.G., Klein, M., 2012. *Kaplan-Meier survival curves and the Log-Rank test*. In: *Survival Analysis. Statistics for Biology and Health*. Springer, New York. (https://doi.org/10.1007/978-1-4419-6646-9_2).
- Knowles, B.H., Dow, J.A.T., 1993. The crystal δ -endotoxins of *Bacillus thuringiensis*: Models for their mechanism of action in the insect gut. *BioEssays* 15, 469–476. <https://doi.org/10.1016/j.toxicon.2006.11.022>.
- Kumar, A., Singh, P., Gupta, V., Prakash, B., 2020. Application of nanotechnology to boost the functional and preservative properties of essential oils. In: Prakash, B. (Ed.), *Functional and preservative properties of phytochemicals*. Academic Press, London, pp. 241–267. <https://doi.org/10.1016/B978-0-12-818593-3.00008-7>.
- Loredo, A.I., Rudd, J.L., Foley, J.E., Clifford, D.L., Cypher, B.L., 2020. Climatic suitability of San Joaquin kit fox (*Vulpes macrotis macrotis*) dens for sarcoptic mange (*Sarcoptes scabiei*) transmission. *J. Wildl. Dis.* 56, 126–133. <https://doi.org/10.7589/2019-02-035>.
- Mellanby, K., Johnson, C.G., Bartley, W.C., Brown, P., 1942. Experiments on the survival and behaviour of the itch mite, *Sarcoptes scabiei* DeG. var. *hominis*. *Bull. Ent. Res.* 33, 267–271. <https://doi.org/10.1017/S0007485300026584>.
- Monnerat, R., Cardoso Batista, A., Telles de Medeiros, P., Soares Martins, É., Melatti, V. M., Botelho Praça, L., Fiúza Dumas, V., Morinaga, C., Demo, C., Menezes Gomes, A. C., Falcão, R., Brod Siqueira, C., Oliveira Silva-Werneck, J., Berry, C., 2007. Screening of Brazilian *Bacillus thuringiensis* isolates active against *Spodoptera frugiperda*, *Plutella xylostella* and *Anticarsia gemmatilis*. *Biol. Control* 41, 291–295. <https://doi.org/10.1016/j.biocontrol.2006.11.008>.
- Montecino-Latorre, D., Cypher, B.L., Rudd, J.L., Clifford, D.L., Mazet, J.A.K., Foley, J.E., 2019. Assessing the role of dens in the spread, establishment and persistence of sarcoptic mange in an endangered canid. *Epidemics* 27, 28–40. <https://doi.org/10.1016/j.epidem.2019.01.001>.
- Moroni, B., Valdeperes, M., Serrano, E., López-Olvera, J.R., Lavín, S., Rossi, L., 2020. Comment on: “The treatment of sarcoptic mange in wildlife: a systematic review. *Parasites Vectors* 13, 471. <https://doi.org/10.1186/s13071-020-04347-0>.
- Mounsey, K.E., Pasay, C.J., Arlian, L.G., Morgan, M.S., Holt, D.C., Currie, B.J., Walton, S. F., McCrath, J.S., 2010. Increased transcription of glutathione S-transferases in acaricide exposed scabies mites. *Parasites Vectors* 3, 43. <https://doi.org/10.1186/1756-3305-3-43>.
- Niedringhaus, K.D., Brown, J.D., Ternent, M.A., Peltier, S.K., Yabsley, M.J., 2019. Effects of temperature on the survival of *Sarcoptes scabiei* of black bear (*Ursus americanus*) origin. *Parasitol. Res.* 118, 2767–2772. <https://doi.org/10.1007/s00436-019-06387-7>.
- Pascoli, M., Lopes-Oliveira, P.J., Fraceto, L.F., Seabra, A.B., Oliveira, H.C., 2018. State of the art of polymeric nanoparticles as carrier systems with agricultural applications: a minireview. *Emerg. Ecol. Environ.* 3, 137–148. <https://doi.org/10.1007/s40974-018-0090-2>.
- Pence, D.B., Ueckermann, E., 2002. Sarcoptic mange in wildlife. *Rev. Sci. Tech. - Int. Epiz* 21, 385–398.
- R Core Team, 2023. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. (<https://www.r-project.org/>).
- Ramirez, L.M., 2004. Technology for encapsulating delta endotoxins of *Bacillus thuringiensis* of the israelensis variety for extending the activity thereof on mosquitoes larvae. *MXPA Patent* 02008705 A, Dec 6, 2004.

- Rowe, M.L., Whiteley, P.L., Carver, S., 2019. The treatment of sarcoptic mange in wildlife: a systematic review. *Parasites Vectors* 12, 99. <https://doi.org/10.1186/s13071-019-3340-z>.
- Ryan, M.F., Byrne, O., 1988. Plant-insect coevolution and inhibition of acetylcholinesterase. *J. Chem. Ecol.* 14, 1965–1975. <https://doi.org/10.1007/BF01013489>.
- Therneau, T., 2023. A package for survival analysis in R. R package version 3.5-5, (<https://CRAN.R-project.org/package=survival>).
- Ureña-Saborío, H., Madrigal-Carballo, S., Sandoval, J., Vega-Baudrit, J.R., Rodríguez-Morales, A., 2017. Encapsulation of bacterial metabolic infiltrates isolated from different *Bacillus* strains in chitosan nanoparticles as potential green chemistry-based biocontrol agents against *Radopholus similis*. *J. Renew. Mater.* 5, 290–299. <https://doi.org/10.7569/JRM.2017.634119>.
- Voronova, N., Horban, V., Bohatkina, V., 2022. The effectiveness of acaricidal drugs based on herbal raw material. *Ecol. Quest.* 33, 55–71. <https://doi.org/10.12775/EQ.2022.003>.
- Walton, S.F., McKinnon, M., Pizzutto, S., Dougall, A., Williams, E., Currie, B.J., 2004. Acaricidal activity of *Melaleuca alternifolia* (tea tree) oil. In vitro sensitivity of *Sarcoptes scabiei* var *hominis* to terpinen-4-ol. *Arch. Dermatol.* 140, 563–566. <https://doi.org/10.1001/archderm.140.5.563>.
- Walton, S.F., Myerscough, M.R., Currie, B.J., 2000. Studies in vitro on the relative efficacy of current acaricides for *Sarcoptes scabiei* var. *hominis*. *Trans. R. Soc. Trop. Med. Hyg.* 94, 92–96. [https://doi.org/10.1016/s0035-9203\(00\)90454-1](https://doi.org/10.1016/s0035-9203(00)90454-1).
- Werdin-González, J.O., Jesser, E.N., Yeguerman, C.A., Ferrero, A.A., Fernández Band, B., 2017. Polymer nanoparticles containing essential oils: New options for mosquito control. *Environ. Sci. Pollut. Res.* 24, 17006–17015. <https://doi.org/10.1007/s11356-017-9327-4>.
- Werdin-González, J.O., Gutiérrez, M.M., Ferrero, A.A., Fernández Band, B., 2014. Essential oils nanoformulations for stored product pest control: Characterization and biological properties. *Chemosphere* 100, 130–138. <https://doi.org/10.1016/j.chemosphere.2013.11.056>.
- Yeguerman, C.A., Urrutia, R.I., Jesser, E.N., Massiris, M., Delrieux, C.A., Murray, A.P., Werdin-González, J.O., 2022. Essential oils loaded on polymeric nanoparticles: bioefficacy against economic and medical insect pests and risk evaluation on terrestrial and aquatic non-target organisms. *Environ. Sci. Pollut. Res.* 29, 71412–71426. <https://doi.org/10.1007/s11356-022-20848-0>.