

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/169101/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Ólafsdóttir Flóvenz, Sigrún, M. Salkovskis, Paul, Gregory, James D., Valdimarsdóttir, Heiðdís, Sigurðsson, Engilbert and Friðrik Sigurðsson, Jón 2024. Persistent physical symptoms and mental health problems among individuals seeking work rehabilitation. *Nordic Psychology* 76 (4) , pp. 534-547.
10.1080/19012276.2024.2342345

Publishers page: <https://doi.org/10.1080/19012276.2024.2342345>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Persistent Physical Symptoms and Mental Health Problems among Individuals seeking Work Rehabilitation

Sigrún Ólafsdóttir Flóvenz^{a*}, Paul M. Salkovskis^b, James D. Gregory^c, Heiðdís
Valdimarsdóttir^{a,d}, Engilbert Sigurðsson^{e,f} and Jón Friðrik Sigurðsson^{a,f},

*^a Department of Psychology, Reykjavik University, Iceland; ^bDepartment of Experimental
Psychology, University of Oxford, England; ^cDepartment of Psychology, University of Bath,
England; ^dMount Sinai School of Medicine, United States of America; ^eLandspítali-The
National University Hospital of Iceland, Iceland; ^fUniversity of Iceland*

Correspondence concerning this article should be addressed to Sigrún Flóvenz, Reykjavik
University, Department of Psychology, Menntavegur 1, 101 Reykjavik Iceland. Tel.: +354-
5996200. E-mail address: sigruno@gmail.com.

Persistent Physical Symptoms and Mental Health Problems among Individuals seeking Work Rehabilitation

Persistent Physical Symptoms (PPS) are symptoms without a known biological cause. They are common and associated with psychological distress, work disability and unemployment. This study aimed to examine the prevalence of PPS and their association with clinical levels of general anxiety, depression, social anxiety, health anxiety or post traumatic symptoms in a work rehabilitation setting and to identify individuals within that setting that might benefit from psychological treatment for PPS. A sample of 324 individuals seeking work rehabilitation filled out questionnaires measuring PPS and mental health problems. To estimate how many service seekers might benefit from psychological treatment for PPS we identified participants who reported both PPS and psychological distress, defined as an elevated score on measures of general anxiety, depression or health anxiety. Eighty point nine percent of participants reported having one or more PPS and 61.7% reported having two or more. Those with PPS were more likely to report clinical levels of depression, anxiety and health anxiety than those without and the number of PPS was positively correlated with symptoms of depression, anxiety and health anxiety. Almost 65% of the participants were identified as having PPS, being psychologically distressed and willing to accept psychological treatment. There is a clear need for specialized treatment for PPS within vocational rehabilitation in Iceland as service seekers commonly have PPS, are psychologically distressed and are willing to accept psychological treatment. Such treatment should preferably be transdiagnostic as the prevalence of multiple PPS is high in this sample.

Keywords: persistent physical symptoms, PPS, work rehabilitation, mental health, medically unexplained symptoms.

Introduction

Persistent Physical Symptoms (PPS) are physical symptoms that cannot be explained by any known biological or physical pathology despite thorough examination. PPS can range in number, duration and severity, from a single mild symptom to multiple chronic symptoms that are severe and disabling. Such symptoms are extremely common across medical and non-medical settings. Many different types of PPS have been described and further defined as syndromes, e.g., irritable bowel syndrome, chronic fatigue, fibromyalgia, non-cardiac chest pain, premenstrual syndrome and chronic pain.

The definition and terminology of PPS is controversial. People commonly meet criteria for more than one PPS condition and there are considerable similarities and overlap between different conditions in case definition, reported symptoms and in non-symptom associations. This has led some researchers to conclude that PPS is better described as one syndrome than many (Dimsdale et al., 2011; Wessely et al., 1999; Wessely & White, 2004). This lack of agreement has led to a lack of consistency in definitions and terminology used for PPS. *Persistent physical symptoms* is a new overarching term which seems to be more acceptable to patients than older terms, such as *Medically unexplained symptoms* or *Functional symptoms* (Marks & Hunter, 2015; Picariello et al., 2015), and is more in accordance with the latest changes made to the Diagnostic and Statistical Manual of Mental Disorders (DSM-5)(American Psychiatric Association, 2013).

PPS are very commonly presented in all healthcare settings. Research indicates that about one third (Roca et al., 2009) of all patients seeking consultation in primary health care have one or more PPS and that PPS comprise between 15% and 30% of all primary care

consultations (Kirmayer et al., 2004). It has been estimated that the prevalence of PPS (Carson et al., 2000; Nimnuan et al., 2001) in secondary care is between one third and two thirds of patients and at about one fifth of patients who frequently attend secondary care specialities (Reid et al., 2001).

PPS are associated with high rates of disability (den Boeft et al., 2016; Harris et al., 2009; Hoedeman et al., 2010; Rask et al., 2017) and psychological distress (Budtz-Lilly et al., 2015; Harris et al., 2009) and many tend to follow a chronic course with high levels of psychiatric comorbidity, although these symptoms can also be present in the absence of a mental disorder (Harris et al., 2009; Löwe et al., 2008; Roca et al., 2009). In addition, PPS carry with them a considerable social cost in the form of: 1) higher health care use (Barsky AJ et al., 2005; Budtz-Lilly et al., 2015; Harris et al., 2009) that tends to increase with number of PPS and symptom severity (Katon et al., 1991) and there is evidence that inappropriate information, over-investigation and over-treatment are common in the management of these patients (Kouyanou et al., 1998; McGorm et al., 2010); 2) increased health care cost (Barsky AJ et al., 2005; Konnopka et al., 2013; Rask et al., 2017; Reid et al., 2001) that is twice as high for those with multiple PPS than for those without (Barsky AJ et al., 2005) and those with well-defined medical symptoms (Rask et al., 2017); 3) increased work disability (den Boeft et al., 2016; Loenggaard et al., 2015; Rask et al., 2017), and 4) increased absence from work due to sick leaves (den Boeft et al., 2016; Harris et al., 2009; Hoedeman et al., 2010; Loenggaard et al., 2015). Not surprisingly PPS are associated with high unemployment rates (Amland et al., 2014; Momsen et al., 2016), an increased risk for becoming unemployed (Hoedeman et al., 2010; Loenggaard et al., 2015) and increased risk of permanent disability

pension even after adjustment for both psychiatric and chronic medical comorbidity (Rask et al., 2017).

The strong association between PPS, work disability, unemployment, and risk for permanent disability pension provides a clear rationale for taking a closer look at PPS in the context of work rehabilitation. This is particularly important as effective treatments are available for many different PPS conditions (Castell et al., 2011; Craske et al., 2011; Dear et al., 2013; Esler et al., 2003; Glombiewski et al., 2010; Ho et al., 2020; Jonsbu et al., 2011; Karlsson et al., 2015; Kim et al., 2020; Kisely et al., 2015; Kleinstäuber et al., 2011; Larun et al., 2017; Ljótsson et al., 2010, 2014; Pike et al., 2016; Tang et al., 2012; Thakral et al., 2020). As little is known about PPS in relation to work rehabilitation, we conducted a study at VIRK – Vocational Rehabilitation Fund in Iceland (<https://www.virk.is/is/english/mission-and-activities-of-virk>), where individuals with employment difficulties linked to ill health or injury are referred. VIRK is a social enterprise of which all the major labour unions and both public and private employers in the labour market are members. VIRK's services are free of charge and all those who have not been able to work because of infirmity, have a doctor's request for service and plan to resume to work or increase their employment, have the right to the fund's service (VIRK Vocational Rehabilitation Fund, 2013, 2014) . The fund operates according to Icelandic law which main objective is to ensure that individuals who have reduced workability following illness or accidents receive proper occupational rehabilitation (VIRK Vocational Rehabilitation Fund, 2013, 2014). The primary goal was to estimate the prevalence of PPS in this particular population and whether there is an association between the presence of PPS and mental health diagnoses as indicated by caseness on self-report scales. Secondly, we wanted to know whether different PPS subtypes differ in relation to the above-mentioned factors.

Thirdly, we wanted to see whether we could identify service seekers at VIRK who might benefit from psychological treatment for PPS.

Method

Participants

The sample consisted of people from all sectors of the labour market who sought work rehabilitation through VIRK, in the capital area of Iceland between June 2017 and May 2018. In Iceland almost everyone in the labour market is a member of a trade union and the participants in this study belonged to unions representing: store and office workers, university graduates, state and municipal employees, manual labourers or trade workers. Service users were not eligible for participation if they were under the age of 18, had started receiving rehabilitation from VIRK or had limitations in their language or reading ability that prevented them from filling out questionnaires. Three hundred eighty-three service seekers were invited to participate in the study of which 324 (84.6%) agreed and 59 (15.4%) declined. Limitations with language and reading abilities were the most common reason for exclusion but a few cases were excluded because the service users started receiving rehabilitation before the study questionnaires could be administered. Not all eligible service seekers were invited to participate in the study. The most common reasons for missed recruitment opportunities were that the counsellors forgot to present the study to their clients or they were never instructed to do so by their supervisors. In addition to that, counsellors within all unions occasionally deemed it to be too invasive for clients to participate, as they were either exhausted or emotionally upset during the counselling session.

Procedure

VIRK's counsellors presented the study to eligible clients during one of their first sessions. They asked for their informed consent and gave a battery of self-report questionnaires to those agreeing to participate. Participants filled out the questionnaires after the counselling session and put them in a sealed mailbox upon completion. Information about participants was retrieved from VIRK's database and matched with the questionnaire data using research identification numbers. This study was approved by the National Bioethics Committee of Iceland (application no. VSN-16-045).

Measures

Background Information

Information on participants' gender, age, marital status and their highest level of education was retrieved from VIRK's database. The response options for 1) marital status were single, married or cohabiting with a partner, divorced or widowed, and for 2) highest education completed were primary school, secondary school, vocational education at secondary school level, other vocational education, university and other.

Psychological Measurements

The Persistent Physical Symptoms Checklist (PPSC). Is a self-report instrument that measures persistent problems with seven types of physical symptoms that clearly interfere with daily life, i.e., sleep problems, pain, chronic tiredness fatigue or muscle problems, gastrointestinal problems, heart and chest symptoms, dizziness and/or related problems and gynaecological problems. Respondents are asked whether they have had problems with each of the symptoms for more than six months (except one month for sleep problems), whether

their symptoms have a known cause, and then indicate on a 9-point scale to what extent each symptom interferes with their lives (0 = Not at all, 2 = Slightly, 4 = Definitely, 6 = Markedly, 8 = Very Severely). The criteria for a particular PPS type are met if the problem has been present for an extended period of time, does not have a medical explanation, and definitely interferes with daily life (≥ 4). Previous validation of the checklists in a small validation sample suggests adequate convergent validity (Flóvenz et al., n.d.). If occasional missing items on the PPSC prevented us from concluding whether the criteria for a particular PPS symptom was met it was counted as absent.

The Work and Social Adjustment Scale (WSAS) (Mundt et al., 2002) measures impaired functioning in five areas of everyday life: the ability to work, manage a home, form and maintain close relationships and engage in social or private leisure activities. The scale consists of five items measured on an 8-point scale with a total score that ranges from 0 to 40. Scores between 10 and 20 have been associated with significant functional impairment and scores over 20 a more severe pathology (Mundt et al., 2002). The scale has been shown to have good psychometric properties (Mundt et al., 2002; Zahra et al., 2014) and in our sample it had a Cronbach's alpha of 0.84.

The Short Health Anxiety Inventory (SHAI) (Alberts et al., 2013; Salkovskis, 2002). Is a 14-item self-report scale that measures health-related anxiety. Answers are rated on a 4-point scale, with a total score between 0 and 54. Each item consists of four statements and the participant is asked to choose the one that best describes his/her feelings, over the past six months. The instrument has good psychometric properties (Alberts et al., 2013) and total scores of 18 or more differentiate reliably between those that meet DSM-IV criteria for

hypochondriasis and those that do not. The Icelandic version also has good psychometric properties (Viðarsson, 2016) and in this study Cronbach's alpha was 0.89.

The Patient Health Questionnaire-9 (PHQ-9) (Kroenke et al., 2001). is a 9-item self-report scale rated on a 4-point scale that measures the participant's severity of depression over the last two weeks (Kroenke & Spitzer, 2002). The scale has cut-off points of 5, 10, 15 and 20, which are interpreted as mild, moderate, moderately severe and severe depression. Both the original and the Icelandic version of the PHQ-9 have good psychometric properties (Kroenke et al., 2001; Kroenke & Spitzer, 2002; Pálsdóttir, 2007) and Cronbach's alpha in our sample was 0.81.

The Generalized Anxiety Disorder-7 (GAD-7) (Spitzer et al., 2006) measures symptoms of general anxiety over the last two weeks. It consists of 7 items rated on a 4-point scale, with a total score that ranges from 0 to 21. The GAD-7 has cut-off points at 5, 10 and 15, which are interpreted as mild, moderate, and severe anxiety. Both the Icelandic and the original version of the GAD-7 have good psychometric properties (Ingólfssdóttir, 2014; Spitzer et al., 2006) and Cronbach's alpha in our sample was 0.88.

The Social Phobia Scale (SPS) and the Social Interaction Anxiety Scale (SIAS) (Mattick & Clarke, 1998) are two companion measures for assessing social anxiety. The SPS measures fear of being observed by others during everyday activities and the SIAS measures fear of engaging socially with others. Both scales consist of 20 items, rated on a 5-point scale with a total score that ranges from 0 to 80. The original and the Icelandic versions of both scales have good psychometric properties (Ólafsdóttir, 2012; Osman et al., 1998) and in our sample SIAS had a Cronbach's alpha of 0.94 and the SPS of 0.96. In this study we considered

social anxiety scores to be in the clinical range if the combined total score of SIAS and SPS met a cut-off of 74 as this scoring method has been shown to best discriminates between those with and without social anxiety disorder in an Icelandic sample (Ólafsdóttir, 2012).

Traumatic experiences. Participants were asked about past traumatic events with the following single, yes or no, question taken from the Icelandic version of MINI International Neuropsychiatric Interview version 5.0 (Lecrubier et al., 1997; Sheehan & Lecrubier, 1998): *Have you ever experienced or witnessed or had to deal with an extremely traumatic event that included actual or threatened death or serious injury to you or someone else?*

The Impact of Event Scale Revised (IES-R) (Beck et al., 2008; Sundin & Horowitz, 2002) measures symptoms of post-traumatic stress. The scale consists of 22 items measured on a 5-point scale with a total score that ranges from 0 to 88. A score of 33 or higher has been shown to discriminate between people with and without Post Traumatic Stress Disorder (PTSD) (Creamer et al., 2003). The IES-R has good psychometric properties (Creamer et al., 2003) and Cronbach's alpha was 0.95 in our sample.

Alcohol Use Disorders Identification Test (AUDIT) (Saunders et al., 1993) measures hazardous or harmful alcohol consumption. The scale consists of 10 items, scored from 0 to 4. Total scores range from 0 to 40 and total scores of 8 or more have been shown to indicate hazardous or harmful alcohol use (Saunders et al., 1993). The AUDIT has been shown to have good psychometric properties (Reinert & Allen, 2007) and in our sample it had a Cronbach's alpha of 0.86.

Drug Use Disorders Identification Test (DUDIT) (Berman et al., 2005) is an 11 item scale that is used to screen for substance-related problems and is intended to be used with the

AUDIT. Total scores range from 0 to 44. According to a Swedish study a total score of 2 for women and 6 for men could be used as a cut-off score for harmful substance use in a general population sample (Berman et al., 2005) whereas an American study indicates that a total score of 8 could discriminate between patients with drug related problems and only alcohol related problems (Voluse et al., 2012) In the current study DUDIT scores were considered to indicate problematic substance use if they met a cut-off score of 6. The scale has been shown to have good psychometric properties (Berman et al., 2005; Voluse et al., 2012) and in our sample it had a Cronbach's alpha of 0.93.

Data Analysis

First, we examined the prevalence of PPS in our sample; χ^2 tests were used to determine whether there were significant associations between having PPS and clinical levels of general anxiety, depression, social anxiety, health anxiety or post traumatic symptoms. This was followed by correlating the number of PPS with continuous measures of those psychological variables found to be related to the presence of PPS. Next it was intended to examine differences between PPS subtypes and the measures of psychological distress. Finally, in order to estimate how many service seekers at VIRK might benefit from psychological treatment for PPS, we identified participants who both reported PPS and psychological distress, defined as an elevated score on the PHQ-9, the GAD-7 or the SHAI.

Results

Prevalence and Descriptives

The majority of participants (262, 80.9%) reported having one or more PPS on the PPSC, while 62 (19.1%) reported no such symptoms. Table 1 shows the numbers and percentages of

participants, with and without PPS, with respect to gender, educational level and relationship status.

[Table 1 near here]

More than three quarters of our participants were female and the gender distribution was similar in both groups. Participants that met criteria for at least one PPS were more likely to be separated than those who did not meet these criteria. They were also younger on average ($M=38.40$, $SD= 11.63$) than participants that did not ($M=43.15$, $SD= 13.16$), $t(305)=2.780$, $p=0.006$, $r= 0.16$. Based on average WSAS scores, participants with PPS were more functionally impaired ($M= 25.73$, $SD= 7.86$) than those without ($M= 16.46$, $SD= 9.70$), $t(76.937)=-6.832$, $p < 0.01$, $r= 0.61$. Seventy-eight percent of participants with PPS, and 35.6% of participants without had WSAS scores in the clinical range (>20), $\chi^2(1) = 40.802$, $p<0.01$. The odds of scoring above cut-off were 6.41 times higher for participants with PPS than without. Based on participants' scores on the AUDIT and the DUDIT, 24.0% ($n=76$) of the sample are likely to have or have had alcohol- or drug-related problems during the past year. Of these, 15.8% ($n=12$) reported that they were abstinent. There was no significant difference in the prevalence of such problems, $\chi^2(1) = 0.043$, $p=0.835$, nor in average scale scores between those with and without PPS, $t_{AUDIT}(317)=0.01$, $p = 0.992$ and $t_{DUDIT}(316)= -0.121$, $p =0.904$.

PPS and Mental Health

Table 2 shows the numbers and percentages of participants with and without PPS who score above cut-off on measures of health anxiety, depression, general anxiety, social anxiety and symptoms of post-traumatic stress.

[Table 2 near here]

As can be seen in Table 2, participants with one or more PPS were more likely to report symptoms of depression, anxiety or health anxiety in the clinical range. They were not more likely to report symptoms of social anxiety and post-traumatic stress disorder in the clinical range nor were they more likely to have experienced a traumatic event in their past.

Number and Types of Persistent Physical Symptoms.

Figure 1 shows the percentage of participants who met criteria for a number of the seven different PPS types. Sixty-one point seven percent of participants reported having two or more PPS.

[Figure 1 near here]

The number of PPS a participant reported was significantly correlated with depressive symptoms, using Pearson's r . $r_{\text{PHQ-9}} = 0.52$, $p < 0.001$, general anxiety symptoms $r_{\text{GAD-7}} = 0.38$, $p < 0.001$, health anxiety symptoms, $r_{\text{SHAI}} = 0.33$, $p < 0.001$. It was not possible to look at the association between the particular types of PPS and psychological factors because too few participants met criteria only for a single PPS type.

[Figure 2 near here]

Figure 2 shows the prevalence of the seven different PPS types. The most common PPS are fatigue and muscle problems, followed by sleep problems and pain. Neurological, gastrointestinal and heart and chest related symptoms are also fairly common, but much less

than the other three. Out of all possible combinations of symptoms, the five most common combinations account for 54.2% of all PPS cases. These are: 1) sleep problems, pain and fatigue and muscle problems accounting for 17.9% of all cases; 2) sleep problems and fatigue and muscle problems, accounting for 9.9% of all cases; 3) fatigue and muscle problems, accounting for 9.9% of all cases; 4) sleep problems, accounting for 9.2% of all cases; and 5) sleep problems, pain, fatigue and muscle problems and neurological symptoms, accounting for 7.3% of all cases.

VIRK Service Seekers who might benefit from Psychological Treatment for PPS

The number and percentages of participants with PPS and elevated scores on the PHQ-9, the GAD-7 (≥ 10 and ≥ 15) or the SHAI (≥ 18) and an elevated score on any of the three scales are presented in Table 3.

[Table 2 near here]

As can be seen in Table 3, a high percentage of those who report PPS have high levels of depression, anxiety and health anxiety symptoms. Crucially, 91.7% of participants with PPS reported that they believed it would be helpful to discuss their PPS and how their PPS affect them with a trained professional and would accept such help if offered. If we use the conventional cut-off score of 10 for the PHQ-9 and the GAD-7, 64.6% of the participants, VIRK service seekers would be identified as having PPS, being psychologically distressed and willing to accept psychological help for their symptoms. If we use the higher cut-off score of 15, 46.2% of the total sample would meet these criteria.

Discussion

In this study, a sample of people attending work rehabilitation were evaluated for prevalence and correlates of PPS. The prevalence was unexpectedly high at 80.9%, the modal number of PPS was three and the presence and number of PPS were associated with depression, anxiety and health anxiety. In addition, most participants with PPS and high levels of psychological distress reported being willing to accept psychological treatment for their symptoms. This suggests there is a need for specialised treatment for PPS within vocational rehabilitation and the high prevalence of multiple PPS indicates that such a treatment should preferably be transdiagnostic to some degree (Salkovskis et al., 2016). It was not possible to look at the association between particular types of PPS and psychological factors as too few participants reported single PPS.

To our knowledge this is the first study of its kind in the context of work rehabilitation. In other settings where prevalence of PPS has been examined the rates found were substantially lower. Studies have indicated that about a third of primary care patients (Roca et al., 2009), between one third and two thirds of secondary care patients (Carson et al., 2000; Nimnuan et al., 2001) and about one fifth of patients who frequently attend secondary care specialities (Reid et al., 2001) have one or more PPS. In two studies, the prevalence of PPS within both English and Icelandic primary care was estimated using *the Persistent Physical Symptom Checklist* (PPSC). In these studies, 31.1% of English and 24.5% of Icelandic primary care patients met criteria for one or more PPS and in both samples the modal number was one ((Flóvenz et al., 2021; Gregory et al., n.d.).

PPS are correlated with common psychiatric disorders such as anxiety and depression (Budtz-Lilly et al., 2015; Löwe et al., 2008) and about 40-60% of primary care patients with PPS have a comorbid depression or anxiety disorder (Harris et al., 2009; Löwe et al., 2008). In our total sample, depression, anxiety and health anxiety were all extremely common, with 80.1% meeting caseness for depression, anxiety or health anxiety on self-report scales. The presence of PPS was nonetheless associated with depression, anxiety and health anxiety and 86.6% of people with PPS met caseness on one or more of those measures.

Our overall sample is broadly similar to the population of people seeking work rehabilitation in Iceland. Comparisons with reports from VIRK Vocational Rehabilitation Fund (VIRK Vocational Rehabilitation Fund, 2018) may show differences in terms of education. This is not surprising because the present study did not include participants who did not speak Icelandic or had insufficient reading skills.

A high number of first-time service seekers at VIRK were not recruited in the study. This happened, most likely, for both random and non-random reasons. The main non-random reason was that counsellors occasionally deemed it too invasive to ask their client to participate in the study because of their emotional state. This might have excluded a small group of VIRK service seekers who possibly have particularly severe symptoms. Despite these limitations, we have a sample that largely represents the targeted population and the response rate was high (85%).

In this study, the prevalence of PPS was high and they were associated with having clinical levels of anxiety, depression and health anxiety. It is not clear which variables are primary and secondary, or whether there is a more complex interplay between them. Consequently, it is

possible that treating the PPS might resolve the psychological distress or it might be the case that depression and anxiety need to be targeted specifically in the treatment of PPS.

Our results are based on self-report including the estimation of whether a particular persistent physical symptom had a known medical cause. In the future, it would be preferable to confirm our results with more robust methods such as using standardized psychiatric diagnostic interviews and including a medical examination to confirm whether the physical symptoms have a medical explanation. It would also be interesting to know to what extent these problems predict work disability and how they influence the outcome and length as well as drop-out rate from the work rehabilitation.

Acknowledgements

Funding.

This study was supported by VIRK – Vocational Rehabilitation Fund and the Icelandic Research Fund under Grant 196395-052.

Declaration of Interests

The authors report there are no competing interests to declare.

Data Availability Statement

The data that support the findings of this study can be made available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

- Alberts, N. M., Hadjistavropoulos, H. D., Jones, S. L., & Sharpe, D. (2013). The Short Health Anxiety Inventory: A systematic review and meta-analysis. *Journal of Anxiety Disorders*, 27(1), 68–78. <https://doi.org/10.1016/j.janxdis.2012.10.009>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders 5th ed.* (Washington, DC). American Psychiatric Association.
- Amland, A., Malterud, K., & Werner, E. L. (2014). Patients with persistent medically unexplained physical symptoms: A descriptive study from Norwegian general practice. *BMC Family Practice*. <https://doi.org/doi:10.1186/1471-2296-15-107>
- Barsky AJ, Orav E, & Bates DW. (2005). Somatization increases medical utilization and costs independent of psychiatric and medical comorbidity. *Archives of General Psychiatry*, 62(8), 903–910. <https://doi.org/10.1001/archpsyc.62.8.903>
- Beck, J. G., Grant, D. M., Read, J. P., Clapp, J. D., Coffey, S. F., Miller, L. M., & Palyo, S. A. (2008). The Impact of Event Scale –Revised: Psychometric properties in a sample of motor vehicle accident survivors. *Journal of Anxiety Disorders*, 22(2), 187–198. <https://doi.org/10.1016/j.janxdis.2007.02.007>
- Berman, A. H., Bergman, H., Palmstierna, T., & Schlyter, F. (2005). Evaluation of the Drug Use Disorders Identification Test (DUDIT) in Criminal Justice and Detoxification Settings and in a Swedish Population Sample. *European Addiction Research*, 11(1), 22–31. <https://doi.org/10.1159/000081413>
- Budtz-Lilly, A., Vestergaard, M., Fink, P., Carlsen, A. H., & Rosendal, M. (2015). Patient characteristics and frequency of bodily distress syndrome in primary care: A cross-sectional study. *Br J Gen Pract*, 65(638), 617–623. <https://doi.org/10.3399/bjgp15X686545>

- Carson, A. J., Ringbauer, B., Stone, J., McKenzie, L., Warlow, C., & Sharpe, M. (2000). Do medically unexplained symptoms matter? A prospective cohort study of 300 new referrals to neurology outpatient clinics. *Journal of Neurology, Neurosurgery, and Psychiatry*, 68(2), 207–210.
- Castell, B. D., Kazantzis, N., & Moss-Morris, R. E. (2011). Cognitive Behavioral Therapy and Graded Exercise for Chronic Fatigue Syndrome: A Meta-Analysis. *Clinical Psychology: Science and Practice*, 18(4), 311–324. <https://doi.org/10.1111/j.1468-2850.2011.01262.x>
- Craske, M. G., Wolitzky-Taylor, K. B., Labus, J., Wu, S., Frese, M., Mayer, E. A., & Naliboff, B. (2011). A Cognitive-Behavioral Treatment for Irritable Bowel Syndrome Using Interoceptive Exposure to Visceral Sensations. *Behaviour Research and Therapy*, 49(6–7), 413–421. <https://doi.org/10.1016/j.brat.2011.04.001>
- Creamer, M., Bell, R., & Failla, S. (2003). Psychometric properties of the Impact of Event Scale—Revised. *Behaviour Research and Therapy*, 41(12), 1489–1496. <https://doi.org/10.1016/j.brat.2003.07.010>
- Dear, B. F., Titov, N., Perry, K. N., Johnston, L., Wootton, B. M., Terides, M. D., Rapee, R. M., & Hudson, J. L. (2013). The Pain Course: A randomised controlled trial of a clinician-guided Internet-delivered cognitive behaviour therapy program for managing chronic pain and emotional well-being. *Pain*, 154(6), 942–950. <https://doi.org/10.1016/j.pain.2013.03.005>
- den Boeft, M., Twisk, J. W. R., Hoekstra, T., Terluin, B., Penninx, B. W. J. H., van der Wouden, J. C., Numans, M. E., & van der Horst, H. E. (2016). Medically unexplained physical symptoms and work functioning over 2 years: Their association and the

influence of depressive and anxiety disorders and job characteristics. *BMC Family Practice*, 17, 46. <https://doi.org/10.1186/s12875-016-0443-x>

Dimsdale, J., Sharma, N., & Sharpe, M. (2011). What do physicians think of somatoform disorders? *Psychosomatics*, 52(2), 154–159.
<https://doi.org/10.1016/j.psych.2010.12.011>

Esler, J. L., Barlow, D. H., Woolard, R. H., Nicholson, R. A., Nash, J. M., & Erogul, M. H. (2003). A brief cognitive-behavioral intervention for patients with noncardiac chest pain. *Behavior Therapy*, 34(2), 129–148. [https://doi.org/10.1016/S0005-7894\(03\)80009-6](https://doi.org/10.1016/S0005-7894(03)80009-6)

Flóvenz, S., Broddadóttir, E., Brynjólfsson, S., Agnarsdóttir, A., Salkovskis, P. M., & Sigurðsson, J. F. (2021). Tíðni þráláttra óútskýrðra líkamlegra einkenna og tengsl við einkenni þunglyndis og kvíða. *Læknablaðið*.

Flóvenz, S., Salkovskis, P. M., Svandsdóttir, E., Karlsson, H. D., & Andersen, K. (n.d.). Non-Cardiac Chest Pain as a Persistent Physical Symptom: Psychological distress and workability. *Manuscript Submitted for Publication*.

Glombiewski, J. A. a, Sawyer, A. T. a, Gutermann, J. a, Koenig, K. a, Rief, W. b, & Hofmann, S. G. a. (2010). Psychological treatments for fibromyalgia: A meta-analysis. *Pain*, 151(2), 280–295. <https://doi.org/10.1016/j.pain.2010.06.011>

Gregory, J., Flóvenz, S., Broddadóttir, E., Sigurðsson, J. F., & Salkovskis, P. M. (n.d.). Managing ‘Persistent Physical Symptoms’ in adults in primary care: Prevalence, co-morbidity with psychological distress and willingness to engage in treatment. *Manuscript in Preperation*.

- Harris, A. M., Orav, E. J., Bates, D. W., & Barsky, A. J. (2009). Somatization Increases Disability Independent of Comorbidity. *Journal of General Internal Medicine*, 24(2), 155–161. <https://doi.org/10.1007/s11606-008-0845-0>
- Ho, F. Y.-Y., Chan, C. S., Lo, W.-Y., & Leung, J. C.-Y. (2020). The effect of self-help cognitive behavioral therapy for insomnia on depressive symptoms: An updated meta-analysis of randomized controlled trials. *Journal of Affective Disorders*, 265, 287–304. <https://doi.org/10.1016/j.jad.2020.01.062>
- Hoedeman, R., Blankenstein, A. H., Krol, B., Koopmans, P. C., & Groothoff, J. W. (2010). The Contribution of High Levels of Somatic Symptom Severity to Sickness Absence Duration, Disability and Discharge. *Journal of Occupational Rehabilitation*, 20(2), 264–273. <https://doi.org/10.1007/s10926-010-9239-3>
- Ingólfssdóttir, R. (2014). *Psychometric Properties of the Icelandic Version of the Generalized Anxiety Disorder-7*. Háskólinn í Reykjavík.
- Jonsbu, E., Dammen, T., Morken, G., Moum, T., & Martinsen, E. W. (2011). Short-term cognitive behavioral therapy for non-cardiac chest pain and benign palpitations: A randomized controlled trial. *Journal of Psychosomatic Research*, 70(2), 117–123. <https://doi.org/10.1016/j.jpsychores.2010.09.013>
- Karlsson, B., Burell, G., Anderberg, U.-M., & Svärdsudd, K. (2015). Cognitive behaviour therapy in women with fibromyalgia: A randomized clinical trial. *Scandinavian Journal of Pain*, 9, 11–21. <https://doi.org/10.1016/j.sjpain.2015.04.027>
- Katon, W., Lin, E., Von Korff, M., Russo, J., Lipscomb, P., & Bush, T. (1991). Somatization: A spectrum of severity. *The American Journal of Psychiatry*, 148(1), 34–40.

- Kim, D.-Y., Lee, J.-S., Park, S.-Y., Kim, S.-J., & Son, C.-G. (2020). Systematic review of randomized controlled trials for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME). *Journal of Translational Medicine*, 18(1), 7.
<https://doi.org/10.1186/s12967-019-02196-9>
- Kirmayer, L. J., Groleau, D., Looper, K. J., & Dao, M. D. (2004). Explaining medically unexplained symptoms. *Canadian Journal of Psychiatry. Revue Canadienne De Psychiatrie*, 49(10), 663–672. <https://doi.org/10.1177/070674370404901003>
- Kisely, S. R., Campbell, L. A., Yelland, M. J., & Paydar, A. (2015). Psychological interventions for symptomatic management of non-specific chest pain in patients with normal coronary anatomy. *Cochrane Database of Systematic Reviews*, 6.
<https://doi.org/10.1002/14651858.CD004101.pub5>
- Kleinstäuber, M., Witthöft, M., & Hiller, W. (2011). Efficacy of short-term psychotherapy for multiple medically unexplained physical symptoms: A meta-analysis. *Clinical Psychology Review*, 31(1), 146–160. <https://doi.org/10.1016/j.cpr.2010.09.001>
- Konnopka, A., Kaufmann, C., König, H.-H., Heider, D., Wild, B., Szecsenyi, J., Herzog, W., Heinrich, S., & Schaefer, R. (2013). Association of costs with somatic symptom severity in patients with medically unexplained symptoms. *Journal of Psychosomatic Research*, 75(4), 370–375. <https://doi.org/10.1016/j.jpsychores.2013.08.011>
- Kouyanou, K., Pither, C. E., Rabe-Hesketh, S., & Wessely, S. (1998). A comparative study of iatrogenesis, medication abuse, and psychiatric morbidity in chronic pain patients with and without medically explained symptoms. *Pain*, 76(3), 417–426.

- Kroenke, K., & Spitzer, R. L. (2002). The PHQ-9: A new depression diagnostic and severity measure. *Psychiatric Annals*, 32(9), 509–515. <https://doi.org/10.3928/0048-5713-20020901-06>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9. *Journal of General Internal Medicine*, 16(9), 606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>
- Larun, L., Brurberg, K. G., Odgaard-Jensen, J., & Price, J. R. (2017). Exercise therapy for chronic fatigue syndrome. *The Cochrane Database of Systematic Reviews*, 4, CD003200. <https://doi.org/10.1002/14651858.CD003200.pub7>
- Lecrubier, Y., Sheehan, D., Weiller, E., Amorim, P., Bonora, I., Harnett Sheehan, K., Janavs, J., & Dunbar, G. (1997). The Mini International Neuropsychiatric Interview (MINI). A short diagnostic structured interview: Reliability and validity according to the CIDI. *European Psychiatry*, 12(5), 224–231. [https://doi.org/10.1016/S0924-9338\(97\)83296-8](https://doi.org/10.1016/S0924-9338(97)83296-8)
- Ljótsson, B., Falk, L., Vesterlund, A. W., Hedman, E., Lindfors, P., Rück, C., Hursti, T., Andréewitch, S., Jansson, L., Lindefors, N., & Andersson, G. (2010). Internet-delivered exposure and mindfulness based therapy for irritable bowel syndrome – A randomized controlled trial. *Behaviour Research and Therapy*, 48(6), 531–539. <https://doi.org/10.1016/j.brat.2010.03.003>
- Ljótsson, B., Hesser, H., Andersson, E., Lackner, J. M., El Alaoui, S., Falk, L., Aspvall, K., Fransson, J., Hammarlund, K., Löfström, A., Nowinski, S., Lindfors, P., & Hedman, E. (2014). Provoking symptoms to relieve symptoms: A randomized controlled dismantling study of exposure therapy in irritable bowel syndrome. *Behaviour Research and Therapy*, 55, 27–39. <https://doi.org/10.1016/j.brat.2014.01.007>

- Loengaard, K., Bjorner, J. B., Fink, P. K., Burr, H., & Rugulies, R. (2015). Medically unexplained symptoms and the risk of loss of labor market participation—A prospective study in the Danish population. *BMC Public Health*, *15*, 844.
<https://doi.org/10.1186/s12889-015-2177-4>
- Löwe, B., Spitzer, R. L., Williams, J. B. W., Mussell, M., Schellberg, D., & Kroenke, K. (2008). Depression, anxiety and somatization in primary care: Syndrome overlap and functional impairment. *General Hospital Psychiatry*, *30*(3), 191–199.
<https://doi.org/10.1016/j.genhosppsych.2008.01.001>
- Marks, E. M., & Hunter, M. S. (2015). Medically Unexplained Symptoms: An acceptable term? *British Journal of Pain*, *9*(2), 109–114.
<https://doi.org/10.1177/2049463714535372>
- Mattick, R. P., & Clarke, J. C. (1998). Development and validation of measures of social phobia scrutiny fear and social interaction anxiety1. *Behaviour Research and Therapy*, *36*(4), 455–470. [https://doi.org/10.1016/S0005-7967\(97\)10031-6](https://doi.org/10.1016/S0005-7967(97)10031-6)
- McGorm, K., Burton, C., Weller, D., Murray, G., & Sharpe, M. (2010). Patients repeatedly referred to secondary care with symptoms unexplained by organic disease: Prevalence, characteristics and referral pattern. *Family Practice*, *27*(5), 479–486.
<https://doi.org/10.1093/fampra/cmq053>
- Momsen, A. H., Nielsen, C. V., Nielsen, M. B. D., Rugulies, R., & Jensen, C. (2016). Work participation and health-related characteristics of sickness absence beneficiaries with multiple somatic symptoms. *Public Health*, *133*, 75–82.
<https://doi.org/10.1016/j.puhe.2015.10.018>

- Mundt, J. C., Marks, I. M., Katherine, M., & Greist, J. M. (2002). The Work and Social Adjustment Scale: A simple measure of impairment in functioning. *The British Journal of Psychiatry*, 180(5), 461–464. <https://doi.org/10.1192/bjp.180.5.461>
- Nimnuan, C., Hotopf, M., & Wessely, S. (2001). Medically unexplained symptoms: An epidemiological study in seven specialities. *Journal of Psychosomatic Research*, 51(1), 361–367. [https://doi.org/10.1016/S0022-3999\(01\)00223-9](https://doi.org/10.1016/S0022-3999(01)00223-9)
- Ólafsdóttir, H. Ó. (2012). *Psychometric properties of the Icelandic translations of the Social Interaction Anxiety Scale (SIAS) and Social Phobia Scale (SPS)*. University of Iceland.
- Osman, A., Gutierrez, P. M., Barrios, F. X., Kopper, B. A., & Chiro, C. E. (1998). The Social Phobia and Social Interaction Anxiety Scales: Evaluation of Psychometric Properties. *Journal of Psychopathology and Behavioral Assessment; New York*, 20(3), 249–264. <http://dx.doi.org/10.1023/A:1023067302227>
- Pálsdóttir, V. E. (2007). *The validity of the Patient Health Questionnaire (PHQ) within Icelandic primary care*. [Cand.psych.]. University of Iceland.
- Picariello, F., Ali, S., Moss-Morris, R., & Chalder, T. (2015). The most popular terms for medically unexplained symptoms: The views of CFS patients. *Journal of Psychosomatic Research*, 78(5), 420–426. <https://doi.org/10.1016/j.jpsychores.2015.02.013>
- Pike, A., Hearn, L., & Williams, A. C. (2016). Effectiveness of psychological interventions for chronic pain on health care use and work absence: Systematic review and meta-analysis. *Pain*, 157(4), 777–785. <https://doi.org/10.1097/j.pain.0000000000000434>
- Rask, Ørnbøl, E., Rosendal, M., & Fink, P. (2017). Long-Term Outcome of Bodily Distress Syndrome in Primary Care: A Follow-Up Study on Health Care Costs, Work

Disability, and Self-Rated Health. *Psychosomatic Medicine*, 79(3), 345–357.

<https://doi.org/10.1097/PSY.0000000000000405>

Reid, S., Wessely, S., Crayford, T., & Hotopf, M. (2001). Medically unexplained symptoms in frequent attenders of secondary health care: Retrospective cohort study. *BMJ*, 322(7289), 767. <https://doi.org/10.1136/bmj.322.7289.767>

Reinert, D. F., & Allen, J. P. (2007). The Alcohol Use Disorders Identification Test: An Update of Research Findings. *Alcoholism: Clinical and Experimental Research*, 31(2), 185–199. <https://doi.org/10.1111/j.1530-0277.2006.00295.x>

Roca, M., Gili, M., Garcia-Garcia, M., Salva, J., Vives, M., Garcia Campayo, J., & Comas, A. (2009). Prevalence and comorbidity of common mental disorders in primary care. *Journal of Affective Disorders*, 119(1–3), 52–58. <https://doi.org/10.1016/j.jad.2009.03.014>

Salkovskis, P. M. (2002). The Health Anxiety Inventory: Development and validation of scales for the measurement of health anxiety and hypochondriasis. *Psychological Medicine*, 32(5), 843–853. <https://doi.org/10.1017/S0033291702005822>

Salkovskis, P. M., Gregory, J., Sedgwick-Taylor, A., White, J., Opher, S., & Olafsdottir, S. (2016). Extending Cognitive-Behavioural Theory and Therapy to medically unexplained symptoms and long-term physical conditions: A hybrid Transdiagnostic/problem specific approach. *Behavioural Change*.

Saunders, J. B., Aasland, O. G., Babor, T. F., de la Fuente, J. R., & Grant, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction (Abingdon, England)*, 88(6), 791–804.

- Sheehan, D. V., & Lecrubier, Y. (1998). The Mini-International Neuro-mental Interview (M.I.N.I.): The Development and Validation of a Structured Diagnostic Mental Interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*, 59 Suppl 20(Suppl 20), 22–33; quiz 34–57.
- Spitzer, R. L., Kroenke, K., Williams, J. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The gad-7. *Archives of Internal Medicine*, 166(10), 1092–1097. <https://doi.org/10.1001/archinte.166.10.1092>
- Sundin, E. C., & Horowitz, M. J. (2002). Impact of Event Scale: Psychometric properties. *The British Journal of Psychiatry*, 180(3), 205–209. <https://doi.org/10.1192/bjp.180.3.205>
- Tang, N. K. Y., Goodchild, C. E., & Salkovskis, P. M. (2012). Hybrid cognitive-behaviour therapy for individuals with insomnia and chronic pain: A pilot randomised controlled trial. *Behaviour Research and Therapy*, 50(12), 814–821. <https://doi.org/10.1016/j.brat.2012.08.006>
- Thakral, M., Von Korff, M., McCurry, S. M., Morin, C. M., & Vitiello, M. V. (2020). Changes in dysfunctional beliefs about sleep after cognitive behavioral therapy for insomnia: A systematic literature review and meta-analysis. *Sleep Medicine Reviews*, 49, 101230. <https://doi.org/10.1016/j.smrv.2019.101230>
- Viðarsson, J. V. (2016). *Health Anxiety and Medically Unexplained Symptoms: Psychometric properties of two Translated Questionnaires*. Reykjavik University.
- VIRK Vocational Rehabilitation Fund. (2013). *Annual Report 2013*. VIRK Vocational Rehabilitation Fund. <https://www.virk.is/static/files/kynningarefni/virk-arsrit-2013-net.pdf>

VIRK Vocational Rehabilitation Fund. (2014). *Annual report 2014*. VIRK-Vocational Rehabilitation Fund.

https://www.virk.is/static/files/_2014/Arsfundur_2014/virk_arsrit_2014_net.pdf

VIRK Vocational Rehabilitation Fund. (2018). *Annual report 2018* (pp. 16–17). VIRK-Vocational Rehabilitation Fund. https://www.virk.is/static/files/virk-a-9568-rsrit-2018_lokaskjal.pdf

Voluse, A. C., Gioia, C. J., Sobell, L. C., Dum, M., Sobell, M. B., & Simco, E. R. (2012). Psychometric properties of the Drug Use Disorders Identification Test (DUDIT) with substance abusers in outpatient and residential treatment. *Addictive Behaviors*, 37(1), 36–41. <https://doi.org/10.1016/j.addbeh.2011.07.030>

Wessely, S., Nimnuan, C., & Sharpe, M. (1999). Functional somatic syndromes: One or many? *Lancet (London, England)*, 354(9182), 936–939. [https://doi.org/10.1016/S0140-6736\(98\)08320-2](https://doi.org/10.1016/S0140-6736(98)08320-2)

Wessely, & White, P. D. (2004). There is only one functional somatic syndrome. *The British Journal of Psychiatry*, 185(2), 95–96. <https://doi.org/10.1192/bjp.185.2.95>

Zahra, D., Qureshi, A., Henley, W., Taylor, R., Quinn, C., Pooler, J., Hardy, G., Newbold, A., & Byng, R. (2014). The work and social adjustment scale: Reliability, sensitivity and value. *International Journal of Psychiatry in Clinical Practice*, 18(2), 131–138. <https://doi.org/10.3109/13651501.2014.894072>

Table 1. Gender, Education and Relationship Status of Participants, with and without PPS

Demographic variables	All participants	Participants with PPS	Participants without PPS	χ^2
Gender				
Male	71 (21.9%)	54 (20.6%)	17 (27.4.0%)	1.358
Female	253 (78.1%)	202 (79.4%)	45 (72.6%)	
Education completed				
Primary school	82 (27.2%)	63 (26.1%)	19 (31.7%)	1.899
Secondary school	103 (34.2%)	81 (33.6.2%)	22 (36.7%)	
Other vocational education	19 (6.3%)	15 (6.2%)	4 (6.7%)	
University degree	97 (32.2%)	82 (34.0%)	15 (25.0%)	
Relationship status				
Single	161 (52.8%)	131 (53.7%)	30 (49.2%)	9.748*
Married or cohabiting	97 (31.8%)	83 (34.0%)	14 (23.0%)	
Separated or widowed	47 (15.4%)	30 (12.2%)	17 (27.9%)	

Note. PPS = Persistent physical symptoms.

* $p < 0.05$

Table 2. Numbers and Percentages of Participants, with and without PPS, with Scores above cut-off on the SHAI, PHQ-9, GAD-7, IES-R and the combined SIAS and SPS

Measure	All participants	Participants with PPS	Participants without PPS	χ^2	OR
SHAI ≥ 18	104 (32.7%)	95 (37.0%)	9 (14.8%)	11.05*	3.39
PHQ-9 ≥ 10	245 (75.9%)	218 (82.2%)	27 (44.3%)	40.97*	6.24
GAD-7 ≥ 10	170 (52.8%)	148 (56.7%)	22 (36.1%)	8.45*	2.32
SIAS/SPS ≥ 74	93 (29.1%)	80 (31.0%)	13 (21.0%)	2.44	-
IES-R ≥ 33	114 (37.9%)	95 (38.9%)	19 (33.3%)	0.34	-

Note. PPS = Persistent physical symptoms; OR = Odds ratio; SHAI = Short Health Anxiety Inventory; PHQ-9 = Patient Health Questionnaire-9; GAD-7=Generalised Anxiety Disorder-7; SPS = Social Phobia Scale; SIAS = Social Interaction Anxiety Scale; IES-R = Impact of Event Scale-Revised;

^a IES-R scores were only calculated for participants who reported a past traumatic experience.

* $p < 0.05$

Table 3. Prevalence of Depression, Anxiety and Health Anxiety Symptoms in the Clinical Range in Participants with PPS

Measure	Participants with PPS N (%)
PHQ-9 _{≥10}	218 (83.2%)
PHQ-9 _{≥15}	119 (45.6%)
GAD-7 _{≥10}	148 (56.7%)
GAD-7 _{≥15}	70 (26.8%)
SHAI _{≥18}	95 (37.0%)
PHQ-9 _{≥10} , GAD-7 _{≥10} or SHAI _{≥18}	226 (86.6%)
PHQ-9 _{≥15} , GAD-7 _{≥15} or SHAI _{≥18}	159 (61.4%)

Note. PPS = Persistent physical symptoms; PHQ-9 = Patient Health Questionnaire-9 GAD-7=Generalised Anxiety Disorder-7; SHAI = Short Health Anxiety Inventory; PHQ-9 scores ≥ 10 indicate moderate depressive symptoms and ≥ 15 indicate moderately severe symptoms; GAD-7 scores ≥ 10 indicate moderate anxiety and ≥ 15 indicates severe anxiety; SHAI scores ≥ 18 indicate probable health anxiety

Figure 1. Percentage of Participants who met Criteria for a Number of PPS Types.





Figure 2. Prevalence of different PPS Types.



Figures

Figure 1. Percentage of Participants who met Criteria for a Number of PPS Types.

Figure 2. Prevalence of different PPS Types.