



Research



Check for updates

Cite this article: Şengül Y. 2024 A variational approach to frame-indifferent quasistatic viscoelasticity of rate type. *Phil. Trans. R. Soc. A* **382**: 20230307.

<https://doi.org/10.1098/rsta.2023.0307>

Received: 12 March 2024

Accepted: 4 June 2024

One contribution of 16 to a theme issue ‘Non-smooth variational problems with applications in mechanics’.

Subject Areas:

analysis, mathematical modelling, materials science, differential equations, mechanics

Keywords:

nonlinear viscoelasticity, quasistatic, weak solution, regularity

Author for correspondence:

Yasemin Şengül

e-mail: sengultezely@cardiff.ac.uk

A variational approach to frame-indifferent quasistatic viscoelasticity of rate type

Yasemin Şengül

School of Mathematics, Cardiff University, Cardiff CF24 4AG, UK

YŞ, 0000-0001-5923-3173

The three-dimensional dynamical model for nonlinear viscoelasticity of strain-rate type is investigated in a quasistatic setting under the assumption of higher-order regularity of the deformation, which in the literature is referred to as the case of non-simple materials. The existence of weak solutions is proven using a time-discretization technique while respecting the condition of dynamical frame indifference. Some observations on frame indifference for strain-rate-type stresses are made, and corrections are proposed for some related work in the literature. Finally, a counterexample is given to show that the assumed higher-order regularity is necessary in order to obtain the required compactness.

This article is part of the theme issue ‘Non-smooth variational problems with applications in mechanics’.

1. Introduction

The requirement for a well-posed qualitative mathematical theory for properly formulated dynamics, based on fundamental physical principles, has been recognized for a long time. In order to realize this purpose, one needs to have answers to some questions that can be stated generally for any evolution equation associated with a non-elliptic variational integral. This paper aims to address the question of the existence of solutions for frame-indifferent viscoelastic models of rate type. An extensive overview of such models can be found in a study by Şengül [1]. These dynamical models have been successfully studied in the literature, including the one-dimensional case [2–5], the general

© 2024 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

three-dimensional case [6–8] and the thermodynamical case when temperature dependence is also taken into account [9].

In a study by Şengül [10], for the first time, quasistatic approximation was considered in the context of nonlinear viscoelasticity of rate type and a variational approach was introduced in a three-dimensional setting for the existence of solutions while handling the dynamical frame indifference of the stress. Following the study by Şengül [10], in Ball & Şengül [11], the one-dimensional problem was studied, which was proved to be equivalent to a gradient flow owing to the quasistatic nature of the governing equations resulting from neglecting the inertia term. Similarly, in Mielke *et al.* [12], the quasistatic case was considered and a variational approach was adopted via metric gradient flows. After the study by Şengül [10], approximating dynamical models by the corresponding quasistatic equations has been adopted by many studies in various contexts. However, none of the studies for such rate-type viscoelastic models have been successful in obtaining well-posedness while dealing with the requirement of frame indifference.

The main difference between the current work and the model investigated by Şengül [10] is that here a higher-order gradient term is included in the equation of linear momentum balance in order to obtain the necessary compactness while passing to the limit after the application of a time-discretization method. More precisely, a term including the second gradient of the deformation is added to the system, representing the *strain gradient*. Materials with such models are called *non-simple materials*. These models have been successfully studied in the literature in various contexts, the most relevant to the current work being by Friedrich & Kružík [13], Badal *et al.* [14], Krömer & Roubíček [15] and Mielke & Roubíček [16]. More precisely, in [13] and [15], the same model considered here is investigated. However, the focus of the first work is more on the linearized case rather than large strains, while the latter one focuses on possible self-contact in deformed configuration. In [16], the same model is coupled with a suitable heat equation so that thermal effects are also taken into account. While this makes the system to be studied more general, the analysis is more complicated, and some restrictions are put into place, including regularization of the mechanical equation with a strain-rate term in order to handle the necessary compactness. More recently, [14] studied thermoviscoelasticity in a quasistatic setting by refining the results obtained in [16].

The present work contributes to the above-mentioned literature in three ways. Firstly, new investigations are proven for frame-indifferent stresses of rate type as well as the corresponding dissipation potentials. Also, some expressions of frame indifference in the literature are proven to be incorrect. Secondly, weak solutions to frame-indifferent quasistatic viscoelasticity in the context of non-simple materials are proven to exist using a variational approach and by applying a time-discretization method. While this can be seen to be less general as a special case of the systems investigated in [16] and [14] owing to being isothermal, the analysis is rather clear, focusing solely on the mechanical equation, in particular, on a specific viscoelastic part of the stress tensor. Thirdly, a counterexample is given in the static case, which can be viewed as the state at each fixed time step, showing that without the inclusion of the strain-gradient term in the model, it is not possible to obtain compactness, leading to the existence of solutions.

The content of the paper is as follows. In §2, we describe some preliminary notions that will be used throughout the paper. Then, in §3, we introduce the governing equations to be studied. §4 is devoted to frame indifference, with some observations about rate-type stresses and dissipation potentials and corrections to some statements in the literature. In §5, we state the assumptions, introduce the time-discretization scheme and prove the main result. Finally, in §6, we give our counterexample before the concluding remarks in the last part.

2. Preliminaries

For a homogeneous elastic body with a reference configuration $\Omega \subset \mathbb{R}^3$ and with unit reference density, a motion is an evolution of diffeomorphisms $y(\cdot, t): \Omega \rightarrow \mathbb{R}^3$, where $t \in [0, T] \subset \mathbb{R}$. The *deformation gradient* at time t is written as $\nabla y(x, t)$, or equivalently F , and can be identified with the $n \times n$ matrix of partial derivatives given as

$$(\nabla y)_{i\alpha} = y_{i,\alpha} = \frac{\partial y_i}{\partial x_\alpha}.$$

We would like our model to be physically realistic, and hence it is necessary to avoid interpenetration of matter so that two distinct material points cannot simultaneously occupy the same position in space. In order to ensure this, it is required that for (almost) every t , the actual position field $y(\cdot, t)$ is injective. This is equivalent to saying that the deformation y is invertible in Ω . We can still allow some cases where, for example, self-contact occurs on the boundary (see [17] for more information). Therefore, it is enough to assume that the admissible deformations satisfy the constraint

$$\det \nabla y(x, t) > 0, \quad (2.1)$$

ensuring that the admissible deformations are *orientation-preserving* and *locally invertible*. As discussed by Ball [17] via some examples, however, local invertibility does not imply global invertibility.

An elastic material is *hyperelastic* if there exists a function $W: \Omega \times \text{GL}^+(3) \rightarrow \mathbb{R}$ differentiable with respect to the variable $F \in \text{GL}^+(3)$ for each $x \in \Omega$ such that the *first Piola–Kirchhoff stress tensor* is given by

$$T_R(x, F) = \frac{\partial W}{\partial F}(x, F). \quad (2.2)$$

Here, $\text{GL}^+(3)$ denotes the set of matrices in $\mathbb{R}^{3 \times 3}$ with positive determinant. The function W is called the *stored-energy function*. Naturally, if the material is *homogeneous*, it is a function of F only (cf. [18,19]), which is the case we consider in this work. As noted by Ball [17], this is more restrictive than saying that Ω is occupied by the same material at each point, since it is possible to have some pre-existing stresses. We can also define the *second Piola–Kirchhoff stress tensor* as

$$\hat{T}(x, F) = F^{-1} T_R(x, F), \quad (2.3)$$

and the *Cauchy stress tensor* as

$$T(x, F) = (\det F)^{-1} T_R(x, F) F^T. \quad (2.4)$$

The elastic energy corresponding to the deformation y is defined as

$$I(y) = \int_{\Omega} W(\nabla y(x, t)) dx. \quad (2.5)$$

Unless stated otherwise, we will make the following convention that the initial free energy is finite, that is,

$$\int_{\Omega} W(\nabla y(x, 0)) dx < \infty.$$

The matrix

$$C = \nabla y^T \nabla y \quad (2.6)$$

is called the *right Cauchy–Green strain tensor*. It is symmetric and is positive-definite where ∇y is non-singular.

3. Modelling

We use the Lagrangian formulation in the domain $\Omega \subset \mathbb{R}^3$ bounded with a smooth boundary Γ . We consider the time variable $t \in [0, T]$. A generalization of Kelvin–Voigt type viscoelasticity can be modelled as

$$\ddot{y} - \text{Div} DW(\nabla y) - \text{Div} S(\nabla y, \nabla \dot{y}) - f(t) = 0, \quad (3.1)$$

where the constitutive equation for the first Piola–Kirchhoff stress tensor reads $T_R = W(\nabla y) + S(\nabla y, \nabla \dot{y})$ with S being the viscoelastic part, and $f(t)$ is the external mechanical loading, which might consist of a dead force and boundary traction. Here, the superposed dot stands for the time derivative.

In this paper, we have two main postulates: firstly, we consider the quasistatic approximation for (3.1), meaning that the inertial effects are neglected; secondly, in order to gain enough regularity to handle the physical nonlinearities, we make the assumption that there is an energy contribution coming from the strain gradient. As a result, the total free energy (2.5) becomes

$$E(y(t)) := \int_{\Omega} (W(\nabla y) + \mathcal{H}(\nabla^2 y)) dx, \quad (3.2)$$

where $\mathcal{H} = \mathcal{H}(\nabla F)$ is the potential corresponding to the elastic hyperstress. As mentioned in the Introduction, materials with such elastic energy are referred to as *second-grade* or *non-simple materials* in the literature. Unfortunately, the problem of the existence of solutions for simple materials, that is, without having the higher-order gradient term, is still open both in the quasistatic and dynamical cases. We can introduce the dissipation potential Ψ as

$$D_H \Psi(F, H) = S(F, H). \quad (3.3)$$

As a result, the balance of linear momentum in the quasistatic case implies

$$\text{Div} (DW(F) - \text{Div} \mathcal{H}'(\nabla F)) + \text{Div} S(F, \dot{F}) = f(t). \quad (3.4)$$

In this paper, we are interested in the existence of solutions of (3.4), which can also be viewed as an abstract gradient flow. In the case of simple materials in one space dimension, it is explicitly shown by Ball & Şengül [11] that this model is equivalent to the equation of gradient flows.

We impose the following initial conditions

$$y(x, 0) = y_0 \quad \text{on } \Omega. \quad (3.5)$$

For boundary conditions, for simplicity and without loss of generality, we only consider $f(t)$ being a time-dependent dead force given as

$$f: [0, T] \times \Omega \rightarrow \mathbb{R}^3. \quad (3.6)$$

The reader is referred to a recent paper by Mielke & Roubíček [16] for the treatment of boundary conditions including traction.

4. Frame indifference

(a) Definition and mathematical expression

The mechanical behaviour of materials is governed by some general principles, one of which is the principle of frame indifference. As a general axiom in physics, it states that the response of a material must be independent of the observer (e.g. [20]). In particular, it restricts the form of the constitutive functions and thus plays an important role in nonlinear continuum mechanics. We state it as follows:

The Principle of Frame Indifference (Objectivity): Constitutive functions are invariant under rigid motions.

In order to express this principle as a mathematical condition, we first note that a change of observer can be seen as the application of rigid-body motions on the current configuration. Since a rigid-body motion consists of a translation and a rotation, in each of these motions, the relative positions of the points of the material remain the same. As the deformation gradient is not affected by the translations of the origin, if the body undergoes a rigid body motion, then the corresponding expression for the stress becomes

$$\tilde{T}_R(x, \tilde{t}) = R(t)T_R(x, t), \tilde{t} = t + a, a \in \mathbb{R}.$$

A formal mathematical statement can be given by the following result.

Lemma 4.1. Any frame-indifferent stress tensor $S(F, \dot{F})$ can be written as

$$S(F, \dot{F}) = R S(U, \dot{U}), \quad (4.1)$$

where $R \in \text{SO}(3)$ and U is the right Cauchy–Green stretch tensor. Here, $\text{SO}(3)$ denotes the set of rotations in \mathbb{R}^3 .

Proof. By the polar decomposition theorem (refer e.g. [21]), we have

$$\begin{aligned} R^T S(F, \dot{F}) &= S(R^T F, \widehat{R^T \dot{F}}) = S(R^T R U, \dot{R}^T \dot{F} + R^T \dot{F}) \\ &= S(U, \dot{R}^T R U + R^T (\dot{R} U + R \dot{U})) \\ &= S(U, (\dot{R}^T R + R^T \dot{R}) U + \dot{U}) = S(U, \dot{U}) \end{aligned}$$

as required. ■

We can obtain a more convenient form of (4.1) by using the second Piola–Kirchhoff stress tensor as follows:

$$T(F, \dot{F}) = F^{-1} S(F, \dot{F}) = U^{-1} S(U, \dot{U}) = :G(C, \dot{C}).$$

Thus, we have

$$S(F, \dot{F}) = F G(C, \dot{C}). \quad (4.2)$$

It is also worth mentioning that rotations are involved in both material symmetry and frame indifference, but they act differently. More precisely, in material symmetry, the rotation acts in the reference configuration, and in frame indifference, the rotation acts in the deformed configuration. Therefore, it is not possible to obtain one variant by rotating another. In other words, given symmetric matrices U_1 and U_2 , it is not possible to find a rotation R such that $R U_1 = U_2$, since this would be inconsistent with the uniqueness property stated in the polar decomposition theorem.

(b) Further observations on frame indifference

This section is devoted to some trivial but crucial observations we make on frame indifference.

Lemma 4.2. Any frame-indifferent stress $S(F, \dot{F})$ should satisfy

$$S(F, \dot{F}) : \dot{F} = \frac{1}{2} G(C, \dot{C}) : \dot{C}.$$

Proof. We know by (4.2) that any frame-indifferent $S(F, \dot{F})$ takes the form $S(F, \dot{F}) = F G(C, \dot{C})$. Therefore, using the fact that $G(C, \dot{C})$ is symmetric, we get

$$\begin{aligned} S(F, \dot{F}) : \dot{F} &= F G(C, \dot{C}) : \dot{F} = G(C, \dot{C}) : F^T \dot{F} = G(C, \dot{C}) : \dot{F}^T F \\ &= G(C, \dot{C}) : \frac{1}{2} (\dot{F}^T F + F^T \dot{F}) = \frac{1}{2} G(C, \dot{C}) : \dot{C} \end{aligned}$$

as required. ■

Lemma 4.3. *The condition*

$$G(C, \dot{C}): \dot{C} \geq \gamma |\dot{F}|^2, \quad \gamma > 0 \quad \text{a constant}, \quad (4.3)$$

contradicts frame indifference.

Proof. Assume for contradiction that there exists a frame-indifferent S satisfying (4.3). By lemma 4.2, we have

$$S(F, \dot{F}): \dot{F} \geq \gamma |\dot{F}|^2 \quad \Leftrightarrow \quad G(C, \dot{C}): \dot{C} \geq 2\gamma |\dot{F}|^2.$$

Choosing $F = R(t) = \exp(Kt) \in \text{SO}(3)$, where K is skew, we get

$$\dot{F} = K \exp(Kt) \quad \text{and} \quad |\dot{F}|^2 = |K|^2 \neq 0.$$

However, $C = F^T F = R^T R = 1$ implies $\dot{C} = 0$, giving a contradiction. ■

Remark 4.1. The condition

$$G(C, \dot{C}): \dot{C} \geq \gamma |\dot{C}|^2, \quad \gamma > 0 \quad \text{a constant},$$

does not contradict frame indifference as can be seen easily by choosing $G(C, \dot{C}) = \dot{C}$ in (4.3).

In contradiction to the claim of Tvedt [7], we have that

Lemma 4.4. *The assumption*

$$(S(F, \dot{F}) - S(F, \dot{H})):(\dot{F} - \dot{H}) \geq \gamma |\dot{F} - \dot{H}|^2, \quad \gamma > 0 \quad (4.4)$$

is incompatible with frame indifference.

Proof. If, for contradiction, the claim was true, then there would exist a frame-indifferent S satisfying (4.2) so that (4.4) would give

$$(F G(C, \dot{C}) - F G(C, \dot{H}^T F + F^T \dot{H})):(\dot{F} - \dot{H}) \geq \gamma |\dot{F} - \dot{H}|^2. \quad (4.5)$$

Let us define $A := G(C, \dot{C}) - G(C, \dot{H}^T F + F^T \dot{H})$ so that we get

$$\begin{aligned} (F G(C, \dot{C}) - F G(C, \dot{H}^T F + F^T \dot{H})):(\dot{F} - \dot{H}) &= A : F^T (\dot{F} - \dot{H}) = \\ &= \frac{1}{2} [A + A^T] : F^T (\dot{F} - \dot{H}) = \frac{1}{2} [A : F^T (\dot{F} - \dot{H}) + A^T : F^T (\dot{F} - \dot{H})] \\ &= \frac{1}{2} [A : F^T (\dot{F} - \dot{H}) + A : (\dot{F} - \dot{H})^T F] = \frac{1}{2} A : [F^T (\dot{F} - \dot{H}) + (\dot{F} - \dot{H})^T F] \\ &= \frac{1}{2} A : [F^T \dot{F} - F^T \dot{H} + \dot{F}^T F - \dot{H}^T F] = \frac{1}{2} A : [\dot{C} - (F^T \dot{H} + \dot{H}^T F)]. \end{aligned}$$

Therefore, (4.5) is now equivalent to

$$\frac{1}{2} (G(C, \dot{C}) - G(C, \dot{H}^T F + F^T \dot{H})) : (\dot{C} - (F^T \dot{H} + \dot{H}^T F)) \geq \gamma |\dot{F} - \dot{H}|^2.$$

However, for any given G , we can choose $F = I$ in this inequality and obtain

$$\frac{1}{2} (G(1, \dot{F}^T + \dot{F}) - G(1, \dot{H}^T + \dot{H})) : (\dot{F}^T + \dot{F} - \dot{H} - \dot{H}^T) \geq \gamma |\dot{F} - \dot{H}|^2.$$

Choosing $\dot{F} = 0$ now gives

$$\frac{1}{2} (G(1, 0) - G(1, \dot{H}^T + \dot{H})) : (-\dot{H} - \dot{H}^T) \geq \gamma |-\dot{H}|^2.$$

Finally, choosing \dot{H} to be a non-zero and skew matrix makes the left-hand side vanish, whereas the right-hand side remains positive. This gives a contradiction proving the claim. ■

Contradicting the claim of Antman [22] we have that

Lemma 4.5. *The following statement is incompatible with frame indifference.*

$$(S(F, \dot{F} + \dot{H}) - S(F, \dot{F})) : \dot{H} > 0 \quad ; \quad \forall \dot{H} \neq 0, \forall \dot{F}. \quad (4.6)$$

Proof. If, for contradiction, the claim was true, then there would exist a frame-indifferent stress tensor S satisfying (4.2), and (4.6) would be equivalent to

$$\begin{aligned} & \left\{ F[G(F^T F, (\dot{F} + \dot{H})^T F + F^T(\dot{F} + \dot{H})) - G(F^T F, \dot{F}^T F + F^T \dot{F})] \right\} : \dot{H} > 0, \\ & \forall \dot{H} \neq 0 \text{ and } \forall \dot{F}. \end{aligned}$$

Taking $F = I$ in this inequality gives

$$\left\{ G(1, (\dot{F} + \dot{H})^T + (\dot{F} + \dot{H})) - G(1, \dot{F}^T + \dot{F}) \right\} : \dot{H} > 0.$$

Letting \dot{F} be skew reduces it further to

$$\left\{ G(1, \dot{H}^T + \dot{H}) - G(1, 0) \right\} : \dot{H} > 0.$$

We can choose \dot{H} to be a non-zero and skew matrix, which will make the left-hand side vanish, giving a contradiction. \blacksquare

(c) Dissipation potentials

In this section, we aim to show that it is possible to have convex (in \dot{F}) potential functions $\Psi(F, \dot{F})$ leading to frame-indifferent viscoelastic stress $S(F, \dot{F})$ as a result of the relation (3.3).

Lemma 4.6. *Let*

$$\Psi(F, H) = \frac{1}{4} |F^T H + H^T F|^2. \quad (4.7)$$

Then, $\Psi(F, H)$ is convex in H , and

$$S(F, \dot{F}) := \frac{\partial \Psi}{\partial \dot{F}}(F, \dot{F}) = F(\dot{F}^T F + F^T \dot{F}) = F \dot{C},$$

which is frame-indifferent.

Proof. The convexity of $\Psi(F, H)$ follows immediately from the fact that it is a non-negative quadratic form in H . We now show that $S(F, \dot{F}) = F \dot{C}$. We have

$$\begin{aligned} \frac{\partial \Psi}{\partial H}(F, H) &= \frac{1}{4} \frac{\partial}{\partial H_{k\beta}} |F^T H + H^T F|^2 = \\ &= \frac{1}{2} (F_{\alpha i} H_{\alpha j} + H_{\alpha i} F_{\alpha j}) (F_{\alpha i} \delta_{k\alpha} \delta_{j\beta} + \delta_{k\alpha} \delta_{i\beta} F_{\alpha j}) \\ &= \frac{1}{2} (F^T H + H^T F)_{ij} (F_{ki} \delta_{j\beta} + \delta_{i\beta} F_{kj}) \\ &= \frac{1}{2} ((F^T H + H^T F)_{i\beta} F_{ki} + (F^T H + H^T F)_{\beta j} F_{kj}) \\ &= \frac{1}{2} ((F^T H + H^T F)_{\beta i}^T F_{ik}^T + (F^T H + H^T F)_{\beta j} F_{jk}^T) \\ &= \frac{1}{2} \left(((F^T H + H^T F)^T F^T)_{\beta k} + ((F^T H + H^T F) F^T)_{\beta k} \right) \\ &= \frac{1}{2} \left[\left((F^T H + H^T F)^T + (F^T H + H^T F) \right) F^T \right]_{k\beta}^T \\ &= \frac{1}{2} \left((H^T F + F^T H + F^T H + H^T F) F^T \right)^T \\ &= ((F^T H + H^T F) F^T)^T = F (F^T H + H^T F)^T \\ &= F (H^T F + F^T H). \end{aligned}$$

As $\dot{C} = \dot{F}^T F + F^T \dot{F}$, putting $H = \dot{F}$ above proves the claim. Frame indifference of $S(F, \dot{F})$ follows immediately from (4.2). ■

Lemma 4.7. *Let*

$$\Psi(F, H) = \frac{1}{4} | H F^{-1} + F^{-T} H^T |^2. \quad (4.8)$$

Then, $\Psi(F, H)$ is convex in H , and

$$S(F, \dot{F}) := \frac{\partial \Psi}{\partial \dot{F}}(F, \dot{F}) = (\dot{F} F^{-1} + (F^T \dot{F})^T) F^{-T} = F C^{-1} \dot{C} C^{-1},$$

which is frame-indifferent.

Proof. Following the proof in lemma 4.6, one can easily show that $\Psi(F, H)$ is convex with respect to H . Therefore, we skip it here and prove only that $S(F, \dot{F}) = F C^{-1} \dot{C} C^{-1}$, whose frame indifference follows from (4.2) again. In order to calculate S , we adopt a different approach from that of the previous result.¹ We have

$$\Psi(F, A + \varepsilon B) = \frac{1}{4} | (A + \varepsilon B) F^{-1} + F^{-T} (A + \varepsilon B)^T |^2.$$

Therefore, we obtain

$$\begin{aligned} \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \Psi(F, A + \varepsilon B) &= \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \frac{1}{4} | (A F^{-1} + F^{-T} A^T) + \varepsilon (B F^{-1} + F^{-T} B^T) |^2 \\ &= \frac{1}{2} (A F^{-1} + F^{-T} A^T) : (B F^{-1} + F^{-T} B^T) \\ &= \frac{1}{2} (A F^{-1} + F^{-T} A^T) : B F^{-1} + \frac{1}{2} (A F^{-1} + F^{-T} A^T) : F^{-T} B^T \\ &= \frac{1}{2} (A F^{-1} F^{-T} + F^{-T} A^T F^{-T}) : B + \frac{1}{2} (F^{-1} A F^{-1} + F^{-1} F^{-T} A^T) : B^T \\ &= \frac{1}{2} (A F^{-1} F^{-T} + F^{-T} A^T F^{-T}) : B + \frac{1}{2} (F^{-T} A^T F^{-T} + A F^{-1} F^{-T}) : B \\ &= (A F^{-1} F^{-T} + F^{-T} A^T F^{-T}) : B. \end{aligned}$$

This gives

$$\frac{\partial \Psi}{\partial A}(F, A) = F^{-T} A^T F^{-T} + A F^{-1} F^{-T}$$

and hence

$$\begin{aligned} \frac{\partial \Psi}{\partial \dot{F}}(F, \dot{F}) &= F^{-T} \dot{F}^T F^{-T} + \dot{F} F^{-1} F^{-T} \\ &= (F F^{-1}) F^{-T} \dot{F}^T (F F^{-1}) F^{-T} + (F F^{-1}) (F^{-T} \dot{F}^T) \dot{F} F^{-1} F^{-T} \\ &= F (F^{-1} F^{-T}) (\dot{F}^T F) (F^{-1} F^{-T}) + F (F^{-1} F^{-T}) (F^T \dot{F}) (F^{-1} F^{-T}) \\ &= F C^{-1} \dot{C} C^{-1} \end{aligned}$$

as required. ■

Remark 4.2. Strict convexity of $\Psi(F, \cdot)$ is incompatible with frame indifference (see [6]).

As a result, we conclude that there exist dissipation potentials Ψ that not only satisfy convexity but also give frame-indifferent S in (3.3).

¹I thank Gero Friesecke for this proof, which is much simpler than the original one.

5. Existence of solutions

In this section, we prove the existence of weak solutions to (3.4) complemented with initial and boundary conditions using a time-discretization method as first introduced in Şengül [10] and Mielke *et al.* [12], and more recently followed in Mielke & Roubíček [16].

(a) Assumptions and auxiliary results

Throughout the paper, we use the standard notation for L^p as Lebesgue spaces and similarly for $W^{k,p}$ Sobolev spaces of functions whose k -th order weak derivatives are in L^p . Spaces $W^{k,p}([0, T]; X)$ denote Banach spaces of mappings from $L^p([0, T]; X)$ whose k -th order weak derivatives with respect to time variable are also in $L^p([0, T]; X)$. Also, $H^k = W^{k,2}$.

We make the following assumptions on the free energy density, the strain-gradient function and the dissipation potential. It is important to note that, as indicated in Friedrich & Kružík [13], as a result of the inclusion of the higher-order gradient term, it is not necessary to have any kind of convexity condition on the stored-energy density function W .

(i) $W: \text{GL}^+(3) \rightarrow \mathbb{R}^+$ is C^2 , frame-indifferent and satisfies

$$W(F) \geq \alpha |F|^s + \frac{\alpha}{(\det F)^q}, \quad \forall F \in \text{GL}^+(3), \quad (5.1)$$

where $\alpha > 0$ is a constant, $s > 2$, $q \geq 3p/(p-3)$ (with $p > 3$) and $\text{GL}^+(3)$ is as before.

(ii) $\mathcal{H}: \mathbb{R}^{3 \times 3 \times 3} \rightarrow \mathbb{R}^+$ is convex, frame-indifferent and C^1 with

$$\alpha |G|^p \leq \mathcal{H}(G) \leq K(1 + |G|^p), \quad \forall G \in \mathbb{R}^{3 \times 3 \times 3}, \quad (5.2)$$

where K is a constant, possibly very large, and $p > 3$.

(iii) $\Psi: \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^+$ is continuous and given by (4.7) so that $S(F, \dot{F}) = F\dot{C}$.

(iv) $f(t) \in L^2([0, T]; \mathbb{R}^3)$ and $y_0 \in W^{1,2}(\Omega)$.

We will make use of the following auxiliary results proven in [16] (see also [23]). We include them here for the convenience of the reader.

Proposition 5.1. *Assume that the components W and \mathcal{H} of the energy function $E: W^{2,p}(\Omega; \mathbb{R}^3) \rightarrow \mathbb{R}$ satisfy assumptions (5.1) and (5.2). Then, for each constant $C > 0$ there is a constant $\hat{C} > 0$ such that all y with $E(y(t)) \leq C$ satisfy*

$$\|y\|_{W^{2,p}} \leq \hat{C}, \quad \|y\|_{C^{1,1-3/p}} \leq \hat{C}, \quad \det \nabla y(x) \geq 1/\hat{C}, \quad \|(\nabla y)^{-1}\|_{C^{1,1-3/p}} \leq \hat{C}.$$

Proposition 5.2. *For fixed $\lambda \in (0, 1)$ and positive constant $K > 1$ define the set*

$$S_K := \left\{ F \in C^\lambda(\Omega; \mathbb{R}^{3 \times 3}); \|F\|_{C^\lambda} \leq K, \min_{x \in \Omega} \det F(x) \geq 1/K \right\}.$$

Then, for all $K > 1$, there exists a constant $c_K > 0$ such that for all $F \in S_K$ we have

$$\forall v \in W^{1,2}(\Omega; \mathbb{R}^3) : \int_{\Omega} |F^T \nabla v + (\nabla v)^T F|^2 dx \geq c_K \|v\|_{W^{1,2}}.$$

Combining propositions (5.1) and (5.2), one can obtain the following corollary.

Corollary 5.1. *Given $C > 0$, there exists a $C_K > 0$ such that for all y with $E(y(t)) \leq C$ we have*

$$\forall v \in W^{1,2}(\Omega; \mathbb{R}^3) : \int_{\Omega} |(\nabla y)^T \nabla v + (\nabla v)^T \nabla y|^2 dx \geq C_K \|v\|_{W^{1,2}}.$$

We give the definition of a weak solution as follows:

Definition 5.1. A function $y: [0, T] \times \Omega \rightarrow \mathbb{R}^3$ is called a weak solution of (3.4) with initial conditions (3.5) and boundary conditions (3.6) if $y \in L^2([0, T]; W^{2,p}(\Omega, \mathbb{R}^3))$ satisfying $\det \nabla y > 0$ as well as the identity

$$\int_0^T \int_{\Omega} (S(F, \dot{F}) + DW(F) : \nabla z + \mathcal{H}(\nabla^2 y) : \nabla^2 z) dx dt = \int_0^T \int_{\Omega} f \cdot z dx dt, \quad (5.3)$$

for all smooth $z: [0, T] \times \Omega \rightarrow \mathbb{R}^3$.

As a result of assumption (iii), testing (3.4) by \dot{y} and integrating over Ω , we obtain the energy equality as

$$\int_{\Omega} DW(\nabla y) : \nabla \dot{y} dx + \int_{\Omega} |\dot{C}|^2 dx + \frac{d}{dt} \int_{\Omega} \mathcal{H}(\nabla^2 y) dx = \int_{\Omega} f(t) \cdot \dot{y} dx.$$

Integrating this equality with respect to time and using (3.2), we obtain

$$E(y(T)) + \int_0^T \int_{\Omega} |\dot{C}|^2 dx dt = E(y(0)) + \int_0^T \int_{\Omega} f(t) \cdot \dot{y} dx dt. \quad (5.4)$$

This relation is vital for the *a priori* estimates we obtain for the solutions in the following sections.

(b) Time-discretization

In this section, we introduce the variational approach we adopt for the existence of weak solutions of the equation

$$\text{Div} (DW(\nabla y) + S(\nabla y, \nabla \dot{y}) - \text{Div} \mathcal{H}'(\nabla^2 y)) - f = 0, \quad (5.5)$$

and define approximate solutions to (5.5) by means of the following implicit time-discretization scheme. For a fixed time step size $\tau > 0$ and initial data $y_0 \in W^{1,2}(\Omega)$, we inductively define

$$\begin{aligned} y_{\tau}^0 &= y_0 \\ y_{\tau}^k &= \text{a minimizer of the functional } J_{\tau}^k(y) \quad (k \in \mathbb{N}), \end{aligned}$$

where

$$J_{\tau}^k(y) = \int_{\Omega} \left(W(\nabla y) + \tau \Psi \left(\nabla y_{\tau}^{k-1}, \frac{\nabla y - \nabla y_{\tau}^{k-1}}{\tau} \right) - \text{Div} \mathcal{H}'(\nabla^2 y) \right) dx - \int_{\Omega} f_{\tau}^k \cdot y dx, \quad (5.6)$$

where $f_{\tau}^k := \frac{1}{\tau} \int_{(k-1)\tau}^{k\tau} f(t) dt$. Note that for $k \geq 1$, the minimizers y_{τ}^k satisfy the Euler–Lagrange equations

$$\int_{\Omega} \left(DW(\nabla y_{\tau}^k) : \nabla z + \Psi_q \left(\nabla y_{\tau}^{k-1}, \frac{\nabla y_{\tau}^k - \nabla y_{\tau}^{k-1}}{\tau} \right) : \nabla z + D\mathcal{H}'(\nabla^2 y_{\tau}^k) : \nabla^2 z - f_{\tau}^k \cdot z \right) dx = 0, \quad (5.7)$$

for all $z \in C_0^{\infty}(\Omega)$, which represent a weak, time-discretized version of (5.5) for $\Psi_q(p, q) = S(p, q)$. For the last term in the integrand, we used Gateaux differentiability of the strain-gradient density (see [16]).

(c) Existence result

We can now state the result of the existence of minimizers for the discrete scheme.

Theorem 5.1. Let the assumptions (i), (ii) and (iii) be satisfied. Let $y_\tau^0 = y_0$ be as in (iv), $N \in \mathbb{N}$ and $\tau = T/N$. Then for $k = 1, 2, \dots, N$, y_τ^k can be found by solving the minimization problem (5.6) whose minimizers satisfy the time-discretized version of the problem (5.5) in the weak sense as in (5.7).

Proof. By assumptions (5.1) and (5.2), the minimization problem (5.6) is coercive on $W^{2,p}(\Omega, \mathbb{R}^3)$. By assumption (iii), we know that Ψ is given as in (4.7). By lemma 4.6, we obtain that

$$\Psi\left(\nabla y_\tau^{k-1}, \frac{\nabla y - \nabla y_\tau^{k-1}}{\tau}\right)$$

is convex in y . This implies lower semi-continuity in $W^{1,2}(\Omega, \mathbb{R}^3)$. As a result, we obtain a minimizer y_τ^k with $E(y_\tau^k) < \infty$. Hence, by proposition 5.1, we know that the minimizer satisfies $\det \nabla y(x) \geq \delta > 0$. So, y_τ^k satisfies the Euler–Lagrange equation (5.7), which completes the proof. ■

We now define piecewise constant interpolants using the discrete approximations y_τ^k for $k = 0, \dots, N$, which is standard in the convergence of discrete schemes. They are given as

$$y^\tau(x, t) := y_\tau^k(x) \quad \text{for } t \in ((k-1)\tau, k\tau). \quad (5.8)$$

Now, we prove the *a priori* estimates for the solutions.

Proposition 5.3. Let assumptions (i)–(iv) be satisfied. Also, assume that $\Psi(\cdot, 0) = 0$. Then, there exists a constant $C > 0$ such that the piecewise constant interpolants $y_\tau \in W^{2,p}$ given in (5.8) satisfy the following *a priori* estimates:

$$\begin{aligned} \|y_\tau\|_{L^\infty([0, T]; W^{2,p}(\Omega; \mathbb{R}^3)) \cap H^1([0, T]; H^1(\Omega; \mathbb{R}^3))} &\leq C \\ \det \nabla y_\tau &\geq 1/C. \end{aligned} \quad (5.9)$$

Proof. Substituting $y = y_\tau^{k-1}$ in (5.6) and using both the assumption that $\Psi(\cdot, 0) = 0$ and the fact that $y = y_\tau^k$ is a global minimum, we obtain

$$\begin{aligned} &\int_\Omega (W(\nabla y_\tau^k) - W(\nabla y_\tau^{k-1})) dx + \int_\Omega \tau \Psi\left(\nabla y_\tau^{k-1}, \frac{\nabla y_\tau^k - \nabla y_\tau^{k-1}}{\tau}\right) dx \\ &- \int_\Omega (\text{Div } \mathcal{H}(\nabla^2 y_\tau^k) - \text{Div } \mathcal{H}(\nabla^2 y_\tau^{k-1})) dx \leq \tau \int_\Omega f_\tau^k \cdot \left(\frac{y_\tau^k - y_\tau^{k-1}}{\tau}\right) dx. \end{aligned}$$

Using $\Psi \geq 0$ and

$$f_\tau^k \cdot \left(\frac{y_\tau^k - y_\tau^{k-1}}{\tau}\right) \leq \|f_\tau^k\|_{H^{-1}} \left\| \frac{y_\tau^k - y_\tau^{k-1}}{\tau} \right\|_{H^1} \leq \hat{C} \|f_\tau^k\|_{H^{-1}} \left\| \frac{\nabla y_\tau^k - \nabla y_\tau^{k-1}}{\tau} \right\|_{L^2},$$

we obtain the recursive estimate

$$E(y_\tau^k) - E(y_\tau^{k-1}) \leq \tau C_1 \|f_\tau^k\|_{H^{-1}}^2 + \tau C_2 \left\| \frac{\nabla y_\tau^k - \nabla y_\tau^{k-1}}{\tau} \right\|_{L^2}^2,$$

where \hat{C}, C_1 and C_2 are generic constants. Using a discrete Grönwall-type estimate for the total energy together with its definition (3.2), we obtain the desired estimate in $L^\infty([0, T]; W^{2,p}(\Omega; \mathbb{R}^3))$. Moreover, by assumption (iii), we know that Ψ is given as in (4.7). By invoking corollary 5.1, we obtain the desired estimate in $H^1([0, T]; H^1(\Omega; \mathbb{R}^3))$. ■

By this proposition, one can conclude that the constant interpolants defined in (5.8) satisfy a suitable discretized version of (5.5). Now we can prove convergence as $\tau \rightarrow 0$.

Theorem 5.2. Let the assumptions (i)–(iv) hold. Then, as $\tau \rightarrow 0$, there exists a limit function y such that

$$y_\tau \rightarrow y \text{ weakly}^* \text{ in } L^\infty([0, T]; W^{2,p}(\Omega; \mathbb{R}^3)) \cap H^1([0, T]; H^1(\Omega; \mathbb{R}^3)). \quad (5.10)$$

Also,

$$\nabla y_\tau \rightarrow \nabla y \text{ strongly in } L^\infty([0, T] \times \Omega; \mathbb{R}^{3 \times 3}), \quad (5.11)$$

and any such y is a weak solution to (5.5) complemented with the initial and boundary values as in (3.5) and (3.6). Moreover, it satisfies the energy balance (5.4).

Proof. By the *a priori* estimates obtained in proposition 5.3, we can extract a subsequence (which we do not relabel for simplicity) such that (5.10) holds. To obtain (5.11), we argue as follows (refer also to [16]). By the continuous embedding $W^{1,p}(\Omega) \subset C^\gamma(\Omega)$ with $\gamma = 1 - 3/p$, we have $\|\nabla y_\tau\|_{C^\gamma} \leq C$. Moreover, (5.9) gives the Hölder estimate

$$\|\nabla y_\tau(t_1) - \nabla y_\tau(t_2)\|_{L^2(\Omega; \mathbb{R}^3)} \leq C_1 |t_1 - t_2|^{1/2},$$

for all $t_1, t_2 \in [0, T]$. Using the interpolation $\|\cdot\|_{C^\beta} \leq C \|\cdot\|_{C^\gamma}^\alpha \|\cdot\|_2^{1-\alpha}$ and the *a priori* estimated in proposition 5.3, we can conclude that ∇y_τ is uniformly bounded in C^α , leading to the desired convergence (5.11) by an application of the Arzelà–Ascoli theorem. For the convergence in the energy balance, we use the form of S given in assumption (iii) and apply Minty’s trick to the strain-gradient part as a result of (5.11). ■

We can state the following corollary.

Corollary 5.2. *Considering time-discretization for $\dot{C} = (\nabla \dot{y})^T \nabla y + \nabla \dot{y}(\nabla y)^T$, where $C = F^T F$ is the right Cauchy–Green stretch tensor, and defining the appropriate constant interpolant C_τ , we obtain*

$$C_\tau \rightarrow C \text{ strongly in } H^1([0, T]; L^2(\Omega)),$$

where $C = (\nabla y)^T \nabla y$.

6. A counterexample

We prove the following result and the counterexample showing that the assumption of strong convergence is necessary for compactness (refer also to [10]).

Theorem 6.1. *Consider the sequence $\{y^{(j)}\}_{j=1}^\infty$ and assume that the following assumptions hold:*

- (i) $y^{(j)} \xrightarrow{*} y$ in $W^{1,\infty}(\Omega; \mathbb{R}^{3 \times 3})$ as $j \rightarrow \infty$
- (ii) $\det \nabla y, \det \nabla y^{(j)} > 0$ for a.e. $x \in \Omega$ and for all j ,
- (iii) $U^{(j)} = \sqrt{\nabla y^{(j)T} \nabla y^{(j)}} \rightarrow U$ for a.e. $x \in \Omega$ as $j \rightarrow \infty$.

Then, $\nabla y^T \nabla y = U^2$ holds.

Proof. Assumption (i) immediately gives, by Theorem 3.4 in [24], that

$$\det \nabla y^{(j)} \xrightarrow{*} \det \nabla y \text{ in } L^\infty(\Omega) \quad (6.1a)$$

$$\text{cof } \nabla y^{(j)} \xrightarrow{*} \text{cof } \nabla y \text{ in } L^\infty(\Omega). \quad (6.1b)$$

By assumption (ii) and polar decomposition theorem, we have $\nabla y^{(j)} = R^{(j)} U^{(j)}$, where $R^{(j)} \in \text{SO}(3)$ and $U^{(j)}$ is the right stretch tensor. This gives

$$\det \nabla y^{(j)} = \det R^{(j)} \det U^{(j)} = \det U^{(j)}.$$

Therefore, by assumption (iii), we obtain

$$\det \nabla y^{(j)} \rightarrow \det U \text{ for a.e. } x \in \Omega. \quad (6.2)$$

Convergences (6.1a) and (6.2) immediately give

$$\det \nabla y = \det U. \quad (6.3)$$

Similarly, we have

$$\operatorname{cof} \nabla y^{(j)} = \operatorname{cof} R^{(j)} \operatorname{cof} U^{(j)} = R^{(j)} \operatorname{cof} U^{(j)},$$

and hence, by (6.1b), we obtain

$$R^{(j)} \operatorname{cof} U^{(j)} \xrightarrow{*} \operatorname{cof} \nabla y \text{ in } L^\infty(\Omega).$$

This shows that $\operatorname{cof} U^{(j)}$ is uniformly bounded, which together with assumption (iii) gives

$$\operatorname{cof} U^{(j)} \rightarrow \operatorname{cof} U \text{ in } L^q(\Omega), \quad 1 \leq q < \infty. \quad (6.4)$$

Without loss of generality, we can say that $R^{(j)} \xrightarrow{*} R$ in $L^\infty(\Omega)$, which implies

$$R^{(j)} \rightharpoonup R \text{ in } L^p(\Omega), \quad 1 \leq p < \infty. \quad (6.5)$$

Choosing $q = p'$ in (6.4) thus gives

$$R^{(j)} \operatorname{cof} U^{(j)} \rightharpoonup R \operatorname{cof} U \text{ in } L^1(\Omega). \quad (6.6)$$

Convergences (6.1b) and (6.6) imply that

$$\operatorname{cof} \nabla y = R \operatorname{cof} U. \quad (6.7)$$

By (6.3) and (6.7) and the fact that $\operatorname{cof} F = (\det F) F^{-T}$, for any $F \in \operatorname{GL}^+(3)$, we obtain

$$(\det U) \nabla y^{-T} = (\det \nabla y) \nabla y^{-T} = \operatorname{cof} \nabla y = R \operatorname{cof} U = R (\det U) U^{-T},$$

giving

$$\nabla y^{-T} = R U^{-T}. \quad (6.8)$$

As we do not know whether R is a rotation and U is symmetric or not, (6.8) is still not enough. However, by boundedness of $\nabla y^{(j)}$ and $R^{(j)}$ and assumption (iii), we deduce that

$$U^{(j)} \rightarrow U \text{ in } L^r(\Omega), \quad 1 \leq r < \infty.$$

Choosing $p = r'$ in (6.5) thus gives

$$R^{(j)} U^{(j)} \rightharpoonup R U \text{ in } L^1(\Omega). \quad (6.9)$$

Assumption (i) and (6.9) imply that

$$\nabla y = R U. \quad (6.10)$$

Therefore, by (6.8) and (6.10), we have

$$\nabla y^T \nabla y = U^T R^{-1} R U = U^T U. \quad (6.11)$$

Equations (6.10) and (6.11) prove that $R \in \operatorname{SO}(3)$. Hence, by assumption (ii) and the polar decomposition theorem, we can conclude that U is symmetric, which, by (6.11), immediately gives the result. ■

We now state the following crucial remark, which shows that without the inclusion of the strain-gradient term in the model, we would not be able to obtain theorem 5.2.

Proposition 6.1. *If assumption (iii) is not satisfied, then the conclusion of theorem 6.1 does not hold.*

Proof. We give the following counterexample in order to prove the claim. Consider the simple laminate in a two-well problem as shown in Figure 1 formed from gradients A, B satisfying $A - B = a \otimes n$ with separating interfaces with normal n , the A layers having thickness λ/j and B layers $(1 - \lambda)/j$ for $0 < \lambda < 1$.

Let $A \in \operatorname{SO}(3) U_1, B \in \operatorname{SO}(3) U_2$ and choose




Figure 1. Laminate in a two-well problem. Here Dy stands for ∇y .

$$A = U_1 \quad \text{and} \quad B = U_1 + a \otimes n.$$

Then, $\nabla y^{(j)}$ satisfies (e.g. [17,25])

$$\nabla y^{(j)} \stackrel{*}{\rightharpoonup} \lambda U_1 + (1-\lambda)(U_1 + a \otimes n) = U_1 + (1-\lambda) a \otimes n =: \nabla y.$$

Therefore,

$$U = \sqrt{\nabla y^T \nabla y} = \sqrt{(U_1 + (1-\lambda) n \otimes a)(U_1 + (1-\lambda) a \otimes n)}.$$

On the other hand, we could also choose

$$U_A^{(j)} = U_1 \quad \text{and} \quad U_B^{(j)} = U_2,$$

where $U_A^{(j)} = U^{(j)}(x)|_{x \in A}$ and similarly for B . In this case, we would get

$$U^{(j)} \stackrel{*}{\rightharpoonup} \lambda U_1 + (1-\lambda)U_2 =: U.$$

However,

$$\lambda U_1 + (1-\lambda)U_2$$

and

$$\sqrt{(U_1 + (1-\lambda) n \otimes a)(U_1 + (1-\lambda) a \otimes n)}$$

are not necessarily equal, contradicting the conclusion of [theorem 6.1](#). ■

7. Conclusion

In this contribution, a very long-standing open problem of the well-posedness of nonlinear viscoelasticity of strain-rate type in high space dimensions while obeying the conditions of frame indifference is revisited. As a result of the adopted modelling postulates, including neglecting the inertia term as well as adding a higher-order regularity term for deformation, the existence of weak solutions is obtained as a result of the application of a time-discretization method together with a minimization argument. However, the problem of the existence of solutions for the fully dynamical case without assuming additional regularity of the deformation is still open. We hope that the observations made, however small, and the counterexample given will be able to shed some light on the discovery of new methods to be developed or new approaches to be adopted in order to tackle this problem.

Data accessibility. This article has no additional data.

Declaration of AI use. I have not used AI-assisted technologies in creating this article.

Authors' contributions. Y.Ş.: conceptualization, formal analysis, investigation, methodology, project administration, resources, writing—original draft, writing—review and editing.

Conflict of interests. I declare I have no competing interests.

Funding. No funding has been received for this article.

Acknowledgements. Y.Ş. thanks John M. Ball for his guidance and valuable contributions to the results obtained in this work.

References

- Şengül Y. 2021 Nonlinear viscoelasticity of strain rate type: an overview. *Proc. R. Soc. A.* **477**, 20200715. (doi:10.1098/rspa.2020.0715)
- Dafermos CM. 1969 The mixed initial-boundary value problem for the equations of nonlinear one-dimensional viscoelasticity. *J. Differ. Equ.* **6**, 71–86. (doi:10.1016/0022-0396(69)90118-1)
- Pego RL. 1987 Phase transitions in one-dimensional nonlinear viscoelasticity: admissibility and stability. *Arch. Ration. Mech. Anal.* **97**, 353–394. (doi:10.1007/BF00280411)
- Andrews G. 1980 On the existence of solutions to the equation $u_{tt} = u_{xxt} + \sigma(u_x)_x$. *J. Differ. Equ.* **35**, 200–231. (doi:10.1016/0022-0396(80)90040-6)
- Greenberg JM. 1969 On the existence, uniqueness, and stability of the equation $p_0 X_{Tt} = E(X_X)X_{Xx} + \Lambda X_{Xxt}$. *J. Math. Anal. Appl.* **25**, 575–591. (doi:10.1016/0022-247x(69)9025-1)
- Demoulini S. 2000 Weak solutions for a class of nonlinear systems of viscoelasticity. *Arch. Ration. Mech. Anal.* **155**, 299–334. (doi:10.1007/s002050000115)
- Tvedt B. 2008 Quasilinear equations for viscoelasticity of strain-rate type. *Arch. Ration. Mech. Anal.* **189**, 237–281. (doi:10.1007/s00205-007-0109-x)
- Friesecke G, Dolzmann G. 1997 Implicit time discretization and global existence for a quasilinear evolution equation with nonconvex energy. *SIAM. J. Math. Anal.* **28**, 363–380. (doi:10.1137/S0036141095285958)
- Watson SJ. 2000 Unique global solvability for initial-boundary value problems in one-dimensional nonlinear thermoviscoelasticity. *Arch. Ration. Mech. Anal.* **153**, 1–37. (doi:10.1007/s002050050007)
- Şengül Y. 2010 Well-Posedness of Dynamics of Microstructure in solids. PhD Thesis, University of Oxford, Oxford, UK.
- Ball JM, Şengül Y. 2015 Quasistatic nonlinear viscoelasticity and gradient flows. *J. Dyn. Diff. Equat.* **27**, 405–442. (doi:10.1007/s10884-014-9410-1)
- Mielke A, Ortner C, Şengül Y. 2014 An approach to nonlinear viscoelasticity via metric gradient flows. *SIAM. J. Math. Anal.* **46**, 1317–1347. (doi:10.1137/130927632)
- Friedrich M, Kružík M. 2018 On the passage from nonlinear to linearized viscoelasticity. *SIAM. J. Math. Anal.* **50**, 4426–4456. (doi:10.1137/17M1131428)
- Badal R, Friedrich M, Kružík M. 2023 Nonlinear and linearized models in thermoviscoelasticity. *Arch. Ration. Mech. Anal.* **247**, 5. (doi:10.1007/s00205-022-01834-9)
- Krömer S, Roubíček T. 2020 Quasistatic viscoelasticity with self-contact at large strains. *J. Elast.* **142**, 433–445. (doi:10.1007/s10659-020-09801-9)
- Mielke A, Roubíček T. 2020 Thermoviscoelasticity in Kelvin–Voigt rheology at large strains. *Arch. Ration. Mech. Anal.* **238**, 1–45. (doi:10.1007/s00205-020-01537-z)
- Ball JM. 1996 Some recent developments in nonlinear elasticity and its applications to material science. In *Nonlinear mathematics and its applications*, pp. 93–119. Cambridge, UK: Cambridge University Press.
- Ciarlet PG. 1987 *Mathematical elasticity: three-dimensional elasticity*. Amsterdam, the Netherlands: North-Holland.
- Ciarlet PG, Nečas J. 1985 Unilateral problems in nonlinear, three-dimensional elasticity. *Arch. Ration. Mech. Anal.* **87**, 319–338. (doi:10.1007/BF00250917)
- Truesdell C, Noll W. 2004 *The non-linear field theories of mechanics*. Berlin, Heidelberg, Germany: Springer-Verlag. (doi:10.1007/978-3-662-10388-3)
- Gurtin ME. 1981 *An introduction to continuum mechanics*. Oakland, CA: Academic Press.
- Antman SS. 2004 *Nonlinear problems of elasticity*. New York, NY: Springer.

23. Healey T, Krömer S. 2009 Injective weak solutions in second-gradient nonlinear elasticity. *ESAIM: COCV* **15**, 863–871. (doi:10.1051/cocv:2008050)
24. Ball JM, Currie JC, Olver PJ. 1981 Null Lagrangians, weak continuity, and variational problems of arbitrary order. *J. Funct. Anal.* **41**, 135–174. (doi:10.1016/0022-1236(81)90085-9)
25. Ball JM. 2004 Mathematical models of martensitic microstructure. *Mater. Sci. Eng. A.* **378**, 61–69. (doi:10.1016/j.msea.2003.11.055)