



Research



Gut microbial community in proboscis monkeys (*Nasalis larvatus*): implications for effects of geographical and social factors

Cite this article: Jose L, Lee W, Hanya G, Tuuga A, Goossens B, Tangah J, Matsuda I, Kumar VS. 2024 Gut microbial community in proboscis monkeys (*Nasalis larvatus*): implications for effects of geographical and social factors. *R. Soc. Open Sci.* **11**: 231756.

<https://doi.org/10.1098/rsos.231756>

Received: 16 November 2023

Accepted: 22 May 2024

Subject Category:

Ecology, conservation, and global change biology

Subject Areas:

ecology, microbiology

Keywords:

Borneo, colobines, foregut-fermenters, gut microbiota, primate

Authors for correspondence:

Ikki Matsuda

e-mail: matsuda.ikki.5a@kyoto-u.ac.jp

Vijay Subbiah Kumar

e-mail: vijay@ums.edu.my

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.7349579>.

Lilian Jose¹, Wanyi Lee^{2,3}, Goro Hanya³, Augustine Tuuga⁴, Benoit Goossens^{4,5,6}, Joseph Tangah⁷, Ikki Matsuda^{8,9,10,11} and Vijay Subbiah Kumar¹

¹Biotechnology Research Institute, Universiti Malaysia Sabah, Jalan UMS, Kota Kinabalu, Sabah 88400, Malaysia

²National Taiwan University, Taipei 10617, Taiwan

³Center for Ecological Research, Kyoto University, Inuyama 484-8506, Japan

⁴Sabah Wildlife Department, and ⁵Danau Girang Field Centre, Sabah Wildlife Department, Wisma Muis, Kota Kinabalu, Sabah 88100, Malaysia

⁶Organisms and Environment Division, Cardiff School of Biosciences, Cardiff University, Cardiff CF10 3AX, UK

⁷Sabah Forestry Department, Forest Research Centre, Sandakan, Sabah, Malaysia

⁸Wildlife Research Center of Kyoto University, 2-24 Tanaka-Sekiden-cho, Sakyo, Kyoto 606-8203, Japan

⁹Chubu Institute for Advanced Studies, Chubu University, 1200, Matsumoto-cho, Kasugai-shi, Aichi 487-8501, Japan

¹⁰Chubu University Academy of Emerging Sciences, 1200, Matsumoto-cho, Kasugai-shi, Aichi 487-8501, Japan

¹¹Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Jalan UMS, Kota Kinabalu, Sabah 88400, Malaysia

IM, 0000-0002-0861-7801; VSK, 0000-0002-0384-1580

Recent technological advances have enabled comprehensive analyses of the previously uncharacterized microbial community in the gastrointestinal tracts of numerous animal species; however, the gut microbiota of several species, such as the endangered proboscis monkey (*Nasalis larvatus*) examined in this study, remains poorly understood. Our study sought to establish the first comprehensive data on the gut microbiota of free-ranging foregut-fermenting proboscis monkeys and to determine how their microbiota are affected locally by environmental factors, i.e. geographical distance, and social factors, i.e. the number of adult females within harem groups and the number of adults and subadults within non-harem groups, in a riverine forest in Sabah, Malaysian Borneo. Using

16S rRNA gene sequencing of 264 faecal samples collected from free-ranging proboscis monkeys, we demonstrated the trend that their microbial community composition is not particularly distinctive compared with other foregut- and hindgut-fermenting primates. The microbial alpha diversity was higher in larger groups and individuals inhabiting diverse vegetation (i.e. presumed to have a diverse diet). For microbial beta diversity, some measures were significant, showing higher values with larger geographical distances between samples. These results suggest that social factors such as increased inter-individual interactions, which can occur with larger groups, as well as physical distances between individuals or differences in dietary patterns, may affect the gut microbial communities.

1. Introduction

The gastrointestinal tracts of animals are colonized by many microorganisms, forming complex microbial ecosystems [1–3]. In general, animal-associated microbial ecosystems have been reported to have a direct effect on host health, contributing not only to daily energy acquisition through the production of vitamins and short-chain fatty acids but also to the host's immune system and resistance to pathogens [4–6]. Understanding how the microbial community, which serves important functions in animals, is formed in the gastrointestinal tracts would shed light on the survival strategies of diverse animal species. Historically, the study of the human gastrointestinal microbial community has been extensive [7,8]. However, with recent advancements in sequencing technology, the ability to analyse gastrointestinal microbial diversity and community structure based on large amplicon libraries of 16S ribosomal RNA (rRNA) genes, primarily using faecal DNA, has prompted additional research in a variety of non-human primates [9–12]. With the recent accumulation of such research findings in non-human primates, it is becoming increasingly evident that their living environment influences and shapes the gastrointestinal microbial community [13]. For example, within the same primate species, a correlation has been observed between the diversity of dietary items and the microbial community [14–17]. Alternately, as in primate species that live in groups, it has been hypothesized that social factors may influence the establishment of the microbial community, as the horizontal transmission of the microbiota may occur via direct social interactions between individuals within the group or even indirect interaction via shared environments [18–21].

Although the gastrointestinal microbial community, particularly the gut microbiota from faecal samples, has been progressively studied in various primate species over the last decade [10,22–25], there are still numerous species in the wild for which even the most fundamental microbial community has not been studied. The foregut-fermenting proboscis monkey (*Nasalis larvatus*), a large, sexually dimorphic, arboreal primate [26], is one of these species for which the hindgut microbiota has not been studied in the wild (but see in captivity [27]), though comprehensive analyses of their foregut microbiota have been reported in the wild [15]. As both the foregut and hindgut have been reported to facilitate the digestive fermentation of dietary fibre in the closely related foregut-fermenting primate species, i.e. *Rhinopithecus roxellana* [23], a comprehensive analysis of hindgut bacteria based on faecal samples in proboscis monkeys is essential for fully understanding their digestive physiology.

Proboscis monkeys inhabit various riparian and coastal forest environments, including riverine, mangrove and peat swamp forests [28], and their dietary patterns have been reported to be flexibly adapted to these environments; diverse diets in riverine forests with higher plant diversity, and low dietary diversity in mangrove forests where plant diversity is extremely low [29–31]. Owing to its adaptable feeding habits, the proboscis monkey is an ideal study species for determining how habitat-specific dietary differences influence gut microbiota. The basic component of social structure in proboscis monkeys is as polygynous single male societies (harem group) that assemble with each other in the trees along rivers [32–35], though all-male groups and, occasionally, groups containing more than one male with multiple females are also found [36]. Consequently, it is a fascinating species with which to investigate the relationship between social factors such as different group types with different sexual compositions of members and the intestinal microbial community.

We sought to establish the first comprehensive data on the gut microbiota of free-ranging proboscis monkeys and to determine how these gut microbiota were affected at the local scale by environmental factors (e.g. geographical location) and social factors (e.g. sexes with different life histories and group size). In particular, we hypothesize that the observed richness and diversity of gut microbial

communities would vary based on the distance of their living areas from the river mouth in the study site and the number of adult females in harem groups. In addition, we propose that beta diversity measures of gut microbiota would correlate with geographical distance between individuals. Previous research has demonstrated that the gut microbiota is frequently influenced by dietary habits. Therefore, we anticipated that there would be differences between the gut microbiota of proboscis monkeys living near the river mouth, where deforestation by oil palm plantations is more pronounced, and those living in the upper river areas (see electronic supplementary material, S1), where relatively large areas of forest remain, at our study site [29]. Indeed, since increased dietary diversity has been reported to increase the diversity of the gut microbial community (e.g. [15,37,38]), it would be predicted that disturbed habitats would generally experience a reduction in plant diversity [39] and that the monkeys inhabiting these habitats would experience a reduction in dietary diversity. Furthermore, sex differences in social behaviour have been associated with sexual biases in the gut microbiota in primates (e.g. [19,40,41]), and female-to-female grooming is the predominant form of grooming in proboscis monkeys, but rarely between males and females [42,43]. Thus, female proboscis monkeys are expected to have more contact with more individuals in the group and, accordingly, alpha diversity and composition of their gut microbiota may be expected to be more diverse and/or of higher similarity than males. Finally, given that several reports in primates have stated that direct inter-individual contact is associated with the transmission of gut microbiota [18,21], it can be predicted that individuals with closer inter-individual distances would have more similar gut microbiota. Additionally, it is likely that individuals in larger groups would have more opportunities for social interaction with more individuals and, as a result, they may possess a more diverse gut microbiota.

2. Methods

2.1. Study site and subjects

The study was carried out in a riverine forest along the Menanggul River, a tributary of the Kinabatangan River, Sabah, Malaysian Borneo (118°30' E, 5°30' N), inhabited by eight species of diurnal primates, including our study species, the proboscis monkey. For more than a decade, this area has been a popular tourist destination that attracts boat tours; as a result, the proboscis monkeys were well habituated to human observers. The study site, in a 4 km stretch from the mouth of the Menanggul River upstream was home to at least 200 proboscis monkeys, organized into 8–10 harem groups with one adult male, multiple adult females and immatures [35,44], and various non-harem groups, including all-male groups (including solitary males), as well as other groups with multiple males and multiple females [36,45]. The southern portion of the Menanggul River is dominated by secondary forests, while the northern part has been cleared for oil palm plantations, excluding a protected zone along the river (electronic supplementary material, S1). Daily temperatures in the area were recorded at approximately 24°C (minimum) and 30°C (maximum), with an average annual rainfall of 2474 mm [29]. The river levels fluctuate by approximately 1 m daily, with seasonal floods causing an average increase of more than 3 m [46].

2.2. Faecal sampling

Proboscis monkeys in the lower Kinabatangan floodplain typically prefer to sleep along the river [47]. Therefore, we conducted a boat survey in the late afternoon to detect proboscis monkeys and record their group composition with GPS coordinates of their sleeping trees. In the early hours of the following morning, while the monkeys were still sleeping, we revisited their sleeping trees. As proboscis monkeys typically defecate shortly before moving into the forest, we carefully searched the ground near their sleeping trees to collect fresh faeces after they had left the sleeping trees [35]. Several harem groups often stayed in close proximity to the trees along the river, and it was sometimes difficult to determine which group the faeces on the forest floor belonged to, but based on the location of the faeces and the location of the group identified during the boat survey on the previous day, we inferred the group to which the individual that had defecated belonged. We only focused on collecting faecal samples presumed to be from adult individuals [48], and between June 2015 and April 2016, a total of 307 samples were opportunistically collected. However, owing to the nature of the faeces collection procedure, it was unclear how many groups in the study area the faeces were collected from. The collection was always carried out immediately after defecation, using a sterilized plastic spoon

attached to the sampling tube. The spoons were inserted into fresh faeces, and only a small amount of the interior was removed and stored in 5 ml lysis buffer (0.5% sodium dodecyl sulfate, 100 mM ethylenediaminetetraacetic acid (pH 8.0), 100 mM Tris-HCl (pH 8.0) and 10 mM NaCl [16] at room temperature. To prevent duplicate analyses after genetic profiling, faeces from the same individual and those samples of unknown sex were excluded, and thus, the analysis of gut microbiota was performed with 264 faecal samples, i.e. 187 females and 77 males [35].

2.3. DNA purification, 16S ribosomal RNA amplification and sequencing

After bead-beating using the bead crusher (TAITEC, μ T-01, Japan) and centrifuged at 4200 r.p.m. for 5 min, 200 μ l of lysis buffer-faecal sample mixture was added with 800 μ l of InhibitEX buffer of the QIAamp DNA Stool Mini Kit (Qiagen GmbH, Hilden, Germany). Next, the mixture was centrifuged at room temperature for 1 min at 13 000 r.p.m. Then, the lysate was transferred to a new 1.5 ml microcentrifuge tube with 25 μ l of proteinase K. This was followed by adding 500 μ l of Buffer AL and the manufacturer's protocols to purify the faecal DNA. Next, the DNA concentration was estimated with a Qubit dsDNA HS Assay Kit and a Qubit fluorometer (Thermo Fisher Scientific). We amplified the V3–V4 region of the 16S ribosomal RNA (rRNA) gene using the primer utilized in [49] with slight modification as follows: S-D-Bact-0341-b-S-17, 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-NS-CCTACGGGNGGCW G-3', and S-D-Bact-0785-a-A-21, 5'-GTCTCGTGGGCTCGGAGATGTGTA-TAAGAGACAG-NS-GACTACHVG GG -3'. Subsequently, 3Ns, 4Ns, 5Ns and 6Ns were inserted in each primer between the specific primer and the adapter to cause an artificial frameshift and improve the sequencing quality [48]. The mixture of these primers was used as forward and reverse primers at a concentration of 1 μ M. KAPA Pure Beads (KAPA Biosystems, Wilmington, MA) were used to purify the polymerase chain reaction amplicons. The Illumina Nextera XT Index Kit (Illumina, Inc., San Diego, CA) was then used to attach specific dual indices and sequencing adapters to the amplicons for each sample. The resulting products were then combined in equal DNA concentrations to form a pooled sequencing library. Subsequently, the size distribution of the library was then estimated using an Agilent 2100 Bioanalyzer (Agilent Technologies, Inc., La Jolla, CA). The library was then diluted to a concentration of 15 pM and sequenced with a 15% PhiX spike-in on an Illumina MiSeq sequencing platform using the MiSeq Reagent Kit v. 3 (600 cycles) (Illumina, Inc., San Diego, CA). The resulting read lengths were 301 bp (forward sequences), 8 bp (forward indices), 8 bp (reverse indices) and 301 bp (reverse sequences).

2.4. Data analysis

2.4.1. Amplicon sequence variants picking and taxonomic identification

The demultiplexed sequences were processed using QIIME2 software [50]. Amplicon sequence variants (ASVs) were generated using the DADA2 pipeline in this software through the *dada2* plugin. In this step, the forward and reverse reads were merged. The unmerged reads were also discarded and the chimera was removed. The parameters used were to exclude demultiplexed sequences with a quality score below 30 from the downstream analysis to ensure high-quality data. The spurious ASVs were also removed by the QIIME2 *dada2* plugin. The ASVs were assigned through the ribosomal database project classifier with GreenGenes v. 13_8 as the reference database for taxonomic identification. Additionally, we used the built-in function *align-to-tree-might-fast* of QIIME2 to construct a phylogenetic tree of the ASVs.

2.4.2. Statistical analysis

Statistical analyses were performed in R v. 4.1.1 [51], with the significance level set at 0.05 using the unrarefied dataset with the singletons being removed. Results were reported as means with standard deviation. Alpha and beta diversity were calculated using the R package *phyloseq* [52]. The Kruskal–Wallis test from the R package *dunn.test* was used to analyse the differences in alpha diversity between sexes and group types. Subsequently, we investigated the effects of social and geographical factors on the microbial alpha diversity of individual samples using a linear model. The alpha diversity of each sample was treated as a normally distributed response variable, while the social factor, i.e. the number of adult females for harem groups and the number of adults and subadults for non-harem

groups, and the geographical factor, i.e. the location of faecal samples collected along the river, which represented as the distance from the river mouth, were treated as explanatory variables. For all models, we verified that the variance inflation factors were smaller than the cut-off value, i.e. less than 10 [53]. Therefore, the collinearity between independent factors (explanatory variables) did not affect the results. For the model selection, the possible combinations of the explanatory variables were examined and ranked using the Akaike information criterion (AIC) from the *MuMIn* package [54]. Following the published guidelines for wildlife research, we generally selected the best-supported models as those with a ΔAIC score of less than 2, where $\Delta\text{AIC} = \text{AIC} - \text{minimum AIC}$ within the candidate models [55], but as suggested by Burnham *et al.* [56] the model with ΔAIC of range from 2 to 7 should not be dismissed; we discussed models that were in those ranges as well.

In our multivariate analysis of microbiota composition, we calculated Bray–Curtis dissimilarity, along with weighted and unweighted UniFrac indices with the R package *vegan*. We conducted the permutational multivariate analysis of variance (PERMANOVA, *adonis2* function in the *vegan* package, R software, v. 4.1.1) tests to estimate the differences in terms of beta diversity. We constructed non-metric multi-dimensional scaling (NMDS) for visualization using Bray–Curtis dissimilarity and principal coordinate analysis (PCoA) plots by weighted and unweighted UniFrac indices with the plot ordination function from package *ggplot2*. In order to further assess the effect of social and geographical factors and to test their correlation with microbial beta diversity, Mantel tests were conducted by using the Mantel function. This function takes three parameters, where the first one is the first distance matrix, followed by the second distance matrix, and the last one is ‘spearman’ which indicates the method that we used to compute the correlation between these two distance matrices. The social factor, known as ‘demographic distance’, was calculated based on the composition of the differences in the group and the number of individuals within the groups, while the geographical factor, known as ‘geographical distances’, was calculated based on the distance between different faecal samples (or individuals). In other words, with respect to these two distances, values are higher with more differences in the pattern of group composition within a group (i.e. differences in the number of males and females) and with greater distances between different faecal samples (or individuals).

3. Results

3.1. Phylogenetic profile of the faecal microbiota

In 264 faecal samples, we detected 16 530 ASV, classified into 29 phyla, 67 classes, 95 orders, 116 families, 134 genera and 50 species. The average number of sequences resulted per sample before filtering through DADA2 pipeline was 124 898, with a maximum of 287 114 and a minimum of 37 712. After filtering, the average was 100 900, with a maximum of 257 018 and a minimum of 10 234.

Figure 1 depicts the general distribution of the top five taxa at the phylum, family and genus levels. At the phylum level, the top five taxa were consistent across sexes; Bacillota dominated the gut microbial community (male: 81.9%; female: 82.3%), followed by Bacteroidota (male: 8.3%; female: 8.0%), Cyanobacteria (male: 1.5%; female: 1.8%), Pseudomonadota (male: 1.5%; female: 1.3%) and Actinomycetota (male: 1.1%; female: 1.1%) (see electronic supplementary material, S2 for details). At the family and genus levels, the top five patterns of the gut microbial community in both sexes were also consistent (see electronic supplementary material, S3 to S4 for details), i.e. family level: Ruminococcaceae (male: 46.3%; female: 44.6%), Lachnospiraceae (male: 14.7%; female: 15.3%), S24-7 (male: 6.8%; female: 6.5%), Christensenellaceae (male: 2.0%; female: 2.2%) and Mogibacteriaceae (male: 1.9%; female: 1.8%); genus level: *Oscillospira* (male: 10.4%; female: 10.2%), *Ruminococcus* (male: 4.7%; female: 4.5%), *Dorea* (male: 2.3%; female: 2.1%) and *Blautia* (male: 1.2%; female: 1.4%). In addition, we found that the pattern of the top five taxa (at the phylum, family and genus levels) was consistent across different group types (harem and non-harem groups) (see electronic supplementary material, S5 to S7). Finally, despite the information available on annotated bacterial taxa at the phylum, family and genus levels, it would be worth noting the proportion of unassigned taxa, which were 3.18–3.38%, 16.02–17.14% and 71.82–72.62%, respectively.




Figure 1. The general pattern of the top five taxa. These were analysed at the phylum (a), family (b) and genus (c) levels.

3.2. Gut microbial diversity

3.2.1. Alpha diversity

The mean observed richness and Shannon diversity index (H') of the gut microbiota in all faecal samples were 400.6 ± 51.5 (range: 161–661) and 5.0 ± 0.18 (range: 4.17–5.48), respectively. There were no significant differences in observed richness between sexes (Kruskal–Wallis $\chi^2 = 2.77$, d.f. = 1, $p = 0.10$; figure 2a), nor between the group types (harem and non-harem groups; Kruskal–Wallis $\chi^2 = 2.97$, d.f. = 1, $p = 0.08$; figure 2a). Shannon diversity index did not differ significantly by sex (Kruskal–Wallis $\chi^2 = 1.81$, d.f. = 1, $p = 0.18$; figure 2b) or group type (Kruskal–Wallis $\chi^2 = 0.25$, d.f. = 1, $p = 0.62$).

The best-fit model to explain the observed richness, as determined by AIC, included both the number of adult females in the harem groups and the location of the collected samples represented as the geographical distance from the mouth of the river (table 1 and figure 3a), although the Δ AIC value of the following model, which included only the number of adult females, was also less than 7.0. The observed richness increased as the number of adult females in the harem groups and the distance from the river mouth increased (upper river). The linear models for Shannon diversity index revealed a similar pattern to the observed richness in harem groups, with a positive effect on the number of adult females and geographical distance (table 1 and figure 3b).

Within the non-harem groups, however, the selection of the model for the investigation of whether the observed richness and Shannon diversity index were affected by social and geographical factors, i.e. the number of individuals, including adult males and females and subadult males, and the location of the collected samples, represented as the geographical distance from the river mouth, indicated that the null model was the best (table 2). The second- and third-best models, with observed richness and Shannon diversity index, incorporated social and geographical factors with Δ AIC values below the




Figure 2. Comparison of the mean observed richness (a) and the Shannon diversity index (b) for different sexes and different group types. The central line of the box represents the median, and the lower and upper bounds of the box represent the first and third quartiles.

cut-off of 7.0. This result indicates that a similar pattern was generally observed in non-harem groups, albeit with weaker effects than in harem groups.

3.2.2. Beta diversity

According to PCoA using weighted and unweighted UniFrac and NMDS plots with Bray-Curtis dissimilarity, individuals of different sexes or group types did not exhibit visually distinguishable patterns (figure 4). Conversely, PERMANOVA analysis revealed weak but significant differences between both sexes (PERMANOVA, Bray-Curtis, $R^2 = 0.0049$, $p = 0.021$; unweighted UniFrac, $R^2 = 0.0045$, $p = 0.009$) and between group types (PERMANOVA, Bray-Curtis, $R^2 = 0.0053$, $p = 0.004$; unweighted UniFrac, $R^2 = 0.0045$, $p = 0.01$) on the gut microbiota.

In order to examine the correlation between beta diversity measures and the two factors, i.e. geographical distances and social factors, the Mantel test was conducted separately on a dataset containing only samples from harem groups and one containing only samples from non-harem groups. Mantel tests performed with different beta diversity measures revealed significant positive correlations for the harem group between weighted UniFrac and geographical distances from different individuals ($r = 0.09$, $p = 0.01$) but not between unweighted UniFrac and geographical distances from different individuals ($r = 0.02$, $p = 0.21$). This indicated that individuals living further apart had a gut microbial community that was less similar, although the explanatory power of geographic distance was relatively weak. In contrast, Bray-Curtis dissimilarity measures of diversity, which do not account for the




Figure 3. Relationship of alpha diversity index, i.e. the observed richness (a) and Shannon diversity index (b), with the number of adult females within the harem groups and the location of faecal samples collected representing the geographical distance from the river mouth, based on the model selection in table 1.

Table 1. Summary of model selection for harem groups using linear models. This method was used to investigate whether observed richness (A) and Shannon diversity index (B) were affected by social factors, i.e. the number of adult females within harem groups, and the geographical factor, i.e. the location of faecal samples collected along the river, which represented the distance from the river mouth.

intercept	adult female	distance from river mouth	d.f.	log-likelihood	AIC	Δ AIC	weight
(A)							
351.5	5.758	0.005165	4	-902.163	1812.3	0.00	0.408
363.0	5.691		3	-903.226	1812.5	0.13	0.383
396.9			2	-905.517	1815.0	2.71	0.105
386.1		0.005031	3	-904.535	1815.1	2.74	0.103
(B)							
4.899		2.721×10^{-5}	3	72.000	-138.0	0.00	0.463
4.834	0.01087	2.746×10^{-5}	4	72.715	-137.4	0.57	0.348
4.958			2	69.568	-135.1	2.86	0.111
4.895	0.01052		3	70.219	-134.4	3.56	0.078




Figure 4. Unweighted UniFrac (a) and weighted UniFrac (b) principal coordinate analysis (PCoA) plots and Bray–Curtis non-metric multi-dimensional scaling (NMDS) plot (c), analysed by group type, i.e. harem and non-harem, and sex, i.e. F: female; M: male.

phylogeny of microbial composition, did not exhibit a significant correlation with geographic distance ($r = 0.05$, $p = 0.09$). In addition, there was no correlation between the different beta diversity measures and the distance of group composition between the various harem groups (weighted UniFrac: $r = 0.10$, $p = 0.09$; unweighted UniFrac: $r = 0.02$, $p = 0.23$; Bray–Curtis: $r = 0.002$, $p = 0.48$).

For samples from non-harem groups, weighted UniFrac ($r = 0.13$, $p = 0.02$) and Bray–Curtis dissimilarity ($r = 0.23$, $p = 0.001$) were significantly correlated with geographical distance, whereas there was no significant correlation between unweighted UniFrac and geographical distance ($r = 0.03$, $p = 0.19$). All the beta dissimilarity indices exhibited a significant correlation with the similarity distance of the group composition between different groups (weighted UniFrac: $r = 0.17$, $p = 0.03$; unweighted UniFrac: $r = 0.10$, $p = 0.04$; Bray–Curtis: $r = 0.13$, $p = 0.04$).

Table 2. Summary of model selection for non-harem groups using linear models. This method was used to investigate whether observed richness (A) and Shannon diversity index (B) were affected by social factors, i.e. the number of individuals, including adult males and females and subadult males within groups, and geographical factors, i.e. the location of faecal samples collected along the river, which represented the distance from the river mouth.

intercept	number of adults and subadults	distance from river mouth	d.f.	log-likelihood	AIC	Δ AIC	weight
(A)							
407.6			2	-504.772	1013.5	0.00	0.525
405.2		0.001834	3	-504.716	1015.4	1.89	0.204
401.7	1.185		3	-504.763	1015.5	1.98	0.195
396.7	1.678	0.001987	4	-504.699	1017.4	3.85	0.076
(B)							
4.969			2	18.998	-34.0	0.00	0.503
4.954		1.146×10^{-5}	3	19.190	-32.4	1.62	0.224
4.946	0.004727		3	19.010	-32.0	1.98	0.187
4.915	0.007743	1.217×10^{-5}	4	19.222	-30.4	3.55	0.085

4. Discussion

4.1. Overall gut microbial composition

The present study was the first comprehensive analysis of the intestinal microbial community based on faeces from more than 250 free-ranging proboscis monkeys. At the phylum level, the composition of their gut microbial community was nearly identical to that of other foregut-fermenting primates. Bacillota and Bacteroidota were predominant in *Colobus polykomos*, *Procolobus badius*, *P. verus*, *Rhinopithecus brelichi* and *R. roxellana* [12,57–59], with the exception of *Pygathrix nemaeus*, which includes Bacteroidota in the top five, but Bacillota and Mycoplasmatota dominate the top two phyla [38]. This trend is not exclusive to foregut-fermenters but is also observed in hindgut-fermenters, such as *Cercocebus atys*, *Cercopithecus campbelli*, *C. diana*, *C. petaurista*, *Chlorocebus sabaeus*, *Lemur catta*, *Macaca fuscata*, *M. mulatta*, *M. thibetana*, *Pan troglodytes*, *Propithecus verreauxi* and *Theropithecus gelada* [12,57,60–64].

At the family level, the predominance of Ruminococcaceae or Lachnospiraceae is comparable to that of the foregut-fermenting *R. brelichi* and *R. roxellanae*. *Macaca fuscata*, a hindgut-fermenting primate, exhibited a similar trend [60], and Ruminococcaceae were frequently the most dominant in other hindgut-fermenting primates [12,61,62]. Although the proportion taxonomically assigned at the genus level was limited, the top two genera, *Oscillospira* and *Ruminococcus*, are listed among the top five in *P. nemaeus*, a foregut-fermenter as well as in the proboscis monkey. *Ruminococcus*, in particular, is a common genus, as it frequently ranks first not only in foregut-fermenters but also in hindgut-fermenters [60,62]. However, *Blautia* is not among the top genera in *P. nemaeus*, but it is among the top genera [60,62] in the hindgut fermenters.

In the light of these findings, the composition of the gut microbial community of the proboscis monkey is unlikely to differ significantly from that of other foregut- and hindgut-fermenting primates. However, when the microbiota is subdivided from the phylum level to the genus level, the degree of overlap obviously decreases, and species-specific microbial composition and bacterial species are found probably owing to more species-specific digestive physiology, dietary patterns related to living environment and/or, sociality of host animals, e.g. lactic acid bacteria, *Lactobacillus nasalidis*, specific only to the proboscis monkey [65,66].

4.2. Microbial patterns in relation to social factors

There were no differences in the alpha diversity index (number of ASV and Shannon diversity index) of the proboscis monkey gut microbiota between sexes or group types, indicating that alpha diversity is not affected by differences in social factors such as sex differences in life history and/or the social

composition of individuals. Conversely, significant differences in beta diversity were observed between the sexes, suggesting that differences in life history and the frequency of social interactions may have influenced the composition of the gut microbiota. Significant differences in beta diversity between group types may also reflect differences of life history in the sexes; the basis for the differences in gut microbiota between group types with different male and female compositions within a group may be related to the differences in such life histories. At maturity, females transfer from their natal harem groups to other harem groups, whereas males disperse from their natal harem groups in the early stages to join all-male groups [32,42,67,68]. Additionally, grooming within harem groups occurs primarily between females, with males rarely participating [43,69]. Consequently, it is likely that these differences in the life histories of the sexes and in the frequency of social interactions between the sexes influence the composition of the gut microbiota in proboscis monkeys. Several other non-human primates have been reported to exhibit sexual biases in the gut microbiota, which could be attributed to differences in such interactions and life histories between sexes as a result of group living, e.g. *Callithrix jacchus* [40], *Rhinopithecus bieti* [41] and *Alouatta pigra* [19], though such sexual differences are little in some species, e.g. *Propithecus verreauxi* [63]. In contrast, it is difficult to determine its ecological significance without knowing how differences in microbiota composition affect factors such as food digestion and immunity. As expected, the positive correlation between the number of females in harem groups and the alpha diversity index suggests that increased individual interactions result in an increase in alpha diversity. This result is consistent with previous research indicating that direct physical contact between social partners is a major factor in the transmission of gut microbiota [19,21,40].

4.3. Microbial patterns in relation to geographical factors

In addition to the number of females within the harem groups, individuals who resided in areas further upstream of the river mouth at our study site had higher alpha diversity indices. This may be owing to differences in the food diversity consumed by proboscis monkeys in upstream and downstream regions. In general, individuals with a more varied diet have a greater variety of symbiotic bacteria in their gastrointestinal tracts [15,37,38]. Indeed, given that the downstream area of the study site is more heavily affected by deforestation [29] and has often lower plant diversity with lower potential food sources for proboscis monkeys, especially on the north side of the river (electronic supplementary material, S9), individuals in the upstream area may have had access to a greater variety of food sources, resulting in a tendency for a higher alpha diversity index in their gut microbiota. Nevertheless, to strengthen this conclusion, differences in vegetation between upstream and downstream areas would require a more quantitative investigation and comparison with the degree of gut microbial diversity obtained in the present study.

In the non-harem groups, a similar relationship was observed between alpha diversity and the number of individuals in the group and the geographical conditions they inhabited, although this trend was weaker with insignificance. This difference may be attributable to the fact that non-harem groups, particularly all-male groups, exploit a broader range of riparian habitats than harem groups [35], which not only facilitate a more diverse diet but also direct/indirect interactions with conspecifics, coupled with increased such interactions with various organisms in the forest elsewhere, possibly leading to the horizontal transmission of the microbiota [70,71]. However, not only the social relationships between individuals within non-harem groups but also those with various organisms remain unclear, making further discussion impossible at this time. The key to advancing this discussion in the future would be the collection of additional ecological and social observation data on non-harem groups.

There was a weaker but significant correlation between the physical distance between individuals and their similarity in the composition of their intestinal microbial communities, regardless of the group type. It should be noted, however, that the significance of the correlation varied slightly between the various group types based on the various indices, namely weighted UniFrac, unweighted UniFrac and Bray–Curtis. As previously mentioned, differences in vegetation between upstream and downstream areas of this study site may influence the alpha diversity of individual proboscis monkeys; it cannot be ruled out that the physical distance between individuals may lead to differences in diets consumed, which may, in turn, lead to differences in the composition of the intestinal microbial community between individuals. However, no correlation was found between similarities in group composition and those in gut microbial community composition. Compared with seasonal and clumped food sources, such as fruits and flowers, there was a higher ubiquitous and abundant

availability of leaves as the main food source for proboscis monkeys in this study site [31,72] and the lack of a clear hierarchy between individuals, which occurs particularly within harem groups [42]. This may have contributed to the lower likelihood of dietary bias across group composition in proboscis monkeys, and hence, there might be no differences in the similarity of the composition of gut microbial community between individuals. Conversely, there have been no studies on the hierarchy among males in non-harem groups, particularly in all-male groups, and the finding that significant differences were observed in non-harem groups by all the beta dissimilarity indices may suggest that there may be a severe hierarchy among males, which resulted in differences in their dietary composition. To verify this, however, more comprehensive behavioural observations of all-male groups are required.

4.4. Outlook

We succeeded not only in determining the general trends in the intestinal microbiota of proboscis monkeys but also in determining how social and geographical factors affected this microbial community. In contrast, while we characterized such differences in the gut microbial community at the local level among individuals, we have not been able to elucidate how these differences act as an advantage to the survival of each individual. Future research must examine how differences in the gut microbial community between individuals influence their feeding strategy through more detailed functional analysis, such as by isolating and cultivating characteristic strains [66]. Furthermore, considering that the gut microbiota also changes with diet in primates (e.g. [60,62,73,74]), there remains a need to compare the dietary patterns and gut microbiota of the individuals in a real-time manner, which we could not demonstrate in the current study, which was a limitation. When analysing gut microbiota from faeces, only a portion of the faeces is used for that analysis, and therefore many other portions are discarded without being used. However, with recent metabarcoding techniques [75–77], it may be possible to estimate dietary patterns from that remaining faecal sample. Hence, the analysis of inter-individual differences in the gut microbial community related to the different dietary patterns of each individual would be relatively feasible in the future study. Finally, the present study suggested that the degree of deforestation may influence the gut microbiota, although quantitative comparisons of these effects were not possible. While traditional vegetation surveys are often beneficial to understanding the relationship between environment and animal ecology [39,78–80], advanced technologies such as the recently developing LiDAR technology [81] may allow more extensive and comprehensive habitat assessments which, when coupled with the analysis of gut microbial data of target animals as in the present study, may reveal more reliable interactions between their living environment and gut microbiota. In sum, with advancements in technology for the analysis of gut microbiota, it may be possible in the future to transition from non-invasively collected free-ranging primate faeces to studies that contribute to animal conservation, such as assessing the effects of forest disturbance. Multiple studies have demonstrated their potential use in animal conservation [15,17,37,82,83].

Ethics. This study was reviewed and approved by the Sabah Biodiversity Council (Access License Reference: JKM/MBS.1000-2/2 JLD.5) and was conducted in compliance with animal care regulations applicable to Malaysian laws. This study is reported in accordance with ARRIVE guidelines (<https://arriveguidelines.org>).

Data accessibility. The original contributions presented in the study are included in the article/supplementary material [84], further inquiries can be directed to the corresponding author/s. The sequencing data have been deposited to NCBI under PRJNA928786.

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. L.J.: data curation, formal analysis, investigation, methodology, visualization, writing—original draft; W.L.: methodology, writing—review and editing; G.H.: methodology, resources, writing—review and editing; A.T.: resources, writing—review and editing; B.G.: methodology, resources, writing—review and editing; J.T.: resources, writing—review and editing; I.M.: conceptualization, data curation, funding acquisition, investigation, methodology, project administration, resources, supervision, validation, visualization, writing—original draft; V.S.K.: conceptualization, methodology, project administration, resources, supervision, validation, visualization, writing—original draft.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This study was partially funded by Japan Science and Technology Agency Core Research for Evolutional Science and Technology 17941861 (no. JPMJCR17A4), Japan Society for the Promotion of Science KAKENHI (nos. 26711027, 15K14605, 19H03308 and 24H00774 to I.M.; no. 23H02563 to G.H.) and Universiti Malaysia Sabah UMSGreat Grant (no. GUG0419-2/2019).

Acknowledgements. We express our sincere thanks to the Sabah Biodiversity Centre and the Sabah Wildlife Department for granting permission to carry out this research. We are also deeply indebted to the Sabah Forestry Department for facilitating the use of their facilities in the field. We are grateful to the researchers at the Biotechnology Research Institute (UMS) and the Inuyama Campus of Kyoto University for the knowledge transfer along the project. L.J. and V.S.K. thank Universiti Malaysia Sabah for managing funds (grant nos. LPA2001 and LPA2101). I.M. thanks the research assistants, especially Asnih Binti Etin and Jasrudy Bin Mandu, for their support in the field.

References

- Ley RE *et al.* 2008 Evolution of mammals and their gut microbes. *Science* **320**, 1647–1651. (doi:10.1126/science.1155725)
- Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI. 2007 The human microbiome project. *Nat. New Biol.* **449**, 804–810. (doi:10.1038/nature06244)
- Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, Henrissat B, Knight R, Gordon JI. 2011 Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science* **332**, 970–974. (doi:10.1126/science.1198719)
- Bergman EN. 1990 Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol. Rev.* **70**, 567–590. (doi:10.1152/physrev.1990.70.2.567)
- Hooper LV, Littman DR, Macpherson AJ. 2012 Interactions between the microbiota and the immune system. *Science* **336**, 1268–1273. (doi:10.1126/science.1223490)
- Buffie CG, Pamer EG. 2013 Microbiota-mediated colonization resistance against intestinal pathogens. *Nat. Rev. Immunol.* **13**, 790–801. (doi:10.1038/nri3535)
- Dave M, Higgins PD, Middha S, Rioux KP. 2012 The human gut microbiome: current knowledge, challenges, and future directions. *Transl. Res.* **160**, 246–257. (doi:10.1016/j.trsl.2012.05.003)
- The Integrative HMP (iHMP) Research Network Consortium. 2019 The integrative human microbiome project. *Nature New Biol.* **569**, 641–648. (doi:10.1038/s41586-019-1238-8)
- Björk JR, Dasari M, Grieneisen L, Archie EA. 2019 Primate microbiomes over time: longitudinal answers to standing questions in microbiome research. *Am. J. Primatol.* **81**, e22970. (doi:10.1002/ajp.22970)
- Amato KR *et al.* 2019 Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut microbiomes. *ISME J.* **13**, 576–587. (doi:10.1038/s41396-018-0175-0)
- Clayton JB *et al.* 2018 The gut microbiome of nonhuman primates: lessons in ecology and evolution. *Am. J. Primatol.* **80**, e22867. (doi:10.1002/ajp.22867)
- Zeng Y *et al.* 2022 Comparison of gastrointestinal microbiota in golden snub-nosed monkey (*Rhinopithecus roxellanae*), green monkey (*Chlorocebus aethiops sabaues*), and ring-tailed lemur (*Lemur catta*) by high throughput sequencing. *Glob. Ecol. Conserv.* **33**, e01946. (doi:10.1016/j.gecco.2021.e01946)
- Sharma AK *et al.* 2022 The primate gut mycobiome-bacteriome interface is impacted by environmental and subsistence factors. *NPJ. Biofilms. Microbiomes.* **8**, 12. (doi:10.1038/s41522-022-00274-3)
- Clayton JB *et al.* 2016 Captivity humanizes the primate microbiome. *Proc. Natl Acad. Sci. USA* **113**, 10376–10381. (doi:10.1073/pnas.1521835113)
- Hayakawa T *et al.* 2018 First report of foregut microbial community in proboscis monkeys: are diverse forests a reservoir for diverse microbiomes. *Environ. Microbiol. Rep.* **10**, 655–662. (doi:10.1111/1758-2229.12677)
- Lee W, Hayakawa T, Kiyono M, Yamabata N, Hanya G. 2019 Gut microbiota composition of Japanese macaques associates with extent of human encroachment. *Am. J. Primatol.* **81**, e23072. (doi:10.1002/ajp.23072)
- McManus N, Holmes SM, Louis EE, Johnson SE, Baden AL, Amato KR. 2021 The gut microbiome as an indicator of habitat disturbance in a critically endangered lemur. *BMC Ecol. Evol.* **21**, 222. (doi:10.1186/s12862-021-01945-z)
- Wikberg EC, Christie D, Sicotte P, Ting N. 2020 Interactions between social groups of colobus monkeys (*Colobus vellerosus*) explain similarities in their gut microbiomes. *Anim. Behav.* **163**, 17–31. (doi:10.1016/j.anbehav.2020.02.011)
- Amato KR, Van Belle S, Di Fiore A, Estrada A, Stumpf R, White B, Nelson KE, Knight R, Leigh SR. 2017 Patterns in gut microbiota similarity associated with degree of sociality among sex classes of a Neotropical primate. *Microb. Ecol.* **74**, 250–258. (doi:10.1007/s00248-017-0938-6)
- Goodfellow CK, Whitney T, Christie DM, Sicotte P, Wikberg EC, Ting N. 2019 Divergence in gut microbial communities mirrors a social group fission event in a black-and-white colobus monkey (*Colobus vellerosus*). *Am. J. Primatol.* **81**, e22966. (doi:10.1002/ajp.22966)
- Tung J *et al.* 2015 Social networks predict gut microbiome composition in wild baboons. *Elife* **4**, e05224. (doi:10.7554/eLife.05224)
- Lee W *et al.* 2021 Stomach and colonic microbiome of wild Japanese macaques. *Am. J. Primatol.* **83**, e23242. (doi:10.1002/ajp.23242)
- Liu R *et al.* 2022 Specialised digestive adaptations within the hindgut of a colobine monkey. *Innov.* **3**, 100207. (doi:10.1016/j.xinn.2022.100207)
- Clayton JB, Shields-Cutler RR, Hoops SL, Al-Ghalith GA, Sha JCM, Johnson TJ, Knights D. 2019 Bacterial community structure and function distinguish gut sites in captive red-shanked doucs (*Pygathrix nemaeus*). *Am. J. Primatol.* **81**, e22977. (doi:10.1002/ajp.22977)
- Hanya G *et al.* 2020 Fermentation ability of gut microbiota of wild Japanese macaques in the highland and lowland Yakushima: *in vitro* fermentation assay and genetic analyses. *Microb. Ecol.* **80**, 459–474. (doi:10.1007/s00248-020-01515-8)

26. Matsuda I, Stark DJ, Saldivar DAR, Tuuga A, Nathan S, Goossens B, van Schaik CP, Koda H. 2020 Large male proboscis monkeys have larger noses but smaller canines. *Commun. Biol.* **3**, 522. (doi:10.1038/s42003-020-01245-0)
27. Hale VL, Tan CL, Niu K, Yang Y, Knight R, Zhang Q, Cui D, Amato KR. 2018 Diet versus phylogeny: a comparison of gut microbiota in captive colobine monkey species. *Microb. Ecol.* **75**, 515–527. (doi:10.1007/s00248-017-1041-8)
28. Meijaard E, Nijman V. 2000 Distribution and conservation of the proboscis monkey (*Nasalis larvatus*) in Kalimantan, Indonesia. *Biol. Conserv.* **92**, 15–24. (doi:10.1016/S0006-3207(99)00066-X)
29. Matsuda I, Nakabayashi M, Otani Y, Yap SW, Tuuga A, Wong A, Bernard H, Wich SA, Kubo T. 2019 Comparison of plant diversity and phenology of riverine and mangrove forests with those of the dryland forest in Sabah, Borneo, Malaysia. In *Primates in flooded habitats: ecology and conservation* (eds K Nowak, AA Barnett, I Matsuda), pp. 15–28. Cambridge, UK: Cambridge University Press. (doi:10.1017/9781316466780)
30. Bernard H, Matsuda I, Hanya G, Phua MH, Oram F, Ahmad AH. 2019 Feeding ecology of the proboscis monkey in Sabah, Malaysia, with special reference to plant species-poor forests. In *Primates in flooded habitats: ecology and conservation* (eds K Nowak, AA Barnett, I Matsuda), pp. 89–98. Cambridge, UK: Cambridge University Press. (doi:10.1017/9781316466780)
31. Boonratana R. 2003 Feeding ecology of proboscis monkeys (*Nasalis larvatus*) in the lower Kinabatangan, Sabah, Malaysia. *Sabah Parks Nature J.* **6**, 1–6.
32. Bennett EL, Sebastian AC. 1988 Social organization and ecology of proboscis monkeys (*Nasalis larvatus*) in mixed coastal forest in Sarawak. *Int. J. Primatol.* **9**, 233–255. (doi:10.1007/BF02737402)
33. Boonratana R. 2002 Social organisation of proboscis monkeys (*Nasalis larvatus*) in the lower Kinabatangan, Sabah. *Mal. Mal. Nat. J.* **56**, 57–75.
34. Yeager CP. 1991 Proboscis monkey (*Nasalis larvatus*) social organization: intergroup patterns of association. *Am. J. Primatol.* **23**, 73–86. (doi:10.1002/ajp.1350230202)
35. Matsuda I, Murai T, Grueter CC, Tuuga A, Goossens B, Bernard H, Yahya NK, Orozco-terWengel P, Salgado-Lynn M. 2024 The multilevel society of proboscis monkeys with a possible patrilineal basis. *Behav. Ecol. Sociobiol.* **78**. (doi:10.1007/s00265-023-03419-2)
36. Matsuda I, Abram NK, Stark DJ, Sha JCM, Ancrenaz M, Goossens B, Lackman I, Tuuga A, Kubo T. 2020 Population dynamics of the proboscis monkey *Nasalis larvatus* in the lower Kinabatangan. *Oryx* **54**, 583–590. (doi:10.1017/S0030605318000467)
37. Amato KR *et al.* 2013 Habitat degradation impacts black howler monkey (*Alouatta pigra*) gastrointestinal microbiomes. *ISME J.* **7**, 1344–1353. (doi:10.1038/ismej.2013.16)
38. Clayton JB *et al.* 2018 Associations between nutrition, gut microbiome, and health in a novel nonhuman primate model. *Sci. Rep.* **8**, 11159. (doi:10.1038/s41598-018-29277-x)
39. Bernard H, Bili R, Matsuda I, Hanya G, Wearn OR, Wong A, Ahmad AH. 2016 Species richness and distribution of primates in disturbed and converted forest landscapes in Northern Borneo. *Trop. Conserv. Sci.* **9**, 194008291668010. (doi:10.1177/1940082916680104)
40. Zhu L, Clayton JB, Suhr Van Haute MJ, Yang Q, Hassenstab HR, Mustoe AC, Knights D, Benson AK, French JA. 2020 Sex bias in gut microbiome transmission in newly paired marmosets (*Callithrix jacchus*). *mSystems* **5**, e00910-19. (doi:10.1128/mSystems.00910-19)
41. Xia W, Zhao M, Wang D, Wang F, Chen H, Liu G, Zhu L, Li D. 2022 Invasion and defense of the basic social unit in a nonhuman primate society leads to sexual differences in the gut microbiome. *Integr. Zool.* **17**, 168–180. (doi:10.1111/1749-4877.12608)
42. Matsuda I, Tuuga A, Bernard H, Furuichi T. 2012 Inter-individual relationships in proboscis monkeys: a preliminary comparison with other non-human primates. *Primates*. **53**, 13–23. (doi:10.1007/s10329-011-0259-1)
43. Yeager CP. 1990 Proboscis monkey (*Nasalis larvatus*) social organization: group structure. *Am. J. Primatol.* **20**, 95–106. (doi:10.1002/ajp.1350200204)
44. Matsuda I. 2008 Feeding and ranging behaviors of proboscis monkey *Nasalis larvatus* in Sabah, Malaysia. Dissertation, [Sapporo, Japan]: Graduate School of Environmental Earth Science, Hokkaido University.
45. Murai, T. 2004 Social structure and mating behavior of proboscis monkey *Nasalis larvatus* (Primates; Colobinae). Sapporo, Japan: Graduate School of Environmental Earth Science, Hokkaido University.
46. Matsuda I, Tuuga A, Higashi S. 2010 Effects of water level on sleeping-site selection and inter-group association in proboscis monkeys: why do they sleep alone inland on flooded days? *Ecol. Res.* **25**, 475–482. (doi:10.1007/s11284-009-0677-3)
47. Matsuda I, Otani Y, Bernard H, Wong A, Tuuga A. 2016 Primate survey in a Bornean flooded forest: evaluation of best approach and best timing. *Mamm. Stud.* **41**, 101–106. (doi:10.3106/041.041.0201)
48. Matsuda I *et al.* 2014 Faecal particle size in free-ranging primates supports a 'rumination' strategy in the proboscis monkey (*Nasalis larvatus*). *Oecologia* **174**, 1127–1137. (doi:10.1007/s00442-013-2863-9)
49. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO. 2013 Evaluation of general 16S Ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* **41**, e1. (doi:10.1093/nar/gks808)
50. Caporaso JG *et al.* 2010 QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* **7**, 335–336. (doi:10.1038/nmeth.f.303)
51. R-Core-Development-Team. 2023 R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. See <https://www.R-project.org/>.
52. McMurdie PJ, Holmes S. 2013 Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* **8**, e61217. (doi:10.1371/journal.pone.0061217)
53. Quinn GP, Keough MJ. 2002 *Experimental design and data analysis for biologists*. Cambridge, UK: Cambridge University Press. (doi:10.1017/CB09780511806384)
54. Barton K. 2012 *MuMIn: multi-model inference*. See <https://cran.r-project.org/web/packages/MuMIn/MuMIn.pdf>.

55. Burnham KP, Anderson DR. 2002 *Model selection and multimodel inference: a practical information-theoretic approach*, 2nd ed. New York, NY: Springer New York.
56. Burnham KP, Anderson DR, Huyvaert KP. 2011 AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behav. Ecol. Sociobiol.* **65**, 23–35. (doi:10.1007/s00265-010-1029-6)
57. Gogarten JF *et al.* 2018 Factors influencing bacterial microbiome composition in a wild non-human primate community in Tai National Park, Cote D'Ivoire. *ISME J.* **12**, 2559–2574. (doi:10.1038/s41396-018-0166-1)
58. Wang X, Wang Z, Pan H, Qi J, Li D, Zhang L, Shen Y, Xiang Z, Li M. 2021 Captivity influences the gut microbiome of *Rhinopithecus roxellana*. *Front. Microbiol.* **12**, 763022. (doi:10.3389/fmicb.2021.763022)
59. Hale VL, Tan CL, Niu K, Yang Y, Zhang Q, Knight R, Amato KR. 2019 Gut microbiota in wild and captive Guizhou snub-nosed monkeys, *Rhinopithecus brelichi*. *Am. J. Primatol.* **81**, e22989. (doi:10.1002/ajp.22989)
60. Sawada A, Hayakawa T, Kurihara Y, Lee W, Hanya G. 2022 Seasonal responses and host uniqueness of gut microbiome of Japanese macaques in lowland Yakushima. *Anim. Microbiome.* **4**, 54. (doi:10.1186/s42523-022-00205-9)
61. Wu Y *et al.* 2020 Characterisation of the gut microbial community of *Rhesus macaques* in high-altitude environments. *BMC Microbiol.* **20**, 68. (doi:10.1186/s12866-020-01747-1)
62. Xia T *et al.* 2021 Seasonal dynamics of gut microbiota in a cohort of wild Tibetan macaques (*Macaca thibetana*) in Western China. *Glob. Ecol. Conserv.* **25**, e01409. (doi:10.1016/j.gecco.2020.e01409)
63. Springer A, Fichtel C, Al-Ghalith GA, Koch F, Amato KR, Clayton JB, Knights D, Kappeler PM. 2017 Patterns of seasonality and group membership characterize the gut microbiota in a longitudinal study of wild Verreaux's sifakas (*Propithecus verreauxi*). *Ecol. Evol.* **7**, 5732–5745. (doi:10.1002/ece3.3148)
64. Trosvik P *et al.* 2018 Multilevel social structure and diet shape the gut microbiota of the gelada monkey, the only grazing primate. *Microbiome* **6**, 84. (doi:10.1186/s40168-018-0468-6)
65. Suzuki-Hashido N, Tsuchida S, Azumano A, Goossens B, Saldivar DAR, Stark DJ, Tuuga A, Ushida K, Matsuda I. 2023 Isolation of bacteria from freeze-dried samples and the functional characterization of species-specific lactic acid bacteria with a comparison of wild and captive proboscis monkeys. *Microorganisms* **11**, 1458. (doi:10.3390/microorganisms11061458)
66. Suzuki-Hashido N, Tsuchida S, Hayakawa T, Sakamoto M, Azumano A, Seino S, Matsuda I, Ohkuma M, Ushida K. 2021 *Lactobacillus nasalis* sp. nov., isolated from the forestomach of a captive proboscis monkey (*Nasalis larvatus*). *Int. J. Syst. Evol. Microbiol.* **71**. (doi:10.1099/ijsem.0.004787)
67. Murai T, Mohamed M, Bernard H, Mahedi PA, Saburi R, Higashi S. 2007 Female transfer between one-male groups of proboscis monkey (*Nasalis larvatus*). *Primates.* **48**, 117–121. (doi:10.1007/s10329-006-0005-2)
68. Murai T. 2004 Social behaviors of all-male proboscis monkeys when joined by females. *Ecol. Res.* **19**, 451–454. (doi:10.1111/j.1440-1703.2004.00656.x)
69. Matsuda I, Zhang P, Swedell L, Mori U, Tuuga A, Bernard H, Sueur C. 2012 Comparisons of intraunit relationships in nonhuman primates living in multilevel social systems. *Int. J. Primatol.* **33**, 1038–1053. (doi:10.1007/s10764-012-9616-1)
70. Bornbusch SL, Greene LK, Rahobilalaina S, Calkins S, Rothman RS, Clarke TA, LaFleur M, Drea CM. 2022 Gut microbiota of ring-tailed lemurs (*Lemur catta*) vary across natural and captive populations and correlate with environmental microbiota. *Anim. Microbiome.* **4**, 29. (doi:10.1186/s42523-022-00176-x)
71. Moeller AH, Peeters M, Ndjango JB, Li Y, Hahn BH, Ochman H. 2013 Sympatric chimpanzees and gorillas harbor convergent gut microbial communities. *Genome Res.* **23**, 1715–1720. (doi:10.1101/gr.154773.113)
72. Matsuda I, Tuuga A, Higashi S. 2009 The feeding ecology and activity budget of proboscis monkeys. *Am. J. Primatol.* **71**, 478–492. (doi:10.1002/ajp.20677)
73. Chen Y, Lai Y, Zheng J, Liu Z, Nong D, Liang J, Li Y, Huang Z. 2023 Seasonal variations in the gut microbiota of white-headed black langur (*Trachypithecus leucocephalus*) in a limestone forest in Southwest Guangxi, China. *Front. Ecol. Evol.* **11**. (doi:10.3389/fevo.2023.1126243)
74. Orkin JD, Campos FA, Myers MS, Cheves Hernandez SE, Guadamuz A, Melin AD. 2019 Seasonality of the gut microbiota of free-ranging white-faced capuchins in a tropical dry forest. *ISME J.* **13**, 183–196. (doi:10.1038/s41396-018-0256-0)
75. Srivathsan A, Ang A, Vogler AP, Meier R. 2016 Fecal metagenomics for the simultaneous assessment of diet, parasites, and population genetics of an understudied primate. *Front. Zool.* **13**, 17. (doi:10.1186/s12983-016-0150-4)
76. Lyke MM, Di Fiore A, Fierer N, Madden AA, Lambert JE. 2019 Metagenomic analyses reveal previously unrecognized variation in the diets of sympatric old world monkey species. *PLoS One* **14**, e0218245. (doi:10.1371/journal.pone.0218245)
77. Osman NA, Abdul-Latiff MAB, Mohd-Ridwan AR, Yaakop S, Nor SM, Md-Zain BM. 2020 Diet composition of the wild stump-tailed macaque (*Macaca arctoides*) in Perlis state park, Peninsular Malaysia, using a chloroplast tRNA DNA metabarcoding approach: a preliminary study. *Animals* **10**, 2215. (doi:10.3390/ani10122215)
78. Matsuda I, Tuuga A, Bernard H. 2011 Riverine refuging by proboscis monkeys (*Nasalis larvatus*) and sympatric primates: implications for adaptive benefits of the riverine habitat. *Mamm. Biol.* **76**, 165–171. (doi:10.1016/j.mambio.2010.03.005)
79. Ruslin F, Matsuda I, Md-Zain BM. 2019 The feeding ecology and dietary overlap in two sympatric primate species, the long-tailed macaque (*Macaca fascicularis*) and dusky langur (*Trachypithecus obscurus obscurus*). *Primates.* **60**, 41–50. (doi:10.1007/s10329-018-00705-w)
80. Sha JCM, Hanya G. 2013 Temporal food resource correlates to the behavior and ecology of food-enhanced long-tailed macaques (*Macaca fascicularis*). *Mamm. Stud.* **38**, 163–175. (doi:10.3106/041.038.0305)

81. Seaman DJI, Bernard H, Ancrenaz M, Coomes D, Swinfield T, Milodowski DT, Humle T, Struebig MJ. 2019 Densities of Bornean orang-utans (*Pongo pygmaeus morio*) in heavily degraded forest and oil palm plantations in Sabah, Borneo. *Am. J. Primatol.* **81**, e23030. (doi:10.1002/ajp.23030)
82. Amato KR, Kuthyar S, Ekanayake-Weber M, Salmi R, Snyder-Mackler N, Wijayathunga L, Vandercone R, Lu A. 2020 Gut microbiome, diet, and conservation of endangered langurs in Sri Lanka. *Biotropica* **52**, 981–990. (doi:10.1111/btp.12805)
83. Barelli C, Albanese D, Stumpf RM, Asangba A, Donati C, Rovero F, Hauffe HC. 2020 The gut microbiota communities of wild arboreal and ground-feeding tropical primates are affected differently by habitat disturbance. *mSystems* **5**, e00061-20. (doi:10.1128/mSystems.00061-20)
84. Jose L, Lee W, Hanya G, Tuuga A, Goossens B, Tangah J. 2024 Data from: Gut microbial community in proboscis monkeys (*nasalis larvatus*): implications for effects of geographical and social factors. Figshare. (doi:10.6084/m9.figshare.c.7349579)