

EurJOC

European Journal of Organic Chemistry

 **Chemistry
Europe**

European Chemical
Societies Publishing



Accepted Article

Title: Adaptable Synthesis of Chondroitin Sulfate Disaccharide Subtypes Preprogrammed for Regiospecific O-Sulfation

Authors: Hannah S. Wootton, Sian S. Berry, Elaine L. Ferguson, Clare S. Mahon, and Gavin J Miller

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* **2024**, e202400587

Link to VoR: <https://doi.org/10.1002/ejoc.202400587>

WILEY-VCH

Adaptable Synthesis of Chondroitin Sulfate Disaccharide Subtypes Preprogrammed for Regiospecific *O*-Sulfation

Hannah S. Wootton,^a Sian S. Berry,^a Elaine L. Ferguson,^b Clare S. Mahon^c and Gavin J. Miller^{a*}

^aCentre for Glycoscience and School of Chemical and Physical Sciences, Keele University, Keele, Staffordshire, ST5 5BG, United Kingdom.

^bAdvanced Therapies Group, School of Dentistry, College of Biomedical and Life Sciences, Cardiff University, Heath Park, Cardiff, UK

^cDepartment of Chemistry, Durham University, South Road, Durham, United Kingdom.

*g.j.miller@keele.ac.uk

Abstract

A divergent synthetic route to chondroitin sulfate (CS) disaccharide precursors, including rarer subtypes such as CS-D, has been developed. From common intermediates, a series of thioglycoside D-glucosyl donors and 4,6-*O*-benzylidene protected D-galactosamine acceptors are utilised in a robust glycosylation reaction, achieving β -selectivity and consistent yields (60-75%) on scales >2.0 g. A post-glycosylation oxidation to D-glucuronic acid and orthogonal protecting groups delivers access to CS-A, CS-C, CS-D, CS-E and CS-O precursor subtypes. Of further note is a 4-*O*-benzyl regioselective reductive ring opening of a 4,6-*O*-benzylidene protected disaccharide using dichlorophenylborane (PhBCl₂) and triethylsilane (Et₃SiH) to access a CS-D precursor, in 73% yield over two steps. Finally, synthesis of a 6-*O*-sulfated CS-C disaccharide containing a conjugable anomeric allyl tether is completed. These materials will provide a benchmark to further synthesise and study chondroitin sulfates.

Introduction

Chondroitin sulfate (CS) is a glycosaminoglycan composed of D-glucuronic acid (D-GlcA) and *N*-Acetyl-D-galactosamine (D-GalNAc), with repeating sequences of -4)- β -D-GlcA-(1,3)- β -D-GalNAc-(1- disaccharides, typically in excess of 100 units.^[1] During biosynthetic assembly, enzymatic modification of the repeating nascent disaccharide backbone gives rise to a number of different sulfation patterns, creating a multitude of CS subtypes and heterogenous structures.^[2] Naturally occurring CS therefore contains an assortment of subtypes, and proportions that also vary depending on the polysaccharide source. CS-A (D-GalNAc 4-*O*-sulfation) and CS-C (D-GalNAc 6-*O*-sulfation) are abundant compared to other subtypes such as CS-D (2'-*O*-D-GlcA-6-*O*-D-GalNAc sulfation) and CS-E (4,6-*O*-D-GalNAc sulfation).^[3-5] CS interacts with growth factors, cytokines, chemokines and adhesion molecules,^[6] regulating physiological processes including embryonic and brain development,^[7,8] anti-inflammatory effects,^[9] wound healing^[10] and signalling.^[11] Whilst it is established that CS sulfation patterns correlate to related biological function, the specificities of such molecular interactions is an underdeveloped field.^[12] Access to homogenous, structurally defined CS oligosaccharides is therefore of utmost importance for the study of related CS-protein interactions, but is hampered by difficulties in obtaining significant amounts of structurally defined sequences from natural

sources, particularly for rarer subtypes such as CS-D. Relatedly, synthetic approaches to heparan sulfate (HS) fragments have advanced more rapidly and proven extremely successful.^[13–17]

Despite recent advances in the synthesis of CS oligosaccharides and related building blocks,^[1,18–23] approaches are required that allow diversification to CS subtypes with varying sulfation patterns, chain lengths and conjugation capabilities. Herein, and as part of a wider program targeting approaches to other glycosaminoglycans,^[24,25] we develop a reliable and versatile synthesis of CS -4)- β -D-GlcA-(1,3)- β -D-GalNAc-(1- building block disaccharides from a small library of D-Glc and D-GalN monosaccharides (*Figure 1*). Careful consideration of protecting groups was made, to enable: β -stereoselectivity using a participating 2-*O*-benzoate (OBz) protecting group, variation of sulfation site programming using regiospecific ring opening of D-GalN 4,6-*O*-benzylidene acetals, orthogonal D-GlcA-C4'-*O*-substituents for elongation potential and a reducing end anomeric allyl group as a conjugable handle or orthogonal group for alternative donor formation.




Figure 1: Top: Various subtypes of the glycosaminoglycan chondroitin sulfate (CS). The standard disaccharide repeat is shown inside the purple dotted box. **Bottom:** A retrosynthetic approach to install capability for regiodefined CS sulfation, conjugation and non-reducing end iterative extension into appropriate disaccharides from monosaccharide precursors.

Results and Discussion


Previous approaches to CS oligosaccharides and associated building blocks have utilised trichloroacetimidate (TCAI) D-GlcA donors, namely a pre-glycosylation oxidation strategy.^[26–28] Whilst proven and undoubtedly useful, issues due to low reactivity of uronate donors and the possibility of donor-derived side product formation (e.g., orthoesters or *N*-trichloroacetamides) are notable considerations.^[29] Here we instead opted to explore a post-glycosylation oxidation approach (D-Glc to D-GlcA) using thioglycoside glucosyl donors

(shelf-stable and generally synthesised in fewer steps than TCAI donors). Previously, the approach has been utilised in oligosaccharide assembly^[30–32] but more relatedly, Lei and co-workers established a post-glycosylation oxidation strategy to synthesise CS-E oligosaccharides using TCAI donors to access a key disaccharide for iterative synthesis.^[33]

Glycosyl Donor and Acceptor Building Block Synthesis

Synthesis towards thioglycoside glucosyl donors commenced from commercial 1,2,3,4,6-penta-*O*-acetyl- β -D-glucose **1**, undergoing thioglycosidation with ethanethiol (EtSH) using boron trifluoro etherate (BF₃·Et₂O) as a Lewis acid and affording the β -thiol in 87% yield (*Scheme 1*). Deacetylation using Na₂CO₃ in MeOH afforded tetrol **2** in 98% yield and an appropriate material to diversify the route to the required building blocks. Benzylidene protection using benzaldehyde dimethyl acetal with 10-camphorsulfonic acid (CSA) in acetonitrile (MeCN) was followed by regioselective C3-*O*-benzylation using dibutyltin oxide (ⁿBu₂SnO) and cesium fluoride (CsF) with benzyl bromide.^[34] Subsequent C2-*O*-benzoylation using benzoyl chloride and pyridine afforded thioglycoside **3** in 49% yield over three steps. From this material two divergent routes were progressed. The first saw regioselective reductive ring opening using borane tetrahydrofuran complex (BH₃·THF) and a catalytic amount of trimethylsilyl trifluoromethanesulfonate (TMSOTf) as Lewis acid to give a 6-OH, 4-OBn protected material in 91% yield. COSY NMR correlation between a broad singlet at $\delta = 1.97$ ppm and H6 environments at $\delta = 3.93$ and 3.72 ppm indicated a C6-OH was present and the desired regiochemistry achieved. Additionally, the ¹³C NMR spectrum displayed a chemical shift at δ 62.1 ppm, which further signified the C6-OH. Reaction of this material with chloroacetyl chloride and pyridine afforded the desired donor **4** in 86% yield. Secondly, thioglycoside **3** was subjected to acidic benzylidene cleavage using CSA in MeOH at 40 °C, followed by regioselective C6-*O*-chloroacetylation at low temperature to afford alcohol **5** in 80% yield. Finally, C4-*O*-levulinoyl (Lev) ester protection proceeded smoothly in 85% yield to generate glycosyl donor **6**, alternatively protected at C4, compared to **4**; the C4-*O*-Lev group enabling access to regioselective deprotection (versus C4 OBn in **4**) and access to a glycosyl acceptor form.

Thioglycoside donor synthesis




Scheme 1: Synthesis of thioglycoside glucosyl donor panel starting from commercial material 1 and delivering orthogonally protected donors 4, 6, 9 and 10.

Furthermore, from tetrol 2 *para*-methoxybenzylidene protection was completed, followed by C2,C3-*O*-benzylation to afford thioglycoside 7 in 88% yield over two steps. Reductive ring opening of *para*-methoxybenzylidene 7 was explored, for the first time, employing BH₃·NMe₃ complex and AlCl₃. Initial yields in forming the desired C6-*O*-*p*-methoxybenzyl (PMB) regioisomer 8 were low (35-44%), noting amounts (22-30%) of returned 7 alongside an unwanted anomeric hydrolysis product formed during the 3 h reaction. By reducing the reaction time to 1 h (expedited by the addition of two equivalents of H₂O),^[35] the isolated yield of 8 increased to 81% and was accessible on 10 g scale. Finally, C4-*O*-protection of 8 was completed using either *tert*-butyldimethylsilyl trifluoromethanesulfonate (TBDMSOTf) or 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC)/levulinic acid to afford orthogonally C4-protected donors 9 (91% yield) and 10 (77% yield), completing the panel.

Relatedly, a scalable route towards *N*-Trichloroethoxycarbonyl (Troc) and *N*-trichloroacetyl (TCA) protected D-GalN acceptors 17 and 18 was developed starting from commercial D-GalN 11. The free amine was temporarily masked using *p*-anisaldehyde under basic conditions (**Scheme 2**). This was followed by global acetylation and subsequent acidic hydrolysis of the aldimine protecting group using HCl in refluxing acetone to afford amine 12 in 78% yield over three steps. This method of temporary amino protection was high yielding, β-selective and chromatography free. It should though be noted that initial large-scale attempts (> 5.0 g) resulted in imine hydrolysis during acetylation, reducing the yields markedly and affording the undesired *N*-acyl by-product. To overcome this, rigorous drying under high

vacuum for 8 h was necessary, but enabled synthesis of **12** to be reproducibly scaled up to >40.0 g.



Scheme 2: Synthesis of D-galactosamine C3-OH acceptors with variant *N*-protecting groups, *N*-Troc **17** and *N*-TCA **18**.


From amine **12** both *N*-Troc and *N*-TCA protected acceptors were synthesised. *N*-Troc protection of **12** was completed on scales up to 15.0 g in 84% yield. Subsequent protection of the anomeric position with a conjugable aglycon (OAllyl) was completed using allyl alcohol and TMSOTf to afford β-**15** in 89% yield and a 19:1 β:α ratio, with the α-anomer separable *via* column chromatography. Deacetylation of **15** using Na₂CO₃ in MeOH was followed by 4,6-*O*-benzylidene acetal protection to deliver acceptor **17** on multigram scale and in 76% yield over two steps. The desired β-anomeric stereochemistry within **17** was confirmed using ¹H NMR and an observed *trans*-diaxial coupling constant of 8.2 Hz for H1 (³*J*_{H1-H2}). For the synthesis of *N*-TCA acceptor **18**, a similar series of reactions were completed (**Scheme 2**), noting however that anomeric allylation of *N*-TCA derivative **14** was achieved *via* the glycosyl bromide, using allyl alcohol and indium chloride to selectively synthesize β-**16** in 57% yield. The required deacetylation and benzylidene protection steps afforded acceptor **18** in 88% yield over two steps.

Glycosylation Methodology Development for CS Disaccharide Synthesis

We next explored glycosylation capability of our donor panel (compounds **4**, **6**, **9** and **10**) with D-GalN acceptors **17** and **18**. Glycosylation of thioglycoside donor **4** using *N*-Troc acceptor **17** and an *N*-iodosuccinimide (NIS)/triflic acid (TfOH) promotor system afforded disaccharide **19** in 75% yield (**Scheme 3**). The reaction proceeded smoothly, and isolated yields were consistent on >2.0 g scale. The desired β-glycosidic linkage was confirmed through a large *J*_{H1'-H2'} coupling constant of 8.0 Hz (¹H δ = 4.83 ppm). A H1'-C3 HMBC correlation confirmed the desired (1,3) linkage had formed. The glycosylation conditions were successfully replicated with C4-*O*-Lev donor **6**, generating disaccharide **20** in 60% yield. Switching to *N*-TCA acceptor **18** and donor **4** furnished disaccharide **21** in 68% yield and

glycosylation of **18** with donor **9** delivered disaccharide **22** in 74% yield. Finally, use of C4-*O*-Lev protected donor **10** afforded disaccharide **23** in a slightly lower 63% yield. From these results the reactivity of an *N*-Troc D-GalN acceptor was established as comparable to an *N*-TCA derivative.

CS Precursor Disaccharide Synthesis




Scheme 3: CS precursor disaccharide synthesis using variously protected donors and acceptors and NIS/TfOH glycosylation conditions. Following column chromatography of the crude reactions only β-anomer was isolated for compounds **19-22**.

Post Glycosylation Oxidation to Access CS Precursor Disaccharides

With access to a robust glycosylation procedure in place, we sought to complete post-glycosylation oxidation of the disaccharide D-Glc component (to D-GlcA) and from there access the required protecting group patterns for CS-A/C/D/E/O. Accordingly, we removed the 6'-*O*-protecting groups from within disaccharides **19** and **23** [thiourea for AcCl cleavage in **19** and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) for oxidative PMB removal in **23**]. From here oxidative conditions using a biphasic 2,2,6,6-tetramethyl-1-piperidinyloxy/(diacetoxyiodo)benzene (TEMPO/BAIB) system were applied to deliver common disaccharides **24** and **25** in 85% and 60% yields respectively (**Scheme 4**). The lower yield for disaccharide **25** was attributed to a sensitivity of the *N*-TCA protecting group to excess K_2CO_3 in the carboxylate methylation step.

Access to CS precursor disaccharide library



Scheme 4: Towards a variably protected series of CS-disaccharide precursors and exemplar synthesis of a CS-C disaccharide.

From disaccharides **24** and **25** we completed access to five CS precursors. Firstly, to access a protecting group pattern towards CS-A (D-GalNAc-4-*O*-sulfation pattern), reductive ring opening of **24** was completed to provide a C6-*O*-benzyl moiety in product **26**. Disaccharide **24** was subjected to triethylsilane (Et₃SiH) and trifluoroacetic acid (TFA) in dichloromethane (DCM) at 0 °C and the crude product from this step was then subject to 6-*O*-chloroacetylation, which proceeded smoothly to afford CS-A precursor **26** in 60% yield over two steps and programmed with orthogonal 4-*O*-protecting group for sulfation. Towards a CS-D disaccharide (D-GalNAc-6-*O*- and D-GlcA-2-*O*-sulfation) we required alternate reductive ring opening conditions. Sakagami and Hamana previously identified a combination of PhBCl₂ and Et₃SiH for regioselective ring opening of benzylidene acetals to 4-OBn/6-OH products.^[36] Reactions were generally performed in DCM at -78 °C with an excess of Et₃SiH and PhBCl₂. However, in DCM, when an excess of Et₃SiH is used, the silane reduces PhBCl₂ to PhBHCl, which can hydroborate alkenes, presenting a potential problem for our anomeric *O*-allyl moiety. To avoid this, use of only 1.1 equivalents of Et₃SiH in DCM at low temperatures has been reported.^[37] Upon adopting these conditions for disaccharide **24** for 15 minutes all starting material was consumed and there was no notable side product formation by TLC analysis. Subsequent C6-*O*-benzoylation of the crude material delivered disaccharide **27** in 84% over two steps. NMR analysis confirmed orthogonal 4-*O*-/6-*O*-protecting group patterns in both **26** and **27** (illustrated for **27** in **Figure 2**).




Figure 2: HMBC NMR (400 MHz) for disaccharide **27**, highlighting the result of regioselective ring opening using PhBCl_2 and Et_3SiH .

Lastly, acidic cleavage of the benzylidene acetal within disaccharide **25** afforded **28** as a precursor to CS-C, CS-E and CS-O. To exemplify the utility of our approach we completed a formal synthesis of a CS-C disaccharide (*Scheme 4*, green box). 6-*O*-Sulfation of disaccharide **28** using 2.5 equivalents of SO_3NMe_3 was achieved in 94% yield. Initially, saponification of all esters and the TCA moiety resulted in disappearance of the distinctive allyl peaks within the ^1H NMR spectrum between δ 5.0 and 6.0 ppm, suggesting an unwanted side-reaction had occurred. Reducing the reaction time and the amount of 30% H_2O_2 used (to 2.0 equivalents) enabled us to isolate, following a final *N*-acetylation, C6-sulfated disaccharide **29** in 19% yield over three steps.

Conclusion

An efficient approach to the synthesis of CS disaccharide precursors programmed towards different natural sulfation patterns has been established. This divergent strategy enables access to protected CS-A, CS-C, CS-D, CS-E and CS-O derivatives from common disaccharide precursors. Optimisation of a central glycosylation reaction using shelf-stable thioglycoside glucosyl donors enables reproducible multigram scale synthesis with β -stereoselectivity and ultimate access to D-GlcA(1,3- β)-D-GalN systems incorporating multiple orthogonal protecting groups. An exemplar synthesis of a free CS-C disaccharide is completed, containing an anomeric *O*-allyl handle, demonstrating capability for conjugation, for example using alkene click chemistry. Overall, this methodology offers an important contribution to

provide the building blocks required for iterative CS oligosaccharide synthesis, for example using a [2+2] glycosylation approach and pre-programmed for regiospecific *O*-sulfation. The synthesis of longer, bespoke CS sequences for biological application is currently underway and will be reported in due course.

Experimental

For full experimental details and characterisation of compounds please refer to the supporting information.

General Experimental

Unless otherwise stated, all reagents used in the following experiments were bought commercially from Acros Organics, Alfa Aesar, Apollo Scientific, Biosynth, Fisher Scientific, Fluorochem, Sigma Aldrich or TCI chemicals and were used without further purification. Dry solvents were dried and stored under N₂ in Young's flasks over 4 Å molecular sieves. Anhydrous DMF, pyridine and THF were purchased from Acros Organics, fitted with AcroSeal™ packaging. For reactions that required heating, DrySyn heating blocks were used as the heat source. Reactions were monitored by thin layer chromatography (TLC) using pre-coated 0.25 mm 60 F₂₅₄ silica gel plates (Merck) and eluent systems outlined in the respective experiments. Visualisation was achieved using UV light ($\lambda = 254$ nm), and 10% H₂SO₄:EtOH or ninhydrin staining followed by heating. Flash column chromatography was performed using silica gel [high purity grade, 60 Å pore size, 40-63 μ m particle size]. NMR spectra were recorded at 400 MHz on a Bruker AVIII400 spectrometer using deuterated solvent. Chemical shifts are reported in parts per million (ppm), coupling constants (*J*) are reported in Hertz (Hz) and multiplicities are abbreviated as; s (singlet), d (doublet), t (triplet) or m (multiplet) or combinations thereof. Chemical shifts were referenced to tetramethylsilane (TMS, where $\delta = 0.00$ ppm). HRMS were recorded on a ThermoScientific LTQ Orbitrap XL at the ESPRC National Mass Spectrometry Facility at Swansea University. Optical rotations were recorded on a Bellingham + Stanley ADP430 (specific rotation, tube length: 50 mm, concentrations in g per 100 mL).

General procedure A – Glycosylation

In a multi-neck flask, glycosyl donor (1.2 equiv.), glycosyl acceptor (1.0 equiv.) and 4 Å M.S. were placed under three cycles of vacuum and N₂. Anhydrous DCM (0.05 M) was added and the solution was pre-dried for 1 h. The solution was cooled to –35 °C (using IMS and dry ice) and stirred for a further 10 min at this temperature. *N*-iodosuccinimide (1.3 equiv.) and TfOH (0.15 equiv.) were added and the reaction was gradually warmed to 0 °C over approximately 2.5 h. TLC analysis (1:1 Hexane:EtOAc) showed full consumption of the glycosyl acceptor. The reaction was quenched with Et₃N (1 mL) and extracted with sat. aq. Na₂S₂O₃ (1 × 50 mL), sat. aq. NaHCO₃ (1 × 100 mL) and brine (1 × 100 mL). The organic phase was dried over MgSO₄ and filtered. Solvents were removed *in vacuo* and the crude foam was purified *via* manual flash column chromatography (10:1 → 3:1 → 1.5:1, hexane:EtOAc) which afforded the target disaccharide.

General Procedure B – Tempo Oxidation

To a vigorously stirred solution of starting material (1.0 equiv.) in DCM:H₂O (2:1 v/v, 0.2 M), 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) (0.2 equiv.) and (diacetoxyiodo)benzene (BAIB) (2.5 equiv.) were added at 0 °C. After 10 min., the solution was warmed to RT and stirred for a further 2-3 h. TLC analysis showed formation of the desired product (7:1:1 EtOAc:MeOH:H₂O). The reaction was diluted with EtOAc, quenched with sat. aq. Na₂S₂O₃ and the layers were separated. The pH of the aqueous phase was tuned to pH 2 with 1 M HCl then extracted with EtOAc (3 ×). The combined organic extracts were dried over MgSO₄, filtered and concentrated *in vacuo*. Under an atmosphere of N₂, the crude material was suspended in anhydrous DMF (0.2 M) and cooled to 0 °C, to which MeI (3.0 equiv.) and K₂CO₃ (3.0 equiv.) were added. The reaction was stirred at RT in the dark for 2 h before dilution with EtOAc and washed with H₂O (2 ×) and brine (1 ×). The organic phase was dried (MgSO₄), filtered and concentrated *in vacuo*. Purification *via* flash column chromatography (10:1 → 3:1 → 1:1 hexane:EtOAc) afforded the desired product as a white solid.

Ethyl 6-*O*-chloroacetyl-3,4-*O*-benzyl-2-*O*-benzoyl-1-thio-β-D-glucopyranoside 4

At 0 °C under an atmosphere of N₂, chloroacetyl chloride (0.4 mL, 5.12 mmol, 1.3 equiv.) was added to a stirred solution of **S4** (2.0 g, 3.94 mmol, 1.0 equiv.) and pyridine (1.3 mL, 15.8 mmol, 4.0 equiv.) in anhydrous DCM (20 mL). The reaction was warmed to RT and stirred for 1.5 h. Following completion, as monitored by TLC analysis (2:1 hexane:EtOAc), the reaction was diluted in DCM (20 mL) and washed with 1 M HCl (2 × 50 mL) and brine (1 × 50 mL) before drying over MgSO₄, filtering and concentrating *in vacuo*. Purification *via* manual column chromatography (10:1 → 7:1 → 5:1, hexane:EtOAc) afforded **4** (1.9 g, 3.39 mmol, 86%) as a white solid.

Ethyl 2,3-di-*O*-benzoyl-6-*O*-*p*-methoxybenzyl-1-thio-β-D-glucopyranoside 8

Under an atmosphere of N₂, compound **7** (10.3 g, 17.7 mmol, 1.0 equiv.) was dissolved in dry THF (140 mL) and cooled to 0 °C. Borane trimethylamine complex (5.16 g, 70.8 mmol, 4.0 equiv.) was added in one portion followed by careful addition of aluminium chloride (14.2 g, 106.2 mmol, 6.0 equiv.). Once the aluminium chloride was completely dissolved, H₂O (0.64 mL, 35.4 mmol, 2.0 equiv.) was added and the reaction was stirred at RT for 1 h. The reaction was diluted using Et₂O (120 mL). After cooling to 0 °C, H₂O (7.5 mL) was added followed by 1M aq. NaOH (7.5 mL) and a further addition of H₂O (22.5 mL). The mixture was vigorously stirred for 15 min. MgSO₄ was added and the mixture was stirred for a further 15 min. After filtration, the solution was concentrated *in vacuo*. Purification *via* flash column chromatography (30:70 to 40:60 EtOAc:hexane + 1% Et₃N) afforded **8** as a colourless oil (7.9 g, 14.3 mmol, 81%).

Allyl 2-deoxy-2,2,2-trichloroethoxycarbonylamino-3,4,6-tri-*O*-acetyl-β-D-galactopyranoside 15

Under an atmosphere of N₂, compound **13** (6.60 g, 11.8 mmol, 1.0 equiv.) and allyl alcohol (2.0 mL, 29.4 mmol, 2.5 equiv.) were dissolved in DCM (80 mL) and stirred in the presence

of 4 Å molecular sieves for 1 h. TMSOTf (2.8 mL, 15.3 mmol, 1.3 equiv.) was added then the reaction was stirred at RT for an additional 1 h. When complete, the solution was washed with sat. aq. NaHCO₃ (2 × 100 mL) and brine (100 mL). The organic phase was dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification *via* manual flash column chromatography (2:1 hexane:EtOAc (α); 1:1 hexane:EtOAc (β)) afforded β-**15** as a white foam (5.4 g, 10.5 mmol, 89%) and α-**15** as a colourless oil (397 mg, 0.826 mmol, 7%).

Allyl 4,6-*O*-benzylidene-2-deoxy-2-*N*-2,2,2-trichloroethoxycarbonylamino-1-β-D-galactopyranoside 17

To a solution of compound **15** (5.40 g, 10.8 mmol, 1.0 equiv.) in MeOH (100 mL), Na₂CO₃ (1.60 g, 3.24 mmol, 0.3 equiv) was added and the reaction was stirred at RT for 2 h. The reaction mixture was neutralised with Dowex® 50W X8 H⁺ resin then filtered, washed with MeOH (3 × 40 mL) and concentrated *in vacuo* to afford a white solid. The crude product was suspended in anhydrous DMF (45 mL) and treated with CSA (400 mg, 2.39 mmol, 0.25 equiv.) and benzaldehyde dimethyl acetal (2.20 mL, 14.3 mmol, 1.5 equiv.). The reaction was stirred at 40 °C for 2 h then diluted with EtOAc (100 mL) before washing with sat. aq. NaHCO₃ (3 × 50 mL) and brine (50 mL). The organic extracts were dried over MgSO₄, filtered and concentrated *in vacuo*. The crude residue was washed with 8:1 petroleum ether:Et₂O (30 mL) to afford **17** as an off-white solid (3.60 g, 8.42 mmol, 78% over 2 steps) as a white solid.

6-*O*-Chloroacetyl-3,4-di-*O*-benzyl-2-*O*-benzoyl-D-glucopyranoside-β(1→3)-Allyl 4,6-*O*-benzylidene-2-deoxy-*N*-2,2,2-trichloroethoxycarbonylamino-β-D-galactopyranoside 19

Disaccharide **19** was prepared according to general procedure A. Acceptor **17** (1.40 g, 2.66 mmol, 1.0 equiv.) and donor **4** (2.00 g, 3.50 mmol, 1.2 equiv.) afforded **19** (2.20 g, 2.20 mmol, 75%) as a white foam.

6-*O*-Chloroacetyl-3-*O*-benzyl-4-*O*-levulinoyl-2-*O*-benzoyl-D-glucopyranoside-β(1→3)-Allyl-4,6-*O*-benzylidene-2-deoxy-*N*-2,2,2-trichloroethoxycarbonylamino-β-D-galactopyranoside 20

Disaccharide **20** was prepared according to general procedure A. Acceptor **17** (50 mg, 0.104 mmol, 1.0 equiv.) and donor **6** (75 mg, 0.125 mmol, 1.2 equiv.) afforded **20** (63 mg, 0.063 mmol, 60%) as a white foam.

6-*O*-Chloroacetyl-3,4-di-*O*-benzyl-2-*O*-benzoyl-D-glucopyranoside-β(1→3)-Allyl 4,6-*O*-benzylidene-2-deoxy-*N*-trichloroacetamido-1-β-D-galactopyranoside 21

Disaccharide **21** was prepared according to general procedure A. Acceptor **18** (600 mg, 1.33 mmol, 1.0 equiv.) and donor **4** (910 mg, 1.60 mmol, 1.2 equiv.) afforded **21** (980 mg, 1.01 mmol, 76%) as a white foam.

2,3-Di-*O*-benzoyl-4-*O*-tert-butylsilyl-6-*O*-*p*-methoxybenzyl- β -D-glucopyranoside-(1 \rightarrow 3)-allyl 4,6-*O*-benzylidene-2-deoxy-2-*N*-trichloroacetamido-1- β -D-galactopyranoside 22

Disaccharide **22** was prepared according to general procedure A. Acceptor **18** (0.469 g, 1.04 mmol) and donor **9** (0.830 g, 1.24 mmol) afforded **22** (0.81 g, 74%) as a white foam.

2,3-Di-*O*-benzoyl-4-*O*-levulinoyl-6-*O*-*p*-methoxybenzyl- β -D-glucopyranoside-(1 \rightarrow 3)-allyl 4,6-*O*-benzylidene-2-deoxy-2-*N*-trichloroacetamido-1- β -D-galactopyranoside 23

Disaccharide **23** was prepared according to general procedure A. Acceptor **18** (1.35 g, 2.07 mmol) and donor **10** (0.78 g, 1.73 mmol) afforded **23** (1.14 g, 63%) as a white foam.

Methyl(3,4-di-*O*-benzyl-2-*O*-benzoyl-D-glucopyranosyl)uronate- β (1 \rightarrow 3)-Allyl-4,6-*O*-benzylidene-2-deoxy-2-*N*,2,2-trichloroethoxycarbonylamino- β -D-galactopyranoside 24

Disaccharide **24** was prepared according to general procedure B. Compound **S7** (745 mg, 0.803 mmol, 1.0 equiv.) afforded **24** (645 mg, 0.674 mmol, 85% over two steps) as a white solid.

Methyl 2,3-Di-*O*-benzoyl-4-*O*-levulinoyl- β -D-glucopyranosyluronate-(1 \rightarrow 3)-allyl 4,6-*O*-benzylidene-2-deoxy-2-*N*-trichloroacetamido-1- β -D-galactopyranoside 25

Disaccharide **25** was prepared according to general procedure B. Compound **S8** (385 mg, 0.420 mmol, 1.0 equiv.) afforded **25** (240 mg, 0.252 mmol, 60%) as a white solid.

Methyl(3,4-di-*O*-benzyl-2-*O*-benzoyl-D-glucopyranosyl)uronate- β (1 \rightarrow 3)-Allyl-4-*O*-chloroacetyl-6-*O*-benzyl-2-deoxy-2-*N*,2,2-trichloroethoxycarbonylamino- β -D-galactopyranoside 26

Under an atmosphere of N₂, a solution of **24** (100 mg, 0.105 mmol, 1.0 equiv.) in DCM (1.1 mL) was cooled to 0 °C and Et₃SiH (84 μ L, 0.523 mmol, 5.0 equiv.) and trifluoroacetic acid (40 μ L, 0.523 mmol, 5.0 equiv.) were added. The reaction was stirred for 1 h then neutralised with Et₃N. The solution was diluted in DCM (10 mL), extracted with sat. aq. NaHCO₃ (10 mL) and brine (10 mL) before drying over MgSO₄, filtering and concentrating *in vacuo*. The crude residue was then placed under an atmosphere of N₂ and redissolved in DCM (1 mL) and pyridine (46 μ L, 79.1 mmol, 6.0 equiv.), to which chloroacetyl chloride (10 μ L, 0.124 mmol, 1.3 equiv.) was added at 0 °C. After 30 min, the reaction was diluted in DCM (10 mL) and washed successively with 1 M HCl (5 mL), sat. aq. NaHCO₃ (5 mL) and brine (5 mL). The organic phase was dried over MgSO₄, filtered and concentrated *in vacuo*. Manual flash column chromatography (100% DCM \rightarrow 5% Et₂O in DCM) afforded **26** (65 mg, 0.063 mmol, 60%) as a white foam.

Methyl(3,4-di-*O*-benzyl-2-*O*-benzoyl-D-glucopyranosyl)uronate- β (1 \rightarrow 3)-Allyl-4-*O*-benzyl-6-*O*-benzoyl-2-deoxy-2-*N*,2,2-trichloroethoxycarbonylamino- β -D-galactopyranoside 27

In a multi-neck flask, disaccharide **24** (1.00 g, 1.05 mmol, 1.0 equiv.) and 4 Å molecular sieves were placed under three cycles of vacuum and N₂. Anhydrous DCM (52.5 mL) was added and the solution was pre-dried for 1 h. The reaction was cooled to – 78 °C (using acetone and dry ice) and Et₃SiH (0.17 mL, 1.05 mmol, 1.0 equiv.) was added. The reaction was stirred for 15 minutes before PhBCl₂ (0.5 mL, 3.68 mmol, 3.5 equiv.) was added dropwise. The reaction was stirred at – 75 °C for 15 min before quenching with Et₃N (1.6 mL) and MeOH (1.6 mL). After diluting with DCM (20 mL), the organic phase was washed successively with sat. aq. NaHCO₃ (50 mL), H₂O (2 × 50 mL) and brine (50 mL). The organic phase was dried (MgSO₄), filtered and concentrated *in vacuo*. The crude residue was dissolved in anhydrous DCM (10 mL) and pyridine (0.5 mL, 6.30 mmol, 6.0 equiv.) was added. The solution was cooled to 0 °C and BzCl (0.2 mL, 1.58 mmol, 1.5 equiv.) was added. The reaction was warmed to RT and stirred for 2 h before diluting with DCM (20 mL) and washing successively with 1 M HCl (30 mL), sat. aq. NaHCO₃ (30 mL) and brine (30 mL). The organic phase was dried (MgSO₄), filtered and concentrated *in vacuo*. The crude syrup was purified *via* manual flash column chromatography (100% DCM → 5% Et₂O in DCM) then washed with 10:1 petroleum ether:Et₂O to afford **27** as a white solid (815 mg, 0.882 mmol, 84% over two steps).

Methyl 2,3-Di-*O*-benzoyl-4-*O*-levulinoyl-β-D-glucopyranosyluronate-(1→3)-allyl 2-deoxy-6-*O*-sulfonato-2-*N*-trichloroacetamido-1-β-D-galactopyranoside **S9**

Under an atmosphere of N₂, diol **28** (137 mg, 0.16 mmol, 1.0 equiv.) was dissolved in anhydrous DMF (4.5 mL). SO₃.TMA (55 mg, 0.40 mmol, 2.5 equiv.) was added and the reaction was heated at 50 °C for 2 h. After cooling to RT, the reaction was concentrated *in vacuo*. Purification via flash column chromatography (10% to 12% MeOH in DCM) afforded **S9** as a flaky white solid (141 mg, 0.150 mmol, 94%).

Sodium β-D-glucopyranosyluronate-(1→3)-allyl 2-deoxy-6-*O*-sulfonato-2-*N*-acetamine-1-β-D-galactopyranoside **29**

Disaccharide **S9** (110 mg, 0.12 mmol, 1.0 equiv.) was dissolved in THF (9 mL) and cooled to 0 °C. 30% aq. H₂O₂ (0.03 mL, 0.24 mmol, 2.0 equiv.) and 0.7 M aq. LiOH (0.34 mL, 0.24 mmol, 2.0 equiv.) were added and the reaction was slowly warmed to RT and stirred overnight. MeOH (4 mL) and 4 M aq. NaOH (0.60 mL, 2.40 mmol, 20 equiv.) was added. After stirring at RT for 24 h, the reaction was neutralised using Amberlyst IR-20 H⁺ resin. The mixture was filtered and concentrated *in vacuo*. The residue was redissolved in MeOH (8 mL) and cooled to 0 °C. Et₃N (0.15 mL, 1.20 mmol, 10 equiv.) and Ac₂O (0.05 mL, 0.48 mmol, 4.0 equiv.) were added and the reaction was stirred at 0 °C for 1 h. The mixture was concentrated *in vacuo* and co-evaporated with toluene (2 × 10 mL). Purification using Sephadex G25 column afforded **29** as a flaky white solid (13 mg, 0.023 mmol, 19% over three steps).

Acknowledgements

UK Research and Innovation are thanked for project grant funding to G.J.M and C.M (Future Leaders Fellowships MR/T019522/1 and MR/V027018/1, respectively). G.J.M, C.M. and E.F thank EPSRC for a New Horizons Grant (EP/X019136/1). We also thank Keele


University for PhD funding to H.W. and the UK National Mass Spectrometry Facility (NMSF) at Swansea University.

References

- [1] M. Mende, C. Bednarek, M. Wawrystyn, P. Sauter, M. B. Biskup, U. Schepers and S. Bräse, *Chem. Rev.*, **2016**, *116*, 8193–8255.
- [2] M. Kusche-Gullberg and L. Kjellén, *Curr. Opin. Struct. Biol.*, **2003**, *13*, 605–611.
- [3] D. D. Cahyadi, K. Warita, N. Takeda-Okuda, J. ichi Tamura and Y. Z. Hosaka, *Anim. Sci. J.*, **2023**, *94*, 13894.
- [4] M. H. Kim, S. R. Park and B. H. Choi, *Tissue Eng. Regen. Med.*, **2021**, *18*, 165–178.
- [5] Q. Shen, Y. Guo, K. Wang, C. Zhang and Y. Ma, *Molecules*, **2023**, *28*, 7093.
- [6] S. Perez, O. Makshakova, J. Angulo, E. Bedini, A. Bisio, J. L. de Paz, E. Fadda, M. Guerrini, M. Hricovini, M. Hricovini, F. Lisacek, P. M. Nieto, K. Pagel, G. Paiardi, R. Richter, S. A. Samsonov, R. R. Vivès, D. Nikitovic and S. Ricard Blum, *JACS Au* **2023**, *3*, 628–656.
- [7] T. Izumikawa, B. Sato and H. Kitagawa, *Sci. Rep.* **2014**, *4*, 1-12.
- [8] C. J. Mycroft-West, A. J. Devlin, L. C. Cooper, P. Procter, G. J. Miller, D. G. Fernig, M. Guerrini, S. E. Guimond, M. A. Lima, E. A. Yates and M. A. Skidmore, *Neural. Regen. Res.*, **2020**, *15*, 1546–1553.
- [9] T. N. Wight, I. Kang, S. P. Evanko, I. A. Harten, M. Y. Chang, O. M. T. Pearce, C. E. Allen and C. W. Frevert, *Front. Immunol.*, **2020**, *11*, 1-12.
- [10] X. H. Zou, Y. Z. Jiang, G. R. Zhang, H. M. Jin, N. T. M. Hieu and H. W. Ouyang, *Acta Biomater.* **2009**, *5*, 1588–1595.
- [11] N. B. Schwartz and M. S. Domowicz, *Front. Cell Dev. Biol.* **2022**, *10*, 1-14.
- [12] A. P. Asimakopoulou, A. D. Theocharis, G. N. Tzanakakis and N. K. Karamanos, *In Vivo.*, **2008**, *22*, 385–390.
- [13] Y. P. Hu, Y. Q. Zhong, Z. G. Chen, C. Y. Chen, Z. Shi, M. M. L. Zulueta, C. C. Ku, P. Y. Lee, C. C. Wang and S. C. Hung, *J. Am. Chem. Soc.*, **2012**, *134*, 20722–20727.
- [14] M. Baráth, S. U. Hansen, C. E. Dalton, G. C. Jayson, G. J. Miller and J. M. Gardiner, *Molecules* **2015**, *20*, 6167–6180.
- [15] A. Ní Cheallaigh, S. E. Guimond, S. Oscarson and G. J. Miller, *Carbohydr. Res.* **2020**, *495*.
- [16] I. Pongener, C. O’Shea, H. Wootton, M. Watkinson and G. J. Miller, *Chem. Rec.*, **2021**, *21*, 3238–3255.
- [17] G. J. Miller, S. U. Hansen, M. Baráth, C. Johannessen, E. W. Blanch, G. C. Jayson and J. M. Gardiner, *Carbohydr. Res.* **2014**, *400*, 44–53.
- [18] J. C. Jacquinet and C. Lopin-Bon, *Carbohydr. Res.* **2015**, *402*, 35–43.
- [19] S. E. Tully, R. Mabon, C. I. Gama, S. M. Tsai, X. Liu and L. C. Hsieh-Wilson, *J. Am. Chem. Soc.*, **2004**, *126*, 7736–7737.

- [20] G. Macchione, S. Maza, M. Mar Kayser, J. L. De Paz and P. M. Nieto, *Eur. J. Org. Chem.*, **2014**, 2014, 3868–3884.
- [21] Y. Ji, S. Zhang, M. Qiao, R. Jiao, J. Li, P. Song, X. Zhang and H. Huang, *Carbohydr. Polym.*, **2020**, 248, 116796.
- [22] X. Zhang, H. Liu, W. Yao, X. Meng and Z. Li, *J. Org. Chem.*, **2019**, 84, 7418–7425.
- [23] W. Yao, Y. Zhu, X. Zhang, M. Sha, X. Meng and Z. Li, *J. Org. Chem.*, **2018**, 83, 14069–14077.
- [24] I. Pongener and G. J. Miller, *J. Org. Chem.*, **2023**, 88, 11130–11139.
- [25] I. Pongener, E. T. Sletten, J. Danglad-Flores, P. H. Seeberger and G. J. Miller, *Org. Biomol. Chem.*, **2024**, 22, 1395–1399.
- [26] Z. W. ei Poh, C. H. eng Gan, E. J. Lee, S. Guo, G. W. Yip and Y. Lam, *Sci. Rep.* **2015**, 5, 14355.
- [27] N. Karst and J. C. Jacquinet, *Eur. J. Org. Chem.*, **2002**, 815–825.
- [28] M. Wakao, R. Obata, K. Miyachi, Y. Kaitsubata, T. Kondo, C. Sakami and Y. Suda, *Bioorg. Med. Chem. Lett.*, **2015**, 25, 1407–1411.
- [29] A. R. De Jong, B. Hagen, V. Van Der Ark, H. S. Overkleeft, J. D. C. Codée and G. A. Van Der Marel, *J. Org. Chem.*, **2012**, 77, 108–125.
- [30] B. Hagen, J. H. M. Van Dijk, Q. Zhang, H. S. Overkleeft, G. A. Van Der Marel and J. D. C. Codée, *Org. Lett.*, **2017**, 19, 2514–2517.
- [31] F. Demeter, F. Veres, M. Herczeg and A. Borbás, *Eur. J. Org. Chem.*, **2018**, 2018, 6901–6912.
- [32] E. Mancuso, C. Romanò, N. Trattinig, P. Gritsch, P. Kosma and M. H. Clausen, *Chem. Eur. J.*, **2021**, 27, 7099–7102.
- [33] S. Yang, Q. Liu, G. Zhang, X. Zhang, Z. Zhao and P. Lei, *J. Org. Chem.*, **2018**, 83, 5897–5908.
- [34] K. Zegelaar-Jaarsveld, H. I. Duynstee, G. A. Van Der Marel and J. H. Van Boom, *Tetrahedron*, **1996**, 52, 3593–3608.
- [35] D. Z. Vinnitskiy, N. E. Ustyuzhanina, A. S. Dmitrenok, A. S. Shashkov and N. E. Nifantiev, *Carbohydr. Res.*, **2017**, 438, 9–17.
- [36] M. Sakagami and H. Hamana, *Tetrahedron Lett*, **2000**, 41, 5547–5551.
- [37] J. Hénault, P. Quellier, M. Mock-Joubert, C. Le Narvor, A. Alix and D. Bonnaffé, *J. Org. Chem.*, **2022**, 87, 963–973.

Table of Contents Entry



A divergent synthetic route to chondroitin sulfate (CS) disaccharide precursors is presented herein. The route employs an orthogonal protecting group pattern and a post-glycosylation oxidation approach to access CS-A, CS-C, CS-D, CS-E and CS-O disaccharide precursors on gram scale. These materials will provide a benchmark to further synthesise and study chondroitin sulfates.