

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/171904/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Akkaif, Mohammed Ahmed, Daud, Nur Aizati Athirah, Noor, Dzul Azri Mohamed, Sha'aban, Abubakar , Kader, Muhamad Ali SK Abdul and Ibrahim, Baharudin 2024. The impact of CYP2C19 genotype on the platelet reactivity index (PRI) among chronic coronary syndromes (CCS) patients undergoing percutaneous coronary intervention (PCI): Affectability of rapid genetic testing. *Cardiovascular Drugs and Therapy* 10.1007/s10557-024-07544-6

Publishers page: <http://dx.doi.org/10.1007/s10557-024-07544-6>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.






Fig 1. Flow Chart Summary of the study. PCR: Polymerase chain reaction; LOF: Loss of function; NMR: Nuclear magnetic resonance; LD: Loading dose; PRI: Platelet reactivity index; PCI: percutaneous coronary intervention; VASP: vasodilator-associated stimulated phosphoprotein. Note: **lack of interaction between ethnicity and both LOF alleles and PRI.**




Fig 2. The population of the study. SCAD: Stable coronary artery disease; LOF: Loss of function; LD: Loading dose; PCI: Percutaneous coronary intervention. **Note: lack of interaction between ethnicity and both LOF alleles and PRL**