

OPEN

Epidemiology Publish Ahead of Print

DOI: 10.1097/EDE.0000000000001826

How do early weight trajectories explain social inequalities in lung function in children with cystic fibrosis? A longitudinal interventional disparity effects analysis with time-varying mediators and intermediate confounders.

Daniela K. Schlüter¹, Ruth H. Keogh², Rhian M. Daniel³, Schadrac C. Agbla⁴, David Taylor-Robinson¹

¹Department of Public Health, Policy and Systems, University of Liverpool, Liverpool, UK

²Department of Medical Statistics and Centre for Statistical Methodology, London School of Hygiene and Tropical Medicine, London, UK

³Division of Population Medicine, School of Medicine, Cardiff University, Cardiff, UK

⁴Department of Health Data Science, University of Liverpool, Liverpool, UK

Corresponding author Daniela K. Schlüter Department of Public Health, Policy and Systems University of Liverpool Liverpool UK e-mail: d.k.schlueter@liverpool.ac.uk

Suggested running head: Early weight trajectories as drivers of inequalities in CF

Sources of Financial Support

DKS and DTR were supported by the Strategic Research Centre “CF-EpiNet: Harnessing data to improve lives” funded by the Cystic Fibrosis Trust. DTR is funded by the MRC on a Clinician Scientist Fellowship (MR/P008577/1). RHK is supported by a UKRI Future Leaders Fellowship (MR/S017968/1). RMD is supported by a Sir Henry Dale Fellowship from the Wellcome Trust and The Royal Society (107617/Z/15/Z).

Conflicts of interest None declared

Process by which someone else could obtain the data and computing code

The data used in this study are from the UK CF Registry and are available from the UK CF Registry on reasonable request. The computing code can be found in the supplementary material.

Acknowledgments

We thank the UK CF Registry team, and the UK CF centers and clinics for submitting data to the Registry. Special thanks to the people with cystic fibrosis and their families who have agreed for their UK CF Registry data to be used for research.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ACCEPTED

Abstract

Background

Children with cystic fibrosis (CF) from socioeconomically deprived areas have poorer growth, worse lung function, and shorter life expectancy than their less-deprived peers. While early growth is associated with lung function around age 6, it is unclear whether improving early growth in the most deprived children reduces inequalities in lung function.

Methods

We used data from the UK CF Registry, tracking children born 2000-2010 up to 2016. We extended the interventional disparity effects approach to the setting of a longitudinally measured mediator. Applying this approach, we estimated the association between socio-economic deprivation (children in the least versus most deprived population quintile; exposure) and lung function at first measurement (ages 6-8, outcome), and the role of early weight trajectories (ages 0-6) as mediators of this relationship. We adjusted for baseline confounding by sex, birthyear, and genotype and time-varying intermediate confounding by lung infection.

Results

The study included 853 children, with 165 children from the least and 172 from the most deprived quintiles. The average lung function difference between the least and most deprived quintiles was 4.5% of predicted forced expiratory volume in 1 second (95% CI: 1.1 - 7.9). If the distribution of early weight trajectories in the most deprived children matched that in the least deprived children, this difference would reduce to 4% (95% CI: 0.57 – 7.4).

Conclusion

Socio-economic deprivation has a strong negative association with lung function for children with CF. We estimate that improving early weight trajectories in the most deprived children would only marginally reduce these inequalities.

Keywords: cystic fibrosis, health inequalities, health disparities, socio-economic conditions, deprivation, interventional disparity measures, interventional effects

ACCEPTED

Introduction

Cystic fibrosis (CF) is a serious inherited condition affecting approximately 1 in 2,500 babies in the UK and over 100,000 people worldwide^{1,2}. CF is a genetic disease caused by variants of the CF transmembrane conductance regulator (CFTR) gene. Symptoms include the build-up of thick, sticky mucus in various organs, in particular those of the respiratory and digestive systems.

There is no cure for CF and most people die prematurely from their disease through respiratory failure. The median life expectancy for a baby born with CF in the UK in 2022 is 56 years³.

Nutritional management is an essential part of the multidisciplinary care people with CF receive to preserve health and slow decline⁴. In the majority of people with CF mucus blocks the passage of enzymes from the pancreas to the small intestine. In addition, an imbalance of the secretion of bicarbonate and acid in the digestive system and reduced gastrointestinal motility all lead to malabsorption and maldigestion of fat, protein, and fat-soluble vitamins⁴⁻⁶. Thus, achieving and maintaining good nutritional status and therefore healthy growth is difficult for many people with CF and is further impacted by increased energy requirements due to infections and worsening lung function⁴.

Child growth and nutritional status in early life as measured by weight-for-age, height-for-age, and percent ideal body weight, are positively associated with lung function at age 6 for children with CF and it has been suggested that interventions aimed at improving child growth and nutrition in early life may improve later pulmonary function⁷. In children with CF diagnosed within 6 months after birth and who were malnourished at diagnosis, weight-for-age improvements to reach their birthweight percentile by age 2 is associated with improved lung function at age 6, while maintenance of weight-for age, height-for age and BMI percentiles from ages 2 to 6 has no strong additional effect on pulmonary function⁸.

Children with CF from socio-economically deprived areas in the UK have worse outcomes than those from more affluent areas⁹⁻¹¹. They have lower nutritional status as measured by weight-for-age, height-for-age, and BMI from infancy compared to children from less deprived areas. Lower lung function is also evident in children from more deprived areas as soon as this can be routinely measured at around age 6⁹. Although there is some evidence from other countries that these inequalities, the unfair differences in outcomes between children with CF from socio-economically more and less deprived backgrounds, may widen with age^{12,13}, no evidence has emerged that this is the case in the UK¹⁴. Therefore, understanding the pathways that lead to inequalities in lung function at age 6 is crucial to identifying potential targets for interventions that could reduce inequalities in CF across the life-course.

In this study we were interested in examining how unequal growth and nutrition in the first 6 years of life can explain inequalities in lung function (when first measured around age 6) between groups of children from areas with different levels of socio-economic deprivation in the UK CF population. More specifically, we estimated the association between socio-economic deprivation at birth and lung function at first measurement between ages 6-8, and the extent to which this could be reduced if we were to intervene to improve the weight trajectories, as markers of growth and nutrition, in the most deprived children such that they have the same distribution as those in the least deprived children.

To estimate the effects of interest, we build on recent developments in the causal inference literature that have led to the definition of “interventional disparity effects”¹⁵ and extend this approach to the setting with a longitudinally measured mediator.

Methods

Data Source

We used data from the UK CF Registry, which has been collecting information since the 1990s and is estimated to cover 99% of the current UK CF population¹⁶. It records data from comprehensive annual reviews of individuals with CF, including assessments of lung function, nutritional status, and the microbiology of respiratory tract secretions since the previous review. Demographic details, including socio-economic status, are also documented. Our dataset includes follow-up data up to the end of 2016.

Setting and Participants

We included children who were born after 2000, were diagnosed by newborn screening, had at least one lung function measure between ages 6-8 years and at least one weight measurement between birth and <7 years of age, i.e. before or at the same time as the lung function measure. We excluded individuals who did not have complete data on sex, genotype, year of birth and socio-economic conditions.

Outcome, exposure, mediator, and confounders

Figure 1 shows for the outcome, exposure, mediator and confounders, their assumed relationships and when they are measured. Our outcome of interest was the first available lung function measurement taken between age 6 and up to but excluding age 9 (i.e. between ages 6-8) given by percent of predicted forced expiratory volume in one second (%FEV1) based on Global Lung Function Initiative reference equations, which adjust for sex, age, height and ethnicity¹⁷. %FEV1 is treated as a continuous variable.

Our exposure of interest was socio-economic conditions at birth. Measures of individual socio-economic conditions are not available in the UK CF Registry. As a proxy for socio-economic conditions at birth, we used the Index of Multiple Deprivation, a relative measure of deprivation at small-area level linked to first home-address postcode recorded in the registry. This index combines information on income, employment, health, education, access to services, crime/safety and environment/housing¹⁸⁻²¹. Since the domains differ between England, Wales, Scotland, and Northern Ireland, we calculated country-specific Index of Multiple Deprivation z-scores¹¹ to obtain a comparable measure of deprivation across these regions. All children were included in the study, but our main interest was the difference in outcome between children in the highest and lowest quintile of deprivation (see eAppendix; <http://links.lww.com/EDE/C210> for details). We therefore converted z-scores to quintiles.

Our mediator of interest was weight measured at annual review visits from birth up to but excluding age 7 as a proxy for growth and nutrition. We used weight-for-age z-scores based on the UK WHO reference²² as a continuous variable. BMI percentile is the most appropriate measure of nutrition in children with CF however it cannot be used in children under 2 years of age as no reference values are available in this age group²³. BMI might also mask the effect of stunting because height is adjusted for. For these reasons we used weight-for-age z-scores in this study.

Confounders were identified based on our assumed causal relationships between socio-economic conditions, weight, and lung function, and their common causes (Figure 1). We included sex (male/female), birthyear (continuous) and genotype (categorical: number of F508del alleles – 0,1,2. F508del is the most common mutation in CF and people with two copies of F508del have generally the most severe and those with no F508del copy the least severe disease). Respiratory

infection may be a time-varying confounder of the mediator–outcome relationship affected by the exposure⁹. We only considered infection by *Pseudomonas aeruginosa*, the most common cause of infection in the UK CF population under study²⁴. At any annual review visit, it is recorded whether an infection with *P. aeruginosa* had been present since the last annual review visit.

Statistical methods

Rationale and effects of interest

We were interested in estimating the association between socio-economic conditions at birth (high versus low) and lung function at first measurement (between ages 6-8) and the extent to which it can be explained by early weight trajectories. Figure 1 illustrates the assumed causal relationships between socio-economic conditions, weight, first lung function measurement and their common causes, including baseline and time-varying confounders of the mediator–outcome relationship that are also affected by exposure (intermediate confounders), and latent variables, which are assumed present but not observed.

Deprivation at birth as measured by the Index of Multiple Deprivation is multi-factorial and determined by a number of historical factors, and it is difficult to conceive of a single intervention (specifically one that would satisfy the consistency condition^{25–27}) that could plausibly be implemented for subsets of the CF population and could shift this exposure. However, one can conceive of interventions that could be implemented to influence weight gain for children with CF, which is downstream of the effect of socio-economic conditions at birth and also related to lung function at around age 6. This motivates defining our effects of interest in terms of interventional disparity effects, which are a variant of interventional effects^{28,29}. They are based on the ideas of counterfactual disparity measures by VanderWeele & Robinson³⁰ and

Naimi et al³¹ and were formally defined in the context of multiple mediators by Micali et al¹⁵. Interventional disparity effect measures define the indirect effects in terms of the effect on the expected outcome in the exposed group if we could intervene on a mediator so as to shift the mediator distribution in the exposed to that in the unexposed; direct effects are defined in terms of the association that would remain between exposure and expected outcome if we could intervene on a mediator so as to shift the mediator distribution in the exposed to that in the unexposed. In our context, this enables us to estimate how the association between socio-economic conditions at birth and lung function at first measurement between ages 6-8 would change if we could intervene on weight trajectory up to age six. In contrast to interventional effects, interventional disparity measures do not give a causal interpretation to the estimate of the effect of the exposure on the outcome. To ensure this is clear, we will refrain from using the term ‘direct effect’ throughout the rest of the article and will rather refer to it as ‘residual association’.

We further extend the approach of interventional disparity measures by considering a hypothetical intervention on a longitudinal mediator trajectory. This is similar to the extension of interventional effects to incorporate time-varying exposures and mediators^{32,33}.

Let X be the exposure of interest, here socio-economic conditions at birth. We are interested in the contrast between those born in the least and most deprived quintiles of the population. Let $X = 0$ indicate that children were born in the least deprived quintile and $X = 1$ indicate that children were born in the most deprived quintile. Let Y be the outcome (first lung function measurement at age [6-9]) and C the vector of baseline covariates (sex, year of birth, genotype). Let M_t be a time-varying mediator (weight) and L_t a time-varying confounder of the mediator–outcome relationship affected by the exposure (infection), where t denotes age. The complete set of

mediator measurements up to age T, when the outcome is measured, is denoted $\bar{M} = (M_1, \dots, M_T)$ and similarly $\bar{L} = (L_1, \dots, L_T)$. Let \bar{M}_c^x denote a random draw from the distribution of mediator trajectories in those where $X=x$ and $\mathbf{C}=\mathbf{c}$. For justification of conditioning the draws on \mathbf{C} , see Supplementary Material. The interventional disparity measure residual association (*IDM-RA*) is the difference in the expected lung function between children from the least and most deprived quintiles, if the weight trajectories of all children were randomly drawn from the distribution of those observed in the least deprived children within strata of baseline confounders, standardized to the distribution of baseline confounders across both groups of children. Formally:

$$IDM - RA = \sum_c \{E[Y(\bar{M}_c^0)|X = 0, \mathbf{C} = \mathbf{c}] - E[Y(\bar{M}_c^0)|X = 1, \mathbf{C} = \mathbf{c}]\}P(\mathbf{C} = \mathbf{c})$$

The interventional disparity measure indirect effect (*IDM-IE*) is the difference in expected lung function amongst the most deprived children if we were to randomly draw their weight trajectories from the distribution of these in least deprived children versus if we were to randomly draw their weight trajectories from the distribution of these in the most deprived children, again standardized to the distribution of baseline confounders across both groups of children. Formally:

$$IDM - IE = \sum_c \{E[Y(\bar{M}_c^0)|X = 1, \mathbf{C} = \mathbf{c}] - E[Y(\bar{M}_c^1)|X = 1, \mathbf{C} = \mathbf{c}]\}P(\mathbf{C} = \mathbf{c})$$

Finally, the total adjusted association (*Adj-TA*) is the difference in expected lung function between the least and most deprived children standardized to the distribution of baseline confounders across the two groups of children. Formally:

$$\begin{aligned} Adj - TA &= \sum_c \{E[Y|X = 0, \mathbf{C} = \mathbf{c}] - E[Y|X = 1, \mathbf{C} = \mathbf{c}]\}P(\mathbf{C} = \mathbf{c}) \\ &= IDM - RA + IDM - IE \end{aligned}$$

In all the estimands corresponding integrals and densities are used for continuous C .

The *IDM-IE* captures the entire effect of socio-economic deprivation on lung function via weight including any effect from deprivation to infection to weight to lung function. However, it does not include any effect of deprivation on lung function via infection if this is not also mediated by weight. For justification of this see Supplementary Material.

Assumptions required for estimation using registry data

We estimate the estimands specified above using UK CF Registry data. This requires several assumptions: 1. no interference, 2. consistency, 3. positivity, and 4. no unmeasured confounding of the mediator–outcome relationship. For explanations of assumptions 1-3 in this context, see eAppendix; <http://links.lww.com/EDE/C210>. The fourth assumption of ‘no unmeasured confounding of the mediator–outcome relationship’ includes both baseline and time-dependent confounders as measured confounders. We include sex, genotype and year of birth as baseline mediator–outcome confounders, and *P. aeruginosa* infection as a time-dependent confounder. However, we are not able to control for development in utero, which we believe could confound the relationship between weight and lung function (Figure 1) but for which we do not have proxy measures in the registry data. We will assess the implications of this in the discussion.

As we do not aim to estimate the causal effect of the exposure on the outcome, the assumptions required for identifiability of these effects are weaker than for other causal mediation estimands¹⁵. In particular, we do not have to assume no unmeasured confounding of the exposure–outcome relationship.

Assuming all four conditions are met, we can identify the effects of interest from the observed data (eAppendix; <http://links.lww.com/EDE/C210>).

Estimation

Full details of the estimation procedure are given in the eAppendix;

<http://links.lww.com/EDE/C210>. We provide an overview here. We discretized time and assumed that one weight and infection measure per year were recorded as is the guideline in the UK. Our discrete time intervals ran from one birthday up to but not including the next birthday. For example, measurements recorded after a child's sixth birthday but before they turned 7, were labelled as measurements at age 6. We used Monte Carlo simulation to estimate the effects of interest^{15,29}. This required a number of parametric models to be specified: (1) for the expected outcome Y conditional on exposure, mediator at ages 0-6, baseline confounders and time-varying intermediate confounders at ages 0-6; (2) for the distribution of the intermediate confounder L_t at each age 0-6 conditional on exposure, baseline confounders, mediator value at the previous age and previous intermediate confounder value; and (3) for the distribution of the mediator M_t at ages 0-6 conditional on exposure, previous mediator value, baseline confounders and intermediate confounder value at the same age. We fitted these models to data from all children in the study. See Supplementary Material for the detailed model specification and goodness of model fit.

For each individual i in the least and most deprived groups, random values were drawn from the fitted distributions for the mediator M_t and the intermediate confounder L_t at each age $t=0, \dots, 6$ sequentially, conditional on baseline confounder values c_i , previous mediator and intermediate confounder values, and $X=0,1$. The expected outcome was then sampled for each individual based on the fitted distribution for Y , conditional on their baseline confounder values c_i , $X=0,1$, and the previously generated draws m_t and l_t for $t=0-6$. The above process was repeated 1200 times to obtain 1200 draws of the mediators, intermediate confounders, and outcome Y (under

$X=0,1$) for each individual. We chose the number of repetitions (1200) to render negligible the Monte Carlo error at two decimal places. We then averaged the conditional mean outcomes to give the standardized means of the potential outcomes, which we then inserted into the equations for the *IDM-RA*, *IDM-IE*, and *Adj-TA* to estimate the effects of interest. We used non-parametric bootstrap (1,000 bootstrap samples) for variance estimation.

Duplicate entries and missing data

In some cases, children had multiple measurements in a year, for example if they moved between clinics. In these cases, we selected the data from the visit closest to the child's next birthday that had complete data on weight and infection status.

We used multiple multivariate imputation by chained equations³⁴ to impute data if weight and infection status were missing for any years (10 imputed datasets). See eAppendix; <http://links.lww.com/EDE/C210> for the imputation model specification and goodness of fit. We applied the estimation algorithm to each imputed dataset and pooled point estimates across all 10 datasets. To estimate the 95% confidence intervals, we applied standard multiple imputation procedures³⁵ using the variance estimated across bootstrap samples as approximation of the within imputation variance³⁶. See eAppendix; <http://links.lww.com/EDE/C210> for full details.

Sensitivity Analyses

Lung function is measured with considerable error, due to natural day-to-day variation and measurement error. In a sensitivity analysis we used all available lung function data up to age 12 for individuals in the study population and fitted a linear mixed effects model with random intercept and random slope adjusted for sex, genotype, year of birth and deprivation quintile.

Based on this model we predicted lung function at age 6 for all individuals which we used as an

alternative outcome. This also explores the sensitivity of the results to the variability in the time point of outcome measurement.

Due to the relatively small sample size, the parametric models specified for the estimation procedure did not include interaction terms or higher-order terms but we conducted a sensitivity analysis, where we included interaction terms between deprivation and all covariates in the parametric models (1)-(3) described above.

Ethical Considerations

NHS research ethics approval (Huntingdon Research Ethics Committee 07/Q0104/2) was granted for the collection of data into the UK CF Registry. The CF Registry committee approved the use of anonymised data in this study.

Results

Study population

In our population, 853 children met the eligibility criteria (Figure 2) including 165 in the least deprived quartile and 172 children in the most deprived quintiles (Table 1).

Only 139 individuals (16%) had complete weight data across all ages (15% and 16% in the least and most deprived quintiles, respectively) (Table1). 766 individuals (90%) had three or more measurements, including 154 (93%) and 155 (90%) individuals from the least and most deprived quintiles, respectively. Missing measurements were primarily due to missing review visits, although a small number of review visits (87) did not record weight data (18 and 17 in the least and most deprived quintiles of children, respectively). Amongst the observed data, children in the most deprived quintile were on average slightly lighter than those in the least deprived quintile across almost all ages 0-6 (Table 1, Figure 3). The proportion of children ever

diagnosed with *Pseudomonas* was similar across deprivation quintiles; the same was the case for mean age at first infection.

There was a social gradient in lung function at first measurement. Mean lung function was 93%FEV1 (SD: 14) and 89 %FEV1 (SD: 18) in the least and most deprived quintiles, respectively (mean difference 4.6, 95% CI: 1.2-7.9 %FEV1).

Estimated total adjusted association and interventional disparity measures

The estimated total adjusted association (*Adj-TA*), given as the average difference in lung function between the least and most deprived quintile of children, was 4.5 percentage points (95% CI: 1.1-7.9) (Table 2). The estimated interventional disparity measure indirect effect (*IDM-IE*) via weight trajectories at ages 0-6 was 0.53 percentage points (95% CI: -0.64-1.7). This is the estimated amount by which lung function would increase on average in children from the most deprived quintile of the population if their early weight could be improved to have the same distribution as that of children from the least deprived quintile of the population. The estimated interventional disparity measure residual association (*IDM-RA*) was 4 percentage points (95% CI: 0.57–7.4). The difference in lung function at first measurement between children from the least and most deprived quintiles of the population would therefore be on average 4 percentage points (95% CI: 0.57–7.4) even if the weight at ages 0-6 was comparable between the least and the most deprived quintiles of children.

Sensitivity analyses

The *Adj-TA*, *IDM-RA*, and *IDM-IE* from the sensitivity analyses are given in Table 2. The results are similar to the main results.

Discussion

We assessed the relationship between socio-economic conditions at birth and lung function at first measurement in children with cystic fibrosis (CF) in the UK, and the extent to which early inequalities in lung function can be explained by differences in weight trajectories from birth.

We estimated that inequalities in early lung function would only be marginally reduced (4 %FEV1 (95% CI: 0.57–7.4) versus 4.5 %FEV1 (95% CI: 1.1-7.9)) if early weight trajectories could be improved for children from the most socio-economically deprived backgrounds.

To address this question, we extended the framework of interventional disparity effects to the setting of a longitudinal mediator and time-varying intermediate confounder. This allowed us to separate the association between socio-economic conditions and lung function into the effects via distinct, longitudinally measured pathways, but without giving a causal interpretation to the estimate of the effect of the exposure. It therefore requires weaker assumptions than more commonly used mediation methods, and may help gain insights in other areas of health inequalities research and social epidemiology where the consistency and no-unmeasured confounding (of the exposure and mediator and or/outcome) assumptions are often not satisfied^{31,37}.

Strength and Limitations

A strength of this study is that it uses the UK CF Registry data, which collects data on 99% of the current CF population. We excluded just under 200 children because of missing data; the majority of these (156) had no lung function measurement primarily because their first lung function measurement had not yet been taken at the end of our follow-up in 2016. We are therefore confident that the exclusion of individuals with missing data did not lead to bias and that our study population is representative of the UK CF population of children born 2000-2010.

The data used in this study are recorded systematically at clinical review visits for the purposes of reporting and research and are therefore more reliable than many observational datasets.

Measurement error in lung function is, however, nevertheless likely to be non-negligible. This error is non-differential and would therefore not be expected to lead to biased estimates³⁸. We also conducted a sensitivity analysis where we used longitudinally collected lung function data to get an estimate of the error-free measure at age six. As expected, the results agreed with the results from our main analysis.

The major limitation of this study is that our results rely on untestable assumptions, namely, “no interference”, “consistency” and “no unmeasured confounding of the mediator–outcome relationship”. In our setting, the “no interference” assumption states that the lung function of one child is not affected by the weight trajectory of another. This is plausible especially as children with CF are encouraged to not meet in person to prevent cross-infections. The second assumption, consistency, makes assumptions about the hypothetical intervention that would shift the weight distribution in the most deprived children. We can imagine here a potentially complex intervention that could combine financial support with aspects of higher levels of training, engagement and support from dieticians for children from socio-economically deprived backgrounds and their families. The consistency assumption states that this intervention would not change the outcome for those children whose weight trajectory was set to the trajectory that was actually observed. However, it is conceivable that stronger engagement and more support for more deprived families would improve several other factors which may lead to improved lung function, such as clinic attendance and self-confidence in other aspects of CF care as well as reduced parental stress and improved mental health. Therefore, although we did not find any evidence that a hypothetical intervention on weight would substantially reduce inequalities in

lung function, it may have positive effects beyond any direct effects on weight. Finally, we had to assume no unmeasured confounding of the mediator–outcome relationship. According to our assumed causal relationships (Figure 1), it is plausible that development in-utero impacts early weight and lung development and there is some evidence that supports that birthweight (but not gestational age) is associated with lung function at age six^{39,40}. Unfortunately, the UK CF registry has only recently started collecting data on birthweight and there was no proxy in the dataset that could be used to assess the impact of ignoring development in-utero in our analysis. However, as the effect of growth in-utero would have positive effects on early growth and lung function, then any hypothetical intervention on weight after birth would likely have even less of an impact than our estimated effect.

Our analysis further relies on the correct specification of the regression models used in the estimation procedure. Due to the small sample size, we did not include interactions and higher order terms in our models for the main analysis. However, we did conduct a sensitivity analysis in which we included interactions between our exposure and all covariates in the models. This did not substantially change our results although the uncertainty around estimates was increased. A further potential limitation of the study is the use of the area-based Index of Multiple Deprivation rather than an individual measure of socio-economic conditions. Some children living in the most deprived areas may be from affluent families and vice versa. Neighborhood-level deprivation is a relevant target for public health action. However, in further work analyses should also be undertaken at individual level, for example by linking family income or maternal education to the CF Registry data or collecting these data in the registry. Finally, there are differences in socio-economic conditions between countries, for example people in Wales are on average more deprived than those in England. Our current approach using within country

deprivation z-scores does not take this into account and therefore our analysis may be subject to exposure misclassification bias. However, approximately 75% of our study population is from England. Therefore, we expect this bias to have only a small effect if any. Other approaches that make the comparison of deprivation scores between countries possible, for example by Abel et al⁴¹, require data that we did not have available in the UK CF Registry.

Comparison with other studies and implications for further research and clinical practice

Weight in the first years of life has been shown to be strongly associated with later lung function in children with CF^{7,8,42-45} which has led to the development of CF clinical practice guidelines for nutrition management⁴⁶. However, this is the first study that we are aware of that has looked at the role of early weight in the generation of socio-economic inequalities in lung function in CF. Although optimal nutrition is undoubtedly important, we found that reducing inequalities in early weight would likely only have a marginal effect on inequalities in later lung function. Other factors that may lie on the pathway from socio-economic conditions to lung function have previously been studied, including the prescription of antibiotics⁹, exposure to environmental tobacco smoke^{47,48}, and age at diagnosis¹¹. However, it remains unclear what drives inequalities in early lung function and further research is needed. There is anecdotal evidence that the regularity of clinic attendance may display a social pattern due to barriers that more socio-economically deprived families face regarding transportation and the ability to take time off work. Also, engagement with the clinical care team and health literacy of parents from more deprived backgrounds may be lower than in less deprived families and may have a role to play^{49,50}. In addition, environmental factors and living conditions need further consideration.

In the absence of well understood mediators of early inequalities, policy and practice should focus on addressing upstream drivers of child health inequalities⁵¹. Our results also highlight that it is important for the clinical care team of children with CF to be mindful of families' circumstances and focus on optimizing care and support for children with CF from the most deprived backgrounds. This involves paying special attention to the children's respiratory health independent of their growth and nutrition.

Conclusion

There is a strong association between socio-economic conditions at birth and early lung function for children with CF and we estimated that this would only be marginally reduced if early weight trajectories could be improved for children from the most deprived backgrounds.

The methods developed and applied here may also help to give insights into the generation of inequalities in other conditions.

ACCEPTED

References

1. Dodge JA, Morison S, Lewis PA, et al. Incidence, population, and survival of cystic fibrosis in the UK, 1968–95. *Arch Dis Child*. 1997;77(6):493-496. doi:10.1136/adc.77.6.493
2. Guo J, Garratt A, Hill A. Worldwide rates of diagnosis and effective treatment for cystic fibrosis. *J Cyst Fibros Off J Eur Cyst Fibros Soc*. 2022;21(3):456-462. doi:10.1016/j.jcf.2022.01.009
3. UK Cystic Fibrosis Registry - 2022 Annual Data Report. Published online 2022.
4. Nutrition Working Group. *Nutritional Management of Cystic Fibrosis*.; 2016:60. <https://www.cysticfibrosis.org.uk/sites/default/files/2020-12/Nutritional%20Management%20of%20cystic%20fibrosis%20Sep%2016.pdf>
5. Davis TA, Miller A, Hachem C, Velez C, Patel D. The current state of gastrointestinal motility evaluation in cystic fibrosis: a comprehensive literature review. *Transl Gastroenterol Hepatol*. 2023;9:10. doi:10.21037/tgh-23-59
6. Sabharwal S. Gastrointestinal Manifestations of Cystic Fibrosis. *Gastroenterol Hepatol*. 2016;12(1):43-47.
7. Konstan MW, Butler SM, Wohl MEB, et al. Growth and nutritional indexes in early life predict pulmonary function in cystic fibrosis. *J Pediatr*. 2003;142(6):624-630. doi:10.1067/mpd.2003.152
8. Lai HJ, Shoff SM, Farrell PM. Recovery of birth weight z-score within two years of diagnosis is positively associated with pulmonary status at age six years in children with cystic fibrosis. *Pediatrics*. 2009;123(2):714-722. doi:10.1542/peds.2007-3089
9. Taylor-Robinson DC, Smyth RL, Diggle PJ, Whitehead M. The effect of social deprivation on clinical outcomes and the use of treatments in the UK cystic fibrosis population: a

longitudinal study. *Lancet Respir Med*. 2013;1(2):121-128. doi:10.1016/S2213-2600(13)70002-X

10. Barr HL, Britton J, Smyth AR, Fogarty AW. Association between socioeconomic status, sex, and age at death from cystic fibrosis in England and Wales (1959 to 2008): cross sectional study. *BMJ*. 2011;343:d4662. doi:10.1136/bmj.d4662
11. Schlüter DK, Southern KW, Dryden C, Diggle P, Taylor-Robinson D. Impact of newborn screening on outcomes and social inequalities in cystic fibrosis: a UK CF registry-based study. *Thorax*. 2020;75(2):123-131. doi:10.1136/thoraxjnl-2019-213179
12. Taylor-Robinson DC, Thielen K, Pressler T, et al. Low socioeconomic status is associated with worse lung function in the Danish cystic fibrosis population. *Eur Respir J*. 2014;44(5):1363-1366. doi:10.1183/09031936.00063714
13. Schechter MS, Shelton BJ, Margolis PA, Fitzsimmons SC. The association of socioeconomic status with outcomes in cystic fibrosis patients in the United States. *Am J Respir Crit Care Med*. 2001;163(6):1331-1337. doi:10.1164/ajrccm.163.6.9912100
14. Taylor-Robinson D, Diggle P, Smyth R, Whitehead M. Health Trajectories in People with Cystic Fibrosis in the UK: Exploring the Effect of Social Deprivation. In: Burton-Jeangros C, Cullati S, Sacker A, Blane D, eds. *A Life Course Perspective on Health Trajectories and Transitions*. Springer; 2015. Accessed May 25, 2021.
<http://www.ncbi.nlm.nih.gov/books/NBK385365/>
15. Micali N, Daniel RM, Ploubidis GB, De Stavola BL. Maternal Prepregnancy Weight Status and Adolescent Eating Disorder Behaviors: A Longitudinal Study of Risk Pathways. *Epidemiol Camb Mass*. 2018;29(4):579-589. doi:10.1097/EDE.0000000000000850

16. Taylor-Robinson D, Archangelidi O, Carr SB, et al. Data Resource Profile: The UK Cystic Fibrosis Registry. *Int J Epidemiol.* 2018;47(1):9-10e. doi:10.1093/ije/dyx196
17. Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J.* 2012;40(6):1324-1343. doi:10.1183/09031936.00080312
18. English indices of deprivation. GOV.UK. Accessed May 27, 2021. <https://www.gov.uk/government/collections/english-indices-of-deprivation>
19. Welsh Index of Multiple Deprivation. Accessed May 27, 2021. <https://statswales.gov.wales/Catalogue/Community-Safety-and-Social-Inclusion/Welsh-Index-of-Multiple-Deprivation>
20. SpatialData.gov.scot SG. Scottish Index of Multiple Deprivation (SIMD) 2016. Published January 15, 2021. Accessed May 27, 2021. <https://data.gov.uk/dataset/a448dd2a-9197-4ea0-8357-c2c9b3c29591/scottish-index-of-multiple-deprivation-simd-2016>
21. Northern Ireland Multiple Deprivation Measures 2017. GOV.UK. Accessed May 27, 2021. <https://www.gov.uk/government/statistics/northern-ireland-multiple-deprivation-measures-2017>
22. Wright CM, Williams AF, Elliman D, et al. Using the new UK-WHO growth charts. *BMJ.* 2010;340:c1140. doi:10.1136/bmj.c1140
23. Cystic Fibrosis Trust. *Nutritional Management of Cystic Fibrosis.*; 2016. <https://www.cysticfibrosis.org.uk/sites/default/files/2020-12/Nutritional%20Management%20of%20cystic%20fibrosis%20Sep%2016.pdf>

24. UK Cystic Fibrosis Trust. *Annual Data Report 2010.*; 2011. Accessed July 6, 2021.
<https://www.cysticfibrosis.org.uk/sites/default/files/2020-12/2010%20Registry%20Annual%20Data%20Report.pdf>
25. Hernán MA, Robins JM. *Causal Inference: What If?* Boca Raton: Chapman & Hall/CRC; 2020.
26. Hernán MA. Does water kill? A call for less casual causal inferences. *Ann Epidemiol.* 2016;26(10):674-680. doi:10.1016/j.annepidem.2016.08.016
27. Hernán MA, Taubman SL. Does obesity shorten life? The importance of well-defined interventions to answer causal questions. *Int J Obes 2005.* 2008;32 Suppl 3:S8-14. doi:10.1038/ijo.2008.82
28. VanderWeele TJ, Vansteelandt S, Robins JM. Effect decomposition in the presence of an exposure-induced mediator–outcome confounder. *Epidemiol Camb Mass.* 2014;25(2):300-306. doi:10.1097/EDE.0000000000000034
29. Vansteelandt S, Daniel RM. Interventional effects for mediation analysis with multiple mediators. *Epidemiol Camb Mass.* 2017;28(2):258-265. doi:10.1097/EDE.0000000000000596
30. VanderWeele TJ, Robinson WR. On the causal interpretation of race in regressions adjusting for confounding and mediating variables. *Epidemiol Camb Mass.* 2014;25(4):473-484. doi:10.1097/EDE.0000000000000105
31. Naimi AI, Schnitzer ME, Moodie EEM, Bodnar LM. Mediation Analysis for Health Disparities Research. *Am J Epidemiol.* 2016;184(4):315-324. doi:10.1093/aje/kwv329

32. VanderWeele T, Tchetgen ET. Mediation Analysis with Time-Varying Exposures and Mediators. *Harv Univ Biostat Work Pap Ser*. Published online March 4, 2014.
<https://biostats.bepress.com/harvardbiostat/paper168>
33. Lin SH, Young J, Logan R, Tchetgen Tchetgen EJ, VanderWeele TJ. Parametric Mediation g-Formula Approach to Mediation Analysis with Time-varying Exposures, Mediators, and Confounders. *Epidemiol Camb Mass*. 2017;28(2):266-274.
 doi:10.1097/EDE.0000000000000609
34. Buuren S van, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained Equations in R. *J Stat Softw*. 2011;45(1):1-67. doi:10.18637/jss.v045.i03
35. Rubin DB. Multiple Imputation After 18+ Years. *J Am Stat Assoc*. 1996;91(434):473-489.
 doi:10.2307/2291635
36. Schomaker M, Heumann C. Bootstrap inference when using multiple imputation. *Stat Med*. 2018;37(14):2252-2266. doi:10.1002/sim.7654
37. Rehkopf DH, Glymour MM, Osypuk TL. The Consistency Assumption for Causal Inference in Social Epidemiology: When a Rose is Not a Rose. *Curr Epidemiol Rep*. 2016;3(1):63-71. doi:10.1007/s40471-016-0069-5
38. VanderWeele T. *Explanation in Causal Inference: Methods for Mediation and Interaction*. Oxford University Press; 2015.
39. Darrah R, Nelson R, Damato EG, Decker M, Matthews A, Hodges CA. Growth Deficiency in Cystic Fibrosis Is Observable at Birth and Predictive of Early Pulmonary Function. *Biol Res Nurs*. 2016;18(5):498-504. doi:10.1177/1099800416643585
40. Atteih SE, Raraigh KS, Blackman SM, Cutting GR, Collaco JM. Predictive effects of low birth weight and small for gestational age status on respiratory and nutritional outcomes in

cystic fibrosis. *J Cyst Fibros Off J Eur Cyst Fibros Soc.* 2020;19(6):888-895.

doi:10.1016/j.jcf.2020.02.003

41. Abel GA, Barclay ME, Payne RA. Adjusted indices of multiple deprivation to enable comparisons within and between constituent countries of the UK including an illustration using mortality rates. *BMJ Open.* 2016;6(11):e012750. doi:10.1136/bmjopen-2016-012750
42. Peterson ML, Jacobs DR, Milla CE. Longitudinal changes in growth parameters are correlated with changes in pulmonary function in children with cystic fibrosis. *Pediatrics.* 2003;112(3 Pt 1):588-592. doi:10.1542/peds.112.3.588
43. Pedreira C c., Robert R g. d., Dalton V, et al. Association of body composition and lung function in children with cystic fibrosis. *Pediatr Pulmonol.* 2005;39(3):276-280. doi:10.1002/ppul.20162
44. Konstan MW, Morgan WJ, Butler SM, et al. Risk Factors For Rate of Decline in Forced Expiratory Volume in One Second in Children and Adolescents with Cystic Fibrosis. *J Pediatr.* 2007;151(2):134-139.e1. doi:10.1016/j.jpeds.2007.03.006
45. Sanders DB, Fink A, Mayer-Hamblett N, et al. Early Life Growth Trajectories in Cystic Fibrosis are Associated with Pulmonary Function at Age 6 Years. *J Pediatr.* 2015;167(5):1081-1088.e1. doi:10.1016/j.jpeds.2015.07.044
46. Stallings VA, Stark LJ, Robinson KA, Feranchak AP, Quinton H. Evidence-Based Practice Recommendations for Nutrition-Related Management of Children and Adults with Cystic Fibrosis and Pancreatic Insufficiency: Results of a Systematic Review. *J Am Diet Assoc.* 2008;108(5):832-839. doi:10.1016/j.jada.2008.02.020

47. Ong T, Schechter M, Yang J, et al. Socioeconomic Status, Smoke Exposure, and Health Outcomes in Young Children With Cystic Fibrosis. *Pediatrics*. 2017;139(2). doi:10.1542/peds.2016-2730
48. Oates GR, Baker E, Rowe SM, et al. Tobacco smoke exposure and socioeconomic factors are independent predictors of pulmonary decline in pediatric cystic fibrosis. *J Cyst Fibros Off J Eur Cyst Fibros Soc*. 2020;19(5):783-790. doi:10.1016/j.jcf.2020.02.004
49. Jackson AD, Kirwan L, Gibney S, Jeleniewska P, Fletcher G, Doyle G. Associations between health literacy and patient outcomes in adolescents and young adults with cystic fibrosis. *Eur J Public Health*. 2020;30(1):112-118. doi:10.1093/eurpub/ckz148
50. Mercer SW, Higgins M, Bikker AM, et al. General Practitioners' Empathy and Health Outcomes: A Prospective Observational Study of Consultations in Areas of High and Low Deprivation. *Ann Fam Med*. 2016;14(2):117-124. doi:10.1370/afm.1910
51. Pearce A, Dundas R, Whitehead M, Taylor-Robinson D. Pathways to inequalities in child health. *Arch Dis Child*. 2019;104(10):998-1003. doi:10.1136/archdischild-2018-314808

Figure captions

Figure 1: Assumed causal relationships between socio-economic conditions at birth, early weight as marker of growth and nutrition, first lung function measurement and their common causes including baseline and time-varying confounders of the mediator-outcome relationship that are also affected by exposure, and latent variables, which are assumed present but not observed (ellipses).

Figure 2: Derivation of the study population.

Figure 3: Weight for age z-scores by age and Index of Multiple Deprivation (IMD) quintile as given in Table 1.

ACCEPTED

Tables

Table 1: Demographics and clinical characteristics of the study population stratified by deprivation quintile.

	1=least deprived	2	3	4	5=most deprived	Overall
n	165	164	180	172	172	853
Male (%)	95 (58)	76 (46)	89 (49)	79 (46)	94 (55)	433 (51)
F508 class (%)						
Heterozygous	72 (44)	74 (45)	72 (40)	67 (39)	67 (39)	352 (41)
Homozygous	77 (47)	74 (45)	91 (51)	90 (52)	86 (50)	418 (49)
other	16 (10)	16 (10)	17 (9)	15 (9)	19 (11)	83 (10)
Cohort 2007-2010 (%)	107 (65)	110 (67)	120 (67)	115 (67)	115 (67)	567 (67)
First measured lung function mean (SD)	93 (14)	91 (14)	91 (15)	89 (17)	89 (18)	91 (16)
Age at first lung function measure (%)						
6	138 (83)	142 (86)	152 (84)	136 (79)	143 (83)	711 (83)
7	24 (15)	18 (11)	23 (13)	32 (19)	25 (15)	122 (14)
8	3 (2)	4 (2)	5 (3)	4 (2)	4 (2)	20 (2)
Ever diagnosed with pseudomonas (%)	98 (59)	105 (64)	108 (60)	101 (59)	95 (55)	507 (59)
Age at first pseudomonas diagnosis mean (SD)	2.8 (1.6)	3.2 (1.7)	3.0 (1.8)	3.0 (1.7)	2.9 (1.7)	3.0 (1.7)
Number of years with missing weight data (%)						
0	25 (15)	25 (15)	30 (17)	31 (18)	28 (16)	139 (16)
1	65 (39)	62 (38)	60 (33)	56 (33)	61 (36)	304 (36)
2	44 (27)	51 (31)	38 (21)	41 (24)	43 (25)	217 (25)
3	20 (12)	12 (7)	27 (15)	24 (14)	23 (13)	106 (12)
4-6	11 (7)	14 (9)	25 (14)	20 (12)	17 (10)	87 (10)
Weight z-scores mean (SD)						
at age [0 – 1)	-0.35 (1.2)	-0.54 (1.5)	-0.50 (1.5)	-0.45 (1.4)	-0.66 (1.4)	-0.49 (1.4)
at age [1 – 2)	0.32 (0.88)	0.40 (1.1)	0.40 (1.0)	0.27 (0.89)	0.08 (1.1)	0.30 (1.0)
at age [2 – 3)	0.10 (0.92)	0.26 (0.95)	0.37 (0.92)	0.17 (0.88)	0.14 (1.1)	0.21 (0.96)
at age [3 -4)	0.11 (0.92)	0.11 (0.97)	0.18 (0.97)	0.09 (0.94)	-0.04 (1.1)	0.09 (0.97)
at age [4 -5)	-0.17 (0.97)	-0.00 (1.1)	0.03 (1.0)	-0.07 (0.93)	-0.17 (1.1)	-0.08 (1.0)
at age [5 -6)	-0.07 (0.95)	-0.08 (1.0)	-0.03 (0.98)	-0.15 (1.0)	-0.14 (1.1)	-0.09 (1.0)
at age [6 -7)	-0.08 (0.96)	0.01 (1.1)	-0.00 (0.98)	-0.03 (1.0)	-0.18 (1.2)	-0.06 (1.1)


SD=Standard deviation.

Table 2: Estimated total adjusted association and interventional disparity measure direct and indirect effects. In sensitivity analysis 1 we used predicted lung function at age 6 rather than observed lung function as outcome; in sensitivity analysis 2 we included interaction terms between deprivation and all covariates in the models for weight, infection and lung function.

	Main analysis	Sensitivity analysis 1	Sensitivity analysis 2
	Point estimate (95% confidence interval)		
Total adjusted association	4.5 (1.1 – 7.9)	4.3 (1.7 – 7)	4.4 (0.95 – 7.9)
Interventional disparity measure indirect effect (IDM-IE)	0.53 (-0.64 – 1.7)	0.47 (-0.46 – 1.4)	0.23 (-1.6 – 2.1)
Interventional disparity measure residual association (IDM-RA)	4 (0.57 – 7.4)	3.9 (1.2 – 6.5)	4.2 (0.43 – 7.9)

ACCEPTED


Figure 1



Downloaded from <http://journals.lww.com/epidem> by BNDMfseP-HKav12Eoun11QIN4a+kULhEZgshH04XXMI0hCwCX1
AMhYQp/IIQH338D00dRy7TvsF14C3VC1y0ab9gQZXd6G52MwLzleI= on 01/20/2025

ACCEPTED

Figure 2



ACCEPTED

Figure 3

