



Importance of solid earth structure for understanding the evolution of the Greenland ice sheet

Jörg Ebbing^{1*}, John R. Hopper^{2,3}, Clinton P. Conrad⁴, Glenn Milne⁵, Rebekka Steffen⁶, Juan Carlos Afonso⁷, Valentina R. Barletta⁸, Ana M. G. Ferreira⁹, Judith Freienstein¹, Samantha E. Hansen¹⁰, Björn H. Heincke², Glenn Jones^{9,11}, Sergei Lebedev¹², Max Moorkamp¹³, Derek L. Schutt¹⁴ and Agnes Wansing¹

¹ Institute of Geosciences, Kiel University, Otto-Hahn-Platz 1, 24118 Kiel, Germany

² Geological Survey of Denmark and Greenland, (GEUS), Øster Voldgade 10, 1350 Copenhagen, Denmark

³ Department of Geosciences and Natural Resources, University of Copenhagen, Øster Voldgade 10, 1350 Copenhagen, Denmark

⁴ Centre for Planetary Habitability, University of Oslo, Oslo, Norway

⁵ Department of Earth and Environmental Sciences, University of Ottawa, Ottawa, ON K1N 6N5, Canada

⁶ Lantmäteriet (The Swedish Mapping, Cadastral and Land Registration Authority), Lantmäterigatan 2C, 801 82 Gävle, Sweden

⁷ University of Twente, Langezijds Building, Hallenweg 8, 7522 NH Enschede, the Netherlands

⁸ Department of Geodesy and Earth Observations, DTU Space, Lyngby, Denmark

⁹ Department of Earth Sciences, University College London, UK

¹⁰ Department of Geological Sciences, The University of Alabama, Tuscaloosa, AL 35487, USA

¹¹ School of Earth and Environmental Sciences, University of Cardiff CF10 3AT, UK

¹² Department of Earth Sciences, University of Cambridge, Cambridge CB3 0EZ, UK

¹³ Institute for Applied Geosciences, Technical University of Berlin, Ernst Reuter Platz 1, 10587 Berlin, Germany

¹⁴ Department of Geosciences, Colorado State University, Fort Collins, CO 80525, USA

JE, 0000-0001-7492-5338; JRH, 0000-0003-3188-7583; CPC, 0000-0003-4314-2351; GM, 0000-0002-5498-9403; RS, 0000-0003-4739-066X; JCA, 0000-0001-9938-6692; VRB, 0000-0003-4427-0830; AMGF, 0000-0002-9492-6415; JF, 0009-0001-2643-1525; SEH, 0000-0001-7608-5715; BHH, 0000-0002-5271-0333; GJ, 0000-0002-7691-332X; SL, 0000-0003-0004-5009; MM, 0000-0002-7879-1346; DLS, 0000-0002-4504-5662; AW, 0000-0002-5164-7256

* Correspondence: Joerg.ebbing@ifg.uni-kiel.de

Abstract: The solid earth structure beneath Greenland, meaning the rocky part of Earth from the ice-bed interface to depth, has gained increased interest in recent years as it provides a critical boundary condition for the dynamic evolution of the Greenland ice sheet (GrIS), one of the largest sources of sea-level rise contributions since the early 2000s. However, no consensus has been reached regarding the key internal or surface earth properties influencing this boundary condition and thus GrIS behaviour. One important surface property is the subglacial heat flow, which affects sliding conditions of the ice sheet including the onset of major ice streams and is related to subglacial geology. Lithospheric architecture and mantle viscosity structure are internal properties that influence ice sheet evolution through changes in the height and slope of the ice-bed interface caused by glacial isostatic adjustment. Because there is no general agreement regarding crustal and lithospheric structures, some glaciological studies use an ensemble of solid earth models to incorporate uncertainties into their GrIS predictions, but it is unclear how these variations ultimately affect estimates of future sea-level rise. Here we describe the main solid earth properties that are important for GrIS evolution (heat flow, temperature, viscosity), from the base of the ice sheet to the upper mantle, and we provide some perspectives on how future collaborative efforts and integrated studies could lead to better agreement regarding these key characteristics.

Received 26 December 2024; **revised** 8 April 2025; **accepted** 9 April 2025

Meltwater from the Greenland ice sheet (GrIS) has been one of the largest sources of sea-level rise since the early 2000s (e.g. [Bamber *et al.* 2018](#); [Fox-Kemper *et al.* 2021](#); [Mankoff *et al.* 2021](#)). Despite the global mean sea-level contribution of up to $+0.89 \text{ mm a}^{-1}$ ([Horwath *et al.* 2022](#)) and a mass loss of more than 247 Gt a^{-1} (2012–16; [Bamber *et al.* 2018](#)), the driving components behind ice-mass loss and their corresponding feedback mechanisms are still not completely understood. Thus, future ice-mass loss remains one of the largest uncertainties for future sea-level projections (e.g. [Bamber *et al.* 2019](#); [Fox-Kemper *et al.* 2021](#)). It has been demonstrated that subglacial conditions can play an important role in ice-sheet behaviour (e.g. [Bell 2008](#); [Smith-Johnsen *et al.* 2020b](#); [McCormack *et al.* 2022](#)). Sub-ice geology, geothermal heat flow (GHF), topography and topographic changes resulting from glacial isostatic adjustment (GIA) can all influence GrIS evolution. These

factors may be especially important for marine-terminating glaciers, where ice interaction with seawater has been shown to accelerate deglaciation in Greenland ([Wood *et al.* 2021](#)). However, rapid GIA-induced bedrock uplift ([Whitehouse *et al.* 2019](#)) or fjord topography resulting from geological structures may block inflow of warm water by ocean currents ([Jakobsson *et al.* 2020](#)) and can exert a stabilizing influence. Such interactions have been demonstrated as important for Antarctica (e.g. [Book *et al.* 2022](#)), but are poorly constrained for Greenland. Therefore, it is essential to enhance our knowledge of solid earth and cryosphere interactions (e.g. [Bell 2008](#); [Whitehouse *et al.* 2019](#)) and to better constrain the solid earth properties that control them ([Fig. 1](#)).

[Karlsson *et al.* \(2021\)](#), for instance, showed that the present GrIS basal melt production amounts to $c. 21.4 \text{ Gt a}^{-1}$ ($c. 10\%$ of the total mass loss) with an uncertainty ranging from $+4.4$ to -4.0 Gt a^{-1} .




Fig. 1. Main elements of the solid earth structure that can affect cryospheric processes and the related datasets that can constrain upper mantle (left) and crustal (right) structure. HPE, heat-producing elements; ΔT and ΔC indicate temperature and compositional anomalies in the upper mantle; LAB, lithosphere–asthenosphere boundary. Source: modified from [Reading *et al.* \(2022\)](#) mainly by adding geophysical observables.

This *c.* 20% uncertainty is mostly related to the heat flow models used in their calculations. Because the uncertainty of individual models is difficult to assess, [Karlsson *et al.* \(2021\)](#) used an ensemble of models to determine an average heat flow estimate. [Rogozhina *et al.* \(2012\)](#) performed sensitivity tests to determine how different input GHF distributions affected simulations of the present state of the GrIS. Their results indicate a high sensitivity of the GrIS to the input GHF, demonstrating the importance of reducing uncertainty in this quantity. Ultimately, [Rogozhina *et al.* \(2012\)](#) questioned whether or how geophysical models should be used, given their large uncertainties. The conclusions of those researchers are supported by a more recent study that quantified the impact of using seven different GHF models to spin up a GrIS model to an equilibrium state ([Zhang *et al.* 2024](#)).

Additional uncertainties stem from the thermal interactions between the Iceland hotspot track and Greenland's lithosphere. Individual observations suggest high heat flow in central Greenland (*c.* 98 mW m⁻²; [Grinsted and Dahl-Jensen 2002](#)), and some models require even higher local heat flow to sustain the NE Greenland Ice Stream (e.g. [Greve 2019](#)). However, reconstructed hotspot tracks and lithospheric-scale models continue to exhibit significant variability and uncertainty regarding the location of potential heat flow anomalies, leaving the expected magnitude of GHF unresolved (e.g. [Heyn and Conrad 2022](#)). Furthermore, variations in GHF at the ice–bedrock interface are probably accompanied by thermal heterogeneities in the underlying mantle. Such an anomalous mantle structure may contribute to the rapid uplift rates, which are observed by global navigation satellite system (GNSS) data (e.g. [Khan *et al.* 2016](#); [Berg *et al.* 2024](#)). These uplift rates are considerably faster than those predicted by 1D (spherically symmetric) GIA models tuned to fit palaeo sea-level data that reflect the response of the solid earth to GrIS deglaciation since the Last Glacial Maximum (e.g. [Lecavalier *et al.* 2014](#); [Khan *et al.* 2016](#)). This has led to the suggestion of a significant transient component in the deformational response ([Adhikari *et al.* 2021](#); [Paxman *et al.* 2023](#)).

The examples above highlight the need for spatially variable (3D) solid earth models with robust uncertainty estimates that can be

utilized when coupling solid earth models to ice-sheet models. Although significant effort has been made to develop such constraints for Antarctica, especially in terms of incorporating geothermal heat flow ([Reading *et al.* 2022](#)) as well as isostatic and erosional changes ([Whitehouse *et al.* 2019](#)), similar initiatives for Greenland are still missing. The upcoming 5th International Polar Year in 2032–33 should provide an opportunity for coordinated international research to tackle the biggest challenges of polar research, including the development of improved solid earth models to more accurately evaluate past and future GrIS evolution on century to millennial timescales.

The role of solid earth structures for the evolution of the Greenland ice sheet

From a solid earth perspective, the conditions along the base of the ice sheet are the most critical for GrIS evolution. [Bell \(2008\)](#) and [Whitehouse *et al.* \(2019\)](#) described some of the main components. These include the role of subglacial water or basal melt, which may result from elevated GHF, as well as local bedrock conditions. Therefore, not only it is important whether the bed is thawed or frozen, but also knowledge about the rugosity, which relates small-scale changes in topography and the geology at the ice-bed interface and the presence of sediments at the ice-bed interface, is critical. Topography modulates outlet glaciers and controls the state of stress for glaciers, critical for determining the dynamics of the ice-sheets ([Catania and Felikson 2022](#)). Even though substantial improvements have been made in recent years in our knowledge of the topography under the ice (e.g. [Morlighem *et al.* 2017, 2022](#)), data coverage is still sparse in some critical regions. For example, knowledge of local bedrock composition, the crustal thickness and tectonic history are needed to understand the distribution of radioactive elements, which affects basal heat flow (see the next section). In addition, variations in lithosphere thickness also allow for large mantle contributions to GHF, although on a different scale. Additionally, the thermal structure of the upper mantle plays a critical role in determining the viscosity distribution and




Fig. 2. Estimates of Greenland's GHF (top row) and associated basal melting (bottom row) for a suite of models. Corresponding GHF model sources are listed above each column. Areas considered as being frozen are masked. (Note the wide range of basal melt estimates, from 3.8 to 5.7 Gt a⁻¹.) Source: all basal melting estimates were calculated following the method of [Karlsson et al. \(2021\)](#).

lithospheric thickness (e.g. [Paxman et al. 2023](#); [Weerdesteijn and Conrad 2024](#)), which influence the rate and amplitude of isostatic responses, affecting the bed topography. These factors, in turn, affect ice-sheet elevation and surface mass balance (e.g. [van den Berg et al. 2008](#); [Zeitzi et al. 2022](#)). We provide further details on these characteristics, and the geophysical datasets available to constrain them, in the following sections.

Basal melt and heat flow

Conditions at the base of an ice sheet play a key role in governing subglacial water and basal melt. Radar imaging is the most effective tool to determine basal water distribution and to assess whether the ice sheet is thawed or frozen at the base (e.g. [MacGregor et al. 2016](#); [Jordan et al. 2018](#)). These data, however, do not necessarily account for changes in bedrock reflectivity (i.e. the reflection amplitude) owing to varying geology.

Of the *c.* 21.4 Gt a⁻¹ of basal ice melt estimated by [Karlsson et al. \(2021\)](#), about half is attributed to basal friction whereas viscous heat dissipation from surface melt water and GHF each contribute about one quarter. For individual drainage basins, the relative contributions vary, reflecting the different basal conditions but also the uncertainties. These uncertainties make up about 20% of the total basal melt estimate and reflect how poorly GHF models agree. The significant differences in GHF amplitude and spatial distribution are apparent when models that have been published over the last decade are compared (e.g. [Colgan et al. 2022](#)). This comparison translates into highly variable basal melt rate estimates from GHF (see [Fig. 2](#)). In general, the discrepancies between these models reflect the fact that only about 10 observation points from boreholes or ice measurements are available from Greenland's interior and that estimates for one of those observation points are highly debated.

That particular point of contention is the heat flow at the onset of the NE Greenland Ice Stream. For the ice stream, basal melt estimates range from 1 to 2 Gt a⁻¹, with variations mainly

depending on the treatment of the North Greenland Ice core Project (NGRIP) GHF determination. For example, [Dahl-Jensen et al. \(2003\)](#) provided upper and lower bound GHF estimates of 160 and 90 mW m⁻², respectively, for the NGRIP data, based on models of the age of ice layers determined from radar echograms. [Smith-Johnsen et al. \(2020a\)](#) instead argued that very high local heat flow (970 mW m⁻²), well above the suggested range from observations, is needed in this area, to sustain the NE Greenland Ice Stream. [Bons et al. \(2021\)](#) contended that such extreme heat flow is unrealistic, as there is no plausible geological origin for it, and [Freienstein et al. \(2024\)](#) provided a statistical analysis of all available observation points and concluded that the NGRIP data point should be considered with caution as it is statistically an outlier, at least on a regional scale. Questions surrounding the NGRIP observation also provided the motivation for the GHF prediction by [Colgan et al. \(2022\)](#), where all observation points have been compiled and (re-) analysed, to present two alternative heat flow maps, one including and the other excluding the NGRIP value ([Fig. 2](#)).

The resulting GHF maps ([Fig. 2](#)) show that, except in the immediate vicinity of NGRIP, one might be able to provide a reasonable regional baseline heat flow estimate. However, local variations, which are needed to describe the coupling conditions for ice-sheet models, might still be obscured (e.g. [McCormack et al. 2022](#)). Therefore, the prediction of local GHF variations remains a challenge for accurate assessment of GrIS changes. It is worth noting that similar challenges are faced in Antarctica. [Reading et al. \(2022\)](#), for instance, discussed how small-scale heat flow variations are difficult to constrain, and [Stål et al. \(2024\)](#) stressed the importance of gaining a better understanding of subglacial geology to link point observations with regional heat flow models.

Sub-ice geology

Sub-ice geology affects the ice sheet in different ways. Rugosity at the ice-bed interface affects basal sliding conditions; for example, a




Fig. 3. (a) Mapped surface geology and interpretation of sub-ice bedrock in terms of major provinces. Dashed grey line denotes the division of Proterozoic crust. (b) Synthesis of apparent subglacial geological provinces from geophysical boundary analysis. Source: (a) major provinces from Dawes (2009); division of Proterozoic crust from Dahl-Jensen *et al.* (2003); (b) MacGregor *et al.* (2024).

sedimentary layer tends to be less dense and less erosion resistant, and hence smoother, facilitating enhanced basal sliding. Geology affects the transfer of heat from the crystalline crust to the ice, reflecting variations in radiogenic heat production. In sedimentary layers and shallow bedrock, porosity and pore fluid compositions affect the thermal conductivities. Variations of either subglacial topography where the thermal conductivity of glacial ice and the solid Earth differ or in thermal conductivity can lead to thermal refraction at the ice-bed interface (e.g. Willcocks *et al.* 2021). Further, neighbouring tectonic units are often associated with different thermal properties (radiogenic heat production and thermal conductivity), which can influence heat flow on a local scale. Hence, detailed knowledge of the sub-ice geology is a prerequisite for describing the basal conditions of the system accurately.

The subglacial geological setting of Greenland was first evaluated by Dawes (2009; Fig. 3a). That study examined exposed geology in the ice-free coastal regions as well as glacial erratics from GrIS outflow streams. Similar and synchronous formations along the coast and plausible tectonic scenarios were often used to propose boundaries between major geological provinces, but these were typically unconstrained across several hundreds of kilometres. It was also recognized that some features identified by geophysical data were inconsistent with some surface-based inferences (Dawes 2009). For example, a SW–NE-striking division of Proterozoic crust was identified based on seismic analyses from Dahl-Jensen *et al.* (2003), but no expression of this boundary was indicated in the geology map by Dawes (2009).

Recently, MacGregor *et al.* (2024) provided a new map (Fig. 3b) that describes the geological provinces beneath the GrIS based on a synthesis of geophysical data, where 19 geological and geophysical datasets were used to delineate major geological provinces. For example, subglacial topography was considered as a potential constraint on subglacial geology. Most of the differences between the two geological province maps (Fig. 3) are located in North Greenland, where MacGregor *et al.* (2024) inferred a less extensive Ellesmere–Ingfield terrane and the Committee–Melville terrane has been eliminated, despite its onshore exposure. In addition, inferred basins, from which exotic glacial erratics could have originated, have also been eliminated. The erratics could plausibly have come from the extensive ‘unknown’ area on the MacGregor *et al.* (2024) map (see grey area in Fig. 3b). Hence, although MacGregor *et al.* (2024) presented an alternative view of the sub-ice geology, a key shortcoming is that it is inconsistent with the exposed

geology in several areas. For example, the East Greenland rift basins are assigned Devonian–Permian ages, despite the fact that significant and thick deposits of Mesozoic sections are well exposed in these areas (e.g. Stoker *et al.* 2017; Fyhn *et al.* 2021, and references therein). Such considerations have important implications for understanding basal palaeo-heat flow during past deglaciations.

A second issue with the MacGregor *et al.* (2024) map is that it relies on the interpretation of geophysical data that are highly heterogeneous in terms of quality, as acknowledged by the authors. For example, the choice of a seismic velocity model is subjective, as competing and contradicting models exist (Fig. 4; Darbyshire *et al.* 2018; Toyokuni *et al.* 2020; Jones *et al.* 2021).

The model of Darbyshire *et al.* (2018), which was used by MacGregor *et al.* (2024), focused on the overall crustal structure using group velocity measurements from regional earthquakes. In contrast, Jones *et al.* (2021) targeted near-surface anomalies using Rayleigh wave ellipticity measurements to describe the subglacial properties. Further, the Jones *et al.* (2021) model is based on a single-station approach that generates 1D velocity models beneath each station, whereas the Darbyshire *et al.* (2018) model was developed using relatively long path-averaged structures. In contrast to these studies, Toyokuni *et al.* (2020) presented a model based on the analysis of P-wave arrival time data that shows a clear seismic anomaly extending from Iceland to eastern Greenland in the crust and upper mantle, but the model is in general focused on the overall lithospheric architecture.

Although the available seismic velocity models (Fig. 4) show some similarities, such as the relatively high velocities in western Greenland, there are also some notable differences, such as in northeastern Greenland, reflecting the differences in methods and data. Given the relatively large inter-station distances in Greenland, the differences between these models are not surprising. Additionally, each study selected different stations for their analyses to make use of the best-quality data for the specific method applied. Interestingly, neither the map by MacGregor *et al.* (2024) nor the tomographic models show an indication of the SW–NE-striking division of Proterozoic crust that was identified in the earlier study by Dahl-Jensen *et al.* (2003) (Fig. 3a).

In general, seismic station coverage across Greenland is rather sparse and uneven (Fig. 4d), and any seismic model would benefit from additional data. It is interesting to note that the unresolved area indicated by MacGregor *et al.* (2024; Fig. 3b) has comparatively




Fig. 4. (a) Relative V_s variations at 5 km depth. (b) Depth slice at 5 km through the S-wave velocity (V_s) model, as used in the map shown in Figure 3b. It should be noted that the Darbyshire *et al.* (2018) model constrains the V_s from 5 km downwards, whereas the Jones *et al.* (2021) model constrains V_s from 5 km upwards. (c) Relative V_p variations at 5 km depth. Each model used different seismic stations (triangles) in their analyses. (d) All available broadband stations across Greenland. Source: (a) from the model by Jones *et al.* (2021), which was not directly considered by MacGregor *et al.* (2024); (b) model from Darbyshire *et al.* (2018), as used in the map developed by MacGregor *et al.* (2024) shown in Figure 3b; (c) from the model by Toyokuni *et al.* (2020); (d) T. Dahl-Jensen, pers. comm., 2024.

dense seismic coverage compared with other regions. That said, other data types that were used by MacGregor *et al.* (2024), such as the magnetic field anomalies acquired from the Earth Magnetic Anomaly Grid (Meyer *et al.* 2017), have relatively crude coverage in this region. Magnetic data are arguably the most sensitive to upper crustal structure and are thus helpful in interpreting sub-ice geology (e.g. Aitken *et al.* 2014; Brethes *et al.* 2018; Golynsky *et al.* 2018). Despite continuing efforts to reprocess the magnetic datasets for Greenland (Heincke *et al.* 2023) to create a new compilation, the lack of adequate, modern high-resolution data hampers the possibility to trace geological structures from the coast and beneath the ice, thereby limiting our ability to use magnetic data for an accurate prediction of variations in subglacial geology.

Crustal and lithospheric architecture

Improved constraints on the crustal and overall lithospheric architecture beneath Greenland are critical for us to gain a better understanding of the solid earth–ice interaction because the lithospheric structure is controlled by both rheological and thermal properties. A number of studies have evaluated lithospheric variability beneath Greenland (e.g. Steffen *et al.* 2018; Artemieva 2019; Wansing *et al.* 2024), using a range of data types and

approaches. Most models are based on gravity measurements, seismic velocities and/or petrophysical data. Considerable differences exist between resulting models that relate to both station (data) coverage and methodological differences.

As an example, seismic estimates of crust–mantle boundary (Moho) depth display large disagreements, even between models based on the same dataset. For instance, Dahl-Jensen *et al.* (2003) analysed P-wave receiver functions for 20 broadband seismic stations to estimate the crustal structure beneath central Greenland and found Moho depths ranging from 23 to 50 km, which were interpreted to reflect different tectonic blocks. Kumar *et al.* (2007) reinterpreted the same P-wave receiver function data and added S-wave receiver functions, as they are less affected by multiples in the ice layer. They found significantly shallower Moho depths beneath central Greenland, which deviate by up to 11 km. The largest discrepancies were found for stations deployed on ice and can probably be attributed to ice layer effects on the seismic signal (for further details see Wansing *et al.* 2024).

Models estimating the depth of the lithosphere–asthenosphere boundary (LAB) are even more diverse (Fig. 5), also owing to differences between employed datasets and different methods. Whereas earlier and especially low-resolution global models (e.g. Priestley and McKenzie 2013; Pasyanos *et al.* 2014; Afonso *et al.* 2019) tend to disagree in the estimated lithospheric thickness, most




Fig. 5. Lithospheric thickness estimates for Greenland highlighting the thermal structure of the lithosphere. Models (e)–(h) are global models, models (a)–(d) are recent Greenland models. Model (a) is based on direct conversion from seismic velocities to lithosphere thickness, while Models (b)–(d) are based on a combination of multiple, yet different, geophysical datasets. Panel (i) highlights the agreement between these models in terms of their predicted depths to the thermal lithosphere–asthenosphere boundary (LAB) by showing the number of models predicting LAB > 150 km depth. Source: Corresponding models are listed above the panels.

recent models (e.g. Steinberger *et al.* 2019; Wansing *et al.* 2024; Salajegheh *et al.* 2025) show similar variation patterns and image the cratonic cores in North and South Greenland in similar locations. In general, the thermal base of the lithosphere appears to be deeper than 150 km for most of Greenland's interior, and significantly thinner lithosphere (<120 km) is found only beneath the south-eastern Greenland coast (see Figs 5 and 6).

Role of the Icelandic hotspot track and its link to lithospheric architecture

Greenland is thought to have passed over the Iceland plume between about 80–65 and 50 myr ago, forming a hotspot track that crosses the island (e.g. Martos *et al.* 2018; Steinberger *et al.* 2019). Several possible tracks have been identified based on geological observations, heat flow constraints and tectonic reconstructions (see Fig. 6, and references cited by Martos *et al.* 2018), and although these paths roughly converge at the southeastern Greenland margin, they diverge towards the west or NW (Fig. 6). This partly relates to the unknown subglacial geology (see above), as any direct expression of volcanism is covered by ice and the geophysical signatures are ambiguous. As outlined by Larsen *et al.* (2015), intraplate volcanism in Greenland and the North Atlantic region occurred simultaneously and over a wide area, particularly around 60 myr ago, but this volcanism shows no resemblance to a hotspot track,

suggesting instead that plume material (head or tail) injected beneath Greenland was channelled into lithospheric thin spots from Mesozoic rift basins along both West and East Greenland (Nielsen *et al.* 2002; Horní *et al.* 2017; Steinberger *et al.* 2019).

However, heat from the Iceland plume may also have thinned and weakened parts of Greenland's lithosphere. For example, Steinberger *et al.* (2019) based their interpretation of the hotspot track on the tomographic model from Lebedev *et al.* (2018), in combination with plate reconstructions and numerical models of mantle flow. These predict east–west palaeo-flow along a corridor of thinned lithosphere, seen in central Greenland by most tomographic models (Fig. 5). In contrast, more recent seismic investigations (e.g. Celli *et al.* 2021) suggest the possibility of alternative plume track pathways to the NW (Fig. 6).

Heyn and Conrad (2022) made numerical models of a mantle plume impinging on continental lithosphere and suggested that the interaction tends to thin the lithosphere along the plume track. The amount of thinning significantly depends on the characteristics of the plume, the lithosphere and the underlying asthenosphere, with stronger plumes and weaker rheologies generating more thinning. The researchers also found that the resulting increase in heat flux depends on the extent of lithospheric thinning, but for Greenland, the heat flux increase is probably limited to at most about 20% of the pre-thinning heat flux. This finding agrees with evidence for only moderately increased heat flow in the interior of the island (e.g. Martos *et al.* 2018). However, the




Fig. 6. Upper mantle velocity structure, uplift rates (red points) and possible Iceland plume-track (green lines). Background colours show variations in seismic velocity at 150 km depth. Faster seismic anomalies generally correlate with colder temperatures but also depend on mineral composition. Red circles mark the positions of GNSS stations, with their size proportional to the average crustal uplift rate. Green lines indicate possible paths of the Iceland hotspot beneath Greenland. Source: variations in seismic velocity at 150 km depth from the model by [Celli et al. \(2021\)](#); red circles from [Berg et al. \(2024\)](#); green lines from summary by [Martos et al. \(2018\)](#).

plume should still significantly affect the thermal structure of the lower lithosphere, especially if the heat is transported by melt ([Heyn et al. 2024](#)), and this added heat may have left a low-viscosity zone in the upper mantle. If so, the plume may significantly accelerate rates of uplift following periods of deglaciation (see the next section and [Weerdesteijn and Conrad 2024](#)).

The importance of Greenland's upper mantle structure for GIA

Greenland GIA is constrained via both geological and geodetic observations. Through modelling these observations, inferences have been made on GrIS evolution and earth viscosity structure (e.g. [Tarasov and Peltier 2002](#); [Fleming and Lambeck 2004](#); [Lecavalier et al. 2014](#); [Khan et al. 2016](#)). This section focuses on the application of geodetic data, and how our knowledge of upper mantle structure is critical to inferring GrIS changes via GIA modelling of these data.

Uplift rates in Greenland reflect the isostatic (GIA) response to both contemporary (during the GNSS monitoring period) and past ice-mass changes. The latter includes signals associated with regional-scale deglaciation since the Last Glacial Maximum (e.g. [Simpson et al. 2011](#); [Khan et al. 2016](#)) as well as signals owing to more recent, lower amplitude changes, such as the Little Ice Age ([Kjeldsen et al. 2015](#); [Adhikari et al. 2021](#)).

Contemporary ice-mass loss across Greenland can be estimated using satellite altimetry ([Simonsen et al. 2021](#); [Khan et al. 2022](#)) and

from satellite observations of temporal changes in the gravity field (e.g. from the GRACE-FO twin satellites). These estimates require a correction for GIA-related earth deformation, which is about 5–10% of the total signal (e.g. [Wake et al. 2016](#); [Barletta et al. 2024](#)). Thus, an accurate estimate of the GIA signal is necessary to produce an accurate determination of contemporary ice-mass changes, which are required for sea-level budget calculations and are an important initial condition for projecting future changes in the GrIS.

Several researchers have noted that areas of lithospheric thinning near Greenland's SE coast coincide with areas of rapid uplift rates observed from GNSS networks (e.g. [Bevis et al. 2012](#); [Khan et al. 2016](#); [Adhikari et al. 2021](#); [Paxman et al. 2023](#); [Berg et al. 2024](#); [Weerdesteijn and Conrad 2024](#); Fig. 6). This highlights the emerging consensus that GIA uplift rates are sensitive to variations in lithospheric thickness and mantle viscosity, which in turn are a result of temperature and compositional variations within the Earth. Lithospheric thickness and mantle viscosity may be reduced in areas that have been heated, as, for example, the Iceland plume may have done for SE Greenland. The 3D rheological viscosity structure beneath Greenland may thus control the rates of coastal uplift following deglaciation. Including 3D viscosity models in GIA calculations is computationally expensive and so relatively few cases have been explored to date. One study showed that uplift rates and postglacial sea-level changes are significantly affected by the 3D viscosity structure (e.g. [Milne et al. 2018](#)). If viscosities are reduced beneath the lithosphere (e.g. 100–200 km wide regions with viscosity of $c. 10^{19}$ Pa s), modern-day deglaciation rates can already induce a rapid viscous uplift of a few centimetres per year ([Weerdesteijn et al. 2022](#)). Furthermore, [Weerdesteijn and Conrad \(2024\)](#) showed that recent deglaciation of Greenland drives unusually rapid uplift when positioned above a low-viscosity plume track beneath SE Greenland.

In addition, recent studies have suggested that the viscosity beneath Greenland may be timescale dependent (e.g. [Adhikari et al. 2021](#); [Paxman et al. 2023](#)). That is, there may be a significant transient component to the deformational response such that the (apparent) viscosity is lower for shorter timescale ice sheet changes. The existence of a transient signal has been proposed to explain why GIA models tuned to fit postglacial sea-level observations produce a poor fit to GNSS-determined uplift rates ([Adhikari et al. 2021](#)). Including a transient component increases the contribution of ice-mass changes during and following the Little Ice Age ([Kjeldsen et al. 2015](#)) to match contemporary uplift rates. On the other hand, [Pan et al. \(2024\)](#) noted that including a thin and weak asthenospheric layer can alternatively explain rapid modern uplift rates, without timescale-dependent rheology. Clearly, improved constraints on the thermal and compositional structure of the Earth from non-GIA approaches (e.g. [Wansing et al. 2024](#)) are necessary to reduce the ambiguity in the interpretation of GIA datasets and thus allow more robust constraints on GrIS evolution.

Future sea-level predictions are sensitive not only to our understanding of the GrIS evolution (e.g. [Höning et al. 2023](#)) but also to global GIA patterns (e.g. [Spada 2017](#)), which in turn cannot be well understood without the Greenland contribution. They are also probably dependent on feedbacks between GIA and ice-sheet evolution (e.g. [Whitehouse et al. 2019](#)). Such feedbacks have been proposed for Antarctica ([Adhikari et al. 2014](#); [Kingslake et al. 2018](#); [Albrecht et al. 2024](#)) and are probably also important for Greenland, but have not been adequately explored.

The way forward

We have outlined some of the key elements from a solid earth perspective that influence the evolution of the GrIS. The question is, where to proceed from here? First of all, better data coverage and improved data processing are essential to advance our knowledge of

Greenland. Some of the critical parameters cannot be measured directly (e.g. viscosity), whereas others (e.g. heat flow) are difficult to acquire beyond measurements in selected spots (boreholes).

However, geophysical data from seismic, magnetotelluric, gravity and magnetic methods are, compared with a large number of boreholes, relatively cheap and feasible to acquire. For example, between the first and second generations of the Antarctic magnetic anomaly map (Golynsky *et al.* 2006, 2018) more than two million line kilometres of airborne data have been acquired, despite the more challenging logistics in Antarctica. If the same amount of data is acquired over Greenland with a regular spacing, the entirety of Greenland would be covered with 1 km profile distance.

With improved spatial coverage of geophysical measurements and their integrated interpretation, the key parameters for the coupling of the solid earth and ice sheet models could be estimated with lower uncertainty, when considered in an integrated manner (e.g. Wansing *et al.* 2024).

Subglacial conditions

The recent map by MacGregor *et al.* (2024) has demonstrated the potential of synthesizing boundaries of different datasets to image the subglacial setting for Greenland. A similar approach has mapped sedimentary basins (Aitken *et al.* 2023) and lithospheric architecture (Stål *et al.* 2019) for Antarctica. Whereas such maps represent a valuable first-order step, these predictions should be coupled with and tested by a physical earth model grounded in geological and tectonic knowledge. Modern approaches that consistently combine different physical models in joint inversions and employ thermodynamic models for the crustal and overall lithospheric architecture hold the potential to enhance our knowledge of the critical parameters beneath the GrIS (e.g. Fullea *et al.* 2021; Afonso *et al.* 2022; Moorkamp 2022; Lebedev *et al.* 2024).

Of course, the accuracy of any model depends on data quality and coverage. Magnetic data are one of the datasets most sensitive to the near-surface geology, but large areas of Greenland are still not well covered. For example, Wansing *et al.* (2024) showed a stepwise inversion for the lithospheric and crustal architecture, and the resulting density and magnetic susceptibility distribution in general agrees with petrophysical data. However, the structure of the model can be described as patchy at best and is not an adequate representation of the complexities of subglacial geology. This is especially true compared with detailed studies in Antarctica (e.g. Aitken *et al.* 2014; Lowe *et al.* 2024), where for large regions the data coverage and quality is far better owing to continuing efforts both in acquisition and coordination (e.g. SCAR Expert Group ADMAP or SCAR RINGS Action Group).

As mentioned above, complete coverage of Greenland would require a dedicated coordinated multi-national effort, but in the short term, a more localized airborne campaign near the onset of the NE Greenland Ice Stream (in the vicinity of the NGRIP stations) might be achievable. This area seems to be critical for deciphering the role of the subglacial geology for geothermal heat flow, but also for understanding the evolution of the ice sheet. Beyond the ice sheet modelling mentioned above (e.g. Smith-Johnsen *et al.* 2020a), a recent local radar study, which argued for a recent onset of the NE Greenland Ice Stream (Jansen *et al.* 2024), would rule out a massive heat flow influence, but still would allow sub-ice geology to exert a controlling influence.

Any dedicated gravity and magnetic airborne mission should be accompanied by further work on petrophysical, seismic and electromagnetic datasets. Petrophysical samples for Greenland can provide constraints on LAB depth (e.g. Fig. 1; Lee *et al.* 2009) and are available mostly from the coastal, ice-free areas. Yet, despite massive collections of rock samples from decades of (geological) mapping or from drilling projects (e.g. Christiansen *et al.* 2024), no

tailored petrophysical database for Greenland is yet available. However, magnetic susceptibility, rock density, seismic velocity, thermal conductivity and heat production are important parameters to benchmark crustal models based on potential field airborne data and to predict thermal parameters that are important for heat flow under the GrIS.

The solid earth architecture and its link to viscosity

In addition to the conditions at the ice-bed interface, an improved knowledge of the lithospheric architecture is also needed. Seismic data provide one of the main constraints on the crustal and lithospheric architecture of Greenland (e.g. Dahl-Jensen *et al.* 2003; Darbyshire *et al.* 2018; Lebedev *et al.* 2018; Mordret 2018; Toyokuni *et al.* 2020; Celli *et al.* 2021; Jones *et al.* 2021; Ajournalou *et al.* 2024; Salajegheh *et al.* 2025). The coverage with broadband stations for Greenland is certainly sparse compared with well-covered regions in Europe or the USA, but not all differences between models can be explained by this. A major difference is that often only a subset of available stations is used, and this selection varies from study to study as seen in the examples above.

Part of this incongruity between seismic studies is related to differences in the acquisition period between stations, as well as where and how the data are available. Here, a common reprocessing format and the establishment of a reference database would be useful, where metadata are properly described and tailored, based on which different processing methods can be easily compared. Such an initiative is currently being carried out but should be complemented by data acquisition in some of the key areas, especially near the coast. In addition, individual seismological studies often use specific types of methods, such as single station techniques, to build 1D models beneath stations (e.g. receiver functions, Rayleigh wave ellipticity) versus multiple station approaches that constrain inter-station averaged seismic structures (e.g. earthquake and ambient noise tomography). In regions with large inter-station distances such as in Greenland, it is unsurprising that using these different approaches leads to conflicting results, as discussed above. Future work should focus on joint inversions of the seismic observables used with these complementary techniques (e.g. dispersion measurements, ellipticity, receiver functions, amplification) to tighten the constraints on subsurface structures. Further, to explore whether the high uplift rates along the southeastern coast of Greenland are related to the Iceland hotspot, a tighter coupling of offshore and onshore studies is also needed.

Methodological improvements may also be helpful, both for seismological methods (e.g. Lebedev *et al.* 2024) as well as for integrated approaches. For example, seismic tomography maps velocity variations in the mantle (e.g. Fig. 6), which depend strongly on temperature, but the conversion factor between velocity and temperature is nonunique (e.g. Lu *et al.* 2020; Lebedev *et al.* 2024). Thermodynamic inversion methods that use computational petrology and thermodynamic databases can avert much of the non-uniqueness and resolve the thermal structure and thickness of the lithosphere based on the Rayleigh and Love surface-wave data, as well as Pn data (e.g. Schutt *et al.* 2018; Porter and Reid 2021; Lebedev *et al.* 2024). The methods are also effective for implementing the joint inversion of seismic and other data for both temperature and composition (e.g. Afonso *et al.* 2013a, b, 2022; Fullea *et al.* 2021).

Other geophysical observables, such as magnetotelluric (MT) data, may become increasingly important for constraining Greenland's upper mantle structures. MT data, which are collected locally in temporary (days-long) deployments, provide constraints on electrical conductivity down to upper mantle depths. Whereas the usefulness of MT data has already been demonstrated for Antarctica (e.g. Peacock and Selway 2016; Wannamaker *et al.* 2017), so far only a few measurement points over Greenland are available; yet

MT data present the possibility, especially in combination with the data discussed above, to provide a more detailed image of the crustal architecture (e.g. Liu and Hasterok 2016; Moorkamp 2022). For example, Ramirez *et al.* (2022) and Manassero *et al.* (2024) have recently shown that the combination of MT and seismic data offers useful constraints to both the temperature and water content of upper mantle rocks. Because these two factors represent the most important controls on rock viscosity, imaging of the subsurface using both MT and seismic methods can provide a new constraint on the viscosity structure of the upper mantle. Initial tests of the joint use of MT and seismic data for Scandinavia, where observations of surface uplift provide independent constraints on viscosity, are promising (Ramirez *et al.* 2024). Thus, a combination of seismic and MT data may complement, or even provide critical input to, numerical studies of GIA. Here, Central and SE Greenland, which were most probably affected by the Iceland plume, represent key areas of interest.

Improvements in the imaging of the Earth structure will improve GIA models. Milne *et al.* (2018) showed the importance of using 3D GIA models for Greenland, but with more recent estimates of seismic velocity variations as well as better constraints on lithospheric thicknesses and improved GIA modelling capabilities (e.g. Weerdesteijn *et al.* 2023) these models can be enhanced. In addition, the determination of certain structures in the lithosphere and mantle based on various geophysical datasets would give GIA modellers a unique set of boundary conditions, in line with upcoming initiatives, such as the GIA Model Intercomparison Project (GIAMIP). Thus, discrepancies between GIA models could be reduced, which in turn affects mass balance as well as sea-level rise estimates.

Modelling and measurement perspectives

The main challenge limiting our understanding of the solid earth beneath the GrIS is data coverage and lack of accurate descriptions of vintage datasets. Still, on the regional scale, models converge towards similar results, as shown here for lithospheric thickness, but also for heat flow estimates, despite the unusual NGRIP measurements. In contrast, much remains unknown with respect to the local structure beneath the ice, which is needed to gain a better understanding and prediction of ice-sheet evolution. As explained above, MacGregor *et al.* (2024) provided a first approach of mapping subglacial geology, which should be extended by more advanced statistical methods (e.g. based on machine learning methods, e.g. Li *et al.* 2022), but also coupled with 3D Earth models (e.g. Lowe *et al.* 2024).

In addition, dedicated efforts should be made by the research community to acquire key datasets in some of the most vulnerable and least understood regions of Greenland. Ideally, a combination of airborne data with ground-based seismic and MT installations should be acquired to fill some of the gaps in Central Greenland, which appears to be the least understood region. This links to efforts in southeastern Greenland, where an improved understanding of ground-based observations (e.g. GNSS) and regional data requires an improved understanding of the lithospheric structure from Iceland towards Greenland. Recent discussion on an AtlanticArray (e.g. Ferreira 2024) spanning this region can only be supported from a Greenland perspective.

In addition to such campaigns, there are also possibilities from upcoming (e.g. Next Generation Gravity Mission) or candidate (e.g. CryoRad) satellite missions. The advantage of satellite missions is the monitoring of the ice sheets in terms of mass, temperature and height changes, which allows us to decipher short-term and long-term effects, providing important information for testing structural models in dynamic modelling. However, such data and models also require detailed information from the solid earth. An example is the ice temperature models based on the SMOS (Soil Moisture and

Ocean Salinity) satellite mission (e.g. Macelloni *et al.* 2019), which rely on geothermal heat flow as constraint, but so far on models with low confidence. The gravity field missions (GRACE and GRACE-FO) provided a unique method to estimate the ice-mass changes for Greenland (for references, see, for example, Velicogna 2009; Harig and Simons 2012; Velicogna *et al.* 2014), but their results are dependent on GIA models (e.g. Caron *et al.* 2018). Although these missions can complement studies of the GrIS as well as Greenland's lithospheric architecture and dynamic processes, they cannot replace stations on the ground.

Conclusions and recommendations

In 2014, Kennicutt *et al.* (2015) defined a roadmap with key questions for Antarctic science for the next two decades, which raised awareness and motivated a number of studies dedicated to Antarctica. Without attempting to reproduce such a detailed analysis, we will highlight some of the key questions identified by Kennicutt *et al.* (2015) that are also relevant for Greenland (it should be noted that we changed the questions by omitting links to the Antarctic ice sheet as these are also valid for the Greenland region).

- (1) *Do variations in geothermal heat flux provide a diagnostic signature of sub-ice geology?*

Yes, they do, but the absence of heat flow measurements means that sub-ice geology is taken as a proxy to derive geothermal heat flow.

- (2) *What is the crust and mantle structure, and how does it affect surface motions owing to glacial isostatic adjustment?*

Whereas recent models of LAB depth show convergence on the regional scale, there is still a disconnect between geophysical and GIA models of the lithosphere and mantle, which has to be addressed to more accurately interpret observations of postglacial sea-level changes and present-day land motion. In addition, future imaging of the crust and mantle should focus not only on elastic, isotropic structures, but also on other properties, such as, for example, anisotropy (to better understand stress in the crust and mantle flow) and attenuation (to obtain independent information on temperature).

- (3) *How does volcanism affect the evolution of the lithosphere, ice sheet dynamics and global climate?*

If we refer here to the Iceland hotspot and its interaction with Greenland, the location of the hotspot track beneath Greenland remains unclear before Greenland moved away from above the hotspot around 50 myr ago. The lithospheric architecture appears to show an imprint of the hotspot track, but the thermal effect at a crustal level is still debated.

- (4) *How do the characteristics of the ice sheet bed, such as geothermal heat flux and sediment distribution, affect ice flow and ice sheet stability?*

Here, we are still at an infancy stage for Greenland, as detailed models of the ice sheet bed are missing. For Antarctica, this has been addressed, for example, by McCormack *et al.* (2022) showing the importance of local variations on subglacial conditions.

- (5) *How do tectonics, dynamic topography, ice loading and isostatic adjustment affect the spatial pattern of sea-level change on all timescales?*

Changes to Greenland's ice load drive glacial isostatic adjustment (GIA), which proceeds more rapidly in areas with thinner lithosphere or reduced sub-lithospheric viscosity (e.g. Milne *et al.* 2018; Weerdesteijn *et al.* 2022). Thus, regions subjected to mantle heating, for example, from the Iceland plume, will experience rapid bedrock uplift soon after deglaciation, and slower uplift later after deglaciation is

completed (Weerdesteijn and Conrad 2024). Because sub-lithospheric viscosity mostly influences uplift rates at relatively short (hundreds of kilometres) wavelengths (Pan *et al.* 2024), these variations in uplift rate are likely to be regionally important for uplift and sea-level change along Greenland's coastline. For coastlines distant from Greenland, Greenlandic deglaciation drives sea-level change through its net mass loss and via spatial variations associated with the 'sea-level fingerprint', which accounts for Earth's gravitational and rotational changes in addition to solid earth deformation (e.g. Mitrović *et al.* 2011). The sea-level fingerprint has already been detected for modern deglaciation in Greenland (Coulson *et al.* 2022) and drove significant sea-level variability following the Last Glacial Maximum (e.g. Lin *et al.* 2021).

(6) *How will the sedimentary record beneath the ice sheet inform our knowledge of the presence or absence of continental ice?*

Mapping the subglacial conditions in Greenland is a prime target to understand the feedback between the solid earth and cryosphere, but requires dedicated campaigns to fill some of the most critical (e.g. seismic and magnetic) data gaps to verify the location and thickness of sedimentary basins.

Although not all of these questions have been answered even in the case of Antarctica, substantial improvement in our understanding of the processes has been made and many of the lessons learned for the Antarctic ice sheet hold for the GrIS. In Antarctica, a number of dedicated international initiatives have been running for a long time or have been initiated in recent years (e.g. Frémand *et al.* 2022; Colleoni *et al.* 2024).

Some of the initiatives almost date back to IPY-3 (1957–58), which is an especially good example for coordinated research activities. IPY-3 was part of the International Geophysical Year and led to the first geophysical traverse from the coast of East Antarctica to the interior of the continent and can be considered as the start of geophysical surveying of Antarctica (Dodds *et al.* 2010), which later led to a number of multinational efforts that have advanced our knowledge of the continent. Similar coordinated efforts for Antarctica were running under the umbrella of IPY-4, which ran from 2007 to 2009. IPY-4 involved over 200 projects examining a wide range of physical, biological and social research topics for both poles. However, research in the Arctic was largely dedicated to processes related to the Arctic Ocean, but none of the projects was dedicated to Greenland or its structure beneath the ice sheets (see list at <https://www.ipy.org/projects>). This omission was at the time certainly related to the fact that the melting of the GrIS was just becoming apparent in satellite data and the discussion of its implication for sea-level rise was at its beginning (e.g. Dowdeswell 2006; Rignot and Kanagaratnam 2006). As we now are aware of the drastic changes in Greenland mass balance in recent years and with the 5th International Polar Year in 2032–33 on the horizon, we emphasize the urgent need for coordinated international research to advance our understanding of the solid earth structure beneath, and its interaction with, the Greenland ice sheet.

Scientific editing by Yildirim Dilek

Acknowledgements We thank the participants of the Greenland workshop held in March 2024, which motivated this paper. We furthermore thank the editor, Y. Dilek, and the reviewer, V. Klemann, for their insightful comments and recommendations.

Author contributions JE: conceptualization (equal), visualization (equal), writing – original draft (lead); JRH: conceptualization (equal), writing – original draft (equal); CPC: visualization (supporting), writing – original draft

(equal); GM: writing – original draft (equal); RS: writing – original draft (equal); JCA: writing – original draft (equal); VRB: writing – original draft (equal); AMGF: writing – original draft (equal); JF: visualization (equal), writing – original draft (supporting); SEH: writing – original draft (equal); BHH: writing – original draft (equal); GJ: writing – original draft (equal); SL: writing – original draft (equal); MM: writing – original draft (equal); DLS: writing – original draft (equal); AW: visualization (equal), writing – original draft (equal).

Funding J.E., A.W. and J.F. acknowledge funding from the ESA STSE 4D Greenland and the German Science Foundation (DFG, project 675325 and 535728086). C.P.C. acknowledges funding from the Research Council of Norway (PHAB Centre of Excellence, project 332523). R.S. acknowledges funding from Rymdstyrelsen (Swedish National Space Board; grant number 2018-00140). D.L.S. acknowledges funding from the National Science Foundation, grant EAR-1925595, and the Fulbright Foundation of Iceland. A.M.G.F. is supported by the UPFLOW project, funded by the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101001601). S.L. is supported by NERC grants NE/X000060/1 and NE/Y000218/1, ESA project 4D Dynamic Earth (4000140327/23/NL/SD) and Project InnerSpace (<https://projectinnerspace.org>).

Competing interests The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability The datasets shown can be downloaded from the original publications. Basal melt rates from Figure 2 are available on request from the authors.

References

- Adhikari, S., Ivins, E.R., Larour, E., Seroussi, H., Morlighem, M. and Nowicki, S. 2014. Future Antarctic bed topography and its implications for ice sheet dynamics. *Solid Earth*, **5**, 569–584, <https://doi.org/10.5194/se-5-569-2014>
- Adhikari, S., Milne, G.A. *et al.* 2021. Decadal to centennial timescale mantle viscosity inferred from modern crustal uplift rates in Greenland. *Geophysical Research Letters*, **48**, e2021GL094040, <https://doi.org/10.1029/2021GL094040>
- Afonso, J.C., Fullea, J., Griffin, W.L., Yang, Y.J., Jones, A.G., Connolly, J.A.D. and O'Reilly, S.Y. 2013a. 3-D multiobservable probabilistic inversion for the compositional and thermal structure of the lithosphere and upper mantle. I: a priori petrological information and geophysical observables. *Journal of Geophysical Research: Solid Earth*, **118**, 2586–2617, <https://doi.org/10.1002/jgrb.50124>
- Afonso, J.C., Fullea, J., Yang, Y.J., Connolly, J.A.D. and Jones, A.G. 2013b. 3-D multi-observable probabilistic inversion for the compositional and thermal structure of the lithosphere and upper mantle. II: general methodology and resolution analysis. *Journal of Geophysical Research: Solid Earth*, **118**, 1650–1676, <https://doi.org/10.1002/jgrb.50123>
- Afonso, J.C., Salajegheh, F., Szwillus, W., Ebbing, J. and Gaina, C. 2019. A global reference model of the lithosphere and upper mantle from joint inversion and analysis of multiple data sets. *Geophysical Journal International*, **217**, 1602–1628, <https://doi.org/10.1093/gji/ggz094>
- Afonso, J.C., Ben-Mansour, W. *et al.* 2022. Thermochemical structure and evolution of cratonic lithosphere in central and southern Africa. *Nature Geoscience*, **15**, 405–410, <https://doi.org/10.1038/s41561-022-00929-y>
- Aitken, A.R.A., Young, D.A. *et al.* 2014. The subglacial geology of Wilkes Land, East Antarctica. *Geophysical Research Letters*, **41**, 2390–2400, <https://doi.org/10.1002/2014GL059405>
- Aitken, A.R.A., Li, L. *et al.* 2023. Antarctic sedimentary basins and their influence on ice-sheet dynamics. *Reviews of Geophysics*, **61**, e2021RG000767, <https://doi.org/10.1029/2021RG000767>
- Ajourlou, P., Darbyshire, F., Audet, P. and Milne, G.A. 2024. Structure of the crust and upper mantle in Greenland and northeastern Canada: insights from anisotropic Rayleigh-wave tomography. *Geophysical Journal International*, **239**, 329–350, <https://doi.org/10.1093/gji/ggac269>
- Albrecht, T., Bagge, M. and Klemann, V. 2024. Feedback mechanisms controlling Antarctic glacial-cycle dynamics simulated with a coupled ice sheet–solid Earth model. *Cryosphere*, **18**, 4233–4255, <https://doi.org/10.5194/tc-18-4233-2024>
- Artemieva, I.M. 2019. Lithosphere thermal thickness and geothermal heat flux in Greenland from a new thermal isostasy method. *Earth-Science Reviews*, **188**, 469–481, <https://doi.org/10.1016/j.earscirev.2018.10.015>
- Bamber, J.L., Westaway, R.M., Marzeion, B. and Wouters, B. 2018. The land ice contribution to sea level during the satellite era. *Environmental Research Letters*, **13**, 063008, <https://doi.org/10.1088/1748-9326/aac2f0>
- Bamber, J.L., Oppenheimer, M., Kopp, R.E., Aspinall, W.P. and Cooke, R.M. 2019. Ice sheet contributions to future sea-level rise from structured expert judgment. *Proceedings of the National Academy of Sciences of the USA*, **116**, 11195–11200, <https://doi.org/10.1073/pnas.1817205116>

- Barletta, V.R., Bordoni, A. and Khan, S.A. 2024. GNET derived mass balance and glacial isostatic adjustment constraints for Greenland. *Geophysical Research Letters*, **51**, e2023GL106891, <https://doi.org/10.1029/2023GL106891>
- Bell, R.E. 2008. The role of subglacial water in ice-sheet mass balance. *Nature Geoscience*, **1**, 297–304, <https://doi.org/10.1038/ngeo186>
- Berg, D., Barletta, V.R. *et al.* 2024. Vertical land motion due to present-day ice loss from Greenland's and Canada's peripheral glaciers. *Geophysical Research Letters*, **51**, <https://doi.org/10.1029/2023GL104851>
- Bevis, M. *et al.* 2012. Bedrock displacements in Greenland manifest ice mass variations, climate cycles and climate change. *Proceedings of the National Academy of Sciences of the USA*, **109**, 11944–11948, <https://doi.org/10.1073/pnas.1204664109>
- Bons, P.D., de Riese, T. *et al.* 2021. Comment on “Exceptionally high heat flux needed to sustain the Northeast Greenland Ice Stream” by Smith-Johnsen *et al.* (2020). *Cryosphere*, **15**, 2251–2254, <https://doi.org/10.5194/tc-15-2251-2021>
- Book, C., Hoffman, M.J., Kachuck, S.B., Hillebrand, T.R., Price, S.F., Perego, M. and Bassis, J.N. 2022. Stabilizing effect of bedrock uplift on retreat of Thwaites Glacier, Antarctica, at centennial timescales. *Earth and Planetary Science Letters*, **597**, 117798, <https://doi.org/10.1016/j.epsl.2022.117798>
- Brethes, A., Guarnieri, P., Rasmussen, T.M. and Bauer, T.E. 2018. Interpretation of aeromagnetic data in the Jameson Land Basin, central East Greenland: structures and related mineralized systems. *Tectonophysics*, **724**, 116–136, <https://doi.org/10.1016/j.tecto.2018.01.008>
- Caron, L., Ivins, E.R., Larour, E., Adhikari, S., Nilsson, J. and Blewitt, G. 2018. GIA model statistics for GRACE hydrology, cryosphere, and ocean science. *Geophysical Research Letters*, **45**, 2203–2212, <https://doi.org/10.1002/2017GL076644>
- Catania, G. and Felikson, D. 2022. Topographic modulation of outlet glaciers in Greenland: a review. *Annals of Glaciology*, **63**, 171–177, <https://doi.org/10.1017/aog.2023.55>
- Celli, N.L., Lebedev, S., Schaeffer, A.J. and Gaina, C. 2021. The tilted Iceland Plume and its effect on the North Atlantic evolution and magmatism. *Earth and Planetary Science Letters*, **569**, 117048, <https://doi.org/10.1016/j.epsl.2021.117048>
- Christiansen, F.G., Whitehead, D., Bojesen-Koefoed, J.A., Boserup, J. and Christiansen, O.C. 2024. Drilling history of Greenland – exploration for minerals and petroleum, and scientific projects. *The Extractive Industries and Society*, **17**, 101417, <https://doi.org/10.1016/j.exis.2024.101417>
- Colgan, W., Wansing, A. *et al.* 2022. Greenland geothermal heat flow database and map (Version 1). *Earth System Science Data*, **14**, 2209–2238, <https://doi.org/10.5194/essd-14-2209-2022>
- Colleoni, F., Naish, T. *et al.* 2024. Instabilities and thresholds in Antarctica. *Past Global Changes Magazine*, **32**, <https://doi.org/10.22498/pages.32.1.53>
- Coulson, S., Dangendorf, S., Mitrovica, J.X., Tamisiea, M.E., Pan, L. and Sandwell, D.T. 2022. A detection of the sea level fingerprint of Greenland ice sheet melt. *Science*, **377**, 1550–1554, <https://doi.org/10.1126/science.abo0926>
- Dahl-Jensen, T., Larsen, T.B. *et al.* 2003. Depth to Moho in Greenland: receiver-function analysis suggests two Proterozoic blocks in Greenland. *Earth and Planetary Science Letters*, **205**, 379–393, [https://doi.org/10.1016/S0012-821X\(02\)01080-4](https://doi.org/10.1016/S0012-821X(02)01080-4)
- Darbyshire, F.A., Dahl-Jensen, T., Larsen, T.B., Voss, P.H. and Joyal, G. 2018. Crust and uppermost-mantle structure of Greenland and the Northwest Atlantic from Rayleigh wave group velocity tomography. *Geophysical Journal International*, **212**, 1546–1569, <https://doi.org/10.1093/gji/ggx479>
- Dawes, P.R. 2009. The bedrock geology under the Inland Ice: the next major challenge for Greenland mapping. *GEUS Bulletin*, **17**, 57–60, <https://doi.org/10.34194/geusbv.17.5014>
- Dodds, K., Gan, I. and Howkins, A. 2010. The IPY-3: the International Geophysical Year (1957–1958). In: Barr, S. and Luedicke, C. (eds) *The History of the International Polar Years (IPYs). From Pole to Pole*. Springer, Berlin, 239–257, https://doi.org/10.1007/978-3-642-12402-0_10
- Dowdeswell, J.A. 2006. The Greenland ice sheet and global sea-level rise. *Science*, **311**, 963–964, <https://doi.org/10.1126/science.1124190>
- Ferreira, A. 2024. AtlanticArray: status and outlook. ORFEUS and GEO-INQUIRE Workshop, Helsinki.
- Fleming, K. and Lambeck, K. 2004. Constraints on the Greenland ice sheet since the Last Glacial Maximum from sea-level observations and glacial rebound models. *Quaternary Science Reviews*, **23**, 1053–1077, <https://doi.org/10.1016/j.quascirev.2003.11.001>
- Fox-Kemper, B., Hewitt, H.T. *et al.* 2021. Ocean, cryosphere and sea level change. In: Masson-Delmotte, V., Zhai, P. *et al.* (eds) *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, 1211–1362, <https://doi.org/10.1017/9781009157896.011>
- Freienstein, J., Szwilius, W., Wansing, A. and Ebbing, J. 2024. Statistical appraisal of geothermal heat flow observations in the Arctic. *Solid Earth*, **15**, 513–533, <https://doi.org/10.5194/se-15-513-2024>
- Frémand, A.C., Fretwell, P. *et al.* 2022. Antarctic Bedmap data: FAIR sharing of 60 years of ice bed, surface and thickness data. *Earth System Science Data Discussions*, **2022**, 1–25.
- Fuller, J., Lebedev, S., Martinec, Z. and Celli, N.L. 2021. WINTERC-G: mapping the upper mantle thermochemical heterogeneity from coupled geophysical–petrological inversion of seismic waveforms, heat flow, surface elevation and gravity satellite data. *Geophysical Journal International*, **226**, 146–191, <https://doi.org/10.1093/gji/ggab094>
- Fyhn, M.B., Hopper, J.R., *et al.* 2021. Three-phased latest Jurassic–Eocene rifting and mild mid-Cenozoic compression offshore NE Greenland. *Tectonophysics*, **815**, 228990, <https://doi.org/10.1016/j.tecto.2021.228990>
- Golynsky, A., Chiappini, M. *et al.* 2006. ADMAP – a digital magnetic anomaly map of the Antarctic. In: Fütterer, D.K. *et al.* (eds) *Antarctic Contributions to Global Earth Science, 9th International Symposium on Antarctic Earth Sciences*. Springer, Berlin, 109–116.
- Golynsky, A.V., Ferraccioli, F. *et al.* 2018. New magnetic anomaly map of the Antarctic. *Geophysical Research Letters*, **45**, 6437–6449, <https://doi.org/10.1029/2018GL078153>
- Greve, R. 2019. Geothermal heat flux distribution for the Greenland ice sheet, derived by combining a global representation and information from deep ice cores. *Polar Data Journal*, **3**, 22–36, <https://doi.org/10.20575/00000006>
- Grinsted, A. and Dahl-Jensen, D. 2002. A Monte Carlo-tuned model of the flow in the NorthGRIP area. *Annals of Glaciology*, **35**, 527–530, <https://doi.org/10.3189/172756402781817130>
- Harig, C. and Simons, F.J. 2012. Mapping Greenland's mass loss in space and time. *Proceedings of the National Academy of Sciences of the USA*, **109**, 19934–19937, <https://doi.org/10.1073/pnas.1206785109>
- Heincke, B.H., Szwilius, W., Freienstein, J. and Ebbing, J. 2023. *A New Magnetic Compilation from Greenland Built by a Combination of Equivalent Source Modeling and Spherical Harmonic Expansion*. Greenland Portal, https://maps.greenmin.gl/geusmap/?mapname=greenland_portal&lang=en
- Heyn, B.H. and Conrad, C.P. 2022. On the relation between basal erosion of the lithosphere and surface heat flux for continental plume tracks. *Geophysical Research Letters*, **49**, e2022GL098003, <https://doi.org/10.1029/2022GL098003>
- Heyn, B.H., Shephard, G.E. and Conrad, C.P. 2024. Prolonged multi-phase magmatism due to plume–lithosphere interaction as applied to the High Arctic Large Igneous Province. *Geochemistry, Geophysics, Geosystems*, **25**, e2023GC011380, <https://doi.org/10.1029/2023GC011380>
- Höning, D., Willeit, M., Calov, R., Klemann, V., Bagge, M. and Ganopolski, A. 2023. Multistability and transient response of the Greenland ice sheet to anthropogenic CO₂ emissions. *Geophysical Research Letters*, **50**, e2022GL101827, <https://doi.org/10.1029/2022GL101827>
- Horní, J.Á., Hopper, J.R. *et al.* 2017. Regional distribution of volcanism within the North Atlantic Igneous Province. *Geological Society, London, Special Publications*, **447**, 105–125, <https://doi.org/10.1144/sp447.18>
- Horwath, M., Gutknecht, B.D. *et al.* 2022. Global sea-level budget and ocean-mass budget, with a focus on advanced data products and uncertainty characterisation. *Earth System Science Data*, **14**, 411–447, <https://doi.org/10.5194/essd-14-411-2022>
- Jakobsson, M., Mayer, L.A. *et al.* 2020. Ryder Glacier in northwest Greenland is shielded from warm Atlantic water by a bathymetric sill. *Communications Earth & Environment*, **1**, 45, <https://doi.org/10.1038/s43247-020-00043-0>
- Jansen, D., Franke, S. *et al.* 2024. Shear margins in upper half of Northeast Greenland Ice Stream were established two millennia ago. *Nature Communications*, **15**, 1193, <https://doi.org/10.1038/s41467-024-45021-8>
- Jones, G.A., Ferreira, A.M.G., Kulesa, B., Schimmel, M., Berbellini, A. and Morelli, A. 2021. Uppermost crustal structure regulates the flow of the Greenland ice sheet. *Nature Communications*, **12**, 1–12, <https://doi.org/10.1038/s41467-020-20314-w>
- Jordan, T.M., Williams, C.N. *et al.* 2018. A constraint upon the basal water distribution and thermal state of the Greenland ice sheet from radar bed echoes. *Cryosphere*, **12**, 2831–2854, <https://doi.org/10.5194/tc-12-2831-2018>
- Karlsson, N.B., Solgaard, A.M. *et al.* 2021. A first constraint on basal melt-water production of the Greenland ice sheet. *Nature Communications*, **12**, 3461, <https://doi.org/10.1038/s41467-021-23739-z>
- Kennicutt, M.C., Chown, S.L. *et al.* 2015. A roadmap for Antarctic and Southern Ocean science for the next two decades and beyond. *Antarctic Science*, **27**, 3–18, <https://doi.org/10.1017/S0954102014000674>
- Khan, S.A., Sasgen, I. *et al.* 2016. Geodetic measurements reveal similarities between post-Last Glacial Maximum and present-day mass loss from the Greenland ice sheet. *Science Advances*, **2**, <https://doi.org/10.1126/sciadv.1600931>
- Khan, S.A., Bamber, J.L. *et al.* 2022. Greenland mass trends from airborne and satellite altimetry during 2011–2020. *Journal of Geophysical Research: Earth Surface*, **127**, <https://doi.org/10.1029/2021JF006505>
- Kingslake, J., Scherer, R.P. *et al.* 2018. Extensive retreat and re-advance of the West Antarctic ice sheet during the Holocene. *Nature*, **558**, 430–434, <https://doi.org/10.1038/s41586-018-0208-x>
- Kjeldsen, K., Korsgaard, N. *et al.* 2015. Spatial and temporal distribution of mass loss from the Greenland ice sheet since AD 1900. *Nature*, **528**, 396–400, <https://doi.org/10.1038/nature16183>
- Kumar, P., Kind, R., Priestley, K. and Dahl-Jensen, T. 2007. Crustal structure of Iceland and Greenland from receiver function studies. *Journal of Geophysical Research: Solid Earth*, **112**, <https://doi.org/10.1029/2005JB003991>
- Larsen, L.M., Pedersen, A.K., Tegner, C., Duncan, R.A., Hald, N. and Larsen, J.G. 2015. Age of Tertiary volcanic rocks on the West Greenland continental margin: volcanic evolution and event correlation to other parts of the North Atlantic Igneous Province. *Geological Magazine*, **153**, 487–511, <https://doi.org/10.1017/S0016756815000515>

- Lebedev, S., Schaeffer, A.J., Fullea, J. and Pease, V. 2018. Seismic tomography of the Arctic region: inferences for the thermal structure and evolution of the lithosphere. *Geological Society, London, Special Publications*, **460**, 419–440, <https://doi.org/10.1144/SP460.10>
- Lebedev, S., Fullea, J., Xu, Y. and Bonadio, R. 2024. Seismic thermography. *Bulletin of the Seismological Society of America*, **114**, 1227–1242, <https://doi.org/10.1785/0120230245>
- Lecavalier, B.S., Milne, G.A. *et al.* 2014. A model of Greenland ice sheet deglaciation constrained by observations of relative sea level and ice extent. *Quaternary Science Reviews*, **102**, 54–84, <https://doi.org/10.1016/j.quascirev.2014.07.018>
- Lee, C.-T.A., Luffi, P., Plank, T., Dalton, H. and Leeman, W.P. 2009. Constraints on the depths and temperatures of basaltic magma generation on Earth and other terrestrial planets using new thermobarometers for mafic magmas. *Earth and Planetary Science Letters*, **279**, 20–33, <https://doi.org/10.1016/j.epsl.2008.12.020>
- Li, L., Aitken, A.R., Lindsay, M.D. and Kulesa, B. 2022. Sedimentary basins reduce stability of Antarctic ice streams through groundwater feedbacks. *Nature Geoscience*, **15**, 645–650, <https://doi.org/10.1038/s41561-022-00992-5>
- Lin, Y., Hibbert, F.D., Whitehouse, P.L., Woodroffe, S.A., Purcell, A., Shennan, I. and Bradley, S.L. 2021. A reconciled solution of Meltwater Pulse 1A sources using sea-level fingerprinting. *Nature Communications*, **12**, 2015, <https://doi.org/10.1038/s41467-021-21990-y>
- Liu, L. and Hasterok, D. 2016. High-resolution lithosphere viscosity and dynamics revealed by magnetotelluric imaging. *Science*, **353**, 1515–1519, <https://doi.org/10.1126/science.aaf6542>
- Lowe, M., Jordan, T. *et al.* 2024. Comparing geophysical inversion and petrophysical measurements for northern Victoria Land, Antarctica. *Geophysical Journal International*, **239**, 276–291, <https://doi.org/10.1093/gji/ggae272>
- Lu, C., Forte, A.M., Simmons, N.A., Grand, S.P., Kajan, M.N., Lai, H. and Garnero, E.J. 2020. The sensitivity of joint inversions of seismic and geodynamic data to mantle viscosity. *Geochemistry, Geophysics, Geosystems*, **21**, e2019GC008648, <https://doi.org/10.1029/2019GC008648>
- Macelloni, G., Leduc-Leballeur, M., Montomoli, F., Brogioni, M., Ritz, C. and Picard, G. 2019. On the retrieval of internal temperature of Antarctica Ice Sheet by using SMOS observations. *Remote Sensing of Environment*, **233**, 111405, <https://doi.org/10.1016/j.rse.2019.111405>
- MacGregor, J.A., Fahnestock, M.A. *et al.* 2016. A synthesis of the basal thermal state of the Greenland ice sheet. *Journal of Geophysical Research: Earth Surface*, **121**, 1328–1350, <https://doi.org/10.1002/2015JF003803>
- MacGregor, J.A., Colgan, W.T. *et al.* 2024. Geologic provinces beneath the Greenland ice sheet constrained by geophysical data synthesis. *Geophysical Research Letters*, **51**, e2023GL107357, <https://doi.org/10.1029/2023GL107357>
- Manassero, M.C., Özyaydin, S. *et al.* 2024. Lithospheric structure and melting processes in southeast Australia: new constraints from joint probabilistic inversions of 3D magnetotelluric and seismic data. *Journal of Geophysical Research: Solid Earth*, **129**, e2023JB028257, <https://doi.org/10.1029/2023JB028257>
- Mankoff, K.D., Fettweis, X. *et al.* 2021. Greenland ice sheet mass balance from 1840 through next week. *Earth System Science Data*, **13**, 5001–5025, <https://doi.org/10.5194/essd-13-5001-2021>
- Martos, Y.M. *et al.* 2018. Geothermal heat flux reveals the Iceland hotspot track underneath Greenland. *Geophysical Research Letters*, **45**, 8214–8222, <https://doi.org/10.1029/2018GL078289>
- McCormack, F.S., Roberts, J.L., Dow, C.F., Stål, T., Halpin, J.A., Reading, A.M. and Siegert, M.J. 2022. Fine-scale geothermal heat flow in Antarctica can increase simulated subglacial melt estimates. *Geophysical Research Letters*, **49**, e2022GL098539, <https://doi.org/10.1029/2022GL098539>
- Meyer, B., Chulliat, A. and Saltus, R. 2017. Derivation and error analysis of the earth magnetic anomaly grid at 2 arc min resolution version 3 (EMAG2v3). *Geochemistry, Geophysics, Geosystems*, **18**, 4522–4537, <https://doi.org/10.1002/2017GC007280>
- Milne, G.A., Latychev, K., Schaeffer, A., Crowley, J.W., Lecavalier, B.S. and Audette, A. 2018. The influence of lateral Earth structure on glacial isostatic adjustment in Greenland. *Geophysical Journal International*, **214**, 1252–1266, <https://doi.org/10.1093/gji/ggy189>
- Mitrovica, J.X., Gomez, N., Morrow, E., Hay, C., Latychev, K. and Tamisiea, M.E. 2011. On the robustness of predictions of sea level fingerprints. *Geophysical Journal International*, **187**, 729–742, <https://doi.org/10.1111/j.1365-246X.2011.05090.x>
- Moorkamp, M. 2022. Deciphering the state of the lower crust and upper mantle with multi-physics inversion. *Geophysical Research Letters*, **49**, e2021GL096336, <https://doi.org/10.1029/2021GL096336>
- Mordret, A. 2018. Uncovering the Iceland hot spot track beneath Greenland. *Journal of Geophysical Research: Solid Earth*, **123**, 4922–4941, <https://doi.org/10.1029/2017JB015104>
- Morlighem, M., Williams, C.N. *et al.* 2017. BedMachine v3: complete bed topography and ocean bathymetry mapping of Greenland from multibeam echo sounding combined with mass conservation. *Geophysical research letters*, **44**, 11051, <https://doi.org/10.1002/2017GL074954>
- Morlighem, M., Williams, C. *et al.* 2022. *IceBridge BedMachine Greenland (IDBMG4, Version 5)* [Dataset]. NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, CO, <https://doi.org/10.5067/GMEVBWFLWA7X> [last accessed 4 April 2025].
- Nielsen, T.K., Larsen, H.C. and Hopper, J.R. 2002. Contrasting rifted margin styles south of Greenland: implications for mantle plume dynamics. *Earth and Planetary Science Letters*, **200**, 271–286, [https://doi.org/10.1016/S0012-821X\(02\)00616-7](https://doi.org/10.1016/S0012-821X(02)00616-7)
- Pan, L., Mitrovica, J.X., Milne, G.A., Hoggard, M.J. and Woodroffe, S.A. 2024. Timescales of glacial isostatic adjustment in Greenland: is transient rheology required? *Geophysical Journal International*, **237**, 989–995, <https://doi.org/10.1093/gji/ggae095>
- Pasyanos, M.E., Masters, T.G., Laske, G. and Ma, Z. 2014. LITHO1.0: an updated crust and lithospheric model of the Earth. *Journal of Geophysical Research: Solid Earth*, **119**, 2153–2173, <https://doi.org/10.1002/2013JB010626>
- Paxman, G.J., Lau, H.C., Austermann, J., Holtzman, B.K. and Havlin, C. 2023. Inference of the timescale-dependent apparent viscosity structure in the upper mantle beneath Greenland. *AGU Advances*, **4**, e2022AV000751, <https://doi.org/10.1029/2022AV000751>
- Peacock, J.R. and Selway, K. 2016. Magnetotelluric investigation of the Vestfold Hills and Rauer Group, East Antarctica. *Journal of Geophysical Research: Solid Earth*, **121**, 2258–2273, <https://doi.org/10.1002/2015JB012677>
- Porter, R. and Reid, M. 2021. Mapping the thermal lithosphere and melting across the continental US. *Geophysical Research Letters*, **48**, e2020GL092197, <https://doi.org/10.1029/2020GL092197>
- Priestley, K. and McKenzie, D. 2013. The relationship between shear wave velocity, temperature, attenuation and viscosity in the shallow part of the mantle. *Earth and Planetary Science Letters*, **381**, 78–91, <https://doi.org/10.1016/j.epsl.2013.08.022>
- Ramirez, F.D.C., Selway, K., Conrad, C.P. and Lithgow-Bertelloni, C. 2022. Constraining upper mantle viscosity using temperature and water content inferred from seismic and magnetotelluric data. *Journal of Geophysical Research: Solid Earth*, **127**, e2021JB023824, <https://doi.org/10.1029/2021JB023824>
- Ramirez, F.D.C., Selway, K., Conrad, C.P., Maupin, V. and Smirnov, M. 2024. Lateral and radial viscosity structure beneath Fennoscandia inferred from seismic and magnetotelluric observations. *Physics of the Earth and Planetary Interiors*, **351**, 107178, <https://doi.org/10.1016/j.pepi.2024.107178>
- Reading, A.M., Stål, T. *et al.* 2022. Antarctic geothermal heat flow and its implications for tectonics and ice sheets. *Nature Reviews Earth & Environment*, **3**, 814–831, <https://doi.org/10.1038/s43017-022-00348-y>
- Rignot, E. and Kanagaratnam, P. 2006. Changes in the velocity structure of the Greenland ice sheet. *Science*, **311**, 986–990, <https://doi.org/10.1126/science.1121381>
- Rogozhina, I., Hagedoorn, J.M., Martinec, Z., Fleming, K., Soucek, O., Greve, R. and Thomas, M. 2012. Effects of uncertainties in the geothermal heat flux distribution on the Greenland ice sheet: an assessment of existing heat flow models. *Journal of Geophysical Research: Earth Surface*, **117**, <https://doi.org/10.1029/2011JF002098>
- Salajegheh, F., Afonso, J.C., Minakov, A., Ajournalou, P., Gaina, C. and Ortega-Gelabert, O. 2025. The lithosphere–asthenosphere system beneath the North Atlantic and surroundings: Results from multi-observable probabilistic inversions. *Geochemistry, Geophysics, Geosystems*, **26**, e2024GC011724, <https://doi.org/10.1029/2024GC011724>
- Schutt, D.L., Lowry, A.R. and Buehler, J.S. 2018. Moho temperature and mobility of lower crust in the western United States. *Geology*, **46**, 219–222, <https://doi.org/10.1130/g39507.1>
- Simonsen, S.B., Barletta, V.R., Colgan, W.T. and Sørensen, L.S. 2021. Greenland ice sheet mass balance (1992–2020) from calibrated radar altimetry. *Geophysical Research Letters*, **48**, e2020GL091216, <https://doi.org/10.1029/2020GL091216>
- Simpson, M.J.R., Wake, L., Milne, G.A. and Huybrechts, P. 2011. The influence of decadal- to millennial-scale ice mass changes on present-day vertical land motion in Greenland: implications for the interpretation of GPS observations. *Journal of Geophysical Research: Solid Earth*, **116**, B02406, <https://doi.org/10.1029/2010JB007776>
- Smith-Johnsen, S., de Fleurian, B., Schlegel, N., Seroussi, H. and Nisancioglu, K. 2020a. Exceptionally high heat flux needed to sustain the Northeast Greenland Ice Stream. *Cryosphere*, **14**, 841–854, <https://doi.org/10.5194/tc-14-841-2020>
- Smith-Johnsen, S., Schlegel, N.-J., de Fleurian, B. and Nisancioglu, K.H. 2020b. Sensitivity of the Northeast Greenland Ice Stream to geothermal heat. *Journal of Geophysical Research: Earth Surface*, **125**, e2019JF005252, <https://doi.org/10.1029/2019JF005252>
- Spada, G. 2017. Glacial isostatic adjustment and contemporary sea level rise: an overview. In: Cazenave, A., Champollion, N., Paul, F. and Benveniste, J. (eds) *Integrative Study of the Mean Sea Level and Its Components*. Springer, Cham, Space Sciences Series of ISSI, **58**, 155–187.
- Stål, T., Reading, A.M., Halpin, J.A. and Whittaker, J.M. 2019. A multivariate approach for mapping lithospheric domain boundaries in East Antarctica. *Geophysical Research Letters*, **46**, 10404–10416, <https://doi.org/10.1029/2019GL083453>
- Stål, T., Halpin, J.A., Goodge, J.W. and Reading, A.M. 2024. Geology matters for Antarctic geothermal heat. *Geophysical Research Letters*, **51**, e2024GL110098, <https://doi.org/10.1029/2024GL110098>

- Steffen, R., Audet, P. and Lund, B. 2018. Weakened lithosphere beneath Greenland inferred from effective elastic thickness: a hot spot effect? *Geophysical Research Letters*, **45**, 4733–4742, <https://doi.org/10.1029/2017GL076885>
- Steinberger, B., Bredow, E., Lebedev, S., Schaeffer, A. and Torsvik, T.H. 2019. Widespread volcanism in the Greenland–North Atlantic region explained by the Iceland plume. *Nature Geoscience*, **12**, 61–68, <https://doi.org/10.1038/s41561-018-0251-0>
- Stoker, M.S., Stewart, M.A., *et al.* 2017. An overview of the Upper Palaeozoic–Mesozoic stratigraphy of the NE Atlantic region. *Geological Society, London, Special Publications*, **447**, 11–68, <https://doi.org/10.1144/SP447.2>
- Tarasov, L. and Peltier, W.R. 2002. Greenland glacial history and local geodynamic consequences. *Geophysical Journal International*, **150**, 198–229, <https://doi.org/10.1046/j.1365-246X.2002.01702.x>
- Toyokuni, G., Matsuno, T. and Zhao, D. 2020. P wave tomography beneath Greenland and surrounding regions: 1. Crust and upper mantle. *Journal of Geophysical Research: Solid Earth*, **125**, e2020JB019837, <https://doi.org/10.1029/2020JB019837>
- Van Den Berg, J., van de Wal, R. and Oerlemans, H. 2008. A mass balance model for the Eurasian Ice Sheet for the last 120,000 years. *Global and Planetary Change*, **61**, 194–208, <https://doi.org/10.1016/j.gloplacha.2007.08.015>
- Velicogna, I. 2009. Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE. *Geophysical Research Letters*, **36**, <https://doi.org/10.1029/2009GL040222>
- Velicogna, I., Sutterley, T.C. and Van Den Broeke, M.R. 2014. Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data. *Geophysical Research Letters*, **41**, 8130–8137, <https://doi.org/10.1002/2014GL061052>
- Wake, L.M., Lecavalier, B.S. and Bevis, M. 2016. Glacial Isostatic Adjustment (GIA) in Greenland: a review. *Current Climate Change Reports*, **2**, 101–111, <https://doi.org/10.1007/s40641-016-0040-z>
- Wannamaker, P. *et al.* 2017. Uplift of the central Transantarctic Mountains. *Nature Communications*, **8**, 1588, <https://doi.org/10.1038/s41467-017-01577-2>
- Wansing, A., Ebbing, J. and Moorkamp, M. 2024. The lithospheric structure of Greenland from a stepwise forward and inverse modelling approach. *Geophysical Journal International*, **238**, 719–741, <https://doi.org/10.1093/gji/ggae183>
- Weerdesteijn, M.F.M. and Conrad, C.P. 2024. Recent ice melt above a mantle plume track is accelerating the uplift of southeast Greenland. *Communications Earth and Environment*, **5**, 791, <https://doi.org/10.1038/s43247-024-01968-6>
- Weerdesteijn, M.F.M., Conrad, C.P. and Naliboff, J.B. 2022. Solid Earth uplift due to contemporary ice melt above low-viscosity regions of the upper mantle. *Geophysical Research Letters*, **49**, e2022GL099731, <https://doi.org/10.1029/2022GL099731>
- Weerdesteijn, M.F.M., Naliboff, J.B., Conrad, C.P., Reusen, J.M., Steffen, R., Heister, T. and Zhang, J. 2023. Modeling viscoelastic solid Earth deformation due to ice age and contemporary glacial mass changes in ASPECT. *Geochemistry, Geophysics, Geosystems*, **24**, e2022GC010813, <https://doi.org/10.1029/2022GC010813>
- Whitehouse, P.L., Gomez, N., King, M.A. and Wiens, D.A. 2019. Solid Earth change and the evolution of the Antarctic ice sheet. *Nature Communications*, **10**, 503, <https://doi.org/10.1038/s41467-018-08068-y>
- Willcocks, S., Hasterok, D. and Jennings, S. 2021. Thermal refraction: implications for subglacial heat flux. *Journal of Glaciology*, **67**, 875–884, <https://doi.org/10.1017/jog.2021.38>
- Wood, M., Rignot, E. *et al.* 2021. Ocean forcing drives glacier retreat in Greenland. *Science Advances*, **7**, eaba7282, <https://doi.org/10.1126/sciadv.aba7282>
- Zeitz, M., Haacker, J.M., Donges, J.F., Albrecht, T. and Winkelmann, R. 2022. Dynamic regimes of the Greenland Ice Sheet emerging from interacting melt-elevation and glacial isostatic adjustment feedbacks. *Earth System Dynamics*, **13**, 1077–1096, <https://doi.org/10.5194/esd-13-1077-2022>
- Zhang, T., Colgan, W., Wansing, A., Lokkegaard, D., Leguy, G., Lipscomb, W.H. and Cunde, X. 2024. Evaluating different geothermal heat-flow maps as basal boundary conditions during spin-up of the Greenland ice sheet. *Cryosphere*, **18**, 387–402, <https://doi.org/10.5194/tc-18-387-2024>