Atacama Cosmology Telescope DR6 and DESI: Structure growth measurements from the cross-correlation of DESI legacy imaging galaxies and CMB lensing from ACT DR6 and *Planck* PR4

Frank J. Qu,^{1,2,3,4,*} Qianjun Hang,⁵ Gerrit Farren,⁶ Boris Bolliet,^{3,4} Jessica Nicole Aguilar,⁶ Steven Ahlen,⁷ Shadab Alam,⁸ David Brooks,⁵ Yan-Chuan Cai,⁹ Erminia Calabrese,¹⁰ Todd Claybaugh,⁶ Axel de la Macorra,¹¹ Mark J. Devlin,¹² Peter Doel,⁵ Carmen Embil-Villagra,³ Simone Ferraro,^{6,13} Andreu Font-Ribera,^{5,14} Jaime E. Forero-Romero,^{15,16} Enrique Gaztañaga,¹⁷ Vera Gluscevic,¹⁸ Satya Gontcho A. Gontcho,⁶ Gaston Gutierrez,¹⁹ Cullan Howlett,²⁰ Robert Kehoe,²¹ Joshua Kim,²² Anthony Kremin,⁶ Andrew Lambert,⁶ Martin Landriau,⁶ Laurent Le Guillou,²³ Michael Levi,⁶ Thibaut Louis,²⁴ Aaron Meisner,²⁵ Ramon Miquel,^{26,14} John Moustakas,²⁷ Jeffrey A. Newman,²⁸ Gustavo Niz,^{29,30} John A. Peacock,⁹ Will Percival,^{31,32,33} Claire Poppett,^{6,34,13} Francisco Prada,³⁵ Ignasi Pérez-Ràfols,³⁶ Graziano Rossi,³⁷ Eusebio Sanchez,³⁸ David Schlegel,⁶ Neelima Sehgal,³⁹ Shabbir Shaikh,⁴⁰ Blake Sherwin,^{3,4} Cristóbal Sifón,⁴¹ Michael Schubnell,^{42,43} David Sprayberry,²⁵ Gregory Tarlé,⁴³ Benjamin Alan Weaver,²⁵ Edward J. Wollack,⁴⁴ and Hu Zou⁴⁵ ¹Kavli Institute for Particle Astrophysics and Cosmology, 382 Via Pueblo Mall Stanford, California 94305-4060, USA ²SLAC National Accelerator Laboratory, 2575 Sand Hill Road, Menlo Park, California 94025, USA ³DAMTP, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 OWA, United Kingdom ⁴Kavli Institute for Cosmology Cambridge, Madingley Road, Cambridge CB3 0HA, United Kingdom ²Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, United Kingdom ⁶Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, California 94720, USA ⁷Physics Department, Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215, USA ⁸Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India ⁹Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, United Kingdom ¹⁰School of Physics and Astronomy, Cardiff University, The Parade, Cardiff, Wales CF24 3AA, United Kingdom ¹¹Instituto de Física, Universidad Nacional Autónoma de México, Cd. de México C.P. 04510, Mexico ¹²Department of Physics and Astronomy, University of Pennsylvania, 209 South 33rd Street, Philadelphia, Pennsylvania 19104, USA ¹³University of California, Berkeley, 110 Sproul Hall No. 5800 Berkeley, California 94720, USA ¹⁴Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra, Barcelona, Spain ¹⁵Departamento de Física, Universidad de los Andes, Carrera 1 No. 18A-10, Edificio Ip, CP 111711, Bogotá, Colombia ¹⁶Observatorio Astronómico, Universidad de los Andes, Carrera 1 No. 18A-10, Edificio H, CP 111711, Bogotá, Colombia ¹⁷Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain ¹⁸Department of Physics and Astronomy, University of Southern California, Los Angeles, California 90007, USA ¹⁹Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, Illnois 60510, USA ²⁰School of Mathematics and Physics, University of Queensland, 4072, Australia ²¹Department of Physics, Southern Methodist University, 3215 Daniel Avenue, Dallas, Texas 75275, USA ²²Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA ²³Sorbonne Université, CNRS/IN2P3, Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), FR-75005 Paris, France ²⁴Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France ²⁵NSF NOIRLab, 950 North Cherry Avenue, Tucson, Arizona 85719, USA ²⁶Institució Catalana de Recerca i Estudis Avançats, Passeig de Lluís Companys, 23, 08010 Barcelona, Spain ²⁷Department of Physics and Astronomy, Siena College, 515 Loudon Road, Loudonville, New York 12211, USA

²⁸Department of Physics and Astronomy and Pittsburgh Particle Physics, Astrophysics, and Cosmology Center (PITT PACC), University of Pittsburgh, 3941 O'Hara Street, Pittsburgh, Pennsylvania 15260, USA
 ²⁹Departamento de Física, Universidad de Guanajuato—DCI, C.P. 37150, Leon, Guanajuato, Mexico
 ³⁰Instituto Avanzado de Cosmología A. C., San Marcos 11—Atenas 202, Magdalena Contreras, 10720, Ciudad de México, Mexico
 ³¹Department of Physics and Astronomy, University of Waterloo,

200 University Avenue West, Waterloo, Ontario N2L 3G1, Canada

³²Perimeter Institute for Theoretical Physics,

31 Caroline Street North, Waterloo, Ontario N2L 2Y5, Canada

³³Waterloo Centre for Astrophysics, University of Waterloo,

200 University Avenue West, Waterloo, Ontario N2L 3G1, Canada

³⁴Space Sciences Laboratory, University of California,

Berkeley, 7 Gauss Way, Berkeley, California 94720, USA

³⁵Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía, s/n, E-18008 Granada, Spain

³⁶Departament de Física, EEBE, Universitat Politècnica de Catalunya,

c/Eduard Maristany 10, 08930 Barcelona, Spain

³⁷Department of Physics and Astronomy, Sejong University, Seoul 143-747, Korea

³⁸CIEMAT, Avenida Complutense 40, E-28040 Madrid, Spain

³⁹Physics and Astronomy Department, Stony Brook University,

Stony Brook, New York 11794, USA

⁴⁰School of Earth and Space Exploration, Arizona State University,

Tempe, Arizona 85287, USA

⁴¹Instituto de Física, Pontificia Universidad Católica de Valparaíso,

Casilla 4059, Valparaíso, Chile

⁴²Department of Physics, University of Michigan, Ann Arbor, Michigan 48109, USA

⁴³University of Michigan, Ann Arbor, Michigan 48109, USA

⁴⁴NASA Goddard Spaceflight Center, 8800 Greenbelt Road, Greenbelt, Maryland 20771, USA

⁴⁵National Astronomical Observatories, Chinese Academy of Sciences, A20 Datun Road, Chaoyang

District, Beijing, 100012, People's Republic of China

(Received 9 November 2024; accepted 10 April 2025; published 2 May 2025)

We measure the growth of cosmic density fluctuations on large scales and across the redshift range 0.3 < z < 0.8 through galaxy clustering and the cross-correlation of the ACT data release 6 cosmic microwave background (CMB) lensing map and galaxies from the Dark Energy Spectroscopic Instrument Legacy Survey, using three galaxy samples spanning the redshifts of $0.3 \leq z \leq 0.45$, $0.45 \leq z \leq 0.6$, $0.6 \leq z \leq 0.8$. We adopt a scale cut where nonlinear effects are negligible, so that the cosmological constraints are derived from the linear regime. We determine the amplitude of matter fluctuations over all three redshift bins using Atacama Cosmology Telescope (ACT) data alone to be $S_8 \equiv \sigma_8 (\Omega_m/0.3)^{0.5} = 0.772 \pm 0.040$ in a joint analysis combining the three redshift bins and ACT lensing alone. Using a combination of ACT and *Planck* data we obtain $S_8 = 0.765 \pm 0.032$. The lowest redshift bin used is the least constraining and exhibits a ~ 2σ tension with the other redshift bins; thus we also report constraints excluding the first redshift bin, giving $S_8 = 0.785 \pm 0.033$ for the combination of ACT and *Planck*. This result is in excellent agreement at the 0.3σ level with measurements from galaxy lensing, but is 1.8σ lower than predictions based on *Planck* primary CMB data. Understanding whether this hint of discrepancy in the growth of structure at low redshifts arises from a fluctuation, from systematics in data, or from new physics is a high priority for forthcoming CMB lensing and galaxy cross-correlation analyses.

DOI: 10.1103/PhysRevD.111.103503

Contact author: jq247@cantab.ac.uk

I. INTRODUCTION

The standard cosmological model, known as Λ cold dark matter (Λ CDM), has provided a remarkably successful framework for understanding the large-scale structure and evolution of the Universe. One of its successes is its predictive power over a wide range of redshifts. The way in which structure grows and clusters from initial primordial

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

seeds depends sensitively on the parameters of the model, on the underlying theory of gravity (e.g., [1]), and on the details of the expansion history (e.g., [2]). Thus, measuring the large-scale structure growth in the Universe and comparing it with an extrapolation from cosmic microwave background (CMB) data is a powerful test for cosmological models.

Multiple observables of cosmological large-scale structure provide a measure of density fluctuations at low redshifts. Some of these observables include gravitational lensing measurements, Sunyaev-Zeldovich effects [3–5], redshift-space distortions [6–8], galaxy cluster counts [9–13], and peculiar velocities [14–16]. These analyses cover a broad range of scales and redshifts and are subject to different systematic effects. For ease of comparison, it is conventional to report the amplitude of matter fluctuations as σ_8 or $S_8 \sim \sigma_8 \sqrt{\Omega_m/0.3}$, where σ_8 is defined as the rootmean-square amplitude of linear fluctuations at present, smoothed on scales of $8h^{-1}$ Mpc.

Several recent measurements have reported values of S_8 approximately $2-3\sigma$ lower than predicted from a Λ CDM fit to the primary CMB [e.g., $S_8 = 0.830 \pm 0.013$ from Planck public release 4 (PR4) [17]], raising questions about a potential breakdown of the ACDM model at low redshift. Examples of such results yielding low values of S_8 include galaxy weak lensing (in particular, the combination of cosmic shear and galaxy clustering): the Dark Energy Survey (DES), the Kilo-Degree Survey (KiDS), and the Hyper Suprime-Cam (HSC) obtain $S_8 = 0.782 \pm 0.019$, $S_8 = 0.765^{+0.017}_{-0.016}$, and $S_8 = 0.775^{+0.043}_{-0.038}$, respectively [18-20]. A joint reanalysis of the DES and KiDS cosmic shear data [21] results in a slightly higher $S_8 = 0.790^{+0.018}_{-0.014}$. Cross-correlations between Planck CMB lensing and Dark Energy Spectroscopic Instrument (DESI) luminous red galaxy (LRG) targets [22] or galaxies from the Baryonic Oscillation Spectroscopic Survey (BOSS) also indicate S_8 values that are $2-3\sigma$ low [23,24]. Results from the crosscorrelation between various datasets from DES and CMB lensing from the South Pole Telescope (SPT) and *Planck* [25,26] have also found discrepancies at the 2.2 and 3σ level. A similar result is found with some cross-correlations with the Atacama Cosmology Telescope (ACT); albeit with large uncertainties, CMB lensing from ACT data release 4 (DR4) and Planck correlations with galaxy shear from KiDS-1000 [27], found $S_8 = 0.64 \pm 0.08$. A similar cross-correlation but using galaxy clustering from DES Year 3 (DES-Y3) [28] instead results in $S_8 = 0.75^{+0.04}_{-0.05}$. Shaikh *et al.* [29] performed cross-correlation of ACT DR4 CMB lensing and DES galaxy shear, resulting in $S_8 = 0.782 \pm 0.059$. This collection of S_8 measurements that consistently return values lower than those measured by the primary CMB is known as the S_8 tension; while none of them are inconsistent with Planck at high statistical significance, it is striking that they all tend to measure a low S_8 .

It is, however, worth pointing out that not all low-redshift measurements find low values of S_8 . Notably, the CMB

lensing autospectrum is in excellent agreement with the primary CMB predictions, CMB lensing is arguably one of the cleanest probes of low-redshift matter fluctuation, given that it probes directly the gravitational potential on mostly linear scales across a large range of redshifts ($z \approx 0.5-5$), and is based on robust and well-understood statistical properties of the CMB. Recent measurements from the Atacama Cosmology Telescope [30,31] and baryonic acoustic oscillation (BAO) data results in $S_8 = 0.840 \pm 0.028$ from ACT alone and $S_8 = 0.831 \pm 0.023$ in combination with the latest *Planck* PR4 lensing [32]. Similar consistent measurements are obtained with SPT-3G [33]: $S_8 = 0.836 \pm 0.039$.

Furthermore, some improved analyses including more data of previous datasets suggesting low S_8 values have found less discrepant results compared to earlier measurements. Examples include the cross-correlation between CMB lensing from ACT DR6 and *Planck* with unWISE (wide-field infrared survey explorer) galaxies [34], which resulted in $S_8 = 0.810 \pm 0.015$, the cross-correlation between *Planck* PR3 lensing and DESI LRGs with $S_8 = 0.763 \pm 0.023$ [35,36], and the cross-correlation between DESI bright galaxy survey (BGS) and LRG, and the DES year 3 galaxy shear in [37], with $S_8 = 0.850^{+0.042}_{-0.050}$. We refer the interested reader to Fig. 19 to a summary of the relevant S_8 measurements.

The present paper is especially concerned with a further CMB lensing cross-correlation measurement that reported a low normalization. Hang et al. [38] measured the crosscorrelation between *Planck* and the DESI legacy galaxy catalog, divided into four tomographic bins; from this, they inferred a constraint on $S_8^{\times} \equiv \sigma_8 (\Omega_m/0.3)^{0.79} =$ 0.758 ± 0.023 , which is 2.8σ lower than the *Planck* CMB prediction. Here, we reexamine the constraints obtained by [38], motivated by the availability of lower noise CMB lensing maps from ACT DR6 and Planck PR4. In addition to these new datasets, we provide improved treatment of mode couplings across bandpowers due to the presence of a mask, a more accurate estimation of the covariance matrix based on simulations, a large suite of systematic tests, and use more conservative scale cut choices. In what follows, we will refer to our combined analysis of the galaxy autospectrum and the galaxy-CMB lensing cross-spectrum as a " 2×2 pt analysis" (i.e., galaxy clustering and lensing cross-correlation).

The paper is structured as follows: In Sec. II we present the datasets used in this analysis. In Sec. III we discuss our measurement pipeline and scale cuts. The verification of the pipeline and the covariance matrices are described in Sec. IV. We present a series of null and consistency tests in Sec. V. In Sec. VI we discuss the modeling of the angular power spectra. Our results are given in Sec. VII, and we place these measurements in context in Sec. VIII.

II. DATA USED

We use the photometric DESI legacy imaging survey galaxy catalog described in [38] and the corresponding

galaxy density contrast maps in four tomographic bins. We conservatively restrict our analysis to linear scales (see Sec. III A) and therefore we exclude the lowest tomographic bin (bin 0) where all bandpowers receive significant nonlinear contributions. In Sec. II A we briefly describe this dataset and refer the interested readers to [38] for more details. We then describe in Sec. II B the lensing convergence map obtained using the ACT DR6 data and in Sec. II C the lensing convergence from *Planck* PR4.

A. DESI legacy imaging surveys DR8

The DESI legacy imaging surveys [39] are the union of three public imaging galaxy surveys: the Dark Energy Camera Legacy Survey (DECaLS) [40,41], the Mayall z-band Legacy Survey (MzLS) [42], and the Beijing-Arizona Sky Survey (BASS) [43], altogether covering a total area of $\sim 14,000 \text{ deg}^2$. The photometry was measured in grz bands, with the addition of the WISE fluxes (W_1 , W_2 , W_3) from the 4-yr NEOWISE-Reactivation imaging, force photometered in the unWISE maps at the location of the legacy survey sources [44,45]. This paper uses the public DR8¹ data, the first release to include both images and catalogs from all three of the legacy surveys. Note that this dataset has been superseded by the subsequent data releases, DR9 and DR10.² The major updates include more recent data releases from the DECaLS, BASS, MzLS, and the NEOWISE-Reactivation, and additionally, DR9 improved in reduction techniques and procedures, and DR10 included additional DECam data from NOIRLab in griz bands. Notably, DR9 was used to select DESI targets. Because the purpose of this paper is to update the measurements from [38] with ACT DR6 CMB lensing, for consistency, we adopt the galaxy data from [38] instead of using a more recent data release.

The galaxy sample was selected with g < 24, r < 22, and $W_1 < 19.5$. Pixels contaminated by bright stars, globular clusters, and incompleteness in optical bands are masked by bitmasks³ supplied by the legacy survey pipeline, with bits = (0, 1, 5, 6, 7, 11, 12, 13). From this, the survey completeness map was computed which indicates the geometric completeness of the observation in the range [0, 1], with 0 indicating no observation and 1 indicating full coverage of the pixel. Regions with completeness < 0.86 were masked. This particular choice of completeness cut is based on the binned relation between the completeness and mean galaxy number density, such that the variation of the average galaxy overdensity, δ_g in bins of completeness is < 0.1.

Photometric redshifts were calibrated using the following spectroscopic datasets: GAMA (DR2 [46]), BOSS LOWZ and CMASS samples (DR12 [47]), eBOSS LRG and emission line galaxy (ELG) samples (DR16 [48]), VIPERS (DR2 [49]), DEEP2 [50], and COSMOS ([51] with a magnitude cut of rMAGAPER2 ≤ 23). Notice that most of the calibration samples overlap with the DECaLS footprint, while the BASS + MzLS footprint is only partially covered by the BOSS sample. The spectroscopic calibration samples were binned in g - r, r - z, and $z - W_1$ color space with a bin width of about 0.03. Additionally, the DESY1A1 redMaGiC sample [52] was used to fill in the color cells that do not have galaxies, thanks to their highly accurate photometric redshifts. The color cells populated with less than 5 objects are excluded. For the remaining cells, we compute the average spectroscopic redshifts and their standard deviation. We do not apply further selections to the cells based on, e.g., the standard deviation of the cells. This could potentially be applied to further increase the photometric redshift accuracy. Finally, the legacy imaging survey galaxies are binned into the same color grid. They are either assigned with the mean redshift of the cell with spectroscopic samples or excluded from the analysis. This process acts effectively as an additional selection, and 78.6% of the above-selected galaxy sample remains. Finally, galaxies were selected with photometric redshifts consistent with those estimated in [53], with a criterion of $|\Delta z| < 0.05$, further removing 23.4% of the galaxy sample. The total number of galaxies after these cuts is 2.0120352×10^7 from DECaLS and 7.117218×10^6 from BASS + MzLS.

The selected galaxies are split into four tomographic bins with redshift bin edges: [0, 0.3, 0.45, 0.6, 0.8]. Maps of the galaxy density contrast, $\delta_g = n/\bar{n} - 1$, were constructed in HEALPix [54] format with pixel resolution Nside = 1024. Here, *n* is the number of galaxies in each pixel, and \bar{n} is the mean number of galaxies per pixel within the mask. The redshift distribution n(z) of each tomographic bin is then further calibrated using a "self-calibration" scheme, where the galaxy cross-power spectra between different tomographic bins were fitted simultaneously.

Hang *et al.* [38] characterized photometric redshift scatter by a modified Lorentzian function,

$$L(z) = \frac{L_0}{[1 + ((z - z_0)/\sigma)^2/2a]^a},$$
 (1)

where z_0 , σ , and a are free parameters controlling the mean, width, and spread of the tail of the n(z) for each tomographic bin, and L_0 is the normalization such that $\int L(z)dz = 1$. They also impose $\sum_i z_0^i = 0$ for each bin *i*, such that the mean redshift of the full sample is not changed. This gives a total of seven nuisance parameters. The calibrated n(z) is computed by the convolution of the raw $n_{\text{raw}}(z)$ with L(z),

¹http://legacysurvey.org/dr8/

²https://www.legacysurvey.org/

³See http://legacysurvey.org/dr8/bitmasks. The different "bits" are flags at pixel level indicating different reasons for masking, e.g., bright stars or saturation at a certain band.



FIG. 1. Overlap of the DESI imaging galaxies with the ACT DR6 lensing map (area within the blue contours) and the *Planck* PR4 lensing footprint (black contours). The overlaid gray scale background is a Galactic dust map from *Planck* [61].

$$n_{\rm cal}(z) = \int_{-\infty}^{\infty} n_{\rm raw}(z') L(z-z') dz'.$$
 (2)

In practice, the convolution can result in nonzero values of $n_{\rm cal}(z)$ at z < 0, but this is in general negligible and thus set to zero $[n_{cal}(z)$ is always normalized to $\int n_{cal}(z)dz = 1$]. In [38], they jointly constrained these nuisance parameters and galaxy biases using the galaxy auto- and cross-spectra. They found that marginalizing over these free parameters gives the same posterior on the lensing amplitude, at fixed cosmology, as using the maximum a posteriori probability estimate. Hence, in this work, we adopt the best-fit model for the n(z), as found in [38]. As shown in [38], the crosscorrelation coefficients, $r_{\ell}^{ij} = C_{\ell}^{ij} (C_{\ell}^{ii} C_{\ell}^{jj})^{-1/2}$, from neighboring tomographic bins *i* and *j* are quite flat up to $\ell_{\rm max} = 500$. This means that the redshift calibration, which takes most information from these cross terms, has little dependence on the scales and is independent of linear bias. Magnification was not included in the calibration, but there is little correlation between the highest and the lowest tomographic bins, suggesting that the effect is small. We also found that marginalizing over the nuisance parameters made little change to the final result compared to simply adopting their best-fit values. Here, we additionally check that the assumption of the fiducial *Planck* cosmology does not affect our redshift calibration process. We verify the n(z) calibration dependence on cosmology via variations in r_{ℓ}^{ij} . We find that, by adopting the Planck 2018 and DES-Y3 best-fit cosmologies, the change in $r_{\ell}^{ij} < 2\%$. We explore

cases where we apply a shift in the mean redshift of all bins and the impact on cosmology in Appendix D.

Figure 2 shows the calibrated redshift distributions of the galaxy samples used in this analysis. In this work, we omit the lowest redshift bin (bin 0) in [38] due to the choice of scale cuts (see Sec. III A) and refer to bins 1–3 as $z_1 - z_3$ hereafter. We also compute and show in Table I the effective redshift for the angular autospectrum C_{ℓ}^{gg} and cross-spectrum C_{ℓ}^{kg}

$$z_{\rm eff}^{xy} = \frac{\int d\chi z(\chi) W^x(z) W^y(z)/\chi^2}{\int d\chi W^x(z) W^y(z)/\chi^2},\tag{3}$$



FIG. 2. The calibrated photometric redshift distribution of each tomographic bin of the DESI legacy imaging galaxy sample from [38].

TABLE I. Summary of the DESI legacy sample properties, with the DESI × ACT mask. The effective redshift z_{eff} is calculated with Eq. (3). The shot noise SN is measured for each redshift bin from $4\pi f_{sky}/N_{gal}$. The mean galaxy number density \bar{n} is shown in unit deg⁻². The best-fit galaxy bias, b_{bfit} , for each bin is shown with the 1σ uncertainty, marginalized over all model parameters. The magnification bias slope *s* is defined by $d \log N/dm$, where *N* is the number of galaxies and *m* is the magnitude limit adopted. ℓ_{max} is the maximum multiple in the galaxy autopower spectra.

Sample	$z_{\rm eff}^{\rm gg}$	$z_{\rm eff}^{\kappa g}$	$10^7 SN$	$\bar{n} (\text{deg}^{-2})$	b_{bfit}	S	$\ell_{\rm max}$
z_1	0.37	0.36	6.58	460.8	$1.39^{+0.94}_{-0.37}$	0.29	154
z_2	0.51	0.51	5.27	579.6	$1.37^{+0.92}_{-0.35}$	0.41	221
<i>z</i> ₃	0.65	0.65	8.41	363.6	$1.72_{-0.48}^{+1.12}$	0.57	283

where $x, y \in \{g, \kappa\}$ and $W^x(z)$ corresponds to the window function of the field that we will define in Sec. VI. Note that these redshifts are quoted to aid the interpretation of the typical distance at which a given galaxy signal arises—but they are not used in the modeling, which employs the full redshift distribution for each tomographic slice.

The shot noise and the effective redshifts of the galaxy samples used are summarized in Table I. The similarity between the z_{eff}^{gg} and z_{eff}^{kg} justifies the use of a constant effective galaxy bias for each redshift bin in Sec. VI.

Two systematic corrections were applied at the level of galaxy density maps for each tomographic bin. First, the correlation between the observed galaxy density and survey completeness was accounted for by including a completeness weight, defined as the inverse of the survey completeness for each pixel within the mask. Because the completeness was purely geometrical, it did not fully account for the number density variation with stellar density or extinction. Therefore, [38] applied a second correction to remove correlations with stellar density using the ALLWISE total density map. The stellar density map was originally in $N_{\rm side} = 512$ and then upgraded to the required $N_{\rm side} = 1024$. Because of the relatively smooth, large-scale variation in number density, they considered the difference in the initial $N_{\rm side}$ to not pose significant bias in their correction procedure. They fitted the correlation between completeness weighted density δ_g and stellar number N_{star} with a fifth-order polynomial.⁴ The particular choice of the functional form is simply to capture the smooth variation between δ_q and the systematics. The mean density of the galaxy field was computed using regions where completeness > 0.95 and $N_{\rm star} < 8.52 \times 10^3 \ {\rm deg^{-2}}$ (about 70% of the total legacy survey footprint). Appendix A shows that the level of stellar density correlation with the galaxy density maps within the DESI \times ACT footprint is about 0.1% at the scales we are interested in. They further excluded pixels with stellar density $N_{\text{star}} > 1.29 \times 10^4 \text{ deg}^{-2}$ where the average $|\delta_g| > 0.05$ in bins of N_{star} . Finally, they checked that the correlation between density fluctuations and the Milky Way extinction E(B - V) map [55] was consistent with zero.

Notice that a possible consequence of applying the corrections directly to the density maps is introducing monopole power, which is coupled and propagated into higher ℓ modes via the mask. We checked that, after applying the correction, the monopole is < 0.3%, and we subtract the monopole before the analysis.

For comparison with [35], Fig. 3 shows the selection of our galaxy sample compared to the DESI LRG target selection cuts. The major difference of the two samples is found in the $r - W_1$ and W_1 space. Our sample from [38] is deeper in W_1 compared to DESI LRGs. Compared to the full LRG sample, our sample is also at slightly lower redshifts—the sample drops off sharply beyond z = 0.8, whereas the DESI LRG has a tail beyond z = 1.2. Our sample is also 2–3 times denser than the DESI LRG. The redshift ranges of our sample, excluding bin 1, are most closely matched to redshift bins 1 and 2 in [35,36]. This shows that we are not in the shot-noise-dominated regime, and if the two samples are tracing the same large-scale structure, we expect the cosmological results to be consistent in the two studies.

B. ACT CMB lensing

Our baseline analysis utilizes the CMB lensing map produced in the sixth data release of the Atacama Cosmology Telescope (hereafter, DR6 lensing map [30,31]). This lensing map is produced using nighttimeonly CMB measurements made between 2017 and 2021 in the frequency bands of 90 and 150 GHz. The lensing map covers 9400 deg² of the sky and is signal dominated on scales of $\ell < 150$. Among many improvements, this map is produced using a cross-correlation-based estimator that makes use of several time-interleaved splits with independent instrument noise in each split to ensure that the lensing bias subtractions are insensitive to the modeling of the noise.

The CMB scales $\ell_{\rm CMB}$ used for the lensing analysis range from 600 < $\ell_{\rm CMB}$ < 3000. The large scales of the input CMB maps are excluded due to the presence of an instrument-related transfer function [56], Galactic foreground, and large atmospheric noise. The cut excluding $\ell_{\rm CMB}$ > 3000 was chosen to minimize contamination from extragalactic foregrounds like the thermal Sunyaev-Zeldovich effect (tSZ), the cosmic infrared background (CIB), and radio sources. Extragalactic foregrounds are further suppressed through the use of a profile-hardened lensing estimator [57,58]. This involves assuming a typical cluster profile for the tSZ and constructing a quadratic estimator that is insensitive to the CMB mode couplings arising from objects with radial profiles similar to the tSZ.

⁴The high-order polynomial was used to provide a smooth interpolation between the δ_q vs N_{star} relation.



FIG. 3. The DESI legacy survey galaxy sample used in this paper in color-color and color-magnitude space. The black lines indicate DESI LRG selection cuts, where the LRG sample is defined to the left of the lines in the first and third panel and to the right of the line in the middle panel. DECaLS galaxies are shown in blue and BASS-MzLS galaxies are shown in orange.

In Sec. VC we use simulations to show that the residual extragalactic contamination is negligible in our cross-correlation measurements.

The baseline ACT DR6 lensing mask is constructed from a Galactic mask that selects 60% of the sky with the lowest dust contamination and is apodized using a cosine roll-off along the edges. For consistency tests, we also employ lensing maps produced using only 40% of the sky, the region with the lowest dust contamination, or a slightly more restrictive mask compared to the baseline 60% mask in order to remove dust clouds along the edges of the 60% baseline mask near the Galactic plane when using CIB deprojection for foreground mitigation. We will subsequently refer to these masks as the 60%, 40%, and CIB-depj masks, respectively.

C. CMB lensing from Planck

The PR4 *Planck* lensing maps utilize CMB scales from $100 \le \ell \le 2048$ using the standard quadratic estimator [32]. This analysis improves over the PR3 analysis by using the reprocessed PR4 NPIPE maps that incorporate around 8% more data than the 2018 *Planck* PR3 release. Pipeline improvements, including optimal anisotropic filtering of both the CMB maps and the reconstructed lensing maps, resulted in an increase of the total signal-to-noise ratio by around 20% compared to the PR3 release.

III. CMB LENSING TOMOGRAPHY MEASUREMENTS

We measure the autospectra of the galaxy samples described in Sec. II A and their cross-correlation with the DR6 lensing map on ~20% of the sky using a pseudo- C_{ℓ} estimator that appropriately accounts for the impact of the mask induced mode coupling on the power spectrum between two fields. These pseudo- C_{ℓ} " \tilde{C}_{ℓ} " differ from the true underlying power spectrum C_{ℓ}^{true} due to the

effects of the mode coupling and their expectation value $\langle \tilde{C}_{\ell} \rangle$ is related to the true spectra as [59]

$$\langle \tilde{C}_{\ell} \rangle = \sum_{\ell'} M_{\ell \ell'} C_{\ell'}^{\text{true}}, \qquad (4)$$

where M is the mode coupling matrix that is purely a function of the mask. One can invert the above relationship approximately to extract the true power spectrum if the power spectrum is assumed to be piecewise constant across several discrete bins. We perform the mode decoupling operation on the binned power spectrum using the NaMaster code [60]. For the galaxy field, we use an apodized mask that contains the joint overlap between the DESI legacy survey and ACT lensing (see Fig. 1). The mode coupling matrix for the cross-correlation is thus computed using the galaxy mask and the square of the analysis mask used for lensing.⁵ This is to account for the fact that the lensing reconstruction with a quadratic estimator takes the product of two filtered, masked CMB fields. We employ HEALPix maps with $Nside = 1024^6$ and run NaMaster with $\ell_{max} =$ 3000 despite using only multipoles $\ell \leq 300$ in our cosmology range.

We show the cross-correlation between the DESI legacy galaxies and ACT DR6 lensing reconstructions in Fig. 4. The best-fit Λ cold dark matter (LCDM) model obtained for the cross-correlation of DESI legacy with ACT DR6 and *Planck* PR4 CMB lensing are shown in red and gray,

⁵This is a good approximation for the regime where the variations of the mask are on much larger scales than the CMB and lensing scales of interest. This approximation is done because the lensing signal is reconstructed using a quadratic estimator where each of the two CMB maps carries one power of the mask.

⁶The original lensing maps come with resolution Nside = 2048 that we downgrade to match the resolution used in the analysis.



FIG. 4. Measurements of C_{ℓ}^{kg} (top row) and C_{ℓ}^{gg} (bottom row) for the three redshift samples of DESI legacy galaxies. Within our analysis range the cross-correlation between galaxies and CMB lensing is detected at SNRs of 5, 9, and 15, respectively. Combined with *Planck* PR4 lensing, the cross-correlation of the joint redshift bins is measured at an SNR of 22. We show in dashed red (gray) the best fit from the joint fit to all redshift samples for the ACT DR6 (Planck PR4) analyses. The difference between the two datasets stems from *Planck* having a larger overlap with the DESI galaxies compared to ACT, resulting in a different overall bias. The subpanels show the model residuals. The total model χ^2 for the joint fit is 25.9 and for 24 bandpowers we estimate to have a PTE of 0.36.

respectively. We obtain a minimum χ^2 of 17.5 for ACT. We estimate a probability to exceed (PTE) of 0.13 for 12 degrees of freedom and a χ^2 of 6.34 with a PTE of 0.9 for the cross-correlation with Planck PR4. For the combination of DR6 + PR4 \times DESI with 24 degrees of freedom, we obtain a minimum χ^2 of 25.9 corresponding to a PTE of 0.36. We note from the sixth panel of Fig. 4 that, for the third redshift bin, the best fit C_{ℓ}^{gg} is significantly different for ACT and *Planck* and this is mainly attributed to the difference in the footprint of the ACT and Planck lensing maps, as marked by the blue and black regions in Fig. 1. The sky fraction for the ACT and Planck cases are $f_{\rm sky} = 0.160$ and $f_{\rm sky} = 0.334$, respectively. We speculate that this may be due to the difference in photometry in the DECaLS and BASS-MzLS regions, leading to a slight difference of galaxy selections in bin 3 that was absorbed into the galaxy nuisance parameters. Hang et al. [35] find a similar, larger than expected difference between the north and south in the DESI LRG samples. However, these differences in biases would not impact on giving consistent

cosmology parameters if appropriately accounted for. In Sec. V we will show null tests of bandpower consistency between the different regions in the ACT footprint showing that the bandpower differences are consistent with null.

We show in Sec. IV that we can recover the theory spectra from simulations to within better than 1% on the scales of interest and give details on the computation of the covariance matrix using simulations.

A. Scale cuts

We measure the cross-correlation between the ACT DR6 lensing reconstruction and the DESI galaxies with signal-to-noise ratios of 5, 9, and 15 for each of the three galaxy samples, respectively, within our cosmology analysis range of $50 \le \ell \le \ell_{\text{max}}^{g_{\text{in}}}$ with $\ell_{\text{max}}^{g_{\text{in}}} = 1.54, 221, 283 \times 10^8$.

of $50 \le \ell \le \ell_{\text{max}}^{g_i}$ with $\ell_{\text{max}}^{g_i} = 1.54, 221, 283 \times 10^8$. The large-scale cut $\ell_{\text{min}} = 50$ is chosen due to potential misestimation of the lensing reconstruction mean-field signal, which can lead to underestimation of error, and to ensure insensitivity to potential large-scale galactic systematics. We further check that on the scales used the

	Hang et al. 2021	This work
$f_{\rm sky}$	0.334 (Planck PR3);	0.160 (ACT); 0.334 (Planck PR4);
Redshift bins	0, 1, 2, 3;	1, 2, 3;
ℓ_{\min}	10;	50;
$\ell_{\rm max}$	100, 200, 300, 400, 500 (all bins);	154 (bin 1); 221 (bin 2); 283 (bin 3);
$\Delta \ell$	10;	60;
Pseudo- C_{ℓ}	HEALPix anafast with $f_{\rm sky}$ correction;	NaMaster;
Model	HALOFIT $P(k)$ with phenomenological nonlinear bias: $P_{am}(k) = b_1 P_{am}^{lin}(k) + b_2 [P^{nl}(k) - P^{lin}(k)]$:	HALOFIT $P(k)$ with linear bias: $P_{em}(k) = bP_{em}^{nl}(k)$:
Cosmology	Fixed to <i>Planck</i> 2018 best-fit parameters with a free lensing amplitude A_{κ} ;	Free (see Sec. VI);
Nuisance parameters	The redshift distribution $n(z)$ and galaxy biases b_1 , b_2 for each bin are fixed from galaxy clustering $C_{\ell}^{g_ig_j}$, shot noise fixed to $SN = 4\pi f_{sky}/N_{gal}$.	Redshift distribution fixed to [38]; galaxy biases and shot noise for each bin are free.

TABLE II.	Summary of different	analysis choices bet	ween [38] and	this work. T	The differences	in measurements a	tre shown in the
upper half of	the table, and the diff	ferences in modeling	g are shown in	the lower h	alf.		

contribution beyond the Limber effect from redshift-space distortions (RSDs) is negligible compared to the uncertainties at a level of at most 0.5%.

The maximum multipole scale cut $\ell_{\max}^{g_i}$ was mainly chosen by requiring an unbiased recovery of the cosmological parameters; given the linear bias model adopted in this work (as described in Sec. VI), this becomes challenging on small scales. Specifically, the $\ell_{\max}^{g_i}$ are chosen by examining the fiducial $C_{\ell,\text{fid}}^{\kappa g}$ and making the cut at the point where the nonlinear part of the matter power spectrum' becomes 10% of the linear power spectrum. These angular scale cuts correspond to spatial scales of $k_{\text{max}} = 0.15, 0.16, 0.17h/\text{Mpc}$ for the three redshift bins, respectively. Given our $\Delta \ell$ binning and the choices for ℓ_{max} , we tested that the actual nonlinear contribution is less than 5% for all data points. The nonlinearity compared to the uncertainties on the data point, $\Delta C_{\ell} / \sigma_{\ell} \equiv (C_{\ell}^{\text{nl}} - C_{\ell}^{\text{lin}}) / \sigma_{\ell}$, is most significant at highest ℓ . For the smallest scale used we find $\Delta C_{\ell}/\sigma_{\ell} = 0.21, 0.30, 0.41$ for tomographic bins 1–3, respectively. The same set of scale cuts are applied to the galaxy autocorrelation, for which we find $\Delta C_{\ell}/\sigma_{\ell} = 0.71$, 1.46, 1.84. Although the nonlinear contribution is larger as expected given the smaller uncertainties in the galaxy autospectrum, the above values are expected to be smaller when we marginalize over the linear bias. The constraining power on S_8 comes mainly from the information of the crosscorrelation-i.e., schematically by taking the ratio of $C_{\ell}^{gg}/(C_{\ell}^{\kappa g})^2$ to break the degeneracy between bias b and σ_8 that C_{ℓ}^{gg} is subjected to. (This ratio is dominated by the uncertainties in the galaxy-lensing cross-spectrum and not limited by the error in C_{ℓ}^{gg} .) In practice, our baseline model when fitting the actual data uses the nonlinear power spectrum, rather than purely linear theory.

To ensure that this choice of ℓ_{max} does not bias the recovery of cosmological parameters, in Appendix B 1, we show that the cosmology constraints on ACT DR6 × DESI legacy galaxies using more conservative ℓ_{max} (discarding z_1 and limiting k_{max} of z_2 and z_3 to $k_{\text{max}} = 0.12, 0.13h/\text{Mpc}$, respectively, results only in a shift of S_8 by 0.03σ , consistent with our baseline measurements).

We also generated theoretical data vectors using the nonlinear power spectrum and obtained fits using the linear matter power spectrum.⁸ We recovered unbiased S_8 within 0.1 σ given the scale cuts. Furthermore, we perform the above consistency test on reanalyzing the previous dataset, using the *Planck* PR3 lensing map. As shown in Appendix B, at the S_8 level the difference between linear and nonlinear modeling is small (0.19 σ), providing further evidence that the scales chosen are insensitive to nonlinear modeling.

Table II summarizes the main measurement differences between [38] and this work.

IV. PIPELINE VERIFICATION

To test the accuracy of our pipeline, we measure the autocorrelation of the simulations described in Sec. IVA and their correlation with the lensing reconstruction simulations in the same manner as we treat the data. We then compare the average of the measured, binned, and modedecoupled spectra to the input power spectra. To make a fair

⁷This is computed using CAMB, assuming a fiducial *Planck* 2018 cosmology by subtracting the linear power spectrum from the total nonlinear power spectrum obtained using HALOFIT [62].

⁸In both cases, we use the linear galaxy bias. In Appendix B, we also test the effect of a scale-dependent bias using *Planck* PR3 and adopting the empirical two-bias model showing that the constraint is consistent with the one obtained with the baseline linear bias.



FIG. 5. Recovery of C^{gg} and $C^{\kappa g}$ on simulations after correcting for the effect of mode coupling and inversion of mode coupling matrix is better than the percent level. The lightly shaded error bars show the measurement errors, and the darker error bars show the error on the mean of 400 Gaussian simulations.

comparison, we account for the fact that the mode decoupling is approximate and convolve the input $C_{\ell,\text{fid}}$ spectra with the appropriate mode coupling matrix before binning them and applying the approximate decoupling matrix applied to the measured \hat{C}_{ℓ} 's. As can be seen in Fig. 5, we can recover the inputs with our measured spectra $\langle \hat{C}_{\ell}^{kg} \rangle$ and $\langle \hat{C}_{\ell}^{gg} \rangle$ to within $\lesssim 1\%$ well beyond our cosmology range.

A. Simulations for the covariance matrix and pipeline verification

We generate a set of 400 ACT DR6 CMB lensing reconstruction simulations with appropriately correlated Gaussian simulations of galaxy number density fields. These simulations are used to test the recovery of the autoand cross-power spectra and to compute the covariance matrices for the baseline measurements and the null tests.

The lensing simulations are obtained by displacing a randomly drawn CMB realization with a Gaussian lensing convergence field. We then add realistic survey noise to the lensed CMB [63] and mask the data, before passing the resulting CMB skies into the same lensing reconstruction pipeline that is applied to the real data [30].

We obtain Gaussian realizations of the galaxy field which are appropriately correlated with the lensing simulations above by filtering the input lensing convergence as follows using fiducial $C_{\ell,\text{fid}}^{gg}$, $C_{\ell,\text{fid}}^{\kappa g-9}$:

$$a_{\ell m}^{g} = \frac{C_{\ell, \text{fid}}^{\kappa g}}{C_{\ell, \text{fid}}^{\kappa \kappa}} a_{\ell m}^{\kappa} + a_{\ell m}^{g, \text{uncorrelated}} + a_{\ell m}^{g, \text{noise}}.$$
 (5)

We added the shot-noise component as white noise on the galaxy spectra. Notice that aliasing and the pixel window function could lead to biases in cosmological parameters, as pointed out by [64]. This effect is most significant for low resolution maps, and at $N_{\text{side}} = 1024$, the impact on the

recovered power spectrum is ~0.5%. The fiducial spectra are convolved with the appropriate pixel window function for our map resolution of Nside = 1024. The part of the galaxy field that is uncorrelated with lensing $a_{\ell m}^{g,\text{uncorr}}$. and the Gaussian noise $a_{\ell m}^{g,\text{noise}}$ is obtained from

$$\langle a_{\ell m}^{g,\text{uncorr.}}(a_{\ell'm'}^{g,\text{uncorr.}})^* \rangle = \delta_{\ell\ell'} \delta_{mm'} \left(C_{\ell,\text{fid}}^{gg} - \frac{(C_{\ell,\text{fid}}^{\kappag})^2}{C_{\ell,\text{fid}}^{\kappa\kappa}} \right), \quad (6)$$

$$\langle a_{\ell m}^{g,\text{noise}}(a_{\ell'm'}^{g,\text{noise}})^* \rangle = \delta_{\ell\ell'} \delta_{mm'} C_{\ell',\text{noise}}^{gg}.$$
 (7)

We do not include correlations between the different galaxy samples, but simply estimate the Gaussian analytic approximation for the covariance between the different samples as described in Sec. IV C.

B. Lensing Monte Carlo transfer function

As discussed in detail in [30,34], the lensing maps obtained from performing lensing reconstruction under the presence of a mask are misnormalized. This is corrected using Gaussian simulations by computing the ratio of the cross-correlation between the appropriately masked input lensing convergence with the lensing reconstruction to the autocorrelation of the known input convergence. Specifically for the case of the cross-correlation this requires computing

$$A_{\ell}^{\mathrm{MC}} = \frac{C_{\ell}^{\hat{\kappa}_{\mathrm{in}, \ell-\mathrm{mask}} \kappa_{\mathrm{in}, g-\mathrm{mask}}}}{C_{\ell}^{\hat{\kappa}_{\mathrm{K}_{\mathrm{in}, g-\mathrm{mask}}}}},$$
(8)

where $\hat{\kappa}$ is the masked CMB lensing reconstruction, κ – mask is the input lensing convergence masked with the lensing mask, and $\kappa_{\text{in},g-\text{mask}}$ is the input lensing convergence masked with the galaxy mask. Since this correction depends on the region of the overlap, we estimate this separately for the analyses that use the ACT DR6, *Planck* PR4 and PR3 footprint. We estimate these using 480 DR6 Gaussian simulations and corresponding lensing

⁹These theory curves are obtained following the same prescription of [38], based on fits to the data fixed to the *Planck* cosmology using a two-bias model.



FIG. 6. Size of the Monte Carlo transfer function correction for the different lensing maps estimated using simulations described in Sec. IVA.

reconstructions provided by [32] to obtain unbiased estimates of CMB lensing cross-correlation in the form of

$$C_{\ell}^{\hat{\kappa}g} \to A_{\ell}^{\mathrm{MC}} C_{\ell}^{\hat{\kappa}g}.$$
 (9)

Figure 6 shows the size of these Monte Carlo corrections for our analyses. On the scales of our analyses, these amount to a change of at most $\sim 3\%$ to the measured cross-correlation.

C. Covariance matrices

We compute the covariance $\text{Cov}(C_{\ell}^{XY}, C_{\ell'}^{AB})$ for $XY, AB \in \{gg, \kappa g\}$, using the suite of 400 Gaussian simulations of the galaxy and lensing fields discussed in Sec. IV A. We take a hybrid approach where the covariance within each tomographic bin is estimated from the simulations, while the intertomographic bins correlation is computed analytically assuming Gaussian signal and noise. The diagonal part of the covariance matrix, including the diagonals between gg and κg between the same galaxy samples, are shown in Fig. 7. The correlation matrix for one of the three galaxy samples is shown in Fig. 8.

Our Gaussian simulations do not capture the correlations between the different galaxy samples. Therefore, to measure the off-diagonal covariance blocks used in the joint analysis of the three galaxy samples, we approximate these analytically using the Gaussian covariance module implemented in NaMaster [60,65], which requires the following as inputs: the fiducial input spectra used for our simulations; theory curves for the galaxy cross-spectra between the different redshift bins obtained using the best-fit biases used in the fiducial input spectra to obtain the input Gaussian simulations (as described in Sec. IVA) and the assumed n(z); and a curve of C_{ℓ}^{κ} including reconstruction noise appropriate to the level of ACT DR6. The different



FIG. 7. The diagonal elements of the covariance matrix, as well as the off-diagonal covariance between C_{ℓ}^{gg} and $C_{\ell}^{\kappa g}$ in dotted lines for the different galaxy samples. Shaded regions show the small-scale cutoff used for the different samples. Red, blue, and green denote the first z_1 , second z_2 , and third z_3 redshift bins, respectively.

samples of C_{ℓ}^{kg} are correlated at the level of (26%–20%) between neighboring bins and (5%–3%) between next to neighbor bins. The correlation between the different C_{ℓ}^{gg} are at the 5% level for neighboring bins and almost negligible for next to neighbor bins.

We verify that the diagonal elements of the covariance matrix computed using simulations agree well with the diagonal covariance estimated using the analytic estimation from the Gaussian covariance module implemented in NaMaster. We additionally validate the error estimates from



FIG. 8. Correlation matrix for z_3 of the ACT × DESI galaxies. The diagonal elements of the correlation matrix have been subtracted for improved legibility. While the correlations between different scales are small, there exist correlations between C_{ℓ}^{gg} and C_{ℓ}^{kg} of up to 20%.



FIG. 9. The theory (solid) and measured (dots) cross-correlation coefficients for galaxy clustering, $C_{\ell}^{ij}/\sqrt{C_{\ell}^{ii}C_{\ell}^{jj}}$, for redshift bins *i* and *j*. The bin combination is shown in the legend as (i, j). The smooth theory curves are generated using the best-fit [38] model to the C_{ℓ}^{gg} autocorrelations with the ACT × DESI joint footprint. Error bars are Gaussian computed using NaMaster assuming the theory curves. The gray region indicates the lower ℓ cut, and the colored regions indicate the upper ℓ cuts, with the colors corresponding to the specific (i, j) combination.

simulations using a jackknife approach. We first downgrade the joint mask to $N_{\rm side} = 8$ and exclude the downgraded mask pixels with value > 0.95 one at a time, giving a total of 86 jackknife samples. We found that, generally, this approach gives errors around 5% larger than the analytical error bars and is consistent with our baseline simulationbased approach. We further verify that the binning introduced with $\Delta \ell = 60$ helps to reduce the off-diagonal correlations of the covariance matrix compared to the $\Delta \ell =$ 10 adopted in [38] and yields a converged estimate of the covariance matrix given the finite number of simulations used. We find correlations of up to 20% between C_{ℓ}^{gg} and $C_{\ell}^{\kappa g}$ at the same ℓ , while the off-diagonal correlations are smaller than 10%. Figure 9 shows the theory input for the off-diagonal covariance in the form of the correlation coefficients, $C_{\ell}^{ij}/\sqrt{C_{\ell}^{ii}C_{\ell}^{jj}}$ for bins *i* and *j*, compared to the actual measured values. These theory inputs are based on the best-fit [38] models fixed at the Planck 2018 cosmology to the autocorrelations C_{ℓ}^{gg} , hence not a fit to the measurements. However, the fact that they match closely means that our covariance estimate is relatively accurate. The only noticeable difference is the (2,3)correlation, where the theory is slightly higher than the measurements. This implies that our uncertainties on the final results may be slightly overestimated.

Wherever we use the inverse of the covariance matrix, we account for the fact that the inverse of the above covariance matrix is not an unbiased estimate of the inverse covariance matrix, and we rescale the inverse covariance matrix by the Hartlap factor [66],

TABLE III. Summary of the lensing null tests described in Sec. VA and galaxy homogeneity null tests of Sec. VB. For each test, we show the PTE values for the baseline range of each redshift bin. The first nine tests are described in Sec. VA and the last three tests are described in Sec. VB.

Null test	PTE z_1	PTE z_2	PTE z_3
$\overline{\operatorname{Curl} \times g}$	0.18	0.14	0.04
$(f090 - f150) \times g \text{ TT}$	0.12	0.03	0.49
$(f090 - f150) \times g \text{ MV}$	0.69	0.14	0.74
$\kappa^{\rm MV} \times g - \kappa^{\rm cib-dpj} \times g$	0.13	0.18	0.18
$\kappa^{\rm MV} \times g - \kappa^{40\%{\rm mask}} \times g$	0.22	0.84	0.44
$\kappa^{f150MV} \times g - \kappa^{f090MV} \times g$	0.70	0.27	0.89
$\kappa^{f150TT} \times g - \kappa^{f090TT} \times g$	0.90	0.89	0.96
$\kappa^{\rm MV} \times g - \kappa^{\rm TT} \times g$	0.58	0.43	0.47
$\kappa^{\rm MV} \times g - \kappa^{\rm MVPOL} \times g$	0.96	0.41	0.96
$C_{\ell}^{gg,\text{north}} - C_{\ell}^{gg,\text{south}}$	0.22	0.88	0.12
$C_{\ell}^{gg,60\%\text{mask}} - C_{\ell}^{gg,40\%\text{mask}}$	0.68	0.79	0.23
$C_{\ell}^{gg,60\%\text{mask}} - C_{\ell}^{gg,cib-dpj\text{mask}}$	0.20	0.20	0.81

$$\alpha_{\rm cov} = \frac{N_s - N_{\rm bins} - 2}{N_s - 1},\tag{10}$$

with $N_s = 400$ simulations and the 12 combined data points for C_{ℓ}^{gg} and $C_{\ell}^{\kappa g}$, this correction is $\alpha_{\rm cov} = 0.97$ for the joint analysis of the three galaxy samples.

In Sec. VII A 1 we also present a combined analysis of the ACT DR6 cross-correlation with the DESI galaxies and the equivalent cross-correlation analysis using *Planck* PR4 lensing reconstructions. We outline the procedure used to estimate the covariance matrix for this joint analysis below. We use the same set of 480 FFP10 CMB simulations used in the *Planck* PR4 lensing analysis [32]. These lensed Gaussian simulations have corresponding lensing reconstructions generated using the PR4 lensing pipeline. In a similar manner as described in Sec. IVA, we obtain a set of correlated Gaussian galaxy realizations allowing us to estimate the covariance for the *Planck* cross-correlation analysis.

As shown in [30], the correlation between the ACT and the *Planck* lensing reconstructions is relatively small. However, there exist large correlations of the order 40%–50% in the cross-correlation despite the partial overlap in survey areas and scales because of the identical galaxy sample used in both cross-correlations. The correlation between the galaxy autocorrelation measured on the ACT and *Planck* footprints is up to 60% (this is accounted for appropriately in the covariance matrix when combining the ACT DR6 and *Planck* PR4 measurements). As in the ACT-only analysis we analytically estimate the covariance between the different galaxy samples; this is again small for the case of C_{ℓ}^{gg} (< 6%) but non-negligible for $C_{\ell}^{\kappa g}$ (up to 30%).

V. TESTS FOR SYSTEMATIC ERRORS

We perform a series of comprehensive null and systematic tests that we discuss below. This test suite establishes that our data are free from significant systematic effects that may affect our measurements. For example, we test for contamination of the lensing reconstruction by extragalactic foregrounds (e.g., tSZ and CIB) that also correlate with the galaxy samples. Furthermore, we also examine contamination from galactic dust by considering different galactic masks. Such contamination could correlate with the galaxy survey data due to correlations of the galaxy sample with galactic structure (e.g., stellar density and dust absorption).

The null tests aimed at systematic errors in the lensing reconstruction correlated with the galaxy samples are found in Sec. VA; tests targeting the spatial inhomogeneity of the galaxy samples are found in Sec. V B. Finally, we estimate the biases in the lensing cross-spectra due to extragalactic foreground contamination of the lensing reconstruction using simulations in Sec. V C.

A. Null tests for contamination of the lensing reconstruction

We perform tests to show that our lensing reconstruction is free from systematic effects that are potentially correlated with the galaxy samples. Such contamination can, in principle, be caused by the tSZ effect, which is produced by the inverse Compton scattering of CMB photons off hot electrons in thermal clusters, and by CIB contamination originating from unresolved dusty galaxies. Both of these astrophysical foregrounds produce nontrivial bispectra and trispectra which can bias the lensing reconstruction if not appropriately mitigated [67]. They are also correlated with the large-scale matter distribution and thus with the galaxy densities, so cross-correlations are also susceptible to biases.

The baseline DR6 lensing map used in this work uses profile hardening to mitigate these biases (see [30,67] for details). We verify here that the contaminants are mitigated sufficiently in cross-correlations using this baseline lensing map. Furthermore, as well as extragalactic foregrounds, the correlation of galaxy density with Galactic structures can also bias the lensing reconstruction In particular, dust contamination has been tested extensively for the lensing reconstruction by using more conservative galactic masks and changing the minimum CMB multipole used for the lensing reconstruction [30]. Their effect on crosscorrelations can be tested with different galactic masks.

Null tests here are divided into two categories: first, cross-correlation of signal-nulled lensing reconstructions with the galaxy samples, where the signal-nulled lensing reconstructions are constructed by taking the difference of the temperature and polarization maps at 150 and 90 GHz to obtain a map containing only noise and foreground



FIG. 10. Distribution of the PTE of the 36 null tests. The distribution is consistent with uniform passing of the K-S statistic with a PTE of 0.15.

residuals on which lensing reconstruction is performed. The second category includes bandpower difference tests where the cross-correlation of the same galaxy samples with lensing reconstructions obtained with different versions of CMB maps are taken and checked for consistency. The results for these null tests are summarized in Table III and the histogram of Fig. 10. We define the criterion for passing a null test to be that it returns a PTE greater than 0.05.¹⁰

The null tests specifically targeting the extragalactic foregrounds leverage the distinctive frequency dependence of these foregrounds. We cross-correlate this signal-nulled lensing map with our galaxy samples. The results for these tests are passing except for a marginal failure with PTE = 0.03 for the cross-correlation of redshift bin 2 with the temperature-only lensing null map. (The bandpowers of this test can be seen in Fig. 25 of Appendix C.) Given that the same test using the temperature and polarization null map passes and no failure is observed with the bandpower-level test discussed below, we attribute this failure to random fluctuations. We also investigate the bandpower difference between the cross-correlations measured using the reconstruction performed only on the 150 and 90 GHz data and find no failures for those tests.

We also explicitly test that profile hardening is effective in mitigating CIB foregrounds by performing a bandpower test difference between the cross-correlation of the DESI galaxies with the minimum variance temperature and polarization (MV) baseline lensing map and a lensing map that explicitly deprojects the CIB (cib-dpj) using the *Planck* high-frequency channels, finding consistency

¹⁰We also flag tests passing with PTE > 0.95 although these are not particularly worrying given that tests like curl × galaxies that specifically test for our covariance matrices give reasonable results.



FIG. 11. We estimate biases due to extragalactic foregrounds using realistic foreground simulations from WebSky [68]. We perform lensing reconstruction on foreground-only maps using different foreground mitigation strategies and cross-correlate them with galaxy number density maps that we obtain by populating the WebSky halo catalog using a HOD. We find that our baseline analysis reduces all biases to $< 0.2\sigma$ while the analysis without any mitigation yields significant biases (up to $\sim 3\sigma$).

in both approaches. In addition, we find consistent bandpower results using a more restrictive Galactic 40% mask compared to our baseline 60% mask that does not significantly affect our measurements.

We also cross-correlate the galaxy samples with the curl modes of the baseline reconstruction. This test is primarily a test of our covariance estimation since we do not expect there to be any physical signal arising from the curl of the lensing reconstruction at the current levels of precision. This test passes for all redshift bins except for bin 3 which has a marginal failure of PTE = 0.04 (see Fig. 26 in Appendix C).

No significant failures are observed apart from two marginal failures of the curl \times galaxies test in redshift bin 3 at the level of 0.04 and the frequency-nulled map in temperature \times galaxy test at redshift bin 2 which fails with 0.03. Further evidence that the above failures are consistent with fluctuations can be seen in Fig. 10, the distribution of the PTE for the null tests is consistent with a uniform distribution and given the fact that we perform 36 null tests, observing 2 null tests at around the 5% level is not unlikely. Indeed, the Kolmogorov-Smirnov (K-S) statistics indicate that this distribution is consistent with uniform¹¹ with a PTE = 0.15.

B. Homogeneity and contamination tests of the galaxy samples

We perform tests on the homogeneity of the galaxy sample by constructing null tests using different masks. This is tested through the cross-correlation test using our 60% and 40% Galactic masks discussed in the previous section as well as bandpower-level autospectra null tests where C_{ℓ}^{gg} is measured across different regions: in particular, we compare the measurement using the 60% and 40% Galactic masks, the 60% mask and the footprint covered by the CIB-deprojection mask and splitting the samples between the north and south galactic caps (i.e., the disjoint regions marked by the blue lines in Fig. 1). All of these tests pass, suggesting that the legacy sample used is uniform within the DESI × ACT footprint on the scales tested without large variations of galaxy bias, shot noise, and redshift distribution across the sky. However, as mentioned in Sec. III, we do observe a fluctuation in the galaxy bias between the DESI × ACT footprint and the DESI × PR4 footprint. (Roughly speaking, the difference between the blue and pink regions in Fig. 1. This is mainly the BASS-MzLS footprint of the DESI legacy survey.)

C. Simulation-based tests for extragalactic foregrounds

We further quantify the potential contamination from extragalactic foregrounds using a simulation-based approach with foreground simulations from WebSky [68] in an approach similar to [67] by making the approximation that all the relevant extragalactic foregrounds are in the temperature channel such that the observed CMB temperature is given by $T = T_{\text{CMB}} + T_{\text{fg}}$. This assumption is valid as we do not expect the polarized tSZ and CIB to cause significant lensing biases at current observation levels [69]. Furthermore, any potential bright polarized sources in DR6 are masked and inpainted before lensing reconstruction [30]. The bias on the cross-correlation of the temperature-only lensing reconstruction with the galaxy samples due to unmitigated foreground contamination is then given by

$$\Delta C_{\ell}^{\kappa g} = \langle \mathcal{Q}(T_{fg}, T_{fg})g\rangle, \tag{11}$$

where $Q(T_A, T_B)$ denotes the quadratic estimator used to reconstruct the lensing convergence field from the two fields T_A and T_B , and the cross-correlation of a field X with the galaxy field g is denoted with the shorthand

¹¹We note that while these 36 tests are not strictly independent (e.g., some involve overlapping subsets of the data), the correlation among them is mild enough that we still expect the K-S result to be a reasonable indicator of whether the null tests, as a whole, behave consistently with uniform passing.

 $C_{\ell}^{Xg} = \langle Xg \rangle$. In Eq. (11) we have assumed that the foregrounds are uncorrelated with the CMB, allowing us to neglect terms of the form $\langle Q(T_{\text{CMB}}, T_{fg}) \rangle$. We quantify the bias due to extragalactic contamination in terms of the bias in the cross-correlation lensing amplitude, A_{lens}^{\times} . The bias A_{lens}^{\times} relative to the uncertainty of this lensing amplitude is given by

$$\frac{\Delta A_{\text{lens}}^{\times}}{\sigma(A_{\text{lens}}^{\times})} = \frac{\sum_{\ell,\ell'} \Delta C_{\ell'}^{\kappa g} \mathbb{C}_{\ell\ell'}^{-1} C_{\ell'}^{\kappa g}}{\sqrt{\sum_{\ell,\ell'} C_{\ell}^{\kappa g} \mathbb{C}_{\ell\ell'}^{-1} C_{\ell'}^{\kappa g}}},$$
(12)

where $C_{\ell'}^{kg}$ is the true, baseline cross-correlation signal and $\mathbb{C}_{\ell\ell'}$ is the associated covariance matrix.

We test the foreground mitigation strategies employed in the lensing maps by cross-correlating lensing reconstructions performed on foreground-only maps with WebSky and galaxy samples prepared by populating the WebSky halo catalog with galaxies using the Halo occupancy distribution (HOD) described in [38]. The resulting galaxy catalog is sampled to match the redshift distribution of our samples.

We find that the baseline foreground mitigation strategy adopted for the ACT DR6 lensing maps is effective in suppressing biases in the cross-spectra, resulting in negligible bias levels of $\Delta A_{\text{lens}}^{\times} / \sigma(A_{\text{lens}}^{\times}) = -0.01$ for all the galaxy samples. Additionally, explicitly deprojecting the CIB does not help in reducing foreground biases further although it is effective in suppressing the foreground biases compared to the no mitigation case. This is consistent with other findings [34,36,67], indicating that profile hardening is effective in suppressing not only tSZ clusters but also diffuse foregrounds from the CIB (see [58] for a detailed explanation). For comparison, the resultant biases when not performing profile hardening are at the $\Delta A_{\text{lens}}^{\times} / \sigma(A_{\text{lens}}^{\times}) =$ -0.19, -0.18, -0.07 level for the three galaxy samples, respectively (see Fig. 11 for a summary). From this test, we have evidence that extragalactic biases are negligible compared to the size of our statistical error, after profile hardening.

VI. MODELING AND ANALYSIS CHOICES

In this section, we introduce the model used to compare to the measured spectra presented in Sec. III when measuring cosmological parameters. We further present our blinding strategy and discuss the likelihood and priors adopted for the analysis.

A. Models for galaxy clustering and CMB lensing power spectra

We use HALOFIT to model the three-dimensional power spectra of the clustering of matter $P_{mm}(k, z)$. We model the linear galaxy bias b(z) in each redshift bin as a single effective number that is scale independent; such that in each

tomographic redshift bin we have $P_{gg} = b(z)^2 P_{mm}$ and $P_{gm} = b(z)P_{mm}$.

We use the Limber approximation [70,71] to project the three-dimensional matter power spectra along the line of sight to obtain the angular galaxy auto- and galaxy-lensing cross-spectra,

$$C_{\ell}^{gg} = \int \frac{d\chi}{\chi^2} [W^g(z)]^2 P_{gg} \left(k = \left(\ell + \frac{1}{2} \right) / \chi, z \right),$$

$$C_{\ell}^{\kappa g} = \int \frac{d\chi}{\chi^2} W^{\kappa}(z) W^g(z) P_{gm} \left(k = \left(\ell + \frac{1}{2} \right) / \chi, z \right), \quad (13)$$

where $z(\chi)$ is the redshift and is implicitly a function of χ , the comoving distance.

As shown in [38], bias evolution is important when considering wide tomographic bins, for example, when all four tomographic bins are combined in a single redshift bin. For the individual redshift bins, however, we tested that a constant mean bias gives consistent lensing amplitude A_{κ} (in the same set up as [38]) to the case of accounting for the bias evolution.

The projection kernels used for the galaxies and CMB lensing are given by

$$\begin{split} W_g(z) &= \frac{H(z)}{c} n(z), \\ W_\kappa(z) &= \frac{3}{2} \Omega_m \left(\frac{H_0}{c}\right)^2 (1+z) \frac{\chi(\chi_* - \chi)}{\chi_*}, \quad (14) \end{split}$$

where n(z) is the normalized redshift distribution for the galaxy sample, Ω_m is the total matter density that includes the density of neutrinos, H(z) is the Hubble rate with $H_0 = H(z = 0)$, and χ_* is the conformal distance to the surface of last scattering.

The observed angular power spectrum also contains contributions from the lensing magnification bias. This effect arises from the gravitational lensing of galaxies by foreground structures, inducing a magnification (or demagnification) affecting the sample selection by artificially increasing (decreasing) the magnitude of a galaxy in a correlated way with the large-scale structure. Denoting quantities related to the magnification bias by μ , we model the contributions from the magnification bias as

$$C_{\mathscr{C}}^{g\mu} = \int d\chi \frac{W_g(z)W_\mu(z)}{\chi^2} P_{gm} \left(k = \left(\ell + \frac{1}{2} \right) / \chi, z \right),$$

$$C_{\mathscr{C}}^{\kappa\mu} = \int d\chi \frac{W_\kappa(z)W_\mu(z)}{\chi^2} P_{mm} \left(k = \left(\ell + \frac{1}{2} \right) / \chi, z \right),$$

$$C_{\mathscr{C}}^{\mu\mu} = \int d\chi \frac{W_\mu^2(z)}{\chi^2} P_{mm} \left(k = \left(\ell + \frac{1}{2} \right) / \chi, z \right), \quad (15)$$

with the lensing magnification kernel given by

$$W_{\mu}(z) = (5s_{\mu} - 2)\frac{3}{2}\Omega_{m}H_{0}^{2}(1+z)$$
$$\times \int_{\chi}^{\chi_{*}} d\chi' \frac{\chi(\chi' - \chi)}{\chi'} H(z')n(z').$$
(16)

The parameter $s_{\mu} \equiv d \log_{10} N/dm$ is the response of the galaxy number density to a change in magnitude and is measured from the data by perturbing the photometry of the DESI galaxies and reapplying the selection criteria [38]. The total, observed, galaxy, and galaxy-CMB lensing spectra are then given by $C_{\ell}^{gg} + 2C_{\ell}^{g\mu} + C_{\ell}^{\mu\mu}$ and $C_{\ell}^{\kappa g} + C_{\ell}^{\kappa \mu}$.

To evaluate Eqs. (13)–(16), we use CLASS_SZ [72,73], a machine-learning accelerated CMB and large-scale structure code written in Python and C, that builds on top of the CLASS [74,75] infrastructure. CLASS_SZ¹² has a parallelized implementation of the Limber integrals and uses neural network emulators for the matter power spectrum. Together, this makes the model evaluation fast (roughly, 0.3 s for evaluation of all C_{ℓ}^{gg} and C_{ℓ}^{kg} in three redshift bins). The emulators are presented in detail in [76] and are based on COSMOPOWER [77].

We find that the impact of the magnification term $C_{\ell}^{\kappa m}$ compared to the signal $C_{\ell}^{\kappa g}$ is about 0.5%–1% for the third redshift bin (where the effect is largest), and including this term only shifts the cosmological parameters by $\sim 0.1\sigma$, given a uniform prior of (0.9s, 1.2s) on the magnification coefficient, where s is given in Table I for bin 3. We thus decide not to include this effect in the subsequent modeling for all the redshift bins. Since we do not correct the effect of pixelization at the measurement level, we forward model the effect of pixelization by convolving the theory curves by two powers of the pixel window function for Nside = 1024 $(p_{w\ell})^2$ for the galaxy autospectra and $p_{w\ell}$ for the galaxy cross-spectra. To compare the theory predictions with the observed spectra, we further convolve the theory spectra with the bandpower window that captures the effect of the approximate mode decoupling applied to the data [34]. The final theory auto- and cross-spectra that we compare with the measurements are given by

$$C_{b}^{\text{th},gg} = \sum_{b'} \widehat{M^{-1}}_{bb'} \sum_{\ell'} w_{\ell'}^{b'} \left[\sum_{\ell'} (M_{\ell\ell'} (p_{w\ell'})^2 C_{\ell'}^{gg} + M_{0,\ell'} N_{\text{shot}}) \right],$$

$$C_{b}^{\text{th},\kappa g} = \sum_{b'} \widehat{M^{-1}}_{bb'} \sum_{\ell'} w_{\ell'}^{b'} \left[\sum_{\ell'} M_{\ell\ell'} p_{w\ell'} C_{\ell'}^{\kappa g} \right], \qquad (17)$$

where $\widehat{M^{-1}}_{bb'}$ is the inverse of the binned mode coupling matrix¹³ and w^b_{ℓ} are the uniform weights associated with

each multipole in bin *b*. The shot-noise level N_{shot} for the data is sampled with a prior centered at the inverse of the galaxy number density in the respective galaxy footprints.

We test the impact of photometric redshift uncertainties on our results in Appendix C showing that they do not affect the constraints on S_8 .

B. Blinding policy

We adopt a blinding policy that is intended to be a reasonable compromise between reducing the effects of confirmation bias and improving our ability to diagnose issues with the data and the pipeline. During the preparation of the analysis presented in this work, constraints on cosmological parameters were blinded until we demonstrated that a sequence of tests described below were passed. We were not blind to the measured spectra which, in the case of C_{ℓ}^{gg} , had already been present in [38]. We followed the procedure below before unblinding the ACT DR6 lensing and DESI galaxies cross-correlations analysis:

- (i) To verify our data are not contaminated by systematic effects, in particular, galactic and extragalactic foregrounds, we run a series of null tests described in Sec. VA. We also perform some tests for the galaxygalaxy autospectra in Sec. V B although the bulk of this work was already presented in [38]. We classify a test to be passing if it yields a PTE greater than 0.05. We qualify this stage to pass and that our mitigation strategies are sufficient if the number of failures is consistent with what is expected from random statistical fluctuations given the number of tests performed.
- (ii) Before unblinding the results using the ACT DR6 and *Planck* PR4 lensing maps, we freeze all the baseline analysis choices. These include the scale cuts used in C_{ℓ}^{gg} and C_{ℓ}^{kg} , the priors on the cosmological parameters and the nuisance parameters for the galaxies' linear biases and shot noise.

Note that, after passing the null tests and verifying the parameter recovery using simulations with our model, we perform a reanalysis of the *Planck* PR3 lensing map cross-correlated with the DESI galaxies discussed in Appendix B. We do not consider this to affect our blindness as this measurement has already been looked at in [38]. Using PR3 × DESI we investigate the consistency of the parameters recovered with different analysis choices, including different sky masks and consistency of parameters when using subsets of the data.

C. Likelihood and priors

We provide constraints on cosmological parameters by constructing a Gaussian likelihood

$$-2\ln\mathcal{L} \propto \sum_{bb'} \begin{bmatrix} \Delta \hat{C}_{b}^{gg}(\boldsymbol{\theta}) \\ \Delta \hat{C}_{b}^{\kappa g}(\boldsymbol{\theta}) \end{bmatrix} \mathbb{C}^{-1} \begin{bmatrix} \Delta \hat{C}_{b'}^{gg}(\boldsymbol{\theta}) \\ \Delta \hat{C}_{b'}^{\kappa g}(\boldsymbol{\theta}) \end{bmatrix}$$
(18)

¹²https://github.com/CLASS-SZ

¹³Obtained under the assumption that the power spectrum is piecewise constant and is the default approximation implemented in NaMaster.

where $\Delta \hat{C}_{b}^{gg}$ and $\Delta \hat{C}_{b}^{\kappa g}$ are the residuals between the binned observed galaxy-galaxy auto- and galaxy lensing crossspectra \hat{C}_{b}^{gg} and $\hat{C}_{b}^{\kappa g}$ and the respective window convolved theory spectra C_{b}^{gg} and $C_{b}^{\kappa g}$. The covariance matrix \mathbb{C} has the form

$$\mathbb{C} = \begin{bmatrix} \mathbb{C}_{bb'}^{gg,gg} & \mathbb{C}_{bb'}^{gg,kg} \\ (\mathbb{C}_{bb'}^{gg,kg})^T & \mathbb{C}_{bb'}^{kg,kg} \end{bmatrix}$$
(19)

where $\mathbb{C}_{bb'}^{gg,gg}$, $C_{bb'}^{\kappa_{g,\kappa_{g}}}$, and $\mathbb{C}_{bb'}^{gg,\kappa_{g}}$ are the galaxy-autospectrum covariance, the galaxy lensing cross-spectrum covariance, and the cross-covariance between them. These covariance blocks are estimated from simulations as described in Sec. IV C.

We use the Markov chain Monte Carlo (MCMC) code COBAYA [78] to perform the sampling and infer parameters from our galaxy-galaxy and galaxy-lensing data using the model described in Sec. VI evaluated with CLASS_SZ for a fast and accurate inference. We consider the chains to be converged when the Gelman-Rubin statistic [79,80] satisfies $R - 1 \le 0.01$ (which is reached within roughly one hour¹⁴).

The dataset used here is insensitive to the optical depth to reionization and, similar to [30], we fix this at the best-fit value of *Planck* [81]. Table IV shows the whole set of cosmological priors used, which largely follows from the priors assumed in the most recent *Planck* and ACT lensing analyses, except for slightly more restrictive A_s and H_0 priors. Similar also to the analysis of [34], we fix $\Omega_b h^2$ to the central value from *Planck* of $\Omega_b h^2 = 0.2242$ [82]. We also fix the tilt of the primordial power spectrum to $n_s = 0.9665$ [82] and assume the minimum neutrino mass allowed in the normal hierarchy ($\sum m_{\nu} = 0.06$ eV).¹⁵

For the shot-noise term, we choose a Gaussian prior on SN, the shot-noise amplitude, centered on the Poisson value with a width of 30%. We have checked that changing the width to 60% has no significant effect (at the 0.08σ level) on the mean of the posteriors. Notice that, due to our conservative binning scheme and hence the small number of data points, large values of shot noise could become degenerate with the signal amplitude, hence affecting the σ_8 values. This happens when the fitted shot noise is about 20 times that from the $1/N_{gal}$ expectation. Since these shotnoise values are unrealistic, we limit our shot noise prior to the above range, rather than using a wide uniform prior. The linear bias is sampled with a uniform distribution from 0 to 3.

TABLE IV. Priors used in the cosmological analysis of this work. Uniform priors are shown in square brackets and Gaussian priors with mean μ and standard deviation σ are denoted $\mathcal{N}(\mu, \sigma)$. Nuisance parameters are marginalized. The expected shot noise for each redshift bin is centered on the values listed in Table I.

Parameter	Prior
Cos	mological
$\ln(10^{10}A_s)$	[2.5, 3.5]
H_0	[50, 80]
n_s	0.9665
$\Omega_b h^2$	0.02242
$\Omega_c h^2$	[0.08, 0.20]
τ	0.055
$\sum m_{ u}$	0.06 eV
N	uisance
b_1	[0, 3]
b_2	[0, 3]
b_3	[0, 3]
<i>s</i> ₁	\mathcal{N} (Table I, 0.3)
<i>s</i> ₂	\mathcal{N} (Table I, 0.3)
<i>s</i> ₃	\mathcal{N} (Table I, 0.3)

Table II summarizes the main modeling differences between [38] and this work.

VII. RESULTS

A. DESI galaxies × ACT DR6

We jointly analyze the autocorrelation of the three DESI legacy galaxy samples and the cross-correlation of each with the ACT DR6 lensing reconstruction to obtain a 5.1% (68%) constraint on $S_8 \equiv \sigma_8 (\Omega_m / 0.3)^{0.5}$ of

$$S_8 = 0.772 \pm 0.040$$
 (DESI × ACT DR6). (20)

The posterior in the σ_8 - Ω_m plane is shown in the left panel of Fig. 12 (purple contours). The constraints from the individual redshift bins are summarized in Table V.

1. Combination of ACT DR6 and Planck

In Appendix B, we present a reanalysis of the 2 × 2 pt analysis with *Planck* PR3 CMB lensing and the four DESI legacy galaxy samples using our more conservative linear model, new pipeline, and previously neglected Monte Carlo lensing norm corrections. Figure 21 shows in detail how the above changes affected the parameter $S_8^{\times} \equiv \sigma_8 (\Omega_m/0.3)^{0.78}$ constrained by [38]. Below, we update the analysis results using our scale cuts and the new *Planck* PR4 lensing map. The σ_8 - Ω_m contours are shown on the right panel of Fig. 12. We find a 4.8% constraint on S_8 ,

¹⁴These analyses were run on a single node of the Niagara supercomputer, each node having 40 Intel Skylake/CascadeLake cores and 188 GB of RAM.

¹⁵We verified that when we enable those parameters fixed to the *Planck* value to vary with an uncertainty set by the *Planck* measurement only parameters degrade S_8 by ~10%.



FIG. 12. Cosmological constraints of the DESI Imaging galaxies. Left: the 1 and 2σ contours using C_{ℓ}^{gg} and C_{ℓ}^{kg} with ACT DR6 only. Right: the equivalent constraints for *Planck* PR4. The black solid line shows the best fit S_8 . As a reference, we also show as dashed lines of constant $S_8 = 0.85$ and $S_8 = 0.75$. Our measurement comfortably sits in between these two lines as can be seen as well with the best fit of $S_8 = 0.772$ and $S_8 = 0.783$ for ACT and *Planck*, respectively.

$$S_8 = 0.776 \pm 0.037$$
 (DESI × *Planck* PR4). (21)

The reconstruction noise in ACT is significantly lower than in *Planck*, and the higher resolution of ACT DR6 helps break the σ_8 - Ω_m degeneracy more effectively, producing less elongated contours (left panel of Fig. 12) than those obtained with *Planck* (right panel). However, because the *Planck* lensing map covers a larger fraction of the DESI footprint (33% vs 18%), it ultimately yields slightly tighter constraints on S_8 than the ACT-based analysis discussed in Sec. VII A.

The consistency between the ACT and *Planck* results motivates us to present the joint 2×2 pt analysis of the

TABLE V. Summary of the 1 σ constraints on cosmological parameters obtained from the 2 × 2 pt analysis with DESI legacy galaxies and ACT DR6 lensing reconstruction. We also present constraints from the joint analysis of *Planck* and ACT. BAO data is incorporated to break the degeneracy between the matter density Ω_m and the amplitude of fluctuations σ_8 as shown in the third and fourth blocks of the table.

	Ω_m	σ_8	S_8
	ACT DR6	× DESI legacy only	
z_1	0.207 ± 0.063	0.712 ± 0.125	0.579 ± 0.095
<i>z</i> ₂	0.246 ± 0.060	0.863 ± 0.107	0.768 ± 0.062
<i>z</i> ₃	0.324 ± 0.108	0.814 ± 0.142	0.812 ± 0.048
Joint	$0.240\substack{+0.014\\-0.046}$	$0.872\substack{+0.088\\-0.061}$	0.772 ± 0.040
	(ACT DR6 + Pla)	anck PR4) \times DESI legacy	
z_1	0.237 ± 0.067	0.714 ± 0.124	0.619 ± 0.077
z_2	0.247 ± 0.060	0.850 ± 0.111	0.756 ± 0.051
Z3	0.304 ± 0.080	0.824 ± 0.120	0.807 ± 0.039
Joint	$0.271\substack{+0.029\\-0.075}$	$0.821^{+0.11}_{-0.095}$	0.765 ± 0.032
	ACT DR6 ×	CDESI legacy + BAO	
z_1	0.290 ± 0.015	$0.644^{+0.041}_{-0.065}$	$0.633^{+0.053}_{-0.067}$
<i>z</i> ₂	0.309 ± 0.015	$0.717^{+0.046}_{-0.064}$	$0.727^{+0.054}_{-0.062}$
Z3	0.315 ± 0.015	$0.770^{+0.049}_{-0.056}$	0.788 ± 0.051
Joint	0.319 ± 0.013	$0.708_{-0.041}^{+0.031}$	0.731 ± 0.038
	(ACT DR6 + Planck	$x PR4) \times DESI legacy + BAO$	
z_1	0.291 ± 0.014	$0.645^{+0.037}_{-0.057}$	$0.635^{+0.047}_{-0.059}$
<i>z</i> ₂	0.310 ± 0.014	$0.714_{-0.053}^{+0.038}$	$0.726^{+0.043}_{-0.050}$
Z3	0.318 ± 0.014	$0.764_{-0.045}^{+0.040}$	0.785 ± 0.040
Joint	0.322 ± 0.011	$0.714\substack{+0.026\\-0.034}$	0.739 ± 0.029



FIG. 13. We show the S_8 (top) and $\sigma_8 1\sigma$ constraints when combined with BAO (bottom) using different data and analysis variations. Gold points correspond to constraints using the baseline analysis for the combination of ACT and *Planck* lensing (top) and ACT DR6 only (second row). Black data points show the joint ACT DR6 + *Planck* PR4 constraints cross-correlated with each redshift bin. We see that the mean of the first bin lies outside the 1σ interval of the baseline analysis although given that we have only bandpower for this redshift bin, it has a small weighting in the overall constraint. This can be seen from the gray point where removing the first redshift bin causes a ~0.45 σ shift with respect to the baseline in gold. Shown as reference in red are the S_8 and σ_8 values obtained from the *Planck* primary CMB.

DESI legacy galaxies with the ACT DR6 and *Planck* PR4 lensing. We describe in Sec. IV C the estimation of the joint covariance, accounting for the correlation between the ACT and *Planck* cross-correlations. Given that we find the bias preferred by the region of the *Planck* footprint is different to that of ACT in Fig. 4, we do not assume them to be identical in the ACT and *Planck* footprints. Hence we include C_{ℓ}^{gg}

measured using both the ACT and *Planck* masks in our analysis. We also include the significant cross-covariance between $C_{\ell}^{gg,ACT}$ and $C_{\ell}^{gg,Planck}$, which we also estimate from our Gaussian simulations.

The combination of ACT DR6 and *Planck* PR4 CMB lensing with DESI legacy results in a 4.2% joint constraint of

$$S_8 = 0.765 \pm 0.032$$
 (DESI × *Planck* PR4 + ACT DR6). (22)

This constitutes a 24% and a 13% improvement compared to ACT DR6 and *Planck* PR4 alone, respectively. The joint analysis in black is shown alongside the constraints from ACT and *Planck* in the top panel of Fig. 13. The best-constrained parameter in our analysis differs slightly from S_8 due to the redshifts and scales used and we determine this empirically to be closer to $S_8^{\times} \equiv \sigma_8 (\Omega_m/0.3)^{0.58}$, which we constrain to 4.1% $S_8^{\times} = 0.757 \pm 0.031$.

A feature that can be seen in Fig. 13 is that the lowest redshift bin is ~1.8 σ lower than the baseline constraint mean for both the ACT, *Planck*, and combined ACT + *Planck* results. This corresponds to a shift in S_8 at the 2.21 and 2.08 σ using the subset test proposed by [83] given that the constraints from z_1 are a subset of that of the joint analysis. We thus also proceed to report joint constraints of ACT + *Planck* × DESI excluding the first redshift bin

$$S_8 = 0.785 \pm 0.033. \tag{23}$$

This only degrades the baseline S_8 constraint by 3% but results in a shift of S_8 upward by 0.45 σ .

Overall, our 2 × 2 pt analysis is robust to the data and analysis choices used. A summary of the S_8 constraints using different analysis variations is shown in the top panel of Fig. 13 and we see that, excluding the analysis using only the first redshift bin, all the other choices are within the 1σ level of the baseline measurements. In green we show that we obtain consistent results when adding a prior on the angular size of the sound horizon $\theta_{\rm MC}$, that is predominantly sensitive to the combination of $\Omega_m h^3$ [84]. Taking the mean value¹⁶ measured by *Planck* [82] of $\Omega_m h^3 =$ 0.09635 we obtain a S_8 value of

$$S_8 = 0.747 \pm 0.028. \tag{24}$$

2. Combination with BAO

The tomographic lensing measurements provide information on a three-dimensional volume comprising the amplitude of matter fluctuations σ_8 , the matter density Ω_m , and the Hubble constant H_0 . To reduce the degeneracies of σ_8 with the other parameters and enable comparisons with other probes of weak lensing, CMB lensing autospectra, and CMB primary information, we include expansion information from the 6dF and SDSS surveys. The included data measure the BAO in the clustering of galaxies up to $z \approx 1$; the data are from 6dFGS [85], SDSS DR7 main galaxy sample [86], BOSS DR12 luminous red



FIG. 14. Distribution of MCMC samples for our constraints with the DESI legacy × ACT DR6 galaxy clustering and galaxy-CMB lensing cross-correlation in the $\sigma_8 - \Omega_m - H_0$ space in black. BAO samples are shown in orange, the intersection between the BAO and the lensing plane (red) provides constraint on σ_8 .

galaxies [87], and eBOSS DR16 LRGs [88]. Additionally, we include the higher-redshift ELGs [89], Lyman- α forest [90], and quasar samples [88] from eBOSS DR16. We do not include additional information that constrains structure growth from RSD. This choice allows us to isolate information on structure formation purely from the galaxy samples and its cross-correlation with CMB lensing. Figure 14 shows a breakdown of the constraints of the lensing-galaxy cross-correlation, BAO, and their intersection in the three-dimensional $\sigma_8 - H_0 - \Omega_m$ space.

The lower panel of Fig. 13 demonstrates again the robustness of the results against different analysis choices. Figure 15 shows the marginalized contours of Ω_m , σ_8 , and S_8 with BAO. With ACT DR6 we find a 5% constraint of

$$\sigma_8 = 0.708^{+0.031}_{-0.041}.$$
 (25)

Combining with *Planck* PR4 lensing again results in an improvement from the ACT DR6-only measurement by 16% giving

$$\sigma_8 = 0.714^{+0.026}_{-0.034}.$$
 (26)

The combined constraints alongside the individual ACT and *Planck* constraints are shown on the right of Fig. 15. We find that the σ_8 constraint after combining with BAO is 3.2σ lower than the CMB 2 pt value obtained from *Planck* PR4 NPIPE.

Similar to the 2×2 pt measurements (made without the inclusion of BAO), we find that the first redshift bin is

¹⁶This combination is determined to approximately 0.03% by *Planck* and so the uncertainty in this is subdominant to our measurement uncertainties.



FIG. 15. Parameter constraints from the cross-correlation of ACT DR6 lensing, PR4 lensing, their combination, and DESI galaxies (left). We break the degeneracy between Ω_m and σ_8 with additional information on Ω_m from BAO (right).

discrepant at the 2.0σ with respect to the third redshift bin. Excluding this first bin to ensure a more internally consistent dataset results in a slightly larger σ_8 value of

$$\sigma_8 = 0.735 \pm 0.035, \tag{27}$$

improving the consistency with the 2pt *Planck* PR4 NPIPE result to 2.1σ .

It is worth noting that by building a normalization flow distribution for the parameter shift between the BAO and the 2×2 pt dataset using TENSIOMETER [91], we find a discrepancy of 1.94σ between these two datasets. This test quantifies the discrepancy between the orange bands from BAO and the black bands from the ACT DR6 \times galaxy cross-correlation in the 3D space of $\sigma_8 - \Omega_m - H_0$ in Fig. 14. This discrepancy hence cautions the naive combination of the two datasets and the constraints obtained when including BAO. We additionally perform a range of systematic tests in Fig. 13 and find that a test with a larger $\ell_{\rm min} = 140$ causes the posterior mean of σ_8 to move up by 0.5σ in the direction that reduces the tension to 1.1σ . Although the error bars nearly double given that the most signal-dominated scales are discarded, this shift suggests that if large-scale systematics are the cause of the low σ_8 values obtained, they will primarily impact the lowest redshift bin. The scale cut essentially removes the constraint provided by the z_1 bin. As a result, the outcome aligns with the analysis where z_1 is simply discarded, but no scale cut is applied to the other bins. The above instabilities and internal tensions suggest further that



FIG. 16. Results of the measured tomographic structure growth at each effective redshift (orange data points) and the joint redshift measurement (solid orange line). The blacked dashed line corresponds to the predicted growth evolution assuming a *Planck* 2018 cosmology. In gray, we also plot $\alpha dN/dz$ of each galaxy sample, with α being an arbitrary constant for display purposes. We also show in silver for comparison the analysis from [35,36], the cross-correlation of DESI LRG and ACT DR6 and PR4 lensing, the filled gray circles are their constraints using linear theory and extends to higher redshifts than the galaxy samples used in our analysis but are nevertheless consistent with our analysis. The first open circle is the result they obtain using their baseline hybrid effective field theory model. In blue we show the results obtained from the cross-correlation of the blue and green unWISE sample, correlated with ACT DR6 lensing [34].



FIG. 17. Measurements of C_{ℓ}^{kg} (top row) and C_{ℓ}^{gg} (bottom row) for the three redshift samples of DESI legacy galaxies. Shown are also the best-fit lines where the cosmological parameters are fixed to that of *Planck* 2018 CMB anisotropies and only the bias and shot-noise parameters are allowed to vary.

investigation into large-scale systematics is needed before external datasets are added in. In what follows we will proceed with the more robust 2×2 pt results that provide a $\sim 4\%$ constraint on structure growth.

B. Consistency with predictions based on *Planck* CMB anisotropies

Our combined S_8 constraint is 1.9σ lower than the one obtained from *Planck* 2018 CMB anisotropies [92] and 1.9σ lower than preferred by the latest PR4 [17] + SRoll2 EE [93] analysis that is based on an extrapolation assuming the Λ CDM model.

We also assess the consistency of our dataset with the preferred parameters obtained from *Planck* 2018 by fixing the cosmological parameters to those reported by *Planck* 2018, namely with $H_0 = 67.4$, $\omega_c = 0.120$, and $\ln 10^{10}A_s = 3.044$ and comparing this with our baseline model. We find a $\Delta \chi^2 = 11.6$ ($\Delta \chi^2 = 12.4$ with the NPIPE parameters) going from the fixed to the free cosmology case. With a PTE of 0.04, the LCDM model provides a fairly reasonable fit to the data. In Fig. 16 we show the redshift evolution predicted by

Planck 2018 and the values measured by our baseline analysis. We rescale the S_8 at each redshift bin by $\sigma_8(z_{\text{eff}}^i)/\sigma_8(0)$. Apart from the constraint from the first fit bin, deviating from the *Planck* 2018 prediction by $\sim 2\sigma$, the joint constraint (orange line) and the constraints from redshift bins 2 and 3 are consistent with the cosmology preferred by *Planck*.

We show in Fig. 17 the best-fit obtained when fixing the cosmology parameters to the values preferred by *Planck* 2018 and varying only the nuisance shot-noise and bias parameters. Apart from the first redshift bin, our data show a good fit to the *Planck* 2018 cosmology.

C. Comparison with other measurements of large-scale structure

Figure 18 compares our results with other weak lensing surveys and the *Planck* CMB anisotropy results. We show in the Ω_m - σ_8 plane the constraints from cosmic shear measurements from DES-Y3 [94,95], KiDS-1000 [96], and ACT DR6 lensing [30]. Our measurements are consistent with the other weak lensing-only results with a similar degeneracy



FIG. 18. The Ω_m - σ_8 contours for this work (black) compared to other lensing-only results: DES-Y3 cosmic shear (blue), KiDS-1000 cosmic shear (green), and ACT DR6 lensing (red). The black, unfilled contour shows the constraints from the Planck PR4 CMB anisotropy measurements for comparison.

direction. On the same plot, we also show in unfilled black contours the *Planck* constraints from CMB anisotropy that we discussed in Sec. VII B.

We further discuss here comparisons of our results to measurements of structure growth from other large-scale structure observables. Figure 19 shows the compilation of these measurements in terms of S_8 constraints. We include results from other cross-correlations with CMB lensing from ACT, *Planck*, and SPT in purple. Galaxy weak lensing surveys (DES, KiDS, and HSC) are in blue and CMB lensing autospectrum analyses are in red.

We note that the posterior values of HSC, DES, and the CMB lensing autospectra measurements from ACT and *Planck* PR4 are analyzed with consistent prior choices to this work, while the other values are taken from the published results.¹⁷

We compare our results with measurements derived from cross-correlations of CMB lensing and galaxy positions, shown as purple points in Fig. 19. Our results are very consistent with the work in [35,36], which employed the same lensing map, but a different DESI galaxy sample containing only LRGs (thus one expects there to be significant overlap in the samples of galaxies used, in an analysis that extends to smaller scales using an EFT-based model). They report $S_8 = 0.775 \pm 0.021$, differing from

our fiducial analysis by 0.26σ . Our more aggressive scale cuts focusing on linear scales result in error bars that are 34% larger than that obtained from the DESI LRG sample. The linear analysis in [35,36] is comparable to the analysis presented here, albeit differences in galaxy sample and redshift distribution discussed in Sec. II A. In Fig. 16, we show a comparison of the constraints of S_8 as a function of redshift between our measurements and those from [34] in blue and [35,36] in gray. It is worth noting that our analysis extends to lower redshifts than those reported in the unWISE and LRG analyses and at the overlapping redshifts the analyses are in good agreement with each other. The linear analysis of [35,36] (filled gray circles) is in good agreement in terms of redshift and scales probes. This is not surprising since we use the same lensing maps and the scales analyzed are similar for the linear analysis; we expect this level of agreement between the two analyses in the absence of independent systematic effects.

Another similar EFT-based analysis using the unWISE sample found $S_8 = 0.810 \pm 0.015$; this analysis at higher redshifts of 0.6 and 1.1 for the blue and green sample is consistent with our fiducial result within 1.3σ . We find a similar 1.4 σ consistency with the measurement of the crosscorrelation of PR4 with Quaia quasars [97] with $S_8 = 0.841 \pm 0.044$. Our result is in good agreement (0.27σ) with the cross-correlation of the DES-Y3 magnitude-limited (MagLim) galaxies with the ACT DR4 lensing map [28]. Reference [37] measured the galaxy-galaxy lensing of the DESI BGS [98] and LRG targets with the DES-Y3 shear catalog. They found a lensing amplitude of $S_8 = 0.850^{+0.042}_{-0.050}$, consistent with the values from the primary CMB. Interestingly, they do not find the "lensing is low" tension to be redshift dependent, but rather could be due to mismodeling of galaxy bias or the deviation of the evolution of the intrinsic alignment signal from the usually assumed functional form.

Our results are also in good agreement with the measurements from the cross-correlation of CMB lensing and galaxy lensing. The first gray data point shows the 2 × 2 point analysis of the correlation between CMB lensing from *Planck* PR3 and SPT and DES-Y3 galaxy clustering δ_g and galaxy shear $\gamma \langle \delta_g \kappa \rangle + \langle \gamma \kappa \rangle$ [25] which yields $S_8 =$ $0.736^{+0.032}_{-0.028}$ (~0.7 σ). Reference [99] additionally includes correlations between galaxy positions and shear, as well as the autospectrum of the respective tracers, in the "5 × 2" point analysis yielding $S_8 = 0.773 \pm 0.016$ (~0.22 σ). Adding the CMB lensing autospectrum from *Planck* PR3 in the "6 × 2" point analysis [25] results in a S_8 of 0.792 ± 0.012 (0.82 σ).

As already shown in Fig. 18, we also find good agreement between our results and those obtained from galaxy lensing. The blue points in Fig. 19 showing the constraints from a reanalysis of KiDS-1000, DES-Y3, and HSC (Fourier and real space) data agree within 0.2σ with our baseline measurement in gold. Our work is more similar to

¹⁷Reference [31] showed that the impact of using priors matching those in the ACT DR6 lensing measurements is small for the galaxy survey datasets.



FIG. 19. Compilation of S_8 measurements from this work in gold. Other cross-correlation measurements (purple), measurements combining CMB lensing and galaxy lensing (gray), galaxy weak lensing in blue, CMB lensing in red, and extrapolation from CMB anisotropies (black). Gold shaded region consists of the 1σ uncertainty of our baseline ACT DR6 + *Planck* PR4 lensing × DESI legacy measured at 4% level.

the combination of galaxy-galaxy lensing and clustering 2×2 pt analysis, such as presented in [100], although with a different lensing kernel. Our results are again in good agreement with [100] using the DES-Y3 MagLim lens galaxies and shear, where $S_8 = 0.778^{+0.037}_{-0.031}$.

Finally, our results are consistent within 1.65, 1.76, 1.52, and 1.41σ to the DR6 + PR4, ACT DR6, *Planck* PR4, and SPT-3G results of the CMB lensing power spectrum, respectively, shown as the red points in Fig. 19. Comparison with the prediction from the primary CMB values are shown in black; refer to Sec. VII B for the detailed comparison with the cosmology preferred by the

Planck 2018 and NPIPE CMB anisotropies. Our measurement is also in 1.7σ agreement with the CMB anisotropies from ACT DR4 + WMAP.

VIII. DISCUSSIONS AND CONCLUSIONS

We have presented cosmological results from the 2×2 pt analysis of the DESI legacy survey galaxies with the ACT DR6 and *Planck* PR4 CMB lensing maps. In an analysis focused on linear scales and following a "blinding" procedure, we provide a 4.2% constraint on S_8 with the combined ACT + *Planck* correlation.

Our analysis passes a suite of null and systematic tests, ensuring the measurement is robust to extragalactic foreground biases and other galaxy systematic tests. The blinding procedure prevented us from inferring cosmological parameters before the null tests were passed, reducing the risk of confirmation bias. At the price of some reduction in SNR, we restricted the analysis to linear scales k < 0.17h Mpc⁻¹ where linearity of bias and of the matter power spectrum apply.

Our result for S_8 lies 1.9σ below the prediction from primary CMB results, constraining structure growth in the redshift range $0.3 \le z \le 0.8$. In Fig. 16, we exhibit the redshift dependence of S_8 by showing our constraint on this parameter within each redshift bin. While the joint constraint and analysis removing the lowest redshift bin are formally consistent with Planck CMB predictions, hints of suppression of structure growth at lower redshifts probed by the lowest redshift bins are emerging, consistent with other independent analyses (i.e., [35,36]). References [101,102] argued that the lower S_8 measured from weak lensing surveys relative to the *Planck* primary CMB estimate could be due to the small-scale modeling not fully capturing the nonlinearity and astrophysical effects such baryonic feedback, rather than a tension between the late and the early Universe. Our analysis, however, focuses on mostly linear scales, but still finds consistent S_8 values with existing galaxy weak lensing measurements. Although the significance of the S_8 "tension" is lower in our case, due to larger uncertainties, our results might hint at the possibility of a redshift-dependent effect affecting the large linear scales and not purely a modification affecting only the small-scale power spectrum (see also discussions in the literature about scale-independent suppression of growth [103]). If this suppression is indeed due to new physics, it becomes particularly intriguing because the lowest redshift bin lies within the dark energy-dominated regime. Any deviations from expected structure growth on these redshifts could potentially point toward new insights on the behavior of dark energy (i.e, [104]).

More statistical power and improved handling of the galaxy sample used for tomography will be needed before we can make concrete statements about the nature of this suppression on large scales-new physics, statistical fluctuation, or perhaps systematics affecting particularly the lowest redshifts. For example, we have not considered changes to the legacy survey galaxy sample compared to the previous analysis [38]. Despite the sample's robustness, there are several areas of potential improvement that we leave for future work, such as improved photometric redshift calibration using the newly available DESI spectra. Some recent work in this direction was carried out by Saraf et al. [105], who recalibrated the photo-z error distribution for the DESI legacy survey sample. Their cross-correlation results with Planck PR3 lensing map show overall a similar $1-3\sigma$ deviation below the theoretical expectation from the *Planck* 2018 cosmology in bins 0–2, although the highest tomographic bin 3 has a higher cross-correlation amplitude, closely consistent with the *Planck* prediction. Further efforts in understanding the exact calibration of the photo-*z* data in this catalog will clearly be of interest. Correlations of CMB lensing with low-redshift galaxy samples, such as done by Chen *et al.* [37] with the photometric DESI BGS targets, are also an important alternative avenue in verifying our findings regarding S_8 . In the future, the spectroscopic DESI BGS data will no doubt provide more insight into structure growth at low redshifts and its consistency with standard cosmology.

ACKNOWLEDGMENTS

The authors thank the review chairs, Martin White and Joe DeRose, for their thorough and insightful comments. F.J.Q. is grateful for Dr. Anil Seal and Trinity College Henry Barlow Scholarship for their support. The authors are honored to be permitted to conduct scientific research on Iolkam Du'ag (Kitt Peak), a mountain with particular significance to the Tohono O'odham Nation. F. J. Q. and B. D. S. acknowledge support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant Agreement No. 851274). Computations were performed on the Niagara supercomputer at the SciNet HPC Consortium. SciNet is funded by Innovation, Science and Economic Development Canada; the Digital Research Alliance of Canada; the Ontario Research Fund: Research Excellence; and the University of Toronto. C. S. acknowledges support from the Agencia Nacional de Investigación y Desarrollo (ANID) through Basal project FB210003. C. E. V. received the support of a fellowship from "la Caixa" Foundation (ID 100010434). The fellowship code is LCF/BQ/EU22/ 11930099. J. K. acknowledges support from NSF Grants No. AST-2307727 and No. AST-2153201. E. C. acknowledges support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant Agreement No. 849169). Support for ACT was through the U.S. National Science Foundation through Awards No. AST-0408698, No. AST-0965625, and No. AST-1440226 for the ACT project, as well as Awards No. PHY-0355328, No. PHY-0855887, and No. PHY-1214379. Funding was also provided by Princeton University, the University of Pennsylvania, and a Canada Foundation for Innovation (CFI) award to UBC. The development of multichroic detectors and lenses was supported by NASA Grants No. NNX13AE56G and No. NNX14AB58G. Detector research at N.I.S.T. was supported by the National Institute of Standards and Technology (NIST) Innovations in Measurement Science program. ACT operated in the Parque Astronómico Atacama in Northern Chile under the auspices of the Agencia Nacional de Investigación y Desarrollo (ANID). We thank the Republic of Chile for hosting ACT in the Northern Atacama. This material is based upon work supported by the U.S. Department of Energy (DOE), Office of Science, Office of High-Energy Physics, under Contract No. DE-AC02-05CH11231, and by the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility under the same contract. Additional support for DESI was provided by the U.S. National Science Foundation (NSF), Division of Astronomical Sciences under Contract No. AST-0950945 to the NSF's National Optical-Infrared Astronomy Research Laboratory; the Science and Technology Facilities Council of the United Kingdom; the Gordon and Betty Moore Foundation; the Heising-Simons Foundation; the French Alternative Energies and Atomic Energy Commission (CEA); the National Council of Humanities, Science and Technology of Mexico (CONAHCYT); the Ministry of Science, Innovation and Universities of Spain (MICIU/AEI/ 10.13039/501100011033), and by the DESI Member Institutions [106]. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the U.S. National Science Foundation, the U.S. Department of Energy, or any of the listed funding agencies.

DATA AVAILABILITY

The data that support the findings of this article are openly available [106].

APPENDIX A: STELLAR DENSITY CORRECTION

In this appendix, we show the galaxy density map correlation with the stellar density, before and after the correction. This correlation C_{ℓ}^{qS} is shown in Fig. 20 for bins 1–3. Given the DESI × ACT footprint, the impact of stellar density is relatively small, about 1% in the ℓ ranges adopted in the baseline analysis. The correction (see [38] for details) brings down the stellar density correlation by a further order of magnitude.

APPENDIX B: DESI GALAXIES × Planck PR3 LENSING REANALYSIS

In this appendix, we provide a reanalysis of the 2 × 2 pt of the *Planck* PR3 lensing and the DESI imaging galaxies as well as the detailed changes leading to the evolution from the original result in [38] to our baseline constraints. We use the priors listed in Table IV. Among the pipeline improvements we include the correction A_{ℓ}^{MC} for the lensing reconstruction misnormalization due to the presence of the mask to obtain an unbiased CMB lensing crosscorrelation as $C_{\ell}^{\hat{\kappa}^{\text{MC}}g} = A_{\ell}^{\text{MC}}C_{\ell}^{\hat{\kappa}g}$. This normalization correction A_{ℓ}^{MC} is computed with 480 Gaussian simulations and corresponding lensing reconstructions provided by [32].

Using the scale cuts discussed in Sec. III with $\ell_{\min}(z) = 50$, $\ell_{\max}^{cross}(z) = (170, 230, 290)$, and $\ell_{\max}^{auto}(z) = (170, 230, 290)$ we find $S_8 = 0.754 \pm 0.037$ using the full nonlinear power spectrum. The triangle plot is shown on Fig. 23.

Excluding the first redshift bin which only contains one bandpower and is prior dominated on the shot noise and linear bias shifts the constraints up to $S_8 = 0.776 \pm 0.043$. We find a minimum $\chi^2 = 5.6$ and $\chi^2 = 7.79$ for the joint sample and for the fit excluding the first galaxy bin, respectively. These correspond to a PTE of 0.94 and 0.65 for the 12 and 10 bandpowers used. The inclusion of BAO to break the degeneracy of σ_8 and Ω_m results in $\sigma_8 = 0.719 \pm$ 0.031 for the joint galaxy samples and $\sigma_8 = 0.73 \pm 0.03$ for the analysis excluding redshift bin 1. The S_8 and σ_8 whisker plots for the *Planck* PR3 analysis are shown in the left and right subpanels of Fig. 24, respectively.

The original paper [38] measures the combination $S_8^{\times} \sim \sigma_8 (\Omega_m/0.3)^{0.79} = 0.760 \pm 0.023$. This analysis used a pseudo- C_{ℓ} analysis with scales in the range of $10 \leq \ell \leq$ 500 and a theoretical covariance matrix with no off-diagonal covariance matrix. In terms of nonlinear modeling, [38] prescribes a different bias parameter to the linear and nonlinear regimes,



FIG. 20. The correlation between galaxy density maps and the normalized ALLWISE total stellar density C_{ℓ}^{gS} for the three tomographic bins with (blue) and without (orange) correction. We use the same measurement pipeline as the baseline C_{ℓ}^{gg} with the DESI × ACT footprint.



FIG. 21. We show systematically the impact on the changes implemented in this work on S_8^{\times} , smoothly connecting the values obtained in [38] to the constraints obtained in this work.

$$C_{\ell}^{gg} \sim b_{\text{linear}}^2 P_{\ell}^{mm,\text{linear}} + b_{\text{nonlinear}}^2 (P_{\ell}^{mm} - P_{\ell}^{mm,\text{linear}}), \quad (B1)$$

$$C_{\ell}^{\kappa g} \sim b_{\text{linear}} P_{\ell}^{mm,\text{linear}} + b_{\text{nonlinear}} (P_{\ell}^{mm} - P_{\ell}^{mm,\text{linear}}), \quad (B2)$$

where P^{mm} are computed using the CAMB power spectrum.

Since then, many improvements to the original analysis pipeline have been made, as described in the main text and it is instructive to demonstrate how the adopted changes alter the derived values of S_8^{\times} . A summary of these changes can be found in Fig. 21. Starting with the leftmost data point in black, we have the $S_8^{\times} = 0.760 \pm 0.023$ constrained by [38]:

- (i) keeping similar scale cuts to the original analysis with 50 ≤ ℓ ≤ 470, but with broader binning Δℓ = 60 instead of the original Δℓ = 10 to reduce bin to bin correlations, ΔS[×]₈ = +0.027;
- (ii) inclusion of a multiplicative transfer function on the *Planck* PR3 lensing map, $\Delta S_8^{\times} = +0.009$;
- (iii) restrict to linear scales, including discarding altogether the first redshift bin z_0 , $\Delta S_8^{\times} = -0.06$. This corresponds to the fifth gray data point in Fig. 21. The shift in constraint suggests that the model

employed in [38] might not be accurately describing the nonlinear scales; and

(iv) switching the model from the one prescribed in [38] to the baseline HALOFIT model used in this work results in a shift of $\Delta S_8^{\times} = +0.01$ ($S_8^{\times} = 0.733 \pm 0.054$). This small shift compared to the results using the model of [38] that accounts for nonlinear bias shows that within the scale cuts used we are insensitive to the details of the modeling of the nonlinear bias.

1. Testing the effects of nonlinear bias

We further verify that the scale cuts used in this work corresponding to spatial scales of $k_{\text{max}} = 0.15, 0.16, 0.17h/\text{Mpc}$ are insensitive to the effects on nonlinear bias.

First, as mentioned in the previous section, the small shift of $\Delta S_8^{\times} = +0.01$, when switching from the model of [38] to a constant bias model independent of scale, supports that the effect of nonlinear bias is small. We conduct an additional test where we analyze the ACT DR6 × DESI legacy data using more conservative scale cuts that discards the first redshift bin and limits the k_{max} of z_2 and z_3 to $k_{\text{max}} = 0.12, 0.13h/\text{Mpc}$, respectively. The constraint in S_8 shifts by 0.03σ as shown by Fig. 22. Hence, we conclude



FIG. 22. Cosmological constraints of the ACT DR6 × DESI legacy imaging data, using more conservative scale cuts for ℓ_{max} in blue compared to the baseline constraints in green.

that the scales chosen are robust to nonlinear bias modeling.

APPENDIX C: NULL BANDPOWERS FOR THE HIGHLIGHTED NULL TESTS IN THE MAIN TEXT

We show specifically the cross-correlation between the signal-nulled κ map obtained by differencing the 150 and 90 GHz temperature maps and the galaxy fields in Fig. 25, which exhibits a marginal PTE of 0.03 in the second redshift bin. While this bin shows a somewhat low PTE, there is no discernible pattern across the multipole range or redshifts that would suggest systematic effects. As discussed in the main text in Sec. V, an analogous test using both temperature and polarization data shows no such outlier in the null test targeting the foregrounds. Likewise, bandpower-level consistency tests (more sensitive to multiplicative biases) reveal no statistically significant deviations.

We also show in Fig. 26 the cross-correlation of the curl of the lensing field—sometimes called the "pseudo-" or "B-mode" lensing field—with the same galaxy samples, where one of the redshift bins has a marginal PTE of 0.04. Physically, we do not expect the curl to exhibit any measurable signal at current sensitivity levels, so any real correlation would be surprising. Again, since there is no coherent pattern across multipoles or redshifts, this result is consistent with a statistical fluctuation rather than a systematic effect.

APPENDIX D: SHIFTS IN REDSHIFT DISTRIBUTION

In this appendix, we show the impact of the photometric redshift uncertainty of the DESI legacy survey on the cosmological results. Figure 27 shows the posterior of the n(z) calibration nuisance parameters in [38]. The parameters f_0-f_3 scales the tail the redshift distribution, such that $a_i = f_i a_i^{\text{spec}}$ in Eq. (1), where a_i^{spec} is the best-fit value for the calibration sample, and $f_i \leq 1$ such that the tails are larger for the photometric sample. The parameters x_i are the shift parameters of the tomographic bin centers and satisfy $\sum_{i=0}^{4} x_i = 0$ such that mean redshift of the four tomographic bins are unchanged.

From the posteriors of these parameters, we identify the 3σ range of the shift parameters for bins 1–3: $\Delta z = 0.0025$,



FIG. 23. Cosmological constraints of the DESI imaging galaxies and Planck PR3 lensing.



FIG. 24. Whisker plots for the DESI imaging galaxies and *Planck* PR3 lensing. Left: S_8 constraints using C_{ℓ}^{gg} and C_{ℓ}^{kg} only. Right: σ_8 constraints when including BAO.



FIG. 25. Cross-correlation of the individual galaxy redshift bins with the temperature-only lensing null map obtained by performing lensing reconstruction on the difference between the 150 and the 90 GHz CMB maps.



FIG. 26. Cross-correlation of the individual galaxy redshift bins with the reconstructed lensing curl maps.



FIG. 27. The posteriors of the redshift distribution calibration nuisance parameters in [38] using the cross-correlation of galaxies in different tomographic bins with the *Planck* 2018 cosmology.



FIG. 28. The change in cosmological parameter constraints by shifting the mean redshifts of the three tomographic bins. The green contours and curves show the baseline scenario, whereas blue and black show cases where the mean redshift of all bins being shifted downward and upward, by $\Delta z = 0.0025$, 0.005, 0.01 for z1 - z3, respectively.

0.005, 0.01. We test scenarios where the mean redshifts of the three bins are simultaneously shifted upward, i.e., $z \rightarrow z + \Delta z$, and downward, i.e., $z \rightarrow z - \Delta z$. Figure 28 shows the change in cosmological parameter constraints for these two cases, compared to the baseline. We find the shifts are $<0.2\sigma$.

- [1] L. Wenzl, R. Bean, S.-F. Chen, G. S. Farren, M. S. Madhavacheril, G. A. Marques, F. J. Qu, N. Sehgal, B. D. Sherwin, and A. van Engelen, Constraining gravity with a new precision E_G estimator using Planck + SDSS BOSS data, Phys. Rev. D **109**, 083540 (2024).
- [2] N. Sailer, E. Castorina, S. Ferraro, and M. White, Cosmology at high redshift—a probe of fundamental physics, J. Cosmol. Astropart. Phys. 12 (2021) 049.
- [3] J. E. Carlstrom, G. P. Holder, and E. D. Reese, Cosmology with the Sunyaev-Zel'dovich effect, Annu. Rev. Astron. Astrophys. 40, 643 (2002).
- [4] B. Horowitz and U. Seljak, Cosmological constraints from thermal Sunyaev–Zeldovich power spectrum revisited, Mon. Not. R. Astron. Soc. 469, 394 (2017).

- [5] B. Bolliet, Cosmological constraints from the thermal Sunyaev Zeldovich power spectrum?, arXiv:1806.04786.
- [6] M. M. Ivanov, M. Simonović, and M. Zaldarriaga, Cosmological parameters from the boss galaxy power spectrum, J. Cosmol. Astropart. Phys. 05 (2020) 042.
- [7] G. d'Amico, J. Gleyzes, N. Kokron, K. Markovic, L. Senatore, P. Zhang, F. Beutler, and H. Gil-Marín, The cosmological analysis of the SDSS/BOSS data from the effective field theory of large-scale structure, J. Cosmol. Astropart. Phys. 05 (2020) 005.
- [8] T. Colas, G. d'Amico, L. Senatore, P. Zhang, and F. Beutler, Efficient cosmological analysis of the SDSS/ BOSS data from the effective field theory of large-scale structure, J. Cosmol. Astropart. Phys. 06 (2020) 001.

- [9] P. A. R. Ade *et al.*, Planck2015 results: XXIV. Cosmology from Sunyaev-Zeldovich cluster counts, Astron. Astrophys. **594**, A24 (2016).
- [10] T. d. Haan *et al.*, Cosmological constraints from galaxy clusters in the 2500 square-degree SPT-SZ survey, Astrophys. J. 832, 95 (2016).
- [11] T. Abbott *et al.*, Dark energy survey year 1 results: Cosmological constraints from cluster abundances and weak lensing, Phys. Rev. D **102**, 023509 (2020).
- [12] A. N. Salcedo, H.-Y. Wu, E. Rozo, D. H. Weinberg, C.-H. To, T. Sunayama, and A. Lee, Dark energy survey year 1 clusters are consistent with Planck, Phys. Rev. Lett. 133, 221002 (2024).
- [13] S. Bocquet *et al.*, SPT clusters with DES and HST weak lensing. II. Cosmological constraints from the abundance of massive halos, Phys. Rev. D **110**, 083510 (2024).
- [14] C. Howlett, K. Said, J. R. Lucey, M. Colless, F. Qin, Y. Lai, R. B. Tully, and T. M. Davis, The Sloan digital sky survey peculiar velocity catalogue, Mon. Not. R. Astron. Soc. 515, 953 (2022).
- [15] C. Saulder *et al.*, Target selection for the DESI peculiar velocity survey, Mon. Not. R. Astron. Soc. **525**, 1106 (2023).
- [16] B. E. Stahl, T. de Jaeger, S. S. Boruah, W. Zheng, A. V. Filippenko, and M. J. Hudson, Peculiar-velocity cosmology with Types Ia and II supernovae, Mon. Not. R. Astron. Soc. 505, 2349 (2021).
- [17] E. Rosenberg, S. Gratton, and G. Efstathiou, CMB power spectra and cosmological parameters from Planck PR4 with CamSpec, Mon. Not. R. Astron. Soc. 517, 4620 (2022).
- [18] T. M. C. Abbott *et al.* (DES Collaboration), Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing, Phys. Rev. D 105, 023520 (2022).
- [19] C. Heymans *et al.*, KiDS-1000 Cosmology: Multiprobe weak gravitational lensing and spectroscopic galaxy clustering constraints, Astron. Astrophys. **646**, A140 (2021).
- [20] S. Sugiyama *et al.*, Hyper Suprime-Cam Year 3 results: Cosmology from galaxy clustering and weak lensing with HSC and SDSS using the minimal bias model, Phys. Rev. D **108**, 123521 (2023).
- [21] T. Abbott *et al.*, DES Y3 + KiDS-1000: Consistent cosmology combining cosmic shear surveys, Open J. Astrophys. 6, 36 (2023).
- [22] M. White *et al.*, Cosmological constraints from the tomographic cross-correlation of DESI Luminous Red Galaxies and Planck CMB lensing, J. Cosmol. Astropart. Phys. 02 (2022) 007.
- [23] S. Singh, R. Mandelbaum, U. Seljak, S. Rodríguez-Torres, and A. Slosar, Cosmological constraints from galaxy–lensing cross-correlations using BOSS galaxies with SDSS and CMB lensing, Mon. Not. R. Astron. Soc. 491, 51 (2020).
- [24] S.-F. Chen, M. White, J. DeRose, and N. Kokron, Cosmological analysis of three-dimensional boss galaxy clustering and Planck CMB lensing cross correlations via

Lagrangian perturbation theory, J. Cosmol. Astropart. Phys. 07 (2022) 041.

- [25] C. Chang *et al.* (DES and SPT Collaborations), Joint analysis of dark energy survey year 3 data and CMB lensing from SPT and Planck. II. Cross-correlation measurements and cosmological constraints, Phys. Rev. D 107, 023530 (2023).
- [26] T. M. C. Abbott *et al.* (DES and SPT Collaborations), Joint analysis of dark energy survey year 3 data and CMB lensing from SPT and Planck. III. Combined cosmological constraints, Phys. Rev. D **107**, 023531 (2023).
- [27] N. C. Robertson *et al.*, Strong detection of the CMB lensing and galaxy weak lensing cross-correlation from ACT-DR4, PlanckLegacy, and KiDS-1000, Astron. Astrophys. **649**, A146 (2021).
- [28] G. A. Marques *et al.* (ACT and DES Collaborations), Cosmological constraints from the tomography of DES-Y3 galaxies with CMB lensing from ACT DR4, J. Cosmol. Astropart. Phys. 01 (2024) 033.
- [29] S. Shaikh *et al.* (ACT and DES Collaborations), Cosmology from cross-correlation of ACT-DR4 CMB lensing and DES-Y3 cosmic shear, Mon. Not. R. Astron. Soc. **528**, 2112 (2024).
- [30] F. J. Qu *et al.*, The Atacama Cosmology Telescope: A measurement of the DR6 CMB lensing power spectrum and its implications for structure growth, Astrophys. J. 962, 112 (2024).
- [31] M. S. Madhavacheril *et al.* (ACT Collaboration), The Atacama Cosmology Telescope: DR6 gravitational lensing map and cosmological parameters, Astrophys. J. 962, 113 (2024).
- [32] J. Carron, M. Mirmelstein, and A. Lewis, CMB lensing from Planck PR4 maps, J. Cosmol. Astropart. Phys. 09 (2022) 039.
- [33] Z. Pan *et al.*, Measurement of gravitational lensing of the cosmic microwave background using SPT-3G 2018 data, Phys. Rev. D 108, 122005 (2023).
- [34] G. S. Farren *et al.*, The Atacama Cosmology Telescope: Cosmology from cross-correlations of unWISE galaxies and ACT DR6 CMB lensing, Astrophys. J. 966, 157 (2024).
- [35] N. Sailer *et al.*, Cosmological constraints from the crosscorrelation of DESI luminous red galaxies with CMB lensing from Planck PR4 and ACT DR6, arXiv:2407.04607.
- [36] J. Kim *et al.*, The Atacama Cosmology Telescope DR6 and DESI: Structure formation over cosmic time with a measurement of the cross-correlation of CMB lensing and luminous red galaxies, J. Cosmol. Astropart. Phys. 12 (2024) 022.
- [37] S. Chen *et al.*, Not all lensing is low: An analysis of DESI × DES using the Lagrangian effective theory of LSS, Phys. Rev. D **110**, 103518 (2024).
- [38] Q. Hang, S. Alam, J. A. Peacock, and Y.-C. Cai, Galaxy clustering in the DESI legacy survey and its imprint on the CMB, Mon. Not. R. Astron. Soc. 501, 1481 (2021).
- [39] A. Dey *et al.*, Overview of the DESI legacy imaging surveys, Astron. J. **157**, 168 (2019).
- [40] B. Flaugher *et al.* (DES Collaboration), The dark energy camera, Astron. J. **150**, 150 (2015).

- [41] The Dark Energy Survey Collaboration, The dark energy survey, arXiv:astro-ph/0510346.
- [42] A. Dey *et al.*, Mosaic3: A red-sensitive upgrade for the prime focus camera at the Mayall 4m telescope, in *Ground-Based and Airborne Instrumentation for Astronomy VI*, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 9908 (SPIE, Bellingham, Washington, 2016), p. 99082C.
- [43] G. G. Williams, E. Olszewski, M. P. Lesser, and J. H. Burge, 90prime: A prime focus imager for the Steward Observatory 90-in. telescope, in *Ground-based Instrumentation for Astronomy*, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 5492, edited by A. F. M. Moorwood and M. Iye (SPIE, Bellingham, Washington, 2004), pp. 787–798.
- [44] E. Wright *et al.*, The Wide-field Infrared Survey Explorer (WISE): Mission Description and Initial On-o rbit Performance, Astron. J. **140**, 1868 (2010).
- [45] E. F. Schlafly, A. M. Meisner, and G. M. Green, The unwise catalog: Two billion infrared sources from five years of wise imaging, Astrophys. J. Suppl. Ser. 240, 30 (2019).
- [46] J. Liske *et al.*, Galaxy and mass assembly (GAMA): End of survey report and data release 2, Mon. Not. R. Astron. Soc. 452, 2087 (2015).
- [47] S. Alam *et al.*, The eleventh and twelfth data releases of the sloan digital sky survey: Final data from SDSS-III, Astrophys. J. Suppl. Ser. **219**, 12 (2015).
- [48] R. Ahumada *et al.*, The 16th data release of the sloan digital sky surveys: First release from the APOGEE-2 southern survey and full release of eBOSS spectra, Astrophys. J. Suppl. Ser. **249**, 3 (2020).
- [49] M. Scodeggio *et al.*, The VIMOS public extragalactic redshift survey (VIPERS). Full spectroscopic data and auxiliary information release (PDR-2), Astron. Astrophys. **609**, A84 (2018).
- [50] J. A. Newman *et al.*, The DEEP2 Galaxy Redshift Survey: Design, observations, data reduction, and redshifts, Astrophys. J. Suppl. Ser. **208**, 5 (2013).
- [51] O. Ilbert *et al.*, Cosmos photometric redshifts with 30bands for 2-deg², Astrophys. J. 690, 1236 (2009).
- [52] R. Cawthon *et al.* (DES Collaboration), Dark Energy Survey Year 1 results: Calibration of redMaGiC redshift distributions in DES and SDSS from cross-correlations, Mon. Not. R. Astron. Soc. **481**, 2427 (2018).
- [53] R. Zhou *et al.*, Preliminary target selection for the DESI luminous red galaxy (LRG) sample, Res. Not. AAS 4, 181 (2020).
- [54] K. M. Górski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke, and M. Bartelmann, HEALPix: A framework for high-resolution discretization and fast analysis of data distributed on the sphere, Astrophys. J. 622, 759 (2005).
- [55] G. Green, dustmaps: A Python interface for maps of interstellar dust, J. Open Source Softwaare 3, 695 (2018).
- [56] S. Naess and T. Louis, Large-scale power loss in ground-based CMB mapmaking, Open J. Astrophys. 6, 21 (2023).

- [57] N. Sailer, E. Schaan, and S. Ferraro, Lower bias, lower noise CMB lensing with foreground-hardened estimators, Phys. Rev. D 102, 063517 (2020).
- [58] N. Sailer, S. Ferraro, and E. Schaan, Foreground-immune CMB lensing reconstruction with polarization, Phys. Rev. D 107, 023504 (2023).
- [59] E. Hivon, K. M. Górski, C. B. Netterfield, B. P. Crill, S. Prunet, and F. Hansen, MASTER of the cosmic microwave background anisotropy power spectrum: A fast method for statistical analysis of large and complex cosmic microwave background data sets, Astrophys. J. 567, 2 (2002).
- [60] D. Alonso, J. Sanchez, and A. Slosar, A unified pseudocell framework, Mon. Not. R. Astron. Soc. 484, 4127 (2019).
- [61] R. Adam, P. A. R. Ade, N. Aghanim, M. I. R. Alves, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday *et al.* (Planck Collaboration), Planck 2015 results. X. Diffuse component separation: Foreground maps, Astron. Astrophys. **594**, A10 (2016).
- [62] A. J. Mead, J. A. Peacock, C. Heymans, S. Joudaki, and A. F. Heavens, An accurate halo model for fitting non-linear cosmological power spectra and baryonic feedback models, Mon. Not. R. Astron. Soc. 454, 1958 (2015).
- [63] Z. Atkins *et al.*, The Atacama Cosmology Telescope: Mapbased noise simulations for DR6, J. Cosmol. Astropart. Phys. 11 (2023) 073.
- [64] A. Baleato Lizancos and M. White, Harmonic analysis of discrete tracers of large-scale structure, J. Cosmol. Astropart. Phys. 05 (2024) 010.
- [65] C. García-García, D. Alonso, and E. Bellini, Disconnected pseudo-cell covariances for projected large-scale structure data, J. Cosmol. Astropart. Phys. 11 (2019) 043.
- [66] J. Hartlap, P. Simon, and P. Schneider, Why your model parameter confidences might be too optimistic. Unbiased estimation of the inverse covariance matrix, Astron. Astrophys. 464, 399 (2006).
- [67] N. MacCrann *et al.*, The Atacama Cosmology Telescope: Mitigating the impact of extragalactic foregrounds for the DR6 cosmic microwave background lensing analysis, Astrophys. J. 966, 138 (2024).
- [68] G. Stein, M. A. Alvarez, J. R. Bond, A. v. Engelen, and N. Battaglia, The websky extragalactic CMB simulations, J. Cosmol. Astropart. Phys. 10 (2020) 012.
- [69] F. J. Qu, M. Millea, and E. Schaan, Impact and mitigation of polarized extragalactic foregrounds on Bayesian cosmic microwave background lensing, Phys. Rev. D 111, 023503 (2025).
- [70] D. N. Limber, The analysis of counts of the extragalactic Nebulae in terms of a fluctuating density field., Astrophys. J. 117, 134 (1953).
- [71] M. LoVerde and N. Afshordi, Extended Limber approximation, Phys. Rev. D 78, 123506 (2008).
- [72] B. Bolliet, B. Comis, E. Komatsu, and J. F. Macías-Pérez, Dark energy constraints from the thermal Sunyaev– Zeldovich power spectrum, Mon. Not. R. Astron. Soc. 477, 4957 (2018).

- [73] B. Bolliet *et al.*, CLASS_SZ I: Overview, EPJ Web Conf. 293, 00008 (2024).
- [74] J. Lesgourgues, The cosmic linear anisotropy solving system (CLASS) I: Overview, arXiv:1104.2932.
- [75] D. Blas, J. Lesgourgues, and T. Tram, The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes, J. Cosmol. Astropart. Phys. 07 (2011) 034.
- [76] B. Bolliet, A. Spurio Mancini, J. C. Hill, M. Madhavacheril, H. T. Jense, E. Calabrese, and J. Dunkley, High-accuracy emulators for observables in Λ CDM, N_{eff}, Σm_{ν} , and w cosmologies, Mon. Not. R. Astron. Soc. **531**, 1351 (2024).
- [77] A. Spurio Mancini, D. Piras, J. Alsing, B. Joachimi, and M. P. Hobson, CosmoPower: emulating cosmological power spectra for accelerated Bayesian inference from nextgeneration surveys, Mon. Not. R. Astron. Soc. 511, 1771 (2022).
- [78] J. Torrado and A. Lewis, Cobaya: Code for bayesian analysis of hierarchical physical models, J. Cosmol. Astropart. Phys. 05 (2021) 057.
- [79] A. Gelman and D. B. Rubin, Inference from iterative simulation using multiple sequences, Stat. Sci. 7, 457 (1992).
- [80] S. P. Brooks and A. Gelman, General methods for monitoring convergence of iterative simulations, J. Comput. Graph. Stat. 7, 434 (1998).
- [81] P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett *et al.* (Planck Collaboration), Planck 2015 results. XV. Gravitational lensing, Astron. Astrophys. **594**, A15 (2016).
- [82] N. Aghanim *et al.* (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641, A6 (2020).
- [83] S. Gratton and A. Challinor, Understanding parameter differences between analyses employing nested data subsets, Mon. Not. R. Astron. Soc. 499, 3410 (2020).
- [84] W. J. Percival *et al.*, Parameter constraints for flat cosmologies from cosmic microwave background and 2dFGRS power spectra, Mon. Not. R. Astron. Soc. 337, 1068 (2002).
- [85] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, L. Campbell, Q. Parker, W. Saunders, and F. Watson, The 6dF galaxy survey: Baryon acoustic oscillations and the local Hubble constant, Mon. Not. R. Astron. Soc. 416, 3017 (2011).
- [86] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden, and M. Manera, The clustering of the SDSS DR7 main galaxy sample—I. A 4 per cent distance measure at z = 0.15, Mon. Not. R. Astron. Soc. **449**, 835 (2015).
- [87] S. Alam *et al.*, The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: Cosmological analysis of the DR12 galaxy sample, Mon. Not. R. Astron. Soc. **470**, 2617 (2017).
- [88] S. Alam *et al.*, Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at

the Apache Point Observatory, Phys. Rev. D **103**, 083533 (2021).

- [89] J. Comparat *et al.*, SDSS-IV eBOSS emission-line galaxy pilot survey, Astron. Astrophys. **592**, A121 (2016).
- [90] H. du Mas des Bourboux *et al.*, The completed SDSS-IV extended baryon oscillation spectroscopic survey: Baryon acoustic oscillations with Ly α Forests, Astrophys. J. **901**, 153 (2020).
- [91] M. Raveri and C. Doux, Non-Gaussian estimates of tensions in cosmological parameters, Phys. Rev. D 104, 043504 (2021).
- [92] N. Aghanim *et al.* (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641, A6 (2020); 652, C4(E) (2021).
- [93] L. Pagano, J.-M. Delouis, S. Mottet, J.-L. Puget, and L. Vibert, Reionization optical depth determination from Planck HFI data with ten percent accuracy, Astron. Astrophys. 635, A99 (2020).
- [94] A. Amon *et al.* (DES Collaboration), Dark Energy Survey Year 3 results: Cosmology from cosmic shear and robustness to data calibration, Phys. Rev. D 105, 023514 (2022).
- [95] L. F. Secco *et al.* (DES Collaboration), Dark Energy Survey Year 3 results: Cosmology from cosmic shear and robustness to modeling uncertainty, Phys. Rev. D 105, 023515 (2022).
- [96] M. Asgari *et al.*, KiDS-1000 cosmology: Cosmic shear constraints and comparison between two point statistics, Astron. Astrophys. **645**, A104 (2021).
- [97] G. Piccirilli, G. Fabbian, D. Alonso, K. Storey-Fisher, J. Carron, A. Lewis, and C. García-García, Growth history and quasar bias evolution at z < 3 from quaia, J. Cosmol. Astropart. Phys. 06 (2024) 012.
- [98] C. Hahn *et al.*, The DESI bright galaxy survey: Final target selection, design, and validation, Astron. J. 165, 253 (2023).
- [99] T. Abbott *et al.*, Joint analysis of dark energy survey year 3 data and CMB lensing from SPT and Planck. III. Combined cosmological constraints, Phys. Rev. D 107, 023531 (2023).
- [100] A. Porredon *et al.* (DES Collaboration), Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and galaxy-galaxy lensing using the MAGLIM lens sample, Phys. Rev. D **106**, 103530 (2022).
- [101] A. Amon and G. Efstathiou, A non-linear solution to the S_8 tension?, Mon. Not. R. Astron. Soc. **516**, 5355 (2022).
- [102] C. Preston, A. Amon, and G. Efstathiou, A non-linear solution to the S_8 tension—II. Analysis of DES Year 3 cosmic shear, Mon. Not. R. Astron. Soc. **525**, 5554 (2023).
- [103] N.-M. Nguyen, D. Huterer, and Y. Wen, Evidence for suppression of structure growth in the concordance cosmological model, Phys. Rev. Lett. 131, 111001 (2023).
- [104] A. G. Adame *et al.* (DESI Collaboration), DESI 2024 VI: Cosmological constraints from the measurements of baryon acoustic oscillations, J. Cosmol. Astropart. Phys. 02 (2025) 021.

- [105] C. S. Saraf, P. Bielewicz, and M. Chodorowski, Effect of redshift bin mismatch on cross correlation between DESI legacy imaging survey and Planck CMB lensing potential, Astron. Astrophys. 690, A338 (2024).
- [106] https://www.desi.lbl.gov/collaborating-institutions
- [107] F. J. Qu and Q. Hang, The Atacama Cosmology Telescope DR6 and DESI: Structure growth measurements from the cross- correlation of DESI legacy imaging galaxies and CMB lensing from ACT DR6 and Planck PR4 (2024), 10.5281/zenodo.13844390.