



# Lung volumes, gas transfer and oscillometry after preterm birth: systematic review and meta-analysis

James T.D. Gibbons <sup>1,2,3</sup>, Michael L. Beaven <sup>1,3</sup>, Christopher W. Course <sup>4</sup>, Sarah J. Kotecha<sup>4</sup>, Thomas Hixson<sup>5</sup>, Melissa Zuidersma<sup>2</sup>, Andrew C. Wilson<sup>1,2</sup>, Sailesh Kotecha <sup>4</sup> and Shannon J. Simpson<sup>1,3</sup>

<sup>1</sup>Children's Lung Health, Wal-yan Respiratory Research Centre, The Kids Research Institute of Australia, Perth, Australia. <sup>2</sup>Department of Respiratory and Sleep Medicine, Perth Children's Hospital, Perth, Australia. <sup>3</sup>Curtin School of Allied Health, Curtin University, Perth, Australia. <sup>4</sup>Department of Child Health, Cardiff University School of Medicine, Cardiff, UK. <sup>5</sup>Regional Neonatal Intensive Care Unit, University Hospital of Wales, Cardiff, UK.

Corresponding author: Shannon J. Simpson ([shannon.simpson@thekids.org.au](mailto:shannon.simpson@thekids.org.au))



Shareable abstract (@ERSpublications)

**Substantial physiological abnormalities persist following preterm birth when lung function is measured using static lung volumes, gas transfer and oscillometry.** <https://bit.ly/4kl67fU>

**Cite this article as:** Gibbons JTD, Beaven ML, Course CW, *et al.* Lung volumes, gas transfer and oscillometry after preterm birth: systematic review and meta-analysis. *Eur Respir Rev* 2025; 34: 240151 [DOI: 10.1183/16000617.0151-2024].

Copyright ©The authors 2025

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact [permissions@ersnet.org](mailto:permissions@ersnet.org)

Received: 18 July 2024  
Accepted: 11 Feb 2025

## Abstract

**Background** Small airway and lung parenchymal abnormalities frequently occur following preterm birth but are commonly missed by spirometry. Static lung volumes, diffusing capacity of the lung for carbon monoxide ( $D_{LCO}$ ) and oscillometry provide a more precise characterisation of these conditions. We hypothesised that differences in these measures exist between individuals born preterm and at term and we aimed to systematically review the literature to identify and quantify these differences in lung function.

**Methods** This systematic review and meta-analysis, registered with PROSPERO (CRD42022320775) and guided by Preferred Reporting Items for Systematic Reviews and Meta-Analyses standards, searched six databases up to 29 November 2024. We included studies comparing lung function between preterm subjects and term controls *via* static lung volumes, gas transfer or oscillometry. Differences in lung function were analysed using random-effects meta-analysis to compute the standardised mean difference (SMD).

**Results** From 12 143 titles, we analysed 52 cohorts with static lung volumes, 37 with gas transfer and 18 with oscillometry data. While total lung capacity was similar between preterm and term cohorts (SMD  $-0.08$ , 95% CI  $-0.17$  to  $0.004$ ), preterm participants showed increased residual volume (SMD  $0.32$ , 95% CI  $0.19$  to  $0.44$ ) and residual volume/total lung capacity (SMD  $0.45$ , 95% CI  $0.28$  to  $0.63$ ).  $D_{LCO}$  was lower in preterm cohorts (SMD  $-0.51$ , 95% CI  $-0.64$  to  $-0.38$ ). Preterm cohorts also demonstrated increased airway resistance at 5/6 Hz (SMD  $0.44$ , 95% CI  $0.22$  to  $0.67$ ), difference between airway resistance at 5/6 Hz and 20 Hz (SMD  $0.51$ , 95% CI  $0.07$  to  $0.96$ ), resonant frequency (SMD  $0.63$ , 95% CI  $0.12$  to  $1.15$ ) and area under the reactance curve (SMD  $0.62$ , 95% CI  $0.35$  to  $0.88$ ).

**Interpretation** We demonstrate that preterm birth is linked to notable abnormalities in static lung volumes, gas transfer and oscillometry, underscoring the necessity of employing comprehensive pulmonary function tests beyond spirometry to monitor and address long-term respiratory outcomes effectively.

## Introduction

Preterm birth is associated with enduring respiratory consequences [1]. Life-sustaining interventions such as mechanical ventilation and supplemental oxygen use can injure the developing lung, while premature exposure to the extrauterine environment disrupts normal alveolar development [2]. Particularly vulnerable are those born at very (<32 weeks') or extremely (<28 weeks') early gestational ages, and those diagnosed with bronchopulmonary dysplasia (BPD) during infancy. However, prematurity-associated lung disease (also referred to as post-prematurity respiratory disease) is not isolated to this group, with even those born moderate-late preterm demonstrating abnormalities in lung health [3]. Increased rates of wheezing, pulmonary hypertension and respiratory-related hospitalisation are well documented in this population [4–6]. Recent studies have also highlighted preterm birth as a significant risk factor for early-onset COPD [7].



Despite preterm birth representing possibly the earliest origin of chronic respiratory disease, it remains unclear how best to monitor this group using lung function techniques commonly available in clinical practice [8].

Our recent systematic reviews demonstrated persistent lung function deficits measured by spirometry in preterm-born individuals, with decreases in forced expiratory volume in 1 s ( $FEV_1$ ), forced vital capacity (FVC) and  $FEV_1/FVC$  ratios when compared to term-born populations [9, 10]. Additionally, meta-regression analyses support previous findings of longitudinal cohort studies showing a decline in  $FEV_1/FVC$  with increasing age. Together, these studies suggest increasing airway obstruction throughout life, particularly in those with an infant diagnosis of BPD [11, 12]. Spirometry, however, has inherent limitations in elucidating the intricate changes that preterm birth inflicts on lung parenchyma, the pulmonary vascular bed and the peripheral airways. Enlarged, simplified, thick-walled alveoli with dysmorphic growth of the pulmonary vascular bed are consistently reported in histological studies [2, 13], while gas trapping, often suggestive of small airway disease, has also been reported on imaging studies [14].

Alternative pulmonary function tests may offer a more granular perspective of the pathophysiological changes occurring in the lungs following preterm birth and allow for earlier diagnosis of ongoing abnormalities. Total body plethysmography and inert gas washout can assess static lung volumes, offering insights into restrictive lung disease, gas trapping and ventilation inhomogeneity [15]. The diffusing capacity of the lungs for carbon monoxide ( $D_{LCO}$ ) test assesses gas transfer, signalling changes in alveolar permeability and pulmonary capillary blood volume [16]. Oscillometry, meanwhile, can evaluate respiratory system mechanics and provide insight into the respiratory system's resistance and compliance [17]. Each of these tools has promise in the assessment of the peripheral lung, including in the smaller airways that are often labelled "the silent zone" but are increasingly recognised as an important indicator of early lung disease [18].

We systematically reviewed the literature to test our hypothesis that preterm-born individuals have abnormalities in static lung volumes, gas transfer and respiratory mechanics measured using oscillometry. In doing so, we provide a comprehensive assessment of respiratory function in survivors of preterm birth to better characterise the pathophysiological changes observed in this disease and highlight the tools with the most potential for diagnosing and monitoring prematurity-associated lung disease in clinical practice across the lifespan.

## Methods

The protocol for this systematic review and meta-analysis was developed according to the principles of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 statement and registered prospectively using PROSPERO (CRD42022320775, supplement 1) [19]. Our primary aim was to determine if lung function measured using static lung volumes, gas transfer or oscillometry significantly differed between preterm-born individuals and those born at term. Secondary aims were to investigate if extremely/very/moderate-late preterm birth, a diagnosis of BPD, era of neonatal care and/or age at lung function test affected lung function.

The search strategy was designed to locate articles that assessed lung function through static lung volumes, gas transfer or oscillometry in groups with a history of preterm birth, including those with or without a history of BPD. Additionally, search terms were included to capture articles reporting on cardiopulmonary exercise testing in preterm cohorts, which has been reported on separately [20]. Key words were chosen within these two broad categories with appropriate Medical Subject Headings identified (supplement 1, section 3.2.3). Six databases were searched: Embase, MEDLINE, CINAHL, Scopus, Proquest Academic and Web of Science. The databases were searched to include all results up to 29 April 2022 and again on 29 November 2024 for articles published between 1 January 2022 and 29 November 2024. Articles of all languages were accepted, seeking translations if required. Duplicate results were identified and removed first using EndNote's inbuilt "Find Duplicates" feature, then manually using ordered lists.

Studies were eligible for inclusion within this analysis if they defined both a preterm (<37 weeks' gestation) and a healthy term ( $\geq 37$  weeks' gestation) control cohort and reported any of the following: static lung volumes by plethysmography or inert gas washout; gas transfer by  $D_{LCO}$ ; or oscillometry measured by either impulse oscillometry or pseudo-random noise-based oscillometry techniques. Studies were excluded from the analysis if term and preterm cohort data could not be separated; cohorts had comorbidities, other than BPD, which may significantly affect respiratory health; or cohorts included data from children younger than 3 years. A 3-year cut-off was chosen based on differences in technique and

equipment typically found in measuring infant lung function. If required, authors were contacted for additional summary estimates of data where possible.

Titles and abstracts identified in searches were independently screened by two reviewers with backgrounds in medical research (J.T.D. Gibbons, M.L. Beaven, C.W. Course, S.J. Kotecha, M. Zuidersma, T. Hixson). Where either reviewer indicated the title may fit the above predefined eligibility criteria, the full manuscript was obtained and again independently reviewed by two reviewers, with a third reviewer (S.J. Simpson) mediating any disagreements. Where articles were not written in English, translations were sought. Where cohorts had data published in multiple manuscripts, the most complete data set was chosen for each of lung volumes, gas transfer and oscillometry to avoid over-representation within meta-analyses.

Risk of bias was assessed using a modified version of the Newcastle-Ottawa Quality Assessment Scale [21], with each study given a rating of low, medium or high risk of bias based on Agency of Healthcare Research and Quality standards (supplement 1, section 7.1). Subgroup analysis was performed using meta-regression to compare if studies given a rating of low risk of bias differed significantly from those of medium or high risk of bias. Publication bias was assessed if 10 or more cohorts were identified using Funnel plots and Egger's test [22].

Demographic and lung function data for each cohort were extracted into a Research Electronic Data Capture (REDCap) database into predefined fields (supplement 1, section 4.4.2) by a single reviewer (J.T.D. Gibbons, M.L. Beaven, C.W. Course, S.J. Kotecha) [23], with results validated by a second reviewer (J.T.D. Gibbons, M.L. Beaven, C.W. Course, S.J. Kotecha). Preterm cohorts were grouped by BPD status and banded into gestational age groups (extremely preterm: <28 weeks'; very preterm: 28–32 weeks'; moderate-late preterm: >32 weeks') based on the reported mean/median gestational age. Lung function data were extracted in order of preference as standard (Z) scores, per cent predicted or non-standardised data.

Statistical analysis was performed in R (version 4.3.2, [www.r-project.org](http://www.r-project.org)) using the *meta*, *metafor* and *dmatar* packages with previously documented techniques [24]. Where necessary, summary statistics from reported study groups were pooled such that a single group was represented for each of all individuals born preterm, all preterm individuals diagnosed with BPD as infants and all preterm individuals not diagnosed with BPD [25]. If variances were presented as standard error, 95% confidence intervals (95% CI) or interquartile range (IQR), these data were converted into standard deviations (SD) [26]. Where data were presented as medians and not significantly skewed [27], an estimate of the SD was calculated [28, 29].


Random-effects models were chosen for all meta-analyses owing to expected high levels of heterogeneity with variations in the techniques used for lung function testing and changes in the characteristics of prematurity-associated lung disease over time. For our primary outcome, standardised mean differences (SMDs) were calculated comparing preterm and term control groups for all lung function measures. For our secondary outcomes, where possible, the analyses were repeated with preterm groups analysed according to BPD status or banded into gestational age groups (<28 weeks', 28–32 weeks' or >32 weeks') based on the mean/median gestational age.

To analyse the impact of age and changes in neonatal care over time, we used meta-regression moderating lung function against the average age and birth year of the cohort, with significance determined by F-tests. A minimum of 10 cohorts per predictor was required for all meta-regression analyses to avoid over-fitting meta-regression models [25].

Where different techniques were used to measure lung volumes (plethysmography or inert gas washout) and respiratory mechanics using oscillometry (impulse oscillometry or pseudorandom noise), subgroup analyses using meta-regression were performed to determine if substantial differences in results existed between techniques. If the technique for measuring lung volumes was not specified, or if a combination of plethysmography and inert gas washout was used, the cohort was included in the overall analysis comparing lung volumes between preterm and term groups, but not in the subgroup analysis. Oscillometry values for resistance and reactance at 5 and 6 Hz were pooled, under the premise of their equivalence for analytical purposes.

## Results


In total, 12 143 titles were identified from our combined searches, with 1243 articles identified for full review (figure 1). In total, 240 articles met inclusion criteria, comprising 68 unique cohorts (supplement 2, table E1), with 53 reporting static lung volumes or ventilation inhomogeneity (34 measuring plethysmography, 25 measuring inert gas washout) [14, 30–81], 38 reporting gas transfer [7, 14, 31, 35,



**FIGURE 1** Flow diagram of search strategy. Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) flow diagram outlining search strategy for systematic review. Cardiopulmonary exercise testing (CPET) results discussed separately in BEAVEN *et al.* [20]. TLC: total lung capacity; RV: residual volume; FRC: functional residual capacity; LCI: lung clearance index; S<sub>cond</sub>: conductive ventilation heterogeneity; S<sub>acin</sub>: acinar ventilation heterogeneity; D<sub>LCO</sub>: diffusing capacity of the lung for carbon monoxide; K<sub>CO</sub>: transfer coefficient for carbon monoxide; V<sub>A</sub>: alveolar volume; R<sub>5/6 Hz</sub>: airway resistance at 5/6 Hz; R<sub>5/6-20 Hz</sub>: difference between airway resistance at 5/6 Hz and 20 Hz; X<sub>5/6</sub>: airway reactance at 5/6 Hz; F<sub>res</sub>: resonant frequency; AX: area under the reactance curve;  $\dot{V}_{CO_2}$ : carbon dioxide production;  $\dot{V}_{O_2}$ : oxygen production; WR: work rate;  $\dot{V}_E$ : minute ventilation;  $f_b$ : breathing frequency; V<sub>T</sub>: tidal volume; HR: heart rate; RER: respiratory exchange ratio; AT: aerobic threshold.

36, 39, 42–44, 46, 47, 49–52, 54, 56, 57, 59, 60, 64, 66–68, 71, 73–76, 81–90] and 18 reporting oscillometry outcomes [14, 46, 60, 73, 91–104]. In addition, 47 unique cohorts reported cardiopulmonary exercise testing and are discussed in the backing manuscript by BEAVEN *et al.* [20].

Characteristics of included studies are reported in supplement 2, table E2. Definitions of BPD varied widely, with diagnosis variably based on duration of supplemental oxygen or ventilation requirements, radiological abnormalities and/or clinical findings. Because we have previously shown minimal variation in FEV<sub>1</sub> outcomes across different diagnoses of BPD [9], we accepted the authors' reported definition of BPD. The birth year of participants ranged from 1961 to 2014, and gestational age ranged between 22 and 36 weeks. Lung function testing was conducted between 4 and 52 years of age. Study quality varied substantially, though did not appear to have any effect on the results (supplement 2, tables E4 and E5). Publication bias was evident for analyses performed on residual volumes (RVs), RV to total lung capacity (TLC) ratio and D<sub>LCO</sub>, although notably was nonsignificant once BPD status was considered (supplement 3).



**FIGURE 2** Summary of meta-analyses comparing the lung function of preterm and term-born cohorts. k: number of cohorts; nPT: number of preterm individuals in analysis; nFT: number of full-term controls in analysis; SMD: standardised mean difference; 95% CI: 95% confidence interval; FEV<sub>1</sub>: forced expiratory volume in 1 s; FVC: forced vital capacity; FEF<sub>25-75%</sub>: forced expiratory flow at 25–75% of FVC; TLC: total lung capacity; RV: residual volume; FRC: functional residual capacity; LCI: lung clearance index; S<sub>cond</sub>: conductive ventilation heterogeneity; S<sub>acin</sub>: acinar ventilation heterogeneity; D<sub>LCO</sub>: diffusing capacity of the lung for carbon monoxide; K<sub>CO</sub>: transfer coefficient for carbon monoxide; V<sub>A</sub>: alveolar volume; R<sub>5/6 Hz</sub>: airway resistance at 5/6 Hz; R<sub>20 Hz</sub>: airway resistance at 20 Hz; R<sub>5/6-20 Hz</sub>: difference between airway resistance at 5/6 Hz and 20 Hz; X<sub>5/6 Hz</sub>: airway reactance at 5/6 Hz; F<sub>res</sub>: resonant frequency; AX: area under the reactance curve. #: spirometry data provided for context [9, 10].

Summarised results of the primary analysis, including the number of eligible cohorts for each specific lung function test, are presented in figure 2, with differences between preterm and term-control groups expressed as SMD (95% CI). Spirometry values from our previous systematic review and meta-analysis are provided for context [10]. Results of secondary analyses assessing the impact of BPD and gestational age on lung function are presented in tables 1 and 2, respectively. Forest plots of individual meta-analyses are shown in supplement 4.

Preterm groups had altered static lung volumes and ventilation inhomogeneity compared to term controls, evidenced by higher RV (SMD 0.32, 95% CI 0.19 to 0.44), RV/TLC (SMD 0.45, 95% CI 0.28 to 0.63) and functional residual capacity (FRC) (SMD 0.15, 95% CI 0.01 to 0.29). Differences in all three lung function measures were greater in the preterm groups with BPD compared to preterm groups without BPD and term control groups. TLC, however, was not different between preterm and term groups and was not influenced by BPD status (table 1) or gestational age (table 2). Extremely preterm groups had significantly larger differences compared to term controls for RV and for RV as a percentage of TLC than did very preterm groups, although not different to moderate-late preterm groups. It should be noted, however, that fewer studies were available for analysis for the moderate-late preterm groups. No differences were observed between static lung volumes measured by plethysmography or inert gas washout (supplement 2, table E6).

Evidence of ventilation inhomogeneity was detected, with an elevated lung clearance index (LCI) seen in preterm groups (SMD 0.30, 95% CI 0.10 to 0.51). There were more pronounced differences observed in BPD groups (SMD 0.58, 95% CI 0.31 to 0.85) and groups born extremely preterm (SMD 0.54, 95% CI 0.16 to 0.92) when compared with term controls. Conductive ventilation heterogeneity ( $S_{\text{cond}}$ ) was increased in preterm compared to term control groups (SMD 0.81, 95% CI 0.38 to 1.24). No significant differences were noted in acinar ventilation heterogeneity between preterm- and term-born groups.

TABLE 1 Summary of meta-analyses comparing lung function of preterm cohorts with and without a diagnosis of BPD with term cohorts

	BPD versus Term			No BPD versus Term		
	k (nPT/nFT)	SMD (95% CI)	I <sup>2</sup> (%)	k (nPT/nFT)	SMD (95% CI)	I <sup>2</sup> (%)
<b>Static lung volumes</b>						
TLC	29 (952/1435)	-0.06 (-0.16 to 0.03)	19	24 (1135/1281)	-0.04 (-0.15 to 0.07)	35
RV	26 (880/1282)	<b>0.49 (0.30 to 0.67)</b>	66	22 (1050/1155)	<b>0.21 (0.05 to 0.37)</b>	55
RV/TLC	24 (855/1261)	<b>0.69 (0.50 to 0.89)</b>	69	23 (1128/1216)	<b>0.33 (0.17 to 0.48)</b>	58
FRC	15 (497/603)	<b>0.29 (0.11 to 0.48)</b>	39	13 (530/543)	0.12 (-0.04 to 0.28)	19
<b>Ventilation inhomogeneity</b>						
LCI	9 (318/299)	<b>0.58 (0.31 to 0.85)</b>	51	9 (265/309)	0.30 (-0.08 to 0.68)	66
$S_{\text{cond}}$	4 (116/112)	<b>0.61 (0.47 to 0.75)</b>	0	5 (143/150)	0.44 (-0.04 to 0.93)	51
$S_{\text{acin}}$	4 (116/115)	0.40 (-0.22 to 1.01)	40	4 (99/114)	0.02 (-0.74 to 0.77)	66
<b>Gas transfer</b>						
$D_{\text{LCO}}$	24 (680/877)	<b>-0.67 (-0.88 to -0.47)</b>	65	24 (873/839)	-0.43 (-0.63 to -0.24)	60
$K_{\text{CO}}$	17 (517/631)	<b>-0.55 (-0.76 to -0.35)</b>	51	16 (565/566)	-0.50 (-0.75 to -0.25)	66
$V_{\text{A}}$	9 (339/418)	-0.10 (-0.33 to 0.13)	38	10 (426/433)	0.02 (-0.11 to 0.15)	0
<b>Oscillometry</b>						
$R_{5/6}$ Hz	7 (372/391)	<b>0.88 (0.42 to 1.34)</b>	72	6 (224/472)	<b>0.57 (0.19 to 0.94)</b>	58
$R_{20}$ Hz	3 (109/120)	0.68 (-2.39 to 3.74)	92	3 (94/120)	0.64 (-0.45 to 1.73)	67
$R_{5/6-20}$ Hz	5 (261/366)	0.97 (-0.02 to 1.97)	83	3 (102/110)	0.43 (-0.07 to 0.93)	0
$X_{5/6}$ Hz	2 (75/59)	-0.63 (-2.85 to 1.60)	0	3 (115/307)	-0.25 (-0.97 to 0.47)	36
$F_{\text{res}}$	5 (218/224)	<b>1.53 (0.45 to 2.61)</b>	85	5 (191/224)	0.75 (-0.02 to 1.53)	80
AX	5 (292/282)	<b>0.87 (0.57 to 1.17)</b>	16	3 (93/73)	<b>0.61 (0.51 to 0.72)</b>	0

Bold values indicate results where the 95% confidence interval does not cross zero. BPD: bronchopulmonary dysplasia; k: number of cohorts; nPT: number of preterm individuals in analysis; nFT: number of full-term individuals in analysis; SMD: standardised mean difference; 95% CI: 95% confidence interval; TLC: total lung capacity; RV: residual volume; FRC: functional residual capacity; LCI: lung clearance index;  $S_{\text{cond}}$ : conductive ventilation heterogeneity;  $S_{\text{acin}}$ : acinar ventilation heterogeneity;  $D_{\text{LCO}}$ : diffusing capacity of the lung for carbon monoxide;  $K_{\text{CO}}$ : transfer coefficient for carbon monoxide;  $V_{\text{A}}$ : alveolar volume;  $R_{5/6}$  Hz: airway resistance at 5/6 Hz;  $R_{20}$  Hz: airway resistance at 20 Hz;  $R_{5/6-20}$  Hz: difference between airway resistance at 5/6 Hz and 20 Hz;  $X_{5/6}$  Hz: airway reactance at 5/6 Hz;  $F_{\text{res}}$ : resonant frequency; AX: area under the reactance curve.

TABLE 2 Summary of meta-analyses comparing lung function of preterm cohorts separated by degree of prematurity with term cohorts

	EP versus Term			VP versus Term			MLP versus Term		
	k (nPT/nFT)	SMD (95% CI)	<i>i</i> <sup>2</sup> (%)	k (nPT/nFT)	SMD (95% CI)	<i>i</i> <sup>2</sup> (%)	k (nPT/nFT)	SMD (95% CI)	<i>i</i> <sup>2</sup> (%)
<b>Static lung volumes</b>									
TLC	20 (1257/995)	-0.10 (-0.24 to 0.05)	47	19 (1268/930)	-0.11 (-0.25 to 0.04)	55	5 (249/285)	-0.05 (-0.28 to 0.18)	0
RV	20 (1440/953)	<b>0.37 (0.21 to 0.54)</b>	58	17 (1078/862)	<b>0.24 (0.07 to 0.41)</b>	53	5 (208/241)	<b>0.51 (0.004 to 1.02)</b>	64
RV/TLC	20 (1430/1010)	<b>0.61 (0.43 to 0.79)</b>	66	17 (1189/822)	<b>0.41 (0.22 to 0.60)</b>	61	6 (269/282)	<b>0.53 (0.11 to 0.94)</b>	63
FRC	11 (813/432)	<b>0.25 (0.07 to 0.44)</b>	40	10 (722/428)	0.18 (-0.04 to 0.41)	44	6 (240/271)	0.14 (-0.57 to 0.85)	67
<b>Ventilation inhomogeneity</b>									
LCI	6 (321/227)	<b>0.54 (0.16 to 0.92)</b>	67	5 (384/206)	<b>0.34 (0.25 to 0.43)</b>	0	3 (194/176)	0.02 (-0.29 to 0.34)	0
S <sub>cond</sub>	2 (42/57)	0.89 (-0.93 to 2.72)	0	2 (65/57)	0.81 (-1.79 to 3.41)	11	#	#	
S <sub>acin</sub>	2 (47/56)	0.16 (-2.84 to 3.16)	21	2 (67/56)	0.00 (0.00 to 0.00)	0	#	#	
<b>Gas transfer</b>									
D <sub>LCO</sub>	17 (1019/588)	<b>-0.66 (-0.91 to -0.42)</b>	71	21 (924/756)	<b>-0.54 (-0.73 to -0.35)</b>	58	6 (435/1419)	-0.15 (-0.36 to 0.06)	30
K <sub>CO</sub>	13 (948/493)	<b>-0.54 (-0.78 to -0.29)</b>	64	14 (682/486)	<b>-0.46 (-0.68 to -0.24)</b>	54	2 (147/141)	0.10 (-1.93 to 2.13)	30
V <sub>A</sub>	7 (673/311)	-0.04 (-0.20 to 0.13)	0	8 (546/364)	-0.06 (-0.23 to 0.12)	1	2 (147/141)	-0.38 (-1.47 to 0.72)	0
<b>Oscillometry</b>									
R <sub>5/6 Hz</sub>	6 (543/334)	<b>0.62 (0.37 to 0.88)</b>	40	6 (542/302)	<b>0.46 (0.03 to 0.90)</b>	81	5 (284/430)	0.15 (-0.10 to 0.39)	3
R <sub>20 Hz</sub>	3 (306/136)	0.77 (-2.02 to 3.56)	91	3 (181/140)	0.45 (-0.94 to 1.83)	81	3 (215/164)	-0.03 (-0.68 to 0.62)	42
R <sub>5/6-20 Hz</sub>	5 (483/313)	<b>0.65 (0.25 to 1.05)</b>	50	5 (294/222)	0.32 (-0.01 to 0.66)	43	2 (76/89)	-0.05 (-0.46 to 0.36)	0
X <sub>5/6 Hz</sub>	3 (110/109)	-0.31 (-1.47 to 0.85)	74	5 (465/229)	-0.24 (-0.63 to 0.14)	61	4 (249/414)	-0.05 (-0.45 to 0.36)	39
F <sub>res</sub>	3 (339/117)	1.32 (-0.69 to 3.33)	80	5 (505/252)	<b>0.50 (0.01 to 0.99)</b>	70	2 (188/145)	0.06 (-2.87 to 2.98)	74
AX	4 (227/236)	<b>0.70 (0.35 to 1.05)</b>	19	4 (292/133)	<b>0.60 (0.56 to 0.64)</b>	0	1 (49/70)	0.08 (-0.28 to 0.45)	N/A

Bold values indicate results where the 95% confidence interval does not cross zero. EP: extremely preterm (<28 weeks' gestation); VP: very preterm (28–32 weeks' gestation); MLP: moderate-late preterm (32–37 weeks' gestation); k: number of cohorts; nPT: number of preterm individuals in analysis; nFT: number of full-term individuals in analysis; SMD: standardised mean difference; 95% CI: 95% confidence interval; TLC: total lung capacity; RV: residual volume; FRC: functional residual capacity; LCI: lung clearance index; S<sub>cond</sub>: conductive ventilation heterogeneity; S<sub>acin</sub>: acinar ventilation heterogeneity; D<sub>LCO</sub>: diffusing capacity of the lung for carbon monoxide; K<sub>CO</sub>: transfer coefficient for carbon monoxide; V<sub>A</sub>: alveolar volume; R<sub>5/6 Hz</sub>: airway resistance at 5/6 Hz; R<sub>20 Hz</sub>: airway resistance at 20 Hz; R<sub>5/6-20 Hz</sub>: difference between airway resistance at 5/6 Hz and 20 Hz; X<sub>5/6 Hz</sub>: airway reactance at 5/6 Hz; F<sub>res</sub>: resonant frequency; AX: area under the reactance curve. #: S<sub>cond</sub> and S<sub>acin</sub> not included because there were insufficient cohorts to perform analyses for these tests.

Reduced gas transfer, as measured by  $D_{LCO}$  (SMD  $-0.51$ , 95% CI  $-0.64$  to  $-0.38$ ) and the transfer coefficient for carbon monoxide ( $K_{CO}$ ) (SMD  $-0.50$ , 95% CI  $-0.65$  to  $-0.34$ ), were observed in those born preterm compared to term controls. The reduction in  $D_{LCO}$ , but not  $K_{CO}$ , was more pronounced in those with a history of BPD (SMD  $-0.67$ , 95% CI  $-0.88$  to  $-0.47$ ) compared with those without BPD (SMD  $-0.43$ , 95% CI  $-0.63$  to  $-0.24$ ). Extremely preterm (SMD  $-0.66$ , 95% CI  $-0.91$  to  $-0.42$ ) and very preterm (SMD  $-0.54$ , 95% CI  $-0.73$  to  $-0.35$ ) birth were associated with substantial reductions in  $D_{LCO}$ , but this was not evident in those born moderate-late preterm. Alveolar volume ( $V_A$ ) was not different between preterm- and term-born groups.

Analysis of respiratory system mechanics measured using oscillometry showed that resistance at 5/6 Hz was greater in preterm cohorts than term cohorts (SMD  $0.44$ , 95% CI  $0.22$  to  $0.67$ ), particularly in those with BPD (SMD  $0.88$ , 95% CI  $0.42$  to  $1.34$ ) or with a history of extremely preterm birth (SMD  $0.62$ , 95% CI  $0.37$  to  $0.88$ ). Area under the reactance curve (AX) was also elevated in preterm (SMD  $0.62$ , 95% CI  $0.35$  to  $0.88$ ), BPD (SMD  $0.87$ , 95% CI  $0.57$  to  $1.17$ ) and extremely preterm groups (SMD  $0.70$ , 95% CI  $0.35$  to  $1.05$ ) compared to term controls, while resonant frequency was elevated in preterm cohorts (SMD  $0.63$ , 95% CI  $0.12$  to  $1.15$ ) and more substantially in BPD groups (SMD  $1.53$ , 95% CI  $0.45$  to  $2.61$ ). The difference between airway resistance at 5/6 Hz and 20 Hz ( $R_{5/6-20\text{ Hz}}$ ) was also higher in preterm cohorts compared with term cohorts (SMD  $0.51$ , 95% CI  $0.07$  to  $0.96$ ). Airway reactance at 5/6 Hz trended lower in those born preterm (SMD  $-0.15$ , 95% CI  $-0.37$  to  $0.07$ ), although a degree of uncertainty remained. No other measures of oscillometry were different between preterm and term cohorts. No differences were noted on subgroup analysis comparing airway resistance at 5/6 Hz ( $R_{5/6\text{ Hz}}$ ) collected using impulse oscillometry and pseudorandom noise ( $p=0.98$ ).

For lung function measures where meta-regression analysis was possible (TLC, RV, RV/TLC, FRC, LCI,  $D_{LCO}$ ,  $K_{CO}$ ,  $V_A$ ,  $R_{5/6}$ ), we did not identify any association between age or birth year (supplement 2, table E7). This suggests that alterations in these measures of lung function do not change with age, nor have they improved or declined across different eras of neonatal care.

## Discussion

This systematic review and meta-analysis expands on our previous reviews of spirometry outcomes [10], providing the first overview of static lung volumes, gas transfer and oscillometry in survivors of preterm birth. We found preterm birth to be associated with gas trapping on static lung volumes (increased RVs, FRC and RV/TLC), ventilation inhomogeneity (elevated LCI and  $S_{\text{cond}}$ ), impaired gas transfer (reductions in  $D_{LCO}$  and  $K_{CO}$ ) and altered respiratory mechanics suggestive of small airways disease (increased  $R_{5/6}$ ,  $R_{5/6-20}$  and AX). These disparities were often exacerbated where there was a history of either BPD or extremely preterm birth. Unlike our previous reviews of spirometry outcomes, we did not observe temporal changes in lung function, either as age-related changes or with changes in birth year across different eras of neonatal care.

Gas trapping has been recognised as a hallmark feature in obstructive lung diseases such as asthma and COPD [105], and with our findings of increased RVs, FRCs and RV/TLC ratios following preterm birth, this review suggests it warrants similar consideration as a feature of prematurity-associated lung disease. Large airway obstruction, evidenced by the reductions in  $FEV_1/FVC$  we have previously reported, offers one likely explanation for gas trapping following preterm birth [106]. However, alternative mechanisms, such as small airway dysfunction and parenchymal alterations, should also be explored to understand the complexities of gas trapping in the context of prematurity-associated lung disease. Indeed, signs of small airways disease were also evident in reported cohorts, with elevations in LCI on multiple breath washout and increased  $R_{5/6\text{ Hz}}$ ,  $R_{5/6-20\text{ Hz}}$  and AX observed using oscillometry in the preterm group. These findings also align with imaging studies showing small airways dysfunction and parenchymal alterations in survivors of preterm birth aged 9–11 years, which depicted bronchial wall thickening alongside areas of collapse/consolidation, emphysema and bronchiectasis [14]. With worsening airway obstruction observed in many preterm survivors as they age, it may have been expected that we would detect a similar progression in markers of gas trapping; however, no such signal was observed in this study and cohort studies reporting static lung volumes longitudinally are lacking. As concern grows of early-onset COPD in the preterm-born population [1], however, systematic monitoring of pulmonary function in clinical settings may become increasingly important. By focusing on the early identification and management of gas trapping, we can potentially mitigate the risk of developing early-onset COPD and other chronic respiratory conditions in these individuals.

Our analysis also identified impairment in alveolar gas transfer to the pulmonary vasculature following preterm birth, as indicated by significantly lower  $D_{LCO}$  and  $K_{CO}$  values. In the absence of changes to  $V_A$ ,

this could indicate that abnormalities in gas exchange are driven by pulmonary vascular impairments, emphysematous change (with preservation of lung volumes) or anaemia [106]. While there are no reports of persisting anaemia in later life in survivors of preterm birth, emphysematous changes have been described in the previously mentioned computed tomography imaging studies [14]. Likewise, histological samples taken from the lungs of preterm-born individuals often have both a larger simplified alveolar appearance akin to emphysema and impaired capillary growth [2, 13]. This implies that both pulmonary vascular impairments and vascular disease contribute to reductions in gas transfer following preterm birth. To determine which may have a greater impact on gas transfer, techniques have been employed to partition  $D_{LCO}$  into its constituent components: pulmonary capillary blood volume and pulmonary membrane diffusing capacity. However, results from the limited studies have thus far been conflicting for survivors of preterm birth [107, 108]. We did not identify any age-related improvement or decline in gas transfer measurements in this study, matching what has been reported in longitudinal follow-up [108]. The clinical significance of  $D_{LCO}$  measurement in prematurity-associated lung disease is not yet fully established; however, given that a diminished  $D_{LCO}$  is recognised as a predictive marker for pulmonary hypertension in COPD [109], it potentially serves as a vital indicator for the necessity of comprehensive cardiopulmonary assessment in the preterm-born population.

BPD is recognised as an indicator of increased disease severity in neonatal and early childhood respiratory diseases, with our study findings reinforcing the association of BPD with more pronounced abnormalities in gas trapping, small airway resistance and impaired gas transfer. First identified over half a century ago, the definition of BPD has undergone significant evolution and remains a subject of ongoing debate. Previously we have identified that the specific definition of BPD employed did not significantly influence subsequent lung function outcomes [9]. Consequently, in this analysis, we intentionally avoided distinguishing between various definitions of BPD, aiming for a pragmatic assessment of the association between an early-life BPD diagnosis and later impaired lung function. Although BPD correlated with more significant lung function changes, our observations also showed lung function variations in preterm infants without BPD. Similarly, while lower gestational ages were also associated with more substantial impacts on lung function, even individuals born moderate-late preterm demonstrated persistent lung function abnormalities with signs of gas trapping and reduced gas transfer. This implies that preterm birth itself, irrespective of BPD or degree of prematurity, remains a critical risk factor for long-term respiratory complications, highlighting the need for comprehensive monitoring and tailored interventions for all preterm infants.

The strengths of this study lie in its ability to pool lung function data presented in different formats (Z-score, per cent predicted or raw) to maximise the number of studies included within each meta-analysis. Additionally, by performing meta-analyses on lung volumes, gas transfer and oscillometry in this systematic review, and consolidating them with our recent systematic review on spirometry, we have provided a comprehensive overview of the summarised lung function seen following preterm birth and interpreted each finding in the context of term-born controls. Limitations include the exclusion of cohorts without a term-born control population, and the lack of data from children aged under 3 years. Infant lung function techniques were considered too different to techniques beyond infancy and were thus omitted from this systematic review. Furthermore, while some cohorts did report on data longitudinally, analysing these data was outside the scope of this review. Additionally, moderate-high levels of heterogeneity were generally observed across the analyses, which may limit the ability to draw significant conclusions, particularly where fewer cohorts were available for analysis. Factors that may have contributed to heterogeneity include variations in the pathology of preterm lung disease across multiple decades, varying definitions of BPD and differences in methodology in the collection of pulmonary function tests.

In summary, here we have provided the first systematic review and meta-analyses on static lung volumes, gas transfer and oscillometry. When combined with our previous reviews on spirometry, these present the most comprehensive assessment of lung function in survivors of preterm birth to date. While spirometry is the most-performed lung function test, we demonstrate that the use of alternative lung function tests to assess the small airways, lung parenchyma and/or pulmonary vasculature offers valuable insight when diagnosing or monitoring the long-term sequelae of preterm birth. Early markers of respiratory disease are critical to prevent disease progression, and tests like those described in this review that assess the “silent zone” are often used as early markers in asthma and COPD. The concept of pre-COPD, in which individuals develop early physiological dysfunction before developing detectable airway obstruction by spirometry, is particularly relevant here because these individuals may develop reduced  $D_{LCO}$  or signs of gas trapping as the earliest manifestations of disease [110]. Currently, definitive studies establishing how such measures present or progress longitudinally in prematurity-associated lung disease are lacking, although our systematic review suggests that a signal should be expected. Coupled with the underpinning

pathophysiology of simplified alveolar structure (a reduced number of larger alveoli) and abnormal pulmonary vascular development, we suggest that these specialised lung function tests should be considered in a clinical setting to identify specific “treatable traits” in survivors of preterm birth.

#### Points for clinical practice

- Consider incorporating comprehensive pulmonary function testing beyond spirometry where available. Static lung volumes, diffusing capacity of the lung for carbon monoxide and oscillometry may help detect gas trapping, small airways dysfunction and impaired gas transfer in preterm survivors that could be missed by spirometry alone.
- Long-term respiratory follow-up is recommended for preterm survivors, including those born moderate-late preterm. Regular monitoring may help to detect early signs of COPD, particularly in individuals with persistent gas trapping or reduced gas transfer.
- Consider prioritising more frequent monitoring for high-risk groups, particularly those with a history of bronchopulmonary dysplasia or born extremely preterm (<28 weeks), because these individuals often show more pronounced lung function abnormalities and may benefit from closer monitoring and earlier intervention.

Published in volume 34, issue 176 of the *European Respiratory Review* on 28 May 2025; republished 13 June 2025 with amendments to typographical errors in figure 1 and correction of the database search end date, which was correct in the methodology section of the article, in the abstract.

Acknowledgements: The authors wish to acknowledge and thank Professor Shyamali Dharmage and Dr Dinh Bui (University of Melbourne, Australia), and Professor Jeanie Cheong and Dr Cassidy Du Berry (Murdoch Children’s Research Institute, Australia) for contributing additional unpublished data for inclusion within this systematic review.

Provenance: Submitted article, peer reviewed.

Author contributions: J.T.D. Gibbons executed the study searches, conducted the data analysis and contributed to the study design, search strategy, outcome selection, database management, screening, data extraction and validation, data interpretation and manuscript preparation. M.L. Beaven contributed to screening, data extraction, data validation and manuscript review. C.W. Course contributed to the study search strategy, screening, data validation and manuscript review. S.J. Kotecha contributed to the study outcome selection, search strategy, screening, data validation and data interpretation. T. Hixson and M. Zuidersma contributed to the study screening. A.C. Wilson contributed to data interpretation and manuscript review. S. Kotecha contributed to the study outcome selection, search strategy and data interpretation. S.J. Simpson contributed to study outcome selection, search strategy, screening, data analysis, data interpretation and manuscript review. All authors contributed to the article and approved the submitted version.

Conflict of interest: The study authors declare no competing interests in this review.

Support statement: CAHS Research Scholarship, Department of Health Western Australia. The study funders had no role in the outcomes of this review.

#### References

- 1 Simpson SJ, Du Berry C, Evans DJ, *et al.* Unravelling the respiratory health path across the lifespan for survivors of preterm birth. *Lancet Respir Med* 2024; 12: 167–180.
- 3 Du Berry C, Nesci C, Cheong JLY, *et al.* Long-term expiratory airflow of infants born moderate-late preterm: a systematic review and meta-analysis. *eClinicalMedicine* 2022; 52: 101597.
- 4 Fawke J, Lum S, Kirkby J, *et al.* Lung function and respiratory symptoms at 11 years in children born extremely preterm: the EPICure study. *Am J Respir Crit Care Med* 2010; 182: 237–245.
- 5 Kim YJ, Shin SH, Park HW, *et al.* Risk factors of early pulmonary hypertension and its clinical outcomes in preterm infants: a systematic review and meta-analysis. *Sci Rep* 2022; 12: 14186.
- 6 Stevenson PG, Cooper MN, Billingham W, *et al.* Health service utilisation for acute respiratory infections in infants graduating from the neonatal intensive care unit: a population-based cohort study. *BMC Pediatr* 2023; 23: 335.
- 7 Bui DS, Perret JL, Walters EH, *et al.* Association between very to moderate preterm births, lung function deficits, and COPD at age 53 years: analysis of a prospective cohort study. *Lancet Respir Med* 2022; 10: 478–484.

- 8 Duijts L, van Meel ER, Moschino L, *et al.* European Respiratory Society guideline on long-term management of children with bronchopulmonary dysplasia. *Eur Respir J* 2020; 55: 1900788.
- 9 Kotecha SJ, Gibbons JTD, Course CW, *et al.* Geographical differences and temporal improvements in forced expiratory volume in 1 second of preterm-born children: a systematic review and meta-analysis. *JAMA Pediatr* 2022; 176: 867–877.
- 10 Gibbons JTD, Course CW, Evans EE, *et al.* Increasing airway obstruction through life following bronchopulmonary dysplasia: a meta-analysis. *ERJ Open Res* 2023; 9: 00046-2023.
- 11 Doyle LW, Adams AM, Robertson C, *et al.* Increasing airway obstruction from 8 to 18 years in extremely preterm/low-birthweight survivors born in the surfactant era. *Thorax* 2017; 72: 712–719.
- 12 Simpson SJ, Turkovic L, Wilson AC, *et al.* Lung function trajectories throughout childhood in survivors of very preterm birth: a longitudinal cohort study. *Lancet Child Adolesc Health* 2018; 2: 350–359.
- 13 Liu N, Cummings OW, Lagstein A, *et al.* Lung transplantation for bronchopulmonary dysplasia in adults: a clinical and pathologic study of 3 cases. *Am J Surg Pathol* 2020; 44: 509–515.
- 14 Simpson SJ, Logie KM, O’Dea CA, *et al.* Altered lung structure and function in mid-childhood survivors of very preterm birth. *Thorax* 2017; 72: 702–711.
- 15 Bhakta NR, McGowan A, Ramsey KA, *et al.* European Respiratory Society/American Thoracic Society technical standard on standardisation of the measurement of lung volumes - 2023 update. *Eur Respir J* 2023; 62: 2201519.
- 16 Graham BL, Brusasco V, Burgos F, *et al.* 2017 ERS/ATS standards for single-breath carbon monoxide uptake in the lung. *Eur Respir J* 2017; 49: 1600016.
- 17 King GG, Bates J, Berger KI, *et al.* Technical standards for respiratory oscillometry. *Eur Respir J* 2020; 55: 1900753.
- 18 Hogg JC, Paré PD, Hackett T-L. The contribution of small airway obstruction to the pathogenesis of chronic obstructive pulmonary disease. *Physiol Rev* 2017; 97: 529–552.
- 19 Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71.
- 20 Beaven ML, Gibbons JTD, Course CW, *et al.* Physiological responses to exercise in survivors of preterm birth: a meta-analysis. *Eur Respir Rev* 2025; in press [<https://doi.org/10.1183/16000617.0163-2024>].
- 21 Wells G, Shea B, O’Connell D, *et al.* The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Available from: [www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp) Date last updated: 3 May 2021.
- 22 Sterne JAC, Sutton AJ, Ioannidis JPA, *et al.* Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011; 343: d4002.
- 23 Harris PA, Taylor R, Thielke R, *et al.* Research electronic data capture (REDCap) – a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009; 42: 377–381.
- 24 Harrer M, Cuijpers P, Furukawa TA, *et al.* Doing meta-analysis with R: a hands-on guide. 1st ed. Boca Raton, Chapman & Hall/CRC Press, 2021.
- 25 Deeks J, Higgins J, Altman DG. Chapter 10: Analysing data and undertaking meta-analyses. *In:* Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch V, eds. *Cochrane Handbook for Systematic Reviews of Interventions* version 63 (updated February 2022). London, Cochrane, 2022.
- 26 Higgins J, Li T, Deeks J. Chapter 6: Choosing effect measures and computing estimates of effect. *In:* Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch V, eds. *Cochrane Handbook for Systematic Reviews of Interventions* version 63 (updated February 2022). London, Cochrane, 2022.
- 27 Shi J, Luo D, Wan X, *et al.* Detecting the skewness of data from the sample size and the five-number summary. *arXiv* 2020, preprint [<https://doi.org/10.48550/arXiv.2010.05749>].
- 28 Luo D, Wan X, Liu J, *et al.* Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Meth Med Res* 2018; 27: 1785–1805.
- 29 Wan X, Wang W, Liu J, *et al.* Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; 14: 135.
- 30 Amitai N, Staffler P, Blau H, *et al.* Palivizumab following extremely premature birth does not affect pulmonary outcomes in adolescence. *Chest* 2020; 158: 660–669.
- 31 Arigliani M, Stocco C, Valentini E, *et al.* Lung function between 8 and 15 years of age in very preterm infants with fetal growth restriction. *Pediatr Res* 2021; 90: 657–663.
- 32 Bader D, Ramos AD, Lew CD, *et al.* Childhood sequelae of infant lung disease: exercise and pulmonary function abnormalities after bronchopulmonary dysplasia. *J Pediatr* 1987; 110: 693–699.
- 33 Barker M, Merz U, Hertl MS, *et al.* School-age lung function and exercise capacity in former very low birth weight infants. *Pediatr Exerc Sci* 2003; 15: 44–55.
- 34 Bertrand JM, Riley SP, Popkin J, *et al.* The long-term pulmonary sequelae of prematurity: the role of familial airway hyperreactivity and the respiratory distress syndrome. *N Engl J Med* 1985; 312: 742–745.

- 35 Caskey S, Gough A, Rowan S, *et al.* Structural and functional lung impairment in adult survivors of bronchopulmonary dysplasia. *Ann Am Thorac Soc* 2016; 13: 1262–1270.
- 36 Cazzato S, Ridolfi L, Bernardi F, *et al.* Lung function outcome at school age in very low birth weight children. *Pediatr Pulmonol* 2013; 48: 830–837.
- 37 Chan HF, Smith LJ, Biancardi AM, *et al.* Image phenotyping of preterm-born children using hyperpolarized <sup>129</sup>Xe lung magnetic resonance imaging and multiple-breath washout. *Am J Respir Crit Care Med* 2023; 207: 89–100.
- 38 Cousins M, Hart K, Williams EM, *et al.* Impaired exercise outcomes with significant bronchodilator responsiveness in children with prematurity-associated obstructive lung disease. *Pediatr Pulmonol* 2022; 57: 2161–2171.
- 39 Di Filippo P, Giannini C, Attanasi M, *et al.* Pulmonary outcomes in children born extremely and very preterm at 11 years of age. *Front Pediatr* 2021; 9: 635503.
- 40 Doyle LW, Cheung MM, Ford GW, *et al.* Birth weight <1501 g and respiratory health at age 14. *Arch Dis Child* 2001; 84: 40–44.
- 41 Doyle LW, Victorian Infant Collaborative Study Group. Respiratory function at age 8–9 years in extremely low birthweight/very preterm children born in Victoria in 1991–1992. *Pediatr Pulmonol* 2006; 41: 570–576.
- 42 Du Berry C, Mainzer RM, Westrupp N, *et al.* The effect of being born moderate-late preterm on lung function and respiratory morbidity at 9 to 10 years of age. *Ann Am Thorac Soc* 2025; 22: 732–741.
- 43 Evensen KA, Steinshamn S, Tjonaa AE, *et al.* Effects of preterm birth and fetal growth retardation on cardiovascular risk factors in young adulthood. *Early Hum Dev* 2009; 85: 239–245.
- 44 Galdès-Sebaldt M, Sheller J, Groggaard J, *et al.* Prematurity is associated with abnormal airway function in childhood. *Pediatr Pulmonol* 1989; 7: 259–264.
- 45 Gross SJ, Iannuzzi DM, Kveselis DA, *et al.* Effect of preterm birth on pulmonary function at school age: a prospective controlled study. *J Pediatr* 1998; 133: 188–192.
- 46 Hagman C, Bjorklund LJ, Bjermer L, *et al.* Lung function deficits and bronchodilator responsiveness at 12 years of age in children born very preterm compared with controls born at term. *Pediatr Pulmonol* 2023; 58: 3156–3170.
- 47 Hakulinen AL, Jarvenpaa AL, Turpeinen M, *et al.* Diffusing capacity of the lung in school-aged children born very preterm, with and without bronchopulmonary dysplasia. *Pediatr Pulmonol* 1996; 21: 353–360.
- 48 Halvorsen T, Skadberg BT, Eide GE, *et al.* Assessment of lung volumes in children and adolescents: comparison of two plethysmographic techniques. *Clin Physiol Funct Imaging* 2005; 25: 62–68.
- 49 Joshi S, Powell T, Watkins WJ, *et al.* Exercise-induced bronchoconstriction in school-aged children who had chronic lung disease in infancy. *J Pediatr* 2013; 162: 813–818.
- 50 Kaplan E, Bar-Yishay E, Prais D, *et al.* Encouraging pulmonary outcome for surviving, neurologically intact, extremely premature infants in the postsurfactant era. *Chest* 2012; 142: 725–733.
- 51 Karila C, Saulnier J, Elie C, *et al.* Exercise-induced alveolar hypoventilation in long-term survivors of bronchopulmonary dysplasia. *Rev Mal Respir* 2008; 25: e61–e68.
- 52 Karnaushkina MA, Strutinskaya AD, Ovsyannikov DY. Prematurity and early childhood infection of lower respiratory tract as risk factors of developing chronic obstructive bronchopulmonary pathology in adults. *Sovremennye Tehnologii v Medicine* 2017; 9: 129.
- 53 Kennedy JD, Edward LJ, Bates DJ, *et al.* Effects of birthweight and oxygen supplementation on lung function in late childhood in children of very low birth weight. *Pediatr Pulmonol* 2000; 30: 32–40.
- 54 Korhonen P, Laitinen J, Hyodynmaa E, *et al.* Respiratory outcome in school-aged, very-low-birth-weight children in the surfactant era. *Acta Paediatr* 2004; 93: 316–321.
- 55 Kouroukli E, Hatzigorou E, Sarafidis K, *et al.* Chronic lung disease of prematurity vs cystic fibrosis: exercise capacity and ventilation inhomogeneity. *Pediatr Pulmonol* 2021; 56: A52.
- 56 Landry JS, Tremblay GM, Li PZ, *et al.* Lung function and bronchial hyperresponsiveness in adults born prematurely. A cohort study. *Ann Am Thorac Soc* 2016; 13: 17–24.
- 57 Lum S, Kirkby J, Welsh L, *et al.* Nature and severity of lung function abnormalities in extremely pre-term children at 11 years of age. *Eur Respir J* 2011; 37: 1199–1207.
- 58 Lundberg B, Merid SK, Um-Bergstrom P, *et al.* Lung function in young adulthood in relation to moderate-to-late preterm birth. *ERJ Open Res* 2024; 10: 00701-2023.
- 59 MacLean JE, DeHaan K, Fuhr D, *et al.* Altered breathing mechanics and ventilatory response during exercise in children born extremely preterm. *Thorax* 2016; 71: 1012–1019.
- 60 Malmberg LP, Mieskonen S, Pelkonen A, *et al.* Lung function measured by the oscillometric method in prematurely born children with chronic lung disease. *Eur Respir J* 2000; 16: 598–603.
- 61 Molgat-Seon Y, Dominelli PB, Peters CM, *et al.* Analysis of maximal expiratory flow-volume curves in adult survivors of preterm birth. *Am J Physiol Regul Integr Comp Physiol* 2019; 317: R588–R596.
- 62 Narayanan M, Beardsmore C, Owers-Bradley J, *et al.* Evidence for acinar airway injury and alveolar catch-up growth in survivors of neonatal chronic lung disease. *Am J Respir Crit Care Med* 2010; 181: A3755.

- 63 Narayanan M, Beardsmore CS, Owers-Bradley J, *et al.* Catch-up alveolarization in ex-preterm children: evidence from <sup>3</sup>He magnetic resonance. *Am J Respir Crit Care Med* 2013; 187: 1104–1109.
- 64 Northway WH, Jr., Moss RB, Carlisle KB, *et al.* Late pulmonary sequelae of bronchopulmonary dysplasia. *N Engl J Med* 1990; 323: 1793–1799.
- 65 Rideau Batista Novais A, Matecki S, Jaussent A, *et al.* Hyperventilation during exercise in very low birth weight school-age children may implicate inspiratory muscle weakness. *J Pediatr* 2012; 160: 415–420.
- 66 Pianosi PT, Fisk M. High frequency ventilation trial. Nine year follow up of lung function. *Early Hum Dev* 2000; 57: 225–234.
- 67 Ronkainen E, Dunder T, Peltoniemi O, *et al.* New BPD predicts lung function at school age: follow-up study and meta-analysis. *Pediatr Pulmonol* 2015; 50: 1090–1098.
- 68 Ruf K, Thomas W, Brunner M, *et al.* Diverging effects of premature birth and bronchopulmonary dysplasia on exercise capacity and physical activity - a case control study. *Respir Res* 2019; 20: 260.
- 69 Salem Y, Willers CC, Kentgens A, *et al.* Lung function and structure in preterm-born adolescents of the surfactant era. *Eur Respir J* 2022; 60: Suppl. 66, 2881.
- 70 Smith LJ, van Asperen PP, McKay KO, *et al.* Reduced exercise capacity in children born very preterm. *Pediatrics* 2008; 122: e287–e293.
- 71 Sørensen JK, Buchvald F, Berg AK, *et al.* Ventilation inhomogeneity and NO and CO diffusing capacity in ex-premature school children. *Respir Med* 2018; 140: 94–100.
- 72 Todisco T, de Benedictis FM, Iannacci L, *et al.* Mild prematurity and respiratory functions. *Eur J Pediatr* 1993; 152: 55–58.
- 73 Um-Bergstrom P, Hallberg J, Pourbazargan M, *et al.* Pulmonary outcomes in adults with a history of bronchopulmonary dysplasia differ from patients with asthma. *Respir Res* 2019; 20: 102.
- 74 Vollaeter M, Clemm HH, Satrell E, *et al.* Adult respiratory outcomes of extreme preterm birth. A regional cohort study. *Ann Am Thorac Soc* 2015; 12: 313–322.
- 75 Vollaeter M, Skromme K, Satrell E, *et al.* Children born preterm at the turn of the millennium had better lung function than children born similarly preterm in the early 1990s. *PLoS One* 2015; 10: e0144243.
- 76 Vrijlandt EJ, Gerritsen J, Boezen HM, *et al.* Lung function and exercise capacity in young adults born prematurely. *Am J Respir Crit Care Med* 2006; 173: 890–896.
- 77 Vrijlandt E, Reijneveld SA, Aris-Meijer JL, *et al.* Respiratory health in adolescents born moderately-late preterm in a community-based cohort. *J Pediatr* 2018; 203: 429–436.
- 78 Wjst M, Popescu M, Trepka MJ, *et al.* Pulmonary function in children with initial low birth weight. *Pediatr Allergy Immunol* 1998; 9: 80–90.
- 79 Yaacoby-Bianu K, Plonsky MT, Gur M, *et al.* Effect of late preterm birth on lung clearance index and respiratory physiology in school-age children. *Pediatr Pulmonol* 2019; 54: 1250–1256.
- 80 Yammine S, Schmidt A, Sutter O, *et al.* Functional evidence for continued alveolarisation in former preterms at school age? *Eur Respir J* 2016; 47: 147–155.
- 81 Yang J, Kingsford RA, Horwood J, *et al.* Lung function of adults born at very low birth weight. *Pediatrics* 2020; 145: e20192359.
- 82 de Kleine MJ, Roos CM, Voorn WJ, *et al.* Lung function 8–18 years after intermittent positive pressure ventilation for hyaline membrane disease. *Thorax* 1990; 45: 941–946.
- 83 Duke JW, Gladstone IM, Sheel AW, *et al.* Premature birth affects the degree of airway dysanapsis and mechanical ventilatory constraints. *Exp Physiol* 2018; 103: 261–275.
- 84 Hadchouel A, Rousseau J, Roze JC, *et al.* Association between asthma and lung function in adolescents born very preterm: results of the EPIPAGE cohort study. *Thorax* 2018; 73: 1174–1176.
- 85 Haraldsdottir K, Watson AM, Pegelow DF, *et al.* Blunted cardiac output response to exercise in adolescents born preterm. *Eur J Appl Physiol* 2020; 120: 2547–2554.
- 86 Kumari S, Barton GP, Goss KN. Increased mitochondrial oxygen consumption in adult survivors of preterm birth. *Pediatr Res* 2021; 90: 1147–1152.
- 87 Manferdelli G, Narang BJ, Bourdillon N, *et al.* Physiological responses to exercise in hypoxia in preterm adults: convective and diffusive limitations in the O<sub>2</sub> transport. *Med Sci Sports Exerc* 2023; 55: 482–496.
- 88 Mitchell SH, Teague WG. Reduced gas transfer at rest and during exercise in school-age survivors of bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 1998; 157: 1406–1412.
- 89 Narang I, Bush A, Rosenthal M. Gas transfer and pulmonary blood flow at rest and during exercise in adults 21 years after preterm birth. *Am J Respir Crit Care Med* 2009; 180: 339–345.
- 90 Satrell E, Roksund O, Thorsen E, *et al.* Pulmonary gas transfer in children and adolescents born extremely preterm. *Eur Respir J* 2013; 42: 1536–1544.
- 91 Accorsi BF, Friedrich FO, Corso AL, *et al.* Intra-breath oscillometry for the evaluation of lung function in children and adolescents with a history of preterm birth. *J Bras Pneumol* 2022; 48: e20210290.
- 92 Cousins M, Hart K, Radics BL, *et al.* Peripheral airway dysfunction in prematurity-associated obstructive lung disease identified by oscillometry. *Pediatr Pulmonol* 2023; 58: 3279–3292.

- 93 Dantas F, Magalhaes PAF, Hora ECN, *et al.* Lung mechanics and respiratory morbidities in school-age children born moderate-to-late preterm. *Pediatr Res* 2022; 91: 1136–1140.
- 94 Durlak W, Klimek M, Wronski M, *et al.* Multimodal longitudinal respiratory function assessment in very low birth weight 7-year-old children. *Adv Med Sci* 2021; 66: 81–88.
- 95 Er I, Gunlemez A, Uyan ZS, *et al.* Evaluation of pulmonary functions in preschool children born late-preterm. *Turk Pediatri Ars* 2017; 52: 72–78.
- 96 Goth FEM, Green K, Hansen BM, *et al.* From neonatal lung function to lung function and respiratory morbidity at 6-year follow-up. *Pediatr Pulmonol* 2023; 58: 566–576.
- 97 Huang CY, Lai SH, Tseng HJ, *et al.* Pulmonary function in school-age children following intravitreal injection of bevacizumab for retinopathy of prematurity. *Sci Rep* 2022; 12: 18788.
- 98 Kalliola S, Malmberg LP, Kajosaari M, *et al.* Assessing direct and indirect airway hyperresponsiveness in children using impulse oscillometry. *Ann Allergy Asthma Immunol* 2014; 113: 166–172.
- 99 Knihtila H, Kotaniemi-Syrjanen A, Pelkonen AS, *et al.* Sensitivity of newly defined impulse oscillometry indices in preschool children. *Pediatr Pulmonol* 2017; 52: 598–605.
- 100 Said HA. Long term lung function in very low birth weight children in the surfactant era (PhD dissertation). Madison, WI, University of Wisconsin-Madison, 2003. Available from: [www.proquest.com/docview/305298205/](http://www.proquest.com/docview/305298205/)
- 101 Shackleton C, Czovek D, Grimwood K, *et al.* Defining ‘healthy’ in preschool-aged children for forced oscillation technique reference equations. *Respirology* 2018; 23: 406–413.
- 102 Suursalmi P, Kopeli T, Korhonen P, *et al.* Very low birthweight bronchopulmonary dysplasia survivors show no substantial association between lung function and current inflammatory markers. *Acta Paediatr* 2015; 104: 264–268.
- 103 Thunqvist P, Gustafsson PM, Schultz ES, *et al.* Lung function at 8 and 16 years after moderate-to-late preterm birth: a prospective cohort study. *Pediatrics* 2016; 137: e20152056.
- 104 Vrijlandt EJ, Boezen HM, Gerritsen J, *et al.* Respiratory health in prematurely born preschool children with and without bronchopulmonary dysplasia. *J Pediatr* 2007; 150: 256–261.
- 105 Hardin M, Silverman EK, Barr RG, *et al.* The clinical features of the overlap between COPD and asthma. *Respir Res* 2011; 12: 127.
- 106 Stanojevic S, Kaminsky DA, Miller MR, *et al.* ERS/ATS technical standard on interpretive strategies for routine lung function tests. *Eur Respir J* 2022; 60: 2101499.
- 107 Drummond D, Hadchouel A, Le Bourgeois M, *et al.* Decreased pulmonary capillary volume in adolescents born very preterm. *Acta Paediatrica* 2020; 109: 621–622.
- 108 Satrell E, Clemm H, Røksund OD, *et al.* Development of lung diffusion to adulthood following extremely preterm birth. *Eur Respir J* 2022; 59: 2004103.
- 109 Beyhan Sagmen S, Fidan A. Can FVC/D<sub>LCO</sub> predict pulmonary hypertension in patients with chronic obstructive pulmonary disease? *Eur Rev Med Pharmacol Sci* 2022; 26: 6658–6664.
- 110 Dharmage SC, Faner R, Agustí A. Treatable traits in pre-COPD: time to extend the treatable traits paradigm beyond established disease. *Respirology* 2024; 29: 551–562.