

# Geophysical Research Letters®



## RESEARCH LETTER

10.1029/2025GL115727

### Key Points:

- Shifts between magmatic and phreatomagmatic activity over the life span of volcanic fields may reflect fluctuations in groundwater levels
- Long-term changes in hydroclimate influenced water-magma interactions at Flores Island but not on shorter timescales (Holocene)
- At ocean island volcanoes with perched aquifers, variations in the mass eruption rate are crucial for triggering phreatomagmatism

### Supporting Information:

Supporting Information may be found in the online version of this article.

### Correspondence to:

M. Andrade,  
[mandrade@geomar.de](mailto:mandrade@geomar.de)

### Citation:

Andrade, M., Hernández, A., Pimentel, A., Cruz, J. V., Ramos, A. M., Ludwig, P., et al. (2025). Controls on water-magma interactions at hydraulically-charged volcanic islands. *Geophysical Research Letters*, 52, e2025GL115727. <https://doi.org/10.1029/2025GL115727>

Received 7 MAR 2025

Accepted 1 SEP 2025

### Author Contributions:

**Conceptualization:** M. Andrade, A. Hernández, A. Pimentel, R. S. Ramalho  
**Data curation:** J. V. Cruz, A. M. Ramos, P. Ludwig  
**Formal analysis:** M. Andrade  
**Investigation:** M. Andrade  
**Methodology:** M. Andrade, A. M. Ramos, P. Ludwig  
**Supervision:** A. Hernández, A. Pimentel, J. C. Schindlbeck-Belo, R. S. Ramalho  
**Writing – original draft:** M. Andrade  
**Writing – review & editing:** A. Hernández, A. Pimentel, J. V. Cruz, A. M. Ramos, P. Ludwig, J. C. Schindlbeck-Belo, R. S. Ramalho

© 2025. The Author(s).

This is an open access article under the terms of the [Creative Commons Attribution License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## Controls on Water-Magma Interactions at Hydraulically-Charged Volcanic Islands

M. Andrade<sup>1,2</sup> , A. Hernández<sup>3</sup> , A. Pimentel<sup>4</sup> , J. V. Cruz<sup>4,5</sup>, A. M. Ramos<sup>6</sup>, P. Ludwig<sup>6</sup> , J. C. Schindlbeck-Belo<sup>1</sup> , and R. S. Ramalho<sup>2,7</sup>


<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany, <sup>2</sup>IDL - Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal, <sup>3</sup>Universidade da Coruña, GRICA Group, Centro Interdisciplinar de Química e Biología (CICA), A Coruña, Spain, <sup>4</sup>Instituto de Investigação em Vulcanologia e Avaliação de Riscos (IVAR), Universidade dos Açores, Ponta Delgada, Portugal, <sup>5</sup>Faculdade de Ciências e Tecnologia (FCT), Universidade dos Açores, Ponta Delgada, Portugal, <sup>6</sup>Institute of Meteorology and Climate Research Troposphere Research (IMKTRO), Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany, <sup>7</sup>School of Earth and Environmental Sciences, Cardiff University, Cardiff, UK

**Abstract** The interaction of rising magma with groundwater can produce phreatomagmatic explosions, which, given their enhanced explosivity and unpredictability, increase the hazard potential of volcanic eruptions. To investigate the link between groundwater occurrence and phreatomagmatism, we compare the volcanic record of Flores Island (Azores) with regional climate reconstructions. Flores is an ideal case study as it experienced multiple magmatic and phreatomagmatic eruptions during the Holocene and Pleistocene. Our results show that at a broader scale (>10 ka), magmatic volcanism prevailed during dry/colder periods, whereas phreatomagmatism preferentially occurred during wet/warm periods. At a shorter timescale (<10 ka), however, water-magma interactions were primarily controlled by variations in eruption rates, with rainfall variability having a secondary role, as phreatomagmatism occurred even during low precipitation periods. This study reinforces that on island volcanoes with perched aquifers and prone to monogenetic volcanism, eruption rates rather than short-term hydroclimate changes control the triggering of phreatomagmatism.

**Plain Language Summary** The explosivity of volcanic eruptions may increase when magma interacts with groundwater (phreatomagmatism). On islands, freshwater reserves are strongly dependent on rainfall, which can be highly variable over time. Climate changes throughout Earth's history, resulted in periods with more or less rainfall and possibly, to variations in the abundance of water resources in certain regions over time. By combining past climate and volcanic records from Flores Island (Azores), we explore how fluctuations in groundwater may control the explosivity of volcanic eruptions on islands with abundant water resources. Previous studies revealed that Flores Island experienced multiple eruptions in the past 300,000 years, with phreatomagmatic eruptions occurring predominantly during the Holocene (last 11,700 years). Our results show a general correlation between long-term climate variations and eruption explosivity. However, short-term climatic variations during the Holocene cannot explain the occurrence or absence of water-magma interactions. Based on this, we conclude that on islands with abundant aquifers, explosive water-magma interactions are primarily controlled by the rate at which magma reaches the surface and, critically, the rate at which the magma recedes, that is, how rapidly the eruption ends.

## 1. Introduction

When ascending toward the surface, magma can encounter groundwater stored in aquifers or surface water in marine and inland environments. The thermal and hydrodynamic contact between magma and water may increase the efficiency of magma fragmentation (Németh & Kósik, 2020; White, 1996; Zimanowski et al., 2015), exacerbating volcanic explosivity. This is common in monogenetic basaltic volcanoes, which often experience changes between magmatic and phreatomagmatic activity (Martí et al., 2011; Planagumà et al., 2023; Zanon & Viveiros, 2019). Variable water-magma interactions throughout eruptions depend on a complex interplay between intrinsic factors (e.g., mass eruption rate, magma physico-chemical properties, conduit geometry) and environmental conditions (e.g., hydrological characteristics such as rainfall regime, infiltration rate, surface runoff, and groundwater occurrence) (e.g., Cassidy et al., 2018; Smith & Németh, 2017).



**Figure 1.** Geological and climatic settings of the study area. (a) Map of Flores Island showing the location of magmatic (scoria cones) and phreatomagmatic (tuff rings/maars) vents; CVS—Comprida Volcanic System and FVS—Funda Volcanic System; (b) Geotectonic setting of the Azores at the triple junction of the North American (NA), Eurasian (Eu), and Nubian (Nu) lithospheric plates (main morphotectonic structures: MAR—Mid-Atlantic Ridge, EAFZ—East Azores Fracture Zone, TR—Terceira Rift, and GF—Gloria Fault); (c) Map of the North Atlantic showing the main atmospheric and oceanic currents influencing the climate in the Azores (modified from Stewart et al. (2017) and Goslin et al. (2018)). Yellow arrows correspond to warm ocean surface currents. White dashed lines show the preferential position of the jet stream depending on the NAO phases. See Data Availability Statement section for elevation and bathymetry data sources.

Several studies have examined how these different factors influence magma-water interactions in different settings and timescales (e.g., Farquharson & Amelung, 2020; Foote et al., 2022; Kshirsagar et al., 2016; Nowell et al., 2006; Pedrazzi et al., 2022). Environmental controls have been particularly examined in the Transmexican Volcanic Belt, where long-term rainfall variability seems to have influenced the eruptive style of some monogenetic volcanoes (e.g., Agustín-Flores et al., 2021; Kshirsagar et al., 2015). Similar influences have been observed in other settings such as the Bakony-Balaton Highland Volcanic field (Kereszturi et al., 2011) and the Kamchatka Peninsula (Belousov, 2006). Despite its importance for accurate volcanic hazard forecasting, a detailed scientific understanding of how water-magma interactions respond to environmental changes remains limited. This is particularly relevant on volcanic islands, especially the ones with abundant water resources (e.g., Hawaii, Azores, west Pacific or Caribbean Islands). These islands often have a small size, isolated geography, and high sensitivity to climate, orographic and atmospheric variations (tropical or temperate, high and low altitudes, or under the influence of trade winds and westerlies) (Veron et al., 2019).

Volcanic islands have complex hydrogeological functioning that depends on several factors, such as the age and internal structure of the volcanoes, their size and topography, and recharge variability. Most exhibit a low-hydraulic-gradient basal groundwater body in equilibrium with the ocean, and inland upper dike-compartmented or perched aquifers (the “Hawaiian/La Reunion” model, Cruz, 2003; Peterson, 1972; Prada et al., 2005; Won et al., 2006). Others, follow the “Canary Islands” model, which considers a continuous basal aquifer with a hydraulic gradient that follows surface topography (Custódio, 1978, 2004).

Flores Island (Azores) is an ideal setting to study groundwater's role in modulating volcanism as it experienced multiple magmatic and phreatomagmatic monogenetic eruptions during the Holocene and Middle-Late Pleistocene. This is attested by the numerous scoria cones, tuff rings, and maars that shape the island's topography (Figure 1a). The recent stratigraphy of Flores also shows that small-volume and mildly explosive basaltic eruptions can rapidly shift into violent explosive events as a result of water-magma interactions (M. Andrade

et al., 2022, 2023). Flores' edifice contains abundant groundwater and surface water resources, as reflected by the abundance of springs, perennial streams, lakes, peatlands, and its dense vegetation. However, the Azorean hydroclimate displays a large inter-annual variability, which can result in fluctuations in surface and groundwater levels (Hernández et al., 2016, 2017; Sáez et al., 2025). Accordingly, humid periods would allegedly favor the generation of phreatomagmatism when compared to dry periods.

Through the integration of Flores' volcanic and paleohydroclimatic records (see Text S1 in Supporting Information S1), this study attempts to better understand how long-term and short-term (multimillennial to centennial timescales) rainfall variability may influence eruptive styles and consequently the hazard of volcanic eruptions. Moreover, it attempts to determine the dominant factors controlling shifts in magmatic/phreatomagmatic styles in the context of small-volume monogenetic eruptions at hydraulically charged ocean island volcanoes. This study corresponds to a first-order approach to the problem, paving the way for future studies on how climate inter-annual variability modulates volcanism.

## 2. Middle-Late Pleistocene and Holocene Volcanism

Flores is the westernmost island of the Azores Archipelago, which straddles the triple junction of the North American (where Flores sits), Eurasian, and Nubian lithospheric plates (Figure 1b). The island's volcanism dates back to 2.16 Ma and was characterized by major eruptive phases separated by long quiescence periods (50–400 kyr, Azevedo & Portugal Ferreira, 2006). At ~300 ka, a caldera collapse formed a depression in the center of the island, presently at 500 m asl (Hildenbrand et al., 2018). Locally known as *Caldeira das Sete Lagoas* (Figure 1a), this depression is currently filled with lavas and pyroclasts from monogenetic basaltic eruptions of Middle-Late Pleistocene and Holocene ages.

Flores' Holocene eruptive history includes six small-volume basaltic eruptions (Table S1 in Supporting Information S1, M. Andrade et al., 2021, 2022, 2023), Hawaiian and Strombolian in style, which frequently shifted to phreatomagmatic phases. They formed several tuff rings and maars 150 m to ~1 km wide (Figure 1a), most of which are currently occupied by lakes as deep as 114 m. Holocene volcanism was limited to a period of ~3,100 years (~3–6 ka), with the last four eruptions clustering within ~250 years. The recurrence rate is  $1.6 \times 10^{-3}$  eruptions/yr (estimated following Connor and Conway (2000)), however, during periods of clustered volcanic activity, recurrence intervals were <100 years.


Limited volcano-stratigraphic and geochronological data precludes a detailed reconstruction of Flores' Middle-Late Pleistocene eruptive history. Nonetheless, ~30 scoria cones and associated lava flows indicate widespread volcanism in the central uplands during this period (Figure 1a), with predominance of magmatic activity over phreatomagmatism. This is supported by the predominance of lava-flow successions throughout the exposed caldera-filling sequence (M. Andrade et al., 2023). Although ages of individual eruptions are unknown, they can be constrained to  $<314 \pm 30$  ka (K/Ar, Hildenbrand et al., 2018, Table S1 in Supporting Information S1).

## 3. Climatic and Hydrological Setting

Since the early Holocene, rainfall variability in the Azores has been mostly influenced by the NAO (Hernández et al., 2016), which is manifested by the anomaly between the Iceland sub-polar low and the Azores sub-tropical high-pressure poles (Figure 1c). Positive NAO phases shift westerlies northward, causing dry conditions in the Azores, while negative phases bring them south, increasing rainfall (Figure 1c). The Azores Current, a south-eastern branch of the Gulf Stream (Figure 1c), also influences the region by transporting warm equatorial waters to the polar regions (Klein & Siedler, 1989).

At Flores, monthly data sets of accumulated precipitation over 3 hydrological years show that rainfall prevails throughout all seasons. “Drier” months (June and July) experience significant precipitation of up to 108 mm/month, whereas rainier months record up to 861 mm/month, a value that is ~8 times higher (Figure S1a in Supporting Information S1).

The elevation of Flores and its steep topography forces westerlies' moist-rich air upwards, resulting in an uneven distribution of precipitation with relief (Figure S1b in Supporting Information S1). Rip fog (wet-fog-catch or hidden precipitation) captured by the cloud forest in the central uplands further exacerbates this effect. Consequently, rainfall is heavily skewed toward higher elevations, with mean precipitation values at the coast being 2–3 times lower than at the central uplands (Azevedo, 1998, Figure S1b in Supporting Information S1). Part of that



**Figure 2.** (a) Upper (semi-confined/unconfined perched aquifers at altitude) and Lower (fresh and salt water lenses close to sea-level) groundwater bodies delimitation at Flores with the distribution and discharge rates of the springs across the island, (b) preferential groundwater recharge zones delimitation, and (c) number of springs in the Upper unit per 100 m-altitude intervals and respective discharge rates. Green line extends from the minimum to maximum values and green squares mark the median value of discharge rates. Data from Secretaria Regional do Ambiente e do Mar (2022).


water ends up retained in crater lakes, which store  $\sim 1.83 \times 10^7 \text{ m}^3$  of water (C. Andrade et al., 2019), but also by bogs, fens, and peatlands (*Sphagnum*-dominated), which cover 17% of the island and store an estimated volume of water of  $2.2 \times 10^7 \text{ m}^3$  (Pereira et al., 2022). These reservoirs (Figure S2 in Supporting Information S1) regulate infiltration and ensure steady water release year-round, mitigating the effects of summer groundwater recharge reduction.

The Azorean Islands typically follow the “Hawaiian/La Reunion” model, with the perched-water bodies being frequently drained out by springs at higher elevations (Cruz, 2004; Cruz & Soares, 2018; Cruz et al., 2015). At Flores, two superimposing hydrogeologic units are present, the Upper and the Lower groundwater bodies (Figure 2a, Secretaria Regional do Ambiente e do Mar, 2022). Groundwater recharge in the Upper unit was estimated as  $1.54 \times 10^8 \text{ m}^3/\text{yr}$ , 75.5% of which occurs from October to March, being the lowest in July (5.9 mm) and the highest in January (173.6 mm) (Cruz et al., 2021). Estimations of the recharge rates are 27.2% and 31.3% for the Lower and Upper units, respectively. The recharge is higher in the central uplands, where rainfall is higher but also where recent, highly porous, volcanic products prevail (Figure 2b). In contrast, recharge is lower at the drier lowlands (<500 m asl), where less permeable, older geological formations crop out (Azevedo, 1998). The volume of water that is not infiltrated is drained over the island's surface, forming a dense drainage network composed of numerous perennial streams (Figure S1c in Supporting Information S1), with surface runoff estimated at  $1.94 \times 10^8 \text{ m}^3/\text{yr}$  (Cruz et al., 2021). The distribution of springs shows that even at high altitudes (>400 m) there are perched-aquifer discharges, some of which exhibit mean daily flow rates up to 200  $\text{m}^3/\text{day}$  (Figure 2c).

#### 4. Rainfall Variability Versus Volcanism

Present day hydrology of Flores Island, reveals substantial year-round groundwater storage in the uplands despite the important seasonal changes in precipitation. Over the Holocene, rainfall oscillations were stronger and followed trends of centennial to millennial timescales (Figure 3 and Figure S3 in Supporting Information S1). This, means that conditions of more/less rainfall persisted throughout longer periods, having therefore greater potential





**Figure 3.** Paleoclimate reconstructions for the North Atlantic over the Holocene. (a) Simulated mean annual precipitation at the Azores obtained from the global climate model AWI-ESM2 (Table S4, <https://doi.org/10.5281/zenodo.17079274>), (b) aridity index reconstructed using lake sediments from Pico Island, Azores (Björck et al., 2006), (c) NAO index reconstructed using a lake sediment record from southwestern Greenland (Olsen et al., 2012), (d) NAO index reconstruction based on southeastern Norwegian Sea sediment cores (Becker et al., 2020), and (e) precipitation variability based on North Iberian speleothems (Baldini et al., 2019). Red (magmatic) and blue (phreatomagmatic) lines mark the timing of Flores' Holocene eruptions.

to change groundwater storage in the uplands, and so, influence the eruptive style of volcanic eruptions in that area.

Of the six Holocene eruptions recorded at Flores, the three oldest were exclusively magmatic, whereas the most recent ones experienced phreatomagmatic activity (Figure 3). Available proxy-based data covering the timing of the oldest Flores' Holocene eruption (LB1-T1, 6,280 cal yr BP) suggest dry conditions in the North Atlantic at that time (Figures 3d and 3e). Albeit purely magmatic, the second Holocene eruption (LB1-T2, 4,990 cal yr BP) correlates with a humid period. This is attested by our model-based precipitation reconstruction and the proxy-based reconstruction by Björck et al. (2006) (Figures 3a and 3b), even though the NAO indexes obtained for other regions across the North Atlantic point to predominant positive phases (drier conditions, Figures 3c–3e).

The last four eruptions occurred during the Late Holocene, over a continuous dry period, as attested by our modeled precipitation for the Azores (Figure 3a), proxy-based climate reconstructions for North Iberia (Figure 3e), and by lake water level reconstructions across the Azores, which suggest a lowstand phase during this period (Sáez et al., 2025). Yet, our model shows that this period was characterized by a reduction in the precipitation anomalies when compared with the variations before and after, suggesting low amplitude changes of precipitation around the average value. Further climate reconstructions indicate, however, that this period was characterized by short-term (multidecadal/centennial) rainfall variability (Figures 3b–3d), with magmatic and phreatomagmatic eruptions coinciding with both dry and wet phases (Figure 3). The reconstruction by Olsen

et al. (2012) shows positive values of the NAO (low rainfall) during the purely magmatic FVS1 eruption and a negative peak (high rainfall) coincident with the phreatomagmatic FVS2 eruption (Figure 3c). However, this peak is brief and does not overlap with the following phreatomagmatic events (FVS3 and CVS), which occurred under drier conditions according to this reconstruction. The higher resolution NAO reconstruction by Becker et al. (2020) shows much more high-frequency variability during the time of FVS and CVS eruptions, with FVS1 (purely magmatic) and CVS (phreatomagmatic) eruptions coinciding with humid periods, and FVS2 and FVS3 phreatomagmatic eruptions with drier periods, pattern that is also observed in Björck et al. (2006) reconstruction. However, rainfall variability during this period is short-termed and smoother-than-usual, with the maximum and minimum peaks of the NAO index rarely exceeding 0.5 and  $-0.5$ , respectively, suggesting almost neutral NAO phases and therefore relatively stable rainfall conditions during the time of FVS and CVS eruptions.

At a broader scale, when compared with the last glacial cycle, the Holocene was a short, warm, and humid period, characterized by weak climatic oscillations (Figure S3 in Supporting Information S1). Over the last  $\sim 300$  ka (maximum age for the Middle-Late Pleistocene volcanism), the climate has been cooler and drier, with the only exception occurring at  $\sim 120$  ka during the Marine Isotope Stage 5e (Martin-Garcia, 2019; Martrat et al., 2007; Naughton et al., 2016). Despite the limited number of observations, and assuming that magmatic activity concentrated at approximately 0.4–0.2 ka (Azevedo & Portugal Ferreira, 2006, Figure S3 in Supporting Information S1), data suggests that magmatic eruptions have dominated during cold/dry periods (Middle-Late Pleistocene), while phreatomagmatic eruptions prevailed during warmer/wetter periods (the Holocene).

## 5. Discussion


Experimental (Valentine et al., 2014) and field (Planagumà et al., 2023) observations showed that most phreatomagmatic eruptions are sourced from explosions occurring  $<200$  m below the surface. Deeper explosions are usually contained (not erupting) even when magma rises through highly productive aquifers. The lack of drilled wells prevents the establishment of a detailed mapping of the Upper groundwater unit at Flores. However, our data on the spring's spatial distribution show the occurrence of highly productive perched aquifers at elevations greater than 400 m (Figure 2c). Considering that the topographic surface of the *Caldeira das Sete Lagoas* depression ranges around 500–600 m asl, this demonstrates the widespread potential for phreatomagmatism across the central uplands.

Flores topography has changed little during the Late Pleistocene and Holocene, meaning that the island's capacity to accommodate perched aquifers in its uplands remained similar in the last 300 ka. Since geological and hydrogeological conditions of the substrate did not change, the uneven distribution of phreatomagmatism over time (with more frequent phreatomagmatic eruptions during the Holocene) suggests that water reserves may have been lower during the Middle-Late Pleistocene, when climate was colder and drier.

Despite the predominance of phreatomagmatism in the Holocene when compared with the Middle-Late Pleistocene, this period experienced both magmatic and phreatomagmatic events (M. Andrade et al., 2021, 2022, 2023). Such variations cannot be explained by the low-magnitude climatic oscillations of the Holocene. For example, in the FVS area (Figure 1a), a Holocene volcanic edifice from a magmatic eruption is overlapped by a maar (M. Andrade et al., 2022), showing that geochemically and rheologically similar magmas ascending through the same aquifer-hosting volcanic sequence resulted in eruptions with distinct eruptive styles (Figure 1a). This, together with the no-correlation between eruptive styles and the Holocene short-term NAO variations, shows that characteristics of the substrate and hydrological changes, may have had a subordinate role in determining eruption explosivity on Flores during the Holocene.

As in other monogenetic volcanoes (e.g., Montsacopa or Orakei, Martí et al., 2011; Németh et al., 2012), water-magma interactions at Flores always occurred during the final phase of an eruption or during a later stage of a volcanic system's evolution, after it has experienced magmatic activity (M. Andrade et al., 2022, 2023). This shows that Flores' Holocene eruptions usually started as magmatic events but, later on, groundwater was able to enter the shallow conduit. The entrance of groundwater in the volcanic conduit is primarily controlled by the stress equilibrium between the magma and country rocks, which, in turn, is a function of the mass eruption rate (Houghton & Schmincke, 1989; Schliz-Antequera et al., 2024).

A high mass eruption rate ensures high magma overpressure in the conduit/eruptive fissure, which, additionally to a high thermal gradient with the country rock, eventually allows magma to cross an aquifer without significant



**Figure 4.** (a) Schematic diagram showing a conceptual model for water-magma interactions during small-volume basaltic eruptions with initial high-mass eruption rate as a function of the speed of decrease in mass eruption rate. Non-explosive water-magma interactions are suggested for rapid magma retreats, and explosive water-magma interactions for slow/gradual decreases in mass eruption rate; (b) efficiency of thermal energy transfer to mechanical energy, resulting from the interaction of magma with water as the mass ratio function of magma to water. The efficiency percentage controls the explosivity and magma fragmentation degree, and consequently the eruptive style (modified from Wohletz (1983)).

magma-water mingling (Gisbert et al., 2009; Figure 4a). However, when the eruption rate wanes, magma overpressure and thermal gradients decrease, allowing water percolation into the conduit, which will mix with the magma, potentially increasing the explosivity of the eruption (Figure 4). Experimental works suggest maximum kinetic energy for fuel:coolant mass ratios of  $\sim 3:1$  (Wohletz, 1983). Under these conditions, high-energy phreatomagmatic eruptions typically produce maars and tuff rings. In contrast, when larger volumes of water enter the conduit, water-magma interactions often result in lower-energy eruptions. These, commonly form wet and cold ( $<100^\circ\text{C}$ ) surges, originating tuff cone structures (Németh & Kósik, 2020; Wohletz & Sheridan, 1983; Figure 4). However, optimal fragmentation conditions require efficient magma-water mingling, which in the natural environment depends, upon many factors, on the length scale, system geometry, and fluid dynamics (White & Valentine, 2016). Therefore, although indicative, any direct assumptions derived from these ratios should be treated with caution, since intense phreatomagmatic explosions may occur at different ratios if the right mingling and triggering conditions are met.

The geological record shows that phreatomagmatic explosions at Flores were generally preceded by an increase in the extrusion of accidental lithics (M. Andrade et al., 2022, 2023), suggesting excavation of the conduit walls during periods of high-mass eruption rates (Gisbert et al., 2009; Sulpizio et al., 2005). We therefore suggest that the widening of the conduit during an initial pulse of high-mass eruption rate, followed by an eruption rate decrease (negative feedback), resulted in a decline in magma overpressure, as there was a wider conduit to accommodate a smaller volume of magma (Wadge, 1981). Consequently, when the magma fragmentation level falls below the existing water table, groundwater is able to enter the conduit, increasing the efficiency of magma fragmentation (Figure 4), triggering violent phreatomagmatic explosions with the generation of maar and tuff ring structures, as occurred at Flores (M. Andrade et al., 2022, 2023). The rate at which basaltic magma is discharged

may vary substantially during an eruption, and therefore, phreatomagmatism may occur at any eruption stage. However, for most basaltic eruptions, a rapid increase in magma discharges up to a maximum typically occurs at the beginning of the eruption, which is followed by a decay, sometimes exponential, over a longer period (Bonny & Wright, 2017; Wadge, 1981).

Hence, we also suggest that an exceedingly rapid drop in the mass eruption rate (associated with a decrease in magma overpressure) may instead lead to a rapid retreat of magma in the conduit, inhibiting phreatomagmatism (Figure 4a). In this scenario, large amounts of water may flood the conduit, resulting in water/magma mass ratios that are too high to generate phreatomagmatic fragmentation (Figure 4b). In these situations, the eruption may eventually end without generating phreatomagmatism. This might have been the case of the FVS1 eruption, since its geological record only shows a magmatic phase despite its close temporal and spatial proximity to major phreatomagmatic events, namely the FVS2 and FVS3 eruptions (M. Andrade et al., 2022).

Observations suggest that the eruptive style at Flores was influenced by long-term variations in the water availability. However, robust conclusions are limited by the small number of observations. Despite that, our findings are comparable with observations in São Miguel Island (eastern Azores), where a change in eruptive style from predominantly magmatic to predominantly phreatomagmatic is recorded at ~5 kyr ago (Guest et al., 1999; Queiroz et al., 2015; Wallenstein et al., 2015).

Phreatomagmatism has also been reported in many volcanic islands (e.g., Vanuatu, Samoa, New Zealand, Jeju, and Canary Islands), particularly along coastal areas or where rift zones meet the sea (Clarke et al., 2009; Go et al., 2017; Houghton & Nairn, 1991; Németh & Cronin, 2009). Studies that focus on phreatomagmatism triggered by groundwater high on island edifices are rarer, however, highlighting the importance of this study. Similarities to Flores are found, for example, in the observations by McPhie et al. (1990), White and Schmincke (1999), and Geshi et al. (2019) for Hawaii, La Palma and Miyakejima volcanoes, where these authors also attributed variations in the eruption rates to changes in eruptive style from magmatic to phreatomagmatic. These studies, however, focus on single eruptions and do not examine water/magma interactions in the same location, over multiple events, to isolate the parameters controlling water/magma interactions. As a first-order approach, our study provides a basis for further investigation in other volcanic islands worldwide.

## 6. Conclusions

Despite the relatively low number of eruptions recorded at Flores, our study shows that at hydraulically charged volcanic islands with high infiltration rates and perched aquifers within the first 200 m below the surface, magma-water interactions can be influenced by long-term, prominent changes in the rainfall regime (e.g., during glacial-interglacial transitions). At Flores dry/colder periods (Middle-Late Pleistocene) were dominated by magmatic eruptions, whereas phreatomagmatism became more frequent during wet/warm periods (Holocene).

In contrast, short-term, low-amplitude climatic fluctuations within the Holocene appear insufficient to explain changes in eruptive style. At hydraulically charged volcanic islands, groundwater reserves during periods of less rainfall are apparently sufficient for water-magma interactions to occur. At this shorter timescale, water-magma interactions seem instead to be primarily controlled by variations in eruption rates.

Our findings suggest that, under current conditions, future eruptions at Flores—or similar hydraulically charged volcanoes—would likely involve water-magma interactions. These interactions could increase eruptive explosiveness and unpredictability, with major implications for hazard assessment. This is particularly important in settings where past activity was dominated by small basaltic eruptions, which may cause underestimation of future hazards.

This study highlights the complexity of unraveling climate/volcanism interactions and how the availability of well-dated, high-resolution climate and volcanic reconstructions is critical to establish possible causal links between climate and eruptive style. Crucially, to obtain more robust correlations and better tackle the critical problem of climate/volcanism feedback systems, more studies across different settings and timescales are necessary.



## Data Availability Statement

The accumulated monthly precipitation used to analyze the temporal and spatial rainfall variability at Flores Island is from the Rede Hidrometeorológica dos Açores (Hydrometeorological Network of the Azores Regional Government), available on Tables S2 and S3 in Supporting Information S1. Analysis and graphical representation of accumulated monthly precipitation (Figure S1 in Supporting Information S1) were performed using R (v.4.2.0) (R Core Team, 2023), using the “ggplot2” package (Wickham, 2016). The R script used to plot the data is available in Text S2 in Supporting Information S1 and the R project with the script and data sets is available at Zenodo repository: M. Andrade et al. (2025) (<https://doi.org/10.5281/zenodo.17079274>). The earth system model used for paleoprecipitation reconstructions is available at <https://fesom.de/models/awi-esm/>. The resulting data is presented in Supporting Information S1 (Table S4) and is available at Zenodo repository: M. Andrade et al. (2025) (<https://doi.org/10.5281/zenodo.17079274>). The digital elevation model of Flores Island was generated from a 1:25000 scale digital altimetric database and purchased to CIGeoE (<https://www.igeoe.pt>). Bathymetry in Figure 1b from EMODnet Bathymetry Consortium (2018), <https://doi.org/10.12770/18ff0d48-b203-4a65-94a9-5fd8b0ec35f6>.

## Acknowledgments

MA and JSB are funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—509166013. MA acknowledges the financial support given by the Fundação para a Ciência e Tecnologia (FCT) through her doctoral grant (SFRH/BD/138261/2018). AH is funded by the Spanish Ministry of Science and Innovation through the Ramón y Cajal Scheme (RYC2020-029253-I). AP is supported by a CEEC Institutional contract funded by FCT (<https://doi.org/10.54499/CEECINST/00024/2021/CP2780/CT0003>). PL is supported by the Helmholtz “Changing Earth” program. We thank Alberto González Casarrubios for providing the Rscript to plot the rainfall data. We also acknowledge Dr. K. Németh and Dr. D. Pedrazzi for their review and comments that helped to improve the manuscript. Open Access funding enabled and organized by Projekt DEAL.

## References

- Agustín-Flores, J., Siebe, C., Ferrés, D., Sieron, K., & González-Zuccolotto, K. (2021). Monogenetic volcanoes with initial phreatomagmatic phases in the Cebroruco graben, western Mexico: The cases of Potrerillo I, Potrerillo II, and San Juanito. *Journal of Volcanology and Geothermal Research*, 412, 107184. <https://doi.org/10.1016/j.jvolgeores.2021.107184>
- Andrade, C., Cruz, J., Viveiros, F., & Coutinho, R. (2019). CO<sub>2</sub> flux from volcanic lakes in the western group of the Azores archipelago (Portugal). *Water*, 11(3), 599. <https://doi.org/10.3390/w11030599>
- Andrade, M., Hernández, A., Pimentel, A., Cruz, J., Ramos, A., Ludwig, P., et al. (2025). Controls on water-magma interactions at hydraulically-charged volcanic islands, supplementary information [Dataset]. Zenodo. <https://doi.org/10.5281/zenodo.17079274>
- Andrade, M., Pimentel, A., Ramalho, R., Kutterolf, S., & Hernández, A. (2022). The recent volcanism of Flores Island (Azores): Stratigraphy and eruptive history of Funda volcanic system. *Journal of Volcanology and Geothermal Research*, 432, 107706. <https://doi.org/10.1016/j.jvolgeores.2022.107706>
- Andrade, M., Ramalho, R., Pimentel, A., Kutterolf, S., & Hernández, A. (2023). The recent volcanism of Flores Island (Azores), Part II: Stratigraphy and eruptive history of Comprida volcanic system. *Journal of Volcanology and Geothermal Research*, 438, 107806. <https://doi.org/10.1016/j.jvolgeores.2023.107806>
- Andrade, M., Ramalho, R. S., Pimentel, A., Hernández, A., Kutterolf, S., Sáez, A., et al. (2021). Unraveling the Holocene eruptive history of Flores Island (Azores) through the analysis of lacustrine sedimentary records. *Frontiers in Earth Science*, 9, 738178. <https://doi.org/10.3389/feart.2021.738178>
- Azevedo, J. M. M. (1998). *Geologia e Hidrogeologia da Ilha das Flores*. (Doctoral Thesis). Universidade de Coimbra.
- Azevedo, J. M. M., & Portugal Ferreira, M. R. (2006). The volcanotectonic evolution of Flores Island, Azores (Portugal). *Journal of Volcanology and Geothermal Research*, 156(1–2), 90–102. <https://doi.org/10.1016/j.jvolgeores.2006.03.011>
- Baldini, L. M., Baldini, J. U. L., McDermott, F., Arias, P., Cueto, M., Fairchild, I. J., et al. (2019). North Iberian temperature and rainfall seasonality over the Younger Dryas and Holocene. *Quaternary Science Reviews*, 226, 105998. <https://doi.org/10.1016/j.quascirev.2019.105998>
- Becker, L. W. M., Sejrup, H. P., Hjelstuen, B. O., Haflidason, H., Kjennbakken, H., & Werner, J. P. (2020). Palaeo-productivity record from Norwegian Sea enables North Atlantic Oscillation (NAO) reconstruction for the last 8000 years. *npj Climate and Atmospheric Science*, 3(1), 42. <https://doi.org/10.1038/s41612-020-00147-6>
- Belousov, A. B. (2006). Distribution and eruptive mechanism of maars in the Kamchatka Peninsula. *Doklady Earth Sciences*, 406(1), 24–27. <https://doi.org/10.1134/S1028334X06010077>
- Björck, S., Rittenour, T., Rosén, P., França, Z., Möller, P., Snowball, I., et al. (2006). A Holocene lacustrine record in the central North Atlantic: Proxies for volcanic activity, short-term NAO mode variability, and long-term precipitation changes. *Quaternary Science Reviews*, 25(1–2), 9–32. <https://doi.org/10.1016/j.quascirev.2005.08.008>
- Bonny, E., & Wright, R. (2017). Predicting the end of lava flow-forming eruptions from space. *Bulletin of Volcanology*, 79(7), 1–13. <https://doi.org/10.1007/s00445-017-1134-8>
- Cassidy, M., Manga, M., Cashman, K., & Bachmann, O. (2018). Controls on explosive-effusive volcanic eruption styles. *Nature Communications*, 9(1), 2839. <https://doi.org/10.1038/s41467-018-05293-3>
- Clarke, H., Troll, V. R., & Carracedo, J. C. (2009). Phreatomagmatic to strombolian eruptive activity of basaltic cinder cones: Montaña Los Erales, Tenerife, Canary Islands. *Journal of Volcanology and Geothermal Research*, 180(2–4), 225–245. <https://doi.org/10.1016/j.jvolgeores.2008.11.014>
- Connor, C. B., & Conway, F. M. (2000). Basaltic volcanic fields. In *Encyclopedia of volcanoes* (pp. 331–343). Academic Press.
- Cruz, J. V. (2003). Groundwater and volcanoes: Examples from the Azores archipelago. *Environmental Geology*, 44(3), 343–355. <https://doi.org/10.1007/s00254-003-0769-2>
- Cruz, J. V. (2004). Ensaio sobre a água subterrânea nos Açores. In *História, ocorrência e qualidade* (p. 288). SRA.
- Cruz, J. V., Fontiela, J., Prada, S., & Andrade, C. (2015). The chemical status of groundwater and pollution risk in the Azores archipelago (Portugal). *Environmental Earth Sciences*, 73(6), 2763–2777. <https://doi.org/10.1007/s12665-014-3407-2>
- Cruz, J. V., Melo, C., Costa, S., Brito de Azevedo, E., Andrade, C., Rodrigues, M. C., et al. (2021). *HIDROBAL - Avaliação e espacialização do balanço hídrico e caracterização da interação entre as águas de superfície e subterrâneas. Relatório Final* (p. 104). DROTRH.
- Cruz, J. V., & Soares, N. (2018). Groundwater governance in the Azores Archipelago (Portugal): Valuing and protecting a strategic resource in small islands. *Water*, 10(4), 408. <https://doi.org/10.3390/w10040408>
- Custódio, E. (1978). *Geohidrologia de terrenos e ilhas vulcânicas [Hydrogeology of volcanic terrains and islands]*. Centro de Estudos Hidrográficos and Instituto de Hidrologia.
- Custódio, E. (2004). Hydrogeology of volcanic rocks. In *Hydrogeology of volcanic rocks* (pp. 395–425). UNESCO.

- EMODnet Bathymetry Consortium. (2018). EMODnet digital bathymetry (DTM 2018). <https://doi.org/10.12770/18ff0d48-b203-4a65-94a9-5fd8b0ec35f6>
- Farquharson, J. I., & Amelung, F. (2020). Extreme rainfall triggered the 2018 rift eruption at Kīlauea Volcano. *Nature*, 580(7804), 491–495. <https://doi.org/10.1038/s41586-020-2172-5>
- Foot, A., Németh, K., & Handley, H. (2022). The interplay between environmental and magmatic conditions in eruption style transitions within a fissure-aligned monogenetic volcanic system of Auckland, New Zealand. *Journal of Volcanology and Geothermal Research*, 431, 107652. <https://doi.org/10.1016/j.jvolgeores.2022.107652>
- Geshi, N., Németh, K., Noguchi, R., & Oikawa, T. (2019). Shift from magmatic to phreatomagmatic explosions controlled by the lateral evolution of a feeder dike in the Suoana-Kazahaya eruption, Miyakejima Volcano, Japan. *Earth and Planetary Science Letters*, 511, 177–189. <https://doi.org/10.1016/j.epsl.2019.01.038>
- Gisbert, G., Gimeno, D., & Fernandez-Turiel, J. L. (2009). Eruptive mechanisms of the Puig de La Garrinada volcano (Olot, Garrotxa volcanic field, Northeastern Spain): A methodological study based on proximal pyroclastic deposits. *Journal of Volcanology and Geothermal Research*, 180(2–4), 259–276. <https://doi.org/10.1016/j.jvolgeores.2008.12.018>
- Go, S. Y., Kim, G. B., Jeong, J. O., & Sohn, Y. K. (2017). Diatreme evolution during the phreatomagmatic eruption of the Songaksan Tuff Ring, Jeju Island, Korea. *Bulletin of Volcanology*, 79(3), 23. <https://doi.org/10.1007/s00445-017-1103-2>
- Goslin, J., Fruergaard, M., Sander, L., Galka, M., Menviel, L., Monkenbusch, J., et al. (2018). Holocene centennial to millennial shifts in North-Atlantic storminess and ocean dynamics. *Scientific Reports*, 8(1), 12778. <https://doi.org/10.1038/s41598-018-29949-8>
- Guest, J. E., Gaspar, J. L., Cole, P. D., Queiroz, G., Duncan, A. M., Wallenstein, N., et al. (1999). Volcanic geology of Furnas volcano, São Miguel, Azores. *Journal of Volcanology and Geothermal Research*, 92(1–2), 1–29. [https://doi.org/10.1016/S0377-0273\(99\)00064-5](https://doi.org/10.1016/S0377-0273(99)00064-5)
- Hernández, A., Kutiel, H., Trigo, R. M., Valente, M. A., Sigró, J., Cropper, T., & Santo, F. E. (2016). New Azores archipelago daily precipitation dataset and its links with large-scale modes of climate variability. *International Journal of Climatology*, 36(12), 4439–4454. <https://doi.org/10.1002/joc.4642>
- Hernández, A., Sáez, A., Bao, R., Raposeiro, P. M., Trigo, R. M., Doolittle, S., et al. (2017). The influences of the AMO and NAO on the sedimentary infill in an Azores Archipelago lake since ca. 1350 CE. *Global and Planetary Change*, 154, 61–74. <https://doi.org/10.1016/j.gloplacha.2017.05.007>
- Hildenbrand, A., Marques, F. O., & Catalão, J. (2018). Large-scale mass wasting on small volcanic islands revealed by the study of Flores Island (Azores). *Scientific Reports*, 8(1), 13898. <https://doi.org/10.1038/s41598-018-32253-0>
- Houghton, B. F., & Nairn, I. A. (1991). The 1976–1982 Strombolian and phreatomagmatic eruptions of White Island, New Zealand: Eruptive and depositional mechanisms at a “wet” volcano. *Bulletin of Volcanology*, 54(1), 25–49. <https://doi.org/10.1007/BF00278204>
- Houghton, B. F., & Schmincke, H.-U. (1989). Rothenberg scoria cone, East Eifel: A complex Strombolian and phreatomagmatic volcano. *Bulletin of Volcanology*, 52(1), 28–48. <https://doi.org/10.1007/BF00641385>
- Kereszturi, G., Németh, K., Csillag, G., Balogh, K., & Kovács, J. (2011). The role of external environmental factors in changing eruption styles of monogenetic volcanoes in a Mio/Pleistocene continental volcanic field in western Hungary. *Journal of Volcanology and Geothermal Research*, 201(1–4), 227–240. <https://doi.org/10.1016/j.jvolgeores.2010.08.018>
- Klein, B., & Siedler, G. (1989). On the origin of the Azores current. *Journal of Geophysical Research*, 94(C5), 6159–6168. <https://doi.org/10.1029/JC094iC05p06159>
- Kshirsagar, P., Siebe, C., Guilbaud, M. N., & Salinas, S. (2016). Geological and environmental controls on the change of eruptive style (phreatomagmatic to Strombolian-effusive) of Late Pleistocene El Caracol tuff cone and its comparison with adjacent volcanoes around the Zacapu basin (Michoacán, México). *Journal of Volcanology and Geothermal Research*, 318, 114–133. <https://doi.org/10.1016/j.jvolgeores.2016.03.015>
- Kshirsagar, P., Siebe, C., Guilbaud, M. N., Salinas, S., & Layer, P. W. (2015). Late Pleistocene Alberca de Guadalupe maar volcano (Zacapu basin, Michoacán): Stratigraphy, tectonic setting, and paleo-hydrogeological environment. *Journal of Volcanology and Geothermal Research*, 304, 214–236. <https://doi.org/10.1016/j.jvolgeores.2015.09.003>
- Martí, J., Planagumà, L., Geyer, A., Canal, E., & Pedrazzi, D. (2011). Complex interaction between Strombolian and phreatomagmatic eruptions in the quaternary monogenetic volcanism of the Catalan Volcanic Zone (NE of Spain). *Journal of Volcanology and Geothermal Research*, 201(1–4), 178–193. <https://doi.org/10.1016/j.jvolgeores.2010.12.009>
- Martin-Garcia, G. M. (2019). Oceanic impact on European climate changes during the quaternary. *Geosciences*, 9(3), 119. <https://doi.org/10.3390/geosciences9030119>
- Martrat, B., Grimalt, J. O., Shackleton, N. J., de Abreu, L., Hutterli, M. A., & Stocker, T. F. (2007). Four climate cycles of recurring deep and surface water destabilizations on the Iberian margin. *Science*, 317(5837), 502–507. <https://doi.org/10.1126/science.1139994>
- McPhie, J., Walker, G. P., & Christiansen, R. L. (1990). Phreatomagmatic and phreatic fall and surge deposits from explosions at Kīlauea volcano, Hawaii, 1790 AD: Kēanākakoi ash member. *Bulletin of Volcanology*, 52(5), 334–354. <https://doi.org/10.1007/BF00302047>
- Naughton, F., Goñi, M. S., Rodrigues, T., Salgueiro, E., Costas, S., Desprat, S., et al. (2016). Climate variability across the last deglaciation in NW Iberia and its margin. *Quaternary International*, 414, 9–22. <https://doi.org/10.1016/j.quaint.2015.08.073>
- Németh, K., & Cronin, S. J. (2009). Phreatomagmatic volcanic hazards where rift-systems meet the sea, a study from Ambae Island, Vanuatu. *Journal of Volcanology and Geothermal Research*, 180(2–4), 246–258. <https://doi.org/10.1016/j.jvolgeores.2008.08.011>
- Németh, K., Cronin, S. J., Smith, I. E. M., & Agustin Flores, J. (2012). Amplified hazard of small-volume monogenetic eruptions due to environmental controls, Orakei Basin, Auckland Volcanic Field, New Zealand. *Bulletin of Volcanology*, 74(9), 2121–2137. <https://doi.org/10.1007/s00445-012-0653-6>
- Németh, K., & Kósik, S. (2020). Review of explosive hydrovolcanism. *Geosciences*, 10(2), 44. <https://doi.org/10.3390/geosciences10020044>
- Nowell, D. A., Jones, M. C., & Pyle, D. M. (2006). Episodic quaternary volcanism in France and Germany. *Journal of Quaternary Science: Published for the Quaternary Research Association*, 21(6), 645–675. <https://doi.org/10.1002/jqs.1005>
- Olsen, J., Anderson, N. J., & Knudsen, M. F. (2012). Variability of the North Atlantic oscillation over the past 5,200 years. *Nature Geoscience*, 5(11), 808–812. <https://doi.org/10.1038/ngeo1589>
- Pedrazzi, D., Cerda, D., Geyer, A., Martí, J., Aulinas, M., & Planagumà, L. (2022). Stratigraphy and eruptive history of the complex Puig de La Banya del Boc monogenetic volcano, Garrotxa Volcanic Field. *Journal of Volcanology and Geothermal Research*, 423, 107460. <https://doi.org/10.1016/j.jvolgeores.2021.107460>
- Pereira, D., Mendes, C., & Dias, E. (2022). The potential of peatlands in global climate change mitigation: A case study of Terceira and Flores Islands (Azores, Portugal) hydrologic services. *SN Applied Sciences*, 4(6), 184. <https://doi.org/10.1007/s42452-022-05066-0>
- Peterson, F. L. (1972). Water development on tropic volcanic islands-type example: Hawaii. *Ground Water*, 10(5), 18–23. <https://doi.org/10.1111/j.1745-6584.1972.tb03586.x>

- Planagumà, L., Bolós, X., & Martí, J. (2023). Hydrogeologic and magmatic controls on phreatomagmatism at the La Garrotxa monogenetic volcanic field (NE of Iberian Peninsula). *Journal of Volcanology and Geothermal Research*, 441, 107894. <https://doi.org/10.1016/j.jvolgeores.2023.107894>
- Prada, S. N., da Silva, M. O., & Cruz, J. V. (2005). Groundwater behaviour in Madeira, Volcanic Island (Portugal). *Hydrogeology Journal*, 13(5–6), 800–812. <https://doi.org/10.1007/s10040-005-0448-3>
- Queiroz, G., Gaspar, J. L., Guest, J. E., Gomes, A., & Almeida, M. H. (2015). Chapter 7 eruptive history and evolution of Sete Cidades volcano, São Miguel Island, Azores. *Geological Society, London, Memoirs*, 44(1), 87–104. <https://doi.org/10.1144/M44.7>
- R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Sáez, A., Hernández, A., Pimentel, A., Andrade, M., Bao, R., Raposeiro, P. M., et al. (2025). Westerlies migrations and volcanic records over the past 4000 years from the Azores lacustrine sequences. Exploring correlations and impacts on Western Europe. *Global and Planetary Change*, 246, 104698. <https://doi.org/10.1016/j.gloplacha.2025.104698>
- Schliz-Antequera, M., Siebe, C., Salinas, S., & Lerner, G. A. (2024). An inventory of phreatomagmatic volcanoes in the trans-Mexican Volcanic Belt. *Journal of Volcanology and Geothermal Research*, 452, 108136. <https://doi.org/10.1016/j.jvolgeores.2024.108136>
- Secretaria Regional do Ambiente e do Mar. (2022). Plano de Gestão da Região Hidrográfica dos Açores RH9 – Caracterização da situação de referência e diagnóstico. Technical Report (Vol. 8).
- Smith, I. E. M., & Németh, K. (2017). Source to surface model of monogenetic volcanism: A critical review. *Science Progress*, 446, 1–28. <https://doi.org/10.1144/SP446.14>
- Stewart, H., Bradwell, T., Bullard, J., Davies, S. J., Golledge, N., & McCulloch, R. D. (2017). 8000 years of North Atlantic storminess reconstructed from a Scottish peat record: Implications for Holocene atmospheric circulation patterns in western Europe. *Journal of Quaternary Science*, 32(8), 1075–1084. <https://doi.org/10.1002/jqs.2983>
- Sulpizio, R., Mele, D., Dellino, P., & Volpe, L. L. (2005). A complex, Subplinian-type eruption from low-viscosity, phonolitic to tephri-phonolitic magma: The AD 472 (Pollena) eruption of Somma-Vesuvius, Italy. *Bulletin of Volcanology*, 67(8), 743–767. <https://doi.org/10.1007/s00445-005-0414-x>
- Valentine, G. A., Graettinger, A. H., & Sonder, I. (2014). Explosion depths for phreatomagmatic eruptions. *Geophysical Research Letters*, 41(9), 3045–3051. <https://doi.org/10.1002/2014GL060096>
- Veron, S., Mouchet, M., Govaerts, R., Haevermans, T., & Pellens, R. (2019). Vulnerability to climate change of islands worldwide and its impact on the tree of life. *Scientific Reports*, 9(1), 14471. <https://doi.org/10.1038/s41598-019-51107-x>
- Wadge, G. (1981). The variation of magma discharge during basaltic eruptions. *Journal of Volcanology and Geothermal Research*, 11(2–4), 139–168. [https://doi.org/10.1016/0377-0273\(81\)90020-2](https://doi.org/10.1016/0377-0273(81)90020-2)
- Wallenstein, N., Duncan, A., Guest, J. E., & Almeida, M. H. (2015). Eruptive history of Fogo volcano. São Miguel, Azores (Vol. 44, No. (1), pp. 105–123). <https://doi.org/10.1144/M44.8>
- White, J. D. (1996). Impure coolants and interaction dynamics of phreatomagmatic eruptions. *Journal of Volcanology and Geothermal Research*, 74(3–4), 155–170. [https://doi.org/10.1016/S0377-0273\(96\)00061-3](https://doi.org/10.1016/S0377-0273(96)00061-3)
- White, J. D., & Schmincke, H. U. (1999). Phreatomagmatic eruptive and depositional processes during the 1949 eruption on La Palma (Canary Islands). *Journal of Volcanology and Geothermal Research*, 94(1–4), 283–304. [https://doi.org/10.1016/S0377-0273\(99\)00108-0](https://doi.org/10.1016/S0377-0273(99)00108-0)
- White, J. D., & Valentine, G. A. (2016). Magmatic versus phreatomagmatic fragmentation: Absence of evidence is not evidence of absence. *Geosphere*, 12(5), 1478–1488. <https://doi.org/10.1130/GES01337.1>
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag. Retrieved from <https://ggplot2.tidyverse.org>
- Wohletz, K. H. (1983). Mechanisms of hydrovolcanic pyroclast formation: Grain-size, scanning electron microscopy, and experimental studies. *Journal of Volcanology and Geothermal Research*, 17(1–4), 31–63. [https://doi.org/10.1016/0377-0273\(83\)90061-6](https://doi.org/10.1016/0377-0273(83)90061-6)
- Wohletz, K. H., & Sheridan, M. F. (1983). Hydrovolcanic explosions. II. Evolution of basaltic tuff rings and tuff cones. *American Journal of Science*, 283(5), 385–413. <https://doi.org/10.2475/ajs.283.5.385>
- Won, J. H., Lee, J. Y., Kim, J. W., & Koh, G. W. (2006). Groundwater occurrence on Jeju Island, Korea. *Hydrogeology Journal*, 14(4), 532–547. <https://doi.org/10.1007/s10040-005-0447-4>
- Zanon, V., & Viveiros, F. (2019). A multi-methodological re-evaluation of the volcanic events during the 1580 CE and 1808 eruptions at São Jorge Island (Azores Archipelago, Portugal). *Journal of Volcanology and Geothermal Research*, 373, 51–67. <https://doi.org/10.1016/j.jvolgeores.2019.01.028>
- Zimanowski, B., Büttner, R., Dellino, P., White, J. D. L., & Wohletz, K. H. (2015). Magma–water interaction and phreatomagmatic fragmentation. In *The encyclopedia of volcanoes* (pp. 473–484). Elsevier. <https://doi.org/10.1016/B978-0-12-385938-9.00026-2>

## References From the Supporting Information

- Berger, A. (1978). Long-term variations of daily insolation and quaternary climatic changes. *Journal of the Atmospheric Sciences*, 35(12), 2362–2367. [https://doi.org/10.1175/1520-0469\(1978\)035<2362:ltvodi>2.0.co;2](https://doi.org/10.1175/1520-0469(1978)035<2362:ltvodi>2.0.co;2)
- Danilov, S., Sidorenko, D., Wang, Q., & Jung, T. (2017). The finite-volume sea ice–ocean model (FESOM2). *Geoscientific Model Development*, 10(2), 765–789. <https://doi.org/10.5194/gmd-10-765-2017>
- Köhler, P., Nehrbaas-Ahles, C., Schmitt, J., Stocker, T. F., & Fischer, H. (2017). Continuous records of the atmospheric greenhouse gases CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O and their radiative forcing since the penultimate glacial maximum. *EPIC Repository*. Retrieved from <https://epic.awi.de/id/eprint/44527/>
- Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, Z., Yang, L. W. M. J., & Merchant, J. W. (2000). Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data. *International Journal of Remote Sensing*, 21(6–7), 1303–1330. <https://doi.org/10.1080/014311600210191>
- Moreno, J., Ramos, A. M., Raposeiro, P. M., Santos, R. N., Rodrigues, T., Naughton, F., et al. (2023). Identifying imprints of externally derived dust and halogens in the sedimentary record of an Iberian alpine lake for the past~ 13,500 years–Lake Peixão, Serra da Estrela (Central Portugal). *Science of the Total Environment*, 903, 166179. <https://doi.org/10.1016/j.scitotenv.2023.166179>
- Raddatz, T. J., Reick, C. H., Knorr, W., Kattge, J., Roeckner, E., Schnur, R., et al. (2007). Will the tropical land biosphere dominate the climate–carbon cycle feedback during the twenty-first century? *Climate Dynamics*, 29(6), 565–574. <https://doi.org/10.1007/s00382-007-0247-8>
- Repschläger, J., Garbe-Schönberg, D., Weinelt, M., & Schneider, R. (2017). Holocene evolution of the North Atlantic subsurface transport. *Climate of the Past*, 13(4), 333–344. <https://doi.org/10.5194/cp-13-333-2017>

- Schwab, C., Kinkel, H., Weinelt, M., & Repschläger, J. (2012). Coccolithophore paleoproductivity and ecology response to deglacial and Holocene changes in the Azores current system: AZF and PP dynamics DURING last 16 kyr. *Paleoceanography*, 27(3), PA3210. <https://doi.org/10.1029/2012PA002281>
- Sidorenko, D., Goessling, H. F., Koldunov, N. V., Scholz, P., Danilov, S., Barbi, D., et al. (2019). Evaluation of FESOM2.0 coupled to ECHAM6.3: Preindustrial and HighResMIP simulations. *Journal of Advances in Modeling Earth Systems*, 11(11), 3794–3815. <https://doi.org/10.1029/2019MS001696>