

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Ahmed, D. Vitsupakorn, K. D. Hartwell, K. Albalawi, P. Rizkallah, P. Watson and D. D. Jones, *Chem. Sci.*, 2025, DOI: 10.1039/D5SC06589E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Chromophore charge-state switching through copper-dependent homodimerisation of an engineered green fluorescent protein.

Rochelle D Ahmed^{a#}, Danoo Vitsupakorn^{a#}, Kieran D. Hartwell^a, Karma Albalawi^{a,c}, Pierre J. Rizkallah^b, Peter D. Watson^a and D. Dafydd Jones^{a*}.

a. School of Biosciences, Molecular Biosciences Division, Cardiff University, Sir Martin Evans Building, Cardiff, CF10 3AX, UK.

b. School of Medicine, Cardiff University, Cardiff, CF14 4XN, UK.

c. Department of Chemistry, Faculty of Science, University of Tabuk, Tabuk, Saudi Arabia.

* Corresponding author. D. Dafydd Jones, School of Biosciences, Sir Martin Evans Building, Cardiff University, Cardiff CF10 3AX, UK. Email: jonesdd@cardiff.ac.uk

These authors contributed equally.

ORCID

RDA: 000-0002-8714-0297

DV: 0000-0001-8000-4121

KDH: 0009-0004-6613-2708

KA: 0000-0002-3932-0381

PJR: 0000-0002-9290-0369

PDW: 0000-0003-0250-7852

DDJ: 0000-0001-7709-3995



Abstract.

View Article Online
DOI: 10.1039/D5SC06589E

Here, we have linked one of the most common protein-protein interaction events, homodimerisation, to an essential trace metal, copper, through engineering green fluorescent protein. Mutation of H148 to cysteine promotes the neutral phenolic chromophore in the monomer that excites predominantly at ~400 nm. Homodimerisation via a copper-dependent disulphide bridge, switches the chromophore to the charged phenolate that excites at ~490 nm. The result is ~30 fold increase in the fluorescence emission ratio. Homo-dimerisation kinetics are further improved by optimising the sfGFP homodimer interface, generating the variant termed GFP-diS2. Structures of the monomeric and dimeric GFP-diS2 suggests charge switching is through peptide bond flipping and the formation of a buried organised water network around the chromophore that span the interface region. Fusion to a leucine zipper protein dimerisation element greatly increased GFP-diS2 association rate making it a more effective copper sensor in vitro and in vivo with Cu(I) instigating the signal change quicker and at lower ion concentrations than Cu(II). Thus, GFP-diS2 provides the framework for generating a sensitive genetically encoded copper sensor and will eventually be adapted to monitor one of the most important protein-protein interactions in biology, homo-oligomerisation.



Introduction

Fluorescent proteins (FPs) have revolutionised how researchers investigate biological processes^{1–3}. Originally used as passive tags simply to fuse to a protein of interest, there has been a shift to monitoring active and dynamic processes that report on multi-component events^{3–5}. Protein-metal and protein-protein interactions (PPIs) are two such areas. Metals play a critical role in biology with an estimated 30–50% of proteins dependent on metals for their structure and/or function^{6,7}. Copper is a particularly important trace metal in biology having roles ranging from enzyme catalysis, to signalling, to redox processes^{8–11}. Due to its role in the formation of reactive oxygen species, free copper ion concentration is tightly regulated in biological systems¹¹. Given its importance to biology, there has been a push to develop molecular sensors capable of detecting copper^{12,13}. While small molecule synthetic copper sensors have been developed with ultra-high affinity (sub pM¹⁴) they have drawbacks with the most significant being the need to apply them exogenously and to be delivered into cells. This in turn limits their true in situ application. Thus, the development of genetically encoded copper-responsive fluorescent proteins that are directly synthesised by the cell is of significant interest as they can be used in a wide variety of organisms and be targeted to specific locations in the cell^{5,12,13}. However, many current copper sensors have small responses, lack metal ion selectivity or rely on a negative signal response (loss of fluorescence signal) as part of the detection process^{12,15–17}.

PPIs underpin a huge number of biological processes critical to life ranging from gene transcription to energy production, to signal transduction^{18–22}, with PPI dysregulation contributing to many disease states^{21,23}. Indeed, a protein is more likely to exist as part of a complex rather than act alone as a simple monomeric unit, with homodimers being the most commonly observed structural unit in the protein data bank (PDB)^{18,19}. PPI interactions have been used as the basis for constructing FP-based sensors^{3,24,25}, including metal ions¹³, through proximity-based processes such as Förster resonance energy transfer (FRET)²⁶ and fragment complementation (BiC)²⁷. These powerful and popular approaches have several drawbacks. One is the requirement for two different components, so restricting analysis to hetero-oligomerisation events thus missing out on frequently observed and equally important homo-oligomerisation.



The chromophore of the original FP, *Aequorea victoria* GFP (avGFP), exists in two forms: the neutral phenolic state (CRO-OH) with an excitation maximum at ~400 nm and the charged phenolate CRO-O⁻ form with an excitation maximum at ~490 nm^{1,28}. We have recently shown that the superfolder version of GFP (sfGFP)²⁹ can be engineered to change its spectral properties by switching the charged state on the chromophore by mutating H148³⁰, a residue that stabilises the chromophore phenolate anion in the ground state^{31–33}. When we replaced H148 by mutually reactive non-canonical amino acids, namely an azide group in one sfGFP and cycloalkyne in a second sfGFP, the CRO-OH state dominated in both^{34,35}. When the two sfGFP “clicked” together to form a pseudo-homodimer there was a spectral shift with the CRO-O⁻ dominating and with the new dimeric form being brighter per monomer unit than sfGFP³⁴. While the basic premise of GFP monomer association provides a sensing mechanism, each sfGFP monomer unit should be chemically identical and not require genetic code reprogramming to incorporate a non-canonical amino acid.

Using only canonical amino acids we engineered sfGFP so that its spectral properties change on homodimerisation in a copper-dependent manner. By incorporating a cysteine at residue 148 together with a mutation that stabilises the dimer interface, we engineered sfGFP to switch to a highly fluorescent “on” dimeric state (Fig.1a). The dimer complex is 2.9 fold brighter than the starting wild-type monomeric sfGFP and has an up to 31 fold ratiometric switch. Structures of the monomeric and dimeric states reveals a mechanism for switching based on a peptide bond flip and burial of critical water molecules central to GFP fluorescence. We also show that when coupled to naturally homodimerizing proteins, the copper-dependent kinetics of the engineered sfGFP increase, with the Cu⁺ being the preferred dimerisation promoting ion. A leucine zipper fused system allowed the newly engineered sfGFP to sense and respond to copper in bacterial cells.






Fig.1. The effect of introducing cysteine residues at residue 148. (a) Overview of the homodimerization strategy. The starting monomers are dim with the CRO-OH form dominating but homodimerisation via a copper-dependent disulphide molecular bolt results in a brighter system through promotion of CRO-O⁻ state. (b) Modelling of the H148C mutation using short timescale molecular dynamics³⁰. The wt sfGFP is shown as green sticks with a H-bond between H148 and CRO shown as a dashed line. The H148C model is shown as grey sticks. Absorbance spectra (c) and fluorescence emission spectra (d) of wild-type sfGFP (H148, black), mGFP-diS1 (red)³⁰ and mGFP-diS2 (green). In (d), the dashed line represents emission on excitation at 400 nm and full lines emission on excitation at 488-492 nm.

Results and Discussion

Switching the monomer chromophore protonation state.

Modulating the charged state of the GFP chromophore phenol group provides a means to sense biological processes through changes in electronic excitation^{32–35}. In the starting wild-type sfGFP, the deprotonated CRO-O⁻ phenolate dominates over the CRO-OH, with H148 capable of forming a H-bond with the chromophore phenolate group that stabilises the CRO-O⁻ form^{32–35} (Fig. 1b). Modelling previously suggested that mutating H148 to cysteine should remove the interaction with the



chromophore³⁰; the weak Lewis acid behaviour of the cystine thiol group together with larger atomic radius of sulphur sterically blocks a conformation allowing H-bonding with the CRO and exposing the thiol group to the solvent (Fig. 1b). Replacing H148 with cysteine generates the variant termed GFP-diS1, which remains monomeric after purification (Fig. S1). Absorbance spectra shows that the major excitation peak is now 400 nm with a minor peak at 498 nm indicating a shift towards the CRO-OH state. The same is mirrored in fluorescence spectra with emission on excitation at 400 nm dominating over that on excitation at around 490 nm (Fig. 1c and Table 1). Overall, the ratio of emission on excitation at wavelengths equivalent to the CRO-OH (~400 nm) or CRO-O⁻ (~490 nm) state (here on termed the $\text{Ex}^{\text{CRO-O}^-}:\text{Ex}^{\text{CRO-OH}}$ ratio) switches from 7:1 in sfGFP to 1:4 in mGFP-diS1 (prefix m refers to monomeric).

Variant	λ_{max} (nm)	λ_{EM} (nm)	ϵ ($\text{M}^{-1}\text{cm}^{-1}$)	$\Phi^{[c]}$ (%)	Brightness ($\text{M}^{-1}\text{cm}^{-1}$)
mGFP-diS1	400	509	34,000	N/D	N/D
	498	513	11,100	50	5,550
dGFP-diS1 ^[a]	400	509	11,400	N/D	N/D
	488	509	142,600	64	91,264
mGFP-diS2	400	511	27,500	N/D	N/D
	496	512	13,000	49	6,370
dGFP-diS2 ^[a]	400	510	10,000	N/D	N/D
	492	511	147,600	72	106,272
GFP-diS2-ZIP (no Cu ²⁺)	400	510	55,000	N/D	N/D
	496	514	22,400	78	17,472
GFP-diS2-ZIP (plus Cu ²⁺)	400	510	17,700	N/D	N/D
	493	511	188,400	85	160,140
sfGFP ^{WT} [a,b]	400	508	8000	N/D	N/D
	485	511	49,000	75	36,750

Table 1. Spectral properties of sfGFP variants and dimers. [a] Excitation at 400 nm was negligible and so quantum yield was not determined. [b] Value differs from that described in Pedelacq et al²⁹ but similar to those reported by Reddington et al³² and Cranfill et al³⁶. [c] quantum yield.



Copper-dependent dimerisation of GFP-diS1

Incubation of sfGFP-H148C mutants (termed GFP-diS1 from here on in) under aerobic conditions at high protein concentration ($>50 \mu\text{M}$) over a prolonged period ($>5 \text{ hr}$) did not result in the formation of significant amounts of the dimer (Fig. S1b); the absorbance spectra retained the characteristics shown in Fig. 1c. This suggests that the cysteine thiol group cannot form the disulphide bond between individual GFP-diS1 monomers. It has previously been reported that Cu^{2+} can aid in the formation of disulphide bridges³⁷. On incubation with CuSO_4 , mGFP-diS1 dimerised to form dGFP-diS1 (prefix d refers to dimer), with $\sim 55\%$ forming dimer after 1 hr (Fig. S1b), similar to the yield observed for click-based dimerisation³⁴. The exact mechanism by which copper enables dimerisation is unclear but is likely by facilitating the formation of a disulphide bridge between the monomer units (vide infra) The dGFP-diS1 dimer is stable and can be purified from the monomer by size exclusion chromatography (Fig. S1c).




Fig. 2. Dimerisation of sfGFP C148 variants. (a) Absorbance spectra of sfGFP^{WT} (black), dGFP-diS1 (red) and dGFP-diS2 (green). Inset is the change in absorbance on conversion of mGFP-S1 (dashed red line) to dGFP-diS1 (solid red line). The full absorbance spectra are shown in Fig. S2. Emission spectra of (b) GFP-diS1 and (c) GFP-diS2, with red and green representing the dimers, respectively, and black the monomer. Solid and dashed lines represent excitation at either the optimal CRO-B wavelength (490-497nm) or CRO-A wavelength (400 nm), respectively. (d) Rate of dimerisation of GFP-diS1 (grey) and GFP-diS2. The data shown are for $10 \mu\text{M}$ of starting monomer with R/R_0 representing the change in fluorescence emission ratio on excitation at either 388 nm or 483 nm.



Dimerisation results in a shift in maximal absorbance to 488 nm (Fig. 2a and Table 1), suggesting that CRO-O⁻ state now dominates dGFP-diS1. Moreover, molar absorbance for the CRO-O⁻ form increases ~2.9 fold with a ~2.5 fold increase in brightness compared to sfGFP (Fig. 2a-b, Table 1). The greater than 2-fold enhancement in the spectral properties of dGFP-diS1 suggests a synergistic relationship on dimerisation where the dimer is greater than the sum of its parts, as observed previously for the click linked dimers^{34,35}. Taking into account both the increase in emission on excitation at ~488 nm and the decrease on excitation at ~400 nm, we see a 31 fold signal change on dimerisation; the $Ex^{CRO-O^-}:Ex^{CRO-OH}$ emission ratio at 510 nm changes from 0.58 for monomer to 18 for the dimer (Fig. 2b).

The rate of mGFP-diS1 dimerisation is slow (Fig. 2d), with an estimated second order rate constant of 0.1 nM/min. While formation of covalent complex means the system is non-equilibrium, an apparent K_D can be estimated to provide an insight into monomer affinity; for GFP-diS1, the relatively low level of association observed (Fig. 2d) suggests that the K_D is >25 μ M. It has been estimated that the dissociation constant for weak GFP dimerisation is 100 μ M³⁸.

Optimising the sfGFP interaction interface.

The slow dimerisation kinetics of GFP-diS1 and low apparent affinity are likely due to interactions between the monomeric units being too transient and weak³⁸. Ideally, the kinetics should be faster and apparent K_D lower. Using the non-canonical amino acid sfGFP dimer structure linked via residue 148³⁴ (PDB 4nhn) as a guide, we aimed to optimise the GFP-diS1 interaction by targeting residue 206 that lies at the dimer interface (Fig. S3) and has been shown previously to contribute to weak dimerisation of *Aequorea victoria* derived FPs^{36,39}. We introduced the V206F mutation, so generating GFP-diS2. The GFP-diS2 variant is expressed and purified as a monomer (Fig. S4a-b) and has similar spectral characteristics to mGFP-diS1 (Fig. 1b-c and Table 1). The CRO-O⁻ absorbance peak at ~490 nm has a slightly higher molar absorbance in mGFP-diS2 compared to mGFP-diS1 with a concomitant drop in ~400 nm value. This could suggest GFP-diS2 improved the still weak



tendency to self-associate so increasing the population of the CRO-O⁻ state (Fig. 1b).

Again, Cu²⁺ is required to initiate dimerisation. In the presence of CuSO₄ mGFP-diS2 dimerised with an efficiency of ~80%, higher than that observed for mGFP-diS1, with the majority of dimerisation completed in the first 5 minutes (Fig. S4b-c). Replacing CuSO₄ with CuCl₂ did not have any discernible change in the ability of GFP-diS2 to dimerise (Fig. S4d). The switch in the major absorbance peak from 400 nm to 492 nm indicates that the CRO-O⁻ now dominates dGFP-diS2 (Fig. 2a). This is also manifested in the fluorescence emission spectra with the higher emission on excitation at ~490 nm (Fig. 2c). As with dGFP-diS1, dGFP-diS2 was brighter (2.9 fold) than wild-type sfGFP (Table 1). Indeed, dGFP-diS2 is brighter than dGFP-diS1 due to slightly higher molar absorbance and quantum yield (Table 1). The $\text{Ex}^{\text{CRO-O}^-}:\text{Ex}^{\text{CRO-OH}}$ (400 nm:492 nm) ratio on measuring emission at 510 nm changes from 0.56 to 13.4, which equates to a 24 fold increase of signal on dimerisation. Significantly, there is also a large increase in dimerisation rate (Fig. 2d), with the second order rate constant increasing two orders of magnitude to $11 \pm 1 \text{ nM}^{-1}\text{min}^{-1}$. The apparent K_D is estimated to be $4.6 \pm 0.2 \text{ }\mu\text{M}$, equivalent to a moderate protein-protein interaction. The dimerisation process is reversible as dGFP-diS2 can be converted back into its monomer components with monomeric spectral characteristics using DTT (Fig. S5). Thus, GFP-diS2 is a distinct improvement on GFP-diS1 in terms of its copper-dependent homodimerisation properties.




Fig. 3. Structure of mGFP-diS2 (PDB 8c1x). (a) Overall structure mGFP-diS2 with the chromophore shown as grey sticks and C148 shown as sticks and highlighted. (b) Structure of the chromophore and residues 147-148. Dashed lines represent potential H-bonds with the phenolate and S147. A channel through to the chromophore (CRO) (coloured orange) is shown together with a single conserved water molecule (W, red sphere) within the channel. The channel was calculated using CAVER3.0 with a probe size of 1.0 Å. (c) Residues involved in the classical proton transfer network related to promotion of the phenolate chromophore form. The grey residues are the equivalent position of S205 and E222 in sfGFP. Distances shown are for mGFP-diS2 and are in Ångstroms.

Structural basis for fluorescence switching in dimerisation

To fully understand the basis behind dimerization-dependent fluorescence switching, we determined the structure of the GFP-diS2 monomer and dimer (see Table S1 for structural statistics). In mGFP-diS2, H148C has a major impact on the local organisation of the structure that directly impacts on the interactions with the CRO (Fig. 3a-b and Fig. S6). The backbone of residues 147-149 is twisted compared to sfGFP^{WT} resulting in a register shift of S147 and C148 (Fig. 3b and Fig. S6). The change in position of C148 results in the loss of the H-bond to the CRO and becomes solvent exposed, therefore making it amenable to forming a disulphide bond. S147 now points towards the chromophore with the OH groups within H-bonding distance of each other (Fig. 4b). The new potential H-bond between CRO and S147 does not however appear to compensate for the H148C mutation in terms of promoting the phenolate chromophore form. Indeed, compared to wt sfGFP, the water network is less extensive around the phenol group for mGFP-diS2, with the sole water present potentially undergoing dynamic exchange with the bulk solvent via the channel shown in Fig. 3b. Such water networks are known to play a key role in chromophore charge state^{34,40,41}. The side chain orientation of E222 and thus its interaction with S205, a critical interaction involved in proton shuttling that promotes the CRO-O⁻ form^{42,43}, is also altered in mGFP-diS2; as a result, the distance between E222 and S203 increased to 3.7 Å (from 3.0 Å in sfGFP). Thus, the cumulation of these structural changes are likely to promote CRO-OH on the introduction of the H148C mutation. The V206F mutation does not have any significant impact on the local structure of sfGFP with both backbone and side-chain



configurations of the mutated residue and surrounding residues being similar (Fig. S7).




Fig. 4. Structure of dGFP-diS2. (a) Overall structure of dGFP-diS2 showing one monomer (green) in relation to the other (grey). The chromophores are shown as sticks and coloured green and the disulphide bridge is shown as spheres. The electron density of the disulphide bridge is shown in Fig. S8. The distance between the two chromophores is shown by the dashed double arrowhead line. (b) The role of the V206F mutation in stabilising the dimer interface. Residues with superscript A come from one subunit and are coloured green and those with superscript B are from the second subunit and coloured grey. The H77^B is highlighted as part of a pi-cation interaction. (c) The local chromophore environment with water molecules shown as red spheres. The monomeric mGFP-diS2 is overlaid and coloured grey with the conserved W¹ water molecule present in both structures coloured as an orange sphere. (d) Inter-subunit network between CRO^A and Y200^B.

The overall structure of dGFP-diS2 is shown in Fig. 4a. The two dimers have a defined, symmetrical interface comprising the same residues from each monomer unit covering 910 Å² (as defined by PISA⁴⁴). The residues that contribute dimer interface are detailed in Tables S2 and S3. The disulphide bridge has two potential configurations (see Fig. S8 for electron density) and is exposed to the solvent



explaining the ability of DTT to break the dimer apart (Fig. S5). The introduction of the V206F mutation plays an important role in promoting the dimer formation as it makes interactions with 6 residues in the neighbouring monomer, including a π -cation interaction with H77 (Fig. 5b).

The likely mechanism of CRO state switching is due to changes in the local chromophore environment on dimerisation. Monomeric and dimeric GFP-diS2 are very similar with a $C\alpha$ RMSD of 0.119 Å and local side chains around the chromophore occupying similar positions (Fig. S9a). The main changes are increased solvation of the chromophore and a plane flip in the position of the C148-N149 peptide bond (Fig. 4c and S9b). Peptide bond plane flipping is relatively rare and may be the rate-limiting step in the dimerisation processes^{45,46}; it could also provide a rationale for the requirement for Cu^{2+} to promote dimerisation. The water molecule close to the CRO (W^1) observed in mGFP-diS2 (Fig. 3b) is still present in the dimer but 4 additional water molecules are now observed. Together with W^1 , the W^2 and W^3 water molecules form a H-bond network to the CRO with these waters being planar to the CRO (Fig. 4c). W^4 and W^5 are above the plane of the CRO and occupy the space available due to rearrangement of the C148-N149 peptide bond; they form a H-bond network with S147. In the broader picture of the dimer, waters W^1 , W^2 and W^3 form a long-range polar network that spans the dimer interface incorporating Y200 from the second protomer (Fig. 4d). The formation of organised water networks in avGFP-like proteins close to the CRO has been postulated previously to promote the formation of the ground state CRO- O^- form^{33–35}. The water networks observed here (Fig. 4c-d) are likely to play a similar role by facilitating proton abstraction which can then be shuttled through the neighbouring water network at the dimer interface. These local water molecules together with S147 can help stabilise the negative charge so promoting CRO- O^- . Crucially, these water molecules are also largely buried at the dimer interface so restricting their exchange with bulk solvent (Fig. S10). Recent work has shown that the structurally conserved water molecule W^1 rapidly exchanges with the bulk solvent in sfGFP³⁰. The potential to restrict exchange of water molecules may contribute to the improved spectral properties on dimerisation both here and elsewhere^{34,35}.






Fig. 5. Dimerisation of GFP-diS2 fused to a leucine zipper element. (a) Schematic representation of activation mechanism. The leucine zipper (ZIP) is coloured grey, linker sequence orange and the GFP-diS2 component green. (b) Absorbance spectra of GFP-diS2-ZIP in the absence (black) and presence (green) of 1 mM CuSO₄. *In vitro* association kinetics of GFP-diS2-ZIP in the presence of (c) Cu²⁺ (1 mM CuSO₄) or (e) Cu⁺ (250 μM CuSO₄ plus 1 mM ascorbate). The influence of either (d) Cu²⁺ or (f) Cu⁺ on the GFP-diS2-ZIP fluorescence ratio.

Fusing GFP-diS2 to a homo-dimerising component.

The switch from CRO-OH to CRO-O⁻ provides a means to monitor copper levels and also protein dimerisation through ratiometric changes in fluorescence emission on excitation at two distinct wavelengths (~400 nm and ~490 nm). The N- and C-termini are on the same face (Fig. S11), so should not preclude association if fused



to naturally homo-dimeric proteins. To assess the effect of fusing GFP-diS2 natural homodimerizing protein component, we employed the leucine zipper region (ZIP) of transcription factor GCN4⁴⁷. In this scenario, the leucine zipper region should result in GFP-diS2 becoming an obligate homodimer (Fig. 5a). Addition of copper ion should result in rapid switching “on” of fluorescence emission on excitation at ~490 nm due to the localisation of the two GFP-diS2 components.

In the absence of CuSO₄, the homodimer construct now termed GFP-diS2-ZIP, exhibited spectral characteristics similar to mGFP-diS2 (Fig. 5b and Table 1), with the ~400 nm absorbance peak equivalent to CRO-OH dominating. On addition of CuSO₄, GFP-diS2-ZIP quickly switches to CRO-O⁻ as indicated by the dominant absorbance at ~493 nm (Fig. 5b). As with the monomer-to-dimer conversion of GFP-diS2, the chromophore molar absorbance coefficient was enhanced compared to sfGFP and dGFP-diS2 (Table 1). The $\text{Ex}^{\text{CRO-O}^-}:\text{Ex}^{\text{CRO-OH}}$ fluorescence emission ratio was also higher in the obligate homodimer, with a 37 fold increase compared to 27 fold change when forming dGFP-diS2. The rate of conversion was also far quicker (Fig. 5c) with a second order rate constant of $810 \pm 30 \text{ nM}^{-1}\text{min}^{-1}$, ~65 fold quicker than monomer-to-dimer conversion; for a 1 μM sample of GFP-diS2-ZIP conversion was essentially complete within 5 min (Fig. 5c). Thus, coupling GFP-diS2 to a known homodimerisation element results in a much improved GFP association and thus switch in fluorescence. The switch is also reversible as addition of DTT quickly converts copper-treated GFP-diS2-ZIP to a form with spectral properties like that in the absence of copper (Figure S12). The change in emission ratio is also dependent on Cu²⁺ concentration suggesting GFP-diS2-ZIP can be used as a cellular copper sensor (Fig. 5d). At Cu²⁺ concentrations $\leq 10 \mu\text{M}$, the rate of switching is slow (Fig. S13a) with the end point change being relatively small (Fig. 5d). The ratiometric switch also appears to be specific to copper ions as other biologically relevant metal ions did not change the $\text{Ex}^{\text{CRO-O}^-}:\text{Ex}^{\text{CRO-OH}}$ emission ratio (Supporting Table S4).

Inside the reducing environment of the cell, Cu²⁺ is reduced to Cu⁺ and is mainly transported into the cell as the Cu⁺ form^{8,10}. For use inside the cell, we assessed the impact of reducing conditions and Cu⁺ on GFP-diS2-ZIP fluorescence switching. Cu⁺ is commonly produced in the presence of ascorbate, a natural biological reducing agent⁴⁸. The rate of association increased considerably in the presence of Cu⁺, with fluorescence switching occurring in <1 min (Fig. 5e), with an estimated second order



rate constant of $3118 \pm 465 \text{ nM}^{-1} \text{ min}^{-1}$. Cu^+ also displays a concentration dependence in fluorescence switching (Fig. 5f and Fig. S13b), as with Cu^{2+} . Ascorbate itself has little effect on the spectral properties of GFP-diS2-ZIP (Fig. S14). It is not clear why Cu^+ , a reducing agent, should promote bringing the two GFP-diS2 units close to each and instigate switching using the proposed disulphide mechanism above. Cu^+ does have high affinity for thiolates capable of forming a bis-thiolate complex (e.g. in CSP1⁴⁹). Here, Cu^+ could be acting as the temporary bridge and/or a catalyst that facilitates linking together the GFP-diS2 units via the engineered cysteine residues.




Fig. 6. *In situ* response of GFP-diS2-ZIP to copper. (a) Change in fluorescence emission ratio (Ex 390 nm: Ex 490 nm) of bulk *E. coli* cell culture on addition of 1 mM CuSO_4 . Inset is the fluorescence emission spectra (emission at 520 nm) measured before (black) and 3600 s after (green) the addition of 1 mM CuSO_4 . (b) Change in fluorescence emission ratio (Ex 405 nm: Ex 470 nm) of individual *E. coli* cells on addition of 1 mM CuSO_4 . A total of 23 different cells were analysed from widefield imaging. The error bars represent the standard deviation between the 23 different cells. (c) Representative widefield images at the different centre excitation wavelength of individual *E. coli* cells at different time periods. OL refers to the overlay of the two individual excitation wavelengths. The accompanying movies over the full time course can be found as Supplementary Movies 1 and 2.

We then tested the GFP-diS2-ZIP system *in situ* by monitoring changes in fluorescence ratio in *E. coli*. The R_0 ratio (Ex 390 nm:Ex 490 nm at 0 s) ratio was



~0.60, close to the in vitro value of 0.58 reported above. This suggests that free copper ion is very low in ambient conditions, which is in line with previous work^{8,50}. On addition of CuSO₄ to the cell suspensions, there is an initial lag of ~500 s, followed by an increase in fluorescence ratio; a plateau is reached after ~55 min (Fig. 6a). The change in the maximal fluorescence ratio is 7.2 fold, which equates to a sub 1 μ M Cu⁺ concentration and/or 100 μ M Cu²⁺, based on the metal ion dependent concentration curves in Fig. 5.

GFP-diS2-ZIP can also be used as a cell imaging probe. Fluorescence microscopy allows the impact of copper on individual cells to be investigated. As with the bulk cell culture, an initial lag is observed of ~60 s (plus 4-5 min preparation before imaging) after adding CuSO₄ to the cells. After the initial lag phase, the fluorescence emission ratio increases until it reaches an R/R₀ plateau at around 9. This equates to a sub micromolar Cu⁺ concentration, the form of copper that is thought to predominate inside the cell under reducing conditions¹¹. The cells analysed appear to retain their general cellular morphology with the changes in fluorescence levels clearly evident (Fig. 6c).

Our data suggests that copper can enter *E. coli* with available levels likely to be in the 10-100s nM range (10⁻⁸-10⁻⁷ M) based on measured Ex^{CRO-O⁻}:Ex^{CRO-OH} emission ratio correlated to copper concentration in Fig. 5. Copper import into *E. coli* is largely unknown but the initial lag period suggests there is a requirement to induce production of import systems such as the recently reported AZY operon system⁵¹. It was originally thought that only efflux systems were induced in response to high copper concentrations^{8,10,11}. Cellular copper is thought to be ~10 μ M but the vast majority is bound to proteins leaving the available intracellular free copper ion at <10⁻¹⁸ M or just a few atoms per cell⁵² but has been reported to rise to 10⁻¹⁵ M^{11,53}. These published copper concentrations are still much lower than our data here suggests but is in line with reported μ M levels in the periplasm⁵⁴. What is clear is that our GFP-diS2-ZIP system can respond dynamically to changes in cellular copper concentrations generating a positive and large (compared to existing genetically encoded copper sensors) signal change. The requirement of disulphide bond formation as the mechanism for switching could limit the sensor's capacity to monitor fluxes in copper over extended periods by trapping GFP-diS-ZIP in the "on" state. However, as we have shown, the process is potentially reversible under



reducing conditions (Fig. S5 and S12). GFP-diS-ZIP produced in the cells is in the “off” state (engineered C148 residues remain as thiols) suggesting this is the default state under low copper conditions. Bacteria such as *E. coli* maintain a reducing cellular environment, with glutathione playing a key role in this process. Glutathione, which is at a concentration of ~17 mM in *E. coli*⁵⁵, is also thought to play a role in copper chelation^{53,56}. In the presence of high copper concentrations, the conditions inside the bacterial cell (e.g. high glutathione levels) do not appear to impede the disulphide-based GFP-diS-ZIP copper response.

Conclusion

We have engineered a well-utilised GFP to change its spectral characteristics in response to copper-dependent homo-dimerisation. The molecular basis of action suggests that organised water molecules play a key role in promoting proton transfer from the chromophore causing a switch in its charged ground state from the neutral phenolic to negatively charged phenolate. This in turn changes the primary excitation wavelength providing a measurable response on homo-dimerisation. The requirement for CuSO₄ provides a means to measure copper levels without interference from other metal ions and to our knowledge is the first genetically encoded ratiometric copper sensor with a large positive gain of signal. Genetically encoded copper biosensors have a range of potential applications. Our demonstration here in bacteria opens routes for environmental monitoring ranging from copper bioavailability in soil to industrial waste tracking and bioremediation. Adaption for use in mammalian cell systems could lead to ways to understand the role of copper levels in diseases ranging from dementia to cancer. Replacement of the disulphide bridge with a copper-chelating motif¹⁵ could provide an addition route to measure copper flux through metal ion binding and unbinding without the need for forming a covalent link. There is the potential to transfer the system to monitor more generally homodimerisation, arguably the most common of all PPIs in biology, by replacing the leucine zipper component with another homodimerisation element. The need for copper and the disulphide bridge is currently a limiting aspect but further engineering to optimise the dimer interface and promote the 148-149 peptide bond plane flipping may potentially result in a non-covalent system that can monitor reversible and dynamic exchange PPI events.



Methods and Materials.

Protein engineering and purification.

The gene encoding sfGFP was resident in the pBAD plasmid (arabinose inducible)³² or the pCA24N P50-GFP fusion (see Supporting Information for gene fusion construction) were used as the template for whole plasmid inverse Q5 DNA polymerase mutagenesis procedure (NE Biolabs). The H148C and V206F mutations were introduced using the following primer pairs: H148C, 5'-GCGGTAATATACACATTGCAGCTGTTG-3' and 5'-CGATAAACAGAAAAATGGCATCAAAGCG-3'; V206F, 5'-GAGCAAAGATCCGAATGAAAAACGTG-3' and 5'-AGAAAGCTCTGGGTGCTCAGATA-3'. All mutations were confirmed by DNA sequencing (Eurofins Genomics). The GFP-diS2-ZIP construct was synthesised by Twist Bioscience and cloned into the pET28a (IPTG inducible). Proteins were purified essentially as described previously^{32,57} and with details provided in the Supporting Information.

Dimerisation of sfGFP variants

CuSO₄ (Sigma-Aldrich) was added to sfGFP variant protein samples to aid dimerisation through formation of a disulphide bond⁵⁸. Dimerisation of the sfGFP variants in 50 mM TrisHCl, pH 8, was initiated by the addition of CuSO₄ to a final concentration of 1 mM and then left to incubate for varying amounts of time. For association kinetics, protein concentrations ranged from 0.1 μM to 0.5 μM with samples taken periodically for analysis. The dimerisation of sfGFP variants used in this study was a second order rate reaction, as homodimerisation is recorded as $A + A \rightarrow A_2$ ⁵⁹. The second order rate constant was calculated using the equation $v = k[A]^2$ where v is the rate, k is the rate constant ($\mu\text{M}^{-1}\text{sec}^{-1}$) and A is the monomer protein concentration (μM). To calculate the percentage of sfGFP that has dimerised, the emission ratios on excitation of the CRO-OH (390-400 nm) or CRO-O⁻ (485-495 nm) were recorded at various time points as described below. For preparation of large quantities of dimer, samples of up to 50 mg/mL of protein were incubated with 1 mM CuSO₄ for 1 hr with mixing. The monomer and dimer were separated by SEC as described in the Supporting Information.



Absorbance and fluorescence spectroscopy.

UV-visible (UV-vis) absorption spectra were recorded on a Cary 60 spectrophotometer (Agilent) in a 1 cm pathlength quartz cuvette (Hellma). Spectra were recorded from 200 – 600 nm at a rate of 300 nm/min. Extinction coefficients were calculated using the Beer-Lambert law by using known concentration of protein as determined by the DC protein assay (BioRAD) with sfGFP as the calibration standard and verified by comparison with sfGFP's 280 nm absorbance (sfGFP, $\epsilon_{280} = 25,400 \text{ M}^{-1}\text{cm}^{-1}$). Steady state emission and excitation fluorescence spectra were measured using Varian Cary Eclipse Fluorimeter and a 10 mm x 2 mm QS quartz cuvette (Hellma). Spectra were recorded with 5 nm excitation/emission, slit width at a rate of 300 nm/min. Emission spectra were recorded at a fixed excitation wavelength according to the excitation maximum of the variant. Excitation spectra were recorded at the fixed wavelength according to the maximum emission. Unless otherwise stated, spectral scans were recorded for at 0.5 μM in 50 mM Tris-HCl, pH 8.0, for monomers and at 0.25 μM for dimers. For GFP-diS1 and GFPdiS2 dimerisation kinetics, a CLARIOstar Plus plate reader (BMG LabTech) was also used. The bandwidth of filters used was 8 nm. As the minimum distance allowed between filters is 30 nm, the emission was recorded at 516 upon excitation at 388 nm and 483 nm. Each measurement was performed in triplicate using a NuncTM F96 MicrowellTM black polystyrene plate (Thermo fisher). To calculate $\text{Ex}^{\text{CRO-OH}}:\text{EX}^{\text{CRO-OH}}$ ratios, emission values upon excitation at major absorbance peak around 490 nm were divided by emission values when recorded at ~400 nm. Extact wavelengths are stated in the main text. The ratio are presented as R/R_0 where R is the ratio at a particular time point and R_0 the ratio at the start of the measurement. Quantum yield was determined using a fluorescein standard as described previously³³.

Structure determination.

Purified proteins samples in 50 mM Tris-HCl, pH 8 at ~10 mg/mL were screened for crystal formation using sitting drop vapour diffusion across a wide variety of conditions as described by the PACT *premier*TM and JCSG-*plus*TM HT-96 broad crystallisation screens (Molecular Dimensions). Crystallisation plates were stored at 23°C and crystal growth monitored. Crystals were harvested, 1 mM ethylene glycol added and flash frozen in liquid nitrogen. X-ray diffraction data was obtained at



Diamond Light Source, Harwell, UK on beamline I03. Crystals of mGFP-diS2 were in the P 1 2₁ 1 space group grew in condition G12 on the PACT Premier screen (0.2 M Sodium malonate dibasic monohydrate, 0.1 M Bis- Tris propane, pH 7.5, 20 % w/v PEG 3350). The crystals diffracted to the highest resolution of 1.89 Å. Crystals of dGFP-diS2 were in the P 2 2 2₁ space group and grew in condition G7 on the PACT Premier screen (0.2 M sodium acetate trihydrate, 0.1 M Bis-tris propane, pH 7.5, 20% w/v PEG 3350). The crystals diffracted to the highest resolution of 1.79 Å. The structures were determined by molecular replacement, with the sfGFP structure (PDB 2b3p²⁹) as the reference, essentially as described previously^{33,34}. Full diffraction data and refinement statistics can be found in Table S1. The PDB entry for mGFP-diS2 is 8c1x and is 8bxp for dGFP-diS2.

E. coli cell imaging.

Bacteria were induced with 1 mM IPTG and grown to mid log phase before being concentrated by centrifugation tenfold to increase the number of bacteria present within an individual field of view. Cells were mixed with CuSO₄ (final concentration of 1 mM), before 20 uL were placed on a fresh microscope slide and covered with a 25mM #1.5 glass coverslip and immediately placed on the microscopy for imaging. Wide-field epi fluorescence measurements were conducted on an inverted Olympus IX73 microscope and a Prior Lumen200Pro light source using filter set 89000, (Chroma, Vermont, U.S.A.) selecting the ET402 nm/15 nm or ET490 nm/20 nm as the excitation filter and the ET525 nm/36 nm as the emission filter. The fluorescence emission was detected with a Hamamatsu ORCA-flash 4.0 V2 sCMOS Camera operated utilizing the HCImage software package (Hamamatsu). An Olympus UPlanSApo 20× oil immersion objective with an NA of 0.75 was used to collect sequential images at full resolution (2048x2048 pixels at 1xbinning) with exposure times of 500 ms for each channel, with a 10 second interval over 75 timepoints. To reduce any impact of sample drift on quantification, cell images were registered in xy using the "Linear stack alignment with SIFT multichannel" plugin⁶⁰ in FIJI⁶¹. A total of 23 individual bacteria were selected through the use of a 5 pixel diameter circular region of interest, and the intensity of each channel within each region of interest (ROI) was measured. Background measurements were calculated from the average of 5 ROI containing no bacteria at time 0, and subtracted from each measured timepoint. Finally, a ratio was determined between centred 402 and 490 nm



excitation and the R/R_0 calculated for each individual bacteria, before being averaged across all bacteria and plotted against time.

Conflicts of interest.

There are no conflicts of interest.

Acknowledgement

We would like to thank the staff at the Diamond Light Source (Harwell, UK) for the supply of facilities and beam time, especially beamline I03 staff, under beamtime code mx18812. DDJ would like to thank the following funders: KESS2 (Knowledge Economy Skills Scholarship European Regional Development Fund via Welsh Government) studentship in partnership with Tenovus (to RDA), a Royal Thai Embassy studentship to DV, Tabuk University and the Royal Embassy of Saudi Arabia (to KA), BBSRC (BB/Z514913/1), and EPSRC (EP/V048147/1). The authors would like to thank the Cardiff School of Biosciences Protein Technology Hub for helping with the production and analysis of proteins, and the Cardiff University Bioimaging Hub Core Facility (RRID: SCR_022556) for their support and assistance in this work.

Author contributions.

All authors contributed to the writing of the paper and analysing data. RDA designed monomer variants, prepared in the monomer variants, analysed the initial monomer variants (including dimerisation potential) and determined their structure. DV designed and generated the LZ variants, analysed the LZ variants and undertook the in cell analysis. KDH contributed to the LZ variant analysis and cell imaging. KA contributed towards LZ variant analysis and cell imaging. PJR collected structural data and helped with structure determination and refinement. PDW contributed to cell imaging and data analysis. DDJ conceived and directed the project, contributed to variant design and data analysis.

Data availability.

The structures determined have been deposited in the PDB with codes 8c1x and 8bxp. Data will be also be made available via FigShare (<https://figshare.com/s/826745d1a6d6d5b1c668>).



590

View Article Online
DOI: 10.1039/D5SC06589E

Open Access Article. Published on 13 October 2025. Downloaded on 10/14/2025 4:04:41 PM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



References

- 1 R. Y. Tsien, *Annu Rev Biochem*, 1998, **67**, 509–44.
- 2 N. C. Shaner, P. A. Steinbach and R. Y. Tsien, *Nat Methods*, 2005, **2**, 905–909.
- 3 E. A. Rodriguez, R. E. Campbell, J. Y. Lin, M. Z. Lin, A. Miyawaki, A. E. Palmer, X. Shu, J. Zhang and R. Y. Tsien, *Trends Biochem Sci*, 2017, **42**, 111–129.
- 4 M. S. Frei, S. Mehta and J. Zhang, *Annu Rev Biophys*, 2024, **53**, 275–297.
- 5 K. Baek, K. Ji, W. Peng, S. M. Liyanaarachchi and S. C. Dodani, *Protein Engineering, Design and Selection*, 2021, **34**, 1–9.
- 6 W. Shi, C. Zhan, A. Ignatov, B. A. Manjasetty, N. Marinkovic, M. Sullivan, R. Huang and M. R. Chance, *Structure*, 2005, **13**, 1473–1486.
- 7 C. L. Dupont, A. Butcher, R. E. Valas, P. E. Bourne and G. Caetano-Anollés, *Proceedings of the National Academy of Sciences*, 2010, **107**, 10567–10572.
- 8 B.-E. Kim, T. Nevitt and D. J. Thiele, *Nat Chem Biol*, 2008, **4**, 176–185.
- 9 L. Banci, I. Bertini, S. Ciofi-Baffoni, T. Kozyreva, K. Zovo and P. Palumaa, *Nature*, 2010, **465**, 645–648.
- 10 I. Bertini, G. Cavallaro and K. S. McGreevy, *Coord Chem Rev*, 2010, **254**, 506–524.
- 11 C. Rensing and G. Grass, *FEMS Microbiol Rev*, 2003, **27**, 197–213.
- 12 K. P. Carter, A. M. Young and A. E. Palmer, *Chem Rev*, 2014, **114**, 4564–4601.
- 13 G. C. Jensen, M. K. Janis, H. N. Nguyen, O. W. David and M. L. Zastrow, *ACS Sens*, 2024, **9**, 1622–1643.
- 14 M. T. Morgan, D. Bourassa, S. Harankhedkar, A. M. McCallum, S. A. Zlatic, J. S. Calvo, G. Meloni, V. Faundez and C. J. Fahrni, *Proc Natl Acad Sci U S A*, 2019, **116**, 12167–12172.
- 15 Y. A. Choi, J. O. Keem, C. Y. Kim, H. R. Yoon, W. Do Heo, B. H. Chung and Y. Jung, *Chem Sci*, 2015, **6**, 1301–1307.
- 16 J. Liang, L. Guo, Y. Ding, L. Xia, Y. Shen, M. Qin, Q. Xu, Y. Cao and W. Wang, *Biochem Biophys Res Commun*, 2014, **443**, 894–898.
- 17 J. Liu, J. Karpus, S. V. Wegner, P. R. Chen and C. He, *J Am Chem Soc*, 2013, **135**, 3144–3149.



- 625 18 M. H. Ali and B. Imperiali, *Bioorg Med Chem*, 2005, **13**, 5013–5020. View Article Online
DOI: 10.1039/D5SC06589E
- 626 19 D. S. Goodsell and A. J. Olson, *Annu Rev Biophys Biomol Struct*, 2000, **29**,
627 105–153.
- 628 20 J. A. Marsh and S. A. Teichmann, *Annu Rev Biochem*, 2015, **84**, 551–575.
- 629 21 L. Bonetta, *Nature*, 2010, **468**, 851–4.
- 630 22 M. P. H. Stumpf, T. Thorne, E. de Silva, R. Stewart, H. J. An, M. Lappe and C.
631 Wiuf, *Proceedings of the National Academy of Sciences*, 2008, **105**, 6959–
632 6964.
- 633 23 M. Vidal, M. E. Cusick and A.-L. Barabási, *Cell*, 2011, **144**, 986–998.
- 634 24 A. Miyawaki, J. Llopis, R. Heim, J. M. McCaffery, J. A. Adams, M. Ikurak and R.
635 Y. Tsien, *Nature*, 1997, **388**, 822–887.
- 636 25 J. Zhang, R. E. Campbell, A. Y. Ting and R. Y. Tsien, *Nat Rev Mol Cell Biol*,
637 2002, **3**, 906–918.
- 638 26 W. R. Algar, N. Hildebrandt, S. S. Vogel and I. L. Medintz, *Nat Methods*, 2019,
639 **16**, 815–829.
- 640 27 T. K. Kerppola, *Nat Rev Mol Cell Biol*, 2006, **7**, 449–456.
- 641 28 R. Heim and R. Y. Tsien, *Curr Biol*, 1996, **6**, 178–182.
- 642 29 J. D. Pédelacq, S. Cabantous, T. Tran, T. C. Terwilliger and G. S. Waldo, *Nat*
643 *Biotechnol*, 2006, **24**, 79–88.
- 644 30 R. D. Ahmed, W. D. Jamieson, D. Vitsupakorn, A. Zitti, K. A. Pawson, O. K.
645 Castell, P. D. Watson and D. D. Jones, *Commun Chem*, 2025, **8**, 174.
- 646 31 B. Campanini, B. Pioselli, S. Raboni, P. Felici, I. Giordano, L. D'Alfonso, M.
647 Collini, G. Chirico and S. Bettati, *Biochim Biophys Acta Proteins Proteom*,
648 2013, **1834**, 770–779.
- 649 32 S. C. Reddington, P. J. Rizkallah, P. D. Watson, R. Pearson, E. M. Tippmann
650 and D. D. Jones, *Angewandte Chemie - International Edition*, 2013, **52**, 5974–
651 5977.
- 652 33 A. M. Hartley, H. L. Worthy, S. C. Reddington, P. J. Rizkallah and D. D. Jones,
653 *Chem Sci*, 2016, **7**, 6484–6491.
- 654 34 H. L. Worthy, H. S. Auhim, W. D. Jamieson, J. R. Pope, A. Wall, R. Batchelor,
655 R. L. Johnson, D. W. Watkins, P. Rizkallah, O. K. Castell and D. D. Jones,
656 *Commun Chem*, 2019, **2**, 83.

- 657 35 J. R. Pope, R. L. Johnson, W. D. Jamieson, H. L. Worthy, S. Kailasam, R. D.
658 Ahmed, I. Taban, H. S. Auhim, D. W. Watkins, P. J. Rizkallah, O. K. Castell
659 and D. D. Jones, *Advanced Science*, 2021, **8**, 2003167–2003167.
- 660 36 P. J. Cranfill, B. R. Sell, M. A. Baird, J. R. Allen, Z. Lavagnino, H. M. De
661 Gruiter, G. J. Kremers, M. W. Davidson, A. Ustione and D. W. Piston, *Nat*
662 *Methods*, 2016, **13**, 557–562.
- 663 37 I. S. Matsui Lee, M. Suzuki, N. Hayashi, J. Hu, L. J. Van Eldik, K. Titani and M.
664 Nishikimi, *Arch Biochem Biophys*, 2000, **374**, 137–141.
- 665 38 G. N. Phillips, *Curr Opin Struct Biol*, 1997, **7**, 821–827.
- 666 39 D. A. Zacharias, J. D. Violin, A. C. Newton and R. Y. Tsien, *Science (1979)*,
667 2002, **296**, 913–916.
- 668 40 A. Shinobu and N. Agmon, *Journal of Physical Chemistry B*, 2015, **119**, 3464–
669 3478.
- 670 41 A. Shinobu, G. J. Palm, A. J. Schierbeek and N. Agmon, *J Am Chem Soc*,
671 2010, **132**, 11093–11102.
- 672 42 S. J. Remington, *Protein Science*, 2011, **20**, 1509–1519.
- 673 43 J. J. van Thor, *Chem Soc Rev*, 2009, **38**, 2935–2950.
- 674 44 E. Krissinel and K. Henrick, *J Mol Biol*, 2007, **372**, 774–97.
- 675 45 S. Hayward, *Protein Sci*, 2001, **10**, 2219–27.
- 676 46 J. E. Milner-White, J. D. Watson, G. Qi and S. Hayward, *Structure*, 2006, **14**,
677 1369–76.
- 678 47 E. K. O'Shea, J. D. Klemm, P. S. Kim and T. Alber, *Science*, 1991, **254**, 539–
679 44.
- 680 48 H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew Chem Int Ed Engl*, 2001,
681 **40**, 2004–2021.
- 682 49 N. Vita, S. Platsaki, A. Baslé, S. J. Allen, N. G. Paterson, A. T. Crombie, J. C.
683 Murrell, K. J. Waldron and C. Dennison, *Nature*, 2015, **525**, 140–3.
- 684 50 Y. Kang, W. Lee, S. Kim, G. Jang, B.-G. Kim and Y. Yoon, *Appl Microbiol*
685 *Biotechnol*, 2018, **102**, 1513–1521.
- 686 51 C. D. Palmer, Y. Ghnamah, N. Livnat-Levanon, O. Lewinson and A. C.
687 Rosenzweig, *Commun Biol*, 2025, **8**, 458.
- 688 52 A. Changela, K. Chen, Y. Xue, J. Holschen, C. E. Outten, T. V. O'Halloran and
689 A. Mondragón, *Science (1979)*, 2003, **301**, 1383–1387.



- 690 53 M. T. Morgan, L. A. H. Nguyen, H. L. Hancock and C. J. Fahrni, *Journal of* View Article Online
DOI: 10.1039/D5SC06589E
691 *Biological Chemistry*, 2017, **292**, 21558–21567.
- 692 54 A. Giachino and K. J. Waldron, *Mol Microbiol*, 2020, **114**, 377–390.
- 693 55 B. D. Bennett, E. H. Kimball, M. Gao, R. Osterhout, S. J. Van Dien and J. D.
694 Rabinowitz, *Nat Chem Biol*, 2009, **5**, 593–9.
- 695 56 J. H. Freedman, M. R. Ciriolo and J. Peisach, *J Biol Chem*, 1989, **264**, 5598–
696 605.
- 697 57 S. C. Reddington, E. M. Tippmann and D. D. Jones, *Chemical*
698 *Communications*, 2012, **48**, 8419–8421.
- 699 58 S. V. Tillu, R. S. Lokhande and J. Bacardit, *International Journal of Pharma*
700 *Research and Health Sciences*.
- 701 59 R. C. Garratt, F. Valadares, J. Fernando and R. Bachega, *Encyclopedia of*
702 *Biophysics*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2013.
- 703 60 D. G. Lowe, *Int J Comput Vis*, 2004, **60**, 91–110.
- 704 61 J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T.
705 Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J.
706 White, V. Hartenstein, K. Eliceiri, P. Tomancak and A. Cardona, *Nat Methods*,
707 2012, **9**, 676–682.
- 708



Chromophore charge-state switching through copper-dependent homodimerisation of an engineered green fluorescent protein.

View Article Online
DOI: 10.1039/D5SC06589E

Rochelle D Ahmed^{a#}, Danoo Vitsupakorn^{a#}, Kieran D. Hartwell^a, Karma Albalawi^{a,c}, Pierre J. Rizkallah^b, Peter D. Watson^a and D. Dafydd Jones^{a*}.

- a. School of Biosciences, Molecular Biosciences Division, Cardiff University, Sir Martin Evans Building, Cardiff, CF10 3AX, UK.
- b. School of Medicine, Cardiff University, Cardiff, CF14 4XN, UK.
- c. Department of Chemistry, Faculty of Science, University of Tabuk, Tabuk, Saudi Arabia.

* Corresponding author. D. Dafydd Jones, School of Biosciences, Sir Martin Evans Building, Cardiff University, Cardiff CF10 3AX, UK. Email: jonesdd@cardiff.ac.uk

These authors contributed equally.



Data availability statement.

View Article Online
DOI: 10.1039/D5SC06589E

Gene sequence data is available in the Supplementary Information.

Data in figures will be made available via FigShare via

<https://figshare.com/s/826745d1a6d6d5b1c668>

Crystal structures have been submitted to the PDB with ID codes 8c1x and 8bpx.

