



Cross-cultural de-centring of Recovery College tools: a 20-country Delphi study



Yasuhiro Kotera ^{1,2,3}✉, Sara Vilar-Lluch ⁴, Dagmar Narusson⁵, Ramona Hiltensperger ⁶, Peanchanan Leah ⁷, Jose Orsi^{8,9}, Mário César Rezende Andrade¹⁰, Yuki Miyamoto¹¹, Ulrika Bejerholm¹², Jana Pluhaříková Pomajzlová¹³, Vlasta Řezníková¹⁴, Mariam Namasaba ¹⁵, Tesnime Jebara¹⁵, Vanessa Lawrence ¹⁵, Simran Kaur Takhi¹, Amy Ronaldson¹⁵, Vanessa Kellermann ¹⁵, Agnieszka Kapka¹⁵, Peter Bates¹⁶, Claire Henderson^{15,18} & Mike Slade ^{1,17,18}

Recovery Colleges are expanding internationally (264 identified in 31 countries across six continents), yet key tools for defining and assessing them were developed in high-income Anglophone contexts. We conducted a two-round international Delphi study using a cross-cultural de-centring approach to examine the cross-cultural applicability of the RECOLLECT Change Model and RECOLLECT Fidelity Measure, and to develop versions that were more applicable across the participating settings and interoperable versions of these tools. In Round 1, 54 panellists from 20 countries rated item importance and cultural difficulty and provided free-text feedback, analysed using corpus-informed linguistic methods. Core principles were broadly endorsed, but ‘co-production’, ‘recovery’, ‘professionals’ and ‘shifting the balance of power’ were interpreted inconsistently, revealing embedded assumptions about equality, autonomy and open-ended participation. Round 1 findings informed wording refinements tested in Round 2 with 41 panellists from 15 countries. All 11 revised items reached consensus for understandability and cultural appropriateness, strengthening the tools while preserving core Recovery College values. By identifying and revising culturally embedded assumptions while preserving interoperability with the original tools, this study demonstrates a transferable approach to the cross-cultural adaptation of mental health tools.

Recovery Colleges (RCs) are an innovative approach to mental health support that combines co-production and adult learning to promote personal recovery and social inclusion^{1,2}. Established in England in 2009, RCs have since expanded internationally. A previous global review identified 221 RCs in 28 countries in 2022³, and we are aware of further growth to 264 RCs worldwide by 2026. RCs provide inclusive educational environments in which people with mental health problems, carers, staff, and people in the wider community (referred to as ‘students’) learn together, develop skills and build support networks. Through self-directed learning and shared expertise, RCs support fuller social participation, greater self-management and personal recovery⁴.

Emerging evidence indicates that RCs can improve student outcomes, including self-esteem, quality of life and internalised stigma⁵, and benefit staff through increased motivation and skill development⁶. A large retrospective matched cohort study involving 1435 students and 4665 controls found reduced subsequent mental health service use and a £5028 greater reduction in total costs per student at 12 months, equivalent to an estimated

8.4:1 return on investment⁷. Uncontrolled before-and-after analyses in the same evaluation also showed improved symptoms and functioning⁷. These economic findings compare favourably with a 2025 estimate of an 8:1 socio-economic return for preventive healthcare interventions⁸. However, operational guidance for RCs has only recently become more standardised through the Recovery Colleges Characterisation and Testing (RECOLLECT) programme (NIHR200605).

Two widely-used outputs from RECOLLECT are the RECOLLECT Change Model (RCM) and the RECOLLECT Fidelity Measure (RFM). The RCM identifies four mechanisms through which RCs are proposed to work: empowering environment; shifting the balance of power; enabling different relationships; and facilitating personal growth⁹. The RFM is a 12-item tool assessing fidelity to core RC principles and best practice¹⁰. The 12 items comprise seven non-modifiable items, representing core principles expected to be present across RCs, and five modifiable items, representing features that may appropriately vary by local context. Non-modifiable items include equality, adult learning, tailoring to the student, co-production, social

connectedness, community focus and commitment to recovery. Modifiable items concern accessibility, location, distinctiveness of course content, strength-based orientation and progressiveness; for example, whether access is universal or targeted, whether the setting is RC-specific or community-integrated, and whether course content is distinct from or integrated with clinical services (Supplementary Table 1). Developed in England through an empirical programme of work, these tools drew on a literature review, consultation with 77 international experts, interviews with 10 RC managers, refinement with 44 students, trainers and managers across 3 sites, and evaluation involving 60 professional stakeholders.

The international growth of RCs raises an important question: whether tools developed in one cultural context can be applied appropriately across others¹¹. This concern is particularly relevant because most RC research has been conducted in England, where approximately one third of all RCs are based¹², and in other high-income Global North settings, limiting understanding of how RC principles and language operate across more diverse cultural, linguistic and service systems^{3,13}.

International research also indicates that RC models are negotiated within local contexts rather than transferred unchanged. Dutch studies have highlighted the importance of co-creation and peer leadership, alongside challenges involving accessibility, family involvement, collaborative learning and organisational positioning^{14,15}. Danish qualitative research identified practical, relational and course-related barriers to participation, including alienation when course activities assumed a willingness to disclose personal experiences¹⁶. Together, these findings suggest that core RC principles may be retained while their enactment and acceptability vary across cultural and service contexts. Previous research has therefore established the RFM¹⁰, applied it across international RC settings^{3,12}, examined cultural variation in fidelity and public-facing discourse^{17,18}, and documented local adaptation of RC models^{14,19}. However, these studies primarily evaluated existing models, compared settings or examined local implementation. To our knowledge, no previous study has prospectively reconsidered and revised both the RCM and RFM through a multi-country de-centring and consensus process designed to preserve cross-cultural interoperability.

Cultural fit matters. A meta-analysis of 76 studies involving 25,225 participants found that culturally adapted mental health interventions showed a moderately strong advantage over unadapted approaches ($d = 0.45$)²⁰. A direct-comparison meta-analysis also found that culturally adapted psychotherapy outperformed unadapted psychotherapy ($d = 0.32$)²¹. Together, these findings indicate that cultural adaptation is not a peripheral concern: the language, examples and assumptions embedded in mental health interventions and implementation tools may materially affect whether they are understood, accepted and effective. Linguistic translation involves rendering wording from one language into another, whereas local cultural adaptation modifies language, examples or implementation to improve fit within a particular context²². Cross-cultural interoperability goes further by preserving shared conceptual meaning across contexts while allowing wording and examples to remain locally intelligible. Linguistic accessibility within countries is also relevant to equity, because people who use non-dominant languages may face additional barriers to accessing mental health services and information, participating in co-production, and engaging with recovery-oriented concepts²³. Linguistic justice extends beyond access alone: it also concerns whether institutional communication accommodates multilingual language practices in ways that enable meaningful participation, rather than privileging users of dominant languages²⁴. As mental health information and services are increasingly mediated through digital and AI-supported communication, linguistic accessibility also depends on whether these systems adequately accommodate linguistic diversity, rather than reproducing or amplifying existing inequalities^{25,26}. Because the RCM and RFM are intended to be used across diverse RC contexts, the key methodological challenge was not only translation or local adaptation, but cross-cultural interoperability: preserving shared conceptual meaning while allowing wording and examples to fit different cultural and service contexts. A consensus exercise was therefore chosen to integrate international stakeholder judgement on how the RCM mechanisms and

RFM items were interpreted across contexts, where agreement or disagreement indicated culturally different understandings, and how the wording should be refined to improve cross-cultural interoperability.

This study aimed to identify the level of cultural influence on the RCM mechanisms and RFM items, and to develop versions with greater cross-cultural applicability across the participating settings while retaining interoperability with the original tools.

Methods

This study followed the ACcurate CONsensus Reporting Document reporting guideline for consensus methods (Supplementary Table 2)²⁷. A study protocol was published in advance²⁸, and the study materials and anonymised underlying data are openly available on the Open Science Framework (<https://doi.org/10.17605/OSF.IO/CUAES>). The study was conducted in accordance with the Declaration of Helsinki. Ethical approval was obtained from the King's College London Research Ethics Psychiatry, Nursing and Midwifery Subcommittee (reference: MRM-24/25-47085). All participants provided informed consent before participating in the study.

Study design

We conducted a two-round international Delphi study to examine the cross-cultural applicability of the RCM and RFM. Delphi methods are widely used in health research to structure expert judgement where evidence is incomplete or contextual interpretation is required, and established guidance emphasises transparent panel selection, iterative feedback, anonymity and prespecified consensus procedures^{29,30}. The study followed Belton's six-step Delphi methodology³¹ and used a de-centring approach to cross-cultural research³². De-centring seeks to reduce construct and wording bias by treating the source-context version as open to revision rather than as a fixed or culturally neutral standard. Culture-specific assumptions are identified from multiple cultural perspectives, and concepts and wording are iteratively revised while preserving their intended meaning across contexts. In this study, de-centring was operationalised through international ratings of importance and cultural difficulty, qualitative feedback on interpretation and wording, corpus-informed linguistic analysis, research team review and Round 2 consensus testing. The Delphi aimed to identify: (1) the perceived importance of each RCM mechanism and RFM non-modifiable item, (2) the perceived cultural difficulty of achieving them, (3) recommended wording changes to improve cross-cultural applicability, and (4) which response category for each RFM modifiable item best aligned with local cultural context.

Participants and recruitment

Participants were recruited through the RECOLLECT International Research Consortium³³, which, at the time of the study, spanned 31 countries across Africa, Asia, Europe, North America, South America and Oceania. Country leads were invited to participate as panellists and to recruit additional eligible panellists within their national networks.

Eligibility criteria were: age 18 years or older and experience of an RC in any role. To avoid over-representation from individual countries, recruitment was monitored by country, and country leads were asked to stop recruitment once five responses had been received, although any additional responses already submitted were retained. Consent was implied through survey completion, in line with standard Delphi practice³⁴. Panellist anonymity was maintained through an individual online survey design, without access to the identities or responses of other panellists. This reduced the risk that panellists, particularly those with less experience or from countries where RCs are still emerging, might feel inhibited in expressing their views³¹. Names were given optionally for acknowledgement purposes and were accessible only to the Delphi administrator (YK), who had no voting rights in the consensus process.

Survey development and procedure

The draft survey was created by YK, reviewed by the research team, and piloted by country leads from Brazil, Estonia, Germany, Japan and Thailand

to improve clarity, linguistic accessibility, suitability for online completion, and cross-cultural relevance. Revisions following piloting included adding clarifying examples to potentially ambiguous terms, and incorporating optional free-text comment boxes alongside quantitative responses.

The finalised survey was used in Round 1 in English. Panellists rated the four RCM mechanisms and seven RFM non-modifiable items for (a) importance and (b) cultural difficulty of achieving them in their context. Ratings used five-point Likert scales (1 = 'Not at all' to 5 = 'Very important/difficult'), and for each item panellists could also suggest wording changes to improve cultural applicability. Panellists then indicated, for each of the five RFM modifiable items, which of the two response categories better aligned with the cultural context of RCs in their country, and could provide wording suggestions. Round 1 remained open from July to November 2025 (16 weeks), closing after 7 consecutive days without a new response.

The Round 1 survey was then modified as described in the next section to create the Round 2 survey. Round 2 remained open from January to February 2026 (5 weeks). Fortnightly reminders were sent to support retention. No financial incentives were used, as these may undermine the quality of panellist input in Delphi studies³¹. Panellists were offered social recognition through acknowledgement in the final paper³⁵.

Analysis

For quantitative importance and cultural difficulty ratings, consensus was predefined as 70% or more of responses falling within the same category after collapsing the original five-point scale into three categories. There is no universally accepted Delphi consensus threshold, and published studies use heterogeneous definitions and criteria; thresholds should therefore be pre-specified and justified³⁶. The 70% threshold was specified in the published protocol and selected to indicate a clear majority across a culturally diverse international panel. Responses of 1 and 2 were combined, 3 was retained as a middle category, and 4 and 5 were combined. This prespecified approach was used to reduce potential cross-cultural response bias³⁷, particularly variation in the use of extreme response options across cultural groups, and was supported by pilot findings indicating that a three-point structure would preserve meaningful distinctions while improving comparability across cultural groups²⁸.

Descriptive statistics were used to summarise responses, including frequencies, proportions, and medians (interquartile ranges (IQRs)). For RFM modifiable items, frequencies and proportions were calculated for Type 1 versus Type 2 responses. Items not reaching consensus in Round 1, or generating substantial qualitative concern, were revised and re-presented in Round 2. Quantitative findings identified where reconsideration was needed, whereas the free-text and corpus-informed analyses clarified why an item was difficult and informed how it should be revised; no formal numerical weighting was applied between these forms of evidence. Where feedback differed across cultural contexts, divergent interpretations were treated as evidence of ambiguity, and wording was revised to make the intended construct more explicit without adopting one culturally specific formulation as the standard.

Free-text responses in Round 1 were analysed using corpus-informed linguistic analysis, a software-assisted approach for examining large collections of digitised text, to identify recurrent terms and themes, patterns of language use, and ambiguities that informed wording changes to improve cross-cultural applicability. This approach differed from the published protocol²⁸, which had proposed descriptive content analysis, because the responses consisted of brief, item-specific comments from panellists across 20 countries. The corpus-informed approach enabled a more systematic and replicable analysis of language use than the one allowed by traditional manual qualitative thematic analysis of texts, particularly for large datasets. The automatic extraction of descriptive statistics, such as frequencies, together with the examination of terms of interest in their contexts of use, allowed recurrent patterns of meaning to be identified more readily and robustly. The analytic aim was to compare recurrent wordings and contextual meanings systematically across response sets rather than to develop a conventional thematic coding framework. Open-text responses were

grouped into datasets for each question type: cultural difficulty, components and mechanisms; importance, components and mechanisms; and cultural fit and wording.

Keywords, defined as words statistically salient in one dataset when compared to a reference dataset, were retrieved for each response dataset using the enTenTen21 corpus as reference (~52 billion words, available in Sketch Engine, the software used for the analysis). Keyword candidates were generated automatically by Sketch Engine and were selected based on their keyness scores. Sketch Engine calculates keyness using the simple maths method, which involves the comparison of the normalised frequencies of the words in the dataset of interest with those of the dataset of reference; this is an effect size measure, and it makes it possible to compare between corpora of different sizes. Each keyword was examined within its concordance lines (strings of text displaying the term of interest in context) to determine whether it indicated recurrent ambiguity, culturally specific interpretation or a wording preference relevant to item revision. Salient themes and insights into language use and preferences were identified inductively by examining concordance lines.

The initial concordance interpretation was conducted by SVL and YK and reviewed through discussion with the research team, including the developers of the RCM and RFM. No formal inter-rater reliability statistic was calculated because the analysis was used to inform wording revision rather than to produce an independently coded thematic framework. Divergent interpretations were treated as substantive evidence of possible cultural or semantic ambiguity rather than resolved solely through majority ratings. Item revisions drew on recurrent ambiguities and culturally loaded wording identified through keyword and concordance analyses, alternative formulations proposed by panellists, and research team interpretation. Findings and proposed revisions were reviewed by the research team, including the developers of the RCM and RFM, before the revised wording was presented in Round 2. The resulting revised wording was then evaluated by the Round 2 panel for understandability and cultural appropriateness across participating settings, with all 11 revised items reaching the pre-specified consensus threshold for both criteria. In Round 2, panellists were asked whether the proposed revisions were appropriate in their contexts and, if not, to suggest further changes. The limited additional suggestions made in Round 2 were reviewed by YK in discussion with co-authors, and three minor post-Round 2 wording refinements were implemented by the research team. All panellists were sent the finalised Global RCM v1 and Global RFM v1 after Round 2.

Results

A total of 54 panellists from 20 countries completed Round 1. Participants included 22 RC managers (40.7%), 10 RC researchers (18.5%), 9 RC peer trainers (16.7%), 7 RC non-peer trainers (13.0%), and 6 other RC staff (11.1%). In terms of RC experience, 13 panellists (24.1%) had 7+ years' experience, 13 (24.1%) had 4–6 years, 22 (40.7%) had 1–3 years, and 6 (11.1%) had less than 1 year. Geographically, panellists were mainly from Western Europe (42.6%) and Asia (29.6%), with additional representation from Central Europe (11.1%), Oceania (5.6%), North America (5.6%), South America (3.7%), and Eastern Europe (1.9%). In Round 2, 41 panellists (75.9%) from 15 countries participated. Geographical representation remained strongest from Western Europe (46.3%) and Asia (26.8%), followed by Central Europe (9.8%), North America (7.3%), Oceania (4.9%), and South America (4.9%). Table 1 presents the panellist characteristics.

Round 1 quantitative findings

All four RCM mechanisms reached consensus for importance, with 92.6–98.1% of panellists rating them in the 4–5 category after collapsing the five-point scale into three categories (Table 2). Six of the seven RFM non-modifiable items also reached consensus for importance (70.4–100.0%). The exception was tailored to the student, which fell just below the threshold (37/54, 68.5%). Median importance ratings were uniformly high (median 4–5).

By contrast, consensus on cultural difficulty was more limited. Crucially, no RCM mechanism reached consensus for low difficulty (1–2

Table 1 | Panellist characteristics (n = 54)

Country	Primary role	Years of experience	R2
Australia	RC staff	7+ years	✓
Australia	RC manager	1–3 years	✓
Brazil	RC manager	1–3 years	✓
Brazil	RC researcher	1–3 years	✓
Canada	RC researcher	7+ years	✓
Canada	RC manager	1–3 years	✓
Canada	RC manager	4–6 years	✓
Czech Republic	RC manager	4–6 years	✓
Czech Republic/ Slovakia	RC researcher	4–6 years	✓
Denmark	RC manager	7+ years	✓
Denmark	RC non-peer trainer	7+ years	✓
Denmark	RC peer trainer	1–3 years	
England	RC researcher	1–3 years	✓
England	RC manager	7+ years	✓
England	RC manager	1–3 years	
England	RC manager	7+ years	✓
England	RC manager	4–6 years	✓
England	RC staff	4–6 years	✓
England	RC manager	7+ years	
England	RC peer trainer	Less than 1 year	✓
England	RC peer trainer	4–6 years	✓
Estonia	RC researcher	Less than 1 year	
Finland	RC researcher	1–3 years	✓
Finland	RC manager	4–6 years	✓
Finland	RC non-peer trainer	Less than 1 year	✓
France	RC researcher	7+ years	✓
Germany	RC researcher	1–3 years	✓
Germany	RC manager	7+ years	
Germany	RC non-peer trainer	1–3 years	✓
Hong Kong	RC manager	4–6 years	
Hong Kong	RC researcher	7+ years	
Italy	RC manager	1–3 years	✓
Italy	RC manager	7+ years	✓
Japan	RC non-peer trainer	7+ years	✓
Japan	RC non-peer trainer	4–6 years	✓
Japan	RC peer trainer	1–3 years	✓
Japan	RC researcher	1–3 years	✓
Japan	RC manager	4–6 years	✓
Japan	RC manager	1–3 years	
Japan	RC non-peer trainer	1–3 years	✓
Japan	RC non-peer trainer	4–6 years	✓
Netherlands	RC researcher	Less than 1 year	✓
New Zealand	RC staff	1–3 years	
Spain	RC manager	7+ years	
Sweden	RC researcher	1–3 years	✓
Sweden	RC manager	Less than 1 year	✓
Sweden	RC manager	1–3 years	✓
Switzerland	RC manager	1–3 years	
Thailand	RC peer trainer	1–3 years	✓
Thailand	RC peer trainer	7+ years	✓

Table 1 (continued) | Panellist characteristics (n = 54)

Country	Primary role	Years of experience	R2
Thailand	RC peer trainer	Less than 1 year	
Thailand	RC peer trainer	4–6 years	
Thailand	RC peer trainer	4–6 years	✓
Thailand	RC staff	1–3 years	✓

RC non-peer trainer includes roles such as psychologist, nurse, social worker, or educator.
R2 participated in the Round 2 survey, RC Recovery College.

category). Among the RFM non-modifiable items, only Learning, Social connectedness, and Commitment to recovery reached consensus for low difficulty, with 70.4%, 70.4%, and 75.9% of panellists respectively rating them in the 1–2 category. Median difficulty ratings ranged from 2 to 3 across items, indicating that achieving these standards involves significant cultural navigation. This pattern suggests that panellists generally endorsed the underlying RC principles but experienced difficulty interpreting and applying their wording within local cultural and service contexts.

Round 1 qualitative findings informing revision

To understand how panellists interpreted the items across settings and to identify wording that could improve cross-cultural applicability, free-text responses from Round 1 (totalling 9541 words) were analysed using corpus-informed linguistic analysis. This analysis extracted ‘keywords’ (terms significantly salient in the focus dataset in comparison to a larger one of general English language) for each dataset of the free-text responses to identify recurrent terms and patterns of language use, and inductively identify themes. Examination of the keywords in context of use can identify ambiguities and points of tension in how panellists understood and applied the RC wording/terminology. Table 3 presents the main cross-cultural issues identified (Supplementary Table 3 for more details).

Free-text feedback on the five modifiable RFM items showed that the original binary Type 1/Type 2 response format did not adequately reflect all RC models. Panellists noted that some colleges combined both types within the same item, such as being partly community-based and partly located in a dedicated RC space, or offering course content both distinct from and integrated with clinical provision. Therefore, a third response option, ‘Both Type 1 and Type 2 (blended)’, was added for all modifiable items before Round 2.

Supplementary Table 4 presents the wording changes made for Round 2 in response to Round 1 input. Items that had already reached consensus as culturally not difficult (i.e. RFM 2 Learning, 5 Social connectedness, and 7 Commitment to recovery) were retained largely unchanged; where qualitative comments informed revision, these changes were minor and intended only to clarify wording or improve cross-cultural accessibility.

Round 2 evaluation of revised wording

In Round 2, all 11 revised items reached consensus for both understandability and cultural appropriateness (Table 4). For understandability, 73.2–95.1% of panellists rated the revised wording in the 4–5 category, and for cultural appropriateness the corresponding range was 80.5–95.1%. These findings supported the acceptability of the revised wording across settings.

Finalised cross-cultural version of the RCM and RFM items

In Round 2, all items reached consensus as culturally appropriate. Qualitative comments (1157 words) were limited and largely confirmatory. Three minor wording refinements were implemented by the research team (Supplementary Table 5). The final Global RCM Version 1 (Global RCM v1) and Global RFM Version 1 (Global RFM v1), alongside the original England-developed wording, are presented in Supplementary Table 6.

Table 2 | Round 1 ratings for importance and cultural difficulty of RCM mechanisms and RFM non-modifiable items (N = 54)

Item	Importance 4–5 n (%)	Importance Median (IQR)	Importance consensus met	Difficulty 1–2 n (%)	Difficulty Median (IQR)	Difficulty consensus met
RCM mechanisms						
Empowering environment	51 (94.4)	5 (4–5)	Yes	33 (61.1)	2 (2–3)	No
Shifting the balance of power	50 (92.6)	5 (4–5)	Yes	22 (40.7)	3 (2–4)	No
Enabling different relationships	52 (96.3)	5 (4–5)	Yes	28 (51.9)	2 (2–3)	No
Facilitating personal growth	53 (98.1)	5 (5–5)	Yes	34 (63.0)	2 (2–3)	No
RFM non-modifiable items						
1. Valuing equality	54 (100.0)	5 (5–5)	Yes	33 (61.1)	2 (2–3)	No
2. Learning	48 (88.9)	5 (4–5)	Yes	38 (70.4)	2 (2–3)	Yes
3. Tailored to the student	37 (68.5)	4 (3–5)	No	20 (37.0)	3 (2–3)	No
4. Co-production of the RC	49 (90.7)	5 (4–5)	Yes	27 (50.0)	2 (2–3)	No
5. Social connectedness	50 (92.6)	5 (4–5)	Yes	38 (70.4)	2 (2–3)	Yes
6. Community focus	38 (70.4)	4 (3–5)	Yes	24 (44.4)	3 (2–4)	No
7. Commitment to recovery	51 (94.4)	5 (5–5)	Yes	41 (75.9)	2 (1–2)	Yes

Consensus was predefined as ≥70% of responses falling within the same collapsed category. For importance, consensus was assessed in the 4–5 category. For difficulty, consensus was assessed in the 1–2 category. Medians and IQRs (interquartile ranges) are based on the original five-point scale.

Table 3 | Main cross-cultural issues identified in Round 1 free-text responses and corresponding wording revision

Item	Panellist feedback	Revision made
Empowering environment	Lacked a clear equivalent in some contexts; meaning was inconsistent	Emphasised strengths, autonomy, choice and control
Shifting the balance of power	Felt abstract, confrontational or hard to realise; cultural and structural barriers noted	Reframed as equal partnership
Enabling different relationships	Unclear what relationships were meant and with whom	Revised to focus on connecting and working together
Facilitating personal growth	Expectation of upward progress did not fit recovery in some contexts	Recentred on recovery rather than growth
Valuing equality	Tension between equality and individual needs; 'student' label queried	Emphasised accessibility, equity and clearer role descriptors
Learning	Adult learning approaches were unfamiliar in some settings; terminology was too jargonistic	Simplified and clarified in non-jargonistic terms
Tailored to the student	Interpreted variously as person-centred flexibility or cultural adaptation; resource constraints noted	Clarified as adapting to student needs within local context
Co-production of the RC	Meaning unclear; panellists requested explanation or examples	Described shared decision-making and contributor roles more explicitly
Social connectedness	Ambiguity between internal RC relationships and wider community links	Distinguished connectedness within the RC and beyond
Community focus	Unclear what kind of community was meant; organisational barriers noted	Made intended meaning of community more explicit
Commitment to recovery	Often understood as clinical rather than personal or social recovery	Clarified intended meaning of recovery

Discussion

This two-round international Delphi study used a cross-cultural de-centring approach, informed by corpus-informed linguistic analysis of panellist feedback, to examine the cross-cultural applicability of the RCM and RFM. We found that cultural influence operated less at the level of whether RC principles were valued, and more at the level of how those principles were understood, worded and enacted. Cross-cultural de-centring helped make implicit assumptions explicit, distinguish personal and social recovery from clinical recovery, and reframe ideas such as tailoring and partnership in ways that were meaningful and feasible across diverse contexts. This is timely in light of current international emphasis on person-centred, rights-based and community-based mental health care³⁸.

A central finding was that the underlying RC principles travelled better than the original English wording used to express them. In Round 1, all four

RCM mechanisms and six of seven non-modifiable RFM items reached consensus for importance, yet consensus on low cultural difficulty was much more limited. This pattern suggests that panellists across settings broadly endorsed the constructs, but not always the phrasing through which those constructs had been operationalised. In cross-cultural psychiatry terms, the challenge was less about conceptual rejection than about semantic portability and contextual intelligibility². The study therefore supports a distinction between valuing a principle and recognising it in a linguistic form shaped by a particular cultural and service discourse. This distinction is especially important for recovery-oriented frameworks, where apparently familiar terms may carry different institutional, clinical and moral meanings across settings³⁹.

The study also makes visible how tools developed in high-income Anglophone settings can contain assumptions that are locally intelligible but

Table 4 | Round 2 quantitative findings

Revised item	Understandable in context 4–5 <i>n</i> (%)	Median (IQR)	Consensus	Culturally appropriate 4–5 <i>n</i> (%)	Median (IQR)	Consensus
RCM mechanisms						
Empowering, autonomy-supporting environment	30 (73.2)	4 (3–4)	Yes	34 (82.9)	4 (4–4)	Yes
Moving toward an equal partnership	39 (95.1)	5 (4–5)	Yes	38 (92.7)	5 (4–5)	Yes
Connecting with peers and working together	37 (90.2)	5 (4–5)	Yes	37 (90.2)	5 (4–5)	Yes
Facilitating recovery	32 (78.0)	4 (4–5)	Yes	33 (80.5)	4 (4–5)	Yes
RFM non-modifiable items						
1. Valuing equality	39 (95.1)	4 (4–5)	Yes	37 (90.2)	5 (4–5)	Yes
2. Learning	37 (90.2)	4 (4–5)	Yes	36 (87.8)	5 (4–5)	Yes
3. Tailored to the student	36 (87.8)	4 (4–5)	Yes	36 (87.8)	4 (4–5)	Yes
4. Co-production of the RC	33 (80.5)	4 (4–5)	Yes	34 (82.9)	4 (4–5)	Yes
5. Social connectedness	39 (95.1)	4 (4–5)	Yes	39 (95.1)	5 (4–5)	Yes
6. Community focus	39 (95.1)	4 (4–5)	Yes	35 (85.4)	5 (4–5)	Yes
7. Commitment to recovery	39 (95.1)	5 (4–5)	Yes	39 (95.1)	5 (4–5)	Yes

Consensus was predefined as $\geq 70\%$ of responses in the 4–5 category.

not automatically universal. Terms such as ‘co-production’, ‘shifting the balance of power’ and ‘recovery’ are well established within UK recovery-oriented policy and service reform discourse, and can therefore appear self-explanatory to researchers and practitioners working within that context. Internationally, however, these terms were not always interpreted consistently and sometimes carried unintended meanings^{17,18}. For example, ‘shifting the balance of power’ assumes a discourse of explicit power redistribution that may resonate in Anglophone service reform language, but may sound confrontational, abstract or structurally unclear elsewhere⁴⁰. Likewise, although co-production is now widely used in UK health policy and personalised care discourse, it does not travel as a universally shared term without explanation⁴¹. One interpretation is that this wording reflected assumptions common in some Anglophone service cultures, including relatively explicit egalitarian language, individual autonomy and tolerance of open-ended participation^{42,43}. Across countries, panellists repeatedly sought clarification of meaning, role and relationship, suggesting that these assumptions were not consistently shared internationally. De-centring therefore did not replace the underlying RC principles; rather, it made their intended meaning and source-context assumptions more explicit across contexts. This aligns with wider concerns in cross-cultural psychiatry about dominant systems mistaking local categories for universal ones⁴⁴.

The item revisions illustrate how this de-centring process worked in practice. ‘Empowering environment’ became ‘Empowering, autonomy-supporting environment’, making explicit a meaning that had been too dependent on English-language recovery vocabulary. ‘Shifting the balance of power’ became ‘Moving toward an equal partnership’, replacing an abstract and potentially adversarial formulation with wording that was more relational and consistent with shared decision-making language. ‘Enabling different relationships’ became ‘Connecting and working together’, removing ambiguity around what ‘different’ meant. ‘Facilitating personal growth’ became ‘Facilitating recovery’, avoiding a potentially culturally loaded assumption that improvement must take the form of upward growth. Similar refinements occurred within the Fidelity Measure. ‘Professionals’ became ‘staff’, reducing dependence on systems in which ‘professional’ implies formal registration or status. ‘Tailored to the student’ was reframed to emphasise flexibility, reasonable adjustments and feasibility within local context and resources, rather than assuming a uniformly well-resourced service infrastructure. ‘Commitment to recovery’ was expanded to specify that recovery refers to personal and social recovery, not only clinical recovery or symptom reduction. Collectively, these changes moved

the framework away from locally saturated shorthand and toward wording that was more explicit, relational and transferable.

These findings have implications beyond RCs. Mental health tools developed in high-income Anglophone contexts may be presented as broadly applicable while still embedding culturally and institutionally specific assumptions about service organisation, professional identity, rights discourse and interpersonal style⁴⁵. This is a general risk when any national or regional model becomes influential internationally⁴⁶. The same de-centring approach could inform the adaptation of other mental health service tools involving concepts such as empowerment, recovery, shared decision-making, co-production, autonomy and community participation, whose meanings and practical expression may vary across cultural and service contexts. The present study shows why explicit de-centring is needed before such tools are used across diverse international settings. De-centring therefore has both methodological and ethical value: it improves interpretability while reducing the risk that culture-specific assumptions are universalised.

The revised wording provides the Global RCM v1 and Global RFM v1 for research, service development and dialogue across international RC settings. These v1 tools are not intended to replace the original England-developed versions, but to retain interoperability with them while supporting more consistent interpretation across countries. The Global versions are intended for cross-national studies and applications outside England, whereas the original versions may remain appropriate where continuity with England-based research or practice is required. The two versions were designed to preserve conceptual interoperability, but their empirical equivalence requires further comparative testing. Any additional linguistic or local adaptations should therefore be clearly documented and reported. This matters because international comparison requires enough continuity to compare findings, but enough de-centring to avoid imposing source-context assumptions uncritically. The Global RCM v1 and Global RFM v1 may therefore support cross-national RC research, local adaptation work, and policy dialogue in countries seeking to build recovery-oriented services without simply importing Anglophone terminology.

This study has several strengths. To our knowledge, the study is the first to apply an explicit multi-country de-centring process to both the RCM and RFM. It builds on previous RC fidelity, international comparison and local adaptation research by moving from describing cultural variation to prospectively revising the tools through international consensus. Many cross-cultural mental health studies adopt frontal comparison⁴⁷, in which tools or

interventions developed in one source context are assessed against one or more target contexts. This can be useful for identifying adaptation needs, but may also reproduce a source-versus-target logic in which non-Anglophone settings are mainly understood by how they differ from an Anglophone reference point. By contrast, this study enabled a more inclusive lateral comparison across multiple RC settings, examining how concepts were interpreted from multiple standpoints and generating findings with broader international relevance. Its rigour lies in the mixed-method design: an international Delphi across 20 countries, predefined quantitative consensus thresholds, corpus-informed linguistic analysis of free-text responses, and iterative research team consultation to ensure that revisions were both evidence-informed and conceptually coherent.

The study also has limitations. The sample was international but necessarily limited to countries where RCs exist or where relevant expertise was available; therefore, the study should not be interpreted as representing all world regions. However, given that RCs have been identified in 31 countries, participation from 20 countries represents substantial coverage of the current international RC community. Participation was still uneven across regions and countries, and some RC settings were represented by more panellists than others. Although recruitment was monitored by country and capped at five responses where possible, representation remained strongest from Western Europe and Asia. Consequently, the consensus may have been influenced more strongly by regions with larger samples and more established RC traditions, while perspectives from less represented regions may not have been fully captured. The consultation was conducted in English, which may have privileged panellists already familiar with Anglophone recovery discourse. English therefore functioned both as a methodological constraint and as part of the phenomenon under investigation, while also making visible some of the linguistic assumptions that the study sought to examine. Round 2 established consensus on understandability and cultural appropriateness of the revised English wording, but did not test translated versions or implementation in each context. Finally, because panellists were people already engaged with RCs, the study does not capture views from all stakeholders who may encounter these tools in wider policy or service environments.

A priority for future research is direct comparative testing of the empirical equivalence of the Global RCM v1 and Global RFM v1 with the original England-developed versions. Future work should test how the Global RCM v1 and Global RFM v1 perform in translated versions and specific national settings, including whether further adaptation is needed at the level of examples, scoring or guidance for interpretation. Comparative work could also examine whether some items continue to function differently across linguistic regions, service systems or legal-political contexts. More broadly, the study supports a research agenda in cross-cultural psychiatry that treats tool adaptation as a substantive epistemic task rather than a final technical step. In conclusion, this Delphi study suggests that RC principles are broadly valued across the participating settings, but that wording developed in a high-income Anglophone context required de-centring before it could become more internationally usable. The Global RCM v1 and Global RFM v1 therefore represent not a departure from the RC model, but a clearer articulation shown to be understandable and culturally appropriate across the settings represented in this study.

Data availability

Study materials and underlying data are openly available on the Open Science Framework: <https://doi.org/10.17605/OSF.IO/CUAES>.

Code availability

Not applicable.

Received: 23 May 2026; Accepted: 22 September 2026;

Published online: 06 October 2026

References

- Whitley, R., Shepherd, G. & Slade, M. Recovery Colleges as a mental health innovation. *World Psychiatry* **18**, 141–142 <https://doi.org/10.1002/wps.20620> (2019).
- Henderson, C. et al. Social inclusion of people with severe mental illness: a review of current practices, evidence and unmet needs, and future directions. *World Psychiatry* **25**, 56–82 <https://doi.org/10.1002/wps.70031> (2026).
- Hayes, D. et al. Organisational and student characteristics, fidelity, funding models, and unit costs of Recovery Colleges in 28 countries: a cross-sectional survey. *Lancet Psychiatry* **10**, 768–779 [https://doi.org/10.1016/S2215-0366\(23\)00229-8](https://doi.org/10.1016/S2215-0366(23)00229-8) (2023).
- Lin, E. et al. Developing an evaluation framework for assessing the impact of Recovery Colleges: protocol for a participatory stakeholder engagement process and cocreated scoping review. *BMJ Open* **12**, e055289 <https://doi.org/10.1136/bmjopen-2021-055289> (2022).
- Theriault, J., Lord, M. M., Briand, C., Piat, M. & Meddings, S. Recovery Colleges after a decade of research: a literature review. *Psychiatr. Serv.* **71**, 928–940 <https://doi.org/10.1176/appi.ps.201900352> (2020).
- Crowther, A. et al. The impact of Recovery Colleges on mental health staff, services and society. *Epidemiol. Psychiatr. Sci.* **28**, 481–488 <https://doi.org/10.1017/S204579601800063X> (2019).
- Ronaldson, A. et al. The impact of Recovery College enrolment on health service use and patient outcomes: retrospective matched cohort study using routinely collected data. Preprint at *medRxiv* <https://doi.org/10.64898/2025.12.09.25341905> (2025).
- Deloitte. The shift to prevention: Realising the socio-economic potential. <https://www.deloitte.com/uk/en/Industries/life-sciences-health-care/research/the-shift-to-prevention-realising-the-socio-economic-potential.html> (2025).
- Hayes, D. et al. Recovery Colleges Characterisation and Testing in England (RECOLLECT): rationale and protocol. *BMC Psychiatry* **22**, <https://doi.org/10.1186/s12888-022-04253-y> (2022).
- Toney, R. et al. Development and evaluation of a Recovery College Fidelity Measure. *Can. J. Psychiatry* **64**, 405–414 <https://doi.org/10.1177/0706743718815893> (2019).
- Kotera, Y., Ronaldson, A., Henderson, C. & Slade, M. Quantifying cultural distance to inform international transfer of mental health approaches. *World Psychiatry* (2027).
- Hayes, D. et al. Evidence-based Recovery Colleges: developing a typology based on organisational characteristics, fidelity and funding. *Soc. Psychiatry Psychiatr. Epidemiol.* **59**, 759–768 <https://doi.org/10.1007/s00127-023-02452-w> (2024).
- Kotera, Y. et al. Cross-cultural insights from two global mental health studies: self-enhancement and ingroup biases. *Int. J. Ment. Health Addict.* **23**, 3572–3581 <https://doi.org/10.1007/s11469-024-01307-y> (2025).
- Beckers, T. & Koopmans, M. Recovery Colleges as enablers of personal recovery: qualitative evaluation of the development of a Recovery College in the Netherlands. *BMC Psychiatry* **25**, 465 <https://doi.org/10.1186/s12888-025-06917-x> (2025).
- van Wezel, M. M. C. et al. Navigating in a value-driven practice: a study of a Dutch Recovery College as a learning, social, and organizational space. *Front. Psychiatry* **16**, 1625779 <https://doi.org/10.3389/fpsy.2025.1625779> (2025).
- Andersen, D., Waldemar, A. K. & Kirkegaard, S. Recovery College dropout: a qualitative study of external, relational and course-related dropout drivers in co-produced mental health care. *Int. J. Ment. Health Nurs.* **32**, 1346–1354 <https://doi.org/10.1111/inm.13173> (2023).
- Kotera, Y. et al. Cross-cultural comparison of Recovery College implementation between Japan and England: corpus-based discourse analysis. *Int. J. Ment. Health Addict.* **23**, 4362–4386 <https://doi.org/10.1007/s11469-024-01356-3> (2025).

18. Kotera, Y. et al. Recovery College advertising in the Netherlands: a Dutch-English-Japanese corpus-based comparison and its implications for quality-of-life orientation. Preprint at <https://doi.org/10.1101/2025.08.13.25333552> (2025).
19. Orsi, J. A. et al. Cultural adaptation of RECOLLECT fidelity measure and checklists in a recovery-oriented NGO in Brazil. *J. Ment. Health Train. Educ. Pract.* **21**, 125–139 <https://doi.org/10.1108/JMHEP-08-2025-0093> (2026).
20. Griner, D. & Smith, T. B. Culturally adapted mental health intervention: a meta-analytic review. *Psychotherapy* **43**, 531–548 <https://doi.org/10.1037/0033-3204.43.4.531> (2006).
21. Benish, S. G., Quintana, S. & Wampold, B. E. Culturally adapted psychotherapy and the legitimacy of myth: a direct-comparison meta-analysis. *J. Couns. Psychol.* **58**, 279–289 <https://doi.org/10.1037/a0023626> (2011).
22. Kotera, Y. et al. Construction and factorial validation of a Short version of the Attitudes Towards Mental Health Problems Scale (SATMHPS). *Ment. Health Relig. Cult.* (2022).
23. Kurlander, D., Lam, A. G., Dawson-Hahn, E. & de Acosta, D. Advocating for language equity: a community-public health partnership. *Front. Public Health* **11**, 1245849 <https://doi.org/10.3389/fpubh.2023.1245849> (2023).
24. Yi, R. Tech-empowered equity: advancing linguistic justice through digital scholarship. *Digit. Scholarsh. Humanit.* **40**, 381–399 <https://doi.org/10.1093/llc/fqaf007> (2025).
25. Lawrence, H. R. et al. The opportunities and risks of large language models in mental health. *JMIR Ment. Health* **11**, e59479 <https://doi.org/10.2196/59479> (2024).
26. Yi, R., Li, Z. & Zhang, W. *Algorithmic Fairness in AI-Mediated Institutional Communication* (Springer Nature, 2026).
27. Gattrell, W. T. et al. ACCORD (ACcurate COnsensus Reporting Document): a reporting guideline for consensus methods in biomedicine developed via a modified Delphi. *PLoS Med.* **21**, e1004326 <https://doi.org/10.1371/journal.pmed.1004326> (2024).
28. Kotera, Y. et al. Enhancing cross-cultural applicability in Recovery Colleges: a global Delphi study protocol. *PLoS ONE* **20**, e0332729 <https://doi.org/10.1371/journal.pone.0332729> (2025).
29. Niederberger, M. & Spranger, J. Delphi technique in health sciences: a map. *Front. Public Health* **8**, 457 <https://doi.org/10.3389/fpubh.2020.00457> (2020).
30. Jünger, S., Payne, S. A., Brine, J., Radbruch, L. & Brearley, S. G. Guidance on Conducting and REporting DElphi Studies (CREDES) in palliative care: recommendations based on a methodological systematic review. *Palliat. Med.* **31**, 684–706 <https://doi.org/10.1177/0269216317690685> (2017).
31. Belton, I., MacDonald, A., Wright, G. & Hamlin, I. Improving the practical application of the Delphi method in group-based judgment: a six-step prescription for a well-founded and defensible process. *Technol. Forecast. Soc. Change* **147**, 72–82 <https://doi.org/10.1016/j.techfore.2019.07.002> (2019).
32. van de Vijver, F. J. R. & Leung, K. *Methods and Data Analysis for Cross-Cultural Research* 2nd edn (Cambridge University Press, 2021).
33. Recovery Research Team. RECOLLECT International Research Consortium (RIRC). <https://www.researchintorecovery.com/recollect-international-research-consortium-rirc/> (2025).
34. Toma, C. L. & Picioreanu, I. The Delphi Technique: methodological considerations and the need for reporting guidelines in medical journals. *Int. J. Public Health Res.* **4**, 47 (2016).
35. Khodyakov, D., Grant, S., Kroger, J. & Bauman, M. (RAND Corporation, 2023).
36. Diamond, I. R. et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J. Clin. Epidemiol.* **67**, 401–409 <https://doi.org/10.1016/j.jclinepi.2013.12.002> (2014).
37. Kotera, Y., Tsuda-McCaie, F., Maughan, G. & Green, P. Cross-cultural comparison of mental health in social work students between UK and Ireland: Mental health shame and self-compassion. *Br. J. Soc. Work* **52**, 3247–3267 <https://doi.org/10.1093/bjsw/bcab240> (2022).
38. World Health Organization. *Guidance on Community Mental Health Services: Promoting Person-Centred and Rights-Based Approaches* (World Health Organization, 2021).
39. Kuek, J. H. L., Raeburn, T. & Wand, T. Asian perspectives on personal recovery in mental health: a scoping review. *J. Ment. Health* **32**, 517–533 <https://doi.org/10.1080/09638237.2020.1818709> (2023).
40. Kotera, Y. et al. 28-country global study on associations between cultural characteristics and Recovery College fidelity. *npj Ment. Health Res.* **3**, <https://doi.org/10.1038/s44184-024-00092-9> (2024).
41. Kotera, Y. et al. Cultural influences on fidelity components in Recovery Colleges: a study across 28 countries and territories. *Gen. Psychiatry* **38**, e102010 <https://doi.org/10.1136/gpsych-2024-102010> (2025).
42. Hofstede, G., Hofstede, G. J. & Minkov, M. *Cultures and Organizations: Software of the Mind* 3rd edn (McGraw-Hill Education, 2010).
43. Nazarian, A., Ranjbaran, A., Afifi, M., Atkinson, P. & Sadeghilar, N. Cross-cultural examination of effects of ethical leadership of independent hotels based on Social Exchange Theory. *Int. J. Cross Cult. Manag.* **24**, 463–484 <https://doi.org/10.1177/14705958241270817> (2024).
44. *Textbook of Cultural Psychiatry* 2nd edn (Cambridge University Press, 2018).
45. Kotera, Y. Culture as wind: aligning the direction of mental health interventions with cultural values. *Int. J. Ment. Health Addict.* <https://doi.org/10.1007/s11469-025-01592-1> (2025).
46. Kotera, Y. A qualitative investigation into the experience of neuro-linguistic programming certification training among Japanese career consultants. *Br. J. Guid. Couns.* **46**, 39–50 <https://doi.org/10.1080/03069885.2017.1320781> (2018).
47. Schnegg, M. & Lowe, E. D. *Comparing Cultures: Innovations in Comparative Ethnography* (Cambridge University Press, 2020).

Acknowledgements

The authors thank Marijn Antens, Simone Arbour, Ana Aznar Martínez, Anders Thorkil Bechgaard, Anne Mette Billekop, Lily Chan, Danielle Coppleson, Sally Easton, Matthew Jonathan Ellis, Daniela Fischer, Vinuri Fernando, Andrew Foster, Kumpon Kawkome, Kenji Kameda, Megumi Kurokami, Aya Kuroda, Ossi Kukkamäki, Ben Hart, Christine Larsen, Klaudia Lepski, Pernilla Lemvall, Living Recovery Center (Thailand), Sue Macdonald, Chiharu Nakayama, Andrew Nicholls, Jared Omundo, Apirak Paenmunin, Jörn Petersen, Supatchakarn Piroos, Geoff Turner, Flavio Trolese, Paola Santo, Gianpaolo Scarsato, Erica Wan, Louisa Whorrod, Alex Thomas, Aurelie Tinland, Jonna Tolonen, Wilasinee Pinyowit, Kruawon Tiengtom, Asuka Yoshimi and Yoshie Morita for their support with this study. M.S. acknowledges the support of NIHR Nottingham Biomedical Research Centre.

Author contributions

Conceptualisation: Y.K.; methodology: Y.K., S.V.L., V.L., C.H., M.S.; validation: Y.K., S.V.L., D.N., R.H., P.L., J.O., M.C.R.A., Y.M., U.B., J.P.P., V.R., M.N., T.J., V.L., S.T., A.R., V.K., A.K., P.B., C.H., M.S.; investigation: Y.K., S.V.L.; writing—original draft: Y.K., S.V.L.; writing—review and editing: Y.K., S.V.L., D.N., R.H., P.L., J.O., M.C.R.A., Y.M., U.B., J.P.P., V.R., M.N., T.J., V.L., S.T., A.R., V.K., A.K., P.B., C.H., M.S.; funding acquisition: C.H., M.S.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s44184-026-00250-1>.

Correspondence and requests for materials should be addressed to Yasuhiro Kotera.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026

This Article in Press is shared early to give you faster access to new research. It is citable and carries a permanent DOI. The final edited version will replace it automatically.

¹School of Health Sciences, Institute of Mental Health, University of Nottingham, Nottingham, UK. ²Center for Infectious Disease Education and Research, The University of Osaka, Suita, Japan. ³Department of Social Sciences, Azerbaijan University, Baku, Azerbaijan. ⁴Centre for Language and Communication Research, Cardiff University, Cardiff, UK. ⁵Faculty of Social Sciences, University of Tartu, Tartu, Estonia. ⁶Institute for Epidemiology and Medical Biometry, Ulm University, Ulm, Germany. ⁷Ratchasuda Institute, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand. ⁸Schizophrenia Program, Department of Psychiatry, Escola Paulista de Medicina, Universidade Federal de São Paulo (PROESQ - EPM/UNIFESP), São Paulo, Brazil. ⁹Brazilian Schizophrenia Association (ABRE), São Paulo, Brazil. ¹⁰Department of Psychology, Universidade Federal de São João del-Rei (UFSJ), São João del-Rei, Brazil. ¹¹Department of Psychiatric Nursing, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Japan. ¹²ReLife Centre, Lund University, Lund, Sweden. ¹³Centre for Mental Health Care Development, Prague, Czechia. ¹⁴Department of Social Work, College of Polytechnics Jihlava, Jihlava, Czechia. ¹⁵Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ¹⁶RECOLLECT Lived Experience Advisory Panel, London, UK. ¹⁷Health and Community Participation Division, Faculty of Nursing and Health Sciences, Nord University, Namsos, Norway. ¹⁸These authors contributed equally: Claire Henderson, Mike Slade. ✉ e-mail: Yasuhiro.kotera@nottingham.ac.uk