The value and impact of information provided through library services for patient care: A systematic review

Weightman AL^1 & Williamson J^2 ; on behalf of the Library & Knowledge Development Network (LKDN) Quality and Statistics Group.

- 1. WeightmanAL@cf.ac.uk. Information Services, Cardiff University, Cardiff CF14 4XN
- 2. jane.williamson@nclwdc.nhs.uk. North Central London Strategic Health Authority, 4th Floor Victory House, 170 Tottenham Court Road, London, W1T 7HA

ABSTRACT

Objective An updated systematic review was carried out of research studies looking at the value and impact of library services on health outcomes for patients and time saved by health professionals.

Methods A comprehensive systematic search was undertaken of the published literature to September 2003 in ERIC, LISA, Medline, PreMedline, Embase, the Cochrane Controlled Trials Register and Google. Some handsearching was carried out, reference lists were scanned and experts in the field were contacted. 28 research studies of professionally led libraries for healthcare staff, including clinical librarian projects, met the inclusion criterion of at least one health or 'time saved' outcome. Papers were critically appraised using internationally accepted criteria. Data were extracted and results were summarised using a narrative format since the studies were heterogeneous and precluded a statistical analysis.

Results There is evidence of impact from both traditional and clinical librarian services. The higher quality studies of traditional services measured impacts of 37-97% on general patient care, 10-31% on diagnosis, 20-51% on choice of tests, 27-45% on choice of therapy and 10-19% on reduced length of stay. Four studies of clinical librarian projects suggested that professionals saved time as a result of clinical librarian input, and two of these studies showed evidence of cost-effectiveness. However, the clinical librarian studies were generally smaller, with poorer quality standards.

Conclusions Research studies suggest that professionally-led library services have an impact on health outcomes for patients and may lead to time savings for health care professionals. The available studies vary greatly in quality but the better quality studies also suggest positive impacts. Good practice can be gathered from these studies to guide the development of a pragmatic survey for library services that includes the direct effects for patients among the outcome measures.

Key words: impact, value, librar*, information service, clinical librarian, health outcomes, systematic review

Implications for policy

- Evidence of the impact of professional library services on health outcomes for patients and time savings for health care staff is available and can be used to demonstrate the impact of library services to users, managers and funding bodies
- An opportunity to compare results across different types of health library service would be
 enhanced if a methodology, including a core set of questions, were agreed and validated
 at a national, or even an international, level

Implications for practice

- Practical guidance can be gathered from existing impact studies to improve the quality of further research
- A practical reliable survey methodology could also be used at a local level, to provide
 evidence of the impact of the library service on a range of outcomes and to guide further
 developments

The value and impact of information provided through library services for patient care: A systematic review

INTRODUCTION

Within the health service one of the key measures of the benefits accruing from a library service is the impact it has on improving patient care. Identifying the means to demonstrate this benefit to users, managers and funding bodies has long been recognised as a challenge, and there is on-going debate as to whether such benefits can be reliably measured.

It is notable that the Medical Library Association has steered away from attempts to measure the direct impact of library services on health outcomes¹. Their Standards for Hospital Libraries include such measures as 'frequent provision of information on which patient care decisions are based' and 'active membership of the librarian on performance improvement and patient safety teams'. Abels and colleagues² used a comprehensive systematic review to develop a taxonomy (ie measures and their surrogates) to assess the value of library and information services to hospitals and academic health institutions. Pretest interviewees noted that there are significant intervening variables for direct outcomes such as 'reducing length of hospital stay' and 'avoiding unnecessary tests' and suggested that there was no valid way of measuring the contribution from the library & information service. The outcome of such decisions meant that proposals for the measurement of the organisational goal of providing excellent clinical care were indirect measures such as 'support informed and timely decision making' and 'support the development of policies and procedures relating to clinical care'.

Despite the difficulties of measurement however, there is an expanding body of evidence from existing systematic reviews that information provided by a library service can influence patient care outcomes in various ways and assessment of impact at a local level is feasible^{3,4,5,6} even if reliant, as is often the case, on indirect measures. In addition, there is evidence that library services can lead to time savings for healthcare professionals and, thus, to cost savings and healthcare benefits^{4,5,6}. In the most recent systematic review of clinical librarian studies⁶, published during the data gathering stage of this review, the authors concluded that (using weighted averages from 12 studies) 65% of respondents felt that services contributed to better patient care. However, it was also noted that evidence of impact on the outcomes of patient care, cost-effectiveness and qualitative data were limited, and that more high quality research evidence was required.

Measuring a direct impact on patient care is therefore difficult, but tangible evidence of benefit is needed to justify continued expansion and investment in healthcare libraries and patient care outcomes should logically be included, along with the social and educational benefits of the library service.

Given the large amount of recent research in this area, the Quality and Statistics Group of the Library and Knowledge Development Network (LKDN) in the UK undertook to carry out an updated systematic review of the literature. The research objective was to review studies looking at the value and impact of library services on health outcomes for patients and time saved by health professionals. Information provided within traditional library settings, including searches carried out by librarians for users (mediated searches), and clinical librarian programmes are considered in this review since these services represent a continuum of provision and are not clearly distinct. It was decided to report on research studies looking at clinical librarian projects as a separate sub-group, to investigate if any general differences in the type(s) of service provision and outcomes emerge, with indications for good practice.

An additional aim of this review was to provide information on best practice to assist the LKDN Quality Panel in developing a pragmatic user survey tool that librarians might use to obtain a measure of the value and impact of the service in terms of patient care and time saved by health professionals. A paper describing this developmental work is in preparation.

METHODS

Search Strategy

Searches were carried out in January-March 2003 with update searches in September 2003 on the following databases: ERIC, LISA, Medline, PreMedline, Embase, Cochrane Controlled Trials Register, Google.

The search terms used were:

[[Performance indicator* OR performance measure* OR performance standard* OR impact OR value] AND [Health* librar* OR medical librar* OR postgraduate centre librar* or hospital librar* or nursing librar* or (health* close to information service*) or (medical close to information service*) OR virtual librar* OR electronic librar* OR electronic information]]

OR

[(clinical or medical or support or outreach) librarian* OR clinical information (professional* or specialist*) OR informationist* OR informaticist*]

In addition, handsearching of the *Health Information Libraries Journal* and *Journal of the Medical Library Association* for all issues in 2003 was carried out in September 2003.

An email was sent to the JISC-mail discussion lists lis-rlg and lis-medical in March 2003 requesting information on papers describing surveys or other methods to obtain a measure of the impact, or value, of library services in terms of health benefits for patients and members of the public. A summary of the project was published in the CILIP Health Libraries Group Newsletter in September 2003 asking for contributions. Specific requests were also made to centres involved with clinical librarian programmes.

Inclusion/Exclusion Criteria

Research papers and reviews of papers examining the impact of services provided from professionally led libraries for healthcare professionals, including clinical librarian projects, were sought. Studies had to have at least one outcome measure relating to health benefits for patients and/or members of the public, or to time saved by health professionals. Impact studies of 'virtual' (electronic) libraries were excluded unless these included traditional (librarian-run or mediated) services. The specific impact of information skills training


provided within, or outside, the traditional library service was excluded. This topic is covered by a recent systematic review based on a literature search to September 2002⁷. Specialist information services delivered to particular professional groups (eg primary care) outside the traditional library setting⁸ were also excluded.

Although the impact of library services³ and clinical librarian projects⁵ have been studied by recent systematic reviews, and these have provided a large number of relevant studies, it was felt that a sufficient number of additional research studies had been published since the literature search dates of these reviews to merit an update. When the Wagner review was published⁶ during the preparation of this review (literature review to 2001), the reference list was scanned but no further clinical librarian studies with patient outcomes were located.

Critical Appraisal

All types of research study were included. All papers that were potentially relevant according to information in the title and/or abstract were obtained in full. Relevant papers were then critically appraised using established methods ^{9,10,11,12,13} and summarised. The critical appraisal sheets and data extraction forms used can be obtained from the authors.

Additional publications located from reference lists, including those of existing reviews ^{3,4,5,6} were examined individually and appraised if they met the inclusion criteria for this review. Each paper was appraised in detail by one author and overseen by the other, with any concerns being resolved by discussion. A flow chart describing this process is given below.



During critical appraisal, in addition to collecting information on study type, total numbers and response rate, a crude measure of quality based on five additional criteria was established for each study. The criteria, which are based on established good practice for survey design^{10,12,13}, were:

- A 100% or truly random sample was used to ensure that a representative sample of the user group were surveyed.
- 2. The sample was **not** pre-selected. [Where subjects are given the opportunity to refuse to participate at outset, as in the Rochester study, volunteers are more likely to be those that value the library service –leading to the potential for a non-representative sample as a result of 'desirability bias'].
- The responses were anonymous, and the subjects were aware of this, to encourage honesty
- 4. The researcher(s) were independent of the library service under study to ensure an objective view of the data and anonymity of respondents.

5. The survey asked about a specific incidence of library use at, or soon after, the provision of the information (current) or users were asked to request information for a patient care query and to assess its value (prospective). In these circumstances bias from 'recall' errors made as a result of the lapse of time (for example the more likely recall of an influential piece of information) is potentially reduced.

If there was no clear answer to a particular query in the publication the quality measure was considered unmet.

Data Analysis

Data were extracted from each study to provide information on the research question, the study population, the study design, the main results and any comments on the potential sources of bias (confounders). The following specific information was provided on study design:

- 1. The method(s) of data collection questionnaire, interview and/or other techniques;
- 2. Whether questions were asked about library use in general or a specific incidence of library use; This latter method is sometimes called the critical incident technique;
- 3. Whether users were asked to consider information they had used in the past (retrospective), had received during the past seven days (current) or a request was made to respond to the survey following a future use of information (prospective).

Although a meta-analysis is theoretically possible, the research studies varied greatly in research design, setting, user group(s) and the way the questions were asked. Thus, a formal statistical analysis was not considered appropriate. The results are presented as a narrative summary with ranges of responses where possible.

RESULTS

Data from the included studies are given in Tables 1-2.

Table 1: Traditional library services

Paper [First author, date]	Question	Study population & research design	Main results Effects of the information provided as % of 'yes' responses with multiple responses allowed.	Comments/confounders
Ali, 2000 ¹⁴	How information provided by medical libraries can enhance day-to-day management of patients	Total population of medical specialists and registrars in two Canberra hospitals. • Questionnaire • Specific use • Prospective	Response rate 28% (80/288) Information provided did, or would contribute to better quality of care 82%; Would definitely handle some aspects of clinical situation differently 23%; Would probably handle some aspect of the clinical situation differently 52%; Identified important changes to care as a result 54% Used for/Changes made: Choice of management 57%, Advice to patients 47%, Prescribed medication 27%, Choice of lab tests 20%, Choice of other tests 24%, Diagnosis 10%, Reduced length of hospitalisation 10%. Avoidance of: Tests & investigation 23%, Surgical intervention 19%, Additional outpatient visits 12%, Hospitalisation 11%, Risk of hospital acquired infection 8%.	Respondents were asked to remain anonymous though survey may have been closely associated with library staff resulting in desirability bias.
Ashcroft, 1998 ¹⁵ [Excluded from further analysis]	To find out what value physiotherapists place on the information obtained through their hospital	Physiotherapists in 37 NHS Trust hospitals in the Northern and Yorkshire region of England.	Response rate 5% (73/1460) Used for/Changes made: Choice of treatment 53%; Amount of treatment received 35%; Criteria for treatment 39%;	The response rate was too small to generalise from these results and this study was excluded from further analysis. Excellent attempts were employed

	library service and whether the information had an impact on patient care.	QuestionnaireSpecific useProspective	Advice given 59%. Avoidance of: Greater inpatient treatment 16%; Additional outpatient hours 24%; Ineffective/inappropriate treatment 55%; Inappropriate referral 29%.	to reduce 'desirability bias', the questionnaire was piloted and is a potentially strong model for a further study.
Burton, 1995 ¹⁶	Do libraries in regional hospitals in New Zealand influence patient care?	All physicians in the six regional (200-450 bed) hospitals that agreed to participate. Rochester style survey but retrospective. • Questionnaire • General use • Retrospective	Response rate 81% (295/372) (useable responses 80%). Used for/Changes made: Diagnosis 93%; Advice to patients 54%; Lab tests 35%; Radiography requests 19%; Drug regimens 40%; Reduced length of stay 20%. Avoidance of: Additional procedures/tests 30%; Admission to hospital 10%; Transfer to another hospital 12%.	Good response rate. Survey was closely linked to library staff and may have lead to desirability bias. Responses probably anonymous but unclear.
Casado Uriguen, 1994 ¹⁷	What is the importance of library services in Spanish University Hospitals within the global context of the hospital and, in particular, the day to day work of physicians?	All doctors at five hospitals. Rochester style survey. Questionnaire General use (probably) Retrospective	Response rate 20% (799/3877). Used for/Changes made: Interpretation of signs and symptoms 27%; Differential diagnosis 58%; Treatment 61%; Drug regimens 26%. Avoidance of: Hospital admission 17%; Hospital acquired infection 25%; Additional tests 28%; Surgery 34%;	Published as conference proceeding only so limited information and no peer review. Confounding not discussed by authors. Low response rate but very large study. Closely associated with library and may have resulted in 'desirability bias'. Respondents probably anonymous.

			Follow-up outpatient visits 40%, Loss of physicians time (ie time saved) 49%.	
Chamberlain, 2003 ¹⁸ [Excluded from further analysis]	Why do mental health professionals request inter-library loans and to what use(s) do they put the articles received?	10 interviewees (three staff nurses, three psychologists, two student nurses, one SHO, one practice educator) from a psychiatric unit in a UK primary care trust. • Interview • Specific use • Current	Response rate N/A No requests were directly related to current patient care queries although two interviewees changed their clinical behaviour – one claiming that changes put into practice helped to reduce waiting list from two years to six months.	Very small, single-site, pilot study. Self reported. Requests essentially for academic assignments. Only two interviewees commented on patient outcomes – thus excluded from further analysis.
Davies, 1997 ¹⁹	What is the impact of information on the clinical knowledge and practice of nurses, midwives and health visitors?	A random sample of nurses, midwives and health visitors using UK health libraries. Two surveys. The second (impact) survey was attached to information provided by the library (inter-library loans and literature searches) • Questionnaire • Specific use • Current	Response rate 40% (311/776) Improved quality of life for patient/carers 68%; Influenced initial assessment of patient/client 45%.	Random sample, independent evaluation. Based on Urquhart (1995) – the IMPACT study

Johnson, 1995 ²⁰	Did mental health practitioners decide to make changes in patient care as a result of information obtained from library literature searches?	Multidisciplinary mental health professionals (psychiatrists, psychologists, nurses, pharmacists etc.) served by the Missouri Institute of Mental Health – 37% in hospital and remainder in other settings. All professionals who requested searches were included. • Questionnaire • Specific use	Response rate 59% (200/340) Used for/Changes made: Diagnosis 14%; Tests 13%; Choice of medication 13%; Choice of treatment 38%. Wide variations between professional groups (details in paper).	Closely associated with library with questionnaires returned to Library Director – unclear whether respondents could be identified from the questionnaire.
King, 1987 ²¹	What is the ability of the library to meet health professional needs and the impact of the information on the quality of patient care?	Retrospective Physicians, nurses and other health professionals from eight hospitals in the Chicago area requesting information relating to a current case or clinical situation. Questionnaire Specific use Prospective	Response rate: 57% (184/310) Information from the library contributed to higher quality care 94%; Definitely handled some aspect(s) of the case differently 22%; Probably handled some aspect(s) of the case differently 52%.	Attempts to limit bias: eg Librarians did not participate in the assessment process and were not able to identify the health professionals involved. No record was kept of those who declined to be involved and participants more likely to be supporters of the library.
Klein, 1994 ²² [Excluded from further analysis]	To examine the associations between the economic indicators of hospital costs, charges and length of stay for inpatient cases and the use of Medline	Physician requested Medline searches mediated by a professional librarian in three metropolitan hospitals in Detroit. Case comparison Specific use	Response rate N/A When compared to matched control cases, test cases where searches were carried out earlier had lower costs, charges and lengths of stay than those whose searches were carried out later. For those cases where earlier searches (conducted	Excellent attempt at objective assessment. Test cases were more severe than controls (but adjusted in the statistical analysis). The physician characteristics were not examined. A request for a

	searches for such cases.	Prospective and controlled. 192 test cases compared to 10,409 control cases where no identified Medline search was carried out. Only 34 grouped pairs were analysed	during first half of hospitalisation) were carried out, the following reductions were noted: Costs (to hospital) 70%; Charges (to patient) 68%; Length of hospital stay 65%.	search could have been associated with all-round excellent patient care and, thus, one of many factors that could influence outcomes. Not directly comparable to other studies so considered separately.
Lovas, 1991 ²³	To examine the purposes for which health professionals requested interlibrary loans, how the immediacy of need for items affected their usefulness, the effect on the health professional's work and whether the illustrations were an important part of the content of documents.	A stratified random sample of requests in Region Seven of the Inter-Library Loan Network: Arizona, California, Hawaii, Nevada, Pacific basin. 53% of ILLs were requested by physicians • Questionnaire • Specific use • Current	Response rate 82% (1412/1722) Documents were requested to support clinical care 51%; Percentage of documents requested for clinical care that were read 95%; Loan documents deemed to have had a major effect on the health professional's work 47%.	Participating libraries and health professionals were randomly selected. The survey was probably associated with the participating library and there may have been desirability bias. Patient care outcomes nonspecific.
Marshall, 1992 ²⁴	To determine the relevance of library services to patient care.	A systematically sampled set of physicians in 15 hospitals in the Rochester area (seven in the city and eight in the surrounding rural area) asked to request information relating to a current clinical case. • Questionnaire • Specific use • Prospective	Response rate 46% (208/448) 97% of the physicians said that information provided by their hospital librarians contributed to better-informed clinical decisions. Used for/Changes made: Diagnosis 29%; Choice of tests 51%; Choice of drugs 45%; Reduced length of stay 19%; Advice given 72%.	Systematic sampling with a random start from each hospital's lists but refusals to participate were not counted and desirability bias can't be avoided. Built on lessons learnt from Chicago study. Questionnaire was pretested. Many attempts to reduce bias – Research assistant carried out the study, the librarians did

			Avoidance of: Hospital admission 12%; Patient mortality 19%; Hospital acquired infection 8%; Surgery 21%; Additional tests or procedures 49%; Additional outpatient visits 26%.	not know who was selected and the physicians were asked not to reveal their involvement.
Quality Development Team, 2003 ²⁵	To analyse the effectiveness of the literature search service in terms of contribution to patient care and the professional and/or academic development of the user.	Questionnaire survey, following receipt of the results, of all users (from South Staffordshire NHS Trust and local Primary Care Trusts) requesting a literature search during a three month period. • Questionnaire • Specific use • Current	Response rate 66% (51/77) 89% of searches contributed to clinical decision making.	The survey was likely to be closely associated with library staff although the questionnaires were analysed by the Quality Group and the letter accompanying the questionnaire stressed that responses were confidential. Small sample size but good response rate. No specific patient outcomes.
Scolaro, 1995 ²⁶	To examine the impact of inter-library loans (ILLs) and literature searches provided by the library.	All requests for inter-library loans/literature searches at a hospital in Australia during a six month period. Users were from medical, nursing, allied health, administration and research professions. • Questionnaire • Specific use • Current	Response rate 37% (278/745) for ILLs and 36% (82/226) for literature searches. On a scale of one to four (1=low, 4 = high) 62% literature searches and 37% ILLs were graded three for impact on patient care (59% of literature searches and 34% ILLs had been requested specifically for patient care).	Published as conference proceeding. Low response rate but all requestors received a questionnaire. Carried out by library staff so desirability bias is a likely confounder but responses were anonymous. Single centre only. Patient outcome question is broad.
Webster Fisher, 1992 ²⁷	To establish total quality management benchmarks and to	The entire clinical staff of Norton Hospital and Kosair Children's Hospital, Kentucky.	Response rate 26% (543 useable/2101) 50% physicians reported that library information had	The research was associated with the library committee and desirability

	examine the significance of the role of the library in clinical care.	56% nursing, 23% allied professions, 21% clinicians. • Questionnaire • General use • Retrospective	an affect on diagnosis, choice of drugs or other therapy. 48% physicians reported an affect on diagnostic tests. 88% physicians, 58% allied health professionals and 69% nurses reported that information contributed to higher quality care.	bias cannot be excluded. However, the Planning and Marketing Department received and reviewed completed surveys to ensure anonymity. A copy of the questionnaire is available from the authors on request.
Urquhart, 1995 ²⁸	To examine the value to clinical decision making of information provided by NHS library and information services. To examine the factors affecting the delivery of that information to clinicians (the IMPACT study).	Clinicians at 11 hospital sites (over 600 clinicians and 43 health information library services provided data). • Questionnaire, interview and analysis of search requests • Specific use • Current	Response rate: Critical incident survey 46% (519/1133); Library searches and requests 68% (486/713); Dial up Medline 60% (60/100). From the critical incident analyses, patient care purposes were involved in 66% of incidents when clinicians sought information but in only 37% of these was patient care the sole purpose. From the library group Used for/Changes made: One or more categories of clinical decision making 79%; Recognition of abnormal or normal condition 36%; Identification/evaluation of alternative therapies 35%; Improved quality of life for patient or family 33%; Confirmation of proposed therapy 32%; Dfferential diagnosis 31%; Minimisation of risks of treatment 27%; Audit or standards of care 26%; Revision of treatment plan 25%; Choice of diagnostic test 22%.	Informed by Rochester and NLM studies. A variety of methods to gain information were used. Random samples of subjects were selected for the critical incident survey and dial up. The direct participation of site staff was precluded.
Wilson, 1989 ²⁹ also	To understand ways in which computer-mediated searching of MEDLINE affects	1,158 critical incident reports derived from 545 interviews with a purposive sample of professionals, mostly	Response rate: 48% of total known eligible interviewees (545/1135) 476 searches affecting patient care decisions	Huge cross US sample and attempts to reduce bias. Random sample, independent of libraries

published as Lindberg, 1993 ³⁰	patient care and other professional activities.	physicians. 65% were direct users of Medline throughout the US and 35% had Medline searches undertaken for them.	identified. Use of information (included multiple responses): diagnosis & etiology 41%, treatment and prognosis 51%, maintaining effective physician patient relationship 4%, disease prevention 3%, discharging responsibilities to third party payers 1%.	and anonymous. However, participants were asked to recall noteworthy rather than typical search.
		 Questionnaire & interview Specific use Retrospective 	Patient care outcomes. With a positive effect on: Longevity 5%, Abnormalities- 23%, Symptoms- 3%, Function- 1%. Improved diagnosis 16%, Improved treatment- 25%, Improved referral – 11.5%, Received care when otherwise may not have done – 0.3%, Process outcomes-66.5%, of which: Unnecessary.tests/procedures avoided –13%, Risks avoided./ reduced 7%, Hospitalisation reduced- 1.6%, Cost of care minimised or reduced – 1.3%, Insurance benefits – 4%	

Table 2: Clinical Librarian studies

Paper	Question	Study population & research design	Main results	Comments/confounders
Barbour, 1986 ³¹	An evaluation of the effect of the clinical librarian programme to the morning report.	Residents and intems of a general medical and surgical hospital in the US. Questionnaire General use Retrospective	Response rate 76% response (actual numbers not given). Literature of value in diagnosis or treatment. 95%; Saved time 86%.	Very little information provided about the research methods used. The sample size is not given and the study may have been closely associated with the librarian leading to desirability bias.
Booth, 2002 ³²	Evaluating impact on patient care e.g. changes in practice/ treatment/ therapy due to information provided by clinical librarians	Evaluation of two clinical librarian posts in a UK University Hospitals NHS Trust. 10% of baseline population in Directorates targeted by the clinical librarian service were assessed. • Questionnaire & interview • Specific use • Retrospective by questionnaire, based on the Value tool-kit using the critical incident technique, plus specific interviews with volunteers.	Response rate 58% (42/72) 27.8% of information from the literature was identified as having an impact on patient management Information provided assisted either significantly or moderately: In diagnosis 62%; In choice of intervention 85%; Length of stay 29%; Information given to patients 74%; Future clinical decisions 94%; Better understanding of patients condition 94% A cost saving (consultant vs librarian carrying out search) of £26 per hour spent searching was estimated. However the service was seen more as providing literature searching services than to answer clinical queries.	The study aimed to reduce bias. The evaluation was carried out by an independent research assistant and a random stratified sample of subjects were assessed. Confidentiality was assured to respondents.
Freeth, 2002 ³³	Evaluation of the impact of the work of the Clinical Support Librarian (CSL) on clinical practice. Evaluation of impact of	Users of CSL service in a multidisciplinary hospital library. • Questionnaire, interview & field notes	Response rate 35% (38/110) plus 26 semi-structured interviews with volunteers. Reports identifiable cost savings – switches to cheaper drugs and materials; reduced length of	The Evaluator was independent of the CSL but the interviewees were self selected, and recommended by the CSL. Risk of 'desirability bias'.

	the first post holder.	General use Retrospective	stay; reduction in readmissions. Actual cost savings not verified. Above said to result, at least in part, from CSL input Development of protocols, guidelines etc – though difficulties in implementation may reduce impact. Health care professionals' time saved.	Participants were assured that their responses would be confidential and anonymous however.
Freeth, 2003 ³⁴ (report in progress)	Evaluation of the impact of first post holder to the Clinical Effectiveness Librarian (CEL) Service.	Users of CEL service in a London NHS Trust. • Questionnaire plus telephone and face to face interviews • General use • Retrospective	Response rate 33% (49/155 questionnaires returned); plus seven telephone or face-to-face interviews. Staff recognised potential impact of CEL role on patient care, but identified a number of barriers to putting findings into practice. Several respondents noted that the CEL saved them time and that CEL could search more quickly.	The Evaluator was independent of the CEL but the interviewees were self selected, and recommended by the CEL. Risk of 'desirability bias'. Participants were assured that their responses would be confidential and anonymous however.
Giuse, 1998 ³⁵	To determine the extent to which clinicians trust the librarians ability to evaluate, interpret and synthesize the medical literature	38 clinical staff – 30 physicians, 2 nurses and 6 pharmacists – in a US medical library. • Questionnaire • General use • Retrospective	Response rate 100% (n=38) Using a Likert scale where 1=low and 10 = high marks were: 9.4 – accuracy of librarians interpretation of literature 7.8 - avoided some referrals 8.9 - improved patient care	A small sample size but good attempts to reduce bias; the questionnaire was carried out by an independent research assistant and results were blinded.
Royal, 1993 ³⁶	Whether information provided by the clinical librarian changed interpretation of the pathology in some cases	Medical students and pathology residents in a US University medical library. All residents and faculty who had participated were surveyed after six months of the clinical librarian program and a random sample of those requesting individual searches.	Response rate 60% (31/51) Respondents were asked to assess whether the information had changed their interpretation of the pathology in some cases using a Likert scale (1= high, 5=low): 1=45% 2= 29% 3= 19% 4=6%	The exact nature of the study is unclear and sources of bias may exist.

Schall, 1976 ³⁷	Whether the literature provided by the clinical librarian during ward rounds was of any value to clinically in determining the diagnosis or treatment patients?	Mixture of general and specific Current (in terms of literature searches) All staff present on rounds in the Neonatal Intensive Care Unit (NICU) and Dept of Orthopaedics during 8 months of project. Questionnaire General use Retrospective	Response rate 83% (30/36) from NICU, 58% (15/26) from Orthopaedics. Program was of clinical importance in determining the diagnosis and treatment of patents hospitalised (97% NICU, 67% Orthopaedics). In NICU, 60% of respondent said information was of use for 1-5 patients, whilst in Orthopaedics this only applied in 40% of cases.	Potential for desirability bias since the survey may have been closely associated with the service as an independent evaluation was not carried out. All users and potential users of the service were surveyed. Limited information provided as to how the study was carried out.
Scura, 1981 ³⁸	Whether information provided through the clinical librarian service affected patient management and, if yes, how was management influenced?	House Officers on medical and paediatric services in a US university hospital. Random sample of 50 searches from a total or 287 searches over five months. Interview (probably) Specific use Current	Response rate 96% (48/50) Patient management was affected in 20% of cases. Influenced either diagnosis or therapy. The cost of the service per question was estimated at considerably less than the cost of a chest x-ray.	The sample of interviewees was randomly selected. Response rate high. The survey was carried out by the clinical librarian and this may have led to 'desirability bias'.
Tobia 1983 ³⁹ [Excluded from further analysi s]	Can a clinical information consultancy service replace or supplement a clinical librarian service	House staff in the Dept, of Medicine, University of Texas Health Science Center • Questionnaire • General use	Response rate 29% (37/126). 100% said saved time (2.2 hours on average).	Continuation of service would be influenced by outcomes of study. Only one quality criterion confirmed (currency) and thus excluded from further

		Retrospective		analysis.
Ward, 2001 ⁴⁰ [Excluded from further analysis]	To test the feasibility of an outreach clinical librarian service in an acute hospital setting.	Evaluation of a 6 month pilot programme working with two clinical teams in a general hospital. All members of the teams were surveyed regarding the detailed evaluation of the service and individual questionnaires were also attached to each summary sent by the CL. • Questionnaire • General use • Retrospective	Response rate to individual questionnaires 10% (14/136) In another questionnaire asking for a detailed evaluation of the Service there were only 15 responses but 73% (8/11) of those who responded to the question indicated that the information provided had a direct impact on patient care.	Low response rate to individual questionnaire and response rate to more detailed questionnaire unknown. The group may have been self selected (desirability bias) and the evaluation may have been closely associated with the clinical librarian. Only one quality criterion confirmed (100% sample) and thus excluded from further analysis.
Veenstra, 1992a ⁴¹	Whether end user searching has lessened the need for clinical information specialists	1 st year interns, junior and senior residents and transitional 1 st year interns (33 in total) at a US hospital. • Questionnaire • General use • Retrospective	Response rate 91% (30/33) Of 45 physicians selected, 12 were unavailable. 93% respondents said that information provided by clinical librarian impacted on patient care in 40-59% of time. The clinical librarian found information that respondents had failed to find themselves.	Risk of response bias – some events had occurred up to 11 months earlier. Single site study – small number of respondents (n=33). The responses were anonymous but study may have been closely associated with the clinical librarian leading to 'desirability bias'.

Veenstra, 1992b ⁴²	Did a clinical librarian in the intensive care setting may provide information that assists House Officers with diagnosis and therapy in a cost-effective manner	House Officers in medical- coronary ICUs over a 3- month period. 66 patient- related questions were asked. • Questionnaire • Specific use • Current	Response rate 65% (43/66) 37.2% information aided in diagnosis, of which: • 3 – cancelled unnecessary tests • 11- helped with interpretation of tests • 1 – required further tests 51.2% aided therapy 30.2% led to better patient management The cost of finding the information on a question was approximately the same as the cost of a chest radiograph in 1999 (\$45, excluding any fax costs, versus \$50).	Single site study and a small sample size. All who asked questions were surveyed. Questionnaire was attached to the information provided by clinical librarian so risk of 'desirability' bias. However, an established service, no threat of withdrawal if poor results obtained.
----------------------------------	--	--	---	--

Only those studies meeting a response rate of 20% or more and at least two of the quality criteria (see Methods) were included within the results tables. One traditional and two clinical librarian studies were, thus, excluded at this stage^{15, 39,40}. An additional pilot study looking at a traditional library (inter-library loan) service was also excluded since only two respondents assessed the influence of the information provided on patient care¹⁸. It must be stressed that these quality measures are a crude assessment of quality. In addition, although it would have been ideal, authors of individual studies were not contacted to confirm or refute the quality criteria assessments. Details of included studies and measured outcomes are given in Tables 3 and 4. The study by Klein²² is considered separately and is not included in the results tables since this piece of research was an economic analysis as well as the only controlled study reporting patient outcomes (Table 1).

Table 3: Quality Measures of Included Studies.

	Traditional library												
	Urquhart 1995 ²⁸	Davies 1997 ¹⁹	Lovas 1991 ²³	Quality Devel 2003 ²⁵	King 1987 ²¹	Wilson 1989 ²⁹	Marshall 1992 ²²	Scolaro 1995 ²⁶	Ali 2000 ¹⁴	Webster F 1992 ²⁷	Burton 1995 ¹⁶	Johnson 1995 ²⁰	Casado Uriguen 1994 ¹⁷
Sample size	1133 CIT 713 Lit sear 100 Medine	776	1722	77	310	1135	448	745 ILL 226 Lit	288	2101	372	340	3877
Response rate (%)	46% CIT 68% Lit sea	40%	82%	66%	57%	48%	46%	37% ILL 36% Lit	28%	26%	81%	59%	20%
Additional quality criteria	5	<u> </u>				4				3		2	I

		Clinical librarian								
	Giuse 1998 ³⁵	Booth 2002 ³²	Scura 1981 ³⁸	Veenstra 1992a ⁴¹	Veenstra 1992b ⁴²	Royal 1993 ³⁶	Schall 1976 ³⁷	Barbour 1986 ³¹	Freeth 2002 ³³	Freeth 2003 ³⁴
Sample size	38	72	50	33	66	51	36 NICU 26 ortho	-	110	155
Response rate (%)	100%	58%	96%	91%	65%	60%	83% NICU 58% ortho	76%	35%	33%
Additional quality criteria		4 3						2	•	

Table 4: Impact Measures Reported Changes in Patient Care: Traditional Library

% response Multiple responses allowed in studies.	Urquhart 1995 ²⁸ [library searches]	Davies 1997 ¹⁹	Lovas ²³	Qual Devel Team 2003 ²⁵	King 1987 ²¹	Wilson 1989 ²⁹	Marshall 1992 ²⁴	Scolaro 1995 ²⁶	Ali 2000 ¹⁴	Webster Fisher 1992 ²⁷	Burton 1995 ¹⁶	Johnson 1995 ²⁰	Casado Uriguen 1994 ¹⁷	RANGE [Range ir studies meeting 4 quality cri
General impact on clinical care	79	-	47	89	94	-	97	37 ILL 62 Lit s	82	-	-	-	-	37-97% [37-97%
Handled case differently [definitely/probably]	25	-	-	-	74	-	-	-	75	-	-	-	-	25-75% [25-75%
Diagnosis	31	-	-	-	-	16	29	-	10	-	93	14	58	10-93% [10-31%
Choice of tests	22	-	-	-	-	-	51	-	20 lab 24 other	48	35	13	-	13-51% [20-51%
Choice of drugs/therapy	-	-	-	-	-	-	45	-	27	-	40	13	-	13-45% [27-45%
Diagnosis & Treatment/Manage- ment	32	-	-	-	-	25	-	-	57	50	-	38	61	25-61% [25-57%
Reduced length of stay	-	-	-	-	-	-	19	-	10	-	20	-	-	10-20% [10-19%
Advice to patients	-	-	-	-	-	-	72	-	47	-	54	-	-	47-72% [47-72%
Important changes in care/alternative therapies/referral	35	-	-	-	-	11	-	-	54	-	-	-	-	11-54% [11-54%
Initial assessment	-	45	-	-	-	-	-	-	-	-	-	-	-	-
Improved quality of life for patient/carer	-	68	-	-	-	-	-	-	-	-	-	-	-	-

Reported Benefits: Traditional Library

%	Wilson 1989 ²⁹	Marshall, 1992 ²⁴	Ali, 2000 ¹⁴	Burton, 1995 ¹⁶	Casado Uriguen 1994 ¹⁷	RANGE
Avoided:	1		•			•
Hospital admission	1.6	12	11	10	17	1.6-17%
Patient mortality	-	19	-	-	-	-
Hospital acquired infection	-	8	8	-	25	8-25%
Surgery	-	21	19	-	34	19-34%
Additional/unnessesary tests or procedures	13	49	23	30	28	13-49%
Additional outpatient visits	-	26	12	-	40	12-40%
Transfer to another hospital	-	-	-	12	-	-
or reduced risks	7	-	-	-	-	-
Saved:	ı	1			1	
Physicians' time	-	-	-	-	49	-

Table 5: Impact Measures: Clinical Librarian

Reported Changes in Patient Care: Clinical Librarian

Multiple responses allowed in studies. %	Giuse 1998 ³⁵	Booth 2002 ³²	Scura 1981 ³⁸	Veenstra 1992b ⁴²	Schall 1976 ³⁷	Barbour 1986 ³¹	Freeth 2002 ³³	Freeth 2003 ³⁴	RANGE
Impact on clinical care/ Higher quality care/better informed decisions	V	-	20	-	-	-	-	-	-
Diagnosis	-	62	-	37	-	95	-	-	37-95%
Choice of drugs/therapy	-	-	-	51	67 Ortho 97 NICU	-	-	-	51-97%
Diagnosis & Treatment/Management	-	85	-	30	-	-	-	-	30-85%
Reduced length of stay	-	29	-	-	-	-	v	-	-
Advice to patients	-	74	-	-	-	-	-	-	-

Reported Benefits: Clinical Librarian

Multiple responses allowed in studies. %	Giuse 1998 ³⁵	Booth 2002 ³²	Scura 1981 ³⁸	Veenstra 1992b ⁴²	Schall 1976 ³⁷	Barbour 1986 ³¹	Freeth 2002 ³³	Freeth 2003 ³⁴	
Avoided:	Avoided:								
Referral	V	-	-	-	-	-	V	-	
Re-admission	-	-	-	-	-	-	V	-	
Saved:	•	•	•	•	•	•	•		
Health professionals' time	-	V	-	-	-	86	V	V	
Money	 -	vcs	-	vcs	-	-	V	-	

v: Positive but un-quantified effect vCS: Estimates of cost savings were made (see text)

The results suggest clear evidence of an impact of library services on patient outcomes from both traditional and clinical librarian services. This is also the case in the higher quality studies (Tables 4 and 5). It is possible that there was a trend for these higher quality studies to report reduced impacts but because of the limited number of data points a statistical analysis lacked the necessary power to determine this. The higher quality traditional library studies (where four or more of the additional quality criteria were met) suggest effects of impacts of between 37 and 97% on general patient care; 10-31% on diagnosis, 20-51% on choice of tests; 27-45% on choice of therapy and 10-19% on the advice given to patients (Table 4).

The economic analysis by Klein²² concluded that, when compared to control cases, test cases associated with earlier literature searches had lower costs, charges and lengths of stay than those whose searches were carried out later (Table 1).

From the clinical librarian studies there are indications of direct patient benefits and, notably of time-saving for healthcare professionals and cost benefits.

DISCUSSION

This review provides updated evidence that health care libraries can influence patient outcomes in various ways. Evidence of impact is available both for traditional library services as well as for clinical librarian studies. There is also evidence, notably from the clinical librarian studies, of time savings to healthcare professionals and costbenefits. Of two studies attempting to measure the costs of the clinical librarian service, one found that the cost per question was approximately equivalent to that of a chest radiograph⁴²; Another estimated a cost saving of a clinical librarian versus a consultant search of £26 per hour in 2002³².

Study quality was variable within both research areas; Clinical librarian studies were generally smaller, with poorer quality standards and results were less likely to be quantified. Further study is therefore indicated. Criteria for limiting sampling, recall and researcher bias (additional quality criteria) were used, along with sample size and response rate for a crude ranking of studies (Table 3). Although it should be stressed that the relative importance of each of these criteria is not proven, a broad ranking of study quality could be estimated in this way. Good research practice also dictates that measures should be taken to reduce sources of bias within a research project as far as is possible, and this good practice has been established on an international level^{9,10,11}.

Only one randomised controlled trial was found during the literature search and it is notable that this was carried out more than 20 years ago⁴³! No patient outcomes were reported, so the study was excluded from this review. It does, however, provide an example as to how a randomised controlled trial can be carried out to examine a radical change to a library service. One included study contained a control group (Klein)²². Although a potential confounder of this study is that the cases associated with an early literature review were those that were more likely to have excellent all-round care (and thus the shorter stay in hospital could have been associated with a number of influences), it shows that the controlled study is feasible within this setting. Such methods shouldn't be excluded when research projects are under consideration, though the organisation and time required by such a study preclude its inclusion within a pragmatic user survey for general library use.

Conclusions from this study and from earlier literature reviews^{3,4} are that, in designing impact studies, qualitative as well as quantitative information should be considered.

Qualitative data, using appropriate methodologies to ensure rigorous gathering, analysis and presentation of data provide the grounding for quantitative approaches by

identifying useful indicators^{43,44,45,46}. The specific or general use design, with follow-up user interviews, provide a practical and appropriate methodology when all possible attempts to reduce bias are included.

Responses are inevitably reported subjectively by the health professionals and opportunities to acquire 'hard data' from patients are limited. The validity of results can be strengthened, however, by gathering and comparing quantitative data from a survey with supporting qualitative information from user interviews. Given the difficulties in collecting hard data on patient outcomes it is recommended that questions covering the effect of information on other important outcomes should be developed with the input of library users. These could include the effect on professional development, influence on key decision makers and/or in the preparation of local guidelines, and savings to the organisation including staff time^{1,2,4}.

If respondents are asked to reply on the basis of a specific & recent instance of library use/information provision, rather than library use in general, this reduces the complexity of the task being undertaken. The requirement for fixed responses oversimplifies the complex nature of decision making, but this is mitigated somewhat by ensuring that the responder is thinking of a single incident of information use rather than a range and may provide a more accurate reflection of need and use^{47,48,49}. This technique (the critical incident technique) has been validated and used in more than 700 studies²⁹. In addition, there are indications that different user groups should be considered or analysed separately. There may be important differences in benefit to particular user groups that might be excluded in an overall result³.

An opportunity to compare results across different types of health library services would be enhanced if a core set of questions were agreed and validated at the national

or even international level. For the measurement of user views on service quality, this has been achieved by the LibQUAL instrument, based on SERVQUAL developed by Texas A&M University Libraries⁵⁰ (http://www.libqual.org). These could then be shared across studies to allow for direct comparison of results. It can be seen from Table 4 that a range of different questions were asked. Language and terminology varied a great deal from study to study making a comparison of results across studies difficult. A core set of questions will be relevant to studies of both traditional library and clinical librarian services. Additional questions of particular relevance to clinical librarian services include those of time saved by health professionals and hence the effectiveness and cost-effectiveness of the service.

There are several suggestions for future research within the traditional library setting. Library services are moving rapidly towards electronic or 'virtual' services. Thus, it is important to carry out and pool studies that compare traditional (librarian/clinical librarian) searches carried out on behalf of health care staff with searches carried out by the health care staff themselves (end user searches) using networked services provided by the library, and for which support and training have been provided. Studies suggest that librarians usually retrieve more relevant citations than physicians and health professionals themselves have said that searches carried out by clinical librarians have added value compared to those that they carry out themselves³. The issues of efficiency and professional time saved/cost-effectiveness will then be an important outcome along with health outcomes in as far as these can be measured.

It would also be useful to compare the various models of clinical librarian services ranging from those offered to users coming into the library, attendance at ward rounds, on grand rounds and so on, to investigate the most valued and cost-effective methods of delivery in terms of benefits for healthcare staff and for patients.

This review looked only at one aspect, albeit a very important aspect, of the service provided by health care libraries. It is clearly appropriate to measure the impact of a health library service in terms of its effect on patient care despite the inherent difficulties of obtaining objective measures. Traditional library services will continue to provide, in addition, a resource for academic learning, research, administration and management, patient information, continuing professional and personal development, clinical teaching, guideline development and so on. Service quality and impact can be assessed by benchmarking services against measures of good practice as, for example, in the Health Libraries and Information Confederation's (HeLicon's) tool for the accreditation of library and information services in the health sector (http://www.nelh.nhs.uk/librarian/accreditation.asp). One important aspect of such an assessment is the direct impact of the service on the users and those they care for.

The LKDN Quality & Statistics Group is working with librarians in and beyond the UK to develop and validate a pragmatic but high quality user survey instrument to measure the effect of the impact of a library service on its users. The survey will include, among other important measures, questions regarding direct outcomes to patients, to get the full picture of the value and impact of a health library service.

The authors acknowledge with thanks, help from staff at Information Services, CILIP who carried out the database searches of LISA. We are also grateful to Christine Urquhart (University of Aberystwyth) and Patrick O'Connor (Yangulla Centre, Rockhampton, Australia) for providing a number of relevant papers and some very helpful advice during the preparation of this review. Finally we would like to acknowledge the invaluable assistance of Dr Hollie Thomas (University of Wales College of Medicine) for advice on statistics and measures of study quality.

REFERENCE LIST

- Gluck, J.C., Hassig, R.A., Balogh, L., Bandy, M., Donaldson, J.D., Kronenfeld, M.R., Lindner, K.L., Murray, K., Petersen, J., Rand, D.C. Standards for hospital libraries
 Journal of the Medical Library Association 2002, 90 (4), 465-472.
 http://www.pubmedcentral.gov/articlerender.fcgi?artid=128964
- 2. Abels, E.G., Cogdill, K.W., Zach, L. The contributions of library and information services to hospitals and academic health sciences centers: a preliminary taxonomy. *Journal of the Medical Library Association* 2002, **90** (3), 276-284
- 3. O'Connor, P. Determining the impact of health library services on patient care: a review of the literature. *Health Information and Libraries Journal* 2002, **19** (1), 1-14
- 4. Urquhart, C.. How do I measure the impact of my service? (Guideline) pp 210-222 of Booth, A. and Brice, A. (eds). Evidence-based practice for information professionals: a handbook. London: Facet Publishing, 2004
- 5. Winning, M.A., Beverley, C.A. Clinical librarianship: a systematic review of the literature. Health & Information Libraries Journal 2003, **20** (Suppl), 10-21
- 6. Wagner, K.C., Byrd, G.D. (2004) Evaluating the effectiveness of clinical medical librarian programs: a systematic review of the literature. *Journal of the Medical Library Association* 2004, **92** (1), 14-33
- 7. Brettle, A. Information skills training: a systematic review of the literature. *Health Information and Libraries Journal* 2003, **20** (suppl.1), 3-9

- 8. National Institute of Clinical Studies. The effectiveness of clinical information services. Prepared by the Centre for General Practice and the Centre for Evidence-Based Practice at the University of Queensland. Melbourne: National Institute of Clinical Studies, 2002
- NHS Centre for Reviews & Dissemination. Undertaking systematic reviews of research on effectiveness. University of York: NHS Centre for Reviews & Dissemination, 2001
- Polgar, A., Thomas, S.A. Critical evaluation of published research. Chapter 22 <u>In</u>
 Introduction to research in the health sciences. 3rd edition. Melbourne: Churchill
 Livingstone, 1995
- Weightman, A.L., Barker, J., Lancaster, J.. Health Evidence Bulletins Wales. A
 Systematic Approach to Identifying the Evidence. Project Methodology 4. Cardiff:
 University of Wales College of Medicine, 2001
- Burton, D. (ed.) Research training for social scientists. London: Sage Publications, 2000
- 13. de Vaus, D.A. Surveys in social research. 3rd ed. London: Allen & Unwin, 1991
- 14. Ali, I. Library provided information and clinical decision making: a study of two hospitals in Canberra. *Australian Academic and Research Libraries* 2000; **31**, 30-45
- 15. Ashcroft, M. The impact of information use on decision making by physiotherapists. *Library Management* 1998, **19** (3), 174-195

- 16. Burton, J.E. The impact of medical libraries and literature on patient care in New Zealand. *Bulletin of the Medical Library Association* 1995, **83**, 425-430.
- 17. Casado Uriguen, M., Garcia Martin, M.A., La Torre, P.M., Montes del Olmo, M.I., Mas Vilardell, T., Ribes Cot M.. Importance of hospital library facilities to clinical decision making by medical staff. In McSean, T., Van Loo, J. and Coutinho, E., (eds.) Health Information New Possibilities: Proceedings of the 4th European conference of medical and health libraries held on 28 June- 2 July 1994, Oslo, Dordrecht, Kluwer for EAHIL, 1995. pp. 126-128.
- 18. Chamberlain, D. A pilot study on the use of inter-library loans by clinical staff. Health Information and Libraries Journal 2003, **20** (4), 235-239
- 19. Davies, R., Urquhart, C.J., Smith, J., Massiter, C., Hepworth, J.B. Establishing the value of information to nursing continuing education: report of the EVINCE project (British Library Research and Innovation Report 44). Boston Spa: British Library, 1997. Also published as: Urquhart C, Davies R. EVINCE: The value of information in developing nursing knowledge and competence. *Health Libraries Review* 1997, **14** (2), 61-72
- 20. Johnson, M. The library as a resource for decision making in mental health care. *Psychiatric Services* 1995, **46** (5), 493-495
- 21. King, D.N. The contribution of hospital library information services to clinical care: a study in eight hospitals, *Bulletin of the Medical Library Association* 1987, **75** (4), 291-301

- 22. Klein, M.S., Ross, F.V., Adams, D.L., Gilbert, C.M. Effect of online literature searching on length of stay and patient care costs. *Academic Medicine* 1994, **69** (6), 494.
- 23. Lovas, I., Graham, E., Flack, V. Health professionals' use of documents obtained through the Regional Medical Library Network. *Bulletin of the Medical Library*Association 1991, **79** (1), 28-35
- 24. Marshall, J.G. The impact of the hospital library on clinical decision making: the Rochester study. *Bulletin of the Medical Library Association* 1992, **80**, 291-301
- 25. Quality Development Team; Library Service Team at St. Michael's Hospital, St. George's Hospital. A survey to assess the impact of mediated/facilitated literature searches on patient care. Stafford: South Staffordshire NHS Trust, October 2003
- 26. Scolaro, T. Assessing the impact of information provided by the library. In:

 Keyword Editorial Services, eds. Synergy in Sydney 1995. Proceedings of the Sixth

 Asian Pacific Specials, Health and Law Librarians Conference. Canberra: Australian

 Library and Information Association, 1995 pp.95-109.
- 27. Webster Fischer, W., Reel, L.B. Total quality management (TQM) in a hospital library: identifying service benchmarks. *Bulletin of the Medical Library Association* 1992, **80** (4), 347-352
- 28. Urquhart, C.J., Hepworth, J.B. *The value to clinical decision making of information supplied by NHS library and information services. British Library Research and Development Report 6205.* London: British Library Research and Development

Department, 1995.

Also published as: Urquhart, C.J, Hepworth, J.B. Comparing and using assessments of the value of information to clinical decision making. *Bulletin of the Medical Library Association* 1996, **84** (4), 482-489

Also published as: Urquhart, C., Hepworth, J. The value of information supplied to clinicians by health libraries: devising an outcomes-based assessment of the contribution of libraries to clinical decision making. *Health Libraries Review* 1995, **12**, 201-213

- 29. Wilson, S.R., Starr-Schneidkraut, N., Cooper, M.D. National Library of Medicine
 Office of Health Information Programes Development (2001) Use of the critical incident
 technique to evaluate the impact of Medline [online] Bethesda, National Library of
 Medicine, September 1989 http://www.nlm.nih.gov/od/ope/cit.html
- 30. Lindberg, D.A.B, Siegel, E.R., Rapp, B.A., Wallingford, K.T., Wilson, S.R. Use of MEDLINE by physicians for clinical problem solving. *Journal of the American Medical Association* 1993, **269** (24), 3124-3129
- 31. Barbour, G.L., Young, M.N. Morning report: role of the clinical librarian. *Journal of the American Medical Association* 1986, **255** (14), 1921-2
- 32. Booth, A., Sutton, A., Falzon, L. Evaluation of the clinical librarian project: University Hospitals of Leicester NHS Trust. Sheffield ScHARR, 2002
- 33. Freeth, D.S, (2002) Evaluation of the role of Clinical Support Librarian. London, City University. Internal Research Report.

- 34. Freeth, D.S., Smith, H. (2003) Evaluation of the role of clinical effectiveness librarian. London, City University (report in progress), Internal Research Report.
- 35. Giuse, N.B., Kafantaris, S.R., Miller, M.D., Wilder, K.S., Martin, S.L., Sathe, N.A., Campbell JD. Clinical medical librarianship: the Vanderbilt experience. *Bulletin of the Medical Library Association* 1998, **86** (3), 412-416
- 36. Royal, M., Grizzle, W.E., Algermissen, V., Mowry, R.W. The success of a clinical librarian program in an academic autopsy patholody service. *American Journal of Clinical Pathology* 1993, **99** (5), 576-81
- 37. Schall, J.G., Wilson, J.W. Evaluation of a clinical medical librarianship program at a University Health Sciences Library, *Bulletin of the Medical Library Association* 1976, **64** (3), 278-283
- 38. Scura, G., Davidoff, F. Case-related use of the medical literature. Clinical librarian services for improving patient care. *Journal of the American Medical Association* 1981, **245** (1), 50-52
- 39. Tobia, R.C., Kronick, D.A. A clinical information consultation service at a teaching hospital. *Bulletin of the Medical Library Association* 1983, **71** (4), 396-399
- 40. Ward, L.M., Honeybourne, C.J., Harrison, J. A clinical librarian can support clinical governance. *British Journal of Clinical Governance* 2001, **6** (4), 248-251
- 41. Veenstra, R.J. (1992a) Clinical medical librarian impact on patient care: a one year analysis. *Bulletin of the Medical Library Association* 1992, **80** (1), 19-22

- 42. Veenstra, R.J., Gluck, E.H. (1992b) A clinical librarian program in the intensive care unit. *Critical Care Medicine* 1992, **20** (7), 1038-1042
- 43. Marshall, J.G., Neufeld, V.R. A randomized trial of librarian education participation in clinical settings. *Journal of Medical Education* 1981, **56**, 409-416.
- 44. Wavell, C., Baxter, G., Johnson, I., Williams, D. Impact evaluation of museums, archives and libraries: Available evidence project. Aberdeen: Aberdeen Business School, The Robert Gordon University, 2002.

Resource study and impact portal. Impact Evaluation Homepage: http://rgusim.rgu.ac.uk/impact/homepage/sidebar.htm

- 45. Crowley, G.H., Gilreath, C.L. Probing user perceptions of service quality: using focus groups to enhance quantitative surveys. *Performance Measurement and Metrics* 2002, **3** (2), 78-84
- 46. Usherwood, B. Demonstrating impact through qualitative research. *Performance Measurement and Metrics* 2002, **3** (3), 117-122
- 47. Flanagan, J.C. The critical incident technique. *Psychological Bulletin* 1954, 51 (4), 327-58
- 48. Kunz, W., Rittel, H.W.J., Schwuchow, W. Methods of analysis and evaluation of information needs: a critical review. Munich: Verlag Dokumentation, 1977
- 49. Martyn, J., Lancaster, F.W. Investigative studies in library and information science: an introduction. Arlington: Information Resources Press, 1981

- 50. Parasuraman, A., Berry, L.L., Zeithaml, V.A. Refinement and reassessment of the SERVQUAL scale. *Journal of Retailing* 1991, **67** (4), 420-450
- 48. Cook, C., Heath, F., Thompson, R.L., Thompson, B. Score reliability in web- or internet-based surveys: unnumbered graphic rating versus Likert-type scales. Educational and Psychological Measurement 2001, **61**, 697-706

[Word count = 9,776]

Excluded Studies

Paper	Reason for
	exclusion
Urquhart, C., Yeoman, A., Cooper, J., Wailoo, A. NeLH pilot	Electronic library
evaluation project. Final	only
repowww.nhsia.nhs.uk/nelh/pages/documents/aber.doc	
Wyatt, J. Development of an evaluation methodology for the	Electronic library
NSW Health Clinical Information Access Program, Sydney, New	only
South Wales Health Department, 2001, available at	
www.health.nsw.gov.au	
Yeoman, A., Cooper, J., Urquhart, C., Tyler, A. <i>The value and</i>	Virtual outreach
impact of virtual outreach services: report of the VIVOS project,	services.
LIC 111, London, Resource: The Council for Museums, Archives	Electronic library
and Libraries, 2001.	and mailings of

	evidence-based
	topics.
Abels, E.G., Cogdill, K.W., Zach, L. The contributions of library	A taxonomy of
and information services to hospitals and academic health	possible impact
sciences centers : a preliminary taxonomy. Journal of the	measures
Medical Library Association 2002, 90 (3), 276-284	developed by
	consensus
	opinion.
Gluck, J.C., Hassig, R.A., Balogh, L., Bandy, M., Donaldson	Standards
Doyle, J., Kronenfeld, M.R., Lindner, K.L., Murray, K., Petersen,	developed by
J., Rand, D.C. Standards for hospital libraries 2002. Journal of	consensus
the Medical Library Association 2002, 90 (4), 465-472	opinion
[Unpublished Study]	Electronic library
Riley, L. The influence of the use of direct 24-hour access to	resources only
electronically-networked information on clinical decision making.	
Gunning, J.E., Fierberg, J., Goodchild, E., Marshall, J.R. Use of	No health/time
an information retrieval service in an obstetrics/gynecology	saved outcomes
residency program. Journal of Medical Education 1980, 55,	
120-123	
Greenhalgh, T., Hughes, J., Humphrey, C., Rogers, S.,	No health/time
Swinglehurst, D., Martin, P. A comparative case study of two	saved outcomes
models of a clinical informaticist service. British Medical Journal	
2002, 324, 524-529	
Pifalo, V.; Hollander, S.; Henderson C.L. The impact of	Information

consumer health information provided by libraries: the Delaware	provided directly
experience. Bulletin of the Medical Library Association 1997, 85	to consumers
(1), 16-22	rather than
	health care staff
Brassey, J., Elwyn, G., Price, C., Kinnersley, P.	Specialist rapid
Just in time information for clinicians: a questionnaire evaluation	response
of the ATTRACT project. British Medical Journal 2001, 322	information
(7285), 529-30	service for
	primary care
	provided outside
	the traditional
	library setting
Silverstein, J.L. Strengthening the links between health	No health/time
sciences information users and providers. Bulletin of the	saved
Medical Library Association 1995, 83 (4), 407-417	outcomes. User
	survey
	questionnaire
	provided.
D'Alessandro, M., D'Alessandro, D.M., Galvin, J.R., Erkonen,	Electronic library
W.E. Evaluating overall usage of a digital health sciences	resources only
library. Bulletin of the Medical Library Association 1998, 86 (4),	
602-609	
Marshall, J.G., Hamilton, J.D The clinical librarian and the	No health/time
patient: report of a project at McMaster University Medical	saved outcomes
Centre. Bulletin of the Medical Library Association 1978, 66 (4),	

420-425	
Ash, J.S., Gorman, P.N., Lavelle, M., Lyman, J., Delcambre,	No health/time
L.M., Maier, D. Bundles: meeting clinical information needs.	saved outcomes
Bulletin of the Medical Library Association 2001, 89 (3), 294-6	
Reid, L., Ikkos, G., Hopkins, W. Clinical librarians at Barnet	No health/time
Primary Care NHS Trust: addressing the requirements of clinical	saved outcomes
governance. Health Information Libraries Journal 2002, 19, 52-	
55	
Watson, J.A., Weist, A. The Forest Healthcare Clinical Support	No health/time
Librarian: 6 months on. Health Information Libraries Journal	saved outcomes
2000, 17, 219-221	
Martin, M., Kauser, A. informaticist working in primary care: a	Not librarian-led
descriptive study. Health Informatics Journal 2001, 7,66-70	
Shearer, B., Seymour, A., Capitani, C. Bringing the best of	Reviews
medical librarianship to the patient team. Journal of the Medical	potential role of
Library Association 2002, 90 (1), 22-31	informationist.
	No patient care
	outcomes.
Staggers, N., Thomas, C., Happ, B. An operational model for	IT systems
patient-centred information. Computers in Nursing 1999, 17 (6),	model
278-285	
Swinglehurst, D.A., Pierce, M., Fuller, J.C.A A clinical	No health/time
informaticist to support primary care decision making. Quality in	saved outcomes
Health Care 2001, 10 (4), 245-9	

Williams, L., Zipperer, L. Improving access to information:	Personal views
librarians and nurses team up for patient safety. Nursing	of how librarians
Economics 2003, 21 (4), 199	can support
	nurses
Schacher, L.F. Clinical librarianship: its value to medical care	Historical review
Annals of Internal Medicine 2001, 134 (8), 717-720	
Davidoff, F., Florance, V. The informationist: a new health	Background to
profession? Annals of Information Medicine 2003, 132 (12),	development of
996-8.	clinical librarian
	role
Sackett, D.L., Straus, S.E. Finding and applying evidence	Not librarian-led
during clinical rounds: the "evidence cart". Journal of the	
American Medical Association 1998, 280, 1336-1338	
Richwine, M., McGowan, J.J., Lilly, R. A rural virtual health	Electronic only
sciences library project: research findings and implications for	
next generation library services. Bulletin of the Medical Library	
Association 2001, 89 (1), 37-44	
House, M.A. Effectiveness indicators for the RC Laird Health	No health/time
Sciences Library. Bibliotheca Medica Canadiana 1991, 13 (1),	saved outcomes
53-61	
Banks, R. Measuring the impact of a hospital library in terms of	No health/time
value added processes. Bibliotheca Medica Canadiana 1989,	saved outcomes
10 (4), 184-192	
Hardy, M.C., Yeoh, J.W., Crawford, S. Evaluating the impact of	Not a research

library services on the quality and cost of medical care. Bulletin	study
of the Medical Library Association 1985, 73 (1), 43-46	
Marriot. R. Yes but how do we know if it's working? Evidence	Electronic only
regarding impact on clinical practice of access for health service	
staff to bibliographic databases and full text electronic journals.	
Library Review 2002, 51 (7), 358-363	
Miller, N. Effects of cost sharing and end-user searching on a	No health/time
clinical medical librarian program. Bulletin of the Medical Library	saved outcomes
Association 1989, 77 (1), 71-2	
Wilkin, A. Evaluation of a clinical librarian experiment. Report	No health/time
to the British Library Research & Development Department on	saved outcomes
Project S1?G/262. 1982 128pp	
Moore, A. The clinical librarian in the Department of Surgery: a	No health/time
report of two years' experience. Report to the British Library	saved outcomes
Research & Development Department on Project S1/G/262.	
1980. 75pp.	
Lusher, A. Getting evidence to the bedside: the role of the	No health/time
clinical librarian, <u>In</u> Libraries without Limits: changing needs-	saved outcomes
changing roles, edited by S Bakker. Kluwer, 1999.	
Glick, J., Sullivan, M. CML in a satellite library. Clinical Librarian	Description of
Quarterly 1984, 3 (1/2), 5-9	setting up a
	CML service
Miller, N., Kaye, D. The experience of a Department of	No health/time
Medicine with a clinical medical library service. Journal of	saved outcomes
Medical Education 1985, 60 (5), 367-73	
Grieves, M. The impact of information use on decision making.	Non systematic

Studies in five sectors – introduction, summary and conclusions.	review.
Library Management 1998, 19 (2), 78-85	Relevant
	primary studies
	already included
Pasquineli, L.M. Report of a survey: impact of a clinical medical	250 word report
librarian on resident education. Journal of Investigative Medicine	– not enough
1998, 46 (1), 2A	detail