

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/52922/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Brown, D. S., Burger, R., Cole, N., Vencatasamy, D., Clare, E. L., Montazam, A. and Symondson, W. O. C. 2014. Dietary competition between the alien Asian Musk Shrew (*Suncus murinus*) and a re-introduced population of Telfair's Skink (*Leiopisma telfairii*). *Molecular Ecology* 23 (15) , pp. 3695-3705. 10.1111/mec.12445

Publishers page: <http://dx.doi.org/10.1111/mec.12445>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Brown DS<sup>1</sup>, Burger R<sup>1</sup>, Cole N<sup>2,3</sup>, Vencatasamy D<sup>3</sup>, Clare EL<sup>4</sup>, Montazam A<sup>5</sup>, Symondson WOC<sup>1</sup>

<sup>5</sup> Genepool, Ashworth Laboratories, King's Buildings, University of Edinburgh, West Mains Road, Edinburgh EH9 3JT, UK

Cardiff School of Biosciences, Sir Martin Evans Building, Cardiff University, Museum Avenue, Cardiff  
CF10 3AX, UK, Fax +44 (0)29 20874116, E-mail [Symondson@cardiff.ac.uk](mailto:Symondson@cardiff.ac.uk)

**Running title:** Niche overlap - alien vs. native predators

## 26 Abstract

27 Reintroduction of rare species to parts of their historical range is becoming increasingly important as  
28 a conservation strategy. Telfair's Skinks (*Leiolopisma telfairii*), once widespread on Mauritius, were  
29 until recently found only on Round Island. There it is vulnerable to stochastic events, including the  
30 introduction of alien predators that may either prey upon it or compete for food resources.  
31 Consequently skinks have been introduced to Ile aux Aigrettes, another small Mauritian island that  
32 has been cleared of rats. However, the island has been invaded by Asian Musk Shrews (*Suncus*  
33 *murinus*), a commensal species spread by man well beyond its natural Asian range. Our aim was to  
34 use next generation sequencing to analyse the diets of the shrews and skinks to look for niche  
35 competition. DNA was extracted from skink faeces and from the stomach contents of shrews.  
36 Application of shrew and skink-specific primers revealed no mutual predation. The DNA was then  
37 amplified using general invertebrate primers with tags to identify individual predators, then  
38 sequenced by 454 pyrosequencing. 119 prey MOTUs (molecular taxonomic units) were isolated,  
39 though none could be identified to species. Seeding of cladograms with known sequences allowed  
40 higher taxonomic assignments in some cases. Although most MOTUs were not shared by shrews and  
41 skinks, Pianka's niche overlap test showed significant prey overlap, suggesting potentially strong  
42 competition where food resources are limited. These results suggest that removal of the shrews from  
43 the island should remain a priority.

44

45

46

47

48

49

## 50 Introduction

51

52 The introduction of locally extinct species to suitable habitats within their wider geographical range is  
53 an increasingly important component of conservation strategies (Seddon *et al.* 2012). When the  
54 distribution of a threatened native species has contracted to one or a few isolated sites it is highly  
55 vulnerable to stochastic events, such as the introduction of alien species, which could rapidly destroy  
56 a last remaining stronghold. Translocation of such a species to a new habitat becomes a conservation  
57 priority. The habitat of such an alternative refuge should ideally be free of threats from alien species,  
58 providing ecological conditions suitable for reintroductions. However, removal of alien species can  
59 often be physically impossible (for example with many invertebrate species) or prohibitively  
60 expensive. In some cases the effective techniques for removal of an alien need to be developed.  
61 Under such conditions it may be necessary to attempt reintroductions under less than ideal  
62 conditions and pragmatically determine whether a rare species can thrive in sympatry with  
63 remaining aliens. Examples of successful translocations are birds such as the Kakapo (*Strigops*  
64 *habroptilus*) between offshore islands in New Zealand (Elliott *et al.* 2001), and both pink pigeon  
65 (*Columba mayeri*) and Mauritius Fody (*Foudia rubra*) to Ile aux Aigrettes (Seymour *et al.* 2005;  
66 Cristinnace *et al.* 2009), and reptiles including whiptail lizards (*Cnemidophorus vanzoi*) to Praslin  
67 Island, Saint Lucia (Dickinson & Fa 2000), Antiguaan racers (*Alsophis antiguae*) to offshore islands of  
68 Antigua (Daltry *et al.* 2001) and lizards to New Zealand islands (Towns & Ferreria 2001).

69

70 Asian Musk Shrews, *Suncus murinus* (Soricidae), are a highly invasive species spread by man to  
71 numerous locations outside its natural Asian range (Ruedi *et al.* 1996). It is a commensal species with  
72 man, often living in and around houses and spread by us between land masses. It was introduced to  
73 Mauritius in the 18<sup>th</sup> century and has been implicated in the loss of endemic vertebrate and  
74 invertebrate species there (Jones 1993; Cole *et al.* 2005; Cheke & Hulme 2008; Solow *et al.* 2008) as  
75 well as in other parts of the world, such as Guam (Fritts & Rodda 1998). Between 2009 and 2010, the

shrew invaded Flat Island to the north of Mauritius, leading to the localised loss of three endemic reptile species within 18 months (N Cole unpublished data). It is thought to have been introduced to Ile aux Aigrettes (southeast of Mauritius) in the early 20<sup>th</sup> century where it spread rapidly (Cheke & Hume 2008). Seymour *et al.* (2005) calculated that 20 females of *S. murinus* on Ile aux Aigrettes could potentially generate a population of 550 individuals over a five month reproductive season. On Ile aux Aigrettes, eradication programmes appeared to be successful for a while, but it soon became clear that some individuals had survived and population recovery was rapid (Varnham *et al.* 2002; Seymour *et al.* 2005; Solow *et al.* 2008). Cats (*Felis catus*) and brown rats (*Rattus rattus*) were successfully eliminated from Ile aux Aigrettes by 1991 as part of a habitat restoration programme (Jones & Hartley 1995), but this may have simply exacerbated the problem with the alien shrews, releasing them from predation and competition with these equally alien predators.

Telfair's Skinks (*Leiolopisma telfairii*) are one of eight species of endemic Mauritian reptiles that managed to survive on Round Island, where they are thriving in the absence of alien predators (North *et al.* 1994; Pernetta *et al.* 2005). Historically these skinks lived on mainland Mauritius and on a number of surrounding islands (Cheke & Hume 2008). As an insurance against loss of the Round Island population, the skinks were introduced to Ile aux Aigrettes between 2006 and 2010 where the adults are surviving well, but there is strong evidence that juveniles may be directly preyed on by Asian Musk Shrews. There is also evidence that adult skinks prey upon shrews and annual population surveys of terrestrial vertebrates along transect lines on Ile aux Aigrettes demonstrated a 68% decline in the relative abundance of shrews since skinks were released (N. Cole unpublished data). However, the skinks and shrews may also be limited by resource competition. Evidence from the eradication programme, based upon live trapping, showed that as numbers of shrews declined, their mean body mass increased considerably. This suggested that food resources were limiting and that this increase in mass was the result of release from intraspecific competition (Seymour *et al.* 2005). It

follows that interspecific competition, between shrews and skinks, might also therefore have an adverse effect upon the skinks if they share the same prey. Both shrews and skinks are omnivorous, eating both plant and animal foods, which may buffer them against food shortages during the dry season on Ile aux Aigrettes, when invertebrate prey are scarce (Cole & Harris 2011). Little is known about the invertebrate prey species consumed by shrews and skinks, although morphological identification of fragments of larger prey in faecal samples has provided some information but mainly at higher taxonomic levels (Vinson & Vinson 1969; Pernetta *et al.* 2005; Richards 2007; Zuël 2009; Copsey *et al.* 2011). These studies using morphological examination of faecal samples from skinks, revealed predation on Araneae, Blattaria, Chilopoda, Coleoptera, Collembola, Decapoda, Dermaptera, Diptera, Embioptera, Hemiptera, Homoptera, Hymenoptera, Isopoda, Lepidoptera, Opisthopora, Orthoptera, Pseudoscorpionida, Scorpionidae, Stylommatophora and Thysanura. Less information appears to exist on invertebrates in the diets of Asian Musk Shrews, which are generally considered to be highly omnivorous, incorporating significant quantities of arthropods in their diets including Orthoptera, Hymenoptera, Blattaria and Chilopoda (Advani & Rana 1981; Prakash & Singh 1999; Lathiya *et al.* 2008). On Ile aux Aigrettes the African land snail *Achatina fulica* was consumed when used as bait in traps (Varnham *et al.* 2002). Given their current wide geographical distribution and adaptability, the shrews are likely to have very different diets within different regions and ecosystems.

The problem with morphological identification of prey remains in the guts or faeces of vertebrates is that it is biased towards prey with hard parts that resist digestion (Symondson 2002). It requires a high level of taxonomic skill and the diagnostic features, essential for species-level identification, may not survive digestive processes (Ingerson-Mahar 2002; Sunderland *et al.* 2005). An alternative approach is to analyse gut and faecal samples using PCR (Symondson 2002; King *et al.* 2008), which can now be combined with next generation sequencing (NGS) (Pompanon *et al.* 2012). General

invertebrate primers can potentially amplify all invertebrates consumed, generating DNA 'barcodes' (diagnostic sequences from a defined region of a gene) for each prey species (Pompanon *et al.* 2012). In tropical ecosystems, such as on Ile aux Aigrettes, the invertebrate fauna has not been barcoded and few, if any, taxa are likely to be found on databases such as GenBank or BOLD (Barcoding of Life Database). However, the sequence output from NGS analyses can be clustered into MOTUs (molecular operational taxonomic units) (Floyd *et al.* 2002) as a proxy for species and can be used to analyse dietary overlap between predator species (Razgour *et al.* 2011). Two predator species may, for example, be consuming the same families of invertebrates but completely different species, and the MOTU approach will reveal this, even when the Linnaean identities of those species cannot be determined. We therefore used next generation sequencing to analyse the invertebrate diets of the shrews and skinks, then tested the hypothesis that there was significant niche overlap between the alien and native species, potentially leading to competition. Tests such as Pianka's niche overlap test (Pianka 1973) do not necessarily reveal where the most significant dietary overlaps lie. We therefore further tested the hypothesis that many prey species are eaten occasionally, probably opportunistically, while a smaller number of key prey species are shared and form a potentially significant part of the diet. Only competition for these prey might be limiting for predator populations. We also tested the hypothesis that shrews and skinks may be competing in a more direct way, by preying on one another.

## Methods

### *Predator sampling*

Samples were collected over an eight week period from the 10<sup>th</sup> March to the 5<sup>th</sup> May 2011, on Ile aux Aigrettes, Mauritius. This 26 ha coralline island nature reserve is leased to, and managed by, the Mauritian Wildlife Foundation. Shrews were initially caught using Sherman traps. However, trapped



shrews had very little material in their guts by the time they were removed. Any remaining gut contents often included bait, and shrews were observed to eat ants from the bait, creating false trophic links. Shrews with full stomachs were subsequently caught more successfully by hand and killed (using UK Home Office approved techniques, Animals (Scientific Procedures) Act 1986) during surveys across the island, both in the early morning and late afternoon/early evening. They were brought back immediately to the field station, dissected under sterile conditions to obtain stomach samples, sexed and measured. Gender was confirmed by post mortem examination for the presence or absence of testes. The length from nose to base of tail was measured to the nearest mm. The presence or absence of fetuses was recorded for females. For males it was often possible to determine adult or juvenile status based on the development of the testes. Females were classed as juveniles if they were less than 12g. The stomach was stored in 94% ethanol at -20°C.

Telfair's Skinks were caught by hand and induced to defecate by gently massaging the belly. A sterile tube was placed below the cloaca to catch the faeces, which was topped up with 94% ethanol and kept at -20°C. Animals were sexed using morphological characteristics including hemipenal eversion of males. Each individual was identified from a unique subcutaneous PIT (Passive Integrated Transponder) tag number, which had been implanted during translocation from Round Island. Finally, measurements of snout-vent length (SVL) were taken. For a full list of both shrews and skinks caught and analysed, with measurements, refer to Table S4.

#### *DNA extraction*

DNA was extracted from faecal and gut samples using the QIAmp DNA Stool Mini Kit (QIAGEN), according to the manufacturer's instructions. Additionally, DNA was extracted from a range of invertebrate samples collected from Ile aux Aigrettes, along with tissue samples from shrews and



skinks, for primer testing, using the DNeasy tissue kit (QIAGEN), according to the manufacturer's instructions.

#### *Primer selection for pyrosequencing*

Published universal PCR primers were tested in a number of different combinations for their ability to amplify DNA from 29 different taxonomic groups of invertebrates (19 orders) collected from Ile aux Aigrettes. Temperature gradient PCRs were performed for each primer pair to determine the optimal annealing temperature at which the most taxa would amplify. PCRs were run on a Peltier Thermal Cycler (Bio-Rad Laboratories, CA, USA) using Multiplex PCR kit (Qiagen) under the following conditions: 1X Master Mix, 0.2  $\mu$ M each primer and 10ng /  $\mu$ L of DNA with an initial denaturation at 95°C for 15 min, 45 cycles of 94°C for 30 s, a gradient of 45–60°C for 90 s and 72°C for 90 s, and a final extension at 72°C for 10 min. DNA of the shrews and skinks were also included so that primer pairs which did not cross-amplify with the predators could be identified. Water was included in each PCR in place of DNA as a negative control. From the large number of primers tested (some unpublished) the best proved to be the forward primer LCO-1490 (Folmer *et al.* 1994) combined with the reverse primer Uni-MiniBar-R (Meusnier *et al.* 2008), which produced a COI (cytochrome oxidase I) amplicon of 177 bp. These primers were found to amplify 28 of the 29 local taxa at an annealing temperature of 49°C and 42 cycles, with no cross-amplification of the predators (Table S1). A second useful primer pair, combining LCO-1490 with ZBJ-ArTr2c (Zeale *et al.* 2011), produce a COI amplicon of 225 bp, and was found to amplify 27 of the 29 taxa at an annealing temperature of 52°C and 40 cycles (Table S1), but in initial tests weakly cross-amplified the shrew. We therefore used the LCO-1490 / Uni-MiniBar-R for further analysis. All other primer combinations tested co-amplified the shrew and/or skink DNA more strongly or amplified a lower range of invertebrate taxa.

#### *Pyrosequencing*

LCO-1490 and Uni-MiniBar-R, modified with fusion primers and MIDS (Multiplex Identifiers in the form of unique DNA tags), were used to amplify faecal/gut DNA extracts from shrews and skinks using PCR conditions described above. By using a unique combination of MIDS on both the forward and reverse primers for each individual predator, MOTUs could be assigned to each predator later bioinformatically. DNA from 41 shrew stomach samples and 29 skink faecal samples were successfully amplified. PCR products were run through a 2% agarose gel stained with ethidium bromide and quantified using UVP VisionWorks® LS Analysis software by comparing fluorescence with known concentrations using MassRuler Low Range DNA ladder (Fermentas). Samples were then pooled together in differing proportions to obtain an approximately equal amount of DNA in the final mixed sample. The pooled sample was purified using the QIAquick PCR Purification Kit (QIAGEN) and pooled DNA concentration quantified by Nanodrop ND-1000 Spectrophotometer.

The DNA was sent to the Genepool, Edinburgh, for NGS. This was performed using the Roche 454 GS-FLX (Roche Applied Sciences) emPCR Lib-L method.

*Sequence Analysis*

Sequences were analysed using the Galaxy platform (<https://main.g2.bx.psu.edu/root>, Giardine *et al.* 2005; Goecks *et al.* 2010; Blankenberg *et al.* 2010) and Bioedit (T. Hall, <http://www.Mbio.ncsu.edu/bioedit/bioedit.html>). Rare haplotypes (represented by <3 copies) were removed, plus sequences much longer or shorter than expected, and then aligned with the remaining haplotypes using clustal W in Bioedit. We then edited the alignment manually to remove indels and match reference sequences.

The sequences were clustered into MOTUs in the program jMOTU (Jones *et al.* 2011) and tested at thresholds from 1-10 bp. A graph of recovered MOTU vs threshold suggests that a 4 bp cut-off was

most appropriate in this data set (see Razgour *et al.* 2011). Representative sequences for each MOTU were compared to the reference database in BOLD ([www. barcodinglife.org](http://www.barcodinglife.org)) recording highest sequence similarity. A phylogenetic tree was constructed of representative MOTUs and a series of known reference sequences using maximum parsimony (MP) in MEGA 5 (Tamura *et al.* 2011) using 1000 bootstrap replications.

### *Ecological Analysis*

Ecological analyses were performed in EcoSim V.7 (<http://grayentsminger.com/ecosim.htm>) and we compared extents of niche overlap using Pianka's (1973) measure of resource sharing (10000 simulated matrices) between shrews and skinks and between males and females in each predator species (equation 3 in Razgour *et al.* 2011). Null models were used to test whether niche overlap was greater than expected by chance. We then re-ran these analyses excluding prey that were only eaten by a single predator. Such occasional prey species are, individually, unlikely to have a significant effect on nutrition and hence on any prey overlap.

Dietary specialization and diversity were estimated using Levins' standardized measure of niche breadth and Shannon's diversity index (equations 1 and 2 in Razgour *et al.* 2011).

### *Prey groups*

Representative sequences from each MOTU were compared to sequences in the BOLD reference database and then included, with known references sequences, in a neighbour-joining reconstruction (Figure 1) in MEGA 5 (Tamura *et al.* 2011). The main prey groups were defined in the cladogram (Figure 1) into Lepidoptera, Dictyoptera, Diptera, Araneae and Gastropoda based on both similarity to known references (category 3 classification, Clare *et al.* in review) and clustering with known references sequences in the cladogram. Individual MOTUs which we found in more than 10% of

either shrews or skinks were also analyzed separately. The effects of predator species (shrew or skink), length, mass, age class (juvenile or adult), sex, and whether gravid, on consumption of prey groups, were explored within a Generalised Linear Model (GLM) (data in Table S4). Length was treated as a covariate and all other predictors as factors. The second order interaction predator:sex was included. A binomial error distribution was used with a logit link function. All analyses were conducted in the R statistical package version 2.9.2.

#### *Species-specific shrew and skink primers*

As the primers used for 454 sequencing did not, in practice, co-amplify either the shrew or skink DNA, species-specific primers were needed in order to determine whether there was intraguild predation between the two predators.

*Cytochrome b* sequences for the skinks (AF280133) and shrews (JF784171), along with sequences for a broad range of vertebrates known to occur on the island (or their close relatives), were acquired from GenBank and aligned in BioEdit in order to design species-specific primers. NetPrimer (Biosoft International) was used to test primer sequences for potential primer-dimer and hairpins which would reduce primer efficiency. LtF1 (5'-CCG TCC CCT ACA TTG GCA CTG-3') and LtR1 (5'-ACA GGA GGT GAA GGA GAG ATA CC-3') were designed to amplify a 140 bp fragment of the skink while SmF1 (5'-TCG GAA TCT GCT TAA TTG CG-3') and SmR1 (5'-AAT AAC GAA TGA GTC AGC CAT AAT T-3') were designed to amplify a 134 bp fragment of the shrew. Gradient PCRs were initially run to determine an optimal annealing temperature for amplification of each target species.

Primers were tested for cross-amplification against DNA extracted from both shrews and skinks, from a range of invertebrate taxa collected on Ile aux Aigrettes and identified to order (n=14) and

274 additionally from invertebrates (n=13) and vertebrates (n=10) collected in the UK (see  
275 Supplementary Table S2).

276  
277 Using the Multiplex PCR Kit (Qiagen) PCR conditions were: 1X Master Mix, 0.5  $\mu$ M each primer, 10%  
278 Q solution and 5ng /  $\mu$ L of DNA with an initial denaturation at 95°C for 15 min, 40 cycles of 94 °C for  
279 30 s, 64.5 °C (for LtF/R) and 64 °C (for SmF/R) for 45 s and 72 °C for 30 s, and a final extension at 72 °C  
280 for 10 min. DNA samples were each tested twice, with water negatives included. Neither primer pair  
281 cross-amplified with any other taxa. Forty eight skink faecal DNA samples were subsequently  
282 screened with LtF/R primers and 49 shrew gut content DNA samples were screened with SmF/R  
283 primers, using the conditions described above.

## 285 Results

### 286 Sequence Analysis

287 Prey DNA was successfully amplified from 42 shrews and 29 skinks, from which 237,402 sequences  
288 were recovered. After removal of rare haplotypes we also removed those that were <100bp and  
289 >220bp and, using the MID codes, the labelled sequences were assigned to individuals (female  
290 shrews n=14, male shrews n=27, one shrew gender unknown, female skinks n=19, male skinks n=10)  
291 and aligned using ClustalW in BioEdit. We edited this alignment to a reference sequence to remove  
292 indels. This combined screening of data yielded 3001 haplotypes. The primer, MID and adapter  
293 sequences were removed for further analysis.

294  
295 The resulting Fasta files in jMOTU (Jones *et al.* 2011) were analysed following the same procedures  
296 employed by Razgour *et al.* (2011) resulting in the recovery of 119 MOTUs, using the 4bp threshold  
297 for assignment.

299 *Ecological analyses.*

300 Of the 119 recovered MOTUs, 53 were found in the diet of skinks and 76 from the diet of shrews with  
301 14 shared between the two predators. Within the 53 MOTUs recovered for skinks, 44 were  
302 consumed by females, 17 by males and 8 were shared (one could not be assigned to an individual as  
303 sequencing did not recover the full MID). Within the 76 MOTUs recovered for shrews, 34 were  
304 consumed by females, 52 by males and 10 were shared.

305

306 Niche overlap was significantly greater than expected by chance between predator species (Pianka's  
307 measure  $O_{jk}=0.55$ ,  $p=0.012$ ), between shrew males and females ( $O_{jk}=0.58$ ,  $p=0.009$ ) and between  
308 skink males and females ( $O_{jk}=0.70$ ,  $p<0.001$ ) (but see Discussion). We then reanalysed the data,  
309 excluding 95 MOTUs that were only recorded from the diets of one animal (rare prey), leaving 24  
310 MOTUs (out of 119 or 20%) that were consumed at least twice. When prey species detected in only  
311 one shrew or skink were excluded (Table S3), prey overlap was shown to be very strong (shrews vs.  
312 skinks  $O_{jk}=0.80$ ,  $p=0.002$ , male vs female shrews  $O_{jk}=0.80$ ,  $p=0.003$ , male vs female skinks  $O_{jk}=0.91$ ,  $p$   
313  $< 0.0001$ ). Overall, the niche breadth of both predator species was narrow (Levins' measure  $B_A=0.18$   
314 for skinks and  $B_A=0.20$  for shrews) but high in diversity ( $H=3.54$  for skinks and  $H=3.74$  for shrews).  
315 Niche breadth and diversity were similar in shrew females ( $B_A=0.26$ ,  $H=3.27$ ) and males ( $B_A=0.30$ ,  
316  $H=3.53$ ). Niche breadth was larger and higher in diversity in skink females ( $B_A=0.30$ ,  $H=3.46$ ) than in  
317 skink males ( $B_A=0.16$ ,  $H=2.69$ ).

318

319 We could not reliably match any sequences to those in BOLD ([www.Barcodinglife.org](http://www.Barcodinglife.org)). A  
320 phylogenetic reconstruction of representative sequences for each MOTU was seeded with reference  
321 sequences (Figure 1) in order to give an indication of taxonomic groups. This showed a large portion  
322 of MOTUs clustering phylogenetically with the reference sequences, suggesting genetic relationships.  
323 Of these, 36 MOTUs were most similar to lepidopteran sequences in BOLD and were phylogenetically

placed in a clade with known lepidopteran sequences. Similarly, 34 MOTUs showed high sequence similarity to representative Dictyoptera in BOLD (termites, cockroaches and mantids), clustered with known Blattaria in the reconstruction, though a few also showed sequence similarity to reference dipteran sequences.

#### *Analysis of consumption of prey groups*

The following analyses were on the putative prey groups as defined in Figure 1. Consumption of Diptera was significantly greater in skinks than in shrews ( $\chi^2 = 11.9$ ,  $df = 1$ ,  $P < 0.001$ ) (Figure 2a), with 41% of skinks found to have consumed Diptera and only 7% of shrews. There was no significant difference in consumption of Gastropoda between shrews and skinks, but male shrews were significantly more likely to consume them than females ( $\chi^2 = 4.3$ ,  $df = 1$ ,  $P = 0.038$ ) (Figure 2b) with 44% of males having consumed them and only 14% of females. Consumption of Dictyoptera by shrews appeared higher than that of skinks but this was not quite significant ( $\chi^2 = 3.3$ ,  $df = 1$ ,  $P = 0.068$ ) (Figure 2c) with 63% of shrews having consumed them and 41% of skinks. Consumption of individual MOTUs, numbers 8, 12 and 13 (all in the Dictyoptera group), were consumed by 20%, 24% and 22% of shrews respectively, but not by any skinks. Conversely, consumption of MOTU number 10 (a dipteran) was found to be significantly higher in skinks than in shrews ( $\chi^2 = 10.1$ ,  $df = 1$ ,  $P = 0.001$ ), with 38% of skinks having consumed them compared to 7% of shrews. Length, age class, mass and whether gravid had no significant effect on consumption of different prey groups.

#### *Species-specific primers*

No evidence was found for intraguild predation between the shrews and skinks; none of the shrew gut samples contained skink DNA and none of the skink faecal samples contained shrew DNA.



## Discussion

Overall our results demonstrate that prey overlap between the shrews and skinks is strong, particularly so when rare prey, consumed only once (80% of prey species detected), were excluded from the analysis. Both analyses may have been affected by sample size (42 shrews and 29 skinks) but the effects are difficult to predict. Larger samples size would increase the probability that less frequently eaten prey will be shared between predator species, but could also increase the number of new rare MOTU's consumed. Rare species (weak links) in food webs may have little influence individually but collectively can increase stability, and this pattern, of many weak links but a few strong links, is commonly found in generalist predator food webs (e.g. McCann *et al.* 1998). All measures of dietary overlap have been criticised (e.g. Wallace 1981) but when the levels of overlap are so strong they are likely to accurately reflect what is happening in the field. We do not know, however, the degree to which the overlap is driven by prey availability or whether at different times of year prey choices by shrews and skinks change. The fact that so many prey were detected only once implies that both shrews and skinks are adaptable and opportunistic, although more prey species may be shown to be shared by the two predators with more sampling. Similarly, species-level analyses of the diets of bats in previous studies showed rare species comprising approx. 50-90% of recovered MOTUs (Clare *et al.* 2009, 2011; Bohmann *et al.* 2011). Strong niche overlap does not necessarily imply significant competition if prey are numerous and not limiting. However, Seymour *et al.* (2005) provided indirect evidence that prey availability can be limiting, by showing that the mean biomass of shrews increased when their numbers were reduced. It is possible, however, that shrew biomass increased for other reasons, such as reduced intensity of social interactions or changes in abiotic conditions. Our field study coincided with when invertebrate resources are considered to be relatively abundant in comparison to other times of year (Cole & Harris 2011).

Although none of the prey could be conclusively identified to a specific taxon, the MOTU approach provided an elegant means of testing for niche overlap between the two predators and between sexes of each predator species, even without access to a reference collection. Data on precisely which prey species are being exploited, particularly those consumed by both shrews and skinks, would require a major barcoding exercise of taxa within the groups indicated on the tree (Figure 1). This would need to be combined with a major effort by museum taxonomists to identify all the taxa morphologically to species. This would not be difficult in, for example, Europe or North America, where the fauna are less diverse and well-studied, but in tropical systems it would present a significant challenge. Only if this were done could the MOTUs found amongst the diets of the shrews and skinks be retrospectively assigned to species. However, analysis of our putative assignments defined in Figure 1 did show some interesting differences. Although Lepidoptera were eaten by both predators, skinks were approximately six times as likely to have consumed Diptera as shrews (Figure 2a). The near significantly greater consumption of Dictyoptera by shrews may relate to Blattaria (Figure 2c), although these have been reported to be eaten by both skinks (Vinson & Vinson 1969; Pernetta *et al.* 2005; Richards 2007; Zuël 2009; Copsey *et al.* 2011) and shrews (Advani & Rana 1981). Dictyoptera are a superorder containing a large range of ecologically very different taxa (termites, cockroaches and mantids), thus possibly masking dietary differences at the group level.

Shrews and skinks clearly have very different physiologies and it might be predicted that the homeothermic shrews would digest their prey more rapidly than poikilotheric skinks. However, we were able to access the shrew samples from an earlier stage of digestion (the stomach) while the skink diet was analysed from fresh faeces. What combined effects these may have had on prey detection, and the relative abundance of different MOTU consumed, could only be established through captive feeding trials.

Some differences were found between sexes, for example female skinks ate a greater diversity of prey species than males, but the reasons for this, though intriguing, are not known. It may be that the dietary needs of reproducing females are different to those of males. Sex differences in diet are often related to sexual dimorphism, for example in birds and mammals (e.g. Rosalino *et al.* 2009; Phillips *et al.* 2011) and arthropods (e.g. Symondson & Liddell 1993; Pekár *et al.* 2011), where the size difference allows predators to access different prey, allowing intersexual partitioning of resources. Adult male skinks and shrews are larger than females. Male shrews were more than three times as likely to have eaten gastropods than females (Figure 2b). However, all of these results would have been affected by the differences in sample sizes and they would require further work to verify.

Analysis with species-specific primers provided no evidence of direct intraguild predation by shrews on skinks or skinks on shrews. However, this contrasts with observations on the island of juvenile skink remains in the guts of shrews and shrew remains in the faeces of skinks (pelts and hair), plus direct observations of mutual predation (N. Cole and D. Vencatasamy pers. obs.). Unavoidable delays in conducting our work meant that shrews and skinks were sampled well after the peak period when skinks hatch and are at their most vulnerable. The release of Telfair's skinks onto Ile aux Aigrettes coincided with substantial declines in the abundance of shrews, possibly as a result of skinks preying on shrews. However, at the current low shrew density dietary evidence of predation may not be detected unless the number of skinks sampled was greatly increased. If prey are limiting then high prey overlap between shrews and skinks may also have played a role in the decline of the shrews.

Any form of analysis of predation, whether morphological or utilising PCR, must always be qualified by the fact that we cannot distinguish between predation, scavenging and secondary predation. Scavenging of dead material by insect predators has been shown to be a likely source of error using PCR (Foltan *et al.* 2005; Juen & Traugott 2005). Within invertebrate food webs, secondary predation,

where one predator eats another and the prey in the guts of the consumed predator can be detected, is probably a less important source of error (Sheppard *et al.* 2005). In all cases (predation, scavenging, secondary predation) the prey detected are contributing to the nutrition of the predator but the dynamics of the interactions are clearly very different.

The novel combination of existing primers proved to be highly effective at amplifying invertebrate DNA, covering a broad range of invertebrates but with no co-amplification of the predators. They proved to be a significant improvement on the Uni-MiniBar primers of Meusnier *et al.* (2008), UniMinibarF1 / UniMinibarR1, which have been criticised for their low taxonomic coverage (Ficetola *et al.* 2010). However, when is UniMinibarR1 combined with the general invertebrate forward primer LCO-1490 of Folmer *et al.* (1994) specificity and coverage were excellent.

As far as we are aware, this is only the second time that PCR has been used to analyse reptile diets from faecal samples, the first being our previous study of predation on earthworms by slow worms, the legless lizards *Anguis fragilis* (Brown *et al.* 2008). In that instance the primers used for NGS were the earthworm group-specific primers developed by Harper *et al.* (2005). A further paper on the diets of snakes in this special issue reports the vertebrate and invertebrate diet of the smooth snake *Coronella austriaca*, analysed using prey-specific primers (Brown *et al.* submitted). The fact that PCR and NGS can be used to analyse the diets of reptiles from faeces, despite the fact that many species digest their prey to the extent of dissolving bones (Secor 2008), opens up a potentially rich field for future research on reptile trophic ecology. A different molecular approach was taken recently by Goiran *et al.* (2013), who demonstrated that fish eggs palpated from the stomachs of sea snakes could be identified by sequencing their DNA.

Concerted trapping in 1999 to eradicate the shrews from Ile aux Aigrettes was only partially effective. Some individuals are 'trap-shy' and can go on to generate a resurgent population within a short time. It appears to be the case that shrews enter traps through curiosity, rather than responding to baits (which are often left untouched) (Varnham *et al.* 2002; Seymour *et al.* 2005). Thus analysis of their diets in the field provided an opportunity to identify favoured prey that, as bait or food odours, could improve trap efficiency. The results of our analysis suggest that Lepidoptera larvae or cockroaches may provide effective bait. Cockroach frass from laboratory cultures is highly pungent and may be sufficient to attract shrews.

The ethics of killing vertebrates in order to obtain gut samples must be properly justified. Here we caught and killed shrews in the field (using UK Home Office approved techniques), to obtain gut samples. Once caught it was not considered ethically acceptable to release these pests back to the wild, where they would continue to pose a threat to native wildlife. This allowed us to maximise the information obtainable from these animals by analysing their stomach contents (rather than faeces) where prey DNA was likely to be less degraded. A key aim of Mauritian conservationists has been to eradicate shrews from offshore islands to permit further restoration processes. However, to date, eradication attempts have only been successful using traps on topographically simple islands of a few hectares or less (Varnham *et al.* 2002). The problem with the shrews is that traps do not catch them efficiently and suitable poison baits have not been devised (Varnham *et al.* 2002; Seymour *et al.* 2005).

Our conclusion, therefore, is that shrews and skinks are feeding to a large extent on the same species of invertebrate prey, potentially leading to competition. If so then shrew control is likely to be beneficial to the fitness of the skinks. Mutual predation is known to occur, but our analysis failed to find evidence of this outside the period when juvenile skinks are particularly vulnerable. This is

probably because skinks grow too large to be attacked by shrews and similarly, at low densities, shrews increase in biomass (Seymour *et al.* 2005) and may be too large for predation by skinks. Given that the shrews pose a threat to island biodiversity, development of methods to eradicate them from islands such as Ile aux Aigrettes should continue to be a priority.

## Acknowledgements

We thank the Durrell Wildlife Conservation Trust and Cardiff School of Biosciences for funding this work. This research would not have been possible without the full support and assistance of the Mauritian Wildlife Foundation and we are particularly grateful to Zayd Jhumka and Rouben Mootoocurpen who assisted in the collection of skinks, shrews and invertebrates on Ile aux Aigrettes. We thank the National Parks and Conservation Service, Ministry of the Agro-Industry, Mauritius, for permission to conduct this research.

## References

- Advani R, Rana BD (1981) Food of the House Shrew, *Suncus murinus sindensis*, in the Indian desert. *Acta Theriologica*, **26**, 133— 134.
- Blankenberg D, Von Kuster G, Coraor N, Ananda G, Lazarus R, Mangan M, Nekrutenko A, Taylor J. (2010) Galaxy: a web-based genome analysis tool for experimentalists. *Current Protocols in Molecular Biology*, **Unit 19.10.1-21**, DOI: 10.1002/0471142727.mb1910s89, Wiley.
- Bohmann K, Monadjem A, Lehmkuhl Noer C *et al.* (2011) Molecular diet analysis of two African free-tailed bats (Molossidae) using high throughput sequencing. *PLoS One*, **6**, e21441.

- 495 Brown DS, Ebenezer KL, Symondson WOC (submitted for Special Issue) Molecular analysis of the  
 496 diets of snakes: changes in prey selection during development of the rare smooth snake  
 497 *Coronella austriaca*. *Molecular Ecology*.
- 498 Brown DS, Jarman SN, Symondson WOC (2012) Pyrosequencing of prey DNA in reptile faeces:  
 499 analysis of earthworm consumption by slow worms. *Molecular Ecology Resources*, **12**, 259-  
 500 266.
- 501 Cheke AS, Hume JP (2008) *Lost land of the Dodo: an ecological history of Mauritius, Réunion &*  
 502 *Rodrigues*. A & C Black, London.
- 503 Clare EL, Barber BR, Sweeney BW, Hebert PDN, Fenton MB (2011) Eating local: influences of habitat  
 504 on the diet of little brown bats (*Myotis lucifugus*). *Molecular Ecology*, **20**, 1772–1780.
- 505 Clare EL, Fraser EE, Braid HE, Fenton MB, Hebert PDN (2009) Species on the menu of a generalist  
 506 predator, the eastern red bat (*Lasiurus borealis*): using a molecular approach to detect  
 507 arthropod prey. *Molecular Ecology*, **18**, 2532–2542.
- 508 Cole NC, Harris S (2011) Environmentally-induced shifts in behavior intensify indirect competition by  
 509 an invasive gecko in Mauritius. *Biological Invasions*, **13**, 2063-2075.
- 510 Cole N, Jones CG, & Harris S. (2005) The need for enemy-free space: The impact of an invasive gecko  
 511 on island endemics. *Biological Conservation*, **125**, 467-474.
- 512 Copsey JA, Shelbourne G, Grice R, Goder M, Buckland S, Jhumka Z, Nundlall V, Jones C, Cole N (2011)  
 513 Possible control of introduced giant African land snails (*Achatina* spp.) by the reintroduced  
 514 endemic skink *Leiolopisma telfairii*, Ile aux Aigrettes, Mauritius. *Management of Biological*  
 515 *Invasions*, **2**, 39-45.
- 516 Cristinnace A, Handschuh M, Switzer RA, Cole RE, Tatayah V, Jones CG, Bell D (2009) The release and  
 517 establishment of Mauritius Fodies *Foudia rubra* on Ile aux Aigrettes, Mauritius. *Conservation*  
 518 *Evidence*, **6**, 1-5.



- 519 Daltry JC, Bloxam Q, Cooper G, Day ML, Hartley J, Henry M, Lindsay K, Smith BE (2001) Five years of  
520 conserving the 'world's rarest snake', the Antigua racer *Alsophis antiguae*. *Oryx*, **35**, 119-  
521 127.
- 522 Dickinson HC, Fa JE (2000) Abundance, demographics and body condition of a translocated  
523 population of St Lucia whiptail lizards (*Cnemidophorus vanzoi*). *Journal of Zoology*, **251**, 187-  
524 197.
- 525 Elliott GP, Merton DV, Jansen PW (2001) Intensive management of a critically endangered species:  
526 the kakapo. *Biological Conservation*, **99**, 121-133.
- 527 Ficetola F, Coissac E, Zundel S et al. (2010) In silico comparison of primers for DNA barcoding. *BMC*  
528 *genomics*, **11**, e434.
- 529 Floyd R, Abebe E, Papert A, Blaxter M (2002) Molecular barcodes for soil nematode identification.  
530 *Molecular Ecology*, **11**, 839-850.
- 531 Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for the amplification of  
532 mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates.  
533 *Molecular Marine Biology and Biotechnology*, **3**, 294-299.
- 534 Foltan P, Sheppard SK, Konvicka M, Symondson WOC (2005) The significance of facultative  
535 scavenging in generalist predator nutrition: detecting decayed prey in the guts of predators  
536 using PCR. *Molecular Ecology*, **14**, 4147-4158.
- 537 Fritts TH, Rodda GH (1998) The role of introduced species in the degradation of island ecosystems: a  
538 case history of Guam. *Annual Review of Ecology and Systematics*, **29**, 113-140.
- 539 Giardine B, Riemer C, Hardison RC, Burhans R, Elnitski L, Shah P, Zhang Y, Blankenberg D, Albert I,  
540 Taylor J, Miller W, Kent WJ, Nekrutenko A (2005) Galaxy: a platform for interactive large-  
541 scale genome analysis. *Genome Research*, **15**, 1451-1455.

- 542 Goecks, J, Nekrutenko, A, Taylor, J and The Galaxy Team (2010) Galaxy: a comprehensive approach  
543 for supporting accessible, reproducible, and transparent computational research in the life  
544 sciences. *Genome Biology*, **11**, R86.
- 545 Goiran C, Dubey S, Shine R (2013) Effects of season, sex and body size on the feeding ecology of  
546 turtle-headed sea snakes (*Emydocephalus annulatus*) on IndoPacific inshore coral reefs. *Coral*  
547 *Reefs*, **32**, 527-538.
- 548 Harper GL, King RA, Dodd CS, Harwood JD, Glen DM, Bruford MW, Symondson WOC (2005) Rapid  
549 screening of invertebrate predators for multiple prey DNA targets. *Molecular Ecology*, **14**,  
550 819-827.
- 551 Ingerson-Mahar J (2002) Relation diet and morphology in adult carabid beetles. In: *The Agroecology*  
552 *of Carabid Beetles* (ed. Holland JM), pp 111-136. Intercept, Andover, UK.
- 553 Jones CG (1993) The ecology and conservation of Mauritian skinks. *Proceedings of The Royal Society*  
554 *of Arts and Sciences of Mauritius*, **5**, 71-95.
- 555 Jones CG, Hartley J (1995) A conservation project on Mauritius and Rodrigues: An overview and  
556 bibliography. *The Dodo Journal of the Jersey Wildlife Preservation Trust*, **31**, 40-65.
- 557 Jones M, Ghoorah A, Blaxter M (2011) jMOTU and taxonator: turning DNABarcode sequences into  
558 annotated operational taxonomic units. *PLoS One*, **6**, e19259.
- 559 Juen A, Traugott M (2005) Detecting predation and scavenging by DNA gut-content analysis: a case  
560 study using a soil insect predator-prey system. *Oecologia*, **142**, 344-352.
- 561 King RA, Read DS, Traugott M, Symondson WOC (2008) Molecular analysis of predation: a review of  
562 best practice for DNA-based approaches. *Molecular Ecology*, **17**, 947-963.
- 563 Lathiya SB, Ahmed SM, Pervez A, Rizvi SWA (2008) Food habits of rodents in grain godowns of  
564 Karachi, Pakistan. *Journal of Stored Products Research*, **44**, 41-46.
- 565 McCann K, Hastings A, Huxel GR (2008) Weak trophic interactions and the balance of nature. *Nature*,  
566 **395**, 794-798.

- Meusnier I, Singer GAC, Landry JF, Hickey DA, Hebert PDN, Hajibabaei M (2008) A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*, **9**, 214.
- North SG, Bullock DJ, Dulloo ME (1994) Changes in the vegetation and reptile populations on Round Island, Mauritius, following eradication of rabbits. *Biological Conservation*, **67**, 21–28.
- Pekár S, Martišová M, Bilde T (2011) Intersexual trophic niche partitioning in an ant-eating spider (Araneae: Zodariidae). *Plos One*, **6**, e14603.
- Pernetta AP, Bell DJ, Jones CG (2005) Macro- and microhabitat use of Telfair's skink (*Leiolopisma telfairii*) on Round Island, Mauritius: implications for their translocation. *Acta Ecologica*, **28**, 313-323.
- Phillips RA, McGill RAR, Dawson DA, Bearhop S (2011) Sexual segregation in distribution, diet and trophic level of seabirds: insights from stable isotope analysis. *Marine Biology*, **158**, 2199-2208.
- Pianka ER (1973) The structure of lizard communities. *Annual Review of Ecology and Systematics*, **4**, 53-74.
- Pompanon F, Deagle BE, Symondson WOC, Brown DS, Jarman SD, Taberlet P (2012) Who is eating what: diet assessment using next generation sequencing. *Molecular Ecology*, **21**, 1931-1950.
- Prakash I, Singh H (1999) Food of the shrew, *Suncus murinus* inhabiting hill tracks of south and southeastern Rajasthan. *Proceeding of the National Academy of Sciences India*, **69**, 245-250.
- Ruedi M, Courvoisier C, Vogel P, Catzeflis FM, (1996) Genetic differentiation and zoogeography of the Asian house shrew *Suncus murinus* (Mammalia: Soricidae). *Biological Journal of the Linnean Society*, **57**, 307–316.
- Razgour O, Clare EL, Zeale MRK, Hanmer J, Schnell IB, Rasmussen M, Gilbert TP, Jones G (2011) High-throughput sequencing offers insight into mechanisms of resource partitioning in cryptic bat species. *Ecology and Evolution*, **1**, 556-570.

- 591 Richards H (2007) *An investigation into the macro- and microhabitat use and dietary preference of*  
 592 *the translocated population of Telfair's skink (Leiolopisma telfairii) on Ile aux Aigrettes,*  
 593 *Mauritius*. MSc thesis, Department of Biology, University of East Anglia, UK.
- 594 Rosalino LM, Santos MJ, Pereira I, Santos-Reis M (2009) Sex-driven differences in Egyptian  
 595 mongoose's (*Herpestes ichneumon*) diet in its northwestern European range. *European Journal*  
 596 *of Wildlife Research*, **55**, 293-299.
- 597 Secor SM (2008) Digestive physiology of the Burmese python: broad regulation of integrated  
 598 performance. *Journal of Experimental Biology*, **211**, 3767-3774.
- 599 Seddon PJ, Strauss WM, Innes J (2012) Animal translocations: what are they and why do we do  
 600 them? In: *Reintroduction Biology: Integrating Science and Management* (eds Ewen JG,  
 601 Armstrong DP, Parker KA, Seddon PJ), pp. 1-32. Wiley-Blackwell, Oxford.
- 602 Seymour A, Varnham K, Roy S, Harris S, Bhageerutty L, Church S, Harris A, Jennings NV, Jones C,  
 603 Khadun A, Mauremootoo J, Newman T, Tatayah V, Webbon C, Wilson G (2005) Mechanisms  
 604 underlying the failure of an attempt to eradicate the invasive Asian musk shrew *Suncus*  
 605 *murinus* from an island nature reserve. *Biological Conservation*, **125**, 23-35.
- 606 Sheppard SK, Bell JR, Sunderland KD, Fenlon J, Skirvin DJ, Symondson WOC (2005) Detection of  
 607 secondary predation by PCR analyses of the gut contents of invertebrate generalist  
 608 predators. *Molecular Ecology*, **14**, 4461-4468.
- 609 Solow A, Seymour A, Beet A, Harris S (2008) The untamed shrew: on the termination of an  
 610 eradication programme for an introduced species. *Journal of Applied Ecology*, **45**, 424-427.
- 611 Sunderland KD, Powell W, Symondson WOC (2005) Populations and communities. In: *Insects as*  
 612 *Natural Enemies: A Practical Perspective* (ed. Jervis MA), pp. 299-434. Springer, Berlin.
- 613 Symondson, W.O.C. (2002). Molecular identification of prey in predator diets. *Molecular Ecology*, **11**,  
 614 627-641.

- 615 Symondson WOC, Liddell JE (1993) The detection of predation by *Abax parallelepipedus* and  
616 *Pterostichus madidus* (Coleoptera: Carabidae) on Mollusca using a quantitative ELISA.  
617 *Bulletin of Entomological Research*, **83**, 641-647.
- 618 Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular Evolutionary  
619 Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum  
620 Parsimony Methods. *Molecular Biology and Evolution*, **28**, 2731-2739.
- 621 Towns DR, Ferreira SM (2001) Conservation of New Zealand lizards (Lacertilia: Scincidae) by  
622 translocation of small populations. *Biological Conservation*, **98**, 211–222.
- 623 Varnham KJ, Roy SS, Seymour A, Mauremootoo JR, Jones CG, Harris S (2002) Eradicating Indian musk  
624 shrews (*Suncus murinus*, *Soricidae*) from Mauritian offshore islands. In: *Turning the tide: the*  
625 *eradication of invasive species* (eds Veitch CR, Clout MN), pp. 342-349. Invasive Species  
626 Specialist Group, Species Survival Commission, World Conservation Union, Gland,  
627 Switzerland.
- 628 Vinson J, Vinson JM (1969) The saurian fauna of the Mascarene Islands. *The Mauritius Institute*  
629 *Bulletin*, **VI**, 203-320.
- 630 Wallace RK (1981) An assessment of diet overlap indexes. *Transactions of the American Fisheries*  
631 *Society*, **110**, 72-76.
- 632 Wathne JA, Haug T, Lydersen C (2000) Prey preference and niche overlap in ringed seals *Phoca*  
633 *hispida* and harp seals *P. groenlandica* in the Barents Sea. *Marine Ecology Progress Series*,  
634 **194**, 233-239.
- 635 Zeale MRK, Butlin RK, Barker GLA, Lees DC, Jones G (2011) Taxon-specific PCR for DNA barcoding  
636 arthropod prey in bat faeces. *Molecular Ecology Resources*, **11**, 236-244.
- 637 Zuël N (2009) *Ecology and conservation of an endangered reptile community on Round Island,*  
638 *Mauritius*. PhD thesis, Mathematisch-naturwissenschaftliche Fakultät, Universität Zürich.
- 639

640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664

**Data accessibility**

All sequences arising from NGS, fully processed, collapsed and aligned, plus allocated to individual predators and ready for analysis, will be included as Online Supplementary Material after acceptance of the paper. Three files will be included: all ‘sequences pooled.fasta’, ‘all sequences shrew.fas’ and ‘all sequences skink.fas’.

**Author Contributions Box**

Gut and faecal samples were collected from Mauritius by RB and DV, and DNA extracted by RB, who designed and applied species-specific primers for analysing mutual predation between shrews and skinks. Preparation of samples for NGS was conducted by DSB, along with analyses of predation on key prey taxa. Bioinformatics and ecological analyses were conducted by ELC. Supervision of the fieldwork in Mauritius was conducted by NC, who provided the expertise on Mauritian ecology. AM conducted the 454 analysis. Overall supervision of the project and the writing of the paper were primarily conducted by WOCS, with major contributions from co-authors.

**Figure legends****Figure 1**


Cladogram showing reconstructed relationships between all MOTUs retrieved from the guts or faecal samples of Asian Musk Shrew and Telfair's Skinks, colour codes to denote prey consumed by shrews, skinks or by both species.


**Figure 2**

Main significant or near significant differences in diet arising from analysis of putative higher-order classifications, as defined in Figure 1. **a.** Predicted probability of consumption of Diptera ( $\pm$  s.e.) by shrews and skinks, showing significantly higher consumption in skinks ( $p < 0.001$ ). **b.** Predicted probability of consumption of Gastropoda ( $\pm$  s.e.) by shrews, showing significantly higher consumption in males than in females ( $p = 0.038$ ). **c.** Predicted probability of consumption of Dictyoptera ( $\pm$  s.e.) by shrews and skinks, showing a trend towards higher consumption by shrews ( $p = 0.068$ ).




- Shared Prey
- Skink Prey
- Shrew Prey






2a



2b



2c

# 1 SUPPLEMENTARY MATERIAL

## 2 Table S1

3 Invertebrates collected from Ile aux Aigrettes and tested for PCR amplification with the two primers  
4 sets developed for 454 pyrosequencing, LCO-1490 / Uni-MiniBar-R and LCO-1490 / ZBJ-ArtR2c.

5

Potential prey	LCO-1490 / Uni-MiniBar-R	LCO-1490 / ZBJ-ArtR2c
Coleoptera 1	✓	✓
Oligochaeta	✓	✓
Hemiptera 1	✓	✓
Isopoda	✓	✓
Dermaptera	✓	✓
Embioptera	✓	✓
Diplopoda	✓	✓
Hymenoptera ( <i>Vespa sp.</i> )	✓	✓
Araneae 1	✓	✓
Gastropoda 1	✓	✓
Lepidoptera 1	✓	✓
Diptera	✓	✓
Blattaria 1	✓	✓
Odonata	✓	✓
Decapoda	✓	✓
Gastropoda 2	✓	✓

Hymenoptera - Formicoidea	✓	✓
Lepidoptera 2	✓	✓
Scorpiones	✓	✓
Araneae 2	✓	✓
Coleoptera - Cerambycidae	✓	✓
Diptera - Culicidae	✓	✓
Collembola	✓	✓
Orthoptera - Gryllidae		✓
Hemiptera 2	✓	✓
Hemiptera 3	✓	
Chilopoda	✓	✓
Coleoptera 2	✓	
Blattaria 2	✓	✓
<b>Total</b>	<b>28/29</b>	<b>27/29</b>

6

7

8

9

10

11

12

13

14

15

16 Table S2.

17 Non-target species tested for cross-amplification with skink-specific (LtF/R) and shrew-specific

18 (SmF/R) PCR primers. Neither primer set co-amplified any of these taxa.

19

Order	Species	Origin
Coleoptera	spp. x2	Ile aux Aigrettes
Lepidoptera	spp. x2	Ile aux Aigrettes
Blattaria	spp. x2	Ile aux Aigrettes
Hymenoptera	spp. x2	Ile aux Aigrettes
Diptera	spp. x2	Ile aux Aigrettes
Isopoda	spp. x2	Ile aux Aigrettes
Aranaea	spp. x2	Ile aux Aigrettes
Pulmonata	<i>Arion intermedius</i>	UK
	<i>A. distinctus</i>	UK
	<i>Limax flavus</i>	UK
Haplotaxida	<i>Lumbricus terrestris</i>	UK
	<i>L. rubellus</i>	UK
	<i>Aporrectodea caliginosa</i>	UK
	<i>A. longa</i>	UK
Coleoptera	<i>Notiophilus biguttaus</i>	UK
	<i>Adalia bipunctata</i>	UK
	<i>Tachyporus obtusus</i>	UK
Diptera	<i>Tipulidae sp.</i>	UK

Dermaptera	<i>Forficula sp.</i>	UK
Aranaea	<i>Erigone ddentipalpis</i>	UK
Squamata	<i>Zootoca vivipara</i>	UK
	<i>Anguis fragilis</i>	UK
	<i>Coronella austriaca</i>	UK
	<i>Natrix natrix</i>	UK
Rodentia	<i>Myodes glareolus</i>	UK
	<i>Mus musculus</i>	UK
	<i>Apodemus flavicollis</i>	UK
Soricomorpha	<i>Neomys fodiens</i>	UK
	<i>Sorex araneus</i>	UK
Caudata	<i>Lissotriton helveticus</i>	UK

20

21

22

23

24

25

26

27

28

29

30

31 Table S3

32 Numbers of shrews and skinks, of each sex, that contained each prey MOTU, excluding MOTUs that  
 33 were only found in one animal overall. Shrew N/R is an animal not sexed (see text). 'Total detections'  
 34 are the numbers of shrews+skinks testing positive for that MOTU. For a complete list, and to find  
 35 MOTU numbers, see Figure 1.

36

37

MOTU no.	Skinks		Shrews		Shrew N/R	Total
	male	female	male	female		detections
2	3	7	10	6	1	27
3	1	1	0	2	0	4
4	3	3	6	5	1	18
5	1	6	10	5	0	22
6	3	5	5	0	0	13
7	1	5	10	3	1	20
8	0	0	3	5	1	9
10	3	8	3	0	0	14
11	0	0	3	0	0	3
12	0	0	7	3	1	11
13	0	0	6	3	1	10
16	0	0	2	0	0	2
20	0	0	3	0	0	3
21	0	0	4	1	0	5



28	0	0	3	0	0	3
30	0	0	1	1	0	2
31	0	0	3	0	0	3
34	0	1	3	2	0	6
39	1	2	2	2	0	7
44	1	3	0	1	0	5
49	0	1	0	1	0	2
59	0	1	1	0	0	2
71	1	1	0	0	0	2
116	0	2	0	0	0	2

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52 Table S4

53 File 'MOTUs consumed by shrews and skinks.xls'. Spreadsheet providing raw data on the shrews and  
54 skinks from which we successfully amplified invertebrate DNA, including sex, mass, length,  
55 adult/juvenile status and whether gravid.

56

57 Tables S5-S6

58 Spreadsheets including representative sequences for all haplotypes arising from NGS, fully  
59 processed, collapsed and aligned, allocated to individual predators and ready for analysis. Divided  
60 into 'All sequences shrew.fas' and 'All sequences skink.fas'.

MOTU	Skink		Shrew		
	Males	Females	Males	Females	Unknown
2	11, 31, 46	20, 33, 36, 41, 45, 7, 9	11, 13, 21, 26, 33, 36, 44, 6, 8, 9	1, 25, 28, 29, 7, 37	2
3	42	20		7, 32	
4	10, 31, 42	3, 7, 9	12, 22, 26, 40, 41, 44	19, 20, 38, 39, 32	2
5	11	20, 29, 2, 36, 41, 44	13, 15, 21, 26, 33, 35, 44, 6, 8, 9	25, 29, 38, 7, 50	
6	11, 15, 42	12, 18, 20, 36, 41	12, 21, 22, 26, 33		
7	39	18, 20, 33, 41, 48	14, 17, 22, 30, 34, 3, 41, 43, 44, 9	45, 46, 49	2
8			21, 28, 8	1, 25, 7, 32, 37	2
10	11, 43, 6	14, 2, 33, 3, 40, 41, 44, 45	15, 40, 44		
11			41, 44, 9		
12			15, 16, 17, 26, 33, 41, 48	24, 25, 49	2
13			21, 33, 36, 44, 8, 9	25, 28, 37	2
16			40, 41		
20			26, 33, 48		
21			22, 26, 48, 46	49	
28			26, 33, 48		
30			41	29	
31			26, 33, 48		
34		9	40, 44, 9	37, 50	
39	42	4, 9	40, 44	4, 33	
44	37	33, 41, 9		49	
49		18		24	
59		7	6		
71	11	41			
116		12, 41			