

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/53925/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Barlow, M. J., Swinyard, B. M., Owen, P. J., Cernicharo, J., Gomez, Haley Louise, Ivison, R. J., Krause, O., Lim, T. L., Matsuura, Mikako, Miller, S., Olofsson, G. and Polehampton, E. T. 2013. Detection of a noble gas molecular ion,  $36\text{ArH}^+$ , in the Crab Nebula. *Science* 342 (6164), pp. 1343-1345. 10.1126/science.1243582

Publishers page: <http://dx.doi.org/10.1126/science.1243582>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# Detection of a Noble Gas Molecular Ion, $^{36}\text{ArH}^+$ , in the Crab Nebula

M. J. Barlow<sup>1†</sup>, B. M. Swinyard<sup>1,2</sup>, P. J. Owen<sup>1</sup>, J. Cernicharo<sup>3</sup>,  
H. L. Gomez<sup>4</sup>, R. J. Ivison<sup>5</sup>, O. Krause<sup>6</sup>, T. L. Lim<sup>2</sup>,  
M. Matsuura<sup>1</sup>, S. Miller<sup>1</sup>, G. Olofsson<sup>7</sup>, E. T. Polehampton<sup>2,8</sup>

<sup>1</sup>Dept. of Physics & Astronomy, University College London, Gower Street, London WC1E 6BT, UK

<sup>2</sup>Space Science & Technology Department, Rutherford Appleton Laboratory, Didcot OX11 0QX, UK

<sup>3</sup>Laboratory of Molecular Astrophysics, Dept. of Astrophysics, CAB, INTA-CSIC,  
Ctra de Ajalvir, km 4, 28850 Torrejón de Ardoz, Madrid, Spain

<sup>4</sup>School of Physics & Astronomy, Cardiff University, The Parade, Cardiff CF24 3AA, UK

<sup>5</sup>UK Astronomy Technology Centre, Royal Observatory Edinburgh,  
Blackford Hill, Edinburgh EH9 3HJ, UK

<sup>6</sup>Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

<sup>7</sup>Dept. of Astronomy, Stockholm University, AlbaNova University Center,  
Roslagstullsbacken 21, 10691 Stockholm, Sweden

<sup>8</sup>Institute for Space Imaging Science, Dept. of Physics & Astronomy,  
University of Lethbridge, Lethbridge, Alberta T1K 3M4, Canada

<sup>†</sup>email:mjb@star.ucl.ac.uk; Published in *Science*, v.342, pp.1343-1345, 2013

**Noble gas molecules have not hitherto been detected in space. From spectra obtained with the *Herschel Space Observatory*, we report the detection of emission in the 617.5 GHz and 1234.6 GHz  $J = 1-0$  and  $2-1$  rotational lines of  $^{36}\text{ArH}^+$  at several positions in the Crab Nebula, a supernova remnant known to contain both  $\text{H}_2$  molecules and regions of enhanced ionized argon emission.  $^{36}\text{Ar}$  is believed to have originated from explosive nucleosynthesis in massive**

**stars during core-collapse supernova events. Its detection in the Crab Nebula, the product of such a supernova event, confirms this expectation. The likely excitation mechanism for the observed  $^{36}\text{ArH}^+$  emission lines is electron collisions in partially ionized regions with electron densities of a few hundred per centimeter cubed.**

Noble gas compounds have not yet been found in space, despite some examples, such as ionized hydrides, being relatively stable (1). Astronomical searches for the near-infrared and far-infrared lines of  $\text{HeH}^+$ , whose dissociation energy is 1.8 eV (2) have not been successful (3, 4). The Crab Nebula is the product of the supernova of 1054 AD and is thought to have originated from the core-collapse explosion of a star 8-16 times as massive as the Sun (5). We have obtained far-infrared to submillimeter spectra of the Crab Nebula using the *Herschel Space Observatory* (6). We report here the detection of emission lines from the ionized hydride of argon, an element predicted to form by explosive nucleosynthesis in core-collapse supernovae (7).

The Crab Nebula was observed with the Fourier Transform Spectrometer (FTS) of the Spectral and Photometric Imaging Receiver (SPIRE) (8, 9) on Operational Day 466 of the *Herschel* mission, as part of the Mass-loss of Evolved StarS (MESS) Guaranteed Time Key Project (10). The 19 SPIRE Long Wavelength (SLW) detectors, each with a beamwidth of  $\sim 37''$ , covered the 447-989 GHz frequency range (303-671  $\mu\text{m}$ ), while 35 SPIRE Short Wavelength (SSW) detectors, each with a beamwidth of  $\sim 18''$ , covered the 959-1544 GHz frequency range (194-313  $\mu\text{m}$ ) (Fig. 1). The full width half maximum spectral resolution was 1.44 GHz at all frequencies, corresponding to a resolving power of 690 in the middle of the frequency range. The observation consisted of 48 FTS scans, for a total on-source exposure time of 3197s. The data were processed using the extended source calibration in version 11 of the Herschel Interactive Processing Environment (11). The  $J = 2-1$ ,  $F = 5/2-3/2$  line of  $\text{OH}^+$  at 971.8038 GHz (12),

which falls in the SLW and SSW spectral overlap region, is present in emission in many of the spectra (Fig. 2). This line has been observed from a range of astrophysical environments by *Herschel*, both in absorption (13) and emission (14, 15). In addition, two unidentified emission lines were found to be present in some of the Crab spectra, one in the SLW range at  $\sim 618$  GHz, and the other in the SSW range at  $\sim 1235$  GHz.

The knots and filaments of the Crab Nebula are known to exhibit expansion velocities ranging between  $700\text{--}1800 \text{ km s}^{-1}$  (16); in different detectors we measured radial velocities for the  $\text{OH}^+$  971.8038 GHz line that ranged between  $-603$  and  $+1037 \text{ km s}^{-1}$ . Several spectra showed multiple  $\text{OH}^+$  velocity components, some blended, but in most spectra the  $\text{OH}^+$  velocity components were unresolved, exhibiting very different radial velocities from detector to detector, consistent with an origin from differing knots or filaments in the nebula, each with its own discrete velocity. Because  $\text{OH}^+$  was the only identified species in the spectra initially, we used the measured radial velocities of the 971.8038 GHz  $\text{OH}^+$  line, whose centroid frequency could typically be measured to an accuracy of  $\pm(25\text{--}40) \text{ km s}^{-1}$ , as a reference to correct to a ‘rest’ frequency the observed frequencies of the 618 or 1235 GHz line falling in the same spectrum. There were four SLW spectra, those from detectors B3, C3, D3 and D4, in which the  $\text{OH}^+$  line and the 618 GHz line were both present, with emission line surface brightnesses in excess of  $2 \times 10^{-10} \text{ W m}^{-2} \text{ sr}^{-1}$ . These four spectra yielded a mean ‘rest’ frequency for the 618 GHz line of  $617.554 \pm 0.209 \text{ GHz}$ . The 1235 GHz line was detected in five SSW spectra (A2, B1, B2, B3 and D4) but only in the B1 (Fig. 2) and B3 spectra were both it and the 971.8038 GHz  $\text{OH}^+$  line present with a single unresolved component. The  $J = 1\text{--}1$ ,  $F = 1/2\text{--}1/2$  line of  $\text{OH}^+$  at 1032.998 GHz was also detected in emission in the SSW B1 spectrum, enabling a third estimate of the rest frequency of the 1235 GHz line. The mean frequency derived from these three estimates was  $1234.786 \pm 0.643 \text{ GHz}$ .

The ratio of the derived rest frequencies of  $1234.786 \pm 0.643 \text{ GHz}$  and  $617.554 \pm 0.209 \text{ GHz}$

is  $1.9995 \pm 0.0012$ , which suggests that the lines correspond to the 2-1 and 1-0 rotational transitions of a simple diatomic molecule (we can rule out their being 4-3 and 2-1 transitions, with a frequency ratio of 4:2, because of the lack of a corresponding 3-2 transition at  $\sim 926$  GHz). A search using the *Cologne Database for Molecular Spectroscopy* (12) and the MADEX code (17) found the only candidate to be  $^{36}\text{ArH}^+$ , whose 1-0 and 2-1 rotational transitions lie at  $617.52523 \pm 0.00015$  GHz and  $1234.60275 \pm 0.00030$  GHz, respectively, agreeing with the derived frequencies for the Crab Nebula lines within the uncertainties. The 1-0 and 2-1 rotational transitions of  $^{40}\text{ArH}^+$  are at  $615.85813 \pm 0.00005$  GHz and  $1231.27100 \pm 0.00009$  GHz, while the corresponding transitions of  $^{38}\text{ArH}^+$  are at  $616.64871 \pm 0.00004$  GHz and  $1232.85100 \pm 0.00004$  GHz, ruling out these two isotopic variants as identifications. Argon is the third most abundant species in the Earth's atmosphere, 0.93% by number, with  $^{40}\text{Ar}/^{38}\text{Ar}/^{36}\text{Ar}$  isotopic ratios of to 1584/1.00/5.30 (18). However,  $^{40}\text{Ar}$  in the earth's atmosphere is a product of the decay, mainly in rocks, of  $^{40}\text{K}$ , whose half-life is  $1.25 \times 10^9$  yrs. For the solar wind the  $^{40}\text{Ar}/^{38}\text{Ar}/^{36}\text{Ar}$  isotopic ratios have been measured to be 0.00/1.00/5.50 (19).  $^{36}\text{Ar}$  is expected to be the dominant isotope of argon in stars, being an explosive nucleosynthesis product of the  $\alpha$ -particle capture chain that takes place in massive star core-collapse supernovae (7). Regions of enhanced emission from optical forbidden lines of ionized argon have previously been mapped in the Crab Nebula by a number of authors (20–22), almost certainly corresponding to enriched argon abundances. Strongly enhanced infrared forbidden lines of  $\text{Ar}^+$  and  $\text{Ar}^{2+}$  have been detected from the southern filament (22) where we find  $^{36}\text{ArH}^+$  line emission to be strongest (SSW detector B1; Fig. 1 and Table 1). The enhanced lines of ionized argon and  $^{36}\text{ArH}^+$  strongly indicate the presence there of pockets of  $^{36}\text{Ar}$  produced by explosive nucleosynthesis during the supernova event.

The  $^{38}\text{Ar}$  isotope of argon is also predicted to be synthesised in core-collapse supernova events (7). We used Crab Nebula SPIRE-FTS spectra in which the  $^{36}\text{ArH}^+$  velocity components

are strong and narrow to put limits on the isotopic ratios of  $^{40}\text{Ar}$  and  $^{38}\text{Ar}$  relative to  $^{36}\text{Ar}$  in the emitting regions. Because the frequency separation between the isotopic variants of  $\text{ArH}^+$  is a factor of two larger for the  $J = 2-1$  lines than for the  $1-0$  lines, whereas the SPIRE-FTS frequency resolution is constant, the  $2-1$  lines are better suited for placing limits on the isotopic ratios of argon. Relative to the  $^{36}\text{ArH}^+$   $2-1$  line, the separation of the  $^{40}\text{ArH}^+$   $2-1$  line is  $-3.33175$  GHz, while that of the  $^{38}\text{ArH}^+$   $2-1$  line is  $-1.75175$  GHz. To estimate upper limits to the abundances of these species, we added synthetic lines to the spectra having the appropriate frequency offset and the same line width as the  $^{36}\text{ArH}^+$   $2-1$  line, changing the strength of the synthetic lines until the signal-to-noise ratio estimated by the line-fitting routine reached  $3\sigma$ . The SSW B1 (Fig. 2) and SSW B3 spectra both yielded  $3\sigma$  lower limits to the abundance ratio of  $^{36}\text{ArH}^+ / ^{38}\text{ArH}^+$  of  $> 2$ , along with  $^{36}\text{ArH}^+ / ^{40}\text{ArH}^+$  lower limits of  $> 5$  and  $> 4$ , respectively.

$\text{ArH}^+$  is a stable molecular ion (dissociation energy  $D_0 = 3.9 \pm 0.1$  eV (1)) that has been studied extensively in the laboratory. The Crab Nebula consists predominantly of ionized gas, photoionized by synchrotron radiation from the pulsar wind nebula (5, 16). It also contains many  $\text{H}_2$ -emitting neutral clumps (23, 24). Transition zones between fully ionized and molecular gas will exist, where  $\text{ArH}^+$  can be formed by the exothermic reaction  $\text{Ar}^+ + \text{H}_2 \rightarrow \text{ArH}^+ + \text{H}$ , releasing 1.49 eV (25). If the elemental species created by the supernova explosion were still largely unmixed in the remnant, then it is possible that  $\text{ArH}^+$  molecules would be found only at interfaces between H-rich gas and Ar-rich gas where mixing has occurred. Four of the seven FTS SSW detectors in whose spectra  $J = 2-1$   $^{36}\text{ArH}^+$  emission was detected (SSW B1, B2, A1 and A2) are situated on a bright filament south of the center of the nebula (Fig. 1), as is the SLW D4 detector in which the strongest  $J = 1-0$  emission was detected. A cluster of seven near-infrared  $\text{H}_2$ -emitting knots, with a wide range of radial velocities, is coincident with the same bright filament (24). Detectors SLW C3/SSW D4 also show  $^{36}\text{ArH}^+$  emission and are coincident with an  $\text{H}_2$ -emitting knot. The lack of  $\text{ArH}^+$  emission in the NW quadrant of the

nebula is mirrored by a relative lack of  $\text{H}_2$  emission knots in that region. However, there are many  $\text{H}_2$  knots in the NE quadrant, whereas  $\text{ArH}^+$  emission is only detected there in two FTS detectors (SLW E1 and E2).

The reaction rate for the formation of  $\text{ArH}^+$  via  $\text{Ar}^+ + \text{H}_2 \rightarrow \text{ArH}^+ + \text{H}$  is  $8.9 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  at 300 K (25). Reaction rates are known for the  $\text{ArH}^+$  destruction reaction with  $\text{H}_2$  (to yield  $\text{Ar} + \text{H}_3^+$ , with a reaction rate of  $6.3 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  at 300 K (26)) and for its dissociative recombination with electrons ( $\text{ArH}^+ + \text{e}^- \rightarrow \text{Ar} + \text{H}^*$ , with a rate  $\leq 5 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$  at low electron energies (27)). However, because the Crab nebula is photoionized by its pulsar wind nebula (5, 16), photodissociation could be the main  $\text{ArH}^+$  destruction mechanism.

The main excitation mechanism for the observed  $\text{ArH}^+$  emission lines is likely to be collisions with either electrons or  $\text{H}_2$  molecules but rate calculations or measurements do not exist as yet. The  $J = 1$  and 2 levels of  $^{36}\text{ArH}^+$  are situated 29.6 K and 88.9 K above the ground state, respectively, negligible compared to the electron temperatures of 7500-15000 K measured for the ionized gas in the Crab Nebula (5), or even compared to the  $\text{H}_2$  excitation temperatures of 2000-3000 K that have been measured (28). If the electron or  $\text{H}_2$  densities in the transition zones where  $\text{ArH}^+$  is hypothesised to be located should exceed the ‘critical densities’ of the emitting levels (where the sum of the collisional excitation and de-excitation rates from a level exceed the radiative decay rate from the level), then the level populations will be in Boltzmann equilibrium. Using the known molecular parameters of  $\text{ArH}^+$  (12, 17) the 2-1/1-0 line emission ratios should then be of the order 30, for excitation temperatures appreciably exceeding 100 K. The SSW D4 and SLW C3 detectors are centred on the same bright knot (Fig. 1) and yield a 2-1/1-0 line surface brightness ratio of 2.5, while the spectra from the approximately co-located SSW B3 and SLW C4 detectors yield a line surface brightness ratio of 2.0 (Table 1), well below the ratio for Boltzmann equilibrium. The densities of the collision partners in the emitting regions must therefore be well below the corresponding critical densities of the  $\text{ArH}^+$

rotational levels. We used the MADEX code (17) with the molecular parameters of  $\text{ArH}^+$ , together with  $\text{SiH}^+ + \text{He}$  collisional de-excitation rates (29) in place of those of  $\text{ArH}^+ + \text{H}_2$ , and  $\text{CH}^+ + \text{e}^-$  collisional de-excitation rates (30) in place of those for  $\text{ArH}^+ + \text{e}^-$  (with upward rates calculated using detailed balance and the correct values of the energies for the levels of  $\text{ArH}^+$ ) in order to estimate corresponding  $\text{H}_2$  and electron critical densities of  $\sim 10^8 \text{ cm}^{-3}$  and  $\sim 10^4 \text{ cm}^{-3}$ , respectively. The observed 2-1/1-0 line ratios of 2.5 and 2.0 indicate  $\text{H}_2$  densities of a few  $\times 10^6 \text{ cm}^{-3}$ , or electron densities of a few  $\times 10^2 \text{ cm}^{-3}$ . The calculations take into account opacity effects although the line centre optical depths are estimated to be significantly less than unity for line widths larger than  $1 \text{ km s}^{-1}$ . For  $\text{H}_2$  collisions at temperatures between 100 K and 3000 K and densities of  $\sim 10^6 \text{ cm}^{-3}$ , or electron collisions at temperatures of  $\sim 3000 \text{ K}$  and densities of a few  $\times 10^2 \text{ cm}^{-3}$ , we estimate  $^{36}\text{ArH}^+$  column densities of  $10^{12} - 10^{13} \text{ cm}^{-2}$ .

Given that likely electron collisional excitation rates are  $\sim 10^4$  larger than those of  $\text{H}_2$  and that the parent  $\text{Ar}^+$  ion must exist in a region that is at least partially ionized, electron collisions are expected to dominate the excitation of  $\text{ArH}^+$ . A density of  $\sim 10^4 \text{ cm}^{-3}$  has been estimated for the Crab  $\text{H}_2$  knots (28), a factor of 100 below that required for  $\text{H}_2$  collisions to be the excitation mechanism for the  $\text{ArH}^+$  lines. This lends further support to electron collisions in partially ionized transition regions being the main  $\text{ArH}^+$  excitation mechanism.

Our detection of  $^{36}\text{ArH}^+$  in the Crab Nebula suggests that an unidentified multi-component broad absorption feature seen between 617 and 618 GHz in a Herschel HIFI spectrum of Sgr B2(M) (towards the center of our galaxy) (31) can potentially be identified with ground-state absorption in the  $J = 0-1$  617.525 GHz line of  $^{36}\text{ArH}^+$  along the interstellar sightline.

## References and Notes

1. J. R. Wyatt *et al.*, *J. Chem. Phys.* **62**, 2555 (1975).



2. S. G. Lias, J. F. Liebman, R. D. Levin, *J. Phys. Chem. Ref. Data* **13**, 695 (1984).
3. J. M. Moorhead, R. P. Lowe, W. H. Wehlau, J.-P. Maillard, P. F. Bernath, *Astrophys. J.* **326**, 899 (1988).
4. X.-W. Liu *et al.*, *Mon. Not. R. Astron. Soc.* **290**, L71 (1997).
5. K. Davidson, R. A. Fesen, *Annu. Rev. Astron. Astrophys.* **23**, 119 (1985).
6. G. L. Pilbratt *et al.*, *Astron. Astrophys.* **518**, L1 (2010).
7. D. Arnett, *Supernovae and Nucleosynthesis*, (Princeton Univ. Press, 1996).
8. M. J. Griffin *et al.*, *Astron. Astrophys.* **518**, L3 (2010).
9. B. M. Swinyard *et al.*, *Astron. Astrophys.* **518**, L4 (2010).
10. M. Groenewegen *et al.*, *Astron. Astrophys.* **526**, A162 (2011).
11. S. Ott, *Astr. Soc. Pacific Conf. Ser.* **434**, 139 (2010).
12. H. S. P. Müller, F. Schlöder, J. Stutzki, G. Winnewisser, *J. Mol. Struct.* **742**, 215 (2005).
13. D. A. Neufeld *et al.*, *Astron. Astrophys.* **521**, L10 (2010).
14. P. P. van der Werf *et al.*, *Astron. Astrophys.* **518**, L42 (2010).
15. L. Spinoglio *et al.*, *Astrophys. J.* **758**, 108 (2012).
16. J. J. Hester, *Annu. Rev. Astron. Astrophys.* **46**, 127 (2008).
17. J. Cernicharo, *EAS Publ. Ser.* **58** 251 (2012).
18. J.-Y. Lee *et al.*, *Geochim. et Cosmochim. Acta* **70**, 4507 (2006).

19. A. Meshik *et al.*, *Science* **318**, 433 (2007).
20. G. M. MacAlpine *et al.*, *Astrophys. J.* **432**, L131 (1994).
21. E. L. Schaller & R. A. Fesen, *Astron J.* **123**, 941 (2002).
22. T. Temim *et al.*, *Astrophys. J.* **753**, 72 (2012).
23. J. R. Graham, G. S. Wright, A. J. Longmore, *Astrophys. J.* **352**, 172 (1990).
24. E. F. Loh *et al.*, *Astrophys. J. Suppl. Ser.* **194**, 30 (2011).
25. A. C. Roach, P. J. Kuntz, *Chem. Comms.* 1336 (1970).
26. V. G. Anicich, *NASA JPL Publ.* 03-19 (2003).
27. J. B. A. Mitchell *et al.*, *J. Phys. B* **38**, L175 (2005).
28. E. D. Loh *et al.*, *Mon. Not. R. Astron. Soc.* **421**, 789 (2012).
29. C. Nkem *et al.*, *J. Mol. Struct. THEOCHEM* **901**, 220 (2009).
30. A. J. Lim, I. Rabadán, J. Tennyson, *Mon. Not. R. Astron. Soc.* **306**, 473 (1999).
31. P. Schilke, HEXOS Team, in *Proceedings of the Herschel First Results Symposium*, (2010).  
`\protect\vrule width0pt\protect\href{http://herchel.esac.esa.int/FirstResultsSymposium/presentations/A34_Schi`
32. H. L. Gomez *et al.*, *Astrophys. J.* **760**, 96 (2012).
33. We thank the anonymous referees for their constructive reports. We thank Dr Edwin Bergin for drawing our attention to the 2010 HEXOS Herschel HIFI spectrum of SgR B2(M) that shows an unidentified broad absorption feature below 618 GHz. *Herschel* is an ESA space

observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. SPIRE has been developed by a consortium of institutes led by Cardiff University (UK) and including Univ. Lethbridge (Canada); NAOC (China); CEA, LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain); Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, Univ. Sussex (UK); and Caltech, JPL, NHSC, Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MINECO (Spain); SNSB (Sweden); STFC and UKSA (UK); and NASA (USA). JC thanks MINECO for financial support under projects AYA2009-07304, AYA2012-32032 and Consolider program ASTROMOL CSD2009-00038.

Table 1: SPIRE-FTS radial velocity and line surface brightness measurements for the J = 1-0 and 2-1 rotational lines of  $^{36}\text{ArH}^+$  from the Crab Nebula

SLW Detector	J = 1-0 617.525 GHz		SSW Detector	J = 2-1 1234.603 GHz	
	Radial Velocity $\text{km s}^{-1}$	Surface Brightness $10^{-10} \text{ W m}^{-2} \text{ sr}^{-1}$		Radial Velocity $\text{km s}^{-1}$	Surface Brightness $10^{-10} \text{ W m}^{-2} \text{ sr}^{-1}$
B3	$+317 \pm 67$	$2.23 \pm 0.41$	C5	$-1354 \pm 26$	$8.2 \pm 1.2$
C3	$+933 \pm 33$	$4.63 \pm 0.40$	D4	$+743 \pm 26$	$11.7 \pm 1.6$
C4	$-58 \pm 50$	$8.65 \pm 0.55$	B3	$-101 \pm 20$	$17.5 \pm 1.4$
D3	$+826 \pm 32$	$3.13 \pm 0.34$			
D3	$-709 \pm 42$	$2.30 \pm 0.34$			
D4	$+101 \pm 27$	$9.89 \pm 0.52$	A1	$-51 \pm 52$	$13.9 \pm 2.0$
			B2	$-572 \pm 25$	$10.8 \pm 1.7$
			B1	$+140 \pm 34$	$38.4 \pm 1.6$
			A2	$+61 \pm 28$	$10.1 \pm 1.4$
E1	$+278 \pm 46$	$5.69 \pm 0.62$			
E2	$-594 \pm 37$	$4.25 \pm 0.46$			

**Fig. 1.** A broad-band Herschel-PACS image mapping the 70- $\mu\text{m}$  dust emission from the Crab Nebula. North is up and East is to the left. The positions on the nebula of the 19 SLW and 35 SSW detectors of the SPIRE FTS are marked with circles whose angular diameters of 37'' and 18'' correspond to the SLW and SSW beam sizes. The positions of detectors in whose spectra the  $J = 1-0$  or  $2-1$  rotational lines of  $^{36}\text{ArH}^+$  were detected are marked with crosses.

**Fig. 2.** SPIRE FTS spectra of the Crab Nebula, plotting surface brightness in  $\text{W m}^{-2} \text{Hz}^{-1} \text{sr}^{-1}$  against frequency in GHz. Several emission lines are superposed on a continuum attributed to thermal dust emission (32). Upper plot: the spectrum from the SLW D4 detector, whose position on the nebula is marked in Fig. 1. Emission line velocity components attributed to the  $J = 2-1$ ,  $F = 5/2-3/2$  971.8038 GHz rotational line of  $\text{OH}^+$  and the  $J = 1-0$  617.525 GHz rotational line of  $^{36}\text{ArH}^+$  are visible. Lower plot: the SSW B1 spectrum. In addition to  $\text{OH}^+$  971.8038 GHz velocity components, emission in the  $J = 2-1$  1234.603 GHz line of  $^{36}\text{ArH}^+$  is visible. The radial velocities and surface brightnesses of the  $^{36}\text{ArH}^+$  emission lines that are present in the spectra obtained from these and other FTS detectors are listed in Table 1.



