

Department of Chemistry
Adran Chemeg



**Mixed donor carbene pyridyl ligands and
their metal complexes**

by

Aminu Ahmed

Submitted in fulfilment of the requirements for the Degree of

Doctor of Philosophy

School of Chemistry
Cardiff University
Wales, UK

2008

UMI Number: U585278

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI U585278

Published by ProQuest LLC 2013. Copyright in the Dissertation held by the Author.
Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against
unauthorized copying under Title 17, United States Code.



ProQuest LLC
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106-1346

Abstract

This thesis describes the synthesis of a series of Ag(I), Pd(II), Rh (I) and Ir(I) complexes of quinoline functionalised nucleophilic heterocyclic carbene (NHC) ligands. The transmetallation properties of the Ag(I) complexes were utilised to prepare the corresponding Pd(II), Rh (I) and Ir(I) (NHC) complexes.

A series of quinoline based imidazolium, pyrimidinium salts were prepared and characterised as NHC ligand precursors.

Ag(I)(NHC) complexes were prepared by the reaction of the quinoline functionalised salts with Ag₂O in DCM. All complexes were spectroscopically characterised and the results of single X- ray crystallographic studies are reported for two of the complexes and the geometry around the silver cation was observed to be distorted linear.

Two quinoline based palladium (II) (NHC) complexes were prepared via transmetallation Ag(I)(NHC) complexes is reported.

The synthesis of a series of methylene bridged quinoline functionalised Rh (I) and Ir(I) (NHC) complexes through transmetallation of the Ag(I)(NHC) complexes is reported and the results of single X- ray crystallographic studies are reported for most of the complexes showing consistent pattern in term of bond lengths and angles.

Two of the Ir(I) (NHC) complexes were tested as catalysts in transfer hydrogenation reactions , showing good activity at low Ir loadings.

Acknowledgements

First and foremost I would like to thank God for creating me and giving me the health to carry out this research work.

I would like to express my profound gratitude and thanks to Professor Kingsley Cavell for his enthusiastic support and endless stream of ideas that was not diminished throughout the period of this work.

My thanks go to Dr. D. J. Beetstra and Dr. Benson Kariuki at the Cardiff University for their sterling X-ray crystallography efforts. Their input has been essential to the understanding of the systems studied in this project.

I am indebted to members of Cavell group especially Dr. David Nielsen for teaching me how to use my line; Tracy Hamilton for going through my manuscript; Manuel Alonso, Emma Jones, Deborha Bacciu, Adrien Normand, Huw, Vanessa and Abeer Binobaid for their assistance and encouragements.

My profound thanks go to my fellow colleagues Abdullahi Nuhu, Ado Mukhtar Bichi, Y. B. Daraja, Abdullahi Mustapha, Abdulhadi Aminu, Ibrahim Abdullahi, and Bello Y. Makama for their advice and encouragements.

Financial support from Kano University of Science and Technology in the form of foreign postgraduate fellowship is gratefully acknowledged. My sincere gratitude and appreciations go to the Vice Chancellor of the KUST Professor Ibrahim Shehu Diso for taking the pain to see that my study goes on without significant financial difficulty.

I would also like to thank the staff and fellow students of the School of Chemistry for fostering a friendly and inclusive environment for study.

Many people in Cardiff and Bristol deserve acknowledgement especially my friends such as Haruna Musa, Ahmed Ali, Mukhtar Atiku Kurawa, and Dr. Shehu Yakasai. To my friends in Nigeria, Yahaya Ahmed, Ismaila Usman, Ibrahim Kutama and Shuaibu Ahmed for looking after my family.

Lastly, to my family especially my wife: Malama Rabi Sani, children; Salim, Halima, Maryam, and Haruna for their endurance and prayers throughout the course of this study.

To all of you my grateful thanks!

Abbreviations used in this Thesis

| | |
|--------------------------|---------------------------------------------|
| AcOH | acetic acid |
| Ar | aryl group |
| Barf | tetrakis[(3,5-trifluoromethyl)phenyl]borate |
| BAP | 4-bromoacetophenone |
| ⁿ Bu | n-butyl group |
| ^t Bu | tert-butyl group |
| COD or cod | 1, 5-cyclooctadiene |
| DCM | dichloromethane |
| DMF | N, N-dimethylformide |
| DMSO | dimethyl sulfoxide |
| ESMS | electrospray mass spectrometry |
| Et | ethyl group |
| Et ₂ O, ether | diethyl ether |
| GC | gas-liquid chromatography |
| GCMS | gas-liquid chromatography/mass spectrometry |
| HRMS | high resolution mass spectrometry |
| ⁱ Pr | iso-propyl group |
| ⁿ Pr | n-propyl group |
| IR | infra red spectrometry |
| LSIMS | liquid secondary ion mass spectrometry |
| M | metal |
| Me | methyl group |
| Mes | mesityl group |
| MS | mass spectrometry |
| NBS | N-bromosuccinimide |
| NHC | nucleophilic heterocyclic carbene |
| NMR | nuclear magnetic resonance |
| OAc | acetate ion |
| OTf | trifluoromethanesulfonate (triflate) anion |
| Ph | phenyl group |
| Qiun | qiunolyl group |

| | |
|----------|------------------------------------------------------------------------------------|
| R | alkyl group |
| THF, thf | tetrahydrofuran |
| TON | turnover number ($\text{mol}_{\text{product}}/\text{mol}_{\text{Ir}}/\text{hr}$) |
| TMS | tetra methyl silane |

Table of Contents

CHAPTER ONE: INTROUCTION

| | | |
|---------|---------------------------------------------------------------------|----|
| 1.1 | Background..... | 1 |
| 1.2 | Brief History of Carbene..... | 3 |
| 1.3 | Synthesis of diaminocarbenes and pKa..... | 7 |
| 1.4 | NHC Ligand properties..... | 8 |
| 1.5 | Chelating, Pincer and Mixed donor Ligands..... | 10 |
| 1.6 | Carbene complexes..... | 19 |
| 1.6.1 | Synthesis of N-Heterocyclic carbene transition metal complexes..... | 21 |
| 1.6.1.1 | In-situ deprotonation of azolium salts..... | 21 |
| 1.6.1.2 | Carbene Complexes through free carbenes..... | 23 |
| 1.6.1.3 | Carbene complexes through transmetallation..... | 24 |
| 1.6.1.4 | Carbene complexes through oxidative addition reactions..... | 24 |
| 1.7 | Abnormal carbene complexes..... | 25 |
| 1.8 | Catalysis involving NHCs..... | 26 |
| 1.8.1 | Ruthenium metathesis..... | 26 |
| 1.8.2 | Asymmetric catalysis..... | 27 |
| 1.8.3 | Hydrogenation..... | 28 |
| 1.8.4 | Catalyst decomposition..... | 30 |
| 1.9 | Aims and thesis overview..... | 30 |
| 1.10 | References..... | 31 |

CHAPTER TWO: QUINOLINE FUNCTIONALISED IMIDAZOLIUM AND PYRIMIDIUM SALTS

| | | |
|------------|--------------------------------------------------------------------------|-----------|
| 2.1 | Introduction | 40 |
| 2.1.1 | Ligand synthesis..... | 42 |
| 2.2 | Results and Discussion | 44 |
| 2.2.1 | Synthesis and characterisation of symmetrical saturated imidazolium salt | 44 |
| 2.2.2 | Synthesis and characterisation of bis quinoline pyrimidinium salt..... | 46 |

| | | |
|------------|-----------------------------------------------------------------------------------------------------|----|
| 2.2.3 | Unsymmetrical substituted quinoline imidazolium salt..... | 48 |
| 2.2.4 | Synthesis and characterisation of methylene-bridged quinoline functionalised imidazolium salts..... | 52 |
| 2.2.5 | Synthesis and characterisation of octahydroacridine based imidazolium salt | 57 |
| 2.3 | Conclusions | 60 |
| 2.4 | Experimental | 62 |
| 2.4.1 | General comments..... | 62 |
| 2.4.2 | N-Substituted imidazoles..... | 63 |
| 2.4.3 | Symmetrical saturated bis(1,3-quinoline)imidazolium salt..... | 65 |
| 2.4.4 | Bis(1, 3-quinoline)pyrimidinium hexafluoro phosphate salt..... | 66 |
| 2.4.5 | Unsymmetrical substituted quinoline imidazolium salts..... | 67 |
| 2.4.6 | Methylene-bridged quinoline functionalised imidazolium salts..... | 69 |
| 2.4.7 | Acridine based imidazolium salt..... | 73 |
| 2.5 | Crystal structure solution..... | 77 |
| 2.6 | References | 83 |

CHAPTER THREE: SILVER(I) AND PALLADIUM(II) COMPLEXES OF QUINOLINE FUNCTIONALISED HETEROCYCLE CARBENE LIGANDS

| | | |
|------------|-----------------------------------------------------------------------------------------------|-----|
| 3.1 | Silver(I) (NHC) complexes | 86 |
| 3.1.1 | Introduction | 86 |
| 3.1.2 | Structural diversity in Ag ^I (NHC) complexes..... | 90 |
| 3.1.3 | Ag ^I (NHC) complexes as transmetallation reagents..... | 93 |
| 3.2 | Palladium(II) complexes of quinoline functionalised heterocyclic Carbene ligands | 94 |
| 3.3 | Results and Discussions | 97 |
| 3.3.1 | Silver(I) complexes rigid quinoline functionalised NHC ligands..... | 97 |
| 3.3.1.1 | Silver(I) complexes of rigid quinoline functionalised NHC ligands..... | 97 |
| 3.3.1.2 | Silver(I) complexes of methylene-bridged quinoline functionalised NHC ligands..... | 98 |
| 3.3.2 | Pd(II) quinoline functionalised (NHC) carbene complexes..... | 105 |
| 3.4 | Conclusions | 106 |
| 3.5 | Experimental | 107 |
| 3.5.1 | General comments..... | 107 |

| | | |
|------------|------------------------------------------------------------------------------------|------------|
| 3.5.2 | Silver(I) complexes of rigid quinoline functionalised mono-NHC ligands..... | 108 |
| 3.5.3 | Silver(I) complexes of methylene bridged quinoline functionalised NHC ligands..... | 109 |
| 3.5.4 | Pd(II) quinoline functionalised NHC complexes..... | 112 |
| 3.5.5 | X-ray crystallography..... | 114 |
| 3.6 | References | 116 |

CHAPTER FOUR: RHODIUM(I) AND IRIDIUM(I) COMPLEXES OF QUINOLINE FUNCTIONALISED HETEROCYCLIC CARBENE LIGANDS

| | | |
|------------|-------------------------------------------------------------------------------------|------------|
| 4.1 | Introduction | 120 |
| 4.2 | Results and Discussions | 123 |
| 4.2.1 | Synthesis and characterisation of quinoline functionalised Rh(I) NHC Complexes..... | 123 |
| 4.2.2 | Synthesis and characterisation of quinoline functionalised Ir(I) NHC Complexes..... | 134 |
| 4.2.3 | Attempted synthesis of Rh(I) and Ir(I) NHC complexes without methylene Bridged..... | 140 |
| 4.3 | Ir(I) NHC complexes catalysed hydrogen transfer reactions..... | 142 |
| 4.3.1 | Catalysis..... | 142 |
| 4.3.2 | Hydrogen transfer reduction..... | 143 |
| 4.4 | Results and discussion..... | 145 |
| 4.4 | Conclusions | 147 |
| 4.5 | Experimental | 148 |
| 4.5.1 | General comments..... | 148 |
| 4.5.2 | Catalysis..... | 157 |
| 4.5.3 | Transfer hydrogenation..... | 157 |
| 4.5.4 | X-ray crystallography..... | 158 |
| 4.6 | References | 165 |

CHAPTER FIVE: CONCLUSIONS AND FUTURE

| | | |
|-----|-------------------|-----|
| 5.1 | Conclusions | 168 |
| 5.2 | Future work | 170 |
| 5.3 | References | 171 |

APPENDIX: TABLE OF BOND ANGLES ($^{\circ}$) AND BOND DISTANCES (\AA) 172

CHAPTER ONE

1.1 Background

Carbenes are neutral compounds featuring a divalent carbon atom with only six electrons in its valence shell [1]. Due to an incomplete octet of electrons, they are generally very reactive species. The carbenic carbon can be either linear or bent, each geometry describable by a certain degree of hybridization. The sp -hybridized carbon carbene adopts a linear geometry with two non bonding degenerate orbital (P_x and P_y). In sp^2 - type hybridization where the carbon adopts bent geometry, the degeneracy is broken in which case the P_x is stabilised by acquiring some s character (it is usually called σ while the P_y (usually called P_π) remains almost unchanged. It should be noted that most carbenes are bent (sp^2 hybridized).

As a result of this break in degeneracy, carbenes can exist in two states: triplet state (where two non bonding electrons occur in two different orbital with parallel spins) and singlet state (two non bonding electrons can pair in the same σ or P_π orbital). Based on this classification, we can have four possible electronic configurations as depicted in figure 1.1




Figure 1.1. Electronic configuration of carbenes.

The ground state spin multiplicity is a fundamental feature of carbenes that dictates their reactivity [2]. A singlet state is favoured by a large σ - P_π

separation and according to Hoffmann; a value of at least 2 eV is needed to impose this. Anything below 1.5 eV will favour the triplet state [3].

The substituents on the carbene determine whether a singlet or triplet state is formed through a combination of inductive, mesomeric and steric effects. Electron withdrawing groups increase the σ - $P\pi$ gap by inductively stabilising the σ non bonding orbital by increasing its s character and thus favouring the singlet state. On the other hand an σ -electron donating group will induce small σ - $P\pi$ gap and favour the triplet state. For illustration purposes substituents that are electron donating can be termed X, while electron withdrawing groups can be termed Y. XX carbenes are bent singlet, while most of the YY carbenes are predicted to be linear singlet.

N-Heterocyclic carbenes (NHCs) (Figure 1.2, (a)) are the focus of this work and are firmly placed within the singlet state. With two nitrogen substituents next to the carbene carbon atom, the NHCs are predicted to stabilise their singlet state (two paired electrons in the σ -orbital) by push- pull effect (Figure 1.2) [4]. Firstly, the σ - electron withdrawing nitrogen inductively stabilises the σ - non bonding orbital by increasing its s-character. Secondly, the energy of the vacant $P\pi$ -orbital is increased by interaction with the symmetric combination of the nitrogen lone pairs.




Figure 1.2: Electronic stabilisation of NHCs.

The combination of the two effects increases the σ - $P\pi$ gap and favours therefore the singlet state. Additionally, the sp^2 hybridization adopted by the carbene carbon atom in its singlet state matches the bent geometry of the NHC five membered ring. The interaction of the nitrogen lone pair with the π -orbital of the

carbene is reflected by an N-C carbene bond length of 1.365 Å, which is consistent with double bond character. An accurate assessment of the π back bonding was found by analysing dynamic $^1\text{H-NMR}$ behaviour of bis (diisopropylamine) carbene **3** [5]. The major process involves rotation about the N-C carbene bonds; the measured barrier to rotation of 53 kJ/mol was mostly attributed to the substantial π - component of these bonds.

Dimerisation of NHCs has been known since the first attempt to isolate them [6]. Alder recently showed that dimerisation is thermodynamically unfavourable for imidazolin-2-ylidenes **1** (singlet/triplet gap of 354 KJ /mol), but very likely to happen for imidazolidin-2-ylidenes **2** due to lack of aromaticity and acyclic NHCs due to loss of conjugation through twisting around N-C carbene bond [7]. The $^{13}\text{C-NMR}$ chemical shifts [1] range from 210-220 ppm downfield from TMS for aromatic imidazolin-2-ylidenes **1** and to 235-245 ppm for imidazolidin-2-ylidenes **2** and acyclic NHCs **3** (Figure 1.3).




Figure 1.3: Unsaturated, saturated NHCs and acyclic carbenes

1.2 Brief History of Carbenes

Since the pioneering work of Doering in 1954, carbenes have been recognised as a unique type of intermediate with characteristics distinct from radicals already known in the organic chemistry community [8]. A decade later Wanzlick and co-workers started working on saturated N-heterocyclic system where they explored the dimers of electron rich tetraaminoethylenes to which they proposed

an equilibrium existed between the free carbenes and the dimer and was named the 'Wanzlick Equilibrium' [6, 9-11] (Figure 1.4). The Wanzlick equilibrium was recently confirmed following observation of equilibrium mixtures between free carbenes and tetraaminoethylenes for some benzimidazolin-2-ylidenes [11, 12, and 14].




Figure 1.4: The Wanzlick equilibrium.

Wanzlick proposed that the dimer dissociated into two carbenes [13].

Since then research on carbenes has rapidly expanded, but almost no attempts were made to prepare stabilised carbenes until 1980 when Tomoika started to study persisted triplet diarylcarbenes [14].

The first isolable carbenes were reported in 1988 by Bertrand [15] **4** and in 1991 by Arduengo [16] **5**. Phosphinocarbene **4** can be distilled at 80-85 °C/10-2 Torr and N-heterocyclic carbene (NHC) **5** is crystalline solid that melts at above 240-241 °C (Figure 1.5).




Figure 1.5: The first isolated carbenes

Although NHCs have been known since the pioneering work of Wanzlick, who observed their dimerisation and was able to trap them to form mercury salt carbene complexes [17], three decades went by before the first NHC was isolated. When compound **5** was isolated, the stability of the compound was thought to be due to steric hindrance of the bulky adamantyl substituents preventing nucleophilic attack [16]. However in 1995, Arduengo proved using NHC carbene **2** that aromaticity was not needed for stabilisation, [18] and in 1996 Alder isolated acyclic NHC **6** [19]. This research area continues to expand with the isolation of four-membered carbene **7** [20] by Grubbs and alkyl carbene **8** by Bertrand in 2004 [21]. Arduengo reported the synthesis of 1,3,4,5-tetramethylimidazolin-2-ylidene **9** and 1,3-dimethylimidazolin-2-ylidene **10** [22] which are less sterically hindered. Carbenes with only one nitrogen atom have also been isolated as indicated in compounds **8** and **11**. This shows that the presence of one nitrogen atom is adequate enough to stabilise carbenes in certain cases provided the carbene carbon is bound to a tertiary alkyl group. The replacement in compound **11** of one of the electronegative nitrogens by a strong σ -donor alkyl group makes the ligand more electronegative than diamino NHCs and therefore behave as strong σ donor towards transition metal centres [21]. Carbenes with more than two heteroatoms (**12** [22]) and those with a mixture of heteroatoms have also appeared in the literature (**13** [23] and **14** [24]). Hermann and co-workers reported the synthesis of chiral **15** and bis-imidazol-2-ylidenes **16** [25].





Figure 1.6: Stable carbenes and their derivatives

1.3 Synthesis of diaminocarbenes and pKa

Three principal methods have been used to successfully generate diaminocarbenes:

- (i) Deprotonation of imidazolium salt **17** or formamidinium salts **18** with a base [16] (ii) Desulfurisation of thioureas **19** [26] (iii) thermolysis of methanol adducts of type **20** [22,95] (Scheme 1.0)



Scheme 1.0: methods of generating NHCs.

The measured pKa value for diisopropylimidazolin-2-ylidenes on the DMSO scale was found to be 24 by Alder [27, 28]. For di-tert-butylimidazolin-2-ylidenes Streitweiser reported a pKa of 20 on the THF scale [29]. It is therefore not surprising that the principal method used in the synthesis of NHCs is by deprotonation of the corresponding imidazolium or formamidinium salts. To synthesize the first NHCs Arduengo used NaH/KH in THF in the presence of KOtBu and DMSO [16].

Herrmann showed that milder conditions such as the use of sodium amide in liquid ammonia and THF at -40°C, were efficient [30]. When the pKa is

increased by 2 to 6 units, formamidinium salts underwent nucleophilic addition of the base rather than deprotonation [28]. Hindered alkali amide bases such as lithium diisopropylamide or potassium hexamethyldisilazide were used to overcome this drawback.

Kuhn and Kratz reported another pathway to imidazolin-2-ylidene by reduction of thioureas using metallic potassium [30]. Though this has been difficult to reproduce [28], it is an interesting discovery because the only other product, potassium sulphide, is insoluble in THF and therefore, can easily be removed.

In another method triazol-2-ylidene was synthesised in good yield by Enders by thermolysis of the corresponding methanol adducts [22]. However, this method has some disadvantages due to the extreme sensitivity of the methanol adduct.

1.5 NHC Ligand properties

N-Heterocyclic carbenes (NHCs) are ligands formed by the deprotonation of an N, N-disubstituted imidazolium (or other azolium) salts. Binding of a transition metal to the C2 carbon of the NHC leads to the formation of a very strong bond, the strength deriving from the thermodynamic instability of the free NHC [31]. Unlike metal-carbon bonds in general, those to NHCs do not undergo fast insertion or reductive elimination reactions and so NHCs are generally good spectator ligands. The role of spectator ligand is to act as a placeholder by promoting a desired reaction at the metal, while avoiding dissociation or entering directly into the reaction. NHCs being used as spectator ligands for many decades have risen to prominence, having both steric and electronic tenability and capability to promote catalysis of many useful reactions.

The bonding mode of metal carbene in Schrock and Fischer carbene complexes are both described by double a bond, though they differ by the polarity of the electron density. This difference arises from the difference in energy between the $d\pi$ orbital of the metal and the $p\pi$ orbital of the carbene (Figure 1.6). If the $d\pi$ orbital is lower in energy than the $p\pi$ orbital, the metal – carbon is polarised δ^- and δ^+ on the carbene and we would have a Fischer carbene complex. On the contrary, if the $d\pi$ orbital is higher in energy than the $p\pi$ orbital, the metal carbon bond is polarised δ^+ on the metal and δ^- on the carbene and we would

Chapter One: Introduction

expect a Schrock carbene complex. NHCs are firmly placed within the Fischer carbenes, their $p\pi$ orbital have high energy because multiple bonding between the carbene atom and the two nitrogen atoms. As a result the $p\pi$ orbital does not interact well with the $d\pi$ thus preventing almost any π - back bonding from the metal to the carbene. In NHC complexes the metal carbon bond is therefore best represented by a single bond.




Figure 1.7 Partial molecular diagrams for Schrock, Fisher and NHC carbene complexes.

The absence of a requirement for back bonding enable NHCs to form stable metal complexes with a range of metals that do not possess occupied orbitals and so would not be able to participate in back-bonding. Metals such as Li [32, 33], and Be [34, 35] and hexavalent U [36, 37] have been reported to form stable metal complexes with NHCs.

The fundamental difference between a typical Schrock carbene and NHC as ligand is shown in the crystal structure of $[\text{RuCl}_2(\text{NHC})_2(=\text{CHC}_6\text{H}_4\text{Cl})]$ (NHC = 1,3-diisopropylimidazol-2-ylidene) where the two types of the carbenes are linked together to the same metal centre [38]. The ruthenium-carbon bond of the Schrock carbene, generally written as a double bond, has a bond length of $1.821(3)\text{\AA}$, whereas the Ru-C bond length in NHC ($2.107(3)\text{\AA}$ and $2.115(3)\text{\AA}$)

which justifies its representation as a single bond (σ - donor and virtually no π -backbonding).

Measurement of IR carbonyl absorption frequencies of NHC carbonyl metal (Fe, Cr, Rh, Mo and Ir) and their phosphine analogues showed significantly donor capacity of NHC relative to phosphines, even to trialkylphosphines [39, 40, and 4]. Furthermore, experimental investigations [42], and calorimetric studies [43, 44] and experimental calculations [45] agree that the ligand dissociation energy of NHC from Ru complexes is higher than for phosphines. Similar results were obtained from calculations with other metals such as Au, Cu, Ag, Pd and Pt [46, 47].

1.6 Chelating, Pincer and Mixed donor Ligands

One of the attractive features of NHCs is the wide variety of steric [48] and asymmetric environments [49] that are available through modification of the substituents attached to the NHC nitrogen atoms. Furthermore, through the use of appropriate donor groups on the NHC substituents, it is possible to make multidentate NHCs [50]. Such variability makes possible the synthesis of NHC analogues of many traditional phosphine ligands. Through this synthetic modification, a wide range of ligands containing two or more NHCs groups are known. In 1994 Dias and Jin isolated bis-carbene **21** and tri-NHC ligand **22** [51] as shown in figure 1.8.





Figure 1.8: Chelating NHCs ligand

Through the use of appropriate donor groups on the NHC substituents, the coordination sphere of the metal can be tailored by chelating ligands incorporating a strongly bound, robust functional group, (NHC), with additional tethers carrying labile donors [52]. The latter can temporarily dissociate during catalytic reactions creating electronic coordinative unsaturation, which is important for catalysis. Towards this end Danopoulos et al have synthesised NHC precursors functionalised with pyridine rings. Almost simultaneously, related ligand designs were reported by Cavell [53] and Crabtree [54].


Danopoulos stated that the σ -donating pyridine rings tethered to the NHCs are believed to add versatility to the ligand designs for four reasons: (i) The pyridine function is expected to bind weakly to lower, softer oxidation states of the metal. (ii) This in combination with adjustment of the chelate ring size by using variable length linkers, can promote ligand hemilability with possible implications on the catalytic activity. (iii) The electronic asymmetry of the chelating N-functionalised NHC ligands renders the corresponding trans-sites electronically inequivalent, due to large difference in the trans effect of the chelating ends. (iv) The donor and steric characteristic of the pyridine and NHC functional groups are easily tunable by a variety of substituents [55]. It was along these lines that a number of research groups were able to prepare hemilabile pyridine functionalised ligands [53, 56, and 57]. A number of examples are depicted in Figure 1.8.

In order to test the hypothesis advanced for the choice of the pyridine functionalised ligands, the ligands shown below were used to prepare some carbene complexes and complexes tested in some catalysis. The reaction of the corresponding silver carbene Ag(I)NHC of **23** with PdMeCl(cod) gave PdMeCl(NHC). The catalytic activity of the palladium carbene was tested in Heck coupling reaction of 4-bromoacetophenone/4-chlorobenzaldehyde with n-butyl acrylate and the catalyst was found to have good activity and high stability. The Suzuki coupling of 4-bromoacetophenone with $C_6H_5B(OH)_2$ gave good to satisfactory result as well [53]. Danopoulos reported the synthesis of PdMeCl(NHC) from salts analogous to **23** which gave good activities in Heck and amination reactions [55]. Ligand **28** which is an example pyridinyl carbene (denoted as $pyN^{\wedge}C-R$) was used to prepare iridium (I) carbene complexes as


well as catalytic activities towards the hydrogen transfer reduction of nitroarenes under mild conditions [58].




23




24




25 R= iPr



26: R = iPr




27



28

Figure 1.9 Pyridyl functionalised Ligands

The reaction of the corresponding silver carbene **30** with $[\text{Ir}(\text{COD})\text{Cl}]_2$ gave the unchelated compound **31**. Chelation of the $\text{pyN}^{\wedge}\text{C-R}$ toward the iridium centre was achieved by the treatment of **31** with an equimolar amount of AgBF_4 , leading to the ligand substitution of chloride by the pyridine nitrogen to give compound **32** as shown in Scheme 2.





Scheme 2: Reagents and condition (i) Ag₂O, NaI, r.t. 24 hrs. (ii) [Ir (COD) Cl]₂, r.t. 3 hrs. (iii) AgBF₄

The catalytic activities towards reduction of benzophenone and nitroarenes showed that the ligand binding in a monodentate fashion compound **31** has higher catalytic activity compared to their respective chelation complexes **32**. However, in all the NHC iridium complexes described showed higher catalytic activities relative to when [Ir (COD) Cl]₂ were used as catalyst towards reduction of benzophenone and nitroarenes.

The research interest in metal N- Heterocyclic carbene (NHC) is now expanding to the study of new versatile ligand topologies, which have shown promising spectator characteristics with classical functional groups [59]. One of these is the pincer architecture which in the case of pyridyl carbene ligands are mostly tridentate in nature. The pincer ligands provide a preorganised backbone capable of blocking pseudo-meridional coordination sites of metal, leaving the remaining available for catalysis [60]. In line with above principles Crabtree reported the CNC ligand **33** and its corresponding pincer complex of Pd (II) **34** through reaction with Pd (II)(OAc)₂. Complex **34** proved to be a robust catalyst for the Heck reaction at 165°C, showing the resistance of the complex to

thermal decomposition and in air [61]. The planar complex showed low solubility but using R= n-Bu wingtip gave sufficient solubility for convenient study. Introduction of methylene linker result in loss of planarity and greatly improved the solubility. The Ru pincer complexes of the mode CNC have also been reported by the same group (**35**, **36**). Catalytic tests showed **35** to be active in both hydrogen transfer and oxidation of olefins while **36** was inactive in both cases [62]. The same group reported CCC pincer ligand **37** and its conversion to Pd (II) complex **38** by using Pd (0) through oxidative addition cyclometallation. The CCC complex was more rigid than CNC pincer but still fluxional enough to give coalescence at elevated temperature. All the species however appeared to be catalytically active in the usual coupling reactions. The Crabtree pincer ligands and complexes are depicted in Scheme 3.





Scheme 3: Synthesis and example of some Crabtree pincer complexes

The Cavell group has also looked at related Pd (II) complexes (**39**, **40**, and **41**) [63 and 64]. Compound **39** which is a Pd-hydrocarbyl pincer complex was prepared using a one pot transmetalation technique via the $\text{Ag}^{\text{I}}(\text{NHC})$ complexes [64]. The complexes reported have shown good activity in a model Heck coupling reaction using activated aryl bromides with the complexes bearing bulkier N- substituents outperforming the N-Me substituent in the case of complexes **40** analogues.






Figure 1.20: Cavell pincer complexes reported.

The Danopoulos group has also reported analogous carbene complexes as part of his contribution to the investigation of the behaviour of these noble metal complexes. Compounds (**42**, **43** and **44**) are Pd (II) carbene complexes and their tests in catalysis have showed good stability and activity in the Heck reaction [65]. The same group also reported the synthesis of complexes **45** and **46**. Complex **45** was accessed by the reaction of the corresponding imidazolium salt with $[\text{Fe}(\text{N}(\text{SiMe}_3)_2)]$. Complex **46** is the first reported dinitrogen complex


stabilised by NHC [66]. The same synthetic methodology was applied for the synthesis of complex **47** by the reaction of $\text{Co}[\text{N}(\text{SiMe}_3)_2]_2$ with a bis imidazolium salt which on further reaction with $\text{Na}(\text{Hg})$ and MeLi gave complexes **48** and **49**. Also reported in the literature by the same group is complex **50** which was prepared by the reaction of 2,6-[(*o*-dialkyl)phenylimidazolylidene]pyridine with $\text{RuCl}_2(\text{PPh}_3)_3$ in THF. Catalytic tests of complex **50** in hydrogenation of $\text{C}=\text{O}$ and $\text{C}=\text{N}$ groups by hydrogen transfer from isopropyl alcohol in the presence of $\text{KO}^\text{i}^\text{Pr}$ or $\text{KO}^\text{t}^\text{Bu}$ showed remarkable activity [67], though the reactions are slow at room temperature but proceed at good rates at 55°C or 80°C .






42: Ar = mesityl




43



44

45: Ar = 2, 6-*i*Pr₂C₆H₃

47

 $\xrightarrow{\text{Na}(\text{Hg}) \text{ or } \text{MeLi}}$


48 / 49





Figure 1.21 Examples of some Danopoulos Pincer carbene complexes

In 2006 Zeng and Yu reported the synthesis of pyridyl-supported pyrazolyl- N-Heterocyclic carbene ligands (**51**, **52**) and their corresponding palladium complexes (**53**, **54**) and tested in Suzuki-Miyaura reactions [68]. A methylene linker was introduced in ligand **52** to release the steric strain encountered on **51**. All the palladium complexes exhibited good to excellent catalytic activity in Suzuki-Miyaura reaction of phenyl or p-tolylboric acid with aryl halide including iodobenzene, aryl bromides, and activated aryl chlorides under mild conditions, revealing that the new ligands are promising for the construction of highly active transition metal catalysts. The ligands and the complexes are depicted in figure 1.22.







Figure 1.22 Pyridyl Supported pyrazolyl NHC Ligands


A series of other pyridyl functionalised pincer carbene complexes were reported by Gibson and tested in oligomerisation and polymerisation of ethylene [69] as shown in Scheme 4 below. However, of all the reported complexes (Fe, V, Cr, Ti, Co), only Fe complexes showed evidence of alkyl carbene coupling signifying the importance of the early transition metal complexes in this chemistry [69, 70].



Scheme 4: Examples of Gibson pincer NHCs

Pincer PCP complex **55** [71] and tetradentate pyridine functional ligand **56** and its corresponding Pd(II) **57** and Ni(II) **58** [72] have also appeared in the literature. Compound **55** as can be seen contains three strong σ -donors and its use in Heck and Suzuki reaction was found to be effective. In contrast to carbene complexes with mono, bi-, and tridentate ligands widely used in C-C coupling reactions, tetradentate ligands are rarely employed because of restriction on the available coordination sites for the incoming substrates, though the more robust nature of metal complexes with tetradentate ligands can

provide extra stability to the catalytic species. Few reports exist on the use of NHC nickel as catalyst in Suzuki coupling [73]. However preliminary application of the tetradentate nickel complex $[\text{NiL}]^{2+} \cdot 2\text{Br}^-$ **58** in Suzuki coupling of aryl halides with phenylboronic acid has shown effective catalytic activities including aryl chlorides as substrates. This is particularly important as nickel is cheaper than palladium, thus the use nickel catalyst will give access to large-scale of inexpensive compounds.



Scheme 5: Examples of PCP and Tetradentate complexes

1.8 Carbene complexes

Carbene was first introduced into organometallics chemistry in 1964 [74] by Fischer. Fischer complexes exhibited σ - donor/ $\pi\pi$ -acceptor behaviour for the bound carbene, and the metal to carbon bonds were shorter than the usual single bond [75]. Following this discovery it became evidently clear that there were

two distinct types of carbene complexes at that time. Fisher carbene complexes combine weakly donating singlet carbene, which accepts back bonding from low-valent metal [1, 75, 76,], while the already known Schrock carbene complexes combine a covalent triplet carbene and triplet metal fragment. In contrast to NHCs, these carbenes generally contain alkyl substituents and therefore are nucleophilic and coordinate to high oxidation state metals.

With the isolation of free carbene by Arduengo, there were renewed interests in the study of nucleophilic carbene complexes. It was initially thought that Arduengo NHCs carbene would yield Fisher type carbene complexes upon coordination to a metal centre, but the bonding properties showed different characteristics. Due to the back donation from the adjacent nitrogen heteroatoms and their strong capacity as σ -donors to metals, NHCs form only a single σ -bond to metals with negligible π -back donation [1, 76], and therefore these complexes exhibit different reaction chemistry to either Fischer or Schrock carbene complexes.


N-heterocyclic carbenes (NHCs) have attracted much attention because their transition metal complexes display rich coordination chemistry and have wide applicability in catalysis [1]. Recently research efforts have been devoted to the synthesis of polydentate ligands containing NHC moieties. The combination of pyridine and NHC functionalities leads to diverse polydentate ligands, some of which have shown interesting coordination chemistry [54, 77,78, 79], efficient catalytic applications [53,80,81,] and biological activities [82]. The basis of this study was to develop new ligand structure, by substituting pyridine with quinoline substituents, which are expected to provide greater rigidity and hence stability, though these rigid structures also lead to significant steric over crowding. Hopefully, this controlled flexibility/steric crowding parameters will lead to improved catalytic performance in a range of reactions.

1.8.1 Synthesis of N- heterocyclic carbene transition metal complex

What is to be done with the transition metal carbene complexes is probably more important than the complexes themselves and since carbene complexes have found utilization in many industries, scientists have been working to find simple methods to access these very important compounds. A number of routes have been developed, allowing the preparation of complexes bearing carbene ligands with a large variety of electronic and steric properties [83]. This has mainly been achieved as a result of the straight forward methods of synthesis of a range of imidazolium salts thereby allowing for the design of carbene ligands with a variety of electronic and steric properties, ideal for tailoring the properties of the desired complex as catalyst. Of the many synthetic methodologies available in the literature for the preparation of NHC metal complexes, four are more prominent : (i) In- situ deprotonation of azolium salts (ii) complexes via free carbenes (iii) Ligand transfer reactions (iv) Oxidative addition reactions.


1.8.1.1 In- situ deprotonation of azolium salts

This is the most widely used method of accessing carbene complexes. In this method, the isolation of the free carbene is not necessary. In his original work, Ofele formed NHCs by in situ deprotonation of the corresponding imidazolium salts. The basic metalate ion $[\text{HCr}(\text{CO})_5]^-$ serves as base as well as ligand acceptor [84].




Scheme 6: Carbene complexes through basic metalate anion deprotonation

Basic counter ions of the metal precursors can also act as deprotonating agents. For example, a convenient method to synthesise NHC-Pd (II) is by mixing Pd(OAc)₂ with the corresponding imidazolium salt. In a similar way, μ -alkoxo complexes of (η^2 -cod) rhodium(I) or iridium(I), formed in situ by adding μ -chloro bridged analogues to a solution of sodium alkoxide in the corresponding alcohol, will deprotonate an imidazolium salt to form the corresponding NHC complex[85]. Despite being relatively simple method, the imidazolium counterion is generally incorporated into the nascent carbene complexes unless non-coordinating anions are used. Good yields require the use of a solvent, such as THF or DMSO, however solvent free reactions have been reported [26, 86].



Scheme 7: Carbene complexes by basic ligand deprotonation

The use of an external base to generate NHCs in the presence of the metal precursor is also an efficient way of accessing carbene complexes. Popular external bases include potassium and lithium tert-butoxide [87, 88], sodium hydride [89], butyl lithium [90, 91], triethylamine [92, 93] and KN(SiMe₃)₂ [94]




Scheme 8: Synthesis of Carbene Complexes via external base deprotonation


Another method that allows the synthesis of carbene complexes is through the elimination of molecules of methanol and chloroform from the diazaortho-ester [95, 96] and trichloromethyl-substituted relatives of imidazolium salts [97, 98]. Thermal elimination of these two substituents gives free carbene which upon reaction by a suitable metal precursor form the appropriate carbene complex.

1.8.1.2 Carbene Complexes through free carbenes

Following the isolation of 1, 3-diamantylimidazol-2-ylidene by Arduengo, a wide range of new carbene complexes could be synthesised by carefully using the appropriate metal precursor complex. The most popular methods of synthesising free carbenes is by the use of strong bases such as sodium hydride and potassium tert-butoxide in THF [16, 99], or mixture of THF and liquid ammonia [26].



Scheme 9: Synthesis of free carbene




Scheme 10: Carbene complex formation from free carbenes and dimeric cleavage and phosphines exchange

NHCs are very strong σ donors and show dissociation energies higher than phosphines for a large number of metals. Therefore, when their free carbene can be isolated, their complexation is achieved in high yield. Free NHCs have been found to be able to cleave dimeric species such as $[(\eta^4\text{-cod})\text{RhCl}]_2$ [26] and exchange phosphines [38] and pyridine [100] ligands as depicted above in Scheme 10

1.8.1.3 Carbene complexes through transmetallation

Some NHCs are difficult or not possible to synthesise via the free routes especially in a situation where the NHC precursor contains an acidic proton in its linker chain. Gratifyingly it was discovered that Chromium, molybdenum and tungsten complexes could be used for carbene transfer to a variety of metals including rhodium(I), Palladium(II), copper(I), Platinum(II), Silver(I) and gold(I) [101-103]. Recently it has been found that transmetallation reactions using silver carbene have been reported for a wide variety of transition metals: Au(I), Cu(I), Cu(II), Ni(II), Pd(II), Pt(II), Rh(I), Rh(III), Ir(I), Ir(III), Ru(II), Ru(III), and Ru(IV) [104]. The Ag(I) NHC complexes are simply prepared by deprotonation of the imidazolium salt with Ag_2O in a suitable solvent and transmetallation reaction is usually conducted in DMSO or DCM [64].




Scheme 11: Carbene complex through silver transfer reactions

1.8.1.4 Carbene complexes through oxidative addition reactions

C-H oxidative addition of an imidazolium salt is another effective alternative method to obtain carbene complexes. The group of Lappert [105] and Stone et al

used oxidative addition method in the 1970's for creating thiazol-2-ylidene complexes from 2-chlorothiazolium salts [106,107]. Cavell group [108, 109] used a similar method to carry out oxidative addition of imidazolium salts to Ni^0 , Pd^0 and Pt^0 and recently the method was utilised by Peris et al [110] to synthesize Ir(III) carbene complexes through direct reaction of pyridyl functionalised imidazolium salt with $[\text{IrCl}(\text{cod})]_2$ as shown in Scheme 11 below.



Scheme 12: carbene complexes through oxidative addition

Though accessing carbene complexes via this method is generally restricted to nickel, palladium, platinum, rhodium and iridium, these are the commonly used metal in catalysis.

1.9 Abnormal carbene complexes

So far all the carbene complexes reported are those in which the coordination took place at the C(2) position of the NHCs. However, in 2001, Crabtree discovered an expected binding mode of NHCs. Instead of having coordination at the usual C(2) position of the NHC, the metal was attached at C(4) or C(5) [111] as shown in figure 1.23.

After this discovery a lot of other publications of NHC have appeared in the literature [112-114].




Normal binding at C(2)

Abnormal binding at C(4) or C(5)

Figure 1.23 C(2) and C(4) or C(5) binding mode of the NHCs.

The abnormal carbene complex was initially observed by mixing pyridine-substituted imidazolium salts with $[\text{IrH}_5(\text{PPh}_3)_2]$ in refluxing benzene. From theoretical calculation [115], binding through C(4) or C(5) positions is less favoured. It was therefore reasoned that steric effects of the bidentate pyridine-NHC around the metal centre and selection of imidazolium salt counter ion controlled the reaction [115]. With many catalytic reactions involving carbene being prepared in situ, care should be taken when designing reactions as slight changes in reaction condition can affect the properties of the catalyst and the overall reaction. The study of the abnormal carbene complexes is still going on.



1.2.0 Catalysis involving NHCs

1.2.1 Ruthenium metathesis

The trial of Schrock and Fischer type carbene complexes in catalytic reactions showed that they had the tendency to suffer from M-C cleavage thereby making them catalytically inactive [75]. However, NHCs form stable bond with metals and can accommodate a wide range of oxidation states, making them suitable for

many catalytic transformations. Because of σ -donor ability and their strong metal-carbon bond, NHC ligands have been applied as directing ligands in various catalytic transformations [59]. It is however in ruthenium catalysed olefin metathesis type reactions that NHC ligands have proved their efficiency giving access to unprecedented successful catalytic systems.




Figure 1.24 NHCs in ruthenium metathesis

In his catalysis investigations Herrmann showed that having one imidazolin-2-ylidene in place of a phosphine **60** favours the dissociative substitution of the phosphine ligand with olefinic substrate, giving rise to a more active species [42, 116]. Catalyst **60** was found to have good activities in ring opening metathesis of 1, 5-cyclooctadiene. In a similar fashion Grubbs [117] synthesised and tested catalyst **61** containing more basic NHC and results showed excellent activity in ring opening metathesis. The use of imidazolidin-2-ylidene allowed access to more catalysts by introducing chirality at the C(4) and C(5) positions of the NHC. Towards this end complex **62** was made and its application in desymmetrisation of triolefins yielded the ring closing metathesis products in high enantioselectivities [118]

1.2.2 Asymmetric catalysis

Following the success of the use of chiral carbenes in asymmetric catalysis in 1996/1997 by Enders [119] and Herrmann [129], chemists have pursued this area leading to many publications on the use of NHCs for asymmetric homogeneous catalysis [121]. Enders applied the NHC and their derivatives in catalysed asymmetric nucleophilic acylation processes with remarkable success.

Chiral NHC ligands have found applications in the following catalytic processes: Rh-hydrosilylation of ketones [120, 122, 123], olefin metathesis [118, 124], Rh(I) and Ir(I)-transfer hydrogenation of ketones [125].

1.2.3 Hydrogenation

In his pioneering work Nolan used achiral monodentate NHC iridium complex **63** for the hydrogenation of cyclohexene and 1-methyl cyclohexene (Figure 1.24). Catalyst **63** and Crabtree's catalyst **64** were found to have comparable activity at room temperature [126]. However, catalyst **63** was found to be more robust and efficient at higher temperature probably due to the stability of metal-carbene bond relative to metal-phosphine bond.




Figure 1.25 Achiral monodentate NHC ligand and Crabtree's catalyst

In another investigation, Buriak discovered that combining NHC with phosphine ligands led to efficient systems for the hydrogenation of simple olefins [127]. Comparing complex **65** with its analogue **66** in hydrogenation of 1-methylcyclohexene and 2, 3-dimethyl-2-butene showed the superiority of catalyst **65** in activity. While complex **65** fully hydrogenated 2, 3-dimethyl-2-butene in less than an hour at 1 bar H₂ at room temperature, complex **66** gave 19% conversion in 4 hours under the same conditions.



Figure 1.26 Achiral monodentate NHC phosphine and NHC pyridine iridium complexes

There also exist in the literature some reports on the use of chiral NHCs carbene complexes in asymmetric iridium-catalysed hydrogenation. In particular Burgess chiral bidentate oxazoline-NHC ligand **67** gave high enantioselectivities for a range of olefins with best results obtained using phosphine-oxazoline (PHOX) **68** and its derivatives **69** [128, 129].




Figure 1.27 Burgess's bidentate oxazoline-NHC ligand, PHOX ligands and its derivatives

NHC carbene complexes are also known to catalyse a wide range of reactions such as hydroformylation [129], hydrosilylation [26, 42, 88, 115, 132,], olefin metathesis [38, 44, 130, 133-135, 136], and polymerisation of alkynes [137] and C-C coupling reactions including Suzuki, Stille and Heck reactions [53, 80, 139, 140].

1.2.4 Catalyst decomposition

NHCs complexes show remarkable stability in many catalytic reactions are often stable to heat, moisture and oxygen. However, McGuinness et al discovered that the carbene catalyst decomposed during the catalytic reaction giving unsatisfactory results [115]. Further investigation indicated that the decomposition was as a result of reductive elimination of *cis* located carbene and alkyl or acyl ligands [141, 142] (Figure 1.28)




Figure 1.28 Decomposition pathways in carbon monoxide ethylene copolymerisation

It is believed that the reaction is assisted by twist of the carbene with respect to the square planar Pd(II) centre by approximately 60° so that the empty p orbital on the carbene centre is directed towards the alkyl/ acyl group adjacent to it on the metal centre. Since the acyl/ carbene intermediates are necessary intermediates in the CO/ ethylene catalytic cycle, the discovery of the decomposition was quite disturbing, and no reports of success has appeared in the literature of carbene complex catalysis of this reaction since then.

1.2.5 Aims and overview of the thesis

Carbene based ligand systems with functionalised pyridine groups have proved very effective as ligands for catalysis and there has been considerable work on these types of system. The task of this thesis was to develop a new variation on this ligand structure, involving quinoline substituents, which are expected to provide greater rigidity and stability to the complexes. Also as part of this work, the catalytic applications of interesting reactions was explored, primarily the iridium complexes. Ligands of this nature were largely unknown and this opens up an opportunity to develop a whole range of systems with a cross section of properties.

This thesis is composed of chapters which contains introductions, and reviews of relevant literatures to date.

Chapter 2 describes the preparation and characterisation of a range of quinoline functionalised imidazolium salts and a tetrahydropyrimidium salt that were required as precursors to the respective NHC ligands.

Chapter 3 deals with the synthesis and characterisation of a range $\text{Ag}^{\text{I}}(\text{NHC})$ complexes and few $\text{Pd}^{\text{II}}(\text{NHC})$ complexes accessed by transmetallation of the corresponding silver carbenes. Some of the $\text{Ag}^{\text{I}}(\text{NHC})$ complexes were found to be biscarbenes and the nature of the functional group was found to be of no influence to the type of complexes formed.

Chapter 4 presents the synthesis and characterisation of a range $\text{Rh}^{\text{I}}(\text{NHC})$ and $\text{Ir}^{\text{I}}(\text{NHC})$ complexes by transmetallation of $\text{Ag}^{\text{I}}(\text{NHC})$ complexes, with $\text{Rh}^{\text{I}}(\text{NHC})$ complexes giving higher yields. Attempt to prepare the chelated version was not successful as the complexes were found to insoluble in less polar solvents and decomposed in high polar solvent like DMSO. It also covers the catalytic testing of some the $\text{Ir}^{\text{I}}(\text{NHC})$ complexes prepare in reduction of 4-bromoacetophenone by hydrogen transfer reactions. The results show that all the complexes were very active giving a conversion of up to 100% with as little as 0.01 mole % of the catalysts.

Chapter 5 deals with conclusions and further work to be carried out as there a lot of research opportunities in this area.

1.2.6 References

1. D. Bourissou, O. Guerret., F. Gabbai, G. Bertrand, *Chem. Rev.*, **2000**,100,39-91.
2. G.Schuster, *Advanced physical Organic Chemistry*, **1986**, 22, 311.
3. R.Hoffmann. *J. Am.Chem.Soc.*, **1968**, 90, 1475.
4. L. Pauling, *Chem. Commun.* **1980**, 688.
5. R. W. Alder, P. R. Allen, M. Murray, A. G. Orpen, *Angew. Chem. Int. Ed.* **1996**, 35, 1121.
6. H. W. Wanzlick, E. Schikora, *Angew. Chem.* **1960**, 72,494.

7. R. W. Alder, M. E. Blake, L. Chaker, J. N. Harvey, F. Paolini, J. Schuetz, *Angew. Chem. Int. Ed.* **2004**, *43* 5896.
8. W. V. E. Doering, A. K. Hoffmann, *J. Am. Chem. Soc.* **1954**, *76*, 6162.
9. H. Wanzlick, *Angew. Chem. Int. Ed. Engl.*, **1962**, *1*, 75-80.
10. H. Wanzlick, B. Lachmann, E. Schikora, *Chem. Ber.*, **1965**, *98*, 3170-3177.
11. V. Bohn, W. Herrmann, *Angew. Chem. Int. Ed. Engl.*, **2000**, *39*, 4036-4086
12. F. E. Hahn, L. Wittenbecher, D. L. Van, R. Froehlich., *Angew. Chem. Int. Ed.* **2000**, *39*, 3, 541.
13. D. Lemal, Y. Liu, P. Lindner, *J. Am. Chem. Soc.*, **1999**, *121*, 10626.
14. H. Tomoika, *Acc. Chem. Res.* **1997**, *30*, 315
15. A. Igau, H. Grutzmacher, A. Baceiredo, G. Bertrand, *J. Am. Chem. Soc.* **1988**, *110*, 6463.
16. A. J. Arduengo, III, R. L. Harlow, M. Kline, *J. Am. Chem. Soc.* **1991**, *113*, 361.
17. H. J. Schoenherr, H. W. Wanzlick, *Chem. Ber.* **1970**, *103*, 1037.
18. A. J. Arduengo, III, J. R. Marshall, *J. Am. Chem. Soc.* **1995**, *117*, 11027.
19. R. W. Alder, P. R. Allen, M. Murray, A. G. Orpen, *Angew. Chem. Int. Ed.* **1996**, *35*, 1121.
20. E. Despagnet-Ayoub, R. H. Grubbs, *J. Am. Chem. Soc.* **2004**, *126*, 10198.
21. V. Lavallo, J. Mafhouz, Y. Canac, B. Donnadiou, W. Schoeller Wolfgang, G. Bertrand, *J. Am. Chem. Soc.* **2004**, *126*, 8670.
22. D. Ender, Breur, G., J. Runsink, J. Teles, J. Melder, K. Ebel, S. Brode, *Angew. Chem. Int. Ed.*, **1995**, *34*, 9, 1021-1023.
23. A. Arduengo, J. Goerlich, W. Marshall, *Liebigs Ann.*, **1997**, 365-374.
24. D. Martin, A. Baceiredo, H. Gornitzka, W. Schoeller, G. Bertrand, *Angew. Chem. Int. Ed.*, **2005**, *44*, 11, 1700-1203.
25. (a) Herrmann, W. A., Ellison, M., Fisher, Kocher, C., Artus, G. R. *J. Chem. Eur. J.* **1996**, *2*, 772. (b) Herrmann, W. A., Kocher, C., Goossen, L. J. Artus, G. R. *J. Chem. Eur. J.* **1996**, *2*, 1627.

26. F. E. Hahn, L. Wittenbecher, D. L. Van, R. Froehlich, *Angew. Chem.* **2000**, *112*, 551; *Angew. Chem. Int. Ed.* **2000**, *39*, 541.
27. R. W. Alder, M. E. Blake, J. M. Oliva, *J. Phy. Chem.* **1999**, *103*, 11200.
28. R. W. Alder, in *Carbene Chemistry*, **2002**, pp. 153.
29. Y. J. Kim, A. Streitweiser, *J. Am. Chem. Soc.* **2002**, *124*, 5757.
30. N. Kuhn, T. Kratz, *Synthesis*, **1993**, 561.
31. R.H.Crabtree, *Platinum metals rev.*, **2007**, *51*, (4), 176-177.
32. P.Arnold, S. Mungur, A. Blake, C. Wilson, *Angew Chem.Int.Ed.* **2003**, *42*, 5981-5984.
33. A. Arduengo, M. Tam, J. Calabrese, F. Davidson, W. Marshall. *Chem. Lett.*, **1999**, 1021-1022.
34. W. Herrmann. O. Runte, G. Artus. *J. Organomet. Chem.*, **1995**, *501*, C1-C4.
35. A. Arduengo, M. Tam, S. McClain, J. Calabrese, F. Davidson, W. Marshall *J. Am. Chem. Soc.* **1994**, *116*, 7927.
36. W.J. Oldman, S. M. Oldman, B.L. Scott, K. D. Abney, W. H. Smith, D. A. Costa, *Chem. Comm.*, **2001**, *15*, 1348.
37. P. Arnold, *Heteroatom. Chem.*, **2002**, *6*, 534.
38. T. Weskamp, W. C. Schattenmann, M. Spiegler, W. A. Herrmann, *Angew.Chem., Int. Ed.* **1998**, *38*, 2490.
39. C. Koecher, W. A. Herrmann, *J. Organomet. Chem.* **1997**, *532*, 261.
40. W. A. Herrmann, T.Weskamp, V. P. W. Bohm, *Adv. Organomet. Chem.* **2001**, *48*, 1.
41. A.R.Chianese, X. Li. M.C.Jansen, J.W.Faller, R. H. Crabtree, *Organometallics*, **2003**, *22*, 1663.
42. T. Weskamp, F.J.Kohl, W.Hieringer, D. Gleich W. A. Herrmann, *Angew.Chem., Int. Ed.* **1999**, *38*, 2416.
43. J. Huang, H. -J Schanz E. D. Stevens, S. P. Nolan, *Organometallics*, **1999**, *18*, 2370.
44. J. Huang, E. D. Stevens, S. P. Nolan, J. L. Petersen, *J. Am. Chem. Soc.* **1999**, *121*, 2674.

45. J. Schwarz, V. P. W. Bohm, M. G. Gardiner, M. Grosche, W. A. Herrmann, W. Hieringer, G. Raudaschl-Sieber, *Chem. Eur. J.* **2000**, 1773.
46. C. Boehme, G. Frenking, *Organometallics* **1998**, 17, 5801.
47. J.C. Green, R. G. Scurr, P.L. Arnold, F. G. N. Cloke, *Chem. Comm.* **1997**, 1963.
48. Y. Ma., W. Jiang, Q. Wu, Y. Wang, X. Liu, M.B. Andrus. *Org. Lett.* **2003**, 5, 3317-3319.
49. T.J. Seiders, D.W. Ward, R.H. Grubbs, *Org Lett* **2001**, 3, 3225-3228.
50. J.J. Vanveldhuizen, S. B. Garber, J. S. Kingsbury, A.H. *J. Am. Chem. Soc.* **2002**, 124, 4954-4955.
51. H. Dias, W. Jin, *Tetrahedron Let.* **1994**, 35, 1365.
52. (a) F. Naud and P. Braustein, *Angew. Chem., Int. Ed.* **2001**, 40, 680; (b) C. S. Slone, D.A. Weinberger and C. A. Mirkin, *prog. Inorg. Chem.*, **1999**, 48, 233.
53. D.S. McGuinness and K.J. Cavell, *Organometallics*, **2000**, 19, 741.
54. S. Grundemann, M. Albrecht, A. Kovacevic, J. W. Faller and R. H. Crabtree, *J. Chem. Soc., Dalton Trans.*, **2002**, 2163.
55. A.A.D. Tulloch, S. Winston, A.A. Danopoulos, G. Eastham and M. B. Hursthouse, *J. Chem. Soc., Dalton Trans.*, **2003**, 699-708.
56. J. Chen, I. Lin, *Organometallics*, **2000**, 19, 5113-5121.
57. J. Chen, I. Lee, *J. Chem. Soc., Dalton Trans.*, **2000**, 839.
58. C. Wang, C. Fu, Y. Liu, S. Peng and S. Liu. *Inorg. Chem.* **2007**, 46, 5779-5786.
59. W. A. Herrmann, *Angew. Chem., Int. Ed.* **2002**, 41, 1290.
60. M. Albrecht and G. van Koten, *Angew. Chem. Int. Ed.*, **2001**, 40, 3750.
61. R. H. Crabtree, *Pure Appl., Chem.*, **2003**, 75, 4, 435-443.
62. M. Poyatos, J. Mata, E. Falommo. R. H. Crabtree. E. Peris. *Organometallics*, **2003**, 22, 1100-1114.
63. D.S. McGuinness, K.J. Cavell, B. F. Yates, B. W. Skelton, *J. Am. Chem. Soc.* 123, 8317, **2001**.
64. D. J. Nielsen, K.J. Cavell, B. W. Skelton, A. H. White, *Inorganic Chemica Acta*, **2006**, 359, 1855-1869.

65. M. B. Andrus, C. Song, *Org. Lett.*, **2001**, *3*, 3761.
66. A. Danopoulos, N. Tsoureas, J. Wright, M. Wright, *J. Am. Chem. Soc.*, **2004**, *23*(2) 166-168.
67. A. A. Danopoulos, S. Winston and W. B. Motherwell, *Chem. Commun.*, **2002**, 1376-1377.
67. F. Zeng, Z. Yu, *J. Org. Chem.* **2006**, *71*, 5274-5281.
68. D. McGuinness, V. Gibson, D. Wass, J. Stead, *J. Am. Chem. Soc.*, **2003**, *125*, 12716-12717.
69. D. McGuinness, V. Gibson, D. Wass, J. Stead, *Organometallics*, **2004**, *23*, 6288-6292.
70. H. Lee, J. Zeng, C. Hu, M. Lee, *Inorg. Chem.*, **2004**, *43*, 6822-6829.
71. P. L. Chiu, C. Lai, C. Chang, C. Hu, H. M. Lee, *Organometallics* **2005**, *24*, 6169-6178.
- 72.(a) G. Steiner, A. Krajeta, H. Kopacta, K. H. Ongania, K. Wurst, P. Preishuber-Pfluegl, B. Bildstein, *Eur. J. Inorg. Chem.*, **2004**, *14*, 2827.
(b) J. Liu, M. J. Robins, *Org. Lett.* **2004**, *6*, 4321.
73. E.O. Fisher and A. Maasbol, *Angew. Chem. Int. Ed. Engl.*, **1964**, 580.
74. W. A. Herrmann and C. Kocher, *Angew. Chem. Int. Ed. Engl.*, **1997**, *36*, 2162.
75. R. H. Crabtree, 'The Organometallic Chemistry of the Transition Metals', John Wiley & Sons, Inc, **1998**.
76. S. Winston, N. Stylianides, A. A. D. Tulloch, J. A. Wright, A. A. Danopoulos, *Polyhedron*, **2004**, *23*, 2813.
77. (a) V. J. Catalano, M. A. Malwitz, *Inorg. Chem.* **2003**, *42*, 5483. (b) V. J. Catalano, M. A. Malwitz, A. O. Etogo, *Inorg. Chem.* **2004**, *43*, 5714.
78. A. A. Danopoulos, A. A. D. Tulloch, S. Winston, G. Eastham, M. B. Hursthouse, *Dalton Trans*, **2003**, 1009.

79. A. M. Magill, D. S. McGuinness, K. J. Cavell, G. J. P. Britovsek, V. C. Gibson, A. J. P. White, D. J. Williams, A. H. White, B. W. J. Skelton, *Organomet. Chem.* **2001**, *546*, 617-618.
80. (a) D. J. Nielsen, K. J. Cavell, A. H. White, B. W. J. Skelton, *Inorg. Chem. Acta* **2002**, *327*, 116. (b) J. A. Loch, M. Albrecht, E. Peris, J. Mata, J. W. Faller, R. H. Crabtree, *Organometallics*, **2002**, *21*, 700.
81. A. Melaiye, R. S. Simons, A. Milsted, F. Pingitore, C. Wesdemiotis, C. A. Tessier, W. J. Young, *J. Med. Chem.* **2004**, *47*, 973.
82. T. Weskamp, V. P. W. Bohm and W. A. Herrmann, *J. Organometallic Chem.*, **2000**, *600*, 12.
83. K. Oefele, *J. Organomet. Chem.* **1968**, *12*, 42.
84. C. Kocher and W. A. Herrmann, *J. Organomet. Chem.* **1997**, *532*, 261.
85. W. A. Herrmann, J. Fisher, M. Elison, C. Kocher and G. R. J. Artus, *Angew. Chem. Int. Engl.*, **1995**, *34*, 2371.
86. D. Enders, H. Gielen, G. Raabe, J. Runsink and J. H. Teles, *Chem. Ber.*, **1996**, *1483*.
87. W. A. Herrmann, L. J. Gooben and M. Spiegler, *J. Organomet. Chem.* **1997**, *547*, 357.
88. K. Ofele, W. A. Herrmann, D. Milhalious, M. Elison, E. Herdtreck, W. Scherer and J. Mink, *J. Organomet. Chem.*, **1993**, *459*, 177.
89. W. P. Fehlhammer, T. Bliss, U. Kernbach and I. Brudgam, *J. Organomet. Chem.*, **1995**, *490*, 149.
90. U. Kernbach, M. Ramm, P. Luger and W. P. Fehlhammer, *Angew. Chem. Int. Ed. Engl.*, **1996**, *35*, 310.
91. D. Enders, H. Gielen, G. Raabe, J. Runsink and J. H. Teles, *Chem. Ber.*, **1997**, *130*, 1253.
92. D. Enders, H. Gielen, J. Runsink, K. Breuer, S. Brode and K. Boehn, *Eur. J. Inorg. Chem.*, **1998**, 913.
93. M. Iglesias, D. J. Beetstra, A. Stasch, P. N. Horton, M. B. Hursthouse, S. J. Cole, K. J. Cavell, A. Dervisi and I. A. Fallis, *Organometallics*, **2007**, *26*(19), 4800-4809.

94. D. Enders, K. Breuer, G. Raabe, J. Runsink, J.H. Teles, J. P. Melder, K. Ebel S. Brode, *Angew. Chem. Int. Ed. Engl.*, **1995**, *34*, 1021.
95. J.H. Teles, J. P. Melder, K. Ebel, R. Schneider, E. Gehrler, W. Harder, S. Brode, D. Enders, K. Breuer and G. Raabe, *Helv. Chim. Acta*, **1996**, *79*, 61.
96. D. M. Lemal and K. I. Kawano, *J. Am. Chem. Soc.*, **1962**, *84*, 1761.
97. W. Wanzling, F. Esser and H. J. Kleiner, *Chem. Ber.*, **1963**, *96*, 1208.
98. A.J. Arduengo III, H. V. R. Davis, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.* **1992**, *114*, 5530.
99. A. C. Hillier, H. M. Lee, E. D. Stevens, S. P. Nolan, *Organometallics* **2001**, *20*, 4246.
100. C.G. Kreiter, K. Ofele and G. W. Wieser, *Chem. Ber.*, **1976**, *109*, 1749.
101. S.T. Liu, T. Y. Hsoeh, G.H. Lee and S.-M. Peng, *Organometallics*, **1998**, *17*, 993.
102. R.Z. Ku, J. C. Huang, J. Y. Cho, F. M. Kiang, K. R. Reddy, Y. C. Chen, K. J. Lee, G. H. Lee, S. M. Peng and S.T. Liu, *Organometallics*,
103. J. C. Garrison and W. J. Youngs, *Chem. Rev.* **2005**, *105*, 3978-4008.
104. M. F. Lappert, A. J. Oliver, *J. Chem. Soc. Comm.*, **1972**, 274.
105. P. J. Frazer, W. R. Roper and F. G. A. Stone, *J. Chem. Soc. Dalton Trans*, **1974**, 102.
106. P. J. Frazer, W. R. Roper and F. G. A. Stone, *J. Chem. Soc. Dalton Trans*, **1974**, 760.
107. D. S. McGuinness, K. J. Cavell, B. F. Yates, B. W. Skelton and A. H. Whites, *J. Am. Chem. Soc.*, **2001**, *123*, 8317- 8328.
108. N. D. Clement, C. Jones and C. Elsevier, *Angew. Chem. Int. Ed. Engl.*, **2004**, *43*, 1277-1279.
109. E. Mas-Masza, M. Sanau and E. Peris, *Inorg. Chem.* **2005**, *44*, 9961-9967.
110. S. Gruedemann, A. Kovacevic, M. Albrecht, J. W. Faller, R.H. Crabtree, *Chem. Comm.* **2001**, 2274.
111. S. Gruedemann, A. Kovacevic, M. Albrecht, J. W. Faller, R.H. Crabtree, *J. Am. Chem. Soc.* **2002**, *124*, 10473.

112. A.R. Chianese, A. Kovacevic, B. M. Zeglis, J. W. Faller, R. H. Crabtree, *Organometallics* **2004**, *23*, 2461.
113. H. Lebel, M. K. Janes, A. B. Charette, S. P. Nolan, *J. Am. Chem. Soc.* **2004**, *126*, 5046.
114. G. Sini, O. Eisenstein, R. H. Crabtree, *Inorg. Chem.* **2002**, *41*, 602.
115. M. S. Sanford, J. A. Love, R. H. Grubbs, *J. Am. Chem. Soc.* **2001**, *123*, 6543.
116. M. Scholl, S. Ding, C. W. Lee, R. H. Grubbs, *Org. Lett.* **1999**, *1*, 953.
117. T. J. Seiders, D. W. Ward, R. H. Grubbs, *Org. Lett.* **2001**, *3*, 3225.
118. D. Enders, H. Gielen, K. Breuer, *Tetrahedron: Asymmetry* **1997**, *8*, 3571.
119. W. A. Herrmann, L. J. Goossen, C. Koecher, G. R. J. Artus, *Angew. Chem. Int. Ed.* **1996**, *35*, 2805.
120. V. Cesar, S. Bellemin-Lapponnaz, L. H. Gade, *Chem. Soc. Rev.* **2004**, *33*, 619.
121. L. H. Gade, V. Cesar, S. Bellemin-Lapponnaz, *Angew. Chem. Int. Ed.* **2004**, *43*, 1014.
122. W. -L. Duan, M. Shi, G. -B. Rong, *Chem. Comm.* **2003**, 2916.
123. A. H. Hoveyda, D. G. Gillingham, J. J. Van Vedhuizen, O. Kataoka, S. B. Garber, J. S. Kingsbury, J. P. A. Harrity, *Org. Boimol. Chem.*, **2004**, *2*, 8.
124. H. Seo, B. Y. Kim, J. H. Lee, H. -J. Park, S. U. Son, Y. K. Chung, *Organometallics*, **2003**, *22*, 4783.
125. H. M. Lee, T. Jiang, E. D. Stevens, S. P. Nolan, *Organometallics* **2001**, *12*, 1255.
126. L. D. Vazquez-Serrano, B. T. Owens, J. M. Buriak, *Chem. Comm.* **2002**, 2518.
127. A. Pfaltz, J. Blankenstein, R. Hilgraf, E. Hormann, S. McIntyre, F. Menges, M. Scholeber, S. P. Smidt, B. Wustenberg, N. Zimmermann, *Adv. Synth. Catal.* **2003**, *345*, 33.
128. F. Menges, A. Pfaltz, *Adv. Synth. Catal.* **2002**, *344*, 40.
129. C. Kocher, Technische Universitat Munchen, **1997**.
130. M. F. Lappert and R. K. Maskell, *J. Organometallics Chem.* **1984**, *264*, 217.






131. L. J. Gooben. Technische Universitat Munchen, **1997**.
132. T. Weskamp, F. J. Kohl, W. Hieringer, and A. Herrmann, *Angew. Chem. Int. Ed. Engl.*, **1999**, *38*, 2416.
133. T. Weskamp, F. J. Kohl, and W. A. Herrmann, *J. Organometallics Chem.*, **1999**, *582*, 362.
134. L. Ackermann, A. Furstner, T. Weskamp, F. J. Kohl, and W. A. Herrmann, *Tetrahedron Lett.*, **1999**, *40*, 4787.
135. M. Scholl, T. M. Trnka, J. P. Morgan and R. H. Grubbs, *Tetrahedron Letters*, **1999**, *40*, 2247.
136. R. Z. Ku, D. Y. Chen, G. H. Lee, S. M. Peng and S. T. Liu, *Angew. Chem. Int. Ed. Engl.*, **1997**, *36*, 2631.
137. D. S. McGuinness, K. J. Cavell, B. W. Skelton, and A. H. White, *Organometallics*, **1999**, *18*, 572, 177.
138. T. Weskamp, V. P. W. Bohm and W. A. Herrmann, *J. Organometallics Chem.*, **1999**, 348.
139. C. Zhang and M. L. Trudell, *Tetrahedron Lett.* **2000**, *41*, 595.
140. D. S. McGuinness, K. J. Cavell and B. F. Yates, *Chem. Comm.*, **2001**, *4*, 355.
141. D. S. McGuinness and K. J. Cavell, *Organometallics*, **2000**, *19*, 4918.
142. D. S. McGuinness, N. Saendig, B. F. Yates and K. J. Cavell, *J. Am Chem. Soc.* **2001**, *123*, 4029.

CHAPTER TWO

Quinoline functionalised imidazolium and pyrimidinium salts

2.1 Introduction

Imidazolium salts can be defined as planar 5 five membered heterocycle with nitrogens at the 1 and 3 positions and a substituents (H, R, Ar, or X) at each position of the ring. The ring members are sp^2 hybridized and the ring bears a single positive charge that is delocalised around the ring [1]. In most cases the imidazole **2a** which does not have substituents on one of the nitrogens serves as a precursor to the many imidazolium salts possible.

**2a****2b****2c****2d****2e****2f**

The numbering presented in **2b** is used throughout this research work with $R_2 = R_4 = R_5 = H$ in all the imidazolium salts. The structures are named according to the degree of saturation in the heterocycle of the parent compound. As presented above the imidazole **2a** has two double bonds. The imidazolin-2-ylidene **2c** is derived from **2b** by loss of a proton. In contrast to **2b**, **2d** is a dihydroimidazolium salt and is fully saturated and this does not allow delocalisation of the charge beyond the NCN region, therefore, it has the effect of increasing the donor ability of the derived imidazolidin-2-ylidenes, **2e** [2]. All


but one of the imidazolium salts described in this thesis are those based on type **2b**. One type **2d** based imidazolium salt is also presented.

Structure **2f** is a typical example of pyrimidinium salt which is a six membered ring with nitrogens at positions 1 and 3. The two nitrogens and C-2 carbon are sp^2 hybridized while C4, C5 and C6 are sp^3 hybridized. The NHC derived from pyrimidinium would be expected to be more basic with high donor capability due to the absence of delocalisation of charge in the NCN region. Though as stated above the NHCs with an unsaturated backbone are generally more stable as free carbenes, since the normally empty pz orbital is part of an aromatic system, conjugated with the C-C double bond in the backbone. There is however little evidence that the aromaticity of an NHC has much bearing on its properties as a ligand for transition metals [3].

The ease of synthesis of imidazolium salts is one of the chief reasons for the popularity of NHCs. Other attractive features of imidazolium based NHCs are the wide variety of steric and asymmetric environments that are available through modification of the substituents on the nitrogen of the heterocycle. Furthermore, through the use of appropriate donor groups on the nitrogen substituents, it is possible to make multidentate NHC ligands [4]. Such variability makes possible the synthesis of numerous ligands. Along these lines Cavell [5], Crabtree [6] and Danopoulos groups [7] have reported wide variety of multidentate pyridyl functionalised ligands. Multidentate ligands especially those that can behave as a chelating ligand having both strong and weak donors (hemilabile ligands) are particularly important in catalysis. The weak hemilabile part of the ligand is capable of reversible dissociation from metal centre, thereby creating vacant coordinating sites during catalytic cycles and stabilising the metal centre by reCOORDINATING when it is catalytically inactive. In addition to the works so far reported on pyridyl ligands, we envisioned the synthesis of an analogue of the quinoline framework. It was our hope that replacing the relatively small pyridine with large quinoline substituents will provide greater rigidity and hence stability to the complexes, though, the rigid structures also lead to steric crowding. Bearing these in mind methylene linker was introduced in some of the imidazolium salts between the imidazole and the quinoline moiety to reduce the steric strain and improve solubility.

2.1.1 Ligand synthesis

Our initial intention was to synthesize a wide range of quinoline based imidazolium salts. However, while we were successful in the synthesis of symmetrically substituted saturated imidazolium salt, the unsymmetrical substituted analogues could not be accessed via traditional approaches [8].



Scheme 2.1: Traditional approach to the synthesis of symmetrical saturated imidazolium salts


Unsymmetrical substituted imidazolium salts have been successfully synthesized via nucleophilic attack of 1-alkylimidazole or 1-aryl imidazole on an alkyl halide [9-12] (Scheme 2.2). Through this direct quaternization, a lot of functional groups have been attached to the imidazolium moiety. Such functional groups include: hydroxyl group [13], carboxylic groups [14], thiol groups [15], alkyne and alkene groups [16, 17].



Scheme 2.2: Synthesis of unsymmetrical imidazolium via nucleophilic attack of imidazole on alkyl halide

However nucleophilic attack on an aryl ring by an imidazole is difficult, making N, N-diaryl substitution unattainable by this approach.

This is a set back because one of our aims was to create as much as possible some steric environment in our target NHCs precursors. Therefore, N, N'-diaryl substituted imidazolium salt with quinoline as one of the substituents was considered an ideal candidate. Fortunately, flexible approaches allowing for the synthesis of bulky N-substituted imidazoles are available in the literature [18, 19] (Scheme 2.3). Synthetic routes to organohalide compounds with donor functionalised groups are also available, though in most cases they can be obtained from commercial sources. Once the N- substituted imidazole is made, a wide range of imidazolium salts can be prepared by reaction with the appropriate alkyl halides as shown in Scheme 2.2 above.



Scheme 2.3: Synthesis of bulky N- substituted imidazoles

In summary, this chapter reports the syntheses of a wide variety of quinoline functionalised imidazolium salts giving access to bidentate and in some cases tridentate ligands. All the reported imidazolium salts were fully characterised by ¹H and ¹³C NMR spectroscopy, mass spectroscopy, micro analysis and in most cases were structurally characterised using single crystal X-ray crystallography.


2.2 Results and Discussion

2.2.1 Synthesis and characterisation of symmetrical saturated imidazolium imidazolium salt

Prior to that the start of this work there was no reported synthesis of symmetrical quinoline substituted imidazolium salts and several attempts towards accessing this compound were not successful by the established procedure [20]. However, in 2006 Michon et al almost at the same we synthesised our quinoline based salts reported the synthesis of analogue chiral tetradentate diamine and chiral dihydroimidazolium salts, [21]. In their reaction they utilised Buchwald-Hartwig Palladium –catalysed amination [22] involving 2.1 equivalents of 8-bromoquinoline, 3 equivalents of sodium t-butoxide, 5 mol% of Pd₂(dba)₃ and 10mol% of rac-BINAP in toluene at 80 °C under argon affording the chiral amine in 90% (Scheme 2.4). Ring cyclisation of the chiral diamine with triethylortho formate solution at 135°C afforded the desired dihydroimidazolium salts in 60-90% yield.

Previously this type of diamine was synthesised using a Bucherer reaction by refluxing 8-hydroxy quinoline, the desired diamine and sodium pyrosulfite in water for two weeks[23,24], but the yield obtained via this method was low by almost 50% compare to that reported by Michon [21]. This method was also used by T. Okada to make N,N-Di quinolyl-1,3-propanediamine [25].

In our study, we formed our diamine from the considerably cheaper 8-hydroxyquinoline material [23] and the achiral dihydroimidazolium **2** was prepared in 95% yield by reaction with triethyl orthoformate and ammonium tetrafluoroborate salt at 120°C in 2 hours as depicted in (Scheme 2.4) above. Compound **1** was characterised by ¹H and ¹³C NMR spectroscopy while imidazolium salt **2a** was characterised by ¹H, ¹³C NMR, mass spectroscopy and microanalysis. The salt is soluble only in high polar solvents such as DMSO or DMF. Of particular importance is the identification of CH unit between the N atoms which appeared at δ value of 11.4 which is higher than the figure reported by Michon [21]. The signal for the C2- carbon in ¹³C NMR occurs at a δ of 159.21 which is typical of such group [21] (161.90).



In order to carry out any further investigations with the salt there was a need to improve the solubility of the dihydroimidazolium salt. This was achieved by replacing the counter ion tetrafluoroborate with Barf giving access to the imidazolium salt **2b** that is soluble in almost all organic solvents with the exception of petroleum ether and hexane. The characteristic features of **2b** as observed in the ^1H and ^{13}C NMR spectra did not change significantly in relation to what was observed in **2a**. Diffusion of hexane into the DCM solution of compound **2b** gave crystals suitable for X-ray crystallographic determination, figure 2.1.




Figure 2.1: ORTEP projection of the cation of **2b.Barf**, excluding hydrogen atoms for clarity, showing labelling of atoms.

Table 2.1: Selected bond lengths (Å) and bond angles (°) of **2b**


| | | | | | |
|---------|-------|---------|-------|--------------|-------|
| N2 -C8 | 1.428 | N3- C13 | 1.476 | N2- C10 -N3 | 113.4 |
| N2-C10 | 1.315 | N4-C21 | 1.369 | N1- C9- C8 | 119.9 |
| N2- C11 | 1.481 | C8- C9 | 1.480 | N2 -C8- C7 | 119.3 |
| N3- C10 | 1.316 | C13-C21 | 1.420 | C10 -N2 -C8 | 127.5 |
| N3-C12 | 1.476 | N1-C9 | 1.363 | C10 -N3 -C13 | 126.4 |

The planes of the quinoline and imidazolium rings make angle of approximately 36.71° with $\text{imC}_2\text{-H}$ and the nitrogen on the quinoline rings on the same side. Some selected bond angles and bond length are presented in table 2.1 above. The N2-C10-N3 bond angle of 113.4° obtained is higher than that generally reported for imidazolium salts (108°) [31, 32], but there is no significant difference in bond lengths.

2.2.2 Synthesis and characterisation of bis quinoline pyrimidium salt

Reports on the synthesis of 1, 3- dimesityl and 1, 3-dialkyl tetra hydroypyrimidinium salts have appeared in the literature [26]. However, pyrimidium salt **4** is the first six membered ring that is functionalised with quinoline giving a tridentate ligand that may behave as a pincer ligand. The method of Okada was employed to prepare N, N-Di quinolyl-1, 3- propanediamine **3** by refluxing 1, 3 -diaminopropane, 8-hydroxyquinoline and sodium pyrosulfite in water for two weeks [23, 24].

The pyrimidium salt **4** was prepared by the reaction of the corresponding diamine **3** with triethyl orthoformate and ammonium hexafluorophosphate salt at 120°C in 3 hours as depicted in (Scheme 2.5) , giving a white powder in good yield. The synthesis of the salt was confirmed from the NMR data. The ^1H NMR showed a signal at 9.3ppm corresponding to $\text{C}_2\text{-H}$ proton and C_2 carbon was appeared at δ value of 156.72 from the ^{13}C NMR spectrum. The data from the mass spectrum and micro analysis were consistent with the proposed structure.



Scheme 2.5: Synthesis of quinoline functionalised pyrimidium salt

Diffusion of Et₂O into a MeCN of solution of **4** yielded crystals suitable for a single X-ray crystallographic determination, Figure 2.1

Table 2.2: Selected bond lengths (Å) and bond angles (°) of **4**

| | | | | | |
|---------|----------|---------|----------|-----------|-----------|
| C1 -N1 | 1.317(3) | N4-C22 | 1.372(3) | N1-C1-N2 | 124.2(3) |
| C1 -N2 | 1.317(3) | C5-C6 | 1.365(3) | C1-N1-C5 | 120.71(3) |
| N1 -C5 | 1.439(3) | C14-C15 | 1.371(3) | C1-N2-C14 | 120.32(3) |
| N2 -C14 | 1.429(3) | N4-C21 | 1.323(3) | N1-C5-C15 | 118.33(3) |
| C13 -N3 | 1.370(3) | | | N1-C2-C3 | 109.90(3) |

The molecular structure of the tetrahydropyrimidium salt **4** in the solid state is depicted in Fig 2.2 being the first structure of a tetrahydropyrimidium salt with nitrogens of the quinoline substituents on the same side with the imC₂-H. The quinoline and imidazolium rings of **4** make an angle of approximately 53.69° which shows a clear divergence from co-planarity. Search of the available literature revealed that X-ray structures of 1,3-disubstituted 3,4,5,6-tetrahydropyrimidinium salts have been reported only for a few symmetrical:

diisopropyl[33], diethyl [34] and dimesityl [35]. The data obtained for compound **4** is very similar to that of the corresponding diisopropyltetrahydropyrimidinium salt reported by Alder [33]. The N1-C1-N2 angle of $124.2(3)^\circ$ and C1-N2 distance of $1.317(3) \text{ \AA}$ are nearly identical to the ones published by Alder[33] and Herrmann[35] who reported $124.72(15)^\circ$ and $1.3147(14) \text{ \AA}$ respectively.





Figure 2.2: ORTEP projection of the cation of **4.PF₆** excluding hydrogen atoms for clarity, showing labelling of atoms.

2.2.3 Unsymmetrical substituted Quinoline imidazolium salt

Two quinoline based imidazolium salts are herein reported. 8-imidazol-1-yl-quinoline was synthesised following the Zhang modified procedure for the synthesis of 1-arylimidazole [27]. The yield obtained is low (~ 20%) which is a known problem in the literature with most 1-aromatic substituted imidazoles.




Scheme 2.6; Synthesis of quinoline imidazolium salt

The low yield may be connected with the generation of large amounts of unknown by-product during neutralisation. The use of large volume of diethyl ether and vigorous agitation during extraction slightly increased the yield. The use of small scale synthesis is generally more efficient, though the overall yields were not constant. The imidazolium salts were prepared following the standard N-alkylation using methyl iodide or benzyl bromide. The alkylation of the quinoline imidazole follows S_N2 behaviour and is difficult to achieve with nucleophile less reactive than a secondary alkyl bromide precluding access to desired bulky N-^tBu, N- Mes and N-dipp substituents on the resulting imidazolium salt via this route [2]. The quinoline imidazole was reacted with either methyl iodide or benzyl bromide in THF overnight to give the desired imidazolium salts (**6a**, and **6c**) as light brown solid in good yield. While imidazolium **6c** is stable towards air and moisture, compound **6a** is hygroscopic and the anion (bromide) was exchanged for tetrafluoroborate anion to obtain compound **6b**. The anion exchange was accomplished by mixing a solution of the halide salt in acetonitrile with a solution of an excess of sodium tetrafluoroborate in water which on work gave the BF₄ salt in good yield. The imidazolium salts were fully characterised. The characteristic peak in the ¹H NMR is the C₂-H imidazolium proton appearing as singlet between 10.15-10.75ppm. In the ¹³C NMR the C₂ appeared at a δ value of 150.70ppm. Quality crystals for X-ray crystallography were obtained for the BF₄ version of the imidazolium salt **6b** by vapour diffusion of Et₂O into DCM solution. In the ¹H NMR spectra of the BF₄ salt (CDCl₃ solvent) the C₂-H imidazolium proton was observed to move upfield while only little of such change could be observed in ¹³C NMR.

Table 2.3: Selected bond lengths (Å) and bond angles (°) of **6b**

| | | | | | |
|--------|------------|---------|----------|-------------|------------|
| N6-C7 | 1.391(2) | N9-C8 | 1.381(2) | N6-C10-N9 | 108.38(14) |
| N6-C10 | 1.3405(19) | C7-C8 | 1.349(2) | C10-N6-C18 | 127.40(13) |
| N6-C18 | 1.4411(19) | C12-C13 | 1.389(3) | N9-C11-C12 | 111.31(14) |
| N9-C10 | 1.324(2) | C19-N20 | 1.367(2) | C18-C19-N20 | 119.85(14) |
| N9-C11 | 1.4846(19) | N20-C21 | 1.315(2) | N6-C18-C27 | 118.31(14) |

**Figure 2.3:** ORTEP projection of the cation of **6b.BF₄** excluding hydrogen atoms for clarity showing atom labelling scheme.

The quinoline and imidazolium rings make an angle of approximately 36.16° and both the nitrogen of the quinoline ring and imC₂-H are directed on the same sides of the molecules. The crystal structure of **6b** reveals N6-C10-N9 bond angle of 108.38° and N6-C10, N9-C10 bond lengths of 1.3405(19) Å and 1.324(19) Å respectively which is within the expected values of imidazolium salts reported [31, 32]. Crystals suitable for X-ray chromatography was obtained for compound **6c** by diffusion of diethyl ether into the DCM solution of the compound and the crystal structure is depicted in figure 2.4. ¹H and ¹³NMR, MS, and micro analysis data correspond to the proposed structure.

Table 2.4: Selected bond lengths (Å) and bond angles (°) of **6c**

| | | | | | |
|-------|----------|--------|----------|------------|----------|
| Cl-C2 | 1.367(4) | C10-N2 | 1.340(4) | N2-C10-N3 | 108.9(2) |
| Cl-C9 | 1.442(4) | C10-N3 | 1.319(4) | C1-N2-C10 | 127.4(3) |
| C1-N2 | 1.439(4) | C13-N3 | 1.461(4) | C1-C9-N1 | 119.5(2) |
| C5-C9 | 1.423(4) | | | C9-N1-C8 | 116.8(2) |
| C9-N1 | 1.365(4) | | | C10-N3-C13 | 125.3(3) |

The planes of the quinoline and imidazolium rings of **6c** make an angle of approximately 37.34° and gave N2-C10-N3 bond angle of $108.9^\circ(2)$ and N2-C10, N3-C10 bond lengths of 1.340(4) and 1.319(4) Å respectively which is consistent with the values reported in the literature [31, 32].





Figure 2.4: ORTEP projection of the cation of **6c.I** excluding hydrogen atoms for clarity showing atom labelling scheme.

2.2.4 Synthesis and characterisation of methylene-bridged quinoline functionalised imidazolium salts

Synthesis of imidazolium salts with the quinoline methylene bridge was desired, as apart from reducing the steric strain, it also improves solubility. The methylene bridged quinoline imidazole was prepared in good yield following the procedure reported in the literature [28] by refluxing a mixture of imidazole, 2-chromethylquinoline monohydrochloride and KOH in THF for 2 days. The direct quarternization of the methylene bridged quinoline imidazole with appropriate alkyl halides gave the desired salts in good yield.



Scheme 2.7: Synthesis of methylene-bridged imidazolium salts

However two of the imidazolium salts reported in this work could not be obtained via this method because of the difficulty associated with synthesising imidazolium salts by alkylating imidazoles with aryl halides or tertiary alkyl halides. Therefore 2-methoxypropionophenone imidazole and mesityl imidazole were prepared according to the reported literature methods [29] and [30] respectively, which upon alkylation with 2-chromethylquinoline monohydrochloride in the presence of a base gave the desired imidazolium salts.

Imidazolium salts **9b**, **9c** and **9d** are stable to moisture and air while compounds **9a**, **9e** and **9f** are hygroscopic salts. The characteristic features confirming the synthesis of the salts are appearances of a singlet in ^1H NMR between 10.20-11.60 corresponding to the $\text{C}_2\text{-H}$ proton between the two nitrogens and the ^{13}C NMR spectra showed the value of the C_2 carbon between 152.25-153.60 ppm with the highest value of 153.60ppm being observed in compound **9b** and 152.25 ppm in **9c**. Crystals of compounds **9b** and **9c** suitable for X-ray structures were obtained by diffusion of Et_2O into the acetonitrile solutions of the salts. Crystals of compound **9f** suitable for X-ray crystallography were grown by layering hexane onto a DCM solution of the salt. In order to investigate the effect of counter ion on the crystal structure, the counter ion chloride in salt **9b** was exchanged for tetrafluoroborate and crystals suitable for X-ray crystallography of the corresponding salt were obtained by vapour diffusion of Et_2O into the DCM solution of the salt Figure 2.5. The nature of the counter ion, i.e coordinating and non-coordinating, may have little influence on the structure of the cation as the X-ray revealed almost identical data, though peaks in the ^1H NMR spectra of the BF_4 version are shifted slightly upfield from the values observed in **9b**. X-ray quality crystals were also obtained for salts **9c** and **9f** by diffusion of diethyl ether into the DCM solutions of the respective salts and the crystal structures are depicted in Figures 2.6 and 2.7 respectively.




Figure 2.5: ORTEP projection of the cation of **9b.BF₄** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 2.5: Selected bond lengths (Å) and bond angles (°) of **9.BF₄**

| | | | | | |
|--------|----------|----------|----------|------------|----------|
| Cl- N1 | 1.340(3) | C4-C9 | 1.402(3) | N1-C1-N2 | 108.2(2) |
| C1-N2 | 1.323(3) | C9-C12 | 1.500(3) | C1-N1-C4 | 126.2(2) |
| C2-N1 | 1.374(3) | C13-N2 | 1.463(3) | C1-N2-C13 | 125.3(2) |
| C3- N2 | 1.370(3) | C13- C14 | 1.506(3) | N1-C4-C9 | 117.5(2) |
| C4-N1 | 1.446(3) | C14-N3 | 1.306(3) | N2-C13-C14 | 112.9(2) |




Figure 2.6: ORTEP projection of the cation of **9c.I** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 2.6: Selected bond lengths (Å) and bond angles (°) of **9c.I**

| | | | | | |
|-------|-----------|----------|----------|----------|-----------|
| C1-N1 | 1.30(2) | C7-C8 | 1.4888 | N1-C1-N2 | 105.9(12) |
| C1-N2 | 1.369(2) | N3-C8 | 1.3899 | C1-N2-C7 | 129.5(13) |
| C7-N2 | 1.515(15) | C8-C9 | 1.3888 | C1-N1-C4 | 124.4(11) |
| C4-N1 | 1.50(2) | | | C6-C4-C5 | 109.7(14) |
| C4-C5 | 1.53(2) | N2-C7-C8 | 111.2(5) | N3-C8-C7 | 119.2 |




Figure 2.7: ORTEP projection of the cation of **9f.Cl** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 2.7: Selected bond lengths (Å) and bond angles (°) of **9f.Cl**

| | | | | | |
|--------|----------|----------|----------|------------|-----------|
| N2- C6 | 1.335(3) | C8-N9 | 1.316(3) | N2-C6-N5 | 108.3(2) |
| N2-C18 | 1.467(3) | C19-N20 | 1.323(3) | C6-N5-C7 | 125.0(2) |
| N5-C6 | 1.327(3) | N9-C10 | 1.371(3) | C6-N2-C18 | 124.7(2) |
| N5-C7 | 1.461(3) | | | N2-C18-C19 | 113.0(2) |
| C7-C8 | 1.511(3) | C7-C8-N9 | 117.6(2) | N5-C7-C8 | 112.07(2) |

The imidazolium salts **9b**, **9c** and **9f** for which single X-ray structure determinations were performed show relatively consistent parameters in terms of bond distances and internal angles around the imidazolium rings and are all within the data reported in the literature [31, 32]. Important bond angles and bond distances are tabulated in tables 2.5, 2.6 and 2.7. In **9b** the mesityl and imidazolium rings planes are almost perpendicular to each making an angle of approximately 88.69° and the imC₂-H and the nitrogen on the quinoline ring are disposed to the same side of the molecule. The planes of the methylquinoline


and imidazolium ring of **9c** make an angle of approximately 61.83° while those of **9f** are almost perpendicular to each other forming an angle of approximately 85.65° and the imC₂-H and nitrogen on the quinoline moiety are disposed on the side of the molecules in the both compounds (**9c** and **9f**).

2.2.4 Synthesis and characterisation of octahydroacridine based imidazolium salt

In addition to the quinoline based imidazolium salts, it was decided to synthesise an octahydroacridine based imidazolium salt. Octahydroacridine can offer a secondary donor function for chelation as well as sp³ hybridized carbons which offers the ability to make the ligand chiral as well as addition of extra steric strain if required. Octahydroacridine was synthesised following the procedures of Paine [29] and Bell [30]. Bell's method offers fewer synthetic steps as well less forcing conditions but the yield as stated in the paper (50%) could only be achieved by strict adherence to conditions and deviation from the procedure led to the formation of large quantities of by-products.

Although Paine's method requires the use of many steps and forcing reaction conditions, it was found to be more reliable giving access to cleaner products.

Paine's synthetic procedure as well as steps leading to the synthesis of the desired imidazolium salt is depicted in Scheme 2.8 above. To introduce the chlorine onto position 4 it was necessary to functionalise the octahydroacridine ring with hydroxyl group. Paine's procedure involves the reaction of octahydroacridine with 3-chloroperoxybenzoic to form octahydroacridine N-oxide which upon reaction with an excess of boiling acetic anhydride gave 4-hydroxyl substituted octahydroacridine **14** after work up. However, in our work Fontena's method was employed [31] which requires the use trifluoroacetic anhydride instead of large excess of acetic anhydride with the reaction being carried out at room temperature to form the desired 4-hydroxyoctahydroacridine. 4-Chlorooctahydroacridine **15** was prepared by the reaction of 4-hydroxyoctahydroacridine with thionyl chloride which upon work up gave the desired compound as a yellow solid.



Scheme 2.8: Synthesis of octahydroacridine based imidazolium salt

Finally the acridine based salt was prepared by reaction of 4-chlorooctahydroacridine with 1-mesitylimidazole in THF in a pressure tube at 90 °C for 14 days. The yield obtained in this reaction was very low because the reaction follows a typical $\text{S}_{\text{N}}2$ pathway, and with secondary alkylchloride, the reaction would be expected to be very slow even under the forcing conditions used. Attempt to prepare a more reactive 4-iodooctahydroacridine was made by the addition of sodium iodide to an acetone solution of 4-chlorooctahydroacridine and stirred overnight and after work up there was no observable effect from the ^1H NMR spectra. The imidazolium salt prepared was fully characterised by ^1H NMR, mass spectroscopy and micro analysis with the

^1H NMR showing C2-H appearing as singlet at a δ value of 10.15ppm. The imidazolium salt **16** returned satisfactory MS and elemental analysis results.

Due to low yield and time constraints large varieties of the acridine based imidazolium could not be prepared using the secondary alkyl chloride **14**. However, it should be noted that, there is a lot of research opportunities in this area that need to be explored. It is envisaged that once a more reactive halo-octahydroacridine such as bromo or iodo-octahydroacridine can be prepared and following the procedure of Steiner et al [29] 1-acridine imidazole can be obtained thereby opening the way to a range of acridine based imidazolium ligands. Some of the interesting acridine based ligands that can possibly be prepared are presented in figure 1.8 below. It will be interesting to compare these potential pincer ligands with the pincer ligands reported by Gibson, Cavell, Danopoulos and Crabtree.









Figure 1.8: Proposed acridine based pincer ligands







2.3 Conclusions

A range of quinoline based imidazolium salts have been synthesised and characterised as precursors to the corresponding NHC ligands. A range of N-substituents give variable steric bulk to the imidazolium rings. The summary of all the imidazolium salts synthesized in this work are presented below in Table 2.8.

Table 2.8: Ligands synthesized in this project

| S. NO | Ligand | Structure |
|-------|-----------|---------------------------------------------------------------------------------------------------------------------|
| 1 | 2b |  Barf |
| 2 | 4 |  PF ₆ ⁻ |
| 3 | 6a |  Br ⁻ |
| 4 | 6c |  I ⁻ |
| 5 | 9a |  Cl ⁻ |

Chapter Two: Quinoline functionalised imidazolium and pyrimidinium salts

| | | |
|----|-----------|-------------------------------------------------------------------------------------|
| 6 | 9b |  |
| 7 | 9c |  |
| 8 | 9d |  |
| 9 | 9e |  |
| 10 | 9f |  |
| 11 | 16 |  |

The previously unreported bis 1, 3-(quinoline) tetrahydropyrimidinium salt **4** is the first reported quinoline based salt of this nature. Though in 2006 Michon et al reported chiral bis 1,3-quinolinedihydroimidazolium salts [21], salt **2b** is an achiral analogue and therefore is a new compound. Compounds **6a**, **6c** and **16** have earlier been prepared by Cavell group while all the other ligands are new.

Similar ligands functionalised with pyridine have been reported in the literatures [7, 31, 32, and 40]. However in this study, the pyridine moiety has been replaced by quinoline and this change is expected to increase the steric strain of the ligands as well as rigidity which may have important implications in the synthesis of the metal complexes. The acridine based imidazolium salt offers the possibility of accessing chiral ligand as well as varying the steric strain. Synthesis of the acridine based imidazolium salt also open up the opportunity to look into the possibility of synthesising the ligands presented in figure 1.8 above.

2.4 Experimental

2.4.1 General comments

Unless otherwise stated all manipulations were carried out using standard Schlenk techniques under an atmosphere of dry argon or in an MBRAUN M72 glove box (N_2 atmosphere with > 1 ppm O_2 and H_2O). Glassware were dried overnight in an oven at $120^\circ C$ or flame dried prior to use. Tetrahydrofuran (THF), diethyl ether (Et_2O), and hexane were dried and freshly distilled before used. Dichloromethane (DCM), methanol (MeOH) and acetonitrile (MeCN) were dried over calcium hydride. All other anhydrous solvents were obtained by distillation from the appropriate drying agent under dinitrogen. Deoxygenation of solvents and reagents was carried out by freeze- thaw- degassing.


All NMR solvents were purchased from Aldrich and Goss, dried over 3\AA molecular sieves and freeze-thaw degassed three times. All reagents were purchased from commercial sources and used without purification, unless otherwise stated.

All NMR data are quoted δ /ppm. 1H and ^{13}C spectra were recorded on a Bruker 400 MHz DPX Avance, unless otherwise stated, and referenced to $SiMe_4$. Electrospray mass spectrometry (ESMS) was performed on a VG Fisons Platform II instrument by the department of Chemistry, Cardiff University. Micro analysis was performed by Warwick Analytical Service.


2.4.2 N-Substituted imidazoles

The following imidazoles were prepared by established literature methods. 1-(2-methylpropiophenone)imidazole [29], mesitylimidazole [19], imidazole-1-yl-quinoline [27] and 1-(2methylquinoline) imidazole [28].


Imidazol-1-yl-quinoline (5):



8-aminoquinoline (2.27 g, 8.8 mmol) was mixed with 40% glyoxal (1.22 g, 8.8 mmol) in 30 mL MeOH and stirred overnight at room temperature to give a yellow mixture. NH_4Cl (0.94 g, 17.6 mmol) and 37% aq formaldehyde (1.42 g, 17.6 mmol) were then added to the mixture which was then diluted by further addition of 150 ml MeOH. This was refluxed for an hour and H_3PO_4 (1.6 mL, 85%) was slowly added to the mixture before heating to reflux for 12 hours. After removal of the solvent the dark residue was poured on to ice (100g) and neutralised with aqueous 40% KOH until pH 9. The resulting mixture was then extracted with diethyl ether (4 x 100 mL). The ethereal phase was then washed with water, brine and dried with Na_2SO_4 . The solvent was then removed to give a light brown solid product, 0.19 g (10%). ^1H NMR (CDCl_3 400 MHz 298K) : 8.90 (s, 1H, NCHN-H), 8.2 (m, 1H, $J = 6.7 \text{ Hz}$, quin-H), 8.05 (s, 1H, CHHC), 7.8 (m, 1H, $J = 7.9 \text{ Hz}$, quin-H), 7.6 (m, 1H, $J = 1.3 \text{ Hz}$, quin-H), 7.5 (m, 1H, $J = 7.9 \text{ Hz}$), 7.4 (m, 2H, $J = 4.2 \text{ Hz}$, quin-H), 7.2 (s, 1H, HCCH-H). ^{13}C NMR (CDCl_3 100.61 MHz 298K): 159.0, 140.5, 138.5, 136.0, 135.5, 128.5, 127.0, 126.8, 125.0, 123.0, 122.5, 120.5.

1-(2-methylpropiophenone) imidazole (8b).

A round bottomed flask was charged with imidazole (1.70g, 25mmol), 2-bromo-2-methylpropiophenone (2mL, 11.9mmol) and 40mL of ethanol. The yellow solution was refluxed for 3 days. Aqueous work up yielded yellow oil which crystallized in the freezer. Recrystallization from from dichloromethane/n-hexane (1:2) afforded pure colourless product (1.56g, 7.25mmol, 61%). ¹H NMR (CDCl₃ 400 MHz R.T.): 7.60(s, 1H, NCHN), 7.4(m, 1H, J = 2.5 Hz, arom-H), 7.2(m, J = 1.1 Hz, 4H, arom-H), 7.05(s, 1H, CHHC), 6.85(s, 1H, CHHC), 1.8(s, 6H, CH₃). ¹³C NMR (CDCl₃ 72.5MHz R.T): 198.45(PhCOC), 134.78,134.48, 130.17, 128.60, 128.50, 117.39, 64.89, 27.14(CH₃).


1-(2methylquinoline)imidazole (7):

A mixture of imidazole (0.38g, 5.51mmol), 2-(chloromethyl) quinoline and KOH (1.24g, 0.022mol) in THF (20mL) was refluxed for 2 days. The solvent was removed completely under reduced pressure. DCM (20mL) and water (20mL) were added and shaken vigorously in a separatory funnel. The organic layer was extracted and washed thoroughly with 20mL of water. The organic layer was separated and dried with anhydrous MgSO₄. The solution was filtered and removal of the solvent under reduced pressure gave an orange solid as the desired product (0.80g, 69%). ¹H NMR (CDCl₃ 400 MHz 298K): 8.1(d, 1H, J = 8.5 Hz, quin-H), 8.0(d, 1H, J = 8.5 Hz, quin-H), 7.75(d, 1H, J = 8.2 Hz, quin-H), 7.65(t, 1H, J = 5.6 Hz, quin-H), 7.6(s, 1H, NCHN), 7.45(t, 1H, J = 8.6 Hz, quin-H), 7.05(s, 1H, CHHC), 7.0(s, 1H, CHHC), 6.95(d, 1H, J = 10.0 Hz, quin-H), 5.35(s, 2H, NCH₂) ¹³C NMR (CDCl₃ 100.61MHz R.T): 156.13 (NCN),

147.71, 137.80, 137.60, 130.14, 129.18, 127.64, 127.41, 126.93, 119.58, 118.66, 53.25 (NCH₂)


2.4.3 Symmetrical saturated bis(1,3-quinoline) imidazolium salt

N, N'-Diquinolinethane -1, 2-diamine (1):



A mixture of 8-hydroxyquinoline (36.25 g, 0.25 mol), 1,2-diaminethane (9.75 g, 0.125 mol), sodium pyrosulfite (47.5 g, 0.25 mol) and water (250mL) was refluxed and stirred for one week. The solution was made strongly alkaline, cooled and filtered. The solid product was extracted several times with hot 0.2N NaOH until removal of 8-quinolinolate was complete and the residue was recrystallised from ethanol to give the desired product as a light yellow solid (6.00g 16%). ¹H NMR (CDCl₃, 400MHz 298K): 8.6(m, 2H, J = 2.6 Hz, Quin-H), 8.0(m, 2H, J = 6.8 Hz, Quin-H), 7.3(m, 4H, J = 7.7 Hz, Quin-H), 7.0(m, 2H, J = 7.5 Hz, Quin-H), 6.7(m, 2H, J = 8.0 Hz, Quin-H), 6.35(b, 2H, CNH), 3.7(s, 4H, NCH₂), ¹³C NMR (CDCl₃ 72.5MHz R.T): 146.94, 144.68, 138.34, 136.01, 128.73, 127.77, 121.46, 114.23, 104.77(quin-C), 42.79(NCCN).

Synthesis of 1, 3-diquinolin-4, 5-dihydroimidazolium tetrafluoroborate (2):




N, N'-diquinolinethane-1, 2-diamine (0.3g 0.58mmol) and NH₄BF₄ (0.095g, 0.58mmol) in triethyl orthoformate were heated at 120°C for 24 hours. The precipitate which was isolated was washed several times with diethyl ether and recrystallised from CHCl₃/Diethyl ether to afford the desired product 0.3g (85%). Anal. Calcd. for C₂₁H₁₇N₄BF₄: C, 61.19; H, 4.13; N, 13.60; F, 18.46%.

Found: C, 59.06; H, 4.01; N, 13.48; F, 19.04%. HRMS: Calculated for $C_{21}H_{17}N_4BF_4$: 325.1453. Found: 325.1437. 1H NMR (DMSO- d_6 , 400MHz R.T.): 11.40(s, 1H, NCHN), 9.20(m, 2H, $J = 2.6$ Hz, Quin-H), 8.70(m, 2H, $J = 1.2$ Hz, Quin-H), 8.20(m, 2H, $J = 8.2$ Hz, Quin-H), 8.10(m, 2H, $J = 8.0$ Hz, Quin-H), 7.90(m, 2H, $J = 8.0$ Hz, Quin-H), 7.8(m, 2H, $J = 4.2$ Hz, Quin-H), 4.95(s, 4H, NCH_2CH_2N). ^{13}C NMR (DMSO- d_6 , 72.5MHz R.T.): 158.91(NCN), 150.95(quin-C), 139.98(quin-C), 137.35(quin-C), 132.52(quin-C), 128.97(quin-C), 127.87(quin-C), 126.60(quin-C), 122.81(quin-C), 49.99(im-C).


2.4.4 Bis(1,3-quinoline)pyrimidinium hexafluoro phosphate salt

N, N-diquinolinepropane-1, 3-diamine (3):



Following the procedure for preparation of **1**, 7.32g of compound **3** was obtained from 8-hydroxyquinoline (18.13g, 0.125moles), 1,3-diaminopropane (4.64g, 0.125moles), sodiumpyrosulfite (23.77g, .125moles) and 125mL of water as a yellow solid. 1H NMR ($CDCl_3$, 400MHz 298K): 8.60(m, 2H, $J = 2.7$ Hz, quin-H), 8.00(m, 2H, $J = 6.7$ Hz, quin-H), 7.30(m, 4H, $J = 4.4$ Hz, quin-H), 7.00(m, 2H, $J = 8.3$ Hz, quin-H), 6.65(m, 2H, $J = 7.7$ Hz, quin-H), 6.20(broad, 2H, CNH), 3.50(m, 4H, $J = 7.0$ Hz, NCH_2), 2.20(m, 2H, $J = 6.8$ Hz, CCH_2C). ^{13}C NMR ($CDCl_3$ 72.5MHz R.T.): 146.84, 144.80, 138.25, 136.04, 128.70, 127.84, 121.41, 113.85, 104.72(quin-C), 41.30(NCC), 29.01(CCH_2C).


1, 3-diquinoline-3, 4, 5, 6-tetrahydropyrimidinium hexafluoro phosphate (4):



Following the procedure for the synthesis of **2**, the desired pyrimidinium salt **4** was synthesised from N,N-diquinolinepropane-1,3-diamine(3.59g,0.011 moles) NH_4PF_6 (1.78g, 0.011 moles) and triethyl orthoformate(30mL) as a light brown solid (Yield= 5.00g, 94.34%). Anal. Calcd. for $\text{C}_{22}\text{H}_{19}\text{N}_4\text{PF}_6$: C, 54.35; H, 3.93; N, 11.57%. Found: 53.74; H, 3.84; N, 11.37%. HRMS: Calculated for $\text{C}_{22}\text{H}_{19}\text{N}_4\text{PF}_6$: 339.1610. Found: 339.1618. ^1H NMR (DMSO- d_6 , 400MHz 298K): 9.30 (s, 1H, NCHN), 9.20(m, 2H, J = 4.2 Hz, quin-H), 8.70(m, 2H, J = 4.1 Hz, quin-H), 8.30(m, 2H, J = 8.3 Hz, quin-H), 8.25(m, 2H, J = 7.4 Hz, quin-H), 7.90(m, 2H, J = 7.8 Hz, quin-H), 7.80(m, 2H, J = 4.1 Hz, quin-H), 4.30(m, 4H, J = 5.4 Hz, NCH_2C), 2.70(m, 2H, J = 5.1 Hz, CH_2C). ^{13}C NMR (DMSO- d_6 , 72.5MHz R.T.): 156.72 (NCN), 151.57, 141.82, 138.10, 136.93, 129.64, 128.97, 126.65, 126.51, 122.72, 48.03, 19.54.

2.4.5 Unsymmetrical substituted quinoline imidazolium salts


Synthesis of 1-benzyl-3-quinolinimidazolium bromide (6a):



Benzyl bromide (1.41g, 8.25 mmol) was added to a solution 8-imidazol-1-yl-quinoline (0.70 g, 3.59 mmol) in 20 mL THF. The resulting mixture was allowed to stir overnight at r.t. The resulting yellow/brown precipitate was filtered using filter stick and washed with fresh THF to afford a brown powder. This was then recrystallised from DCM/Hexane to give the desired imidazolium salt (light brown solid). ^1H NMR(CDCl_3 :250MHz 298K):10.80(s, 1H, NCHN), 8.90(m, 1H, J = 4.1 Hz, Quin-H), 8.25(m, 1H, J = 8.7 Hz, Quin-H), 8.00(m, 1H, J = 8.3 Hz, Quin-H), 7.90(s, 1H, CHHC), 7.70(m, 1H, J = 5.7 Hz, Quin-H), 7.60(m, 2H, J = 6.2 Hz, Quin-H, CHHC), 7.50(broad 2H, Quin-H), 7.30(m, 3H, J = 7.3 Hz, Ar-H), 5.85(s, 2H, CH_2). ^{13}C NMR (CDCl_3 100MHz R.T.): 152.07(NCN), 140.79, 137.30, 137.10, 133.60, 131.40, 131.03, 129.86, 129.75, 129.70, 129.55, 126.90, 126.00, 124.72, 123.23, 121.95, 53.76(NCH₂).

1-benzyl-3-quinolinimidazoliumtetrafluoroborate (6b): The counter ion bromide in **6a** was exchanged for tetrafluoroborate ion by mixing one equivalent of 1-benzyl-3-quinolineimidazolium bromide (0.5g, 1.75 mmol) in acetonitrile with 1.5equivalent of NaBF₄(0.28g, 2.63mmol) in water. The acetonitrile was removed under reduced pressure and product was washed twice with water. The residue was then dissolved in DCM and the organic and aqueous layer separated. The DCM solution was dried over MgSO₄ and solution concentrated under reduced pressure. Addition of Et₂O precipitates out the product which was filtered and dried in vacuum to give the desired product as brown solid. Crystals suitable for X-ray crystallography were obtained by vapour diffusion of Et₂O into DCM solution of the product. Anal. Calcd. for C₁₉H₁₆N₃BF₄: C, 61.29; H, 4.29; N, 11.27%. Found: C, 60.96; H, 4.31; N, 11.12%. ¹H NMR (CDCl₃, 250MHz R.T.): 9.40(s, 1H, NCHN), 8.85(d, 1H, J = 4.2 Hz, quin-H), 8.35(d, 1H, J = 7.5 Hz, quin-H), 8.20(d, 1H, J = 8.4 Hz, quin-H), 8.00(d, 1H, J = 8.3 Hz, quin-H), 7.85(s, 1H, CHHC), 7.70(t, 1H, J = 4.2 Hz, quin-H), 7.50(m, 3H, arom-H), 7.60(s, 1H, CHHC), 7.50(m, 3H, J = 7.0 Hz, arom and quin-H), 5.50(s, 2H, CH₂). ¹³C NMR (CDCl₃ 72.5MHz R.T): 150.70(NCN), 139.31, 135.92, 135.69, 131.97, 129.96, 129.68, 128.46, 128.41, 128.26, 128.16, 125.42, 124.21, 123.35, 121.85, 120.77, 52.68(NCH₂).

1-methyl-3-quinolineimidazolium iodide (6c):




Methyl iodide (1g, 7mmol) was added to a solution of 8-imidazol-1-yl-quinoline (0.5g, 2.7mmol) in 20ml of THF. The resulting mixture was stirred at room temperature for 48 hrs. The resulting brown precipitate was filtered using filter stick and washed with further amount of THF to afford a brown powder. This was then recrystallised from DCM/Hexane to give the desired imidazolium salt (0.6g, 70%). Crystals suitable for X-ray were obtained by vapour diffusion of Et₂O into the DCM solution of the compound. Anal. Calcd. for C₁₃H₁₂N₃I: C, 46.30; H, 3.56; N, 12.47%. Found: C, 46.09; H, 3.60; N, 12.16%. ¹H NMR(DMSO-d₆:250MHz R.T): 9.80(s, 1H, NCHN), 8.90(d, 1H, J = 1.7 Hz, quin-H), 8.30(d, 2H, J = 1.6 Hz, quin-H), 8.00(d, 1H, J = 7.4 Hz, quin-H), 7.80(s,


1H, CHHC), 7.70(d, 1H, J = 7.9 Hz, quin-H), 7.50(m, 1H, J = 4.2 Hz, quin-H), 7.40 (s, 1H, CHHC), 4.30(s, 3H, CH₃). ¹³C NMR (DMSO-d₆, 72.5MHz R.T.): 152.02(NCN), 140.62, 138.48, 136.95, 131.39, 130.69, 128.77, 126.38, 126.27, 124.41, 123.29, 123.06, 67.00(CH₃)

2.4.6 methylene-bridged quinoline Functionalised imidazolium salts


1-methyl -3-(2-methylquinoline) imidazolium chloride 9a:



A mixture of methyl imidazole (0.58g, 7mmol), 2-chloromethyl quinoline monohydrochloride (1.5g, 7.00 mmol) and K₂CO₃ (0.50g, 3.50 mmol) in acetonitrile was refluxed for 24 hours. The solvent was removed and the remaining residue was dissolved in dichloromethane. The solution was filtered to remove the KCl formed and unreacted K₂CO₃ and the filtrate were concentrated to few millilitres. Addition of THF to the solution precipitated out the product which was repeatedly to give the product as a pale yellow solid (1.20g, 81%). Required: For C₁₃H₁₄N₃Cl: C, 60.12; H, 5.41; N, 15.80; Cl, 13.68%. Found: C, 64.14; H, 5.42; N, 16.00; Cl, 13.93%.HRMS: Anal.Calcd. for C₁₃H₁₄N₃Cl: 224.1188. Found: 224.1193. ¹H NMR (CDCl₃, 400MHZ, R.T.) : 10.67 (s,1H, NCHN), 8.10 (m, 1H, J = 8.4 Hz, Quin-H), 7.90 (d, 1H, J = 8.4 Hz, Quin-H), 7.65 (m, 3H, J = 8.1 Hz Quin-H, CHHC), 7.60 (s, 1H, CHHC), 7.40 (t, 1H, J = 7.1 Hz, Quin-H), 5.90 (s, 2H, CH₂), 4.00 (s, 3H, CH₃). ¹³C NMR (CDCl₃, 100.61MHZ, R.T): 153.14 (NCN), 147.86, 138.32, 138.14, 130.47, 129.06, 127.92, 127.57, 123.61, 123.25, 120.71, 54.57(NCH₂), 36.92(CH₃).

Synthesis of 1-mesityl 3-(2-methylquinoline)imidazolium chloride 9b :


A mixture of mesityl imidazole (0.87g, 4.67mmol), 2-chloromethylquinoline monohydrogen chloride (1.0g, 4.67mmol) and K_2CO_3 (0.33g, 2.39mmol) in acetonitrile was refluxed for 24 hours. The solvent was removed and the residue dissolved in dichloromethane. The solution was filtered to remove the KCl formed and unreacted K_2CO_3 and the filtrate were concentrated to few millilitres. Addition of THF to the solution precipitated the product which was repeatedly to give a white product (1.25g, 73.5%). Anal. Calcd. for $C_{22}H_{22}N_3Cl$: C, 72.63; H, 6.05; N, 11.55; Cl, 9.77%. Found: C, 69.41; H, 6.03; N, 10.98; Cl, 9.57%. HRMS: Calculated for $C_{22}H_{22}N_3Cl$: 328.1814. Found: 328.1807. 1H NMR ($CDCl_3$, 400MHz, R.T.) : 10.50 (s, 1H, NCHN) , 8.20 (m, 1H, J = 8.4 Hz, Quin-H), 8.00 (s, 1H, aromatic-H), 7.85 (m, 2H, J = 8.4 Hz, Quin-H), 7.75 (d, 1H, J = 8.1 Hz, Quin-H), 7.65 (t, 1H, J = 7.0 Hz, Quin-H) , 7.50 (t, 1H, J = 7.1 Hz, Quin-H), 7.05 (s, 1H, aromatic-H) , 6.95 (s, 2H, CHHC), 6.30 (s, 2H, CH_2), 2.30 (s, 3H, P- CH_3), 2.00 (s, 6H, O- CH_3). ^{13}C NMR ($CDCl_3$, 100MHz, 298K): 153.60, 147.77, 141.56, 139.17, 138.36, 134.71, 131.17, 130.48, 130.14, 129.16, 128.23, 128.15, 127.53, 124.43, 123.04, 121.18, 54.57, 21.47, 17.94.

Synthesis of 1-isopropyl -3-(2-methylquinoline) imidazolium iodide (9c)


A mixture 2-iodopropane (1.63g, 9.57mmol) and 1-(2-methylquinolin) imidazole (1g, 4.8mmol) in 50ml of ethyl acetate was refluxed for 3 days. The reaction solution was allowed to cool to room temperature, after which the crystals were filtered off and washed with ethyl acetate and dried in vacuum to give the desired product, (1.25g, 68.68%). Anal. Calcd. for $C_{16}H_{18}N_3I$: C, 50.67; H, 4.75; N, 11.08; I, 33.49%. Found: C, 50.17; H, 4.64; N, 10.89; I, 33.13%. HRMS: Calculated for $C_{16}H_{18}N_3I$: 252.1501. Found: 252.1491. 1H

NMR (CDCL₃, 400MHz, R.T): 10.35 (s, 1H, NCHN), 8.15 (d, 1H, J = 8.4 Hz, quin-H), 7.90 (d, 1H, J = 3.5 Hz, quin-H), 7.90 (t, 2H, J = 8.4 Hz, quin-H), 7.80 (m, 2H, quin-H), 7.70 (m, 2H, J = 6.5 Hz, CHHC, Quin-H), 7.50 (m, 2H, J = 9.0 Hz, quin-H, CHHC), 6.90 (s, 2H, NCH₂C), 4.80 (m, 1H, J = 7.8 Hz, NCHC₂), 1.60 (d, 6H, J = 7.0 Hz, CH₃). ¹³C NMR (CDCL₃, 100MHz, 298K): 152.25 (NCN), 147.93, 138.43, 136.07, 130.63, 129.45, 128.25, 128.15, 127.74, 123.63, 121.28, 120.29, 54.72, 23.59.


1-n-butyl-3-(2-methylquinolin) imidazolium iodide (9d):



A mixture of 1-(2-methylquinoline) imidazole (1.00g, 4.80mmol) and 1-iodobutane (1.76g, 9.60mmol) was refluxed in 50mL of ethyl acetate for two days. The reaction solution was allowed to cool to room temperature and the ethyl acetate decanted off. The residue was dissolved in minimum amount of DCM and the product precipitated out by the addition of hexane. The desired product was filtered and dried in vacuum. (1.20g, 63.83%). MS (ESMS, Da): M/Z 266.16 [M-I]⁺ (100%). Anal. Calcd. for C₁₇H₂₀N₃I: C, 51.92; H, 5.09; N, 10.69; I, 32.30%. Found: C, 52.58; H, 4.99; N, 10.57; I, 31.94%. ¹H NMR (CDCL₃, 400MHz, R.T): 10.2(s, 1H, NCHN), 8.15(d, 1H, J = 8.4 Hz, quin-H), 7.90(d, 1H, J = 6.6 Hz, quin-H), 7.85(m, 2H, J = 8.3 Hz, quin-H, CHHC), 7.65(t, 2H, J = 5.9 Hz, quin-H), 7.50(t, 1H, J = 7.4 Hz, quin-H), 7.40(s, 1H, CHHC), 5.90(s, 2H, NCH₂), 4.20(t, 2H, J = 7.2 Hz, NCH), 1.85(m, 2H, J = 7.5 Hz, CH₂), 1.30(m, 2H, J = 7.4 Hz, CH₂), 0.90(t, 3H, J = 7.1 Hz, CH₃). ¹³C NMR (CDCL₃, 100MHz, 298K): 152.78(NCN), 147.92, 138.29, 137.02, 130.58, 129.44, 128.21, 128.07, 127.68, 123.64, 122.36, 120.93, 54.65(NCC), 50.43, 32.41, 19.79, 13.88(CH₃).

1-(2-methylpropio-phenone)-3-(2-methylquinolin) imidazolium chloride (9e)

2-(chloromethylquinoline)monohydrochloride(1.50g,7.0mmol),1(2methylpropio-phenone) imidazole (1.50g, 7.0mmol) and K_2CO_3 (0.55g, 3.98mmol) in 30mL of acetonitrile was refluxed for 2 days. The solvent was removed and the residue dissolved in dichloromethane. The solution was then filtered to remove KCl and unreacted K_2CO_3 and the dichloromethane completely removed. The residue was repeatedly washed with diethyl ether and dried in a vacuum to afford the desired product. MS (ESMS): M/Z 366.16 $[M-Cl]^+$ (100%). 1H NMR ($CDCl_3$, 400MHz, 298K): 11.60(s, 1H, NCHN), 8.10(d, 1H, $J = 8.40$ Hz, quin-H), 7.90(m, 2H, $J = 3.6$ Hz, quin-H, aromatic-H), 7.80(d, 1H, $J = 8.3$ Hz, quin-H), 7.65(t, 1H, $J = 1.5$ Hz, Quin-H), 7.60(s, 1H, CHHC), 7.50 -7.40(m, 4H, $J = 7.7$ Hz, arom-H), 7.20(m, 2H, $J = 1.8$ Hz, quin-H), 7.25(s, 1H, CHHC), 7.00(s, 1H, CHHC). ^{13}C NMR ($CDCl_3$, 100MHz, R.T): 197.30(CO), 153.45(NCN), 147.91, 138.33, 134.06, 133.75, 130.44, 129.43, 129.26, 129.08, 128.83, 128.22, 128.13, 127.62, 123.78, 121.40, 120.87, 70.08, 54.79, 27.31(CH_3).

Bis-1, 3-(2-methylquinolin) imidazolium chloride (9f)


A mixture of 1-(2-methylquinolin) imidazole (1.00g, 4.79mmol), 2-(chloromethylquinoline) monohydrogen chloride (1.02g, 4.79mmol) and K_2CO_3 (0.50g, 3.62mmol) in 50mL acetonitrile was refluxed for 24 hours. The solvent was removed and the residue dissolved in dichloromethane. The solution was filtered to remove the KCl formed and unreacted K_2CO_3 and the filtrates were concentrated to few millilitres. Addition of THF to the solution precipitated the product which was washed repeatedly with THF to give a pale yellow product. (Yield, 0.98g; 72%). Anal. Calcd. for $C_{23}H_{19}N_4Cl$: C, 71.41; H, 4.92; N, 14.48%. Found: C, 71.08; H, 5.00; N, 14.07%. HRMS: Calculated for

$C_{23}H_{19}N_4Cl$: 351.1610. Found: 351.1606. 1H NMR ($CDCl_3$, 400MHz, 298K): 11.20(NCHN), 8.10(d, 2H, $J = 8.4$ Hz, quin-H), 7.85(d, 2H, $J = 8.4$ Hz, quin-H), 7.70(m, 4H, $J = 8.3$ Hz, quin-H), 7.65(m, 4H, quin-H, CHHC), 7.45(t, 2H, $J = 6.7$ Hz, quin-H), 5.80(s, 4H, NCH₂C). ^{13}C NMR ($CDCl_3$, 100MHz, R.T.): 152.79, 147.58, 138.44, 137.98, 130.12, 129.13, 127.76, 127.69, 127.57, 120.67, 54.57.


2.4.7 Acridine based imidazolium salt

To prepare the desired imidazolium salt, 4-chlorooctahydroacridine was prepared following Paine multi step synthesis [29] and then coupled with mesitylimidazole.


2-Dimethylaminomethylcyclohexanone (10):




A mixture of cyclohexanone (22.30g, 0.23 mol), dimethylamine hydrochloride (9.9 g, 0.12 mol), and formaldehyde (36.90 g of 37% solution in water) was refluxed (oil bath at 130 °C, 30 min) and then cooled to room temperature. Sodium chloride (4.25g) was added, and the mixture was stirred at 23 °C (20 min). The mixture was transferred to a separatory funnel, and the organic and aqueous phases were separated. The aqueous phase was extracted with Et_2O (4 X 10 mL) to further remove unreacted cyclohexanone, and adjusted to pH= 13.5 by addition of 9.50 g of KOH in 22.5 mL of water. The mannich base was separated as yellow oil, which exhibited a strong amine odour. The aqueous phase was extracted with Et_2O (3 x 20 mL), and the yellow oil and the ether extracts were combined and then dried over anhydrous sodium sulphate. After removal of the ether at 23 °C under reduced pressure, the remaining oil was distilled under vacuum (42-43 °C/100 mTorr): yield 16.00 g (90%). 1H NMR ($CDCl_3$ -d₃:400 MHz 298K): $\delta = 2.65$ (m, 1H, $J = 6.0$ Hz), 2.40-2.20(m,3H, $J = 3.4$ Hz), 2.10(m, 8H, $J = 6.2$ Hz), 1.9(m, 1H, $J = 2.7$ Hz,), 1.8(m, 1H, $J = 4.8$ Hz,), 1.6(m, 2H, $J = 3.0$ Hz), 1.3(m, 1H, $J = 11.1$ Hz,) ^{13}C NMR (DCM, 400MHz R.T.): 24.30, 27, 80, 32.23, 41.69, 45.57, 48.73, 58.80, 212.27.

2, 2'-Dicyclohexanoylmethane (11):

2-Dimethylaminomethylcyclohexanone (19.5 g, 0.125 mol) and cyclohexanone (37 g, 0.38 mol) were mixed and refluxed (oil bath at 205 °C, 1.5 h), and the resulting mixture was distilled under vacuum. The fraction collected at 90-100 °C/ 100 mTorr was colourless oil (20.75g). Hexane (25mL) was added to the oil and mixture was cooled to -30 °C overnight. A white solid was collected and washed with cold hexane (3 x 15 mL). The colourless solid product was obtained: yield 14.63 (56.2%). ¹H NMR (CDCl₃:400 MHz 209K): δ = 1.0-2.4(m, 20H): ¹³C NMR (DCM, 400MHz R.T.): 24.37, 24.60, 27.56, 29.11, 29.88, 33.88, 34.68, 41.44, 41.77, 47.18, 48.33, 212.33, 212.83.

Sym-Octahydroacridine (12):


Hydroxylamine hydrochloride (7.63 g, 0.11 mol) was added with stirring to a boiling solution of 2, 2-Dicyclohexanone (14.13g, 0.068mol) in EtOH (100 ML). The mixture was refluxed (20 min), after cooling, the ethanol was removed under reduced pressure, and 25 mL of water was added to dissolve the residue. A solution of NaOH (5.00g) in water (25mL) was added at 0 °C to bring the pH of the solution to 13.5. The solid that appeared was collected, washed with water and dried. Yield 11.7 g (92. 2%). ¹H NMR (CDCl₃-d₃:400 MHz R.T): δ = 7.00 (s 1H, aromatic-H), 2.80(m, 4H, J = 6.2 Hz), 2.6(m, 4H, J = 6.3 Hz), 1.8(m, 4H J = 6.2 Hz), 1.6(m, J = 6.2 Hz), ¹³C NMR (DCM, 400MHz R.T.): 22.53, 22.95, 27.98, 31.75, 128.85, 137.14, 153.51.

Sym-Octahydroacridine N-Oxide (13):

A mixture of sym-Octahydroacridine (11.50 g, 0.062 mol) and 3-chloroperoxybenzoic acid (17.75 g, 77% max) in CHCl₃ (100ml) was stirred (17h) at 23 °C. The resulting reaction mixture was extracted with NaHCO₃ (32.50 g) and Na₂CO₃ (15 g) in water (350 mL). The aqueous phase (pH = 8.5)


was washed with CH_2Cl_2 (2 x 75 mL), and the chloroform and dichloromethane solutions were combined and dried with anhydrous sodium sulphate. After removal of the solvent at 23 °C under reduced pressure, the product sym-octahydroacridine N-oxide was obtained as a light yellow solid: yield 12.0 g (96.3%). ^1H NMR (CDCl_3 -d₃: 400 MHz R.T): 6.7(s, 1H), 2.80(m 4H, J = 6.5 Hz), 2.6(m, 4H, J = 6.2 Hz), 1.8(m, 4H, J = 6.4 Hz), 1.6(m, 4H, J = 2.4 Hz); ^{13}C NMR (CDCl_3 , 100MHz R.T.): 21.19, 21.44, 24.11, 27.58, 125.58, 131.10.

1, 2, 3, 4, 5, 6, 7, 8-octahydroacridin-4-ol (14):



Trifluoroacetic anhydride (10.2mL, 0.072mol, 2.5 equivalent) was slowly added to a stirred solution of octahydroacridine N-oxide (5.85g, 0.029mol) in dry DCM (50ml) causing a slight increase in temperature of the solution. The solution was allowed to stir for a further hour at room temperature. The volatiles were then removed under reduced pressure to leave a yellow viscous residue, which was taken up in 20ml DCM. This was then saponified by the addition of a 2M solution of sodium carbonate, the biphasic mixture being vigorously stirred for 3 hours. The organic phase was then separated and the aqueous phase washed twice with DCM. The combined organic extract extracts were then washed with water and brine and then dried over magnesium sulphate. The solvent was removed to give octahydroacridine-4-ol (4.98g, 85%). ^1H NMR (CDCl_3 , 400MHz, δ): 7.05(s, 1H, ArH), 4.55(m, 1H, J = 5.5 Hz CHOH), 4.05(broad, 1H, CHOH), 2.8(m, 2H, J = 6.3 Hz CH_2), 2.7(m, 4H, J = 5.9 Hz CH_2), 2.2(m, 1H, J = 3.2 Hz CH_2), 1.6-2.0(m, 7H, J = 4.8 Hz CH_2). ^{13}C NMR (CDCl_3 , 100MHz, δ): 154.86, 154.35, 137.5, 130.97, 128.55, 68.36, 31.88, 31.15, 28.42, 27.99, 23.16, 22.79, 19.39.


4-chlorooctahydroacridine (15):



Thionyl chloride (15mL) in CHCl_3 (15mL) was added to a solution of octahydroacridin-4-ol (7g, 0.0345mol) in 25mL DCM. The mixture was stirred at

room temperature for 10 minutes, and then refluxed for an hour at 80 °C. The excess thionyl chloride and volatile by-products were then removed under reduced pressure and the residue dissolved in DCM (75mL). The solution was then extracted with Na₂CO₃ (7.5g) in water (125mL) and the aqueous phase (pH 8.5) was washed with DCM (2x100mL). The DCM extract was then dried with Na₂CO₃ and the solvent removed under reduced pressure to give a yellow solid of 4-chlorooctahydroacridine (4.90g, 64%). ¹H NMR (CDCl₃, 400MHz, 298K): 7.05 (s, 1H, ArH), 5.20 (m, 1H, J = 2.8 Hz CHCl), 2.55-2.95(m, 6H, J = 4.7 Hz, CH₂), 2.05-2.35(m, 3H, J = 2.9 Hz CH₂), 1.65-1.85 (m, 5H, J = 4.3 Hz CH₂). ¹³CNMR (CDCl₃, 100MHz, δ): 155.53, 151.22, 137.96, 132.35, 128.97, 59.44, 32.65, 32.24, 28.60, 27.54, 23.15, 22.65, 17.42.

1-mesityl-3-octahydroacridinimidazolium chloride (16):



4-chlorooctahydroacridine (1.00g, 4.5mmol) and mesityl imidazole (0.84g, 4.5mmol) were placed in an ACE pressure tube with 10 mL THF. The mixture was refluxed at 90 °C for 10 days as a dark precipitate formed. The precipitate was filtered and washed with Et₂O (4x10mL) to give the imidazolium salt as an off-white solid (0.15g, 8%). The filtrate was placed back under reflux where it continued to react further over time. Anal. Calcd. for C₂₅H₂₉N₃Cl: C, 73.80; H, 7.13; N, 10.33%. Found: C, 65.65; H, 7.11; N, 8.98%. ¹H NMR (CDCl₃, 400MHz, 298K): 10.15 (s, 1H, NCHN), 8.18(s, 1H, CHHC), 7.65(s, 1H, CHHC), 7.20(s, 1H, ArH), 6.95 (s, 2H, ArH), 6.55 (d, 1H, J = 9.7 Hz CHN), 1.80-3.40 (m, 23H, CH₂, CH₃). MS (ES) m/z (%): 372.3(29) [M-Cl]⁺; MS (ESI) m/z (%): found 372.3424 [M-Cl]⁺; expected: 372.5333.

2.4.8 Crystal structure solution

All single crystal X-ray chromatographic determinations presented in this study were kindly performed by Dr. Dirk Beetstra and Dr. Benson Kariuki at the Cardiff University.

X-ray data collection was carried out at 150K on a Bruker/Nonius Kappa CCD diffractometer using graphite monochromated Mo-K α radiation, equipped with an Oxford Cryostream cooling apparatus. The data was corrected for Lorentz and polarization effects and for absorption using SORTAV [36]. Structure solution was achieved by direct methods [37] and refined by full matrix least – square on F2 with all non- hydrogen atoms assigned anisotropic displacement parameters. Hydrogen atoms attached to carbon atoms were placed in idealised positions and allowed to ride on the relevant carbon atom. In the final cycles of refinement a weighting scheme that gave a relatively flat analysis of variance was introduced and refinement continued until convergence was reached. Structure refinement and final geometrical calculations were carried out with the SHELXL-97 [38] program implemented in the WinGX and the diagrams were generated using the ORTEP-3 for windows [39] program.

Table 2.9: Crystal data and structure refinement for 4

| | | |
|------------------------|-----------------------------------------------------------------|------------------------------|
| Empirical formula | C ₂₂ H ₁₉ F ₆ N ₄ P | |
| Formula weight | 484.38 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Triclinic | |
| Space group | P -1 | |
| Unit cell dimensions | a = 7.8550(3) Å | $\alpha = 75.6640(10)^\circ$ |
| | b = 8.300(3) Å | $\beta = 82.7570(10)^\circ$ |
| | c = 16.4760(8) Å | $\gamma = 80.586(2)^\circ$ |
| Volume | 1022.59(7) Å ³ | |
| Z | 2 | |
| Density (calculated) | 1.573Mg/m ³ | |
| Absorption coefficient | 0.208mm ⁻¹ | |

| | |
|-----------------------------------|--------------------------------------------|
| F000 | 496 |
| Crystal size | 0.30 x 0.10 x 0.10 mm ³ |
| Theta range for data collection | 2.56 to 35.01 ° |
| Index ranges | -12<=h<=12, -13<=k<=13, -26<=l<=26 |
| Reflections collected | 12823 |
| Independent reflection | 8815 [R _{int} = 0.745] |
| Completeness of theta = 35.01° | 97.7% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | .9795 and 0.9403 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 8815/0/298 |
| Goodness of fit on F ² | 1.029 |
| Final R indices [I>2 sigma(I)] | R1 = 0.0802, wR2 = 0.1579 |
| R indices (all data) | R1 = 0.1888, wR2 = 0.1992 |
| Largest diff. peak hole | 0.364 and -0.573e Å ⁻³ |

Table 2.10: Crystal data and structure refinement for 6b

| | | |
|------------------------|----------------------------------------------------------------|------------------|
| Empirical formula | C ₁₉ H ₁₆ BN ₃ F ₄ | |
| Formula weight | 373.16 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Triclinic | |
| Space group | P -1 | |
| Unit cell dimensions | a = 7.1833(2) Å | α = 82.4233(9)° |
| | b = 7.9742(2) Å | β = 79.9094(10)° |
| | c = 16.0924(4) Å | γ = 76.9424(11)° |
| Volume | 879.97(4) Å ³ | |
| Z | 2 | |
| Density (calculated) | 1.408Mg/m ³ | |
| Absorption coefficient | 0.114mm ⁻¹ | |
| F000 | 384 | |
| Crystal size | 0.08 x 0.25 x 0.35 mm ³ | |

| | |
|-----------------------------------|--------------------------------------------|
| Theta range for data collection | 2.945 to 27.506 ° |
| Index ranges | -9<=h<=9, -10<=k<=10, -20<=l<=20 |
| Reflections collected | 14794 |
| Independent reflection | 3998 [R _{int} = 0.120] |
| Completeness of theta = 26.13° | 99.50% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | 0.9900 and 0.9700 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 3998/0/245 |
| Goodness of fit on F ² | 1.2019 |
| Final R indices [I>2 sigma(I)] | R1 = 0.0793, wR2 = 0.1677 |
| R indices (all data) | R1 = 0.0578, wR2 = 0.1560 |
| Largest diff. peak hole | 0.400 and -0.370e Å ⁻³ |

Table 2.11: Crystal data and structure refinement for 6C

| | | |
|---------------------------------|-------------------------------------------------|-------------------|
| Empirical formula | C ₁₃ H ₁₂ IN ₃ | |
| Formula weight | 337.16 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Monoclinic | |
| Space group | P 21/C | |
| Unit cell dimensions | a = 7.2140(10) Å | α = 90.00° |
| | b = 19.3630(3) Å | β = 126.8350(10)° |
| | c = 11.5400(2) Å | γ = 90.00° |
| Volume | 1290.16(3) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.736Mg/m ³ | |
| Absorption coefficient | 2.464mm ⁻¹ | |
| F000 | 656 | |
| Crystal size | 0.68 x 0.25 x 0.25 mm ³ | |
| Theta range for data collection | 3.01 to 27.50 ° | |

| | |
|-----------------------------------|--------------------------------------------|
| Index ranges | -9<=h<=9, -25<=k<=25, -14<=l<=14 |
| Reflections collected | 18959 |
| Independent reflection | 2943 [R _{int} = 0.1107] |
| Completeness of theta = 27.50° | 99.50% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | 0.5779 and 0.2851 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 2943/0/153 |
| Goodness of fit on F ² | 1.071 |
| Final R indices [I>2 sigma(I)] | R1 = 0.0360, wR2 = 0.0737 |
| R indices (all data) | R1 = 0.0305, wR2 = 0.0702 |
| Largest diff. peak hole | 0.643 and -1.089e Å ⁻³ |

Table 2.12: Crystal data and structure refinement for 9b

| | | |
|---------------------------------|----------------------------------------------------------------|----------------|
| Empirical formula | C ₂₂ H ₂₂ BF ₄ N ₃ | |
| Formula weight | 415.24 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Monoclinic | |
| Space group | P 21/C | |
| Unit cell dimensions | a = 9.8460(4) Å | α = 90.00° |
| | b = 12.5190(6) Å | β = 93.507(3)° |
| | c = 17.2380(7) Å | γ = 90.00° |
| Volume | 2120.81(16) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.300Mg/m ³ | |
| Absorption coefficient | 0.01mm ⁻¹ | |
| F000 | 864 | |
| Crystal size | 0.30 x 0.30 x 0.20 mm ³ | |
| Theta range for data collection | 2.663 to 27.47 ° | |
| Index ranges | -12<=h<=12, -14<=k<=16, -22<=l<=22 | |
| Reflections collected | 8027 | |

| | |
|---------------------------------------|--------------------------------------|
| Independent reflection | 4825 [$R_{int} = 0.559$] |
| Completeness of theta = 27.47° | 99.30% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | 0.9800 and 0.9702 |
| Refinement method | Full matrix least square on F^2 |
| Data/restraints/parameters | 4825/267/302 |
| Goodness of fit on F^2 | 1.028 |
| Final R indices [$I > 2 \sigma(I)$] | $R1 = 0.1366$, $wR2 = 0.1814$ |
| R indices (all data) | $R1 = 0.0650$, $wR2 = 0.1476$ |
| Largest diff. peak hole | 0.263 and -0.320 e \AA^{-3} |

Table 2.13: Crystal data and structure refinement for 9c

| | | |
|---------------------------------|------------------------------------------|---------------------------|
| Empirical formula | $C_{16}H_{18}IN_3$ | |
| Formula weight | 379.23 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 \AA | |
| Crystal system | Monoclinic | |
| Space group | P C | |
| Unit cell dimensions | $a = 11.2770(4) \text{\AA}$ | $\alpha = 90.00^\circ$ |
| | $b = 12.300(5) \text{\AA}$ | $\beta = 90.100(5)^\circ$ |
| | $c = 11.5370(4) \text{\AA}$ | $\gamma = 90.00^\circ$ |
| Volume | 1600.26(10) \AA^3 | |
| Z | 4 | |
| Density (calculated) | 1.574 Mg/m^3 | |
| Absorption coefficient | 1.996 mm^{-1} | |
| F000 | 752 | |
| Crystal size | 0.30 x 0.22 x 0.15 mm^3 | |
| Theta range for data collection | 3.02 to 27.42 ° | |
| Index ranges | -14 ≤ h ≤ 11, -10 ≤ k ≤ 15, -14 ≤ l ≤ 14 | |
| Reflections collected | 8017 | |
| Independent reflection | 5779 [$R_{int} = 0.287$] | |
| Completeness of theta = 27.42° | 92.30% | |
| Absorption correction | Semi-empirical from equivalents | |

| | |
|-----------------------------------|--------------------------------------------|
| Max and min. transmission | 0.7339 and 0.5859 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 5779/50/306 |
| Goodness of fit on F ² | 1.055 |
| Final R indices [I>2 sigma(I)] | R1 = 0.0627, wR2 = 0.1040 |
| R indices (all data) | R1 = 0.0434, wR2 = 0.0939 |
| Largest diff. peak hole | 1.046 and -0.934e Å ⁻³ |

Table 2.14: Crystal data and structure refinement for 9f

| | | |
|---------------------------------|--------------------------------------------------|-----------------|
| Empirical formula | C ₂₃ H ₁₀ ClN ₄ | |
| Formula weight | 386.86 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Monoclinic | |
| Space group | P 1 21/n 1 | |
| Unit cell dimensions | a = 8.34480(10) Å | α = 90.00° |
| | b = 20.0690(4) Å | β = 95.5741(8)° |
| | c = 11.4953(2) Å | γ = 90.00° |
| Volume | 1916.04(6) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.341 Mg/m ³ | |
| Absorption coefficient | 0.216 mm ⁻¹ | |
| F000 | 808 | |
| Crystal size | 0.20 x 0.25 x 0.25 mm ³ | |
| Theta range for data collection | 2.030 to 27.422 ° | |
| Index ranges | -10 ≤ h ≤ 10, -24 ≤ k ≤ 26, -14 ≤ l ≤ 14 | |
| Reflections collected | 30860 | |
| Independent reflection | 4358 [R _{int} = 0.189] | |
| Completeness of theta = 27.442° | 99.70% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.9600 and 0.9500 | |
| Refinement method | Full matrix least square on F ² | |

| | |
|---------------------------------------|---------------------------------|
| Data/restraints/parameters | 2307/0/254 |
| Goodness of fit on F ² | 0.9465 |
| Final R indices [$I > 2 \sigma(I)$] | R1 = 0.0422, wR2 = 0.1019 |
| R indices (all data) | R1 = 0.0792, wR2 = 0.1145 |
| Largest diff. peak hole | 0.29 and -0.29e Å ⁻³ |

2.5 References

1. W. A. Herrmann and C. Kocher, *Angew. Chem. Int. Ed. Engl.*, **1997**, *36*, 2162-2187.
2. W. A. Herrmann, T. Weskamp and V. P. W. Bohm, *Adv. Orgmet. Chem.*, **2001**, *48*, 1-69.
3. D. Bourissou, O. Guerret., F. Gabbai, G. Bertrand, *Chem. Rev.*, **2000**, *100*, 39-91.
4. A. Danopoulos, N. Tsoureas, J. Wright, M. Wright, *J. Am. Chem. Soc.*, **2004**, *23*(2), 166-168.
5. D. J. Nielsen, K.J. Cavell, B. W. Skelton, A. H. White, *Inorganic Chimica Acta*, **2006**, *359*, 1855-1869.
6. A.R.Chianese, X. Li. M.C.Jansen, J.W.Faller, R. H. Crabtree, *Organometallics*, **2003**, *22*, 1663.
7. A.A.D. Tulloch, S. Winston, A.A. Danopoulos, G. Eastham and M. B. Hursthouse, *J. Chem. , Dalton Trans.*, **2003**, 699-708.
8. A. W. Waltman and R. H. Grubbs, *Organometallics*, **2004**, *23*, 3105-3107.
9. M. M. Viciu, R. F. Germaneau, S. P. Nolan, *Org. Lett.* **2002**, *4*, 4053-4056.
10. D. McGuinness, K. J. Cavell, *Organometallics* **1999**, *18*, 1596-1605.
11. D. McGuinness, K. J. Cavell, *Organometallics*, **2000**, *19*, 741-748.
12. J. Schwarz, V. P. W. Bohm, M. G. Gardner, M. Grosche, W.A. Herrmann. G. Rausdaschl-Sieber, *Chem. Eur. J.*, **2000**, *6*, 1773-1780.
13. L. C. Branco, J. N. Rosa, R. J. J. Moura, C. A. M. Alfonso, *Chem. Eur. J.* **2002**, *18*, 3671.
14. J. Fraga-Dubreuli, J. P. Bazureau, *Tetrahedron Letters*, **2001**, *42*, 6097.
15. K. -S. Kim, D. Dembereinyamba, H. Lee, *Longmuir* **2004**, *20*, 556.

16. H. Schottenberger, K. Wurst, U. E. I. Horvath, S. Cronje, J. Lukasser, J. Polin, J. M. McKenzie, H. Raubenheimer, *Dalton Trans.* **2003**, 4275.
17. D. Zhao, Z. Fei, T. J. Geldbach, R. Scopelliti, G. Laurenczy, P. J. Dyson, *Helv. Acta*, **2005**, *88*, 665.
18. A. A. Gridnev and I. M. Mihaltseva, *Synth. Commun.* **1994**, *24*, 1547.
19. A. J. Arduengo III, F. P. Gentry, P. K. Taverkere and H. E. Simmons; E. I. Du Pont de Nemours and Company: **US6177575**, **2001**.
20. M. Zettlizer, T. H. Dieck, E. Haupt, L. Stamp, *Chem. Ber.* **1986**, *119*, 1868-1875.
21. C. Michon, A. Ellern, R. J. Angelici, *Inorganic Chimica Acta* **2006**, *359*, 4549-4556.
22. (a) J. P. Wolfe, S. Wagaw, J. -F. Marcoux, S. L. Buchwald, *Acc. Chem. Res.* **1998**, *31*, 805.
(b) J.F. Hartwig, *Acc. Chem. Res.* **1998**, *31*, 852.
23. K. A. Jensen, P. H. Nielsen, *Acta Chem. Scand.* **1964**, *18*, 1
24. P.H. Nielsen, O. Dahl, *Acta Chem. Scand.* **1966**, *20*, 1113.
25. T. Okada, M. Yoshida, T. Hirose, K. Kasuga, T. Yu, M. Yuasa, I. Sekine, *Electrochimica Acta*, **2000**, *45*, 4419-4428.
26. N. Imlinger, M. Mayr, D. Wang, K. Wurst and M. R. Buchmeiser; *Adv. Synth. Catal.* **2004**, *346*, 1836-1843.
27. J. Liu, J. Chen, J. Zhao, Y. Zhao, L. Li, H. Zhang; *Synthesis* **2003**, No.17, 2661-2666.
28. P. L. Chiu, C. Lai, C. Chang, C. Hu and H. M. Lee, *Organometallics*, **2005**, *24* 6169-6178.
29. G. Steiner, H. Kopacka, K. Ongania, K. Wurst, P. Preishuber-Pflugl and B. Bildstein; *Eur. J. Inorg. Chem.* **2005**, 1325-1333.
30. C. Fontenas, E. Bejan, H. Haddou, G. G. Balavoine, *Synth. Comm.*, **1995** *25*(5)629-633.
31. A. Danopoulos, S. Winston, T. Gelbrich, M. Hursthouse, R. Tooze, *Chem. Comm.*, **2002**, 482-483.
32. D. Nielsen, K.J. Cavell, B. Skelton, A. White, *Inorg., Chim., Acta*, **2002**, *327*, 116-125.
33. R. W. Alder, M. E. Blake, C. Bortolotti, S. Bufali, C. P. Butts, E. Linehan, J. M. Oliva, G. A. Orpen, M. J. Quayle, *Chem. Commun.*, **1999**, 241-242.

34. R. W. Alder, M. E. Blake, C. Bortolotti, S. Bufali, C. P. Butts, G. A. Orpen, J. Schutz, S. J. Williams, *J. Chem. Soc., Perkin Trans.* **2001**, *1*, 1586-1593.
35. W.A. Herrmann, S. K. Schneider, K. Ofele, M. Sakamoto, E. J. Herdweck, *J. Organomet. Chem.* **2004**, *689*, 2441-1449.
36. R.H. Blessing, *Acta Crystallogr. Sect A* **1995**, *51*, 33.
37. A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, Sir-92, *J. Appl. Crystallogr.* **1993**, *26*, 343.
38. G. M. Sheldrick, SHEXL-97, *Institut fur Anorganische Chemie der Universitat, Tammanstrasse 4, D-3400 Gottingen, Germany* **1998**.
39. L. J. Farrugia, *J. Appl. Cryst.*, 1997, *30*, 565.
40. C. Wang, C. Fu, Y. Liu, S. Peng, S. Liu, *Inorg. Chem.* **2007**, *46*, 5779-5786.


CHAPTER THREE

Silver(I) and Palladium(II) complexes of quinoline functionalised Heterocyclic Carbene ligands.

3.1 Silver(I) (NHC) complexes

3.1.1 Introduction

The synthesis of Ag^I(NHC) was first reported by Arduengo in 1993 using the free carbene route [1]. This was accomplished by deprotonation of the imidazolium salt **3.01** to make the free carbene (1, 3-dimesitylimidazolin-2-ylidene) **3.02** and subsequent reaction of **3.02** with silver triflate to yield the desired homoleptic Ag^I(NHC) **3.03**. Several other Ag^I(NHCs) have been synthesised using this method [1, 2, 3,4,5,]. However, this method has been applied to the synthesis of only a limited number Ag^I(NHC) due to the difficulty of generating most free carbenes, which have other acidic protons especially azolium salts with methylene linkers [6].




Scheme 3.0: Synthesis of Arduengo homoleptic Ag carbene complex

Ag^I(NHC) can also be accessed by transmetalation. Transmetalation of the NHC ligands to AgPF₆ yielding homoleptic imidazolindin-2-ylidene complexes **3.04-3.06** was also reported [7].

The *in situ* deprotonation of imidazolium salts with basic silver precursors is the most commonly used method to synthesise Ag^I(NHC) complexes. In their reaction Bertrand et al demonstrated that Ag(OAc) reacts with 1,2,4-triazolium

salts producing polymeric Ag^I(1,2,4-triazolin-3,5-diylidene) complexes with alternating Ag^I and 1,2,4-triazolin-3,5-diylidene units [8]. The use of Ag(OAc) protocol has not been widely utilised with the reports of Ag^I(NHC) complexes being restricted to a few examples with symmetrically alkyl substituted NHCs.




3.04: R = Et

3.05: R = allyl

3.06: R = benzyl

Scheme 3.2: Synthesis of Ag^I(NHC) via transmetalation

The most commonly used base is silver (I) oxide, pioneered by Lin and Wang [9]. These workers reported that stirring 1, 3-diethylbenzimidazolium bromide with Ag₂O in DCM, or with AgBr and NaOH under phase transfer conditions gave the Ag^I(benzimidazol-2-ylidene)₂ complexes **3.07** and **3.08** in high yield, Scheme 3.3.



3.07: X = AgBr₂⁻

3.08: X = PF₆⁻

Scheme 3.3: Wang and Lin's preparation of Ag^I(benzimidazol-2-ylidene)₂ complexes via silver(I) oxide.

The principle advantage of the Ag₂O protocol is its tolerance to oxygen and moisture, indeed water is the by-product of the reaction. There are in fact reports

of the reactions being carried out at ambient temperature, in a variety of solvents including water [10-12]. The formation of silver complexes in water suggests that the deprotonation of the imidazolium salt and coordination to metal centre is a concerted process because free carbenes are water sensitive [13]. Several groups have successfully prepared a variety Ag^I(NHC) complexes through the application of the Ag₂O method of Wang and Lin with different kinds of imidazolium salts [14-21]. Modification of Wang and Lin's original method used dichloroethane as solvent [17], thereby allowing the reaction to be carried out at elevated temperatures for less reactive imidazolium salts.

In the cases where the imidazolium salts are insoluble in DCM, the use of solvent mixtures such as DCM-MeCN [22] and DCM – EtOH [23] has been found to be useful. The reaction can also be performed in DMSO at 55°C as reported recently [24]. Furthermore, addition of molecular sieves to the reaction has also been reported to facilitate the formation of Ag^I(NHC) complexes by removing the water generated in the reaction as shown in Scheme 3 above [17]. Silver(I) carbonate was also reported to have been used in the synthesis of Ag^I(NHC), though less effective than the Ag₂O protocol as a longer reaction time is required for the reaction to reach completion [17]

McGuinness and Cavell reported the synthesis of a series of mono donor-functionalised Pd^{II}(NHC) by the use of Ag₂O protocol via the transmetalation Ag^I(NHC) complexes **3.09** – **3.10** with appropriate palladium precursors [16].





Figure 3.10: Cavell carbonyl and pyridyl functionalised Ag^I(NHC) used to prepare Pd^{II}(NHC) complexes

The Cavell group also reported the application of the Ag₂O protocol to synthesise phenoxy- functionalised [25], pyridyl-functionalised [26] and thiophene and furan-functionalised [27]Ag^I(NHC) complexes. A range of imidazolium salts bearing a diverse variety of functionalised groups have been shown to be compatible with the formation of Ag^I(NHC) complexes via Ag₂O. Thus, Ag^I(NHC) complexes with amine [18], ferrocenyl [15], imines[17], amide [17] and alkoxy functionalised bis-NHC have been prepared through the Ag₂O route. Other Ag^I(NHC) prepared were not isolated but used directly as transmetallation reagents [18].

The wide applicability and ease of preparation of stable Ag^I(NHC) complexes via the Ag₂O route and the good transmetallation properties of these complexes promised ready access to donor-functionalised NHC complexes of catalytically interesting transition metals [28]. The ability to obtain Ag^I(NHC) complexes from the reaction of Ag₂O with imidazolium, saturated imidazolidinium and benzimidazolium salts indicates that their formation is relatively unaffected by the electronics of the heterocyclic ring [28]. The reaction is also tolerant of steric group and a wide range functional groups on the NHC N-substituents.

Halogen exchange reactions have been found to occur when NHC complexes are synthesised in chlorinated solvents. Danopoulos and co-workers reported halide exchange reactions when synthesising Ag^I(NHC) in 1,2-dichloroethane or dichloromethane [17]. An example of the halide exchange is depicted in Scheme 3. 4. Similarly Lin and co-workers reported the salt metathesis of imidazolium iodide salts with chloride from methylene chloride during the synthesis of Ag^I(NHC) complexes [22].



Scheme 3.4: Halide exchange in Ag^I(NHC) complex synthesis

Mixtures of halogeno and halide anions of silver NHCs have been reported [18, 29]. To reduce the complications due to inorganic halogeno complexes and cluster formation, anion exchange of the imidazolium halide salt for a noncoordinating anion has been widely employed. After the anion exchange, the synthesis of the Ag^I(NHC) proceeds cleanly to one product. Anion exchange of the halide after the synthesis of the Ag^I(NHC) has also been performed using reagents such as AgBF₄, to obtain clean products without the complexities of the halide [26,29,30,31].

Theoretical calculations of group 11 NHC carbenes showed that the metal-carbene bond strengths follow the pattern Au > Cu > Ag [13]. While the bond strength of Ag^I(NHC) complexes were shown to be relatively the weakest, the overall strength of 56.5 Kcal/mol was considered strong enough to stabilise the Ag^I(NHC) [32].

3.1.2 Structural diversity in Ag^I(NHC) complexes


The structural characterisation of N-heterocyclic carbene complexes of silver has led to very complex bonding motifs in the solid state, especially in complexes with halide anions. The ability of silver to form complex anions of the formula [AgX₂]⁻ (X = halogen), coordinate to either one or two NHC moieties and engage in Ag(I)...Ag(I) interactions in the solid state appear to account for most of this structural diversity. The interactions between Ag(I) and functional groups present on the NHC ligand are weak compared to most of the Ag(I)...Ag(I) interactions observed in Ag(I)NHCs.

Ag^I(NHC) complexes with non coordinating anions exist as biscarbene salts with the cationic silver bound by two carbene moieties and the noncoordinating anion balancing out the charge (C₂- Ag).


Studying the crystal structure of their non functionalised Ag^I(NHC) complex Wang and Lin [9] reported a mononuclear complex with two NHC ligands coordinated to Ag(I) and the counter ion [AgBr₂]⁻ coordinating through a Ag(I)...Ag(2) interaction above the plane of the cation. The Ag(I)...Ag(2) distance of 2.954 Å was found to be significantly shorter than the van Waals contact distance of 3.44 Å. and relatively short for an unsupported Ag(I)...Ag(I) interaction [9].

Another study of non functionalised Ag^I(NHC) complex **3.11** revealed a linear NHC-Ag(I)-NHC motif with coordinated [AgBr₂]⁻ counterion almost perpendicularly aligned [22]. Other structural studies of Ag^I(NHC) complexes revealed mono NHC complexes with coordination anions or uncoordination anions. For example, in **3.12** the C₂-Ag-X strings deviate from linearity [5, 21]. **3.13** [19] and **3.14** [22] may therefore be considered to be aggregated **3.12** motifs formed due to Ag...X and weak Ag...Ag interactions between adjacent molecules. As a result of alternating Ag-Br and Br-Ag units in the solid state, the bond length around Ag₂Br₂ rhomboids in **3.14** are unequal.


Ag^I(NHC) complexes that have donor functionalised group feature most of the structural diversities observed in the non functionalised counter parts. In **3.15** the [AgBr₂]⁻ counter ion is coordinated to the pyridine functionality of one of the NHC ligands that make up the linear NHC-Ag(I)-NHC unit [17]. Other reported examples are shown in structures **3.16** and **3.17** where they exhibited rhomboid Ag₂Br₂ and linear NHC-Ag(I)-X geometries respectively [17]. Also reported is a dinuclear Ag^I(NHC) complex **3.16** with Ag₂L₂ formulation [3]. The structure of **3.18** is that of double helical unit with the C₂-Ag-C₂ strings distorted by 14.5° from the linear orientation because of relatively rigid ligands Ag...Ag' bonding interaction is present and brings the these atoms to within 3.158 Å. The pyridyl group of each ligand is equidistant to Ag, Ag' at a distance of 3.02 Å. Structure **3.19** is a trinuclear pyridyl functionalised Ag^I(NHC) complex obtained by the reaction of 3-methyl-1-picolylimidazolium iodide with 2.5 eq mole of Ag₂O in DCM [33]. The geometry at the silver centre is planar, with every Ag coordinated by two carbene atom and one triply bridging iodine. The three Ag(I) cations are linked by the bridging I(1) anion symmetrically with an Ag(1)-I(1) distance of 3.04 Å which is more longer than those of the Ag-I(bridging) bonds 2.83 and 2.78 Å [37], and the net charge of 2+ is balanced by two non interacting iodide ions [33].




3.07




3.11




3.12




3.13




3.14




3.15




3.16



3.17



3.18



3.19

Figure 3.11: Bonding motifs in Ag^I(NHC) complexes

Ag...Ag interactions have not been observed in Ag^I(NHC) complexes that have been prepared in the absence of coordinating halide anions. In Ag^I(NHC) complexes with triflate [1,3,8], barf[15], nitrate[22], and carborane [4] anions, homoleptic [Ag^I(NHC)₂]X complexes are formed with quasi-linear C₂-Ag-C₂ strings.

3.1.3 Ag^I(NHC) complexes as transmetallation reagents

The use of Ag^I(NHC) in homogenous catalysis has recently appeared in the literature: such as in the preparation of 1,2-bis(borane) esters [34], ring opening polymerization of lactides [35 , 36] and olefin polymerization [37]. However, the increased interest in the chemistry of Ag^I(NHC) complexes has been mainly due to their role as transfer agents in the development of many important metal-NHCs. The fact that active hydrogen atoms other than C₂-H can be protected effectively by this method solved difficulties encountered in the synthesis of metal- NHCs by other methods [16,18, 26,38]. NHC ligands have thus been transferred from Ag^I(NHC) complexes to a variety of metals including Cr^{III}, Fe^{III}, Co^{II}, Ni^{II} [38], Cu^I [1], Cu^{II} [39], Pd^{II} [9,14, 16,18,39,40,], Au^I [9,23], Rh^I and Ir^I [41].


Transmetallation may be carried out using isolated Ag^I(NHC) complexes or conveniently performed in a one pot protocol where the Ag^I(NHC) complex is

generated from an imidazolium salt and Ag₂O and then reacted in situ with a precursor of the desired metal. The transfer of NHC depends on the nature of Ag^I(NHC), the receiving metal precursors and the reaction conditions.

3.2.0 Palladium (II) complexes of quinoline functionalised heterocyclic carbene ligands


Pd^{II}(NHC) carbene complexes were first reported in 1995 by Herrmann [42] by the reaction of Pd(OAc)₂ with 1,3,-dimethylimidazolium iodide to give **3.20** and 3, 3-dimethyl-1,1-methylenediimidazolium diiodide to give **3.21** as shown in scheme 3.5. It was discovered that these new palladium complexes were stable to heat, air and moisture. In addition to this, these complexes were found to be excellent catalysts in Heck reactions.

As enumerated in chapter 1 NHCs exhibit properties that are complimentary to their application as ligands in catalysis. They are more strongly bonded to transition metals such as Pd than the widely used phosphine ligands [42, 43], and thus required only in stoichiometric amounts, and in general Pd^{II}(NHC) complexes are less toxic, easier to handle and exhibit better thermal stability than typical phosphine ligands.



Scheme 3.5: Herrmann's synthesis of **3.20** and **3.21**.

Following the discovery of the qualities of complexes **3.20** and **3.21**, there was renewed interest to explore catalytic applications of the Pd^{II}(NHC) system through modification of the NHC ligands. Among the important ligand design was the combination of the strongly bound NHC moiety with more weakly nucleophilic functional groups allowing access to ligands with hemilabile donor groups, as these can increase catalytic activities by stabilising the low-valent metal centres formed during catalysis. Along this line several Pd^{II}(NHC) complexes were reported bearing one or two functionalised substituents with carbamoyl (**3.22** [44,45]), ester (**3.23**[46] and **3.24**[47]), ethers (**3.25**[48]), hydroxyl(**3.26** [42], **3.27** [48]), oxazoline(**3.28**[49]), picolyl(**3.29** [50]), pyridyl (**3.30** and **3.31** [51], **3.32** [52]),imine **3.33** [53] groups Figure 3.12.






Figure 3.12: Structurally characterised examples of some reported Pd(II) complexes bearing functionalised NHC ligands.

Many synthetic routes have been employed to synthesise Pd^{II}(NHC) complexes and these have been outlined in chapter one. The routes to be used depend on the nature of available imidazolium salts. The Pd(OAc)₂ route is among the many routes available and is suitable for simple mono and bis imidazolium salts as shown by Herrmann in Scheme 3.5 above but does not give access to catalytically important Pd-hydrocarbyl species. Pd^{II}(NHC) complexes can also be prepared via the free carbene route by the use of strong bases such as NaH with catalytic amounts of KOtBu and amides (K[N(SiMe₃)₂]). Once the free carbene is formed, it can be reacted with appropriate metal source such as PdCl₂(COD) to produce the desired Pd^{II}(NHC) complexes. However free carbenes are known to be unstable especially in the absence of bulky aryl or alkyl groups on the 1 and 3 positions of the ring thereby making the route unsuitable in the synthesis of some Pd^{II}(NHC) complexes. The use of strong bases have been found to be unsuitable in accessing functionalised carbenes because of acidic protons commonly associated with functional group and any methylene linkers.

The chapter also reports the synthesis and characterisation of quinoline functionalised Ag^I(NHC) complexes. X-ray crystallographic studies have been

carried on two of the Ag^I(NHC) complexes prepared in this work and the data obtained compared with the published examples. The reported Ag^I(NHC) complexes were synthesised primarily as transmetallation reagents to access the catalytically important quinoline functionalised Pd^{II}(NHC) complexes and Ir^I and Rh^I(NHC) complexes.

Also described in this chapter are the Pd^{II}(NHC) complexes synthesised via transmetallation of the desired Ag^I(NHC) complexes and the complexes were characterised spectroscopically and by micro analysis.

3.3 Results and Discussion


3.3.1 Silver (I) complexes of quinoline functionalised NHC ligands

General comments

Quinoline functionalised imidazolium salts described in chapter two were reacted with Ag₂O to give stable complexes with the general formula [Ag^I(NHC)₂][AgX₂] or Ag^I(NHC)X based on the two x-ray structures of the complexes obtained. No attempt was made to abstract halide to isolate the Ag^I(NHC) complexes as salts of non coordinating anions such as tetrafluoroborate, hexafluorophosphate or triflate to improve solubility or purity of the complexes. For the purpose of clarity the quinoline based Ag^I(NHC) complexes described below are divided into two: the methylene bridged and the rigid quinoline Ag^I(NHC) complexes.

3.3.1.1 Silver(I) complexes of rigid quinoline functionalised NHC ligands

Two quinoline functionalised imidazolium salts (**6b** and **6c**) were reacted with Ag₂O to give the corresponding Ag^I(NHC) complexes. Two equivalent of the imidazolium salt **6b** was reacted in DCM with one equivalent of Ag₂O overnight at room temperature until the black suspension of Ag₂O disappeared. The reaction was performed free of light as most Ag^I(NHC) complexes are light sensitive. After work up, the Ag^I(NHC) complex **3.34** was obtained as a light brown powder. The room temperature ¹H NMR of complex **3.34** was consistent with the proposed structure with no evidence of residual imC₂-H resonance.




Scheme 3.6: Synthesis of rigid quinoline functionalised Ag^I(NHC) complexes

The proton peak of the methylene linker of the benzyl moiety was shifted upfield (5.35 ppm) relative to those of the corresponding imidazolium salt (5.85 ppm). The carbenic carbon could not be observed in the ¹³ NMR spectra and this is consistent with some of the reports that appeared in the literature as a significant number of silver^I(NHC) complexes were reported with no observable carbene resonances [8]. Elemental analysis returned a satisfactory result for complex **3.34**. Crystals suitable for X-ray chromatography were not obtained but the analytical data as well spectroscopic data are consistent with the proposed structure. Complex **3.35** was characterised by ¹H NMR, mass spectroscopy and micro analysis. The characteristic feature confirming the formation of the silver^I(NHC) complex is the disappearance of C₂-H in the ¹H NMR spectra.

3.3.1.2 Silver(I) complexes of methylene-bridged quinoline functionalised NHC ligands

The methylene-bridged quinoline functionalised Ag^I(NHC) complexes were prepared following the method reported by Wang and Lin [9], i.e. by interaction of the imidazolium salts with Ag₂O as shown in scheme 3.7.



Scheme 3.7: Synthesis of methylene bridged quinoline functionalised Ag^I(NHC) complexes

It was found that: (i) with the relatively unreactive sterically hindered imidazolium salts (**9e** and **9f**) the reaction was carried out in refluxing dichloromethane, whereas for all other imidazolium salts the reaction occurred at room temperature; (ii) synthesis in refluxing dichloromethane increases the formation of by-product (iii) the purity of the product is improved by addition of activated molecular sieves to the reaction medium.

Compound **3.36** was characterised by ¹H NMR, ¹³C NMR, microanalysis and X-ray crystallography. The room temperature ¹H NMR of Complex **3.36** was consistent with the proposed structure showing complete disappearance of imC₂-H resonance and the protons of the methylene linker was observed to move up field. In the ¹³C NMR there was sharp singlet peak at 181.18 ppm which is assignable to the carbene carbon of the Ag^I(NHC) complex. Microanalysis returned satisfactory results for complex **3.36**.

Crystals suitable for X-ray structural determination were obtained by diffusing Et₂O into a DCM solution of the complex. The crystal structure is depicted in figure 3.13 below.




Figure 3.13: ORTEP projection of complex **3.36** excluding hydrogen atoms for clarity showing atom labelling scheme.

Complex **3.36** as shown in the above figure consists of a linear $[\text{Ag}(\text{NHC})_2]^+$ cation and a linear $[\text{AgCl}_2]^-$ with the two ions associating through Ag^1 - Ag^1 interaction. The Ag-C bond distance ($\text{Ag}(1)\text{-C}(1) = 2.106(12)$) are comparable to those reported by Wang and Lin (2.073 \AA) [8]. The geometry of $\text{C}(1)\text{-Ag}(1)\text{-C}(1_2)$ is close to linear ($170.8^\circ(8)$) though it is lower than what was obtained by Wang and Lin (175.6°) [8]. The $\text{Ag}(1)\text{-Ag}(2)$ separation of 3.201 \AA is smaller than the contact van de Waals distance of 3.44 \AA and is at upper range of the ligand-unsupported Ag-Ag bond lengths (range $2.80\text{-}3.30 \text{ \AA}$) [54]. The $\text{Cl}(1)\text{-Ag-Cl}(1)$ angle of 176.6° deviates only slightly from that of the coordinated linear species. Some selected bond lengths (A) and angles (o) are presented in table 3.1 below.

Table 3.1: Selected bond lengths (Å) and bond angles (°) of **3.36**

| | | | | | |
|----------|-----------|-----------|-----------|-------------|-----------|
| Ag1- Ag2 | 3.201(4) | N2-C5 | 1.48(2) | C1-Ag1-Ag2 | 94.6(4) |
| Ag2-Cl1 | 2.290(5) | N2-C1-N1 | 104.0(10) | C1-Ag1-Ag2 | 85.4(4) |
| Ag1-C1 | 2.106(12) | N2-C1-Ag1 | 130.6(8) | Cl1-Ag1-Cl1 | 176.6(11) |
| C1-N1 | 1.353(16) | N1-C1-Ag1 | 125.2(9) | C3-C2-N1 | 107.8(11) |
| C1-N2 | 1.351(14) | C1-Ag1-C1 | 170.8(8) | C1-N2-C5 | 119.3(10) |

The ¹H NMR spectrum of complex **3.37** shows the disappearance of C₂-H signal with other signals consistent with the proposed structure. The ¹³C NMR spectrum reveals the absence of C_{carbene} resonance and the elemental analysis returned satisfactory results. A ES-MS showed a peak corresponding to cation [M⁺-Cl = 328.15] with intensity of 100% similar to that of the corresponding imidazolium salt indicating that under these conditions the silver complex undergoes decomposition. This decomposition of silver(NHC) complexes and the observation of the corresponding of the corresponding salts has been reported using ES-MS [55]. Crystals suitable for X-ray chromatography were obtained by diffusion of Et₂O into a saturated DCM solution of the complex and the crystal structure is depicted in Figure 3.14 below. The geometry of the silver in the carbene complex is that of distorted linear with C13- Ag1-Cl2 bond angle of 169.03° (10) which is similar to the silver carbene complex reported by Cesar and Gade

(169.4° (1)) [56] and lower than those reported by Danopoulos et al (176.1°(2)) [17], Pytkowicz et al (175.2°(5)) [39] and Paas et al (173.5°(2)) [57]. All other bond lengths and bond angles are consistent with the reported values [17,39, 56, 57]. Some selected bond length and bond angles are presented in table 3.2 below.




Figure 3.14: ORTEP projection of complex **3.37** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 3.2: Selected bond lengths (Å) and bond angles (°) of **3.37**

| | | | | | |
|---------|----------|-------------|------------|-------------|----------|
| Ag1-Cl2 | 2.350(9) | N5-C14 | 1.320(5) | Ag1-C13-N11 | 123.7(3) |
| Ag1-C13 | 2.090(3) | C4-C16 | 1.344(5) | N3-C13-N11 | 104.9(3) |
| N3-C13 | 1.342(5) | C4-N3 | 1.384(4) | N3-C6-C14 | 111.6(3) |
| N11-C13 | 1.354(4) | C13-Ag1-Cl2 | 169.03(10) | C13-N11-C22 | 122.6(3) |
| N3-C6 | 1.474(4) | Ag1-C13-N3 | 131.4(2) | C6-C14-N5 | 116.2(3) |

The formation of complexes **3.38**, **3.39**, **3.40** and **3.41** were confirmed by ¹H NMR and ¹³C NMR spectroscopy. The ¹H NMR revealed complete disappearance of the C₂-H protons while the C_{carbene} of complexes **3.38**, **3.39** and **3.41** appeared at 184.10, 183.23 and 180.37 ppm respectively, that of complex **3.40** was not observed. High resolution mass spectroscopy of complex **3.41** indicated only the decomposition of the complexes as only the its corresponding imidazolium salt ions was observed at M/z = 351.16 (100% intensity) equivalent

to [(NHC)+H]⁺. ¹H NMR and ¹³C NMR spectra of complex **3.38** are consistent with the bis carbene formulation with the C_{carbene} of the complex appearing at 184.10 ppm. HRMS showed cluster at M/z = 609.1886 (100% intensity) attributable to [Ag(NHC)₂]⁺ of the bis carbene. The ¹H NMR and ¹³C NMR data of complexes **3.36-3.41** is presented in Table 3.3 below.

Table 3.3: ¹H and ¹³C NMR data of complexes 3.36- 3.41

| Compound | ¹ H NMR at 298K | ¹³ C NMR(Ag-C) |
|---------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------|
| [Ag(NHC) ₂] ⁺ [AgCl ₂] ⁻ 3.36 | H, 3.55; N, 11.46; Cl, 9.67%. Found: C, 45.71; H, 3.45; N, 11.33; Cl, 9.60%. ¹ H NMR (CDCl ₃ , 400MHz, 298K): 8.10 (d, 2H, J = 8.5 Hz, Quin-H), 8.00 (d, 2H, J = 8.4 Hz, Quin-H), 7.80 (d, 2H, J = 8.1 Hz, CHHC), 7.5 (d, 2H, J = 8.4 Hz, Quin-H), 7.35 (d, 2H, J = 1.7 Hz, Quin-H) 7.15 (d, 2H, J = 1.8 Hz, Quin-H), 6.95 (s, 1H, Quin-H), 5.50 (s, 4H CH ₂), 3.85 (s, 6H, CH ₃). | 181.18 |
| [Ag(NHC)Cl] 3.37 | ¹ H NMR (CDCl ₃ , 400MHz, 298K) : 8.10 (d, 1H, J = 8.4 Hz, Quin-H), 8.00 (d, 1H, J = 8.5 Hz, Quin-H), 7.80 (d, 2H, J = 8.1 Hz, CHHC), 7.70 (t, 1H, J = 1.4 Hz, Quin-H), 7.50 (t, 1H, J = 7.0 Hz, Quin-H), 7.35 (t, 2H, J = 8.7 Hz, Quin-H), 6.90 (s, 3H, Aromatic-H), 5.60 (s, 2H, CH ₂), 2.25 (s, 3H, m-CH ₃), 1.95 (s, 6H, o-CH ₃). | N.A. |
| [Ag(NHC)Cl] 3.38 | ¹ H NMR (CDCl ₃ , 400MHz, 298K): | 184.10 |

Chapter Three: Ag^I and Pd^{II} complexes of quinoline functionalised NHC ligands


| | | |
|-------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
| | 8.05(d, 2H, J = 8.5 Hz, quin-H), 7.65(t, 2H, J = 7.0 Hz, quin-H), 7.50(d, 2H, J = 5.2 Hz, quin-H), 7.10(s, 1H, CHHC), 6.90(s, 1H, CHHC), 5.60(s, 2H, NCH ₂ C), 4.85(m, 1H, NCHC), 1.35(d, 6H, J = 6.8 Hz, CH ₃). | |
| [Ag(NHC)Cl] 3.39 | ¹ H NMR (CDCl ₃ , 400MHz, 298K): 8.00(d, 1H, J = 8.5 Hz, quin-H), 7.90(d, 1H, J = 8.5 Hz, quin-H), 7.60(m, 2H, J = 8.1 Hz, quin-H), 7.55(d, 1H, J = 8.5 Hz, quin-H), 7.40(t, 1H, J = 7.2 Hz, quin-H), 7.00(s, 1H, CHHC), 6.80(s, 1H, CHHC), 5.60(s, 2H, methylene linker-H), 4.10(t, 2H, NCH ₂), 1.70(m, 2H, J = 7.5 Hz, CH ₂), 1.20(m, 2H, J = 7.0 Hz, CH ₂), 0.80(t, 3H, J = 7.3 Hz, CH ₃). | 183.23 |
| [Ag(NHC)Cl] 3.40 | ¹ H NMR (CDCl ₃ , 400MHz, 298K): 8.00(d, 2H, J = 10.7 Hz, quin-H), 7.75(d, 1H, J = 7.1 Hz, quin-H), 7.60(d, 1H, J = 7.5 Hz, arom-H), 7.50(t, 1H, J = 7.3 Hz, arom-H), 7.35(s, 2H, arom-H), 7.25(s, 2H, CHHC), 7.20(d, 1H, J = 6.8 Hz, quin- H), 7.10(t, 2H, J = 8.4 Hz, quin-H), 6.90(d, 1H, J = 11.8 Hz, arom-H), 5.50(s, 2H, NCH ₂ quin), 2.00(s, 6H, CH ₃). | N.A. |
| [Ag(NHC)Cl] 3.41 | ¹ H NMR (CDCl ₃ , 400MHz, 298K): 8.10(d, 2H, J = 8.4 Hz, quin-H), 8.00(d, 2H, J = 8.5 Hz, quin-H), 7.75(d, 2H, J = | 180.37 |

| | | |
|--|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| | <p>8.1 Hz, quin-H), 7.65(t, 2H, J = 5.6 Hz, quin-H), 7.5(t, 2H, J = 7.1 Hz, quin-H), 7.35(d, 2H, J = 8.5 Hz, quin-H), 7.15(s, 2H, HHC), 5.5(s, 4H, NCH₂C).</p> | |
|--|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|

A look at the data presented in Table 3.3 indicate that there is no significant difference between complexes 3.36 -3.41 in terms of the chemical shift of quinoline protons and that of the methylene linkers and the C- Ag carbene resonances were observed between 180.37 -184.10 which are within the range reported in the literature (180-243 ppm) [13]. Elemental analysis of complex **3.38** returned a satisfactory results consistent with the proposed structure as suitable crystal for X-ray crystallography could not be obtained. Complex **3.39** revealed a cluster at M/z = 637.2189 Da (100% intensity) attributable to the cation of biscarbene [Ag(NHC)₂]⁺. The ¹H NMR and ¹³C NMR spectra as well as the elemental analysis of complex **3.40** are consistent with the proposed silver carbene complex. MS analysis revealed a cluster at M/z = 356.1642 Da (100% intensity) and 817.2445 Da (36%) attributable to the cation of the corresponding salt [(NHC)+H]⁺ and biscarbene ion [Ag(NHC)₂]⁺ respectively. The observation of the peak attributable to the bis carbene ion may not necessarily indicate that the complex is indeed a biscarbene. Danopoulos and colleagues reported that silver NHCs with solid-state motifs of C-Ag-X and C-Ag-X₂ formed biscarbenes (C₂-Ag) in the gas phase [17].

3.3.2 Pd(II) Quinoline functionalised (NHC) carbene complexes

Two quinoline based Pd(II) (NHC) complexes were prepared according to steps in Scheme 3.8 by the reaction of quinoline functionalised Ag(I)(NHC) complex with one equivalent of PdCl₂(COD) in DCM, over night, the PdCl₂(COD) being synthesised by standard procedures.



Scheme 3.8: Synthesis of quinoline functionalised Pd(II)(NHC) complexes

Complex **3.42** was characterised by ¹H and ¹³C NMR. Relative to the silver (I) (NHC) complex from which the Pd complex was prepared, the protons were observed to have moved considerably down field indicating that chelation has indeed taken place. Crystal suitable for X-ray crystallography was not obtained but elemental analysis gave satisfactory results in conformity with the proposed structure. Complex **3.43** was characterised by ¹H NMR and elemental analysis. The solubility of the complex in CDCl₃ was not good enough to get a satisfactory ¹³C NMR spectrum. Elemental analysis returned unsatisfactory results.

3.4 Conclusions

Ag^I(NHC) complexes have been valuable intermediates in the preparation of functionalised Pd^{II}(NHC) complexes that are difficult to be accessed via other methodologies. Thus a range of imidazolium salts synthesised in chapter two was reacted with Ag₂O to form the corresponding silver (I) carbene complexes. The syntheses of the silver complexes were confirmed by the absence of C₂-H in the ¹H NMR spectrum, and other analytical techniques.

Compounds **3.36** and **3.37** are the first examples of quinoline functionalised Ag^I(NHC) complexes and were structurally characterised by X-ray

crystallography and formulated as $[\text{Ag}(\text{NHC})_2]^+[\text{AgCl}_2]^-$ and $[\text{Ag}(\text{NHC})\text{Cl}]$ respectively. Results of MS data indicated that complexes **3.38**, **3.39** and **3.40** may have a similar formulation to that of **3.36** because of the presence of cluster attributable to $[\text{Ag}(\text{NHC})_2]$. However the MS is not enough to formulate the formula of Ag^I(NHC) complexes as Danopoulos and colleagues reported that silver NHCs with solid-state motifs of C-Ag-X and C-Ag-X₂ formed biscarbenes (C₂-Ag) in the gas phase [17]. All the Ag^I(NHC) complexes showed stability to air and moisture and were not exposed to light.

In order to extend the synthetic applicability of Ag^I(NHC) complexes as transmetallation agents complexes **3.34** and **3.39** were reacted with PdCl₂·COD in DCM to obtain quinoline functionalised Pd(II)(NHC) complexes (**3.42** and **3.43**) respectively. The formation of the Pd(II)(NHC) complexes was confirmed by ¹H NMR spectrum as relative to the corresponding Ag^I(NHC) complexes, the protons signals of the quinoline moiety of the Pd complexes were observed to have moved down field indicating the chelation between the nitrogen of the quinoline moiety and Pd metal. Attempts were made to synthesise the Pd complexes of by reacting the other described quinoline functionalised Ag^I(NHC) complexes in this work with PdCl₂·COD but a lot of difficulties were encountered in separating the complexes formed from the silver halides.


3.5 Experimental

3.5.1 General comments

All reactions were performed under the atmosphere of dry dinitrogen or argon using standard Schlenk techniques, and solvents were purified and dried by usual means [58], unless otherwise indicated. Imidazolium salts were prepared as detailed in Chapter 2, and all other reagents were used as received. All NMR data are quoted δ/ppm. ¹H and ¹³C (proton decoupled) spectra NMR were recorded on a Bruker DPX Advance 400 (¹H at 400MHz, ¹³C at 100.61 MHz) at ambient temperature, unless otherwise stated, and referenced to SiMe₄. Electrospray mass spectrometry (ESMS) was performed on a VG Fisons Platform II instrument by the department of Chemistry, Cardiff University. Micro analysis was performed by Warwick Analytical Service. All reactions involving silver compounds were performed with the exclusion of light.


3.5.2 The silver (I) complexes of rigid quinoline functionalised mono-NHC ligands

[Ag(1-benzyl-3-quinolineimidazol-2-ylidene)Br] 3.34:



A mixture of 1-benzyl-3-quinolineimidazolium bromide (0.7g, 2mmols) and silver (I) oxide (0.222g, 1mmol in 10ml of DCM) was stirred over night at 40°C. The resulting solution was filtered through celite and the solvent removed under reduced pressure to leave a brown coloured solid. The solid was recrystallised from DCM / Hexane to give the desired Ag complex. 0.55g (60.84%). Anal. Calcd. for C₁₉H₁₅N₃AgBr: C, 48.23; H, 3.17; N, 8.88%. Found: C, 49.45; H, 3.22; N, 8.96%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.85 (d, 1H, J = 1.7 Hz, quin-H), 8.25 (d, 1H, J = 6.7 Hz, quin-H), 7.90 (d, 2H, J = 7.7 Hz, quin-H), 7.55 (t, 1H, J = 7.8 Hz, quin-H), 7.45 (m, 2H, J = 3.0 Hz, CHHC), 7.25-7.35 (m, 5H, J = 5.4 Hz, Ar-H), 7.10 (s, 1H, quin-H), 5.35 (s, 2H, CH₂-Ph). ¹³C NMR (CDCl₃, 100 MHz, 298K): 151.75, 142.86, 136.99, 135.60, 130.04, 129.68, 129.60, 129.19, 128.59, 127.54, 126.72, 125.74, 122.77, 120.64, 56.49(CH₂).

[Ag(1-methyl-3-quinolineimidazol-2-ylidene)I] 3.35:




A mixture of 1-methyl-3-quinolineimidazolium iodide (0.44g, 1.31mmol) and silver (I) oxide (0.15g, 0.66mmol) in 10ml of CH₃CN was stirred over night at 40°C. The resulting solution was filtered to remove silver halide through celite and the solvent removed under reduced pressure to leave a tan coloured solid. The solid was then recrystallised from DCM/Hexane to give the desired Ag

complex (0.26g, 63%). ¹H NMR (CDCl₃, 400MHz, 298K): 8.85 (d, 1H, J = 2.7 Hz, Quin-H), 8.25(d, 1H, J = 7.3 Hz, quin-H), 7.95 (d, 1H, J = 8.2 Hz, quin-H), 7.80(d, 1H, J = 7.4 Hz, quin-H), 7.5 (t, 1H, J = 7.6 Hz, quin-H), 7.4 (d, 1H, J = 4.2 Hz, quin-H), 7.35 (s, 2H, CHHC). 3.90 (s, 3H, CH₃). ¹³C NMR (d₂-DCM, 100 MHz, 298K): 163.00, 151.3, 147.6, 145.2, 139.4, 138.5, 136.9, 134.7, 133.2, 131.8, 44.0(CH₃).


3.5.3: Silver (I) complexes of methylene bridged quinoline functionalised NHC ligands

[Ag (1-methy-3-(2-methylquinoline) imidazolin-2-ylidene) ₂][AgCl₂] 3.36




A mixture of 1-methyl-3-(2-methylquinoline) imidazolium chloride (0.30g, 1.16mmol) and Ag₂O (0.14g, 0.59mmol) in dichloromethane was stirred for 12 hours. The solution was filtered through celite and the filtrate concentrated. Diethyl ether was added to precipitate the carbene complex which was repeatedly washed with diethyl ether and dried under vacuum to give the desired silver carbene complex. Crystals suitable for X- ray structure were grown by layering diethyl ether on dichloromethane. Anal. Calcd. for C₁₄H₁₃N₃AgCl: C, 45.86; H, 3.55; N, 11.46; Cl, 9.67%. Found: C, 45.71; H, 3.45; N, 11.33; Cl, 9.60%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.10 (d, 2H, J = 8.5 Hz, Quin-H), 8.00 (d, 2H, J = 8.4 Hz, Quin-H), 7.80 (d, 2H, J = 8.1 Hz, CHHC), 7.5 (d, 2H, J = 8.4 Hz, Quin-H), 7.35 (d, 2H, J = 1.7 Hz, Quin-H) 7.15 (d, 2H, J = 1.8 Hz, Quin-H), 6.95 (s, 1H, Quin-H), 5.50 (s, 4H CH₂), 3.85 (s, 6H, CH₃). ¹³C NMR (CDCl₃, 100MHz, 298K): 181.18 (C-Ag), 155.45, 148.09, 138.12, 130.55, 129.62, 128.11, 127.92, 127.50, 123.01, 122.31, 120.20, 58.14, 39.27. MS (HRMS, Da): M/z (224.09) [(NHC)+H]⁺ (100%).

Synthesis of [Ag (1-mesityl-3-(2-methylquinoline) imidazolium-2-ylidene) Cl] 3.37



A mixture of 1-mesityl 3-(2-methylquinoline) imidazolium chloride 4 (0.50g, 1.38mmol) and Ag₂O in dichloromethane was stirred at room temperature for 12 hours. The solution was filtered through celite and the filtrate concentrated. Diethyl ether was added to precipitate the carbene complex which was repeatedly washed with diethyl ether and dried under vacuum to give the desired silver carbene complex. Crystals suitable for X-ray structure were grown by layering diethyl ether on dichloromethane. Anal. Calcd. for C₂₂H₂₁N₃AgCl: C, 56.13; H, 4.46; N, 8.93; Cl, 7.55%. Found: C, 56.23; H, 4.53; N, 8.77; Cl, 7.55%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.10 (d, 1H, J = 8.4 Hz, Quin-H), 8.00 (d, 1H, J = 8.5 Hz, Quin-H), 7.80 (d, 2H, J = 8.1 Hz, CHHC), 7.70 (t, 1H, J = 1.4 Hz, Quin-H), 7.50 (t, 1H, J = 7.0 Hz, Quin-H), 7.35 (t, 2H, J = 8.7 Hz, Quin-H), 6.90 (s, 3H, Aromatic-H), 5.60 (s, 2H, CH₂), 2.25 (s, 3H, m-CH₃), 1.95 (s, 6H, o-CH₃). ¹³C NMR (CDCl₃, 100 MHz, R.T.): 155.40, 148.0, 139.20, 138.18, 135.08, 130.59, 129.84, 129.64, 128.16, 127.95, 127.50, 123.57, 122.29, 119.74, 58.16, 21.47, 18.11. MS(HRMS, Da): M/z (328.15) [(NHC)+H]⁺ (100%).


[Ag (1-isopropyl-3-(2-methylquinolin) imidazolium-2-ylidene) I] 3.38



Following the method for the synthesis of 3.37, compound 3.38 was synthesized from 1-isopropyl-3-(2-methylquinoline) imidazolium iodide (0.10g, 0.26mmol) and Ag₂O(32mg, 0.14mmol) in dichloromethane.(Yield =0.12g, 73.00%). Anal. Calcd. for C₁₆H₁₇N₃AgI: C, 39.52; H, 3.50; N, 8.65; I, 7.55%. Found: C, 39.70; H, 3.42; N, 8.57; I, 25.92%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.05(d,


2H, J = 8.5 Hz, quin-H), 7.65(t, 2H, J = 7.0 Hz, quin-H), 7.50(d, 2H, J = 5.2 Hz, quin-H), 7.10(s, 1H, CHHC), 6.90(s, 1H, CHHC), 5.60(s, 2H, NCH₂C), 4.85(m, 1H, NCHC), 1.35(d, 6H, J = 6.8 Hz, CH₃). ¹³C NMR (CDCl₃, 100 MHz, R.T.): 184.10(C-Ag), 156.55, 137.91, 130.18, 129.47, 128.12, 127.91, 121.71, 121. MS (HRMS, Da): M/z = 609.1886 [Ag(NHC)₂]⁺ (100%).

[Ag(1-n-butyl-3-(2-methylquinolin)imidazolin-2-ylidene)I] 3.39



Following the method for the synthesis of **3.38**, compound **3.39** was synthesized from 1-nbutyl-3-(2-methylquinoline)imidazoliumiodide (0.50g, 1.27mmol) and Ag₂O(0.15g, 0.64mmol) in dichloromethane.(Yield = 0.40g, 62.50%). Required for C₁₇H₁₉N₃AgI: C, 40.82; H, 3.80; N, 8.40%. Found: C, 39.90; H, 3.67; N, 7.79%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.00(d, 1H, J = 8.5 Hz, quin-H), 7.90(d, 1H, J = 8.5 Hz, quin-H), 7.60(m, 2H, J = 8.1 Hz, quin-H), 7.55(d, 1H, J = 8.5 Hz, quin-H), 7.40(t, 1H, J = 7.2 Hz, quin-H), 7.00(s, 1H, CHHC), 6.80(s, 1H, CHHC), 5.60(s, 2H, methylene linker-H), 4.10(t, 2H, NCH₂), 1.70(m, 2H, J = 7.5 Hz, CH₂), 1.20(m, 2H, J = 7.0 Hz, CH₂), 0.80(t, 3H, J = 7.3 Hz, CH₃). ¹³C NMR (CDCl₃, 100 MHz, R.T.): 183.23(C-Ag), 155.02, 146.52, 136.50, 128.80, 128.07, 126.70, 126.50, 126.14, 125.76, 120.30, 119.96, 64.84, 56.56, 50.68, 32.50, 18.75. MS (HRMS, Da): M/z (637.2189) [Ag(NHC)₂]⁺ (100%)


[Ag (1-(2-methylpropiophenone)-3-(2-methylquinolin) imidazolin-2-ylidene) Cl] 3.40.



Following the method for the synthesis of **3.39**, compound **3.40** was synthesized from 1-(2-methylpropiophenone)-3-(2-methylquinolin) imidazolium chloride (0.30g, 0.77mmol) and Ag₂O (90mg, 0.39mmol) in dichloromethane. (Yield = 0.32g, 90.00%). Anal. Calcd. for C₂₃H₂₁N₃OAgCl: C, 55.38; H, 4.14; N, 8.43%. Found: C, 55.78; H, 4.18; N, 8.22%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.00(d,

2H, $J = 10.7$ Hz, quin-H), 7.75(d, 1H, $J = 7.1$ Hz, quin-H), 7.60(d, 1H, $J = 7.5$ Hz, arom-H), 7.50(t, 1H, $J = 7.3$ Hz, arom-H), 7.35(s, 2H, arom-H), 7.25(s, 2H, CHHC), 7.20(d, 1H, $J = 6.8$ Hz, quin-H), 7.10(t, 2H, $J = 8.4$ Hz, quin-H), 6.90(d, 1H, $J = 11.8$ Hz, arom-H), 5.50(s, 2H, NCH₂quin), 2.00(s, 6H, CH₃). ¹³C NMR (CDCl₃, 100MHz, 298K): 196.71(PhCOC), 153.72, 146.53, 136.77, 136.59, 131.72, 129.18, 129.12, 128.17, 127.94, 127.60, 126.72, 126.66, 126.46, 126.12, 120.91, 118.72, 118.43, 118.38, 66.99, 57.61, 27.56. MS (HRMS, Da): M/z (356.1642) [(NHC)+H]⁺ (100%), 817.2445 [Ag(NHC)₂]⁺ (36%).


[Ag (bis-1, 3(2-methylquinolin) imidazolin-2-ylidene) Cl] 3.41



Following the method for the synthesis of **3.40**, compound **3.41** was synthesized from bis-1, 3-(2-methylquinoline) imidazolium chloride (0.30g, .78mmol) and Ag₂O (93mg, 0.40mmol) in 20 ml of dichloromethane. (Yield = 0.12g, 40%). Anal. Calcd. for C₂₃H₁₈N₄AgCl: C, 55.94; H, 3.65; N, 11.35%. Found: C, 56.39; H, 3.61; N, 11.21%. ¹H NMR (CDCl₃, 400MHz, 298K): 8.10(d, 2H, $J = 8.4$ Hz, quin-H), 8.00(d, 2H, $J = 8.5$ Hz, quin-H), 7.75(d, 2H, $J = 8.1$ Hz, quin-H), 7.65(t, 2H, $J = 5.6$ Hz, quin-H), 7.5(t, 2H, $J = 7.1$ Hz, quin-H), 7.35(d, 2H, $J = 8.5$ Hz, quin-H), 7.15(s, 2H, HHC), 5.5(s, 4H, NCH₂C). ¹³C NMR (CDCl₃, 100 MHz, 298K): 180.37(C-Ag), 154.09, 146.58, 136.71, 129.08, 128.09, 126.70, 126.47, 126.01, 121.31, 118.75, 56.74. MS (HRMS, Da): M/z (351.16) [(NHC)+H]⁺ (100%).

3.5.4: Pd(II) quinoline functionalised NHC complexes


[Pd(1-benzyl-3-quinoline-imidazolin-2-ylidene)Cl₂] 3.42



A solution of PdCl₂(COD) (0.092g, 0.32 mmol) in 10ml DCM was added to solution [Ag(1-benzyl-3-quinolineimidazolin-2-ylidene)Br] (0.30g, 0.32mmol) in 10mL DCM and stirred at room temperature for 2 hours. The solution was

then filtered through celite to remove the silver bromide formed and the solvent was concentrated to 5ml. Hexane was added to precipitate the Pd carbene complex which was repeatedly washed with hexane to any 1,5-cyclooctadiene. Anal. Calcd. for C₁₉H₁₅N₃PdCl₂: C, 49.31; H, 3.24; N, 9.08%. Found: C, 46.08; H, 3.30; N, 7.78%. ¹H NMR (CDCl₃, 400MHz, 298K): 9.70(d, 1H, quin-H), 8.40(d, 1H, J = 1,4 Hz, quin-H), 8.00(d, 1H, J = 6.7 Hz, quin-H), 7.90 (d, 1H, J = 7.2 Hz, quin-H), 7.70 (t, 1H, J = 7.9.5 Hz, quin-H), 7.50(t, 1H, J = 6.3 Hz, quin-H), 7.40(d, 2H, J = 6.1 Hz, Ar-H), 7.35(s, 1H, CHHC), 7.30 (m, 3H, J = 7.7 Hz, Ar-H), 6.95(s, 1H, CHHC), 5.25(s, 2H, Ph-CH₂)

[Pd(1-nbutyl-3-(2-methylquinoline)imidazolin-2-ylidene)Cl₂] 3.43



1-nbutyl-3-(2-methylquinoline)imidazolium (0.15g, 0.38mmol) and Ag₂O (44mg, 0.2mmol) in 15mL of DCM were stirred over night and the reaction mixture was filtered via cannula to a solution PdCl₂·COD (0.11g, 0.38mmol) in 15mL DCM and the reaction mixture stirred overnight. The reaction mixture was filtered through celite and the filtrate was concentrated over vacuum.

Hexane was added to precipitate the desired Pd complex which repeated recrystallised from hot CH₃CN and diethyl ether. Anal. Calcd. for C₁₇H₁₉N₃PdCl₂: C, 46.21; H, 4.30; N, 9.51%. Found: C, 43.92; H, 4.05; N, 6.72%. ¹H NMR (DMSO, 400MHz, R.T): 9.20(d, 1H, J = 8.8 Hz quin-H), 8.70(d, 1H, J = 8.2 Hz quin-H), 8.10(d, 1H, J = 8.0 Hz, quin-H), 7.90(t, 2H, J = 5.2 Hz, Quin-H), 7.70(t, 1H, J = 7.5 Hz quin-H), 7.60(s, 1H, CHHC), 6.35(s, 1H, CHHC), 6.10(d, 1H, J = 15.6 Hz methylene linker-H), 5.80(d, methylene J = 15.7 Hz linker-H), 4.60(m, 1H, J = 6.6 Hz NCH₂), 4.10(m, 1H, J = 6.6 Hz NCH₂), 2.90(m, 2H, J = 7.0 Hz CH₂), 1.15(m, 2H, J = 8.5 Hz, CH₂), 0.80(t, 3H, J = 7.3 Hz, CH₃).

3.5.4 X-Ray crystallography

Standard conditions as outlined in section 2.4.7.

Table 3.4: Crystal data and structure refinement for 3.36

| | | |
|-----------------------------------|-----------------------------------------------------|-----------------|
| Empirical formula | C ₁₄ H ₁₃ AgN ₃ Cl | |
| Formula weight | 366.59 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Monoclinic | |
| Space group | C2 | |
| Unit cell dimensions | a = 15.9530(14) Å | α = 90.00° |
| | b = 6.5010(8) Å | β = 111.937(8)° |
| | c = 13.8760(13) Å | γ = 90.00° |
| Volume | 1334.90(2) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.8424 Mg/m ³ | |
| Absorption coefficient | 1.698 mm ⁻¹ | |
| F000 | 728 | |
| Crystal size | 0.30 x 0.15 x 0.01 mm ³ | |
| Theta range for data collection | 2.61 to 27.40 ° | |
| Index ranges | -19 ≤ h ≤ 20, -8 ≤ k ≤ 7, -17 ≤ l ≤ 16 | |
| Reflections collected | 2473 | |
| Independent reflection | 2473 [R _{int} = 0.0714] | |
| Completeness of theta = 27.4° | 93.20% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.6299 and 0.9832 | |
| Refinement method | Full matrix least square on F ² | |
| Data/restraints/parameters | 2473/144/133 | |
| Goodness of fit on F ² | 1.094 | |
| Final R indices [I > 2σ(I)] | R1 = 0.1141, wR2 = 0.2350 | |
| R indices (all data) | R1 = 0.1015, wR2 = 0.2261 | |
| Largest diff. peak hole | 1.763 and -1.552 e Å ⁻³ | |

Table 3.5: Crystal data and structure refinement for 3.37

| | | |
|--------------------------------------------|-----------------------------------------------------|------------------------------|
| Empirical formula | C ₂₂ H ₂₁ AgN ₃ Cl | |
| Formula weight | 470.75 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Triclinic | |
| Space group | P -1 | |
| Unit cell dimensions | a = 8.9956(3) Å | α = 79.2749(17) ^o |
| | b = 9.3915(3) Å | β = 81.9409(16) ^o |
| | c = 12.4646(4) Å | γ = 74.7489(16) ^o |
| Volume | 993.57(6) Å ³ | |
| Z | 2 | |
| Density (calculated) | 1.573Mg/m ³ | |
| Absorption coefficient | 1.160mm ⁻¹ | |
| F000 | 476 | |
| Crystal size | 0.20 x 0.20 x 0.20 mm ³ | |
| Theta range for data collection | 1 to 27 ^o | |
| Index ranges | -10<=h<=11, -12<=k<=12, -16<=l<=16 | |
| Reflections collected | 24373 | |
| Independent reflection | 4440 [R _{int} = 0.145] | |
| Completeness of theta = 25.18 ^o | 99.30% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.7900 and 0.7900 | |
| Refinement method | Full matrix least square on F ² | |
| Data/restraints/parameters | 3644/0/244 | |
| Goodness of fit on F ² | 1.0042 | |
| Final R indices [I>2 sigma(I)] | R1 = 0.0418, wR2 = 0.11129 | |
| R indices (all data) | R1 = 0.0505, wR2 = 0.1206 | |
| Largest diff. peak hole | 0.69 and -1.07e Å ⁻³ | |

3.6 References

1. A. J. Arduengo III, H. V. R. Dias, J. C. Calabrese and F. Davidson, *Organometallics*, **1993**, *12*, 3405-3409.
2. A. Caballero, E. Diez-Barra, F. A. Jalon, S. Merino, J. Tejada, *Organomet.Chem.* **2001**, *395*, 617-618.
3. A. Caballero, E. Diez-Barra, F. A. Jalon, S. Merino, A. M. Rodriguez, J. Tejada, *J. Organomet. Chem.* **2001**, *263*, 627.
4. M. A. Fox, M. F. Mahon, N. J. Patmore, A. S. Weller, *Inorg. Chem.* **2002**, *41*, 4567.
5. M. C. Chung, *Bull. Korean Chem. Soc.* **2002**, *23*, 921.
6. I. J. B. Lin, C.S. Vasam, *Inorg. Chem.*, **2004**, *25*, 75.
7. R. -Z. Ku, -C. Huang, J. -Y. Cho, F. -M.Kiang, K. J. Reddy, Y. -C.Chen, K. -J. Lee, J. -H. Lee, G. -H. Lee, S. -M. Peng and S. -T. Liu, *Organometallics*, **1999**, *18*, 2145-2154.
8. O. Guerret, S. Sole, H. Gornitzka, G. Trinquier and G. Bertrand, *J. Organomet.Chem.*, **2000**, *600*, 112-117.
9. H. M. J. Wang and I. J. B. Lin, *Organometallics*, **1998**, *17*, 972-975.
10. J. C. Garrison, R. S. Simons, C. A. Tessier, W. J. Youngs, *J. Organomet. Chem.* **2003**, *673*, 1.
11. A. Kascatan- Nebioglu, M. J. Panzner, J. C. Garrison, C. A. Tessier, W. J. Youngs, *Organometallics*, **2004**, *23*, 1928.
12. C. A. Quezada, J. C. Garrison, M. J. Panzner, C. A. Tessier, W. J. Youngs, *Organometallics*, **2004**, *23*, 4846.
13. J. C. Garrison, W. J. Youngs, *Chem. Rev.* **2005**, *25*, 3978.
14. C. K. Lee, J. C. C. Chen, K. M. Lee, C. W. Liu and I. J. B. Lin, *Chem Mater*, **1999**, *11*, 1237-1242.
15. B. Bildstein, M. Maulaun, H. Kopacka, K. Wurst, Mitterbock, K. -H. Ongania, G. Opromolla and P. Zanello, *Organometallics*, **1999**, *18*, 4325- 4336.
16. D. S. McGuinness and K. J. Cavell, *Organometallics*, **2000**, *19*,741- 748.
17. A. A. D. Tulloch, A. A. Danopoulos, S. Winston, S. Kleinhenz and G. Eastham, *J. Chem. Soc., Dalton Trans.*, **2000**, 4499-4506.

18. A. M. Magill, D. S. McGuinness and K. J. Cavell, G. J. P. Britovsek, V. C. Gibson, A. J. P. White, D. J. Williams, A.H. White and B. W. Skelton, *J.Organomet. Chem.*, **2001**, 546, 617-618.
19. J. Pytkowicz, S. Roland and P. Mangeney, *J. Organomet. Chem.*, **2001**, 631,157-163.
20. C. K. Lee, K. M. Lee and I. J. B. Lin, *Organometallics*, **2002**, 21, 10-12.
21. W. Chen, B. Wu and K. Matsumoto, *J. Organomet. Chem.*, **2002**, 654, 233-236.
22. K. M. Lee, H. M. J. Wang and I. J. B. Lin, *J. Chem. Soc., Dalton Trans.*,**2000**, 2852-2856.
23. H. M. J. Wang, C. Y. L. Chen and I. J. B. Lin, *Organometallics*, **1999**, 18, 1216-1223.
24. J. C. Garrison, R. S. Simons, J. M. Tally, C. Wesdemiotis, C. A. Tessier and Youngs, *Organometallics*, **2001**. 20, 1276.
25. D. J. Nielsen, K.J. Cavell, B. W. Skelton and A. H. White, *Organometallics*, **2001**, 20, 995.
26. D. J. Nielsen, K.J. Cavell, B. W. Skelton and A. H. White, *Inorg. Chim. Acta*, **2002**, 327, 116.
27. D. J. Nielsen, K.J. Cavell, B. W. Skelton and A. H. White, *Inorg. Chim. Acta*, **2003**, 352, 143-150.
28. D. J. Nielsen, PhD Thesis, University of Tasmania, March **2004**.
29. D. J. Nielsen, K. J. Cavell, B. W. Skelton, A. H. White, *Inorg. Chim. Acta* **2003**, 143, 352.
30. A.R.Chianese, B. M. Zeglis, R. H. Crabtree, *Chem. Commun.* **2004**, 2176.31. P. L. Chiu, H. M. Lee, *Organometallics* **2005**, 24, 1692.
32. C. Boehme and G. Frenking, *Organometallics*, **1998**, 17, 5801-5809.
33. X. W. S. Liu, L. Weng, G. Jin, *Organometallics*, **2006**, 25, 3565-3569.
34. R. Corberan, J. Rammirez, M. Poyatos, E. Fernandez, *Tetrahedron: Asymmetry* **2006**, 17,1759.
35. A.C. Sentman, S. Csihony, R. M. Waymouth, J. L. Hedrick, *J. Org. Chem.***2005**, 70, 2391.
36. M. K. Samantaray, V. Katiyar, D. Roy, K.Pang, H. Nanavati, R. Stephen, R.B. Sunoj, P. Ghosh, *Eur. J. Inorg. Chem.* **2006**, 2006, 2975.

37. X. Wang, S. Liu, L. -H, Weng, G. -X, Jin, *Organometallics* **2006**, *25*, 3565.
38. M. Tilset, O. Andell, A. Dhindsa and M. Froseth, Borealis Tech Oy: WO **2002**, 0249758.
39. J. Pytkowicz, S. Roland and P. Mangeney, *Tetrahedron: Asymmetry*, **2001**, *12*, 2087-2089.
40. A. A. D. Tulloch, A. A. Danopoulos, G. J. Tizzard, S. J. Coles, M. B. Hursthouse, R. S. Hay- Motherwell and W. B. Motherwell, *Chem. Commun.* **2001**, 1270.
41. S. McGrandle, G. C. Saunders, *J. Fluor. Chem.* **2005**, *126*, 449.
42. J. Schwarz, V. P. W. Bohm, M. G. Gardner, M. Grosche, W. A. Herrmann, W. Hieringer and G. Rausdaschl-Sieber, *Chem. Eur. J.*, **2006**, 1773.
43. J.C.C.Chen and I.J. B. Lin, *Organometallics*, **2000**, *19*, 5113.
44. R. A. Batey, T. D. Quach, M. Shen, A. N. Thadani, D. V. Smil, S. -W. Li. and D. B. Mackay, *Pure appl. Chem.*, **2002**, *74*, 43-55.
45. R. A. Batey, M. Shen and A. J. Lough, *Org. Lett.*, **2002**, *4*, 1411-1414.
46. W. A. Herrmann, L. J. Goossen and M. Spiegler, *J. Organomet. Chem.*, **1997**, *547*, 357-366.
47. H. Glas, E. Herdtweck, M. Spiegler, A. -K.Pleier and W. R. Thiel, *J. Organomet. Chem.*, **2001**, *626*, 100-105.
48. N. Kuhn, C. Mossmer-Maichle, E. Niquet and I. Walker, *Z. Naturforsch.*, **2002**, *57B*, 47-52.
49. W. A. Herrmann, L. J. Goossen and M. Spiegler, *Organometallics*, **1998**, *17*, 2162.
50. A. A. D. Tulloch, A. A. Danopoulos, R.P. Tooze, S. M. Cafferkey, S. Kleinhenz and M. B. Hursthouse, *Chem. Commun.*, **2000**, 1247- 1248.
51. S. Grundemann, M. Albrecht, A. Kovacevic, J. W. Faller and R. H. Crabtree, *J. Chem. Soc. Dalton Trans.*, **2002**, 2163-2167.
52. J. A. Loch, M. Albrecht, E. Peris. J. Mata, J. W. Faller and R. H. Crabtree, *Organometallics*, **2001**, *21*, 700-706.
53. W. A. Herrmann, M. Ellison, J. Fischer, C. Kocher, G. Artus, *Angew. Chem. Int. Ed. Eng.*, **1995**, *34(12)*, 2371-2374.
54. N.Marsich, G.Pellizer, A. Camus, A. M. M. Lanfredi, F. Ugozzoli,

Inorg. Chim. Acta **1990**, *169*, 171.


55. L.G. Bonnet, R. E. Douthwaite, R. Hogson, J. Houghton, B. M. Kariuki, S. Simonovic, *Dalton Trans.* **2004**, 3528.
56. V. Cesar. S. Bellemin-Lsponnaz, L. H. Gade, *Organometallics*, **2002**, *21*, 5204.
57. M. Paas, B. Wibbeling, R. Frohlich and F. E. Hahn, *Eur. J. Inorg. Chem.* **2006**, 158-162.
58. W. L. F. Armarego and D. D. Perrin, *Purification of Laboratory Chemicals*, 4th ed., Butterworth Heinemann: Bath, **1996**.

CHAPTER FOUR


Rhodium(I) and Iridium(I) complexes of quinoline functionalised Heterocyclic Carbene ligands

4.1 Introduction


Prior to the isolation of the first stable crystalline NHC **4.5** in 1991 by Arduengo et al [1], metal complexes of N-heterocyclic carbenes (NHCs) were reported concurrently in 1968 by Ofele **4.1** [2] and by Wanzlick and Shonherr **4.2** [3] both being prepared directly from imidazolium salts. In his contribution Lappert prepared a wide range of transition metal- NHC compounds including **4.3** and **4.4** from electron rich olefins in 1970 [4] and complex **4.3** was found to be an active catalyst for hydrosilylation.




4.1




4.2



4.3



4.4




4.5

Figure 5.1: Early metal complexes of NHCs **4.1- 4.4** and first crystalline free Carbene **4.5**

Encouraged by the promise of these early applications utilising Rh(I) complexes of simple monodentate NHC ligands, there was a natural evolution to expand the catalytic applications of Rh and Ir NHC complexes systems


through modification of the NHC ligands. The desired qualities of modified NHC ligands include the incorporation of functional groups leading to easily recoverable catalyst, water or methanol soluble catalyst, and catalyst containing flexible steric bulk as well as chiral and bidentate and pincer ligands [5] and combination of the strongly bound NHC moiety with more weakly nucleophilic functional groups to furnish ligands with hemilabile donor groups as these can increase catalytic activities by stabilising the low valent centres formed during catalysis. These functional groups may be incorporated at one or both of the ring nitrogens to give access to bidentate or tridentate NHC ligands.

Examples of Rh(I) and Ir(I) complexes of functionalised NHC ligands are limited but have increased considerably in the past few years. Functionalised Rh(I) and Ir(I) (NHC) complexes have appeared in the literature with picolyl (4.5 and 4.6) [6], 4.17 and 4.18[11], ether, ester, amide and ketone(4.7, 4.8, 4.9 and 4.10) [7], amino (4.11 , 4.12) [8], pyridyl (4.13, 4.14, 4.15, 4.16 [9], imino 4.19 [10] Figure 4.2. As this thesis was nearing completion a publication by Webster et al reported quinoline functionalised NHC complexes of rhodium and iridium.



4.5: M = Rh

4.6: M = Ir





4.7: R¹, R² = ether


4.8: R¹, R² = ester

4.9: R¹, R² = amide

4.10: R¹, R² = ketone




4.12



4.13: M = Rh

4.14: M = Ir



4.15: M = Rh

4.16: M = Ir

carbene ligands to other metals such Pd [14, 15], Au [16] including Rh and Ir [6, 7, 8, 9, 10, 11, 16]. Realising the simple way of obtaining our desired Rh(I) and Ir(I) through transmetallation of silver carbene, all the Rh(I) and Ir(I) presented herein were prepared using the silver carbene protocol. However it is worth noting that the reaction yield via silver carbene is depended on the imidazolium salt/ metal ratio and the starting metal complex used. Mas-Marza et al [6] reported that Rh(III) and Ir(III) NHC can also be prepared by the transmetallation, as the silver carbene can play dual role (i) in NHC transfer and (i) as oxidizing agent.

In summary, this chapter presents work pertaining to Rh(I) and Ir(I) NHC complexes of quinoline functionalised imidazolin-2-ylidene ligands and includes a review of literature pertaining to this work.


The synthesis and characterisation of a series of [Rh(COD)Cl] NHC and [Ir(COD)Cl] NHC complexes of quinoline functionalised via $\text{Ag}^{\text{I}}(\text{NHC})$ is outlined. The ligands have range of steric bulk at their 1, 3 positions. Also described are chelated complexes of the corresponding [Rh(COD)Cl] NHC and [Ir(COD)Cl]NHC complexes obtained by treatment of the neutral complexes with AgBF_4 , although only one such of complexes is presented because of solubility problems encountered in the characterisation of other complexes. All the Rh (I) and Ir NHC complexes obtained are found to be stable to moisture and air.

4.2 Results and Discussions

4.2.1: Synthesis and characterisation of quinoline functionalised Rh(I) NHC complexes.

The use of free carbene route by deprotonation of the imidazolium salts with strong base, which was expected to produce the corresponding free carbene, was not successful presumably because of interference from deprotonation of benzylic methylene protons [17]. Therefore, all the Rh(I) NHC complexes presented in this work were prepared from the $\text{Ag}^{\text{I}}(\text{NHC})$ complexes presented in chapter 3. Treatment of $[\text{Rh}(\text{COD})\text{Cl}]_2$ with the methylene bridged quinoline functionalised $\text{Ag}^{\text{I}}(\text{NHC})$ complexes in dichloromethane at ambient temperature

gave the desired rhodium carbene complexes as yellow solids in good yield after work up as shown in Scheme 4.1 below. All the prepared rhodium complexes were characterised by ^1H NMR, ^{13}C NMR and mass spectroscopy, elemental analysis and X-ray crystallography. The ^{13}C NMR data for the coordinating carbene carbons for complexes **4.20**, **4.22**, **4.23** and **4.24** appear at δ values of 181.60, 180.38, 178.66 and 183.45 ppm respectively, suggesting the formation of the Rh-C bond which are in the usual range for other Rh(I)-NHC complexes [18, 19, 20-22]. The carbenic carbon in complex **4.21** was not observed in the ^{13}C NMR spectrum. The ^1H NMR shifts corresponding to the protons of the quinoline ring are essentially similar to those of the silver complexes from which they were made, indicating that the quinoline nitrogen donor remains uncoordinated. Furthermore, ^1H NMR spectra show diastereotopic protons for the CH_2 linker ($\delta = 6.5$ and 5.6 for complex **4.20**, 6.5 and 6.1 for complex **4.21**, 6.5 and 5.75 for complex **4.22**, 6.5 and 5.6 for complex **4.23** and 6.5 and 5.8 for complex **4.24**) which suggests that this group is out of the coordination plane of the molecule thus reducing its symmetry.



3.36: R = Me, X = Cl

3.37: R = Mes, X = Cl

3.38: R = iPr, X = I

3.39: R = n-Bu, X = I

Bu

3.41: R = QuinCH₂, X = Cl

QuinCH₂

4.20: R = Me

4.21: R = Mes

4.22: R = iPr

4.23: R = n-

4.24: R =

Scheme 4.1: Synthesis of quinoline functionalised Rh(I) NHC complexes.

Elemental analysis results for complex **4.20** were satisfactory but no reasonable data could be obtained from the mass spectrum. Suitable crystals for X-ray single crystal diffraction were obtained from a DCM/ n-pentane solution at ambient temperature. The detailed solid state coordination sphere around the rhodium centre of complex **4.20** is confirmed by the X-ray crystal structural analysis. The complete molecular structure of complex **4.20** is depicted in Figure 4.3 below. Selected bond distances and bond angles are listed in table 4.1




Figure 4.3: ORTEP projection of complex **4.20** excluding hydrogen atoms for clarity, showing atom labelling scheme.

Table 4.1: Selected bond lengths (Å) and bond angles (°) of **4.20**

| | | | | | |
|-----------|----------|--------------|-----------|----------------|-----------|
| C1 - N1 | 1.356(6) | C16 - Rh1 | 2.126(5) | N1 - C1 - Rh1 | 131.4 (4) |
| C1 - Rh1 | 2.028(5) | C19 - Rh1 | 2.206(5) | C1 - Rh1 - C11 | 87.99(14) |
| C1 - N2 | 1.362(7) | Rh1 - C11 | 2.386(13) | N2 - C5 - C6 | 113.90(4) |
| N2 - C3 | 1.394(9) | C20 - Rh1 | 2.233(6) | C1 - Rh1 - C15 | 92.00(2) |
| C15 - Rh1 | 2.123(5) | N1 - C1 - N2 | 103.8(4) | C1 - N2 - C3 | 111.00(4) |

The structural arrangement of complex **4.20** shows that the molecular geometry around the rhodium ion is a square planar arrangement with two coordination

sites occupied by carbene and chloride in a *cis* fashion. The distances of Rh-C(COD) *trans* to the carbene donor appear to be longer than those in the *cis* arrangement, suggesting that the σ -donor nature of the diaminocarbene is stronger than that of the chloride. No major deviations were observed in the bond lengths and bond angles of complex **4.20** compared with those reported in the literature [23-25].

The ^1H NMR shifts for complex **4.21** corresponding to the protons of the quinoline ring are essentially similar to those of the silver complexes from which they were made, indicating that the quinoline nitrogen donor remains uncoordinated with the carbenic carbon not observed in the ^{13}C NMR spectrum. The high resolution MS measurement were consistent with the proposed formulation with $[\text{M}-\text{Cl}]^+$ observed at $M/z = 538.18$ Da (100% intensity). Crystals suitable for X-ray crystallography were obtained for complex **4.21** by diffusion of *n*-pentane into the DCM solution of the compound enabling elucidation of the solid state structure as depicted in Figure 4.4. Selected bond distances and bond angles are listed in table 4.2




Figure 4.4: ORTEP projection of complex **4.21** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 4.2: Selected bond lengths (Å) and bond angles (°) of **4.21**.

| | | | | | |
|----------|----------|-----------|-----------|-------------|-----------|
| C1 - N1 | 1.355(5) | C24 - Rh1 | 2.179(4) | N1-C1-Rh1 | 130.8 (4) |
| C1 - Rh1 | 2.045(5) | C27- Rh1 | 2.123(4) | C1-Rh1- Cl1 | 87.67(11) |
| C1- N2 | 1.355(5) | Rh1- Cl1 | 2.3865(9) | C1- N1- C4 | 124.3(3) |
| N2 - C3 | 1.379(9) | C28 - Rh1 | 2.112(4) | C1- Rh1-C28 | 94.63(2) |
| C23 -Rh1 | 2.202(4) | N1- C1-N2 | 103.7(3) | C24-Rh1-Cl1 | 90.55(14) |

Elemental analysis of complex **4.21** returned satisfactory results. There are no significant deviations in terms of the bond lengths and bond angles of complex **4.21** with complex **4.20** and all are within the range reported in the literature [20-22, 23,].

Complex **4.22** was characterised by spectroscopic analysis, elemental analysis and X- ray crystallography. The ^1H NMR spectrum for complex **4.22** displays similar pattern with that of complexes **4.20** and **4.21**. The synthesis of complex was confirmed by ^{13}C NMR, which reveals Rh-C resonance at a σ value of 180.38 ppm. The high resolution MS measurement was consistent with the proposed formulation with $[\text{M}- \text{Cl}]^+$ observed at $M/z = 462.14$ Da (100% intensity) and elemental analysis results were in agreement with the proposed formulation. Crystals suitable for X-ray structural determination were obtained by diffusion of n-pentane into the saturated DCM solution of the compound and Ortep crystal structure is as depicted in Figure 4.5. Selected bond distances and bond angles are listed in table 4.3.




Figure 4.6: ORTEP projection of complex **4.23** excluding hydrogen atoms for clarity, showing atom labelling scheme.

Table 4.4: Selected bond lengths (Å) and bond angles (°) of **4.23**.

| | | | | | |
|---------|----------|----------|------------|------------|------------|
| C1-N1 | 1.355(5) | C19-Rh1 | 2.114(4) | N1-C1-Rh1 | 126.4(3) |
| C1-Rh1 | 2.032(4) | C22-Rh1 | 2.197(4) | C1-Rh1-Cl1 | 88.03(11) |
| C1-N2 | 1.368(5) | Rh1-Cl1 | 2.3786(10) | C3-N1-C14 | 123.7(3) |
| N1-C3 | 1.387(5) | C23-Rh1 | 2.208(4) | C1-Rh1-C18 | 90.62(15) |
| C18-Rh1 | 2.097(4) | N1-C1-N2 | 103.8(3) | C1-Rh1-C22 | 163.07(15) |




Figure 4.7: ORTEP projection of complex **4.24** excluding hydrogen atoms for clarity, showing atom labelling scheme.

Table 4.5: Selected bond lengths (Å) and bond angles (°) of **4.24**.

| | | | | | |
|---------|-----------|-------------|----------|--------------|------------|
| C11-N12 | 1.358(3) | Rh1-C3 | 2.105(3) | C11-Rh1-Cl2 | 87.46 (7) |
| C11-Rh1 | 2.016(3) | Rh1-C4 | 2.121(3) | C11- Rh1- C8 | 167.02(10) |
| Rh1-Cl2 | 2.3794(7) | C11-N26 | 1.358(3) | C11- Rh1 -C3 | 92.99(10) |
| Rh1-C8 | 2.230(3) | | | C11-N12 -C13 | 125.20(2) |
| Rh1-C7 | 2.196(2) | N12-C11-N26 | 103.7(2) | C3-Rh1-Cl2 | 157.31(8) |

Due to variations in the steric bulk around the carbene carbon there was the need to compare the structure of all the rhodium(I) carbene complexes **4.20-4.24**. In doing so two planes were chosen to calculate the inter planer angles. The planes chosen are the NHC consisting of the five atoms in the heterocycle carbene ring and the rhodium coordination plane consisting of C1-Rh1-Cl2. The inter planer angles and other important angles are listed in Table 4.6 below.

Table 4.6: Bond angles ($^{\circ}$) for compounds **4.20- 4.24** of rhodium complexes of general formula $[\text{Rh}(\text{Cod})\text{Cl}(\text{NHC})]$

| Compound | Θ (interplaner angle) | N1-C1-N2 | C1-Rh1-Cl1 | C11-Rh1-(C18-C19) |
|-------------|------------------------------|-----------|------------|-------------------|
| 4.20 | 76.87 | 103.80(4) | 87.99(14) | 165.30(2) |
| 4.21 | 76.77 | 103.70(3) | 87.67(11) | 165.92(11) |
| 4.22 | 80.31 | 104.70(4) | 89.25(3) | 159.6(2) |
| 4.23 | 76.49 | 103.80(3) | 88.03(11) | 163.07(15) |
| 4.24 | 82.69 | 103.70(2) | 87.46(7) | 167.02(10) |

From the table presented above, the main structural features of these compounds are:

- A distorted square planar geometry with Cl-Rh-L angles within the range of 87.46 - 89.25° , the highest being observed in compound **4.22**.
- An average value of 76.71° inter planar angles for compounds **4.20**, **4.21** and **4.23**, 80.31° for **4.22** and 82.69° for **4.24**, thus orientation of the NHC with respect to the coordination approaches perpendicular probably to reduce steric interactions.
- The N1-C1-N2 angle of 103.70° observed around the carbene carbon in all complexes. Only **4.22** slightly differ with a value of 104.70° .
- C1-Rh1-(Cod) angles for the complexes ranges from $159.6(2)$ - $167.02(10)^{\circ}$, with the highest being observed in **4.24**, a significant variation from the normal square planar(180°).

Some important bond distances are listed in Table 4.7 in order to see the variations in the complexes.

Table 4.7: Other bond distances (Å) for complexes of general formula [Rh(Cod)Cl(NHC)]


| Compound | Rh-C | Rh-Cl | N1-C1 | Rh-(C15-C16) | Rh-(C19-C20) |
|----------|-----------|-----------|----------|--------------|--------------|
| 4.20 | 2.028(5) | 2.386(13) | 1.336(6) | 2.123(5) | 2.206(5) |
| 4.21 | 2.045(5) | 2.387(9) | 1.355(5) | 2.112(4) | 2.202(4) |
| 4.22 | 2.037(5) | 2.377(13) | 1.343(6) | 2.113(5) | 2.213(5) |
| 4.23 | 2.032(4) | 2.379(10) | 1.355(5) | 2.114(4) | 2.197(4) |
| 4.24 | 2.0163(3) | 2.379(7) | 1.358(3) | 2.121(3) | 2.105(3) |

-The distances between Rh and the NHC ligands are shorter than distances between Rh and the Cod, owing to the strong electron donating ability of NHCs.

- There is no significant difference in the C-N bond distance of NHCs, with an average value of 1.35 Å (Table 4.7), this indicates a degree of C-N double character. This parameter may vary depending on the degree of back donation from Rh to the NHC (more back donation decreases the push mesomeric effect in the NHC and increases the C-N distance), based on these observations, it can be deduced that all the complexes display the same level of back donation. Similarly, from Table 4.6 the N1-Ccarbene-N2 angles (103.95° on average) are very close.

The chelation of **quinN⁺C-R** toward the rhodium centre was achieved by the treatment of **4.20** and **4.24** with an equimolar amount AgBF₄, leading to chloride abstraction and ligand substitution of chloride by the quinoline nitrogen. All of the ¹H NMR signals of the quinoline hydrogen atoms in **4.25** are shifted downfield relative to those in the non chelated rhodium complex **4.20** indicating chelation of the **quinN⁺C-R** ligand. The ¹H NMR data for the non chelated complex **4.20** and that of the corresponding chelated complex **4.25** are δ= 8.10 (d, 1H, quin-H), 8.00 (d, 1H, quin-H), 7.75 (d, 1H, quin-H), 7.70 (t, 2H, quin-H), 7.45 (t, 1H, quin-H), 6.80 (s, 2H, CHHC), 6.50 (d, 1H, CH_{2linker}), 5.60 (d, 1H, CH_{2linker}), 5.00 (m, 2H, COD), 4.10 (s, 3H, CH₃), 3.40 (m, 1H, COD), 3.20 (m, 1H, COD), 2.30 (m, 4H, COD), 1.90 (m, 4H, COD) and δ 8.75 (d, 1H, quin-

H), 8.25 (d, 1H, quin-H), 8.10 (d, 1H, quin-H), 7.85 (t, 2H, quin-H), 7.60 (d, 2H, CHHC, quin-H), 6.65 (s, 1H, CHHC), 6.25 (d, 1H, CH_{2linker}), 6.05 (d, 1H, CH_{2linker}), 5.50 (m, 1H, COD), 4.70 (m, 1H, COD), 4.40 (m, 2H, COD), 4.10 (m, 1H, COD), 3.65 (s, 3H, CH₃), 2.80 (m, 1H, COD), 2.40 (m, 4H, COD), 2.10 (m, 2H, COD) respectively. High resolution MS of complex **4.25** displays a cluster of peak $M/z = 434.1117$ Da (100% intensity) assignable to the $[M - BF_4]^+$. Elemental analysis gave unacceptable results probably because of contamination from silver halides which were difficult to remove. ¹³C NMR spectrum for complex **4.25** could not be obtained because the complex is not soluble in most solvents. Suitable crystals for X-ray crystallography were not obtained for complex **4.25**. Complex **4.26** was not characterised because it is insoluble in most solvents such as DCM, acetonitrile and methylene chloride. The use of DMSO did help solve the problem of solubility but the ¹H NMR spectra of both chelated and non chelated rhodium complexes are essentially similar possibly due to competitive coordination of DMSO. Elemental analysis returned a slightly low percentage of carbon presumably due contamination from silver halide.




Scheme 4.2: Synthesis of chelated quinoline functionalised Rh(I) NHC complexes.

4.2.2: Synthesis and characterisation of quinoline functionalised Ir(I) NHC complexes.

The method for the synthesis of quinoline functionalised Ir(I) NHC complexes is similar to that of Rh(I) NHC. Complexes were prepared from the silver carbene complexes prepared in chapter 3. Stirring of silver carbene complexes with 0.5 equivalent of $[\text{Ir}(\text{COD})\text{Cl}]_2$ in DCM at ambient temperature gave the corresponding Ir(I)NHC complexes in good yield after work up as shown in Scheme 4.3 below.

Characterization of the iridium complexes, **4.27**, **4.28**, **4.29**, **4.30** and **4.31** was performed by NMR spectroscopy, elemental analysis and in most cases by X-ray crystallography. The ^{13}C NMR data for the coordinating carbene carbons appear at δ 180.02 ppm for **4.27**, 178.66 for **4.29**, 179.56 for **4.30** and 180.57 for **4.31**, indicating the formation of the Ir-C bond. All these signals are within the typical range for Ir-C(carbene) observed for analogous complexes [11,23, 24]. There is no evidence of coordination between the nitrogen of the quinoline with the iridium ion as the ^1H NMR shifts corresponding to the proton of the quinoline ring are essentially the same to those of the silver complexes. In all the iridium complexes prepared, the ^1H NMR spectra show diastereopic protons for the CH_2 linker ($\delta = 6.35$ and 5.50 for **4.27**, $\delta = 6.25$ and 5.95 for **4.28**, $\delta = 6.35$ and 5.60 for **4.29**, $\delta = 6.35$ and 5.55 for **4.30** and $\delta = 6.25$ and 5.65 for **4.31** indicating that this group is out of coordination plane of the molecule thus reducing its symmetry [11].



3.36: R = Me, X = Cl

4.27: R = Me

Table 4.8: Selected bond lengths (Å) and bond angles (°) of **4.28**.

| | | | | | |
|---------|-----------|-------------|--------------|-------------|-----------|
| C11-Ir1 | 2.051(11) | Ir1-C8 | 2.177(11) | C11-Ir1-C4 | 95.13(4) |
| C11-N12 | 1.367(15) | Ir1-C7 | 2.159(11) | C11-Ir1-C12 | 87.70(9) |
| Ir1-C12 | 2.372(2) | C25-N12 | 1.433(14) | C11-Ir1-C8 | 157.9(5) |
| Ir1-C3 | 2.125(13) | C11-N15 | C16 124.7(9) | C4-Ir1-C12 | 158.2(4) |
| Ir1-C4 | 2.096(11) | N12-C11-N15 | 105.6(9) | C11-N12-C25 | 125.30(9) |

The elemental analysis of complex **4.29** gave results consistent with the formulation of the compound and crystals suitable for X-ray crystallography were obtained by diffusion of n-pentane into a saturated DCM solution of the compound. The molecular structure of the complex was unequivocally confirmed by means of single X-ray crystallography. Figure 4.9 shows the molecular structure of **4.29** which is virtually identical to that of **4.28**. The Ir centre is in square planar geometry and the Ir-C of 2.045(6) Å is in the usual range for Ir^I-NHC complexes [25, 26]. Selected bond distances and bond angles are presented in table 4.9.


**Figure 4.9:** ORTEP projection of complex **4.29** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 4.9: Selected bond lengths (Å) and bond angles (°) of **4.29**.

| | | | | | |
|----------|------------|-------------|----------|--------------|------------|
| C1- Ir1 | 2.045(6) | Ir1-C21 | 2.196(7) | C1-Ir1- C17 | 93.10(3) |
| C1- N1 | 1.357(8) | Ir1-C22 | 2.195(6) | C1- Ir1-C11 | 90.04 (19) |
| Ir1- C11 | 2.3719(16) | C1-N2 | 1.367(8) | C1-Ir1-C18 | 88.5(3) |
| Ir1-C17 | 2.112(6) | C11-N15-C16 | 124.7(9) | C1- Ir1- C21 | 160.0(3) |
| Ir1-C18 | 2.104(7) | N1-C1-N2 | 103.7(5) | C1-N1-C4 | 124.60(5) |

Crystals suitable for X-ray crystallography were not obtained for complex **4.30** but elemental analysis returned satisfactory results in agreement with the proposed structure.

Elemental analysis of complex **4.31** gave results that are in agreement with the formulation of the complex and crystals suitable for X-ray crystallography were obtained by diffusion of n- pentane into saturated DCM solution of the complex. The crystal structure and selected bond distances and angles are shown in Figure 4.10 and Table 4.8 respectively. The iridium exhibits a square planar geometry and the Ir-C11(carbene) bond is typical for an Ir-C single bond [25, 26]. Other parameters in terms of bond lengths and bond angles are similar to that of **4.28** and **4.29**.


**Figure 4.10:** ORTEP projection of complex **4.31** excluding hydrogen atoms for clarity showing atom labelling scheme.

Table 4.10: Selected bond lengths (Å) and bond angles (°) of **4.31**.

| | | | | | |
|---------|------------|--------------|----------|-------------|------------|
| C11-Ir1 | 2.022(4) | Ir1-C8 | 2.196(5) | C11-Ir1-C4 | 91.42(11) |
| C11-N12 | 1.354(5) | Ir1-C7 | 2.160(4) | C11-Ir1-Cl2 | 87.54 (12) |
| Ir1-Cl2 | 2.3653(11) | C11-N26 | 1.361(5) | C11-Ir1-C8 | 168.44(18) |
| Ir1-C3 | 2.082(4) | C11-N12- C13 | 123.7(3) | C8-Ir1-Cl2 | 82.45(19) |
| Ir1-C4 | 2.121(4) | N12-C11-N26 | 104.0(3) | C11-Ir1-C7 | 154.57(19) |

To compare the structure of the three Ir(I) complexes the interplanar angles between the planes of the carbene ring and Ir coordination plane were calculated as Θ and tabulated in Table 4.11.

Table 4.11: Bond angles (°) for compounds **4.28**, **4.29** and **4.31** of iridium complexes of general formula [Ir(Cod)Cl(NHC)].

| Compound | Θ (interplaner angle) | N1-C1-N2 | C1-Ir1-Cl1 | C11-Ir1-(C18-C19) |
|-------------|------------------------------|-----------|------------|-------------------|
| 4.28 | 76.43 | 105.6(9) | 87.70(14) | 158.20(4) |
| 4.29 | 77.79 | 103.70(5) | 90.04(19) | 160.00(3) |
| 4.31 | 86.76 | 104.03(3) | 87.54(3) | 168.44(2) |

In general, neutral complexes display slight variation in their geometries and the main structural features of these compounds are:

- Distorted square planar geometry with Cl-Ir-L of around 87.62° on the average for compounds **4.28** and **4.31**, and 90.04° for **4.29** (Table 4.11).
- A value of 77.11° (on average) with respect to the plane of carbene ring and the Ir coordination plane (Cl-Ir-C11) for complexes **4.28** and **4.29** and 86.76° for complex **4.29** which is almost perpendicular, probably to reduce steric interactions.
- C1-Rh1-(Cod) angles for the complexes ranges from $158.20(4)$ - $168.44(2)^\circ$, with the highest being observed in **4.31**, a significant variation from the normal square planar (180°).


Some important bond distances are listed in Table 4.12 in order to see the variations in the complexes.

Table 4.12: Other bond distances (Å) for complexes of general formula [Ir(Cod)Cl(NHC)]

| Compound | Ir-C | Ir-Cl | N1-C1 | Ir-(C15-C16) | Ir-(C19-C20) |
|-------------|-----------|-----------|-----------|--------------|--------------|
| 4.28 | 2.051(11) | 2.372(2) | 1.367(15) | 2.096(11) | 2.177(11) |
| 4.29 | 2.045(6) | 2.372(9) | 1.357(8) | 2.104(7) | 2.196(7) |
| 4.31 | 2.022(5) | 2.365(11) | 1.354(5) | 2.082(4) | 2.196(5) |

-The distances between Rh and the NHC ligands are shorter than distances between Rh and the Cod, owing to the strong electron donating ability of NHCs.

- There is no significant variation in the C-N bond distance of NHCs, with an average value of 1.36 Å indicating double bond character (Table 4.12). This parameter may vary depending on the degree of back donation from Ir to the NHC (more back donation decreases the push mesomeric effect in the NHC and increases the C-N distance), based on these observations, it can be deduced that all the complexes display the same level of back donation. Similarly, from Table 4.11 the N1-Ccarbene-N2 angles (104.44° on average) are very close.



4.27: R = Me

4.31: R = CH₂Quin

4.32: R = Me

4.33: R = CH₂Quin

Scheme 4.3: Synthesis of chelated quinoline functionalised Ir(I) NHC complexes

The chelation of **quinN⁺C-R** with the iridium centre was achieved by the treatment of **4.27** and **4.31** with an equimolar amount AgBF_4 , leading to the ligand substitution of chloride by the quinoline nitrogen as shown in Scheme 4.3 above. All of the ^1H NMR signals of the quinoline hydrogen atoms in **4.32** are shifted downfield relative to those in the silver complex **4.20** indicating chelation of the **quinN⁺C-R** ligand. The methylene linker protons shifted from 6.35 and 5.50 ppm for complex **4.20** to 6.80 and 5.90 ppm for complex **4.32** and the protons from the imidazole moiety shifted from 6.80, 6.70 ppm to 7.55 and 6.80 ppm respectively.

Complex **4.33** was not characterized by NMR spectroscopy due to lack of suitable solvent for analysis because it is insoluble in most common solvents. In high polar solvent such as DMSO, the spectrum of chelated and the non chelated complexes are virtually similar possibly due to competitive coordination of DMSO. Elemental analysis of complexes **4.32** and **4.33** returned unacceptable results with low percentage of carbon, nitrogen and hydrogen probably as a result of contamination from silver halides.

4.2.3 Attempted synthesis of Rh(I) and Ir(I) complexes of ligands without methylene separating the quinoline

Attempts were made to prepare the Rh(I) and Ir(I) (NHC) complexes of ligands **2** (1,3-diquinolin-4,5-dihydroimidazolium tetrafluoroborate) and **4** (1,3-diquinoline-3,4,5,6-tetrahydropyrimidinium hexafluoro phosphate) by the treatment the ligands with the following reagents:


- i) Ag_2O in dichloromethane, acetonitrile and DMSO
- ii) $\text{K}[\text{N}(\text{SiMe}_3)_2]$ and $[\text{Rh}(\text{COD})\text{Cl}]_2$ in THF
- iii) $\text{K}[\text{N}(\text{SiMe}_3)_2]$ and $[\text{Ir}(\text{COD})\text{Cl}]_2$ in THF

The reaction of the ligands with Ag_2O did not yield the expected carbene complexes as NMR spectra revealed only the starting salts. Performing the reaction at elevated temperature and for longer times did not give the desired carbene complexes. However treatment of the ligands with $\text{K}[\text{N}(\text{SiMe}_3)_2]$ in THF showed that the $\text{C}_2\text{-H}$ has been removed but could not coordinate as addition of $[\text{Rh}(\text{COD})\text{Cl}]_2$ or $[\text{Ir}(\text{COD})\text{Cl}]_2$ did not give the desired carbene metal complexes.

Solubility was thought to be responsible for the negative results obtained in the preparation of the silver(I) carbene complexes. With this in mind, it was decided to change the tetrafluoroborate and hexafluorophosphate counter ions to sodium tetrakis [(3, 5-trifluoromethyl) phenyl] borate (NaBArF_{24}). The replacement of the counter ion solved the problem of solubility but upon reaction with silver oxide followed by $[\text{Rh}(\text{COD})\text{Cl}]_2$ or $[\text{Ir}(\text{COD})\text{Cl}]_2$ the desired carbene complexes were not formed. After several problems preparing the complexes with ligands **2** and **4** we decided to diversify and were successful in the synthesis of a variety of compounds by slightly varying the structure of the ligands.

A variation on this type of structure has been to add a methylene linker between the carbene and the quinoline moiety. This will provide extra flexibility and reduce steric strain.


Attempt was also made to synthesise the Rh(I) and Ir(I) (NHC) complexes by the reaction of silver(I) (NHC) complex **3.34** prepared in chapter 3 with $[\text{Rh}(\text{COD})\text{Cl}]_2$ and $[\text{Ir}(\text{COD})\text{Cl}]_2$ in DCM as shown in Scheme 4.4 below.



Scheme 4.4: Attempted of rigid quinoline functionalised Rh(I) and Ir(I) (NHC) carbene complexes.

The reaction did not give the desired product, the NMR looked messy and therefore could not be interpreted.

C-H oxidative addition was thought to be another feasible way to obtain the rhodium and iridium complexes, though, this would lead to the synthesis of Rh(III) and Ir(III) (NHC) carbene complexes. Therefore, following the methods reported in the literature [8], 1-benzyl-3-quinolinimidazoliumtetrafluoroborate (**6b**) was reacted with $[\text{Rh}(\text{COD})\text{Cl}]_2$ and $[\text{Ir}(\text{COD})\text{Cl}]_2$ in refluxing acetonitrile for 24 hours. However the reaction gave the original mixture, even when the reaction was carried out for a longer time (one week). Modification of the reaction conditions (refluxing toluene and DMSO) did not produce any of the expected products.



Scheme 4.5: Attempted of Rh and Ir carbene complex via oxidative addition

4.3 Ir(I) NHC catalysed hydrogen transfer reactions

4.3.1 Catalysis

A catalyst may be defined as a substance that increases the rate at which a chemical reaction approaches equilibrium; thus a catalyst affects the kinetics of a reaction rather than the overall thermodynamics [27]. Different catalysts will also accelerate one reaction with respect to another, and are thus especially valued for their ability to influence product selectivity [28]. In a catalytic cycle, the catalyst remained unchanged, aside from degradation or poisoning by side reactions.

The importance of the catalyst can not be over emphasised due to the fact that many chemical reactions will not proceed to appreciable extent unless in the presence of a suitable catalyst. Apart from the importance of catalysts to the world economy, it is also crucial to the existence of all living organisms through the actions of naturally occurring enzymes. It is worth noting that 75% of all chemicals are produced with the application of some sort of catalyst; and it is up to 90% when only those more modern processes are considered [28].

Catalytic reactions are mainly divided into heterogeneous and homogeneous systems dependent on the phase relationship of the catalyst to the substrate. Enzymatic catalysis as seen in biological reactions constitutes an additional and separate group [27] and these systems are sometimes classified as immobilised catalyst [28].

The current work involves homogeneous systems utilising iridium complexes as precatalysts, i.e. precursors to the actual catalytic species. The essential

characteristics of homogeneous systems have been broadly defined by Cornils and Herrmann [29]:

- i) the catalyst is moderately dispersed in the same phase as the reactants;
- ii) the catalyst is able to be unequivocally characterised chemically and spectroscopically, and thus able to be synthesised in a reproducible manner;
- iii) new catalysts are able to be rationally designed for specific purposes according to known chemical principles;
- iv) unequivocally reaction kinetics may be related to each metal atom of the catalyst.

Although the catalytic reactions take place at the metal atom, the supporting ligands bound to the metal are important for promoting and modifying catalytic activity through the prevention of metal aggregation, stabilisation of intermediates, provision of vacant coordination sites at the metal via dissociation equilibria, and modification of the steric and electronic environment about the metal ion [29].

While the market share of homogeneous catalysis is only 10- 15%, recent developments in the application of transition metal complexes as catalysts has led to the prospect of a wide variety of new, high value organic molecules being accessed using relatively simple and affordable substrates and procedure [30]. Progress in these areas of homogeneous catalysis will ultimately lead to the synthesis of highly active, selective, and robust catalyst that use cheap substrates to produce important compounds in an efficient manner.


Due to time constraints, catalyst testing of the complexes synthesised in this chapter was confined to the hydrogen transfer reduction.

4.5.2 Hydrogen-transfer reduction

The conventional methods for the hydrogenation of unsaturated bonds have involve the use of molecular hydrogen which has its draw backs due to the difficult handling procedures involved with working at high pressures. Efforts have since been devoted to developing safer, more cost effective methods and transfer hydrogenation represents an ideal alternative.

Transfer hydrogenation hydrogenates a double bond (e.g. C=C in olefins [31], N=O [11] in nitro groups and C=O in carbonyls [33]) by abstracting hydrogen from a proton donor source such as isopropanol, usually present in excess as the reaction solvent. The reaction is carried out in the presence of the catalyst in addition to the base used to assist deprotonation.

The hydrogen transfer reduction of a carbonyl function catalysed by transition metal complexes is well documented [34] and an accepted mechanism for the catalytic transfer hydrogenation on the reduction of benzophenone is given in Scheme 5 below.



Scheme 4.6: mechanism of transfer hydrogenation of benzophenone

The attractiveness of this method lies in its inexpensive starting materials and simple experimental procedure, which has been exploited in the extensive level of development in this area. It is generally agreed that this research area was spearheaded by Noyori et al in 1995 [35]. They discovered that chiral Ru(II) complexes were capable of asymmetric transfer hydrogenation at reflux temperature and had sufficient catalytic activity on aryl ketones at ambient temperatures. The Noyori catalyst is a Ru(II)Cl₂ centre complexed with chiral (1S, 2S)-N-(p-toluenesulfonyl)-1, 2-diphenylethylenediamine [(S,S)-TsDPEN]. Buriak et al explored the efficiency of N-heterocyclic carbene ligands in combination with phosphines


in hydrogenation of olefins [36]. Pyridinyl N-Heterocyclic carbene complexes have also been used in the reduction of nitroarenes and carbonyls [11].

Rh and Ir are known to be effective catalysts for the transfer hydrogenation of unsaturated substrate by hydrogen donors (e.g. cyclohexene or 2-propanol) [32]. Significantly, transfer hydrogenation of carbonyl compounds in iPrOH is the most widely used reaction to test the catalytic properties of Rh and Ir because of its simplicity. It has been established that iridium carbene complexes are more active than their rhodium analogues in transfer hydrogenation [33].

The catalysis presented in this work involves the use quinoline functionalised iridium carbene complexes in transfer hydrogenation.


4.5.3 Results and discussions

The catalytic transfer hydrogenation was examined using 2-propanol as the hydrogen source with a KO^tBu base promoter.



Scheme 4.7: Catalytic hydrogen transfer of 4-bromoacetophenone

Two quinoline based iridium carbene complexes were used in catalytic hydrogen transfer of 4-bromoacetophenone:



The activity of iridium complexes **4.27** and **4.31** was tested at different concentrations which enable a direct comparison between the effects of the alkyl and the quinoline on the N-substituents. The results of the catalytic tests are presented in table 4.13 below.

Table 4.13: Catalysis with **4.27** and **4.31** in hydrogen transfer reaction

| Entry | Catalyst | Concentration (mole %) | Conversion (%) | TON |
|-------|-------------|------------------------|----------------|------|
| 1 | 4.27 | 1 | 99 | 99 |
| 2 | 4.27 | 0.1 | 99 | 990 |
| 3 | 4.27 | 0.01 | 65 | 6500 |
| 4 | 4.31 | 1 | 99 | 99 |
| 5 | 4.31 | 0.1 | 99 | 990 |
| 6 | 4.31 | 0.01 | 78 | 7800 |

Conditions: substrate: 1.00 mmol; KO^tBu: 1.00 mmol; 80°C; 24hr. Determined by NMR.

As presented in the table above, both of the catalysts showed good catalytic activity towards transfer hydrogenation with no significant change when the catalyst loading was reduced from 1 mole % to 0.1 mole %. When 0.01 mole % of the catalysts was used, the catalysts still showed good activity with virtually no difference

In order to evaluate the differences between catalysts **4.27** and **4.31** carbene complexes, a low catalyst loading (0.01 mol %) at 80 °C. The yield for the transfer hydrogenation of 4-bromoacetophenone was studied over a period of 3 hours by taking and running the samples after a given time and the results are presented in Figure 4.11 below.

As shown in the diagram below, there is no significant difference between catalysts **4.31** and **4.27** as both catalysts appeared to be highly efficient in the transfer hydrogenation of 4-bromoacetophenone to 1-phenylethanol both giving over 90% conversion after 180 minutes. The performances of the iridium carbene complexes are comparable to the ones reported by Hahn et al [39], though it required longer time to achieve the desired results. It also compares

well with the pyridinyl Ir(I) NHC carbene complexes reported by Peris et al [38].




Figure 4.11: Time dependence of the catalytic transfer hydrogenation of 4-bromoacetophenone; 0.01 mol % of cat. (4.27 and 4.37), KO^tBu (10 mol%), 10 mol% of 4-bromoacetophenone, solvent 2-propanol (5mL), T = 80 °C

The steric properties of the carbene may have played important roles in the selectivity and reactivity of the systems. 4.31 is significantly more sterically hindered than the 4.27 analogue because of the presence of the bulky quinoline moiety. However this steric differences does not seem to influence the activity of the catalyst. Another consideration is partial chelation of the ligand during catalysis.

4.4 Conclusions

In this chapter both of the rhodium(I) carbene complexes and iridium(I) carbene complexes were synthesized from the silver(I) carbene described in chapter 3


by transmetallation reactions. This procedure was earlier utilised to synthesise the iridium pyridinyl NHC complexes [11] producing both chelated and non chelated complexes. The replacement of the pyridine moiety with quinoline increases the steric bulk which may have significant implications in catalysis. While the non chelated carbene complexes were obtained in good yield and were fully characterized, the chelated analogues were found to be insoluble in most solvents rendering full characterization difficult. To our knowledge the rhodium (I) and iridium(I) NHC complexes presented in this work are the first of their kind that are functionalised with quinoline except the recent publication by Webster et al. The reaction of quinoline ligands without methylene linker under different reaction condition did not give the desired carbene complexes. The Ir(I) NHC Complexes tested in transfer hydrogenation reactions showed good catalytic activity even at low catalyst loading

4.5 Experimental

4.5.1 General comments


All reactions were performed under the atmosphere of dry dinitrogen or argon using standard Schlenk techniques, and solvents were purified and dried by usual means [40], unless otherwise indicated. Silver(I) NHC complexes were prepared as detailed in Chapter 3, and all other reagents were used as received. All NMR data are quoted δ /ppm. ^1H and ^{13}C (proton decoupled) spectra NMR were recorded on a Bruker DPX Advance 400 (^1H at 400MHz, ^{13}C at 100.61 MHz) at ambient temperature, unless otherwise stated, and referenced to SiMe_4 . Electrospray mass spectrometry (ESMS) was performed on a VG Fisons Platform II instrument by the department of Chemistry, Cardiff University. Micro analysis was performed by Warwick Analytical Service. All reactions involving silver compounds were performed with the exclusion of light.

[1-methyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride 4.20



[RhCl(cod)]₂ (99mg, 0.2mmol) in 10ml of DCM was added to 20 ml of DCM solution of [Ag (1-methy-3-(2-methylquinoline) imidazolin-2-ylidene)₂][AgCl₂] **3.36** (147mg, 0.40mmol). The reaction mixture was stirred over night and filtered through celite to remove silver chloride and any insoluble residue. The filtrate was concentrated and hexane was added to precipitate out the desired carbene complex as a yellow powder (152mg, 80.42%). Crystals suitable for X-ray crystallography were grown by layering diethyl ether on dichloromethane. Anal. Calcd. for C₂₂H₂₅N₃RhCl: C, 56.20; H, 5.33; N, 8.95%. Found: C, 54.80; H, 5.23; N, 8.66%. ¹H NMR (CDCl₃, 400MHz, 298K): δ 8.10(d, 1H, J = 8.5 Hz, quin-H), 8.00(d, 1H, J = 8.5 Hz, quin-H), 7.75(d, 1H, J = 8.0 Hz, quin-H), 7.70(t, 2H, J = 6.7 Hz quin-H), 7.45(t, 1H, 7.5 Hz, quin-H), 6.80(s, 2H, CHHC), 6.50(d, 1H, J = 14.8 Hz CH₂linker), 5.60(d, 1H, J = 14.8 Hz CH₂linker), 5.00(broad, 2H, COD), 4.10(s, 3H, CH₃), 3.40(broad, 1H, COD), 3.20(broad, 1H, COD), 2.30(m, 4H, j= 4.4 Hz, COD), 1.90(m, 4H, J = 4.2 Hz COD). ¹³C NMR (CDCl₃, 100MHz, 298K): 181.60 (C-Rh), 154.88, 145.53, 135.57, 127.82, 127.05, 125.86, 125.73, 124.78, 120.78, 118.99, 118.87, 96.93(N-CH₃), 66.32, 54.96, 35.83, 31.33, 30.65, 27.20, 26.66.


[1-mesityl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride 4.21



Following the procedure for the synthesis of complex **4.20**, **4.21** was obtained from **3.37** (110 mg, 0.23mmol) and [RhCl(cod)]₂ (58 mg, 0.12mmol). Yield = 95 mg (71.00%). Crystals suitable for X-ray crystallography were grown by

layering diethyl ether on dichloromethane. Anal. Calcd. for $C_{30}H_{33}N_3RhCl$: C, 62.78; H, 5.76; N, 7.32%. Found: C, 61.44; H, 5.48; N, 7.06%. 1H NMR ($CDCl_3$, 400MHZ, 298K): δ 8.15(m, 1H, J = 8.4 Hz, quin-H), 8.00(d, 1H, J = 8.1 Hz, quin-H), 7.80(d, 1H, J = 5.7 Hz, quin-H), 7.70(s, 1H, Ar-H), 7.50(t, 2H, J = 7.0 Hz, quin-H), 7.05(t, 2H, quin-H, Ar-H), 6.90(s, 1H, CHHC), 6.70(s, 1H, CHHC), 6.50(d, 1H, J = 15.4 Hz, $CH_{2linker}$), 6.10(d, 1H, J = 15.4 Hz, $CH_{2linker}$), 4.80(broad, 2H, COD), 3.30(broad, 1H, COD), 3.00(broad, 1H, COD), 2.40(broad, 3H, COD), 2.30(s, 3H, p-CH₃), 1.80(s, 6H, o-CH₃), 1.50(broad, 4H, COD). ^{13}C NMR ($CDCl_3$, 100MHZ, 298K): 155.03, 145.70, 136.61, 135.05, 134.97, 134.01, 132.36, 127.68, 127.52, 126.16, 125.69, 125.48, 124.62, 121.19, 119.73, 118.75, 95.44(NCH₂), 55.14, 31.56, 29.58, 26.93, 25.95(p-CH₃), 19.06(o-CH₃), 17.06, 15.75. HR-MS for $[M - Cl]^+$ = 538.18(100%): calculated, 538.1730($C_{30}H_{33}N_3Rh$), found, 538.1751.


[1-isopropyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride 4.22



Following the procedure for the synthesis of complex **4.20**, **4.22** was obtained from **3.38** (134mg, 0.276mmol) and $[RhCl(cod)]_2$ (68 mg, 0.14mmol). Yield = 110 mg (80.29%). Crystals suitable for X-ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $C_{24}H_{29}N_3RhCl$: C, 57.90; H, 5.83; N, 8.4%. Found: C, 53.85; H, 5.59; N, 7.47%. 1H NMR ($CDCl_3$, 400MHZ, 298K): δ 8.10(m, 1H, J = 8.5 Hz, quin-H), 8.00(d, 1H, J = 8.4 Hz, quin-H), 7.70(d, 1H, J = 8.1 Hz, quin-H), 7.65(t, 2H, J = 4.1 Hz, quin-H), 7.45(t, 1H, J = 7.0 Hz, quin-H), 6.80(s, 2H, CHHC), 6.50(d, 1H, J = 14.8 Hz, $CH_{2linker}$), 5.75(m, 1H, J = 6.8 Hz, iPr-H), 5.60(d, 1H, J = 14.8 Hz, $CH_{2linker}$), 5.00(m, 1H, J = 7.6 Hz, COD), 4.90(m, 1H, J = 5.2 Hz, COD), 3.35(m, 1H, J = 2.6 Hz, COD), 2.1-2.40(m, 4H, J = 4.5 Hz, COD), 1.7-2.00(m, 4H, J = 6.6 Hz, COD), 1.5(d, 6H, J = 4.2 Hz, CH₃). ^{13}C NMR ($CDCl_3$, 100MHZ, 298K): 180.38(C-Rh), 155.00,


145.53, 135.60, 127.80, 127.04, 126.74, 125.87, 125.73, 124.76, 119.22, 115.25, 97.17, 66.24, 55.19, 50.17, 31.50, 30.46, 27.38, 26.45, 22.19, 21.36, 20.72. HR-MS for $[M - Cl]^+ = 462.1415(100\%)$: calculated, 462.1417($C_{24}H_{29}N_3Rh$), found, 462.1415.

[1-n-butyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride 4.23




Following the procedure for the synthesis of complex **4.20**, **4.23** was obtained from **3.39** (150mg, 0.300mmol) and $[RhCl(cod)]_2$ (74 mg, 0.15mmol). Yield = 121 mg (79.08%). Crystals suitable for X-ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $C_{25}H_{31}N_3RhCl$: C, 58.66; H, 6.06; N, 8.21%. Found: C, 58.44; H, 6.48; N, 7.37%. 1H NMR ($CDCl_3$, 400MHz, 298K): δ 8.10(m, 1H, $J = 9.5$ Hz, quin-H), 8.00(d, 1H, $J = 8.4$ Hz, quin-H), 7.75(d, 1H, $J = 7.9$ Hz, quin-H), 7.65(t, 2H, $J = 4.8$ Hz, quin-H), 7.45(t, 1H, quin-H), 6.75(s, 2H, CHHC), 6.50(d, 1H, $J = 14.8$ Hz, $CH_{2linker}$), 5.60(d, 1H, $J = 14.8$ Hz, $CH_{2linker}$), 5.00(broad, 2H, COD), 4.50(m, 2H, $J = 6.1$ Hz, CH_2), 3.30(m, 2H, $J = 2.3$ Hz, COD), 2.40(m, 2H, $J = 3.3$ Hz, CH_2), 1.8(m, 4H, $J = 2.4$ Hz, COD), 1.40(m, 2H, $J = 7.4$ Hz, CH_2), 1.0(t, 3H, $J = 7.4$ Hz, CH_3). ^{13}C NMR ($CDCl_3$, 100MHz, 298K): 178.66(C-Rh), 155.72, 146.45, 136.45, 128.73, 127.97, 126.76, 126.62, 125.67, 119.86, 119.73, 115.80, 84.35, 83.04, 55.73, 52.42, 50.97, 33.20, 31.87, 29.14, 28.03, 22.97, 22.08. HR-MS for $[M-Cl]^+ = 476.1578(100\%)$: calculated, 476.1573($C_{24}H_{29}N_3Rh$), found, 476.1578.

[bis-1,3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride 4.24



Following the procedure for the synthesis of complex **4.20**, **4.24** was obtained from **3.41** (300mg, 0.600mmol) and $[\text{RhCl}(\text{cod})]_2$ (150 mg, 0.30mmol). Yield = 250 mg (68.87%). Crystals suitable for X-ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $\text{C}_{31}\text{H}_{30}\text{N}_4\text{RhCl}$: C, 62.37; H, 5.03; N, 9.39%. Found: C, 61.59; H, 5.13; N, 8.967%. ^1H NMR (CDCl_3 , 400MHz, 298K): δ 8.10(m, 2H, $J = 8.5$ Hz, quin-H), 8.00(d, 2H, $J = 8.4$ Hz, quin-H), 7.75(d, 2H, $J = 8.1$ Hz, quin-H), 7.65(m, 4H, $J = 4.8$ Hz, quin-H), 7.50(t, 2H, $J = 7.0$ Hz, quin-H), 6.85(s, 2H, CHHC), 6.50(d, 1H, $J = 14.9$ Hz CH_2 linker), 5.80(d, 1H, $J = 14.8$ Hz CH_2 linker), 5.00(broad, 2H, COD), 4.50(broad, 2H, CH_2), 3.30(m, 2H, $J = 2.7$ Hz, COD), 2.30(m, 4H, $J = 4.0$ Hz, COD), 1.90(m, 4H, $J = 8.5$ Hz, COD). ^{13}C NMR (CDCl_3 , 100MHz, R.T): 183.45(C-Rh), 155.64, 146.52, 136.52, 128.78, 128.03, 126.75, 126.61, 125.74, 120.58, 119.78, 98.38(CH_2) 67.93, 55.92, 32.84, 30.56, 27.81, 26.98, 21.63. HR-MS for $[\text{M}-\text{Cl}]^+ = 561.1548(100\%)$: calculated, 561.1526($\text{C}_{31}\text{H}_{30}\text{N}_4\text{Rh}$), found, 561.1548.


[1-methyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) tetrafluoroborate **4.25**



AgBF_4 (26 mg, 0.132mmol) was added to a stirred solution of complex **4.20** (62 mg, 0.132mmol) in 10 ml of DCM. The reaction mixture was stirred for 1 hour and reaction filtered through celite to remove silver chloride and any insoluble residue. The filtrates was then added drop by drop to a stirring solution of hexane to precipitate the compound which was then dried in vacuum to afford the desired compound as a deep yellow powder(42 mg, 60.87%). Anal. Calcd. for $\text{C}_{22}\text{H}_{25}\text{N}_3\text{RhBF}_4$: C, 50.70; H, 4.80; N, 8.07%. Found: C, 42.18; H, 4.26; N, 6.20%. ^1H NMR (CDCl_3 , 400MHz, 298K): δ 8.75(d, 1H, $J = 8.5$ Hz, quin-H), 8.25(d, 1H, $J = 8.3$ Hz, quin-H), 8.10(d, 1H, $J = 8.3$ Hz, quin-H), 7.85(t, 2H, $J = 8.2$ Hz, quin-H), 7.60(d, 2H, CHHC, $J = 5.3$ Hz, quin-H), 6.65(s, 1H, CHHC), 6.25(d, 1H, $J = 15.4$ Hz, CH_2 linker), 6.05(d, 1H, $J = 15.4$ Hz, CH_2 linker), 5.50(broad, 1H, COD), 4.70 (broad, 1H, COD), 4.40(broad, 2H, COD), 4.10(broad, 1H, COD), 3.65(s, 3H, CH_3), 2.80(broad, 1H, COD),


2.40(broad, 4H, COD), 2.10(broad, 2H, COD). HRMS for $[M-BF_4]^+$: calculated for $C_{22}H_{25}N_3Rh$ (M^+)434.1104, found, 434.1117.

[1-methyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) Chloride 4.27




$[IrCl(cod)]_2$ (180mg, 0.27mmol) in 10ml of DCM was added to 20 mL of DCM solution of $[Ag(1-methyl-3-(2-methylquinoline)imidazolin-2-ylidene)_2][AgCl_2]$ **3.36** (196mg, 0.54mmol). The reaction mixture was stirred over night and filtered through celite to remove silver chloride and any insoluble residue. The filtrate was concentrated and hexane was added to precipitate the desired carbene complex as a yellow powder (210mg, 70.23%). Crystals suitable for X-ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $C_{22}H_{25}N_3IrCl$: C, 47.25; H, 4.48; N, 7.52%. Found: C, 46.83; H, 4.51; N, 6.14%. 1H NMR ($CDCl_3$, 400MHz, 298K): δ 8.10(d, 1H, $J = 8.5$ Hz, quin-H), 8.00(d, 1H, $J = 8.4$ Hz, quin-H), 7.75(d, 1H, $J = 8.1$ Hz, quin-H), 7.65(t, 1H, $J = 5.6$ Hz, quin-H), 7.60(t, 1H, $J = 8.5$ Hz, quin-H), 7.45(d, 1H, $J = 5.6$ Hz, quin-H), 6.80(s, 1H, CHHC), 6.70(s, 1H, CHHC), 6.35(d, 1H, $J = 14.7$ Hz, CH_2 linker), 5.50(d, 1H, $J = 14.7$ Hz, CH_2 linker), 4.60(m, 2H, $J = 4.6$ Hz, COD), 3.90(s, 3H, CH_3), 3.00(m, 1H, $J = 4.4$ Hz, COD), 2.80(m, 1H, $J = 4.10$ Hz, COD), 2.20(m, 4H, $J = 3.2$ Hz, COD), 1.70(m, 4H, $J = 3.3$ Hz, COD). ^{13}C NMR ($CDCl_3$, 100MHz, R.T): 180.02 (C-Rh), 155.56, 146.43, 136.42, 128.75, 127.96, 126.75, 126.60, 125.70, 121.38, 119.68, 119.43, 83.40(N- CH_3), 55.48, 51.00(, 36.44), 32.91, 32.12, 30.56, 28.18, 21.63.

1-mesityl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) Chloride 4.28



Following the procedure for the synthesis of complex was obtained from **3.37** (200 mg, 0.425mmol) and $[\text{IrCl}(\text{cod})]_2$ (140 mg, 0.21mmol) . Yield: 151 mg (53.55%). Crystals suitable for X- ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $\text{C}_{30}\text{H}_{33}\text{N}_3\text{IrCl}$: C, 54.32; H, 4.98; N, 6.34; Cl, 5.37%. Found: C, 53.59; H, 5.08; N, 6.07; Cl, 5.25%. ^1H NMR (CDCl_3 , 400MHZ, 298K): δ 8.15(m, $J = 8.4$ Hz, 1H, quin-H), 7.90(d, 1H, $J = 8.0$ Hz, quin-H), 7.80(d, 1H, $J = 8.1$ Hz, quin-H), 7.70(d, 2H, $J = 7.0$ Hz, Ar-H), 7.50(t, 1H, $J = 7.7$ Hz, quin-H), 7.40(t, 1H, $J = 10.3$ Hz, quin-H), 7.30(t, 1H, $J = 7.8$ Hz, quin-H,), 7.1(s, 1H, CHHC), 6,70(s, 1H, CHHC), 6.25(d, 1H, $J = 15.2$ Hz, $\text{CH}_{2\text{linker}}$), 5.95(d, 1H, $J = 15.3$ Hz, $\text{CH}_{2\text{linker}}$), 4.40(broad, 2H, COD), 2.90(m, 1H,COD),2.70(m, 1H, $J = 4.8$ Hz, COD), 2.30(broad, 4H, COD), 2.2(s, 3H, p-CH₃), 1.80(s, 6H, o-CH₃), 1.50(m, 4H, $J = 5.9$ Hz, COD). ^{13}C NMR (CDCl_3 , 100MHZ, 298K):


[1-isopropyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) Chloride 4.29



Following the procedure for the synthesis of complex was obtained from **3.38** (120mg, 0.250mmol) and $[\text{IrCl}(\text{cod})]_2$ (83 mg, 0.13mmol) . Yield 121 mg (73.78%). Crystals suitable for X- ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $\text{C}_{24}\text{H}_{29}\text{N}_3\text{IrCl}$: C, 49.09; H, 4.94; N, 7.16%. Found: C, 47.98; H, 5.44; N, 6.98%. ^1H NMR (CDCl_3 , 400MHZ, 298K): δ 8.10(m, 1H, $J = 8.5$ Hz, quin-H), 8.00(d, 1H, $J = 8.4$ Hz, quin-H), 7.70(d, 1H, $J = 8.2$ Hz, quin-H), 7.65(t, 1H, $J = 5.8$ Hz, quin-H), 7.60(d, 1H, $J =$


8.5 Hz, quin-H), 7.45(d, 1H, J = 7.0 Hz, quin-H), 6.80(s, 2H, CHHC), 6.35(d, 1H, J = 14.8 Hz, CH₂linker), 5.60(m, 1H, J = 6.8 Hz, iPr-H), 5.50(d, 1H, J = 14.8 Hz, CH₂linker), 4.60(m, 1H, J = 3.5 Hz, COD), 4.50(m, 1H, J = 4.2 Hz, COD), 3.0(m, 1H, J = 5.2 Hz, COD), 2.80(m, 1H, J = 4.0 Hz, COD), 2.20(m, 4H, J = 6.9 Hz, COD), 1.6(m, 4H, J = 6.6 Hz, COD), 1,3(d, 6H, J = 8.0 Hz, CH₃). ¹³C NMR (CDCl₃, 100MHz, R.T): 178.66(C-Ir), 155.72, 146.45, 136.45, 128.73, 127.97, 126.76, 126.62, 125.67, 124.76, 119.86, 115.80, 84.35, 83.04, 55.73, 52.42, 51.23, 33.20, 31.87, 29.14, 28.03, 22.97, 22.08. LRMS-ES for [M – Cl]⁺ = 552.23(100%).

[1-n-butyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene) iridium(I) Chloride 4.30




Following the procedure for the synthesis of complex was obtained from **3.39** (150mg, 0.30mmol) and [IrCl(cod)]₂ (74 mg, 0.15mmol) . Yield:125 mg (69.44%). Anal. Calcd. for C₂₅H₃₁N₃IrCl: C, 49.94; H, 5.16; N, 6.99%. Found: C, 49.42; H, 5.43; N, 4.86%. ¹H NMR (CDCl₃, 400MHz, 298K): δ 8.1(m, 1H, J = 8.5 Hz quin-H), 8.0(d, 1H, J = 8.5 Hz, quin-H), 7.75(d, 1H, J = 8.0 Hz, quin-H), 7.65(t, 1H, J = 7.1 Hz quin-H), 7.55(d, 1H, J = 8.5 Hz, quin-H), 7.45(d, 1H, J = 7.1 Hz quin-H), 6.80(s, 2H, CHHC), 6.35(d, 1H, J = 14.8 Hz, CH₂linker), 5.55(d, 1H, J = 14.8 Hz, CH₂linker), 4.60(broad, 2H, COD), 4.40(m, 2H, J = 6.0 Hz, CH₂), 2.90(m, 2H, J = 7.3 Hz COD), 2.20(m, 2H, J = 5.5 Hz CH₂), 1.60(m, 4H, J = 8.3 Hz, COD) 1.4(m 2H, J = 7.3 Hz, CH₂), 1.00(t, 3H, J = 7.3 Hz, CH₃). ¹³C NMR (CDCl₃, 100MHz, R.T): 179.56(C-Ir), 155.67, 146.44, 136.43, 128.73, 127.97, 126.75, 126.61, 125.68, 119.73, 119.57, 119.47, 83.93, 83.64, 55.65, 51.07, 50.79, 49.27, 32.58, 31.91, 30.56, 29.94, , 28.55, 21.63, 19.04, 13.12.

[bis-1,3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) Chloride 4.31 5-



Following the procedure for the synthesis of complex was obtained from **3.41** (300mg, 0.600mmol) and $[\text{IrCl}(\text{cod})]_2$ (200 mg, 0.30mmol) . Yield: 261 mg (62.14%). Crystals suitable for X- ray crystallography were grown by layering hexane on dichloromethane. Anal. Calcd. for $\text{C}_{31}\text{H}_{30}\text{N}_4\text{IrCl}$: C, 54.23; H, 4.37; N, 8.16%. Found: C, 53.73; H, 4.30; N, 7.71%. ^1H NMR (CDCl_3 , 400MHz, 298K): δ 8.1(m, 2H, $J = 8.5$ Hz, quin-H), 8.00(d, 2H, $J = 8.5$ Hz, quin-H), 7.75(d, 2H, $J = 8.2$ Hz, quin-H), 7.65(m, 4H, $J = 4.6$ Hz, quin-H), 7.50(t, 2H, $J = 7.0$ Hz, quin-H), 6.85(s, 2H, CHHC), 6.25(d, 1H, $J = 14.9$ Hz, CH_2 linker), 5.65(d, 1H, $J = 14.9$ Hz, CH_2 linker), 4.60(m, 2H, $J = 2.8$ Hz, COD), 2.90(m, 2H, $J = 3.0$ Hz, COD), 2.10(m, 4H, $J = 3.4$ Hz, COD), 1.70(m, 4H, $J = 11.6$ Hz, COD). ^{13}C NMR (CDCl_3 , 100MHz, 298K): 180.57(C-Ir), 155.43, 146.52, 136.48, 128.81, 128.03, 126.74, 126.60, 125.75, 120.25, 119.60, 84.73(CH_2) 64.85, 55.53, 51.48, 32.47, 28.48, 14.26.

1-methyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) tetrafluoroborate 4.32



AgBF_4 (18 mg, 0.092mmol) was added to a stirred solution of complex **4.27**(50 mg, 0.089mmol) in 10 ml of DCM. The reaction mixture was stirred for 1 hour and reaction filtered through celite to remove silver chloride and any insoluble

residue. The filtrates was then added drop by drop to a stirring solution of hexane to precipitate the compound which was then dried in vacuum to afford the desired compound as a deep yellow powder(42.00 mg, 60.87%). Anal. Calcd. for $C_{22}H_{25}N_3IrBF_4$: C, 43.33; H, 4.09; N, 6.89%. Found: C, 35.11; H, 3.10; N, 5.09%. 1H NMR ($CDCl_3$, 400MHz, 298K): δ 8.80(d, 1H, $J = 8.6$ Hz, quin-H), 8.25(d, 1H, $J = 8.4$ Hz, quin-H), 8.10(d, 1H, $J = 8.4$ Hz, quin-H), 7.90(t, 1H, $J = 8.2$ Hz, quin-H), 7.85(t, 1H, $J = 8.2$ Hz, quin-H), 7.60(t, 1H, $J = 7.7$ Hz, quin-H), 7.55(s, 1H, CHHC), 6.80(s, 1H, CHHC), 5.90(d, $J = 15.4$ Hz, 1H, $CH_{2linker}$), 5.70(d, 1H, $J = 15.4$ Hz, $CH_{2linker}$), 5.50(broad, 1H, COD), 4.70(m, 1H, $J = 5.2$ Hz, COD), 4.00(m, 2H, $J = 8.4$ Hz, COD), 3.80(m, 1H, $J = 4.2$ Hz, COD), 3.70(s, 3H, CH₃), 2.75(m, 1H, $J = 10.2$ Hz, COD), 2.3(m, 4H, $J = 7.4$ Hz, COD), 2.0(m, 2H, $J = 6.7$ Hz, COD).

4.5.2: Catalysis

General Comments. All air sensitive experiments were performed under nitrogen atmosphere in an MBraun glove box or under dinitrogen by standard Schlenk techniques. Isopropanol was distilled from calcium hydride under N_2 atmosphere. The iridium complexes were synthesised as described above and 1H NMR spectra were recorded using a Bruker Advance DPX₄₀₀ spectrometer.

4.5.3 Transfer hydrogenation

The iridium catalyst precursor was dissolved in a solution of K^tBuO (1 mmol) in 2-propanol and 4-bromoacetophenone (1 mmol) was added in a Schlenk tube. The solution heated to 353K for 24 hours, volatiles were evaporated and the percentage conversion was calculated by 1H NMR.

The progress of the reaction was monitored by GC-MS analysis in order to calculate the time dependence of the transfer hydrogenation of 4-bromoacetophenone. Aliquots of 0.1 mL were taken every 10 minutes for the first 1 hour and every 30 minutes for the next hours. The samples were filtered through a short pad of silica, and the silica was washed with DCM.

4.5.4 X-Ray crystallography

Standard conditions as outlined in section 2.4.7 were used.

Table 4.14: Crystal data and structure refinement for 4.20

| | | |
|--------------------------------------------|-----------------------------------------------------|------------------------------|
| Empirical formula | C ₂₂ H ₂₅ RhN ₃ Cl | |
| Formula weight | 469.81 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Triclinic | |
| Space group | P-1 | |
| Unit cell dimensions | a = 7.5160(2) Å | α = 94.2310(10) ^o |
| | b = 11.4580(3) Å | β = 99.2930(10) ^o |
| | c = 11.6750(4) Å | γ = 99.204(2) ^o |
| Volume | 974.31(5) Å ³ | |
| Z | 2 | |
| Density (calculated) | 1.602Mg/m ³ | |
| Absorption coefficient | 1.025mm ⁻¹ | |
| F000 | 480 | |
| Crystal size | 0.13 x 0.07 x 0.01 mm ³ | |
| Theta range for data collection | 3.03 to 27.47 ° | |
| Index ranges | -9<=h<=9, -14<=k<=14, -15<=l<=15 | |
| Reflections collected | 15668 | |
| Independent reflection | 4449[R _{int} = 0.1609] | |
| Completeness of theta = 27.47 ^o | 99.50% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.8782 and 0.9898 | |
| Refinement method | Full matrix least square on F ² | |
| Data/restraints/parameters | 4449/0/245 | |
| Goodness of fit on F ² | 1.044 | |
| Final R indices [I>2 sigma(I)] | R1 = 0.1609, wR2 = 0.1153 | |
| R indices (all data) | R1 = 0.641, wR2 = 0.1488 | |
| Largest diff. peak hole | 1.426 and -2.812e Å ⁻³ | |

Table 4.15: Crystal data and structure refinement for 4.21

| | | |
|-----------------------------------|-----------------------------------------------------|---------|
| Empirical formula | C ₃₀ H ₃₃ RhN ₃ Cl | |
| Formula weight | 573.95 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Orthorhombic | |
| Space group | P212121 | |
| Unit cell dimensions | a = 9.9380(2) Å | α = 90° |
| | b = 14.6240(3) Å | β = 90° |
| | c = 17.8520(4) Å | γ = 90° |
| Volume | 2594.49(9) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.469Mg/m ³ | |
| Absorption coefficient | 0.785mm ⁻¹ | |
| F000 | 1184 | |
| Crystal size | 0.10 x 0.05 x 0.05 mm ³ | |
| Theta range for data collection | 3.03 to 27.51 ° | |
| Index ranges | -10 ≤ h < 12, -18 ≤ k ≤ 18, -23 ≤ l ≤ 23 | |
| Reflections collected | 43795 | |
| Independent reflection | 5932[R _{int} = 0.0689] | |
| Completeness of theta = 27.51° | 99.60% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.9618 and 0.9256 | |
| Refinement method | Full matrix least square on F ² | |
| Data/restraints/parameters | 5932/0/319 | |
| Goodness of fit on F ² | 1.053 | |
| Final R indices [I > 2 sigma(I)] | R1 = 0.0537, wR2 = 0.0882 | |
| R indices (all data) | R1 = 0.0423, wR2 = 0.0836 | |
| Largest diff. peak hole | 0.560 and -0.722e Å ⁻³ | |

Table 4.16: Crystal data and structure refinement for 4.22

| | | |
|-----------------------------------|-----------------------------------------------------|---------|
| Empirical formula | C ₂₄ H ₃₀ RhN ₃ Cl | |
| Formula weight | 540.33 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Orthorhombic | |
| Space group | P212121 | |
| Unit cell dimensions | a = 13.1930(10) Å | α = 90° |
| | b = 20.8640(2) Å | β = 90° |
| | c = 8.5280(4) Å | γ = 90° |
| Volume | 2347.41(11) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.529Mg/m ³ | |
| Absorption coefficient | 0.972mm ⁻¹ | |
| F000 | 1108 | |
| Crystal size | 0.25 x 0.22 x 0.10 mm ³ | |
| Theta range for data collection | 3.01 to 27.61 ° | |
| Index ranges | -17<=h< 17, -26<=k<=27, -10<=l<=11 | |
| Reflections collected | 38012 | |
| Independent reflection | 5371[R _{int} = 0.0669] | |
| Completeness of theta = 27.61° | 98.70% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.9090 and 0.7931 | |
| Refinement method | Full matrix least square on F ² | |
| Data/restraints/parameters | 5371/3/292 | |
| Goodness of fit on F ² | 1.057 | |
| Final R indices [I>2 sigma(I)] | R1 = 0.0669, wR2 = 0.1103 | |
| R indices (all data) | R1 = 0.0457, wR2 = 0.0996 | |
| Largest diff. peak hole | 0.840 and -0.902e Å ⁻³ | |

Table 4.17: Crystal data and structure refinement for 4.23

| | |
|-------------------|-----------------------------------------------------|
| Empirical formula | C ₂₅ H ₃₁ RhN ₃ Cl |
| Formula weight | 511.89 |

| | |
|-----------------------------------|----------------------------------------------------------------------------------------------------------------------------|
| Temperature | 150(2) K |
| Wavelength | 0.71073 Å |
| Crystal system | Triclinic |
| Space group | P-1 |
| Unit cell dimensions | a = 10.0680(3) Å α = 95.0950(10)° b = 10.6960(3) Å β = 91.6400(10)° c = 10.8350(4) Å γ = 98.0910(10)° |
| Volume | 1149.61(6) Å ³ |
| Z | 2 |
| Density (calculated) | 1.479Mg/m ³ |
| Absorption coefficient | 0.876mm ⁻¹ |
| F000 | 528 |
| Crystal size | 0.50 x 0.50 x 0.05 mm ³ |
| Theta range for data collection | 3.01 to 27.53 ° |
| Index ranges | -12 ≤ h < 13, -13 ≤ k ≤ 13, -14 ≤ l ≤ 14 |
| Reflections collected | 19594 |
| Independent reflection | 5209[R _{int} = 0.0956] |
| Completeness of theta = 27.53° | 98.50% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | 0.9575 and 0.6686 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 5371/3/292 |
| Goodness of fit on F ² | 1.060 |
| Final R indices [I > 2σ(I)] | R1 = 0.0851, wR2 = 0.1124 |
| R indices (all data) | R1 = 0.0504, wR2 = 0.0978 |
| Largest diff. peak hole | 0.747 and -1.059e Å ⁻³ |

Table 4.18: Crystal data and structure refinement for 4.24

| | |
|-------------------|-----------------------------------------------------|
| Empirical formula | C ₃₁ H ₃₀ RhN ₄ Cl |
| Formula weight | 600.90 |
| Temperature | 150(2) K |
| Wavelength | 0.71073 Å |
| Crystal system | Triclinic |

| | | |
|-----------------------------------|--------------------------------------------|------------------------------|
| Space group | P-1 | |
| Unit cell dimensions | a = 11.8769(2) Å | $\alpha = 89.9829(10)^\circ$ |
| | b = 11.9874(2) Å | $\beta = 118.0475(10)^\circ$ |
| | c = 12.3176(2) Å | $\gamma = 106.1171(9)^\circ$ |
| Volume | 1469.26(4) Å ³ | |
| Z | 2 | |
| Density (calculated) | 1.541Mg/m ³ | |
| Absorption coefficient | 0.883mm ⁻¹ | |
| F000 | 696 | |
| Crystal size | 0.25 x 0.25 x 0.38 mm ³ | |
| Theta range for data collection | 3.114 to 27.643 ° | |
| Index ranges | -15<=h< 15, -15<=k<=15, -15<=l<=16 | |
| Reflections collected | 24833 | |
| Independent reflection | 11368[R _{int} = 0.052] | |
| Completeness of theta = 27.643° | 97.20% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.800 and 0.800 | |
| Refinement method | Full matrix least square on F ² | |
| Data/restraints/parameters | 11368/0/362 | |
| Goodness of fit on F ² | 0.9877 | |
| Final R indices [I>2 sigma(I)] | R1 = 0.0526, wR2 = 0.1120 | |
| R indices (all data) | R1 = 0.0461, wR2 = 0.1073 | |
| Largest diff. peak hole | 1.63 and -1.88e Å ⁻³ | |

Table 4.19: Crystal data and structure refinement for 4.28

| | | |
|----------------------|-----------------------------------------------------|---------------------|
| Empirical formula | C ₃₀ H ₃₃ IrN ₃ Cl | |
| Formula weight | 663.28 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Orthorhombic | |
| Space group | P212121 | |
| Unit cell dimensions | a = 9.9463(2) Å | $\alpha = 90^\circ$ |
| | b = 14.5997(3) Å | $\beta = 90^\circ$ |

| | | |
|----------------------------------------|-----------------------------------------------------------|---------------------|
| | $c = 17.8669(5) \text{ \AA}$ | $\gamma = 90^\circ$ |
| Volume | $2594.51(10) \text{ \AA}^3$ | |
| Z | 4 | |
| Density (calculated) | 1.698 Mg/m^3 | |
| Absorption coefficient | 5.273 mm^{-1} | |
| F000 | 1312 | |
| Crystal size | $0.04 \times 0.17 \times 0.38 \text{ mm}^3$ | |
| Theta range for data collection | $3.014 \text{ to } 27.394^\circ$ | |
| Index ranges | $-12 \leq h < 10, -18 \leq k \leq 18, -23 \leq l \leq 23$ | |
| Reflections collected | 42001 | |
| Independent reflection | 5841 [$R_{\text{int}} = 0.199$] | |
| Completeness of theta = 27.394° | 99.30% | |
| Absorption correction | Semi-empirical from equivalents | |
| Max and min. transmission | 0.8100 and 0.4100 | |
| Refinement method | Full matrix least square on F^2 | |
| Data/restraints/parameters | 5841/0/317 | |
| Goodness of fit on F^2 | 0.5948 | |
| Final R indices [$I > 2 \sigma(I)$] | $R1 = 0.0740, wR2 = 0.1219$ | |
| R indices (all data) | $R1 = 0.0422, wR2 = 0.997$ | |
| Largest diff. peak hole | $2.85 \text{ and } -3.59 \text{ e \AA}^{-3}$ | |

Table 4.20: Crystal data and structure refinement for 4.29

| | | |
|----------------------|---------------------------------------------------|---------------------|
| Empirical formula | $\text{C}_{24}\text{H}_{30}\text{IrN}_3\text{Cl}$ | |
| Formula weight | 629.61.28 | |
| Temperature | $150(2) \text{ K}$ | |
| Wavelength | 0.71073 \AA | |
| Crystal system | Orthorhombic | |
| Space group | P212121 | |
| Unit cell dimensions | $a = 8.562(2) \text{ \AA}$ | $\alpha = 90^\circ$ |
| | $b = 13.267(3) \text{ \AA}$ | $\beta = 90^\circ$ |
| | $c = 21.012(5) \text{ \AA}$ | $\gamma = 90^\circ$ |
| Volume | $2386.80(10) \text{ \AA}^3$ | |
| Z | 4 | |

| | |
|-----------------------------------|--------------------------------------------|
| Density (calculated) | 1.752Mg/m ³ |
| Absorption coefficient | 5.835mm ⁻¹ |
| F000 | 1236 |
| Crystal size | 0.20 x 0.20 x 0.10 mm ³ |
| Theta range for data collection | 2.57 to 33.73 ° |
| Index ranges | -13<=h< 13, -20<=k<=20, -32<=l<=32 |
| Reflections collected | 9511 |
| Independent reflection | 9511[R _{int} = 0.0589] |
| Completeness of theta = 33.73° | 99.80% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | 0.5930 and 0.3882 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 9511/9/291 |
| Goodness of fit on F ² | 1.036 |
| Final R indices [I>2 sigma(I)] | R1 = 0.065, wR2 = 0.1242 |
| R indices (all data) | R1 = 0.0500, wR2 = 0.1144 |
| Largest diff. peak hole | 3.528 and -2.850e Å ⁻³ |

Table 4.21: Crystal data and structure refinement for 4.31

| | | |
|------------------------|-----------------------------------------------------|------------------|
| Empirical formula | C ₃₁ H ₃₀ IrN ₄ Cl | |
| Formula weight | 686.28 | |
| Temperature | 150(2) K | |
| Wavelength | 0.71073 Å | |
| Crystal system | Triclinic | |
| Space group | P - 1 | |
| Unit cell dimensions | a = 6.59420(10) Å | α = 104.4358(8)° |
| | b = 12.1294(2) Å | β = 100.9896(8)° |
| | c = 17.2628(3) Å | γ = 97.8041(10)° |
| Volume | 1298.61(4) Å ³ | |
| Z | 4 | |
| Density (calculated) | 1.755Mg/m ³ | |
| Absorption coefficient | 5.272mm ⁻¹ | |
| F000 | 676 | |

| | |
|-----------------------------------|--------------------------------------------|
| Crystal size | 0.15 x 0.20 x 0.38 mm ³ |
| Theta range for data collection | 3.158 to 27.503 ° |
| Index ranges | -8<=h< 7, -15<=k<=15, -21<=l<=22 |
| Reflections collected | 22471 |
| Independent reflection | 21713[R _{int} = 0.073] |
| Completeness of theta = 23.503° | 99.50% |
| Absorption correction | Semi-empirical from equivalents |
| Max and min. transmission | 0.45 and 0.35 |
| Refinement method | Full matrix least square on F ² |
| Data/restraints/parameters | 10018/0/335 |
| Goodness of fit on F ² | 0.9719 |
| Final R indices [I>2 sigma(I)] | R1 = 0.0469, wR2 = 0.1073 |
| R indices (all data) | R1 = 0.0417, wR2 = 0.1029 |
| Largest diff. peak hole | 2.13 and -1.79e Å ⁻³ |

References:

1. A. J. Arduengo, R. L. Harlow, M. J. Kline, *J. Am. Chem. Soc.*, **1991**, *113*, 361.
2. K. J. Ofele, *Organomet. Chem.* **1968**, *21*, P42.
3. H. W. Wanzlick, H. J. Schonherr, *Angew. Chem. Int. Edit.* **1968**, *7*, 141.
4. M. F. Lappert, *Organomet. Chem.* **1975**, *100*, 139.
5. R. S. Bon, F. J. J. de Kanter, M. Lutz, A. L. Spek, M. C. Jahnke, F. E. Hahn, M. B. Groen and R. V. Orru, *Organometallics*, **2007**, *26*, 3639-3650.
6. E. Mas-Marza, M. Sanau and E. Peris, *Inorganic Chem.* **2005**, Vol. *44*, No. *26*, 9962.
7. W. A. Herrmann, L. J. Gooben and M. Spiegler, *J. Organomet. Chem.* **1997**, *547*, 357-366.

8. M. V. Jimenez, J. J. Perez-Torrente, M. I. Bartolome, V. Gierz, F. J. Lahoz and L. A. Oro, *Organometallics*, **2008**, *27*, 224- 234.
9. N. Stylianides, A. A. Danopoulos, N. Tsoureas, *J. Organomet. Chem.*, **2005**, *690*, 5948-5958.
10. K. S. Coleman, H. T. Chambarlayne, S. Turberville, M. L.H. Green and A. R. Cowley, *J. Chem. , Dalton Trans.*, **2003**, 2917-2922.
11. C. Wang, C. Fu, Y. Liu, S. Peng and S. Liu, *Inorganic Chem.*, 2007.
12. C. H. Leung, C. D. Incarvito and R. H. Crabtree, *Organometallics*, **2006**, *25*, 6099-6107.
13. D. S.McGuinness, K. J. Cavell, B. F. Yates, B. W. Skelton, A. H. White, *J. Am. Chem. Soc.* **2001**, *123*, 8317.
14. 15. H. M.Wang and I. J. B. Lin, *Organometallics*, **1998**, *19*, 972
15. A. A. Tulloch, A. A. Danopoulos, G. J. Tizzard, S. J. Coles, M. B. Hursthouse, R. S.Hay-motherwell, W. B. Motherwell, *Chem. Commun.***2001**, 1270.
16. R. S. Simons, P. Custer, C. A. Tessier, W. J. Youngs, *Organometallics*, **2003**, *22*, 1979.
17. D. S.McGuinness and K. J. Cavell , *Organometallics* **2000**, *19*, 741.
18. M. Payatos, E. Mas-Marza, J. A. Mata, M. Sanau, E. Peris, *Eur. J. Inorg. Chem.***2003**, 1215.
19. M. Payatos, M. Sanau, E. Peris, *Inorg. Chem.* **2003**, *42*, 2572.
20. M. Payatos, E. Mas-Marza, M. Sanau, E. Peris, *Inorg. Chem.* **2004**, *43*, 2213.
21. E. Mas-Marza, E. Peris, I. Castro-Rodriguez, K. Meyer, *Organometallics*, **2005**,*24*, 3158.
22. J, A. Mata, A. R. Chianese, J. R. Miecznikowski, M. Payatos, E. Peris, J. W. Faller, R. H. Crabtree, *Organometallics* **2004**, *23*, 1253. 23. X. Yu, B. O. Patrick and B. R. James, *Organometallics*, **2006**, *25*, 2359-2363.
24. A.R. Chianese, X. Li, M. C. Janzen, J. W. Faller, R. H. Crabtree,

- Organometallics* **2003**, *22*, 1663.
25. M. Prinz, L. F. Veiros, M. J. Calhorda, C. C. Romao, E. Herdtweck, F. E. Kuhn and W. A. Herrmann, *J. Organomet. Chem.* **2006**, *691*, 4446-4458.
26. P. J. Alaimo, R. G. Bergman, *Organometallics*, **1999**, *18*, 2707.
27. J. T. Richardson, *Principle of catalyst development*; Plenum Press: New York, **1989**.
28. J. Hagen, *Industrial catalysis: a practical approach*; Wiley- VCH: Weinheim, **1999**.
29. B. Cornils and W. A. Herrmann, Eds. *Applied homogeneous catalysis with Organometallics compounds: A comprehensive textbook in two volumes*; VCH: Weinheim, **1996**.
30. A. Zapf and M. Beller, *Topics in catalysis*, **2002**, *19*, 101-109.
31. H. M. Lee, T. Jiang, E. D. Stevens, S. P. Nolan, *Organometallics*, **2001**, *20*, 1255.
32. A. T. Normand and K. J. Cavell, *Eur. J. Inorg. Chem.*, **2008**, 2781-2800.
33. M. Albrecht, J. R. Miecznikowski, A. Samuel, J.W. Faller, R. H. Crabtree, *Organometallics*, **2002**, *21*, 3596.
34. C. Sauzzo, M. Lemaire, *Adv. Synth. Catal.* **2002**, *344*, 915.
35. S. Hashiguci, A. Fujii, J. Takehara, T. Ikariya, R. Noyori, *J. Am. Chem. Soc.* **1995**, *117*, 7562-7563.
36. L. D. Vazquez-Serrano, B. T. Owens, J. M. Buriak, *Chem. Commun.*, **2002**, 2518-2519.
37. M. Viciano, E. Mas-Marza, M. Poyatos, M. Sanau, R.H. Crabtree, E. Peris, *Angew. Chem., Int. Ed.* **2005**, *44*, 444.
38. E. Mas-Marza, M. Sanau, and E. Peris, *Inorg. Chem.* **2005**, *44*, 9961-9967.
39. H. Turkmen, T. Pape, E. Hahn and B. Cetinkaya, *Organometallics* **2008**, *27*, 571-575.
40. W. L. F. Armarego and D. D. Perrin, *Purification of Laboratory Chemicals* 4th ed., Butterworth: Bath, **1996**.

CHAPTER FIVE

5.1 Conclusions

Ligands containing both phosphorus and nitrogen donor have been shown to form strong metal phosphorus bonds and weak nitrogen bonds [1] and are important in many catalytic reactions [2, 3]. In realisation of this many groups have made the natural progression from unidentate mono carbenes towards mixed donor chelating carbenes. The interest stems from the potential advantages a hemilabile ligand may offer to a catalytic reaction. A number of research groups have successfully incorporated a second donor to the carbene ligand, with early progress being made with pyridine functions as the N-substituents [1, 4, and 5]. In furtherance of the use of a hemilabile group on N-substituents quinoline based imidazolium salts were thought to be ideal candidates. Towards this end, a series of quinoline based imidazolidinium salts have been synthesised and characterised as precursors to the corresponding NHC ligands. A range of N-substituents imparts variable steric bulk to the imidazolium rings.

Crystallographically characterised examples include the saturated 1, 3-diquinolin-4, 5-dihydroimidazolium tetrafluoroborate **2b**, the unsaturated imidazolium salts **6a-6c**, and the analogous methylene bridged quinoline-functionalised imidazolium salts **9a- 9f**.

An additional example reported includes the acridine based imidazolium salt **16** which can offer a secondary donor group for chelation as well as sp³ hybridized carbons. The sp³ hybridized carbon can make the ligand chiral and increase the steric bulk if required.

Quinoline based pyrimidinium salt **4** was also prepared following the same procedure for **2b** was crystallographically characterised. All the quinoline based salts are new compound as there is no reported synthesis of any of them

Ag^I(NHC) complexes have reported to be a versatile transmetallation reagents for the preparation of transition metal- NHC complexes. Accordingly, a range of Ag^I complexes of methylene bridge quinoline functionalised NHC ligands were prepared by reaction of the corresponding imidazolium salts with Ag₂O in

DCM. Of these $\text{Ag}^{\text{I}}(\text{NHC})$ complexes **3.36-3.41**, two (**3.36** and **3.37**) were Crystallographically characterised with the silver geometry in both cases being a quasi linear. Additional Ag^{I} complexes of quinoline functionalised NHC ligands were synthesised in a similar manner and utilised as transmetallation agent in the synthesis of $\text{Pd}^{\text{II}}(\text{NHC})$ complexes **3.42** and **3.43**. Efforts to prepare other $\text{Pd}^{\text{II}}(\text{NHC})$ complexes were not successful due to high insolubility of the complexes in most solvents. Indeed difficulty was encountered in separating the desired complexes from the silver halides.

The synthesis of a series of $\text{Rh}(\text{I})$ (NHC) of the methylene bridged quinoline functionalised NHC ligands via transmetallation from $\text{Ag}(\text{I})$ complexes is reported. All the rhodium complexes were fully characterised and the crystallographic data show consistent pattern. The reaction of the $[\text{Rh}(\text{cod})\text{Cl}]_2$ with Ag_2O gave the neutral rhodium complexes **4.20-4.24** and there was no evidence of chelation between the nitrogen of the quinoline ring and the rhodium metal as evidenced by the ^1H NMR data being essentially similar to that of the corresponding $\text{Ag}(\text{I})$ complexes from which they were made.


Chelation was achieved by the reaction of $\text{Rh}(\text{I})$ (NHC) complexes with one equivalent of AgBF_4 in DCM. However the chelated $\text{Rh}(\text{I})$ (NHC) complexes could not be fully characterised due to high insolubility of the compounds in most solvent. Attempts to improve the solubility by replacing the tetraflouoroborate counter ion with Barf solve the problem of solubility but difficulty was encountered in separation.

In a similar fashion $\text{Ir}(\text{I})$ (NHC) complexes **4.27-4.32** were synthesised via transmetallation of the corresponding $\text{Ag}(\text{I})$ complexes with $[\text{Ir}(\text{cod})\text{Cl}]_2$ in DCM. The trend observed for the rhodium complexes are essentially similar to that of the iridium complexes. Thus both chelated and non chelated $\text{Ir}(\text{I})$ (NHC) complexes were prepared.

Finally two of the neutral $\text{Ir}(\text{I})$ (NHC) complexes prepared **4.27** and **4.31** were catalytically tested towards transfer hydrogenation of 4-bromoacetophenone. The two iridium complexes have shown good activity upon hydrogen transfer reduction of carbonyl in 4-bromoacetophenone at different catalyst concentrations.

5.2 Future work

Attempts to prepare the unsymmetrical quinoline based unsaturated imidazolium and pyrimidinium salts following the established procedures [5] were not successful. However it will be interesting if the following quinoline and acridine based ligands will be prepared and investigated.



It will be interesting to compare these potential pincer ligands with the pincer ligands reported by Gibson, Cavell, Danopoulos and Crabtree.

Also to be investigated are the metal complexes **2b** and **4** which we were unable to prepare in this work.

It is recommended that the quinoline based palladium complexes prepared in this work be tested in some important catalytic reactions such as Heck coupling reactions.

The Rh(I) and Ir(I) (NHC) complexes should be tested in different catalytic reaction such as reduction of alkenes via direct hydrogenation and transfer hydrogenation. Efforts should be to isolate pure soluble chelated versions of the

Rh(I) and Ir(I) (NHC) complexes, their catalytic activities compared with that of the unchelated complexes and the results obtained compared with that of the iridium pyridinyl N-heterocyclic carbene complexes reported by Wang et al [6]. Other interesting future work will be to look into the metal complexes of the ligands reported in chapter two such as Pt, Ni, Co, and Fe.

5.3 References

1. J. Chen, I. Lin, *Organometallics*, **2000**, *19*, 5113-5121.
2. D. McGuinness, N. Saendig, B. Yates, K. J. Cavell, *J. Am. Chem. Soc.*, **2001**, *123*, 4029-4040.
3. D. McGuinness, W. Mueller, P. Wasserscheid, K. J. Cavell, B. Skelton, A. White, U. Englert, *Organometallics*, **2002**, *21*, 4318-4319.
3. D. McGuinness, K.J. Cavell, *Organometallics*, **2000**, *19*, 741.
4. J. Chen, I. Lee, *J. Chem. Soc., Dalton Trans*, **2000**, 839.
5. R. Jazzar, H. Liang, B. Donnadieu, G. Bertrand, *J. Organomet. Chem.* **2006**, *691*, 3201-3205.
6. Wang, C. Fu, Y. Liu, S. Peng and S. Liu, *Inorganic Chem.*, **2007**.

APPENDIX

Tables of bond distances and angles

Table A1.1: Bond lengths for 1, 3-diquinoline-3, 4, 5, 6-tetrahydropyrimidinium hexafluoro phosphate (4):

| Bond lengths (Å) | | Bond lengths (Å) | | Bond lengths (Å) | |
|------------------|----------|------------------|----------|------------------|------------|
| C1- N2 | 1.317(3) | C9- C13 | 1.419(3) | C21- N4 | 1.323(3) |
| C1- N1 | 1.317(3) | C10 -C11 | 1.351(3) | C21- H21 | 0.9500 |
| C1- H1 | 0.9500 | C10- H10 | 0.9500 | C22- N4 | 1.372(3) |
| C2- N1 | 1.479(3) | C11- C12 | 1.413(3) | F1-P1 | 1.6054(16) |
| C2- C3 | 1.513(3) | C11- H11 | 0.9500 | F2-P1 | 1.5933(18) |
| C2- H2A | 0.9900 | C12- N3 | 1.322(3) | F3-P1 | 1.5912(16) |
| C2 -H2B | 0.9900 | C12- H12 | 0.9500 | F4-P1 | 1.5876(18) |
| C3- C4 | 1.507(3) | C13 -N3 | 1.370(3) | F5-P1 | 1.5784(18) |
| C3- H3A | 0.9900 | C14-C15 | 1.371(3) | F6-P1 | 1.5853(17) |
| C3- H3B | 0.9900 | C14- C22 | 1.425(3) | | |
| C4 -N2 | 1.485(3) | C14 -N2 | 1.429(3) | | |
| C4- H4A | 0.9900 | C15- C16 | 1.405(3) | | |
| C4- H4B | 0.9900 | C15- H15 | 0.9500 | | |
| C5 -C6 | 1.365(3) | C16- C17 | 1.365(3) | | |
| C5- C13 | 1.422(3) | C16 -H16 | 0.9500 | | |
| C5- N1 | 1.439(3) | C17- C18 | 1.415(3) | | |
| C6- C7 | 1.417(3) | C17 -H17 | 0.9500 | | |
| C6- H6 | 0.9500 | C18- C22 | 1.415(3) | | |
| C7 -C8 | 1.356(4) | C18- C19 | 1.422(3) | | |
| C7- H7 | 0.9500 | C19- C20 | 1.353(4) | | |
| C8- C9 | 1.417(3) | C19 -H19 | 0.9500 | | |
| C8- H8 | 0.9500 | C20- C21 | 1.403(4) | | |
| C9- C10 | 1.418(3) | C20- H20 | 0.9500 | | |
| Bond angles (°) | | Bond angles (°) | | Bond angles (°) | |
| N2- C1- N1 | 124.2(2) | C7- C8- H8 | 119.6 | C20-C19-C18 | 119.4(2) |
| N2- C1- H1 | 117.9 | C9- C8- H8 | 119.6 | C20- C19- H19 | 120.3 |

Appendix

| | | | | | |
|--------------|------------|--------------|------------|---------------|------------|
| N1- C1- H1 | 117.9 | C8-C9-C10 | 122.8(2) | C18- C19- H19 | 120.3 |
| N1-C2-C3 | 109.03(18) | C8-C9-C13 | 119.3(2) | C19-C20-C21 | 119.2(2) |
| N1- C2- H2A | 109.9 | C10-C9-C13 | 117.9(2) | C19-C20-H20 | 120.4 |
| C3- C2- H2A | 109.9 | C11-C10-C9 | 119.5(2) | C21-C20-H20 | 120.4 |
| N1- C2- H2B | 109.9 | C11-C10-H10 | 120.3 | N4-C21-C20 | 124.6(2) |
| C3- C2- H2B | 109.9 | C9- C10- H10 | 120.3 | N4-C21-H21 | 117.7 |
| H2A- C2- H2B | 108.3 | C10-C1-C12 | 118.7(2) | C20-C21-H21 | 117.7 |
| C4-C3-C2 | 109.48(19) | C10-C11-H11 | 120.7 | N4-C22-C18 | 123.2(2) |
| C4- C3- H3A | 109.8 | C12-C11-H11 | 120.7 | N4-C22-C14 | 118.6(2) |
| C2- C3- H3A | 109.8 | N3-C12-C11 | 124.6(2) | C18-C22-C14 | 118.2(2) |
| C4- C3- H3B | 109.8 | N3-C12-H12 | 117.7 | C1-N1-C5 | 120.71(18) |
| C2- C3- H3B | 109.8 | C11-C12-H12 | 117.7 | C1-N1-C2 | 120.41(18) |
| H3A- C3 -H3B | 108. | N3-C13-C9 | 122.2(2) | C5-N1-C2 | 118.73(18) |
| N2-C4-C3 | 108.91(17) | N3-C13-C5 | 119.63(19) | C1-N2-C14 | 120.32(18) |
| N2- C4- H4A | 109.9 | C9-C13-C5 | 118.18(19) | C1-N2-C4 | 120.78(19) |
| C3- C4- H4A | 109.9 | C15-C14-C22 | 120.7(2) | C14-N2-C4 | 118.88(17) |
| N2- C4- H4B | 109.9 | C15-C14-N2 | 120.38(19) | C12-N3-C13 | 117.06(19) |
| C3- C4 -H4B | 109.9 | C22-C14-N2 | 118.9(2) | C21-N4-C22 | 116.5(2) |
| H4A- C4- H4B | 108.3 | C14-C15-C16 | 120.3(2) | F5-P1-F6 | 91.17(11) |
| C6- C5 -C13 | 121.4(2) | C14-C15-H15 | 119.8 | F5-P1-F4 | 90.55(12) |
| C6- C5- N1 | 120.3(2) | C16-C15-H15 | 119.8 | F6-P1-F4 | 90.78(11) |
| C13-C5-N1 | 118.33(19) | C17-C16-C15 | 120.6(2) | F5-P1-F3 | 89.22(10) |
| C5- C6- C7 | 119.7(2) | C17-C16-H16 | 119.7 | F6-P1-F3 | 179.52(11) |
| C5- C6- H6 | 120.2 | C15-C16-H16 | 119.7 | F4-P1-F3 | 89.49(10) |
| C7 -C6 -H6 | 120.2 | C16-C17-C18 | 120.3(2) | F5-P1-F2 | 179.18(11) |
| C8- C7- C6 | 120.6(2) | C16-C17-H17 | 119.8 | F6-P1-F2 | 89.24(11) |
| C8- C7 -H7 | 119.7 | C18-C17-H17 | 119.8 | F3-P1-F2 | 90.36(10) |
| C6- C7- H7 | 119.7 | C22-C18-C17 | 119.8(2) | F5-P1-F1 | 90.66(10) |
| C7- C8 -C9 | 120.9(2) | C22-C18-C19 | 117.1(2) | F6-P1-F1 | 90.50(10) |
| | | C17-C18-C19 | 123.1(2) | F4-P1-F1 | 178.22(11) |
| | | | | F3-P1-F1 | 89.22(9) |
| | | | | F2-P1-F1 | 88.64(9) |

Appendix

Table A1.2: Bond Lengths and angles for 1-benzyl-3-quinolinimidazolium tetrafluoroborate (**6a**):

| Bond lengths (Å) | | Bond lengths (Å) | |
|------------------|------------|------------------|----------|
| F1-B2 | 1.386(2) | C15-H151 | 0.994 |
| B2-F3 | 1.386(2) | C16-C17 | 1.404(3) |
| B2-F4 | 1.387(2) | C16-H161 | 0.942 |
| B2-F5 | 1.387(2) | C17-H171 | 0.944 |
| N6-C7 | 1.391(2) | C18-C19 | 1.419(2) |
| N6-C10 | 1.3405(19) | C18-C27 | 1.369(2) |
| N6-C18 | 1.4411(19) | C19-N20 | 1.367(2) |
| C7-C8 | 1.349(2) | C19-C24 | 1.424(2) |
| C7-H71 | 0.961 | N20-C21 | 1.315(2) |
| C8-N9 | 1.381(2) | C21-C22 | 1.416(2) |
| C8-H81 | 0.963 | C21-H211 | 0.942 |
| N9-C10 | 1.324(2) | C22-C23 | 1.361(2) |
| N9-C11 | 1.4846(19) | C22-H221 | 0.946 |
| C10-H101 | 0.961 | C23-C24 | 1.411(2) |
| C11-C12 | 1.498(2) | C23-H231 | 0.964 |
| C11-H111 | 1.005 | C24-C25 | 1.417(2) |
| C11-H112 | 0.922 | C25-C26 | 1.365(2) |
| C12-C13 | 1.389(3) | C25-H251 | 0.972 |
| C12-C17 | 1.376(3) | C26-C27 | 1.410(2) |
| C13-C14 | 1.381(3) | C26-H261 | 1.000 |
| C13-H131 | 0.922 | C27-H271 | 0.956 |
| C14-C15 | 1.360(4) | | |
| C14-H141 | 0.955 | | |
| C15-C16 | 1.373(4) | | |
| Bond angles (°) | | Bond angles (°) | |
| F1-B2-F3 | 108.2815) | C15-C16-H161 | 122.2 |
| F1-B2-F4 | 109.03(16) | C17-C16-H161 | 117.7 |
| F3-B2-F4 | 110.49(16) | C16-C17-C12 | 119.8(2) |
| F1-B2-F5 | 110.07(15) | C16-C17-H171 | 123.0 |

Appendix

| | | | |
|--------------|------------|--------------|------------|
| F3-B2-F5 | 109.55(16) | C12-C17-H171 | 117.1 |
| F4-B2-F5 | 109.40(15) | C14-C15-H151 | 118.7 |
| C7-N6-C10 | 108.30(13) | C16-C15-H151 | 120.9 |
| C7-N6-C18 | 124.30(13) | C15-C-16-C17 | 120.0(2) |
| C10-N6-C18 | 127.40(13) | N6-C18-C19 | 119.76(14) |
| N6-C7-C8 | 106.97(14) | N6-C18-C27 | 118.31(14) |
| N6-C7-H71 | 124.4 | C19-C18-C27 | 121.88(14) |
| C8-C7-H71 | 128.6 | C18-C19-N20 | 119.85(14) |
| C7-C8-N9 | 107.09(15) | C18-C19-C24 | 117.31(14) |
| C7-C8-H81 | 131.2 | N20-C19-C24 | 122.83(15) |
| N9-C8-H81 | 121.7 | C19-N20-C21 | 117.12(14) |
| C8-N9-C10 | 109.25(13) | N20-C21-C22 | 124.60(15) |
| C8-N9-C11 | 125.68(14) | N20-C21-H211 | 116.6 |
| C10-N9-C11 | 124.91(14) | C22-C21-H211 | 118.8 |
| N6-C10-N9 | 108.38(14) | C21-C22-C23 | 118.32(16) |
| N6-C10-H101 | 124.8 | C21-C22-H221 | 123.0 |
| N9-C10-H101 | 126.8 | C23-C22-H221 | 118.7 |
| N9-C11-C12 | 111.31(14) | C22-C23-C24 | 119.98(15) |
| N9-C11-H111 | 110.0 | C22-C23-H231 | 119.9 |
| C12-C11-H111 | 110.2 | C24-C23-H231 | 120.1 |
| N9-C11-H112 | 100.3 | C19-C24-C23 | 117.13(15) |
| C12-C11-H112 | 111.1 | C19-C24-C25 | 119.96(15) |
| H11-C11-H112 | 113.6 | C23-C24-C25 | 122.91(14) |
| C11-C12-C13 | 119.83(18) | C24-C25-C26 | 120.57(15) |
| C11-C12-C17 | 121.41(18) | C24-C25-H251 | 117.2 |
| C13-C12-C17 | 118.75(19) | C26-C25-H251 | 122.3 |
| C12-C13-C14 | 121.2(2) | C25-C26-C27 | 120.31(15) |
| C12-C13-H131 | 118.5 | C25-C26-H261 | 121.8 |
| C14-C13-H131 | 120.3 | C27-C26-H261 | 117.8 |
| C13-C14-C15 | 119.8 | C26-C27-C18 | 119.95(15) |
| C13-C14-H14 | 117.5 | C26-C27-H27 | 118.2 |
| C15-C14-H141 | 122.5 | C18-C27-H271 | 121.8 |
| C14-C15-C16 | 120.4(2) | | |

Table A1.3: Bond length for 1-methyl-3-quinolinimidazolium iodide (**6c**)

| Bond lengths (Å) | | Bond lengths (Å) | |
|------------------|----------|------------------|----------|
| C1-C2 | 1.367(4) | C8-N1 | 1.325(4) |
| C-C9 | 1.428(4) | C8-H8 | 0.9500 |
| C1-N2 | 1.439(4) | C9-N1 | 1.365(4) |
| C2-C3 | 1.411(4) | C10-N3 | 1.319(4) |
| C2-H2 | 0.9500 | C10-N2 | 1.340(4) |
| C3-C4 | 1.364(4) | C10-H10 | 0.9500 |
| C3-H3 | 0.9500 | C11-C12 | 1.347(4) |
| C4-C5 | 1.409(4) | C11-N2 | 1.386(4) |
| C4-H4 | 0.9500 | C11-H11 | 0.9500 |
| C5-C6 | 1.420(4) | C12-N3 | 1.382(4) |
| C5-C9 | 1.423(4) | C12-H12 | 0.9500 |
| C-C7 | 1.356(4) | C13-N3 | 1.461(4) |
| C6-H6 | 0.9500 | C13-H13A | 0.9800 |
| C7-C8 | 1.399(4) | C13-H13B | 0.9800 |
| C7-H7 | 0.9500 | C13-H13C | 0.9800 |

Table A1.4: Bond angles for 1-methyl-3-quinolinimidazolium iodide (**6c**)

| Bond angles (°) | | Bond angles (°) | |
|-----------------|----------|-----------------|----------|
| C2-C1-C9 | 121.3(3) | N1-C9-C1 | 119.5(2) |
| C2-C1-N2 | 118.7(2) | C5-C9-C1 | 117.2(2) |
| C9-C1-N2 | 119.8(2) | N3-C10-N2 | 108.9(2) |
| C1-C2-C3 | 120.5(3) | N3-C10-H10 | 125.5 |
| C1-C2-H2 | 119.8 | N2-C10-H10 | 125.5 |
| C3-C2-H2 | 119.8 | C12-C11-N2 | 106.9 |
| C4-C3-C2 | 120.0(3) | C12-C11-H11 | 126.6 |
| C4-C3-H3 | 120.0 | N2-C11-H11 | 126.6 |
| C2-C3-H3 | 120.0 | C11-C12-N3 | 107.5(3) |
| C3-C4-C5 | 120.7(3) | C11-C12-H12 | 126.3 |
| C3-C4-H4 | 119.7 | N3-C12-H12 | 126.3 |

| Appendix | | | |
|----------|----------|--------------|----------|
| C5-C4-H4 | 119.7 | N3-C13-H13A | 109.5 |
| C4-C5-C6 | 123.0(3) | N3-C13-H13B | 109.5 |
| C4-C5-C9 | 120.3(2) | H13-A13-13B | 109.5 |
| C6-C5-C9 | 116.7(3) | N3-C13-H13C | 109.5 |
| C7-C6-C5 | 119.5(3) | H13-C13-H13 | 109.5 |
| C7-C6-H6 | 120.2 | H13-C13-H13C | 109.5 |
| C5-C6-H6 | 120.2 | C8-N1-C9 | 116.8(2) |
| C6-C7-C8 | 119.4(3) | C10-N2-C11 | 108.1(2) |
| C6-C7-H7 | 120.3 | C10-N2-C1 | 127.4(2) |
| C8-C7-H7 | 120.3 | C11-N2-C1 | 124.4(2) |
| N1-C8-C7 | 124.3(3) | C10-N3-C12 | 108.6(2) |
| N1-C8-H8 | 117.9 | C10-N3-C13 | 125.3(3) |
| C7-C8-H8 | 117.9 | C12-N3-C13 | 125.9(3) |
| N1-C9-C5 | 123.2(2) | | |

Table A1.5: Bond lengths for 1-mesityl 3-(-2-methylquinoline)imidazolium chloride **9b**:

| Bond lengths (Å) | | Bond lengths (Å) | | Bond lengths (Å) | |
|------------------|----------|------------------|----------|------------------|----------|
| C1-N2 | 1.323(3) | C7-C8 | 1.389(4) | C19-H19 | 0.9500 |
| C1-N1 | 1.340(3) | C7-C11 | 1.515(4) | C20-C21 | 1.373(4) |
| C1-H1 | 0.9500 | C8-C9 | 1.390(4) | C20-H20 | 0.9500 |
| C2-C3 | 1.356(3) | C8-H8 | 0.9500 | C21-C22 | 1.409(4) |
| C2-N1 | 1.374(3) | C9-C12 | 1.500 | C21-H21 | 0.9500 |
| C2-H2 | 0.9500 | C10-H10A | 0.9800 | C22-N3 | 1.374(3) |
| C3-N2 | 1.370(3) | C13-N2 | 1.463(3) | B1-F2 | 1.333(4) |
| C3-H3 | 0.9500 | C13-C14 | 1.506(3) | B1-F4A | |
| C4-C5 | 1.392(3) | C13-H13A | 0.9900 | 1.360(10) | |
| C4-C9 | 1.402(3) | C13-H13B | 0.9900 | B1-F3 | 1.383(3) |
| C4-N1 | 1.446(3) | C14-N3 | 1.306(3) | B1-F1 | 1.393(4) |
| C5-C6 | 1.377(3) | C14-C15 | 1.406(3) | B1-F4 | 1.402(3) |
| C5-C10 | 1.507(4) | C15-C16 | 1.354(3) | B1-F2A | 1.414(9) |

| Appendix | | | | | |
|----------|----------|---------|----------|--------|----------|
| C6-C7 | 1.382(4) | C15-H15 | 0.9500 | B1-F1A | 1.425(8) |
| C6-H6 | 0.9500 | C16-C17 | 1.410(4) | F1-F4A | 1.06(3) |
| C10-H10B | 0.9800 | C16-H16 | 0.9500 | F- F1A | 1.57(2) |
| C10-H10C | 0.9800 | C17-C22 | 1.414(3) | F1A-F2 | 1.17(2) |
| C11-H11A | 0.9800 | C17-C18 | 1.418(3) | F2-F2A | 1.23(2) |
| C11-H11B | 0.9800 | C18-C19 | 1.350(4) | F2A-F4 | 1.58(3) |
| C11-H11C | 0.9800 | C18-H18 | 0.9500 | F4-F4A | 1.37(3) |
| C12-H12A | 0.9800 | C19-C20 | 1.402(4) | | |
| C12-H12B | 0.9800 | | | | |
| C12-H12C | 0.9800 | | | | |

Table A1.6: Bond angles for 1-mesityl 3-(-2-methylquinoline)imidazolium chloride **9b**:

| Bond angles (°) | | Bond angles (°) | |
|-----------------|----------|-----------------|------------|
| N2-C1-N1 | 108.2(2) | C19-C18-H18 | 119.7 |
| N2-C1-H1 | 126.5 | C17-C18-H18 | 119.7 |
| N1-C1-H1 | 125.9 | C18-C19-C20 | 120.5(2) |
| C-C2-N1 | 107.0(2) | C18-C19-H19 | 119.7 |
| C3-C2-H2 | 126.5 | C20-C19-H19 | 119.7 |
| N1-C2-H2 | 126.5 | C21-C20-C19 | 120.5(3) |
| C2-C3-N2 | 107.0(2) | C21-C20-H20 | 119.8 |
| C2-C3-H3 | 126.5 | C19-C20-H20 | 119.8 |
| N2-C3-H3 | 126.5 | C20-C21-C22 | 120.5(2) |
| C5-C4-C9 | 123.0(2) | C20 C21 H21 | 119.8 |
| C5-C4-N1 | 119.3(2) | C22-C21-H21 | 119.8 |
| C9-C4-N1 | 117.6(2) | N3-C22-C21 | 119.0(2) |
| C6-C5-C4 | 117.5(2) | N3-C22-C17 | 122.4(2) |
| C6-C5-C10 | 120.8(2) | C21-C22-C17 | 118.7(2) |
| C4-C5-C10 | 121.7(2) | C1-N1-C2 | 108.5(2) |
| C5-C6-C7 | 122.5(2) | C1-N1-C4 | 126.2(2) |
| C5-C6-H6 | 118.8 | C2-N1-C4 | 109.3(2) |
| C7-C6-H6 | 118.8 | C1-N2-C13 | 125.3(2) |
| C6-C7-C8 | 118.0(2) | C3-N2-C13 | 125.0(2) |
| C6-C7 -C11 | 121.1(3) | C14-N3-C22 | 117.13(19) |

| Appendix | | | |
|-----------------|-----------|------------|-----------|
| C8-C7-C11 | 120.9(3) | F2-B1-F4A | 120.3(10) |
| C7-C8-C9 | 122.8(3) | F2-B1-F3 | 111.9(3) |
| C7-C8-H8 | 118.6 | F4A-B1-F3 | 126.9(11) |
| C9-C8-H8 | 118.6 | F2-B1-F1 | 111.1(3) |
| C8-C9-C4 | 116.1(2) | F4A-B1-F1 | 45.1(12) |
| C8-C9-C12 | 122.2(2) | F3-B1-F1 | 107.8(3) |
| C4-C9-C12 | 121.6(2) | F2-B1-F4 | 111.4(3) |
| C5-C10H-10A | 109.5 | F4A-B1-F4 | 59.4(12) |
| C5-C10-0H10B | 109.5 | F3-B1-F4 | 110.2(2) |
| H10A-C10-H10B | 109.5 | F1-B1-F4 | 104.1(3) |
| C5-C10-H10C | 109.5 | F2-B1-F2A | 52.9(11) |
| H10A-C10-H10C | 109.5 | F4A-B1-F2A | 118.1(16) |
| H10B-C10-H10C | 109.5 | F3-B1-F2 | 100.0(7) |
| C7-C11-H11A | 109.5 | F1-B1-F2A | 152.0(7) |
| C7-C11-H11B | 109.5 | F4-B1-F2A | 68.5(12) |
| H11A-C11-H11B | 109.5 | F2-B1-F1A | 50.2(10) |
| C7-C11-H11C | 109.5 | F4A-B1-F1A | 102.8(13) |
| H11A-C11-H11C | 109.5 | F3-B1-F1A | 102.8(6) |
| H11B-C11-H11C | 109.5 | F1-B1-F1A | 67.7(10) |
| C9-C12-H12A | 109.5 | F4-B1-F1A | 146.8(7) |
| C9-C12-H12B | 109.5 | F2A-B1-F1A | 103.0(15) |
| H12A-C12-H12B | 109.5 | F4A-F1-B1 | 65.8(7) |
| C9-C12-H12C | 109.5 | F4A-F- F1A | 110.5(12) |
| H12A-C12-H12C | 109.5 | B1-F1-F1A | 57.1(6) |
| H12B-C12-H12C | 109.5 | F2-F1A-B1 | 60.8(5) |
| N2-C13-C14 | 112.9(2) | F2-F1A-F1 | 109.2(8) |
| N2-C13-H13A | 109.0 | B1-F1A-F1 | 55.2(6) |
| C14-C13-H13A | 109.0 | F1A-F2-F2A | 135.6(10) |
| N2-C13-H13B | 109.0 | F1A-F2-B1 | 68.9(7) |
| C14-C13-H13B | 109.0 | F2A-F2-B1 | 66.9(7) |
| H13A-C13-H13B | 107.8 | F2-F2A-B1 | 60.1(6) |
| N3-C1-C15 | 124.4(2) | F2-F2A-F4 | 106.3(8) |
| N3-C14-C13 | 119.7(19) | B1-F2A-F4 | 55.4(7) |
| C15-C14-C13 | 116.5 | F4A-F4-B1 | 58.8(6) |

| Appendix | | | |
|-------------|----------|------------|-----------|
| C16-C15-C14 | 119.2(2) | F4A-F4-F2A | 107.1(10) |
| C16-C15-H15 | 120.4 | B1-F4-F2A | 56.1(7) |
| C14-C15-H15 | 120.4 | F1-F4A-B1 | 69.0(8) |
| C15-C16-C17 | 119.2(2) | F1-F4A-F4 | 130.1(10) |
| C15-C16-H16 | 120.4 | B1-F4A-F4 | 61.8(8) |
| C17-C16-H16 | 120.4 | | |
| C16-C17-C22 | 117.7(2) | | |
| C22-C17-C18 | 119.3(2) | | |
| C19-C18-C17 | 120.6(2) | | |

Table A1.7 : Bond lengths for 1-isopropyl -3-(2-methylquinoline) imidazolium iodide (**9c**)

| Bond lengths (Å) | | Bond lengths (Å) | | Bond lengths (Å) | |
|------------------|-----------|------------------|---------|------------------|---------|
| N1-C1 | 1.30(2) | C4-C5 | 1.53(2) | C20-C21 | 1.52(2) |
| N1-C2 | 1.42(2) | C7-C8 | 1.4888 | C20-C22 | 1.57(2) |
| N1-C4 | 1.50(2) | N3-C12 | 1.3899 | C23-C24 | 1.4359 |
| N2-C3 | 1.34(2) | N3-C8 | 1.3899 | N6-C28 | 1.3899 |
| N2-C1 | 1.369(19) | C8-C9 | 1.3888 | N6-C24 | 1.3899 |
| N2-C7 | 1.515(15) | C9-C10 | 1.3898 | C24-C25 | 1.3888 |
| N4-C18 | 1.31(2) | C10-C11 | 1.3899 | C25-C26 | 1.3898 |
| N4-C17 | 1.37(2) | C11-C12 | 1.3888 | C26-C27 | 1.3899 |
| N4-C20 | 1.475(19) | C11-C16 | 1.3899 | C27-C28 | 1.3888 |
| N5-C17 | 1.30(2) | C12-C13 | 1.3899 | C27-C32 | 1.3899 |
| N5-C19 | 1.41(2) | C13-C14 | 1.3898 | C28-C29 | 1.3899 |
| N5-C23 | 1.421(16) | C14-C15 | 1.3888 | C29-C30 | 1.3898 |
| C2-C3 | 1.35(3) | C15-C16 | 1.3899 | C30-C31 | 1.3888 |
| C4-C6 | 1.46(2) | C18-C19 | 1.34(3) | C31-C32 | 1.3888 |

Appendix

Table A1.8 : Bond angles for 1-isopropyl -3-(2-methylquinoline) imidazolium iodide (**9c**)

| Bond angles (°) | | Bond angles (°) | |
|-----------------|-----------|-----------------|-----------|
| C1-N1-C2 | 111.4(11) | C11-C12-C13 | 120.0 |
| C1-N1-C4 | 124.4(11) | N3-C12-C13 | 120.0 |
| C2-N1-C4 | 124.2(12) | C14-C13-C12 | 120.0 |
| C3-N2-C1 | 109.8(15) | C15-C14-C13 | 120.0 |
| C3-N2-C7 | 120.7(13) | C14-C15-C16 | 120.0 |
| C1-N2-C7 | 129.5(13) | C11-C16-C15 | 120.1 |
| C18-N4-C17 | 104.7(15) | N5-C17-N4 | 111.2(14) |
| C18-N4-C20 | 128.7(15) | N4-C18-C19 | 112.5(16) |
| C17-N4-C20 | 126.5(14) | C18-C19-N5 | 104.8(15) |
| C17-N5-C19 | 106.8(15) | N4-C20-C21 | 109.9(13) |
| C17-N5-C23 | 124.5(11) | N4-C20-C22 | 107.6(14) |
| C19-N5-C23 | 128.5(13) | C21-C20-C22 | 114.3(15) |
| N1-C1-N2 | 105.9(12) | N5-C23-C24 | 120.1(5) |
| C3-C2-N1 | 103.6(13) | C28-N6-C24 | 120.1 |
| N2-C3-C2 | 109.3(16) | C25-C24-N6 | 120.0 |
| C6-C4-N1 | 110.5(14) | C25-C24-C23 | 125.0 |
| C6-C4-C5 | 109.7(14) | N6-C24-C23 | 114.8 |
| N1-C4-C5 | 109.7(13) | C24-C25-C26 | 120.0 |
| C8-C7-N2 | 111.2(5) | C25-C26-C27 | 120.0 |
| C12-N3-C8 | 120.1 | C28-C27-C32 | 120.0 |
| C9-C8-N3 | 120.0 | C28-C27-C26 | 120.0 |
| C9-C8-C7 | 120.7 | C32-C27-C26 | 120.1 |
| N3-C8-C7 | 119.2 | C27-C28-N6 | 120.0 |
| C8-C9-C10 | 120.0 | C27-C28-C29 | 120.0 |
| C9-C10-C11 | 120.1 | N6-C28-C29 | 120.0 |
| C12-C11-C16 | 120.0 | C30-C29-C28 | 120.0 |
| C12-C11-C10 | 120.0 | C31-C30-C29 | 120.0 |
| C16-C11-C10 | 120.1 | C30-C31-C32 | 120.0 |
| C11-C12-N3 | 120.0 | C27-C32-C31 | 120.1 |

Table A1.9: Bond lengths for Bis-1, 3- (2-methylquinolin) imidazolium chloride (9f)

| Bond lengths (Å) | | Bond lengths (Å) | | Bond lengths (Å) | |
|------------------|----------|------------------|----------|------------------|----------|
| N2-C3 | 1.474(3) | C10-C17 | 1.415(3) | C19-C24 | 1.40(3) |
| N2-C6 | 1.335(3) | C11-C12 | 1.413(3) | N20-C21 | 1.410(4) |
| N2-C18 | 1.467(3) | C11-C14 | 1.420(4) | C21-C28 | 1.415(3) |
| C3-C4 | 1.350(3) | C12-C13 | 1.362(4) | C22-C23 | 1.414(3) |
| C3-H31 | 0.943 | C12-H121 | 0.942 | C22-C25 | 1.416(4) |
| C4-N5 | 1.383(3) | C13-H131 | 0.940 | C23-C24 | 1.360(3) |
| C4-H41 | 0.959 | C14-C15 | 1.370(3) | C23-H231 | 0.938 |
| N5-C6 | 1.327(3) | C14-H141 | 0.933 | C24-H241 | 0.962 |
| N5-C7 | 1.461(3) | C15-C16 | 1.400(3) | C25-C26 | 1.366(4) |
| C6-H61 | 0.940 | C15-H151 | 0.956 | C25-H251 | 0.963 |
| C7-C8 | 1.511(3) | C16-C17 | 1.360(4) | C26-C27 | 1.394(4) |
| C7-H72 | 0.991 | C16H161 | 0.966 | C26-H261 | 0.916 |
| C7-H71 | 0.962 | C17-H171 | 0.938 | C27-C28 | 1.365(4) |
| C8-N9 | 1.316(3) | C18-C19 | 1.514(3) | C27-H271 | 0.945 |
| C8-C13 | 1.407(3) | C18-H181 | 0.975 | C28-H281 | 0.971 |
| N9-C10 | 1.371(3) | C19-N20 | 1.323(3) | | |
| C10-C11 | 1.41(3) | | | | |

Appendix

Table A1.10: Bond angles for Bis-1, 3- (2-methylquinolin) imidazolium chloride (**9f**)

| Bond angles (°) | | Bond angles (°) | | Bond angles (°) | |
|-----------------|------------|-----------------|----------|-----------------|----------|
| C3-N2-C6 | 108.7(2) | C13-C12-H121 | 118.9 | C19-C18-H181 | 108.1 |
| C3-N2-C18 | 126.6(2) | C8-C13-C12 | 119.2(2) | N2-C18-H182 | 108.3 |
| C6-N2-C18 | 124.7(2) | C8-C13-H131 | 119.5 | C19-C18-H182 | 107.7 |
| N2-C3-C4 | 107.4(2) | C12-C13-H131 | 121.3 | H181-C18 H182 | 111.4 |
| N2-C3-H31 | 124.9 | C11-C14-C15 | 120.1(2) | C18-C19-N20 | 113.7(2) |
| C4-C3-H31 | 127.7 | C11-C14-H141 | 118.3 | C18-C19-C24 | 122.4(2) |
| C3-C4-N5 | 106.6(2) | C15-C14-H141 | 121.6 | N20-C19-C24 | 123.9(2) |
| C3-C4-H41 | 127.3 | C14-C15-C16 | 120.4(2) | C19-N20-C21 | 117.2(2) |
| N5-C4-H41 | 126.0 | C11-C10-C17 | 118.7(2) | N20-C21-C22 | 123.1(2) |
| C4-N5-C6 | 109.0(2) | C10-C11-C12 | 117.7(2) | N20-C21-C28 | 118.4(3) |
| C4-N5-C7 | 125.6(2) | C10-C11-C14 | 119.2(2) | C22-C21-C28 | 118.5(2) |
| C6-N5-C7 | 125.0(2) | C12-C11-C14 | 123.1(2) | C21-C22-C23 | 117.2(2) |
| N2-C6-N5 | 108.3(2) | C11-C12-C13 | 119.2(2) | C21-C22-C25 | 119.7(2) |
| N2-C6-H61 | 124.7 | C11-C12-H121 | 121.9 | C23-C22-C25 | 123.2(3) |
| N5-C6-H61 | 127.0 | C13-C12-H121 | 118.9 | C22-C23-C24 | 119.8(3) |
| N5-C7-C8 | 112.07(19) | C8-C13-C12 | 119.2(2) | C22-C23-H231 | 118.8 |
| N5-C7-H72 | 109.8 | C8-C13-H131 | 119.5 | C24-C23-H231 | 121.3 |
| C8-C7-H72 | 109.0 | C12-C13-H131 | 121.3 | C19-C24-C23 | 118.8(2) |
| N5-C7-H71 | 107.4 | C11-C14-C15 | 120.1(2) | C19-C24-H241 | 121.5 |
| C8-C7-H71 | 108.5 | C11-C14-H141 | 118.3 | C23-C24-H241 | 119.7 |
| H72-C7-H71 | 110.0 | C15-C14-H141 | 121.6 | C22-C25-C26 | 120.3(3) |
| C7-C8-N9 | 117.6(2) | C14-C15-C16 | 120.4(2) | C22-C25-H251 | 117.4 |
| C7-C8-C13 | 118.4(2) | C14-C15-H151 | 117.9 | C26-C25-H251 | 122.2 |
| N9-C8-C13 | 124.0(2) | C16-C15-H151 | 121. | C25-C26-C27 | 119.8(3) |
| C8-N9-C10 | 117.5(2) | C15-C16-C17 | 120.7(2) | C25-C26-H261 | 119.3 |
| N9-C10-C11 | 122.5(2) | C15-C16-H161 | 119.2 | C27-C26-H261 | 120.9 |
| N9-C10-C17 | 118.8(2) | C17-C16-H161 | 120.2 | C26-C27-C28 | 121.5(3) |
| C11-C10-C17 | 118.7(2) | C10-C17-C16 | 120.8(2) | C26-C27-H271 | 118.2 |
| C10-C11-C12 | 117.7(2) | C10C17H171 | 120.3 | C28-C27-H271 | 120.3 |
| C10-C11-C14 | 119.2(2) | C16-C17-H171 | 118.9 | C21-C28-C27 | 120.2(3) |

Appendix

| | | | | | |
|--------------|----------|-------------|----------|--------------|-------|
| C12-C11-C14 | 123.1(2) | N2-C18-C19 | 113.0(2) | C21-C28-H281 | 120.3 |
| C11-C12-C13 | 119.2(2) | N2-C18-H181 | 108.4 | C27-C28-H281 | 119.5 |
| C11-C12-H121 | 121.9 | | | | |

Table A1.11: Bond lengths and angles for [Ag (1-methy-3-(-2-methylquinoline)imidazolin-2-ylidene) ₂][AgCl₂] (**3.36**)

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|-----------|-----------------|-----------|-----------------|-----------|
| C1-N2 | 1.351(14) | N2-C1-N1 | 104.0(10) | C12-C13-C14 | 120.0 |
| C1-N1 | 1.353(16) | N2-C1-Ag1 | 130.6(8) | C9-C14-C13 | 120.0 |
| C1-Ag1 | 2.106(12) | N1-C1-Ag | 125.2(9) | C1-Ag1-C1 | 170.8(8) |
| C2-C3 | 1.32(2) | C3-C2-N1 | 107.8(11) | C1-Ag1-Ag2 | 94.6(4) |
| C2-N1 | 1.378(16) | C2-C3-N2 | 105.8(12) | C1-Ag1-Ag2 | 94.6(4) |
| C3-N2 | 1.392(19) | N2-C5-C6 | 112.7(13) | C1-Ag1-Ag2 | 85.4(4) |
| C4-N1 | 1.480(19) | C1-N1-C2 | 110.1(11) | C1-Ag1-Ag2 | 85.4(4) |
| C5-N2 | 1.446(18) | C1-N1-C4 | 125.4(12) | Ag2-Ag1-Ag2 | 180.0(3) |
| C5-C6 | 1.48(2) | C2-N1-C4 | 122.4(12) | Cl1-Ag2-Cl1 | 176.6(11) |
| N3-C6 | 1.3900 | C1-N2-C3 | 109.9(9) | Cl1-Ag2-Ag1 | 88.3(5) |
| N3-C10 | 1.3900 | C1-N2-C5 | 119.3(10) | Cl1-Ag2-Ag1 | 88.3(5) |
| C6-C7 | 1.3900 | C3-N2-C5 | 130.2(10) | Cl1-Ag2-Ag1 | 91.7(5) |
| C7-C8 | 1.3900 | C6-N3-C10 | 120.0 | Cl1-Ag2-Ag1 | 91.7(5) |
| C8-C9 | 1.3900 | N3-C6-C7 | 120.0 | Ag1-Ag2-Ag1 | 180.0 |
| C9-C14 | 1.3900 | N3-C6-C5 | 116.5(11) | | |
| C9-C10 | 1.3900 | C7-C6-C5 | 123.5(10) | | |
| C10-C11 | 1.3900 | C6-C7-C8 | 120.0 | | |
| C11-C12 | 1.3900 | C9-C8-C7 | 120.0 | | |
| C12-C13 | 1.3900 | C14-C9-C10 | 120.0 | | |
| C13-C14 | 1.3900 | C14-C9-C8 | 120.0 | | |
| Cl1-Ag2 | 2.290(5) | C10-C9-C8 | 120.0 | | |
| Ag1-C1 | 2.106(12) | C9-C10-C11 | 120.0 | | |
| Ag1-Ag2 | 3.201(4) | C9-C10-N3 | 120.0 | | |
| Ag1-Ag2 | 3.300(4) | C11-C10-N3 | 120.0 | | |
| Ag2-Cl1 | 2.290(5) | C12-C11-C10 | 120.0 | | |

Table A1.12: Bond lengths and angles for [Ag (1-mesityl-3-(2-methylquinoline) imidazolin-2-ylidene) Cl] 3.37

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|------------|-----------------|------------|-----------------|----------|
| Ag1-Ag1 | 3.8010(5) | Ag1-Ag1-Cl2 | 37.997(17) | C4-C16-H161 | 128.2 |
| Ag1-Cl2 | 3.2126(10) | Ag1-Ag1-Cl2 | 57.31(2) | C8-C17-C9 | 119.1(3) |
| Ag1-Cl2 | 2.3501(9) | Cl2-Ag1-Cl2 | 95.30(3) | C8-C17-N5 | 118.0(3) |
| Ag1-C13 | 2.090(3) | Ag1-Ag1-C13 | 133.23(10) | C9-C17-N5 | 123.0(3) |
| N3-C4 | 1.384(4) | Cl2-Ag1-C13 | 95.29(10) | C15-C18-C10 | 120.5(4) |
| N3-C6 | 1.474(4) | Cl2-Ag1-C13 | 169.03(10) | C15-C18-C22 | 122.0(4) |
| N3-C13 | 1.342(5) | Ag1-Cl2-Ag1 | 84.70(3) | C10-C18-C22 | 117.5(4) |
| C4-C16 | 1.344(5) | C4-N3-C6 | 123.9(3) | C21-C19-H193 | 112.7 |
| C4-H41 | 0.950 | C4-N3-C13 | 111.1(3) | C21-C19-H191 | 108.8 |
| N5-C14 | 1.320(5) | C6-N3-C13 | 125.0(3) | H193-C19-H191 | 107.4 |
| N5-C17 | 1.373(5) | N3-C4-C16 | 106.9(3) | C21-C19-H192 | 110.3 |
| C6-C14 | 1.516(5) | N3-C4-H41 | 124.8 | H193-C19-H192 | 108.2 |
| C6-H62 | 0.974 | C16-C4-H41 | 128.2 | H191-C19-H192 | 109.4 |
| C6-H61 | 0.967 | C14-N5-C17 | 117.2(3) | C9-C20-C25 | 120.0(4) |
| C7-C21 | 1.396(5) | N3-C6-C14 | 111.6(3) | C9-C20-H201 | 119.4 |
| C7-C23 | 1.390(5) | N3-C6-H62 | 108.1 | C25-C20-H201 | 120.6 |
| C7-H71 | 0.957 | C14-C6-H62 | 107.4 | C19-C21-C7 | 120.6(4) |
| C8-C17 | 1.422(5) | N3-C6-H61 | 109.2 | C19-C21-C10 | 120.3(4) |
| C8-C27 | 1.366(6) | C14-C6-H61 | 109.7 | C7-C21-C10 | 119.1(3) |
| C8-H81 | 0.932 | H62-C6-H61 | 110.9 | N11-C22-C18 | 118.9(3) |
| C9-C12 | 1.421(6) | C21-C7-C23 | 121.6(3) | N11-C22-C23 | 118.0(3) |
| C9-C17 | 1.410(6) | C21-C7-H71 | 119.2 | C18-C22-C23 | 123.1(3) |
| C9-C20 | 1.417(5) | C23-C7-H71 | 119.2 | C22-C23-C7 | 117.3(3) |
| C10-C18 | 1.399(5) | C17-C8-C27 | 120.1(4) | C22-C23-C26 | 121.5(3) |
| C10-C21 | 1.384(6) | C17-C8-H81 | 120.2 | C7-C23-C26 | 121.2(3) |
| C10-H101 | 0.948 | C27-C8-H81 | 119.8 | C14-C24-C12 | 119.2(4) |
| N11-C13 | 1.353(4) | C12-C9-C17 | 117.6(3) | C14-C24-H241 | 120.4 |

Appendix

| | | | | | |
|----------|----------|---------------|----------|---------------|----------|
| N11-C16 | 1.389(5) | C12-C9-C20 | 122.8(4) | C12-C24-H241 | 120.3 |
| N11-C22 | 1.445(4) | C17-C9-C20 | 119.6(4) | C20-C25-C27 | 120.7(4) |
| C12-C24 | 1.371(6) | C18-C10-C21 | 121.3(4) | C20-C25-H251 | 119.6 |
| C12-H121 | 0.934 | C18-C10-H101 | 119.0 | C27-C25-H251 | 119.7 |
| C14-C24 | 1.407(5) | C21-C10-H101 | 119.7 | C23-C26-H262 | 109.5 |
| C15-C18 | 1.510(6) | C13-N11-C16 | 110.9(3) | C23-C26-H263 | 111.4 |
| C15-H152 | 0.945 | C13-N11-C22 | 122.6(3) | H262-C26-H263 | 108.2 |
| C15-H153 | 0.960 | C16-N11-C22 | 126.5(3) | C23-C26-H261 | 111.2 |
| C15-H151 | 0.961 | C9-C12-C24 | 118.9(4) | H262-C26-H261 | 106.3 |
| C16-H161 | 0.945 | C9-C12-H121 | 119.3 | H263-C26-H261 | 110.1 |
| C18-C22 | 1.389(5) | C24-C12-H121 | 121.8 | C25-C27-C8 | 120.6(4) |
| C19-C21 | 1.509(5) | N11-C13-N3 | 104.9(3) | C25-C27-H271 | 119.4 |
| C19-H193 | 0.950 | N11-C13-Ag1 | 123.7(3) | C8-C27-H271 | 120.0 |
| C19-H191 | 0.942 | N3-C13-Ag1 | 131.4(2) | | |
| C19-H192 | 0.945 | C6-C14-N5 | 116.2(3) | | |
| C20-C25 | 1.366(7) | C6-C14-C24 | 119.7(3) | | |
| C20-H201 | 0.931 | N5-C14-C24 | 124.1(3) | | |
| C22-C23 | 1.391(5) | C18-C15-H152 | 109.5 | | |
| C23-C26 | 1.498(5) | C18-C15-H153 | 111.6 | | |
| C24-H241 | 0.936 | H152-C15-H153 | 105.3 | | |
| C25-C27 | 1.409(7) | C18-C15-H151 | 113.5 | | |
| C25-H251 | 0.946 | H152-C15-H151 | 107.4 | | |
| C26-H262 | 0.954 | H153-C15-H151 | 109.1 | | |
| C26-H263 | 0.962 | N11-C16-C4 | 106.2(3) | | |
| C26-H261 | 0.954 | N11-C16-H161 | 125.6 | | |
| C27-H271 | 0.930 | | | | |

Appendix

Table 1.13: Bond lengths and angles for 1-methyl-3-(2-methylquinoline)imidazolin-2-ylidene][1,5-cyclooctadiene]rhodium(I) Chloride **4.20**

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|----------|-----------------|----------|-----------------|----------|
| C1-N1 | 1.356(6) | N1-C1-N2 | 103.8(4) | Rh1-C16-H16 | 86.6 |
| C1-N2 | 1.362(7) | N1-C1-Rh1 | 131.4(4) | C18-C17-C16 | 112.8(4) |
| C1-Rh1 | 2.028(5) | N2-C1-Rh1 | 124.9(4) | C18 C17 H17A | 109.0 |
| C2-C3 | 1.345(8) | C3-C2-N1 | 107.1(5) | C16 C17 H17A | 109.0 |
| C2-N1 | 1.371(7) | C3-C2-H2 | 126.5 | C18 C17 H17B | 109.0 |
| C2-H2 | 0.9500 | N1-C2-H2 | 126.5 | C16-C17-H17B | 109.0 |
| C3-N2 | 1.394(7) | C2-C3-N2 | 106.3(5) | H17A-C17-H17B | 107.8 |
| C3-H3 | 0.9500 | C2-C3-H3 | 126.9 | C19-C18-C17 | 113.3(4) |
| C4-N1 | 1.466(7) | N2-C3-H3 | 126.9 | C19-C18-H18A | 108.9 |
| C4-H4A | 0.9800 | N1-C4-H4A | 109.5 | C17-C18-H18A | 108.9 |
| C4-H4B | 0.9800 | N1-C4-H4B | 109.5 | C19-C18-H18B | 108.9 |
| C4-H4C | 0.9800 | H4A-C4-H4B | 109.5 | C17-C18-H18B | 108.9 |
| C5-N2 | 1.461(6) | N1-C4-H4C | 109.5 | H18A-C18-H18B | 107.7 |
| C5-C6 | 1.508(7) | H4A-C4-H4C | 109.5 | C20-C19-C18 | 125.8(5) |
| C5-H5A | 0.9900 | H4B-C4-H4C | 109.5 | C20-C19-Rh1 | 73.0(3) |
| C5-H5B | 0.9900 | N2-C5-C6 | 113.9(4) | C18-C19-Rh1 | 106.6(4) |
| C6-N3 | 1.320(7) | N2-C5-H5A | 108.8 | C20-C19-H19 | 117.1 |
| C6-C7 | 1.420(7) | C6-C5-H5A | 108.8 | C18-C19-H19 | 117.1 |
| C7-C8 | 1.378(8) | N2-C5-H5B | 108.8 | Rh1-C19-H19 | 90.5 |
| C7-H7 | 0.9500 | C6-C5-H5B | 108.8 | C19-C20-C21 | 125.3(5) |
| C8-C9 | 1.412(7) | H5A-C5-H5B | 107.7 | C19-C20-Rh1 | 70.9(3) |
| C8-H8 | 0.9500 | N3-C6-C7 | 123.5(5) | C21-C20-Rh1 | 111.6(4) |
| C9-C10 | 1.408(7) | N3-C6-C5 | 114.7(5) | C19-C20-H20 | 117.3 |
| C9-C14 | 1.417(7) | C7-C6-C5 | 121.8(5) | C21-C20-H20 | 117.3 |

Appendix

| | | | | | |
|----------|----------|-------------|----------|---------------|----------|
| C10-C11 | 1.372(8) | C8-C7-C6 | 118.8(5) | Rh1-C20-H20 | 87.4 |
| C10-H10 | 0.9500 | C8-C7-H7 | 120.6 | C20-C21-C22 | 112.5(5) |
| C11-C12 | 1.405(8) | C6-C7-H7 | 120.6 | C20-C21-H21A | 109.1 |
| C11-H11 | 0.9500 | C7-C8-C9 | 119.8(5) | C22-C21-H21A | 109.1 |
| C12-C13 | 1.377(8) | C7-C8-H8 | 120.1 | C20-C21-H21B | 109.1 |
| C12-H12 | 0.9500 | C9-C8-H8 | 120.1 | C22-C21-H21B | 109.1 |
| C13-C14 | 1.416(7) | C10-C9-C8 | 123.5(5) | H21A-C21-H21B | 107.8 |
| C13-H13 | 0.9500 | C10-C9-C14 | 119.1(5) | C15-C22-C21 | 113.5(4) |
| C14-N3 | 1.378(7) | C8-C9-C14 | 117.4(5) | C15-C22-H22A | 108.9 |
| C15-C16 | 1.410(7) | C11-C10-C9 | 120.8(5) | C21-C22-H22A | 108.9 |
| C15-C22 | 1.499(8) | C11-C10-H10 | 119.6 | C15-C22-H22B | 108.9 |
| C15-Rh1 | 2.123(5) | C9-C10-H10 | 119.6 | C21-C22-H22B | 108.9 |
| C15-H15 | 0.9500 | C10-C11-C12 | 120.1(5) | H22A-C22-H22B | 107.7 |
| C16-C17 | 1.526(7) | C10-C11-H11 | 120.0 | C1-N1-C2 | 111.9(4) |
| C16-Rh1 | 2.126(5) | C12-C11-H11 | 120.0 | C1-N1-C4 | 124.5(5) |
| C16-H16 | 0.9500 | C13-C12-C11 | 120.8(5) | C2-N1-C4 | 123.5(4) |
| C17-C18 | .526(8) | C13-C12-H12 | 119.6 | C1-N2-C3 | 111.0(4) |
| C17-H17A | 0.9900 | C11-C12-H12 | 119.6 | C1-N2-C5 | 124.1(4) |
| C17-H17B | 0.9900 | C12-C13-C14 | 119.8(5) | C3-N2-C5 | 124.7(4) |
| C18-C19 | 1.524(8) | C12-C13-H13 | 120.1 | C6-N3-C14 | 117.9(5) |
| C18-H18A | 0.9900 | C14-C13-H13 | 120.1 | C1-Rh1-C15 | 92.0(2) |
| C18-H18B | 0.9900 | N3-C14-C13 | 117.8(5) | C1-Rh1-C16 | 93.2(2) |
| C19-C20 | 1.375(8) | N3-C14-C9 | 122.8(5) | C15-Rh1-C16 | 38.8(2) |
| C19-Rh1 | 2.206(5) | C13-C14-C9 | 119.4(5) | C1-Rh1-C19 | 158.3(2) |
| C19-H19 | 0.9500 | C16-C15-C22 | 125.7(5) | C15-Rh1-C19 | 97.2(2) |
| C20-C21 | 1.506(8) | C16-C15-Rh1 | 70.7(3) | C16-Rh1-C19 | 82.0(2) |
| C20-Rh1 | 2.233(6) | C22-C15-Rh1 | 111.4(4) | C1-Rh1-C20 | 165.5(2) |

Appendix

| | | | | | |
|----------|------------|-------------|----------|-------------|------------|
| C20-H20 | 0.9500 | C16-C15-H15 | 117.2 | C15-Rh1-C20 | 80.7(2) |
| C21-C22 | 1.544(8) | C22-C15-H15 | 117.2 | C16-Rh1-C20 | 88.8(2) |
| C21-H21A | 0.9900 | Rh1-C15-H15 | 87.8 | C19-Rh1-C20 | 36.1(2) |
| C21-H21B | 0.9900 | C15-C16-C17 | 123.8(5) | C1-Rh1-C11 | 87.99(14) |
| C22-H22A | 0.9900 | C15-C16-Rh1 | 70.5(3) | C15-Rh1-C11 | 161.31(15) |
| C22-H22B | 0.9900 | C17-C16-Rh1 | 113.0(4) | C16-Rh1-C11 | 159.89(15) |
| Rh1-C11 | 2.3862(13) | C15-C16-H16 | 118.1 | C19-Rh1-C11 | 89.44(15) |
| | | C17-C16-H16 | 118.1 | C20-Rh1-C11 | 95.03(15) |

Table A1.14: Bond lengths and angles for [1-mesityl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride **4.21**

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|----------|-----------------|----------|-----------------|----------|
| C1-N1 | 1.355(5) | N1-C1-N2 | 103.7(3) | C22-C21-H21 | 119.8 |
| C1-N2 | 1.355(5) | N1-C1-Rh1 | 130.8(3) | N3-C22-C17 | 122.7(4) |
| C1-Rh1 | 2.043(4) | N2-C1-Rh1 | 125.3(3) | N3-C22C21 | 118.2(3) |
| C2-C3 | 1.336(6) | C3-C2- N1 | 106.4(4) | C17-C22-C21 | 119.1(4) |
| C2-N1 | 1.397(5) | C3-C2-H2 | 126.8 | C24-C23-C30 | 123.6(5) |
| C2-H2 | 0.9500 | N1-C2-H2 | 126.8 | C24-C23-Rh1 | 70.9(3) |
| C3-N2 | 1.379(5) | C2-C3-N2 | 106.8(4) | C30-C23-Rh1 | 111.2(3) |
| C3-H3 | 0.9500 | C2-C3-H3 | 126.6 | C24-C23-H23 | 118.2 |
| C4-C9 | 1.394(6) | N2-C3-H3 | 126.6 | C30-C23-H23 | 118.2 |
| C4-C5 | 1.396(6) | C9-C4-C5 | 122.8(3) | Rh1-C23-H23 | 88.0 |
| C4-N1 | 1.447(5) | C9-C4-N1 | 117.3(3) | C23-C24-C25 | 125.3(5) |
| C5-C6 | 1.406(6) | C5-C4-N1 | 119.8(4) | C23-C24-Rh1 | 72.7(3) |
| C5-C10 | 1.497(6) | C4-C5-C6 | 116.8(4) | C25-C24-Rh1 | 109.6(3) |

Appendix

| | | | | | |
|----------|----------|---------------|----------|---------------|----------|
| C6-C7 | 1.385(7) | C4-C5-C10 | 121.3(4) | C23-C24-H24 | 117.3 |
| C6-H6 | 0.9500 | C6-C5-C10 | 121.9(4) | C25-C24-H24 | 117.3 |
| C7-C8 | 1.378(6) | C7-C6-C5 | 122.1(4) | Rh1-C24-H24 | 87.6 |
| C7-C11 | 1.512(6) | C7-C6-H6 | 118.9 | C26-C25-C24 | 115.7(5) |
| C8-C9 | 1.410(6) | C5-C6 H6 | 118.9 | C26-C25-H25A | 108.4 |
| C8-H8 | 0.9500 | C8-C7-C6 | 119.1(4) | C24-C25-H25A | 108.4 |
| C9-C12 | 1.505(5) | C8-C7-C11 | 119.7(4) | C26-C25-H25B | 108.4 |
| C10-H10A | 0.9800 | C6-C7-C11 | 121.2(4) | C24-C25-H25B | 108.4 |
| C10-H10B | 0.9800 | C7-C8-C9 | 121.6(4) | H25A-C25-H25B | 107.4 |
| C10-H10C | 0.9800 | C7-C8-H8 | 119.2 | C25-C26-C27 | 115.3(5) |
| C11-H11A | 0.9800 | C9-C8-H8 | 119.2 | C25-C26-H26A | 108.5 |
| C11-H11B | 0.9800 | C4-C9-C8 | 117.4(4) | C27-C26-H26A | 108.5 |
| C11-H11C | 0.9800 | C4-C9-C12 | 121.8(3) | C25-C26-H26B | 108.5 |
| C12-H12A | 0.9800 | C8-C9-C12 | 120.8(4) | C27-C26-H26B | 108.5 |
| C12-H12B | 0.9800 | C5-C10-H10A | 109.5 | H26A-C26-H26B | 107.5 |
| C12-H12C | 0.9800 | C5-C10-H10B | 109.5 | C28-C27-C26 | 122.7(5) |
| C13-N2 | 1.451(5) | H10A-C10-H10B | 109.5 | C28-C27-Rh1 | 70.4(3) |
| C13-C14 | 1.521(6) | C5-C10-H10C | 109.5 | C26-C27-Rh1 | 112.9(3) |
| C13-H13A | 0.9900 | H10A-C10-H10C | 109.5 | C28-C27-H27 | 118.7 |
| C13-H13B | 0.9900 | H10B-C10-H10C | 109.5 | C26-C27-H27 | 118.7 |
| C14-N3 | 1.320(5) | C7-C11-H11A | 109.5 | Rh1-C27-H27 | 86.7 |
| C14-C15 | 1.419(6) | C7-C11-H11B | 109.5 | C27-C28-C29 | 126.8(5) |
| C15-C16 | 1.360(6) | H11A-C11-H11B | 109.5 | C27-C28-Rh1 | 71.3(3) |
| C15-H15 | 0.9500 | C7-C11-H11C | 109.5 | C29-C28-Rh1 | 110.8(3) |
| C16-C17 | 1.403(6) | H11A-C11-H11C | 109.5 | C27-C28-H28 | 116.6 |
| C16-H16 | 0.9500 | H11B-C11-H11C | 109.5 | C29-C28-H28 | 116.6 |
| C17-C22 | 1.415(5) | C9-C12H-12A | 109.5 | Rh1-C28-H28 | 87.9 |

Appendix

| | | | | | |
|----------|----------|---------------|----------|---------------|------------|
| C1-C18 | .417(6) | C9-C12-H12B | 109.5 | C30-C29-C28 | 116.9(6) |
| C18-C19 | 1.352(7) | H12A-C12-H12B | 109.5 | C30-C29-H29A | 108.1 |
| C18-H18 | 0.9500 | C9-C12-H12C | 109.5 | C28-C29-H29A | 108.1 |
| C19-C20 | 1.408(6) | H12A-C12-H12C | 109.5 | C30-C29-H29B | 108.1 |
| C19-H19 | 0.9500 | H12B-C12-H12C | 109.5 | C28-C29-H29B | 108.1 |
| C20-C21 | .366(6) | N2-C13-C14 | 112.6(3) | H29A-C29-H29B | 107.3 |
| C20-H20 | 0.9500 | N2-C13-H13A | 109.1 | C29-C30-C23 | 113.8(5) |
| C21-C22 | 1.421(6) | C14-C13-H13A | 109.1 | C29-C30-H30A | 108.8 |
| C21-H21 | 0.9500 | N2-C13-H13B | 109.1 | C23-C30-H30A | 108.8 |
| C22-N3 | 1.365(5) | C14-C13-H13B | 109.1 | C29-C30-H30B | 108.8 |
| C23-C24 | 1.369(6) | H13A-C13-H13B | 107.8 | C23-C30-H30B | 108.8 |
| C23-C30 | 1.514(7) | N3-C14-C15 | 122.9(4) | H30A-C30-H30B | 107.7 |
| C23-Rh1 | 2.202(4) | N3-C14-C13 | 118.6(3) | C1-N1-C2 | 111.1(3) |
| C23-H23 | 0.9500 | C15-C14-C13 | 118.5(4) | C1-N1-C4 | 124.3(3) |
| C24-C25 | 1.509(7) | C16-C15-C14 | 118.8(4) | C2-N1-C4 | 123.7(3) |
| C24-Rh1 | 2.179(4) | C16-C15-H15 | 120.6 | C1-N2-C3 | 111.9(3) |
| C24-H24 | 0.9500 | C14-C15-H15 | 120.6 | C1-N2-C13 | 124.0(3) |
| C25-C26 | 1.480(7) | C15-C1-C17 | 120.2(4) | C3-N2-C13 | 123.9(3) |
| C25-H25A | 0.9900 | C15-C16-H16 | 119.9 | C14-N3-C22 | 118.1(3) |
| C25-H25B | 0.9900 | C17-C16-H16 | 119.9 | C1-Rh1-C28 | 94.63(17) |
| C26-C27 | 1.515(7) | C16-C17-C22 | 117.2(4) | C1-Rh1-C27 | 93.45(15) |
| C26-H26A | 0.9900 | C16-C17-C18 | 124.5(4) | C28-Rh1-C27 | 38.31(15) |
| C26-H26B | 0.9900 | C22-C17-C18 | 118.3(4) | C1-Rh1-C24 | 157.59(17) |
| C27-C28 | 1.389(6) | C19-C18-C17 | 121.6(4) | C28-Rh1-C24 | 94.88(19) |
| C27-Rh1 | 2.123(4) | C19-C18-H18 | 119.2 | C27-Rh1-C24 | 81.68(17) |
| C27-H27 | 0.9500 | C17-C18-H18 | 119.2 | C1-Rh1-C23 | 165.92(17) |
| C28-C29 | 1.499(7) | C18-C19-C20 | 120.1(4) | C28-Rh1-C23 | 81.26(16) |

Appendix

| | | | | | |
|----------|-----------|-------------|----------|-------------|------------|
| C28-Rh1 | 2.112(4) | C18-C19-H19 | 120.0 | C27-Rh1-C23 | 91.58(18) |
| C28-H28 | 0.9500 | C20-C19-H19 | 120.0 | C24-Rh1-C23 | 36.42(16) |
| C29-C30 | 1.476(7) | C21-C20-C19 | 120.5(4) | C1-Rh1-Cl1 | 87.67(11) |
| C29-H29A | 0.9900 | C21-C20-H20 | 119.8 | C28-Rh1-Cl1 | 159.29(13) |
| C29-H29B | 0.9900 | C19-C20-H20 | 119.8 | C27-Rh1-Cl1 | 162.22(13) |
| C30-H30A | .9900 | C20-C21-C22 | 120.3(4) | C24-Rh1-Cl1 | 90.55(14) |
| C30-H30B | 0.9900 | C20-C21-H21 | 119.8 | C23-Rh1-Cl1 | 91.55(13) |
| Rh1-Cl1 | 2.3865(9) | | | | |

Table A1.15: Bond lengths and angles for [1-isopropyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride **4.22**

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|----------|-----------------|----------|-----------------|----------|
| C1-N1 | 1.343(6) | N1-C1-N2 | 104.7(4) | C20-C19-Rh1 | 112.9(4) |
| C1-N2 | 1.345(6) | N1-C1-Rh1 | 129.8(3) | C18-C19-H19 | 117.8 |
| C1-Rh1 | 2.037(5) | N2-C1-Rh1 | 125.1(4) | C20-C19-H19 | 117.8 |
| C2-C3 | 1.351(8) | C3-C2-N1 | 105.8(5) | Rh1-C19-H19 | 87.1 |
| C2-N1 | 1.389(7) | C3-C2-H2 | 127.1 | C21-C20-C19 | 114.5(4) |
| C2-H2 | 0.9500 | N1-C2-H2 | 127.1 | C21-C20-H20A | 108.6 |
| C3-N2 | 1.369(7) | C2-C3-N2 | 106.9(5) | C19-C20-H20A | 108.6 |
| C3-H3 | 0.9500 | C2-C3-H3 | 126.5 | C21-C20-H20B | 108.6 |
| C4-N1 | 1.477(6) | N2-C3-H3 | 126.5 | C19-C20-H20B | 108.6 |
| C4-C6 | 1.506(8) | N1-C4-C6 | 109.8(4) | H20A-C20-H20B | 107.6 |
| C4-C5 | 1.521(7) | N1-C4-C5 | 110.2(5) | C22-C21-C20 | 114.2(5) |
| C4-H4 | 1.0000 | C6-C4-C5 | 112.4(5) | C22-C21-H21A | 108.7 |

Appendix

| | | | | | |
|---------|----------|-------------|----------|---------------|----------|
| C5-H5A | 0.9800 | N1-C4-H4 | 108.1 | C20-C21-H21A | 108.7 |
| C5-H5B | 0.9800 | C6-C4-H4 | 108.1 | C22-C21-H21B | 108.7 |
| C5-H5C | 0.9800 | C5-C4-H4 | 108.1 | C20-C21-H21B | 108.7 |
| C6-H6A | 0.9800 | C4-C5-H5A | 109.5 | H21A-C21-H21B | 107.6 |
| C6-H6B | 0.9800 | C4-C5-H5B | 109.5 | C23-C22-C21 | 125.4(6) |
| C6-H6C | 0.9800 | H5A-C5-H5B | 109.5 | C23-C22-Rh1 | 72.6(3) |
| C8-N2 | 1.467(7) | C4-C5-H5C | 109.5 | C21-C22-Rh1 | 108.8(3) |
| C8-C9 | 1.525(7) | H5A-C5-H5C | 109.5 | C23-C22-H22 | 117.3 |
| C8-H8A | 0.9900 | H5B-C5-H5C | 109.5 | C21-C22-H22 | 117.3 |
| C8-H8B | 0.9900 | C4-C6-H6A | 109.5 | Rh1-C22-H22 | 88.5 |
| C9-N3 | 1.311(7) | C4-C6-H6B | 109.5 | C22-C23-C24 | 123.0(6) |
| C9-C10 | 1.413(7) | H6A-C6-H6B | 109.5 | C22-C23-Rh1 | 71.5(3) |
| C10-C11 | 1.381(7) | C4-C6-H6C | 109.5 | C24-C23-Rh1 | 110.4(4) |
| C10-H10 | 0.9500 | H6A-C6-H6C | 109.5 | C22-C23-H23 | 118.5 |
| C11-C12 | 1.422(7) | H6B-C6-H6C | 109.5 | C24-C23-H23 | 118.5 |
| C11-H11 | 0.9500 | N2-C8-C9 | 110.7(4) | Rh1-C23-H23 | 88.1 |
| C12-C17 | 1.396(7) | N2-C8-H8A | 109.5 | C23-C24-C25 | 113.2(5) |
| C12-C13 | 1.422(7) | C9-C8-H8A | 109.5 | C23-C24-H24A | 108.9 |
| C13-C14 | 1.371(8) | N2-C8-H8B | 109.5 | C25-C24-H24A | 108.9 |
| C13-H13 | 0.9500 | C9-C8-H8B | 109.5 | C23-C24-H24B | 108.9 |
| C14-C15 | 1.405(8) | H8A-C8-H8B | 108.1 | C25-C24-H24B | 108.9 |
| C14-H14 | 0.9500 | N3-C9-C10 | 24.6(5) | H24A-C24-H24B | 107.8 |
| C15-C16 | 1.364(7) | N3-C9-C8 | 115.7(5) | C18-C25-C24 | 113.9(5) |
| C15-H15 | 0.9500 | C10-C9-C8 | 119.7(5) | C18-C25-H25A | 108.8 |
| C16-C17 | 1.425(7) | C11-C10-C9 | 118.5(5) | C24-C25-H25A | 108.8 |
| C16-H16 | 0.9500 | C11-C10-H10 | 120.8 | C18-C25-H25B | 108.8 |
| C17-N3 | 1.378(6) | C9-C10-H10 | 120.8 | C24-C25-H25B | 108.8 |

Appendix

| | | | | | |
|----------|------------|-------------|----------|---------------|------------|
| C18-C19 | 1.410(9) | C10-C11-C12 | 118.7(5) | H25A-C25-H25B | 107.7 |
| C18-C25 | 1.496(9) | C10-C11-H11 | 120.6 | C1-N1-C2 | 111.1(4) |
| C18-Rh1 | 2.113(5) | C12-C11-H11 | 120.6 | C1-N1-C4 | 125.0(4) |
| C18-H18 | 0.9500 | C17-C12-C13 | 120.0(4) | C2-N1-C4 | 123.7(4) |
| C19-C20 | 1.523(7) | C17-C12-C11 | 118.0(5) | C1-N2-C3 | 111.4(5) |
| C19-Rh1 | 2.127(5) | C13-C12-C11 | 122.0(5) | C1-N2-C8 | 125.7(5) |
| C19-H19 | 0.9500 | C14-C13-C12 | 119.1(5) | C3-N2-C8 | 122.8(5) |
| C20-C21 | 1.522(8) | C14-C13-H13 | 120.5 | C9-N3-C17 | 117.1(5) |
| C20-H20A | 0.9900 | C12-C13-H13 | 120.5 | C1-Rh1-C18 | 88.1(2) |
| C20-H20B | 0.9900 | C13-C14-C15 | 121.0(5) | C1-Rh1-C19 | 93.1(2) |
| C21-C22 | 1.495(7) | C13-C14-H14 | 119.5 | C18-Rh1-C19 | 38.8(2) |
| C21-H21A | 0.9900 | C15-C14-H14 | 119.5 | C1-Rh1-C22 | 164.3(2) |
| C21-H21B | 0.9900 | C16-C15-C14 | 120.8(5) | C18-Rh1-C22 | 96.6(2) |
| C22-C23 | 1.367(8) | C16-C15-H15 | 119.6 | C19-Rh1-C22 | 81.5(2) |
| C22-Rh1 | 2.213(5) | C14-C15-H15 | 119.6 | C1-Rh1-C23 | 159.6(2) |
| C22-H22 | 0.9500 | C15-C16-C17 | 119.5(5) | C18-Rh1-C23 | 81.2(2) |
| C23-C24 | 1.506(9) | C15-C16-H16 | 120.2 | C19-Rh1-C23 | 89.4(2) |
| C23-Rh1 | 2.226(5) | C17-C16-H16 | 120.2 | C22-Rh1-C23 | 35.9(2) |
| C23-H23 | 0.9500 | N3-C17-C12 | 123.1(5) | C1-Rh1-C11 | 89.24(13) |
| C24-C25 | 1.516(10) | N3-C17-C16 | 117.4(5) | C18-Rh1-C11 | 159.1(2) |
| C24-H24A | 0.9900 | C12-C17-C16 | 119.5(4) | C19-Rh1-C11 | 162.05(18) |
| C24-H24B | 0.9900 | C19-C18-C25 | 125.7(6) | C22-Rh1-C11 | 91.45(14) |
| C25-H25A | 0.9900 | C19-C18-Rh1 | 71.1(3) | C23-Rh1-C11 | 94.52(17) |
| C25-H25B | 0.9900 | C25-C18-Rh1 | 110.5(4) | C13-C26-C12 | 106.8(6) |
| Rh1-C11 | 2.3755(13) | C19-C18-H18 | 117.1 | C13-C26-H26A | 110.4 |
| C13-C26 | 1.758(9) | C25-C18-H18 | 117.1 | C12-C26-H26A | 110.4 |
| C12-C26 | 1.788(9) | Rh1-C18-H18 | | C13-C26-H26B | 110.4 |
| | | 88.3 | | | |

Appendix

| | | | | | |
|----------|--------|-------------|----------|---------------|-------|
| C26-H26A | 0.9900 | C18-C19-C20 | 124.5(5) | Cl2-C26-H26B | 110.4 |
| C26-H26B | 0.9900 | C18-C19-Rh1 | 70.0(3) | H26A-C26-H26B | 108.6 |

Table A1.16: Bond lengths and angles for [1-n-butyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)rhodium(I) Chloride **4.23**

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|----------|-----------------|----------|-----------------|----------|
| C1-N1 | 1.355(5) | N1-C1-N2 | 103.8(3) | C25-C18-Rh1 | 111.3(3) |
| C1-N2 | 1.368(5) | N1-C1-Rh1 | 126.4(3) | C19-C18-H18 | 117.8 |
| C1-Rh1 | 2.032(4) | N2-C1-Rh1 | 129.6(3) | C25-C18-H18 | 117.8 |
| C2-C3 | 1.340(6) | C3-C2-N2 | 106.7(4) | Rh1-C18-H18 | 87.3 |
| C2-N2 | 1.379(5) | C3-C2-H2 | 126.6 | C18-C19-C20 | 123.6(4) |
| C2-H2 | 0.9500 | N2-C2-H2 | 126.6 | C18-C19-Rh1 | 70.0(2) |
| C3-N1 | 1.387(5) | C2-C3-N1 | 107.0(4) | C20-C19-Rh1 | 113.4(3) |
| C3-H3 | 0.9500 | C2-C3-H3 | 126.5 | C18-C19-H19 | 118.2 |
| C4-N2 | 1.467(5) | N1-C3-H3 | 126.5 | C20-C19-H19 | 118.2 |
| C4-C5 | 1.508(6) | N2-C4-C5 | 112.4(3) | Rh1-C19-H19 | 86.6 |
| C4-H4A | 0.9900 | N2-C4-H4A | 109.1 | C19-C20-C21 | 114.2(3) |
| C4-H4B | 0.9900 | C5-C4-H4A | 109.1 | C19-C20-H20A | 108.7 |
| C5-N3 | 1.320(5) | N2-C4-H4B | 109.1 | C21-C20-H20A | 108.7 |
| C5-C6 | 1.405(6) | C5-C4-H4B | 109.1 | C19-C20-H20B | 108.7 |
| C6-C7 | 1.361(6) | H4A-C4-H4B | 107.9 | C21-C20-H20B | 108.7 |
| C6-H6 | 0.9500 | N3-C5-C6 | 124.6(4) | H20A-C20-H20B | 107.6 |
| C7-C8 | 1.416(6) | N3-C5-C4 | 115.6(4) | C22-C21-C20 | 113.9(3) |
| C7-H7 | 0.9500 | C6-C5-C4 | 119.8(4) | C22-C21-H21A | 108.8 |
| C8-C13 | 1.407(6) | C7-C6-C5 | 118.9(4) | C20-C21-H21A | 108.8 |

Appendix

| | | | | | |
|----------|----------|--------------|----------|---------------|----------|
| C8-C9 | 1.415(6) | C7-C6-H6 | 120.5 | C22-C21-H21B | 108.8 |
| C9-C10 | 1.364(6) | C5-C6-H6 | 120.5 | C20-C21-H21B | 108.8 |
| C9 -C9 | 0.9500 | C6-C7-C8 | 119.5(4) | H21A-C21-H21B | 107.7 |
| C10-C11 | 1.403(6) | C6-C7-H7 | 120.3 | C23-C22-C21 | 123.2(4) |
| C10-H10 | 0.9500 | C8-C7-H7 | 120.3 | C23-C22-Rh1 | 72.3(2) |
| C11-C12 | 1.360(6) | C13-C8-C9 | 119.1(4) | C21-C22-Rh1 | 109.3(3) |
| C11-H11 | 0.9500 | C13-C8-C7 | 117.4(4) | C23-C22-H22 | 118.4 |
| C12-C13 | 1.416(5) | C9-C8-C7 | 123.4(4) | C21-C22-H22 | 118.4 |
| C12-H12 | 0.9500 | C10-C9-C8 | 120.0(4) | Rh1-C22-H22 | 88.4 |
| C13-N3 | 1.380(5) | C10-C9-H9 | 120.0 | C22-C23-C24 | 123.5(4) |
| C14-N1 | 1.464(5) | C8 -9-H9 | 120.0 | C22-C23-Rh1 | 71.4(2) |
| C14-C15 | 1.518(6) | C9-C10-C11 | 120.9(4) | C24-C23-Rh1 | 111.5(3) |
| C14-H14A | 0.9900 | C9-C10-H10 | 119.6 | C22-C23-H23 | 118.2 |
| C14-H14B | 0.9900 | C11-C10-H10 | 119.6 | C24-C23-H23 | 118.2 |
| C15-C16 | 1.509(6) | C12-C11-C10 | 120.3(4) | Rh1-C23-H23 | 87.1 |
| C15-H15A | 0.9900 | C12-C11-H11 | 119.8 | C23-C24-C25 | 112.9(3) |
| C15-H15B | 0.9900 | C10-C11-H11 | 119.8 | C23-C24-H24A | 109.0 |
| C16-C17 | 1.522(6) | C11-C12-C13 | 120.2(4) | C25-C24-H24A | 109.0 |
| C16-H16A | 0.9900 | C11-C12-H12 | 119.9 | C23-C24-H24B | 109.0 |
| C16-H16B | 0.9900 | C13-C12-H12 | 119.9 | C25-C24-H24B | 109.0 |
| C17-H17A | 0.9800 | N3-C13-C8 | 123.2(4) | H24A-C24-H24B | 107.8 |
| C17-H17B | 0.9800 | N3-C13-C12 | 117.4(4) | C18-C25-C24 | 113.4(3) |
| C17-H17C | 0.9800 | C8-C13-C12 | 119.4(4) | C18-C25-H25A | 108.9 |
| C18-C19 | 1.395(5) | N1-C14-C15 | 113.3(3) | C24-C25-H25A | 108.9 |
| C18-C25 | 1.520(5) | N1-C14-H14A | 108.9 | C18-C25-H25B | 108.9 |
| C18-Rh1 | 2.097(4) | C15-C14-H14A | 108.9 | C24-C25-H25B | 108.9 |
| C18-H18 | 0.9500 | N1-C14-H14B | 108.9 | H25A-C25-H25B | 107.7 |

Appendix

| | | | | | |
|------------|----------|---------------|----------|-------------|------------|
| C19-C20 | 1.517(5) | C15-C14-H14B | 108.9 | C1-N1-C3 | 111.1(3) |
| C19-Rh1 | 2.114(4) | H14A-C14-H14B | 07.7 | C1-N1-C14 | 125.2(3) |
| C19-H19 | 0.9500 | C16-C15-C14 | 111.4(4) | C3-N1-C14 | 123.7(3) |
| C20-C21 | 1.530(6) | C16-C15-H15A | 109.3 | C1-N2-C2 | 111.3(3) |
| C20-H20A | 0.9900 | C14-C15-H15A | 109.3 | C1-N2-C4 | 124.3(3) |
| C20-H20B | 0.9900 | C16-C15-H15B | 109.3 | C2-N2-C4 | 124.3(3) |
| C21-C22 | 1.502(6) | C14-C15-H15B | 109.3 | C5-N3-C13 | 116.4(3) |
| C21-H21A | 0.9900 | H15A-C15-H15B | 108.0 | C1-Rh1-C18 | 90.62(15) |
| C21-H21B | 0.9900 | C15-C16-C17 | 113.8(4) | C1-Rh1-C19 | 95.45(15) |
| C22-C23 | 1.373(6) | C15-C16-H16A | 108.8 | C18-Rh1-C19 | 38.70(15) |
| C22-Rh1 | 2.197(4) | C17-C16-H16A | 108.8 | C1-Rh1-C22 | 163.07(15) |
| C22-H22 | 0.9500 | C15-C16-H16B | 108.8 | C18-Rh1-C22 | 97.47(16) |
| C23-C24 | 1.508(6) | C17-C16-H16B | 108.8 | C19-Rh1-C22 | 81.99(15) |
| C23-Rh1 | 2.208(4) | H16A-C16-H16B | 107.7 | C1-Rh1-C23 | 160.58(15) |
| C23-H23 | 0.9500 | C16-C17-H17A | 109.5 | C18-Rh1-C23 | 81.65(16) |
| C24-C25 | 1.530(6) | C16-C17-H17B | 109.5 | C19-Rh1-C23 | 89.62(15) |
| C24-H24A | 0.9900 | H17A-C17-H17B | 109.5 | C22-Rh1-C23 | 36.32(14) |
| C24-H24B | 0.9900 | C16-C17-H17C | 109.5 | C1-Rh1-Cl1 | 88.03(11) |
| C25-H25A | 0.9900 | H17A-C17-H17C | 109.5 | C18-Rh1-Cl1 | 159.47(11) |
| C25-H25B | 0.9900 | H17B-C17-H17C | 109.5 | C19-Rh1-Cl1 | 161.73(11) |
| 2.3786(10) | | C19-C18-C25 | 124.3(4) | C22-Rh1-Cl1 | 89.46(11) |
| | | C19-C18-Rh1 | 71.3(2) | C23-Rh1-Cl1 | 93.01(11) |

Table A1.17: Bond lengths and angles for [bis-1,3-(2-methylquinoline)imidazolin-2-ylidene](1, 5-cyclooctadiene)rhodium(I) Chloride 4.24

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|-----------|-----------------|------------|-----------------|----------|
| Rh1-Cl2 | 2.3794(7) | Cl2-Rh1-C3 | 158.31(8) | C14-N15-C16 | 117.8(2) |
| Rh1-C3 | 2.105(3) | Cl2-Rh1-C4 | 162.93(8) | N15-C16-C17 | 118.8(2) |
| Rh1-C4 | 2.121(3) | C3-Rh1-C4 | 38.70(10) | N15-C16-C21 | 122.5(2) |
| Rh1-C7 | 2.196(2) | Cl2-Rh1-C7 | 89.99(8) | C17-C16-C21 | 118.8(2) |
| Rh1-C8 | 2.230(3) | C3-Rh1-C7 | 97.74(10) | C16-C17-C18 | 120.3(3) |
| Rh1-C11 | 2.016(3) | C4-Rh1-C7 | 82.16(11) | C16-C17-H171 | 118.7 |
| C3-C4 | 1.400(4) | Cl2-Rh1-C8 | 93.69(8) | C18-C17-H171 | 120.9 |
| C3-C10 | 1.514(4) | C3-Rh1-C8 | 81.19(10) | C17-C18-C19 | 121.0(3) |
| C3-H31 | 0.979 | C4-Rh1-C8 | 88.89(10) | C17-C18-H181 | 119.9 |
| C4-C5 | 1.523(4) | C7-Rh1-C8 | 36.16(10) | C19-C18-H181 | 119.1 |
| C4-H41 | 0.989 | Cl2-Rh1-C11 | 87.46(7) | C18-C19-C20 | 120.0(3) |
| C5-C6 | 1.531(4) | C3-Rh1-C11 | 92.99(10) | C18-C19-H191 | 119.6 |
| C5-H52 | 0.972 | C4-Rh1-C11 | 93.80(10) | C20-C19-H191 | 120.4 |
| C5-H51 | 0.976 | C7-Rh1-C11 | 156.82(11) | C19-C20-C21 | 120.6(3) |
| C6-C7 | 1.505(4) | C8-Rh1-C11 | 167.02(10) | C19-C20-H201 | 119.0 |
| C6-H62 | 0.970 | Rh1-C3-C4 | 71.26(15) | C21-C20-H201 | 120.4 |
| C6-H61 | 0.980 | Rh1-C3-C10 | 110.76(18) | C16-C21-C20 | 119.3(2) |
| C7-C8 | 1.374(4) | C4-C3-C10 | 126.0(2) | C16-C21-C22 | 117.2(2) |
| C7-H71 | 0.975 | Rh1-C3-H31 | 110.3 | C20-C21-C22 | 123.5(2) |
| C8-C9 | 1.514(4) | C4-C3-H31 | 117.0 | C21-C22-C23 | 120.0(2) |
| C8-H81 | 0.977 | C10-C3-H31 | 112.4 | C21-C22-H221 | 120.6 |
| C9-C10 | 1.528(4) | C3-C4-Rh1 | 70.04(15) | C23-C22-H221 | 119.4 |

Appendix

| | | | | | |
|----------|----------|------------|------------|---------------|----------|
| C9-H92 | 0.985 | C3-C4-C5 | 125.5(3) | C14-C23-C22 | 118.9(2) |
| C9-H91 | 0.969 | Rh1-C4-C5 | 112.58(19) | C14-C23-H231 | 119.8 |
| C10-H102 | 0.967 | C3-C4-H41 | 115.5 | C22-C23-H231 | 121.3 |
| C10-H101 | 0.982 | Rh1-C4-H41 | 110.8 | N12-C24-C25 | 106.8(2) |
| C11-N12 | 1.358(3) | C5-C4-H41 | 113.4 | N12-C24-H241 | 124.4 |
| C11-N26 | 1.358(3) | C4-C5-C6 | 112.7(2) | C25-C24-H241 | 128.7 |
| N12-C13 | 1.466(3) | C4-C5-H52 | 108.3 | C24-C25-N26 | 106.3(2) |
| N12-C24 | 1.388(3) | C6-C5-H52 | 108.1 | C24-C25-H251 | 129.0 |
| C13-C14 | 1.512(3) | C4-C5-H51 | 108.5 | N26-C25-H251 | 124.7 |
| C13-H132 | 0.986 | C6-C5-H51 | 109.7 | C25-N26-C11 | 111.7(2) |
| C13-H131 | 0.982 | H52-C5-H51 | 109.5 | C25-N26-C27 | 124.2(2) |
| C14-N15 | 1.313(3) | C5-C6-C7 | 113.5(2) | C11-N26-C27 | 124.1(2) |
| C14-C23 | 1.426(4) | C5-C6-H62 | 106.9 | N26-C27-C28 | 114.2(2) |
| N15-C16 | 1.372(3) | C7-C6-H62 | 108.5 | N26-C27-H272 | 107.2 |
| C16-C17 | 1.413(4) | C5-C6-H61 | 109.2 | C28-C27-H272 | 108.9 |
| C16-C21 | 1.421(4) | C7-C6-H61 | 109.5 | N26-C27-H271 | 107.1 |
| C17-C18 | 1.367(4) | H62-C6-H61 | 109.1 | C28-C27-H271 | 109.6 |
| C17-H171 | 0.944 | C6-C7-Rh1 | 106.55(18) | H272-C27-H271 | 109.8 |
| C18-C19 | 1.409(4) | C6-C7-C8 | 126.7(3) | C27-C28-N29 | 114.6(2) |
| C18-H181 | 0.940 | Rh1-C7-C8 | 73.24(15) | C27-C28-C37 | 122.0(2) |
| C19-C20 | 1.364(4) | C6-C7-H71 | 113.8 | N29-C28-C37 | 123.3(2) |
| C19-H191 | 0.951 | Rh1-C7-H71 | 108.4 | C28-N29-C30 | 117.7(2) |
| C20-C21 | 1.416(4) | C8-C7-H71 | 116.4 | N29-C30-C31 | 123.0(2) |
| C20-H201 | 0.954 | Rh1-C8-C7 | 70.60(15) | N29-C30-C35 | 118.2(3) |
| C21-C22 | 1.416(3) | Rh1-C8-C9 | 110.60(18) | C31-C30-C35 | 118.8(3) |
| C22-C23 | 1.349(4) | C7-C8-C9 | 124.7(3) | C30-C31-C32 | 119.7(3) |
| C22-H221 | 0.944 | Rh1-C8-H81 | 108.1 | C30-C31-C36 | 117.0(2) |

Appendix

| | | | | | |
|-----------|----------|---------------|------------|---------------|----------|
| C23 -H231 | 0.941 | C7-C8-H81 | 116.6 | C32-C31-C36 | 123.3(3) |
| C24-C25 | 1.343(4) | C9-C8-H81 | 114.8 | C31-C32-C33 | 120.1(3) |
| C24-H241 | 0.945 | C8-C9-C10 | 112.6(2) | C31-C32-H321 | 118.9 |
| C25-N26 | 1.391(3) | C8-C9-H92 | 108.1 | C33-C32-H321 | 121.0 |
| C25-H251 | 0.952 | C10-C9-H92 | 107.9 | C32-C33-C34 | 120.4(3) |
| N26-C27 | 1.462(3) | C8-C9-H91 | 109.6 | C32-C33-H331 | 120.0 |
| C27-C28 | 1.516(4) | C10-C9-H91 | 109.9 | C34-C33-H331 | 119.6 |
| C27-H272 | 0.985 | H92-C9-H91 | 108.7 | C33-C34-C35 | 120.8(3) |
| C27-H271 | 0.974 | C9-C10-C3 | 112.7(2) | C33-C34-H341 | 120.4 |
| C28-N29 | 1.317(3) | C9-C10-H102 | 108.3 | C35-C34-H341 | 118.7 |
| C28-C37 | 1.414(4) | C3-C10-H102 | 108.1 | C30-C35-C34 | 120.2(3) |
| N29-C30 | 1.375(3) | C9-C10-H101 | 110.0 | C30-C35-H351 | 119.9 |
| C30-C31 | 1.406(4) | C3-C10-H101 | 109.3 | C34-C35-H351 | 119.9 |
| C30-C35 | 1.421(4) | H102-C10-H101 | 108.3 | C31-C36-C37 | 119.9(3) |
| C31-C32 | 1.418(4) | Rh1-C11-N1 | 129.86(18) | C31-C36-H361 | 120.3 |
| C31-C36 | 1.415(4) | Rh1-C11-N26 | 126.25(18) | C37-C36-H361 | 119.8 |
| C32-C33 | 1.361(4) | N12-C11-N26 | 103.7(2) | C28-C37-C36 | 119.0(3) |
| C32-H321 | 0.950 | C11-N12-C13 | 125.2(2) | C28-C37-H371 | 120.1 |
| C33-C34 | 1.404(5) | C11-N12-C24 | 111.5(2) | C36-C37-H371 | 120.9 |
| C33-H331 | 0.939 | C13-N12-C24 | 122.9(2) | Cl38-C39-Cl40 | 117.3(3) |
| C34-C35 | 1.364(4) | N12-C13-C14 | 110.8(2) | Cl38-C39-H392 | 107.3 |
| C34-H341 | 0.953 | N12-C13-H132 | 105.9 | Cl40-C39-H392 | 106.7 |
| C35-H351 | 0.951 | C14-C13-H132 | 108.1 | Cl38-C39-H391 | 108.7 |
| C36-C37 | 1.364(4) | N12-C13-H131 | 109.9 | Cl40-C39-H391 | 105.0 |
| C36-H361 | 0.955 | C14-C13-H131 | 110.5 | H392-C39-H391 | 112.0 |
| C37-H371 | 0.943 | H132-C13-H131 | 111.5 | | |
| Cl38-C39 | 1.713(4) | C13-C14-N15 | 116.6(2) | | |

Appendix

| | | | |
|----------|----------|-------------|----------|
| C39-C140 | 1.676(4) | C13-C14-C23 | 119.8(2) |
| C39-H392 | 0.994 | N15-C14-C23 | 123.6(2) |
| C39-H391 | 0.993 | | |

Table A1.18 Bond lengths and angles for 1-mesityl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) Chloride **4.28**

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|-----------|-----------------|-----------|-----------------|-----------|
| Ir1-Cl2 | 2.372(2) | Cl2-Ir1-C3 | 161.2(4) | C14-N15-C16 | 125.0(9) |
| Ir1-C3 | 2.125(13) | Cl2-Ir1-C4 | 158.2(4) | C11-N15-C16 | 124.7(9) |
| Ir1-C4 | 2.096(11) | C3-Ir1-C4 | 40.4(4) | N15-C16-C17 | 117.0(9) |
| Ir1-C7 | 2.159(11) | Cl2-Ir1-C7 | 91.5(3) | N15-C16-C22 | 120.4(9) |
| Ir1-C8 | 2.177(11) | C3-Ir1-C7 | 92.5(5) | C17-C16-C22 | 122.5(10) |
| Ir1-C11 | 2.051(11) | C4-Ir1-C7 | 79.9(4) | C16-C17-C18 | 121.1(10) |
| C3-C4 | 1.456(16) | Cl2-Ir1-C8 | 89.6(4) | C16-C17-C19 | 117.5(9) |
| C3-C10 | 1.532(18) | C3-Ir1-C8 | 82.5(5) | C18-C17-C19 | 121.4(9) |
| C3-H31 | 0.991 | C4-Ir1-C8 | 95.3(5) | C17-C18-H181 | 109.3 |
| C4-C5 | 1.509(18) | C7-Ir1-C8 | 37.4(4) | C17-C18-H182 | 109.9 |
| C4-H41 | 0.984 | Cl2-Ir1-C11 | 87.7(3) | H181-C18-H182 | 108.9 |
| C5-C6 | 1.477(15) | C3-Ir1-C11 | 93.1(4) | C17-C18-H183 | 110.7 |
| C5-H51 | 0.970 | C4-Ir1-C11 | 95.3(4) | H181-C18-H183 | 109.3 |
| C5-H52 | 0.970 | C7-Ir1-C11 | 164.6(5) | H182-C18-H183 | 108.8 |
| C6-C7 | 1.510(17) | C8-Ir1-C11 | 157.9(5) | C17-C19-C20 | 120.9(10) |
| C6-H61 | 0.975 | Ir1-C3-C4 | 68.7(8) | C17-C19-H191 | 119.5 |
| C6-H62 | 0.969 | Ir1-C3-C10 | 113.8(9) | C20-C19-H191 | 119.7 |
| C7-C8 | 1.389(16) | C4-C3-C10 | 118.7(13) | C19-C20-C21 | 119.2(10) |
| C7-H71 | 0.984 | Ir1-C3-H31 | 115.0 | C19-C20-C24 | 119.0(11) |

Appendix

| | | | | | |
|----------|-----------|------------|-----------|---------------|-----------|
| C8-C9 | 1.512(18) | C4-C3-H31 | 116.0 | C21-C20-C24 | 121.7(11) |
| C8-H81 | 0.981 | C10-C3-H31 | 116.0 | C20-C21-C22 | 123.1(11) |
| C9-C10 | 1.512(16) | C3-C4-Ir1 | 70.9(7) | C20-C21-H211 | 117.8 |
| C9-H91 | 0.968 | C3-C4-C5 | 128.6(13) | C22-C21-H211 | 119.1 |
| C9-H92 | 0.976 | Ir1-C4-C5 | 113.7(7) | C16-C22-C21 | 116.5(11) |
| C10-H101 | 0.972 | C3-C4-H41 | 111.7 | C16-C22-C23 | 120.5(11) |
| C10-H102 | 0.971 | Ir1-C4-H41 | 112.2 | C21-C22-C23 | 122.9(11) |
| C11-N12 | 1.367(15) | C5-C4-H41 | 112.5 | C22-C23-H231 | 109.5 |
| C11-N15 | 1.347(14) | C4-C5-C6 | 113.3(14) | C22-C23-H232 | 108.8 |
| N12-C13 | 1.389(14) | C4-C5-H51 | 108.1 | H231-C23-H232 | 109.5 |
| N12-C25 | 1.433(14) | C6-C5-H51 | 108.2 | C22-C23-H233 | 111.7 |
| C13-C14 | 1.330(16) | C4-C5-H52 | 108.6 | H231-C23-H233 | 108.3 |
| C13-H131 | 0.935 | C6-C5-H52 | 108.9 | H232-C23-H233 | 109.1 |
| C14-N15 | 1.402(13) | H51-C5-H52 | 109.8 | C20-C24-H241 | 108.2 |
| C14-H141 | 0.936 | C5-C6-C7 | 113.8(12) | C20-C24-H242 | 108.8 |
| N15-C16 | 1.426(13) | C5-C6-H61 | 108.0 | H241-C24-H242 | 110.1 |
| C16-C17 | 1.413(14) | C7-C6-H61 | 108.8 | C20-C24-H243 | 109.9 |
| C16-C22 | 1.416(14) | C5-C6-H62 | 107.7 | H241-C24-H243 | 110.5 |
| C17-C18 | 1.489(14) | C7-C6-H62 | 108.5 | H242-C24-H243 | 109.3 |
| C17-C19 | 1.406(15) | H61-C6-H62 | 110.0 | N12-C25-C26 | 114.3(9) |
| C18-H181 | 0.964 | Ir1-C7-C6 | 113.4(7) | N12-C25-H251 | 107.9 |
| C18-H182 | 0.960 | Ir1-C7-C8 | 72.0(7) | C26-C25-H251 | 108.1 |
| C18-H183 | 0.966 | C6-C7-C8 | 123.6(14) | N12-C25-H252 | 108.4 |
| C19-C20 | 1.380(15) | Ir1-C7-H71 | 112.7 | C26-C25-H252 | 107.8 |
| C19-H191 | 0.936 | C6-C7-H71 | 114.2 | H251-C25-H252 | 110.3 |
| C20-C21 | 1.401(18) | C8-C7-H71 | 113.6 | C25-C26-N27 | 116.9(9) |
| C20-C24 | 1.527(16) | Ir1-C8-C7 | 70.6(7) | C25-C26-C35 | 119.9(10) |

Appendix

| | | | | | |
|----------|-----------|---------------|-----------|--------------|-----------|
| C21-C22 | 1.372(16) | Ir1-C8-C9 | 109.2(8) | N27-C26-C35 | 123.2(10) |
| C21-H211 | 0.930 | C7-C8-C9 | 123.5(14) | C26-N27-C28 | 116.9(9) |
| C22-C23 | 1.497(16) | Ir1-C8-H81 | 113.6 | N27-C28-C29 | 117.3(9) |
| C23-H231 | 0.966 | C7-C8-H81 | 115.7 | N27-C28-C33 | 123.3(10) |
| C23-H232 | 0.964 | C9-C8-H81 | 115.0 | C29-C28-C33 | 119.4(10) |
| C23-H233 | 0.964 | C8-C9-C10 | 117.0(11) | C28-C29-C30 | 119.8(10) |
| C24-H241 | 0.961 | C8-C9-H91 | 107.8 | C28-C29-H291 | 119.6 |
| C24-H242 | 0.963 | C10-C9-H91 | 109.0 | C30-C29-H291 | 120.6 |
| C24-H243 | 0.959 | C8-C9-H92 | 106.9 | C29-C30-C31 | 120.3(12) |
| C25-C26 | 1.502(15) | C10-C9-H92 | 107.3 | C29-C30-H301 | 119.4 |
| C25-H251 | 0.970 | H91-C9-H92 | 108.7 | C31-C30-H301 | 120.3 |
| C25-H252 | 0.975 | C3-C10-C9 | 113.0(12) | C30-C31-C32 | 119.9(11) |
| C26-N27 | 1.348(13) | C3-C10-H101 | 108.7 | C30-C31-H311 | 120.4 |
| C26-C35 | 1.412(15) | C9-C10-H101 | 108.8 | C32-C31-H311 | 119.7 |
| N27-C28 | 1.379(13) | C3-C10-H102 | 108.8 | C31-C32-C33 | 121.4(11) |
| C28-C29 | 1.413(14) | C9-C10-H102 | 107.8 | C31-C32-H321 | 119.9 |
| C28-C33 | 1.413(15) | H101-C10-H102 | 109.6 | C33-C32-H321 | 118.7 |
| C29-C30 | 1.371(14) | Ir1-C11-N12 | 123.7(8) | C28-C33-C32 | 119.2(11) |
| C29-H291 | 0.936 | Ir1-C11-N15 | 130.6(9) | C28-C33-C34 | 117.7(11) |
| C30-C31 | 1.421(17) | N12-C11-N15 | 105.6(9) | C32-C33-C34 | 123.1(11) |
| C30-H301 | 0.938 | C11-N12-C13 | 110.2(9) | C33-C34-C35 | 119.3(12) |
| C31-C32 | 1.356(16) | C11-N12-C25 | 125.3(9) | C33-C34-H341 | 119.5 |
| C31-H311 | 0.938 | C13-N12-C25 | 123.9(9) | C35-C34-H341 | 121.2 |
| C32-C33 | 1.395(16) | N12-C13-C14 | 106.8(10) | C26-C35-C34 | 119.6(11) |
| C32-H321 | 0.935 | N12-C13-H13 | 125.6 | C26-C35-H351 | 119.2 |
| | | C14-C13-H131 | 127.6 | C34-C35-H351 | 121.2 |
| | | C13-C14-N15 | 107.9(10) | | |

Appendix

| | | | |
|----------|-----------|--------------|-----------|
| C33-C34 | 1.415(17) | C13-C14-H141 | 126.7 |
| C34-C35 | 1.374(17) | N15-C14-H141 | 125.4 |
| C34-H341 | 0.934 | C14-N15-C11 | 109.5(10) |
| C35-H351 | 0.937 | | |

Table A1.19: Bond lengths and angles for [1-isopropyl-3-(2-methylquinoline)imidazolin-2-ylidene](1,5-cyclooctadiene)iridium(I) Chloride **4.29**

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|-----------|-----------------|----------|-----------------|----------|
| C1-N1 | 1.357(8) | N1-C1-N2 | 103.7(5) | C1-N1-C2 | 111.0(6) |
| C1-N2 | 1.367(8) | N1-C1-Ir1 | 130.1(5) | C1-N1-C4 | 124.6(5) |
| C1-Ir1 | 2.045(6) | N2-C1-Ir1 | 125.8(5) | C2-N1-C4 | 124.1(6) |
| C2-C3 | 1.340(10) | C3-C2-N1 | 106.9(6) | C1-N2-C3 | 111.6(6) |
| C2-N1 | 1.390(8) | C2-C3-N2 | 106.8(6) | C1-N2-C7 | 124.5(6) |
| C3-N2 | 1.374(9) | N1-C4-C5 | 110.1(6) | C3-N2-C7 | 123.9(6) |
| C4-N1 | 1.471(9) | N1-C4-C6 | 109.7(6) | C8-N3-C16 | 118.0(6) |
| C4-C5 | 1.518(12) | C5-C4-C6 | 112.6(7) | C1-Ir1-C18 | 88.5(3) |
| C4-C6 | 1.531(10) | N2-C7-C8 | 111.2(6) | C1-Ir1-C17 | 93.1(3) |
| C7-N2 | 1.472(9) | N3-C8-C9 | 123.1(7) | C18-Ir1-C17 | 39.3(3) |
| C7-C8 | 1.499(10) | N3-C8-C7 | 116.8(7) | C1-Ir1-C22 | 163.6(3) |
| C8-N3 | 1.324(9) | C9-C8-C7 | 120.1(7) | C18-Ir1-C22 | 97.1(3) |
| C8-C9 | 1.432(10) | C10-C9-C8 | 119.2(7) | C17-Ir1-C22 | 81.8(3) |
| C9-C10 | 1.358(11) | C9-C10-C11 | 119.8(7) | C1-Ir1-C21 | 160.0(3) |
| C10-C11 | 1.411(10) | C10-C11-C16 | 117.8(6) | C18-Ir1-C21 | 81.0(3) |
| C11-C16 | 1.429(9) | C10-C11-C12 | 123.8(7) | C17-Ir1-C21 | 89.4(3) |
| C11-C12 | 1.439(10) | C16-C11-C12 | 118.5(6) | C22-Ir1-C21 | 36.3(3) |

Appendix

| | | | | | |
|---------|------------|--------------|----------|-------------|-----------|
| C12-C13 | 1.368(12) | C13-C12-C11 | 120.4(7) | C1-Ir1-C11 | 90.09(18) |
| C13-C14 | 1.432(12) | C12-C13-C14 | 120.4(7) | C18-Ir1-C11 | 159.1(2) |
| C14-C15 | 1.366(10) | C15-C14-C13 | 120.2(7) | C17-Ir1-C11 | 161.5(2) |
| C15-C16 | 1.413(9) | C14-C15-C16 | 120.9(7) | C22-Ir1-C11 | 90.04(19) |
| C16-N3 | 1.379(9) | N3-C16-C15 | 118.4(6) | C21-Ir1-C11 | 93.8(2) |
| C17-C18 | 1.419(11) | N3-C16-C11 | 122.1(6) | C25-C12-C25 | 41.5(10) |
| C17-C24 | 1.531(10) | C15-C16-C11 | 119.5(6) | C13-C13-C25 | 85.6(6) |
| C17-Ir1 | 2.112(6) | C18-C17-C24 | 124.2(7) | C13-C13-C25 | 56.5(5) |
| C18-C19 | 1.509(11) | C18-C17-Ir1 | 70.0(4) | C25-C13-C25 | 35.5(9) |
| C18-Ir1 | 2.104(7) | C24-C17-Ir1 | 114.4(5) | C12-C25-C13 | 105.2(9) |
| C19-C20 | 1.513(12) | C17-C18-C19 | 124.6(7) | | |
| C20-C21 | 1.532(11) | C17-C18-Ir1 | 70.6(4) | | |
| C21-C22 | 1.367(11) | C19-C18-Ir1 | 112.2(5) | | |
| C21-Ir1 | 2.196(7) | C18-C19-C20 | 113.3(6) | | |
| C22-C23 | 1.533(10) | C19-C20-C21 | 112.3(6) | | |
| C22-Ir1 | 2.195(6) | C22-C21-C20 | 122.9(7) | | |
| C23-C24 | 1.547(11) | C22-C21-Ir1 | 71.8(4) | | |
| C11-Ir1 | 2.3719(16) | C20-C21-Ir1 | 111.8(5) | | |
| C12-C25 | 1.648(17) | C21-C22-C23 | 124.3(7) | | |
| C12-C25 | 1.786(18) | C21-C22-Ir1 | 71.9(4) | | |
| C13-C13 | 1.296(13) | C23-C22-Ir1 | 110.0(4) | | |
| C13-C25 | 1.759(14) | C22-C23-C240 | 112.6(6) | | |
| C13-C25 | 2.103(16) | C17-C24-C23 | 113.9(5) | | |

Appendix

Table A1.20: Bond lengths and angles for [bis-1,3-(2-methylquinoline)imidazolin-2-ylidene](1, 5-cyclooctadiene)iridium(I) Chloride 4.31

| Bond lengths (Å) | | Bond angles (°) | | Bond angles (°) | |
|------------------|------------|-----------------|------------|-----------------|----------|
| Ir1-C12 | 2.3653(11) | C12-Ir1-C3 | 158.86(14) | H132-C13-H131 | 112.3 |
| Ir1-C3 | 2.082(4) | C12-Ir1-C4 | 161.77(14) | C13-C14-N15 | 116.5(4) |
| Ir1-C4 | 2.121(4) | C3-Ir1-C4 | 39.36(19) | C13-C14-C19 | 119.7(4) |
| Ir1-C7 | 2.160(4) | C12-Ir1-C7 | 90.72(14) | N15-C14-C19 | 123.8(4) |
| Ir1-C8 | 2.196(5) | C3-Ir1-C7 | 98.15(19) | C14-N15-C16 | 117.3(4) |
| Ir1-C11 | 2.022(4) | C4-Ir1-C7 | 82.45(19) | N15-C16-C17 | 123.4(4) |
| C3-C4 | 1.416(7) | C12-Ir1-C8 | 94.70(14) | N15-C16-C23 | 117.8(4) |
| C3-C10 | 1.518(6) | C3-Ir1-C8 | 81.47(18) | C17-C16-C23 | 118.7(4) |
| C3-H31 | 0.960 | C4-Ir1-C8 | 89.95(18) | C16-C17-C18 | 116.9(4) |
| C4-C5 | 1.515(6) | C7-Ir1-C8 | 36.9(2) | C16-C17-C20 | 119.1(4) |
| C4-H41 | 0.998 | C12-Ir1-C11 | 87.54(12) | C18-C17-C20 | 124.0(4) |
| C5-C6 | 1.543(8) | C3-Ir1-C11 | 92.36(17) | C17-C18-C19 | 119.6(4) |
| C5-H52 | 0.987 | C4-Ir1-C11 | 91.42(17) | C17-C18-H181 | 120.5 |
| C5-H51 | 0.977 | C7-Ir1-C11 | 154.57(19) | C19-C18-H181 | 119.9 |
| C6-C7 | 1.531(7) | C8-Ir1-C11 | 168.44(18) | C14-C19-C18 | 118.9(4) |
| C6-H62 | 0.982 | Ir1-C3-C4 | 71.8(3) | C14-C19-H191 | 121.6 |
| C6-H61 | 0.965 | Ir1-C3-C10 | 113.7(3) | C18-C19-H191 | 119.5 |
| C7-C8 | 1.379(8) | C4-C3-C10 | 124.7(4) | C17-C20-C21 | 120.3(5) |
| C7-H71 | 0.990 | Ir1-C3-H31 | 114.7 | C17-C20-H201 | 120.0 |
| C8-C9 | 1.534(8) | C4-C3-H31 | 112.8 | C21-C20-H201 | 119.7 |
| C8-H81 | 0.996 | C10-C3-H31 | 112.9 | C20-C21-C22 | 121.2(5) |
| C9-C10 | 1.554(7) | Ir1-C4-C3 | 68.8(2) | C20-C21-H211 | 119.4 |
| C9-H92 | 0.981 | Ir1-C4-C5 | 113.5(3) | C22-C21-H211 | 119.4 |

Appendix

| | | | | | |
|----------|----------|------------|----------|---------------|----------|
| C9-H91 | 0.984 | C3-C4-C5 | 124.5(4) | C21-C22-C23 | 119.1(5) |
| C10-H101 | 0.968 | Ir1-C4-H41 | 111.6 | C21-C22-H221 | 119.3 |
| C10-H102 | 0.965 | C3-C4-H41 | 116.5 | C23-C22-H221 | 121.7 |
| C11-N12 | 1.354(5) | C5-C4-H41 | 113.1 | C16-C23-C22 | 121.5(5) |
| C11-N26 | 1.361(5) | C4-C5-C6 | 112.8(4) | C16-C23-H231 | 117.8 |
| N12-C13 | 1.461(5) | C4-C5-H52 | 108.6 | C22-C23-H231 | 120.7 |
| N12-C24 | 1.383(5) | C6-C5-H52 | 106.7 | N12-C24-C25 | 106.7(4) |
| C13-C14 | 1.512(6) | C4-C5-H51 | 110.6 | N12-C24-H241 | 125.3 |
| C13-H132 | 0.971 | C6-C5-H51 | 108.1 | C25-C24-H241 | 128.0 |
| C13-H131 | 0.978 | H52-C5-H51 | 109.9 | C24-C25-N26 | 106.6(4) |
| C14-N15 | 1.318(5) | C5-C6-C7 | 113.0(4) | C24-C25-H251 | 127.7 |
| C14-C19 | 1.424(6) | C5-C6-H62 | 108.2 | N26-C25-H251 | 125.7 |
| N15-C16 | 1.372(6) | C7-C6-H62 | 108.8 | C25-N26-C11 | 111.1(3) |
| C16-C17 | 1.414(6) | C5-C6-H61 | 108.4 | C25-N26-C27 | 126.3(4) |
| C16-C23 | 1.420(6) | C7-C6-H61 | 111.6 | C11-N26-C27 | 122.5(3) |
| C17-C18 | 1.426(6) | H62-C6-H61 | 106.6 | N26-C27-C28 | 114.2(3) |
| C17-C20 | 1.418(6) | C6-C7-Ir1 | 108.6(3) | N26-C27-H271 | 107.6 |
| C18-C19 | 1.362(7) | C6-C7-C8 | 25.2(5) | C28-C27-H271 | 109.0 |
| C18-H181 | 0.939 | Ir1-C7-C8 | 72.9(3) | N26-C27-H272 | 108.6 |
| C19-H191 | 0.955 | C6-C7-H71 | 113.7 | C28-C27-H272 | 110.1 |
| C20-C21 | 1.365(7) | Ir1-C7-H71 | 114.9 | H271-C27-H272 | 107.2 |
| C20-H201 | 0.965 | C8-C7-H71 | 114.1 | C27-C28-N29 | 114.5(4) |
| C21-C22 | 1.416(8) | Ir1-C8-C7 | 70.1(3) | C27-C28-C33 | 121.4(4) |
| C21-H211 | 0.931 | Ir1-C8-C9 | 111.9(3) | N29-C28-C33 | 123.9(4) |
| C22-C23 | 1.364(7) | C7-C8-C9 | 123.8(5) | C28-N29-C30 | 117.0(4) |
| C22-H221 | 0.954 | Ir1-C8-H81 | 112.1 | N29-C30-C31 | 122.6(4) |
| C23-H231 | 0.942 | C7-C8-H81 | 115.0 | N29-C30-C37 | 118.8(4) |

Appendix

| | | | | | |
|----------|----------|---------------|----------|--------------|----------|
| C24-C25 | 1.344(6) | C9-C8-H81 | 114.8 | C31-C30-C37 | 118.6(4) |
| C24-H241 | 0.951 | C8-C9-10 | 112.3(4) | C30-C31-C32 | 117.8(4) |
| C25-N26 | 1.388(5) | C8-C9-H92 | 107.7 | C30-C31-C34 | 119.2(5) |
| C25-H251 | 0.947 | C10-C9-H92 | 110.8 | C32-C31-C34 | 123.0(5) |
| N26-C27 | 1.460(5) | C8-C9-H91 | 111.4 | C31-C32-C33 | 119.3(4) |
| C27-C28 | 1.521(6) | C10-C9-H91 | 109.2 | C31-C32-H321 | 119.9 |
| C27-H271 | 0.981 | H92-C9-H91 | 105.2 | C33-C32-H321 | 120.8 |
| C27-H272 | 0.969 | C9-C10-C3 | 111.7(4) | C28-C33-C32 | 119.2(4) |
| C28-N29 | 1.327(6) | C9-C10-H101 | 111.3 | C28-C33-H331 | 120.2 |
| C28-C33 | 1.415(6) | C3-C10-H101 | 111.3 | C32-C33-H331 | 120.6 |
| N29-C30 | 1.374(6) | C9-C10-H102 | 106.3 | C31-C34-C35 | 121.0(5) |
| C30-C31 | 1.415(7) | C3-C10-H102 | 106.5 | C31-C34-H341 | 119.1 |
| C30-C37 | 1.422(7) | H101-C10-H102 | 109.5 | C35-C34-H341 | 119.9 |
| C31-C32 | 1.416(7) | Ir1-C11-N12 | 130.5(3) | C34-C35-C36 | 120.0(5) |
| C31-C34 | 1.415(7) | Ir1-C11-N26 | 125.4(3) | C34-C35-H351 | 118.8 |
| C32-C33 | 1.357(7) | N12-C11-N26 | 104.0(3) | C36-C35-H351 | 121.1 |
| C32-H321 | 0.956 | C11-N12-C13 | 123.7(3) | C35-C36-C37 | 120.3(5) |
| C33-H331 | 0.940 | C11-N12-C24 | 111.5(3) | C35-C36-H361 | 120.5 |
| C34-C35 | 1.366(8) | C13-N12-C24 | 124.7(3) | C37-C36-H361 | 119.2 |
| C34-H341 | 0.944 | N12-C13-C14 | 111.2(4) | C30-C37-C36 | 120.8(5) |
| C35-C36 | 1.412(9) | N12-C13-H132 | 107.2 | C30-C37-H371 | 119.5 |
| C35-H351 | 0.933 | C14-C13-H132 | 109.4 | C36-C37-H371 | 119.7 |
| C36-C37 | 1.367(8) | N12-C13-H131 | 107.7 | | |
| C36-H361 | 0.946 | C14-C13-H131 | 108.9 | | |
| C37-H371 | 0.935 | | | | |

