

**Effect of plant *WEE1* on the cell  
cycle and development in *Arabidopsis  
thaliana* and *Nicotiana tabacum***

**A thesis submitted for the degree of Doctor of Philosophy**

**by**

**Ilario Siciliano**

**School of Biosciences  
University of Wales, Cardiff  
Park Place  
Cardiff**

**April 2006**

UMI Number: U584855

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI U584855

Published by ProQuest LLC 2013. Copyright in the Dissertation held by the Author.  
Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against  
unauthorized copying under Title 17, United States Code.



ProQuest LLC  
789 East Eisenhower Parkway  
P.O. Box 1346  
Ann Arbor, MI 48106-1346

## **Acknowledgements**

I would like to acknowledge Cardiff University and Worcester University for providing funding for this project.

I would like to thank my supervisors, Dr Dennis Francis, Dr Hilary Rogers and Dr Robert Herbert for their advice and support throughout the different phases of this project.

I am grateful to Mr Mike O'Reilly for his help and technical support.

I would also like to thank the rest of the lab members for their help and companionship.

I am deeply indebted to my parents and siblings who supported and encouraged me during this time.

This thesis is dedicated to my wife Miguella. Without her moral support, wise opinions and suggestions it would never have been completed.

# **Contents**

## **Chapter 1: General introduction**

|   |    |
|---|----|
| 1.1. The eukaryotic cell cycle and its regulation   | 1  |
| 1.1.1. Cyclin-dependent kinase (CDK)  | 3  |
| 1.1.2. Cyclins  | 8  |
| 1.1.3. CDK-Activating Kinases   | 13 |
| 1.1.4. Cyclin-dependent kinase inhibitor (CKIs)   | 15 |
| 1.2. Comparison between the CDK/cyclin complexes during the different phases of the cell cycle in yeast, animals and plants | 16 |
| 1.2.1. Budding yeast cell cycle   | 16 |
| 1.2.2. Animal cell cycle  | 18 |
| 1.2.3 Plant cell cycle  | 20 |
| 1.3. Retinoblastoma protein and the E2F transcription factor regulate G1/S transition                                       | 23 |
| 1.4. Cell-cycle regulation at the G2/M transition in fission yeast, animals and plants                                      | 25 |
| 1.5. The role of <i>WEE1</i> during the G2/M transition   | 29 |
| 1.5.1. <i>WEE1</i> plays an important role in checkpoint control  | 29 |
| 1.5.2. The <i>WEE1</i> protein kinase is regulated at multiple levels   | 30 |
| 1.5.3. <i>WEE1</i> protein levels fluctuated during the cell cycle  | 34 |
| 1.6. Tobacco BY-2 cells   | 34 |
| 1.7. Aims   | 37 |

## **Chapter 2: Materials and Methods**

|   |    |
|---|----|
| 2.1. BIN HYG TX and pTA7002 plasmids  | 39 |
| 2.1.1. Construction of plasmid  | 39 |
| 2.2. Transformation into bacterial cells and tobacco cells                        | 43 |
| 2.2.1. <i>Escherichia coli</i> DH5 $\alpha$ transformation                        | 43 |
| 2.2.2. <i>Agrobacterium tumefaciens</i> LBA4404 transformation                    | 45 |
| 2.2.3. Transformation of tobacco BY-2 cells with <i>AtWEE1</i>                    | 45 |
| 2.3. Tobacco BY-2 cells synchronization, mitotic index and cell area measurements | 46 |
| 2.3.1. Tobacco BY-2 cell culture propagation                                      | 46 |
| 2.3.2. Synchronization of tobacco BY-2 cells                                      | 48 |
| 2.3.3. Mitotic index measurements   | 49 |
| 2.3.4. Cell area  | 50 |
| 2.3.5. Growth rate measurement  | 51 |
| 2.3.6. Tobacco BY-2 cells containing pTA7002                                      | 51 |
| 2.4. RT-PCR and semi quantitative RT-PCR  | 51 |
| 2.4.1. RNA extraction   | 51 |
| 2.4.2. DNase treatment of cDNA  | 52 |
| 2.4.3. cDNA synthesis   | 53 |
| 2.4.4. PCR amplification of cDNA  | 54 |
| 2.4.5. Design of degenerate and non-degenerate primers                            | 54 |
| 2.4.6. Estimation of optimum PCR cycles for semi-quantitative RT-PCR              | 56 |
| 2.4.7. Semi-quantitative RT-PCR   | 56 |



**Chapter 3: Expression of *Arabidopsis thaliana* WEE1 (*AtWEE1*) in TBY-2 cells induces a shortened G2 phase, a premature entry into mitosis and a smaller mitotic cell area**

|  |    |
|--|----|
| 3.1. Introduction  | 58 |
| 3.2. Materials and methods   | 60 |
| 3.3. Results: Production of tobacco BY-2 cell lines transformed with <i>AtWEE1</i> constructs and empty vector controls        | 60 |
| 3.4. Results: Analysis of tobacco BY-2 cells transformed with BIN HYG TX- <i>AtWEE1</i>  | 62 |
| 3.4.1. Comparison of growth rate between tobacco BY-2 cell lines expressing <i>AtWEE1</i> , wild type and empty vector         | 64 |
| 3.4.2. Mitotic cell area is smaller in TBY-2 cells expressing <i>AtWEE1</i> over a 6 day period                                | 65 |
| 3.4.3. Expression of <i>AtWEE1</i> in TBY-2 cell lines induces a premature entry into mitosis through a shortening of G2 phase | 69 |
| 3.4.4. Can hygromycin affect the cell cycle?   | 75 |
| 3.4.5. Mitotic cell area is smaller in the constitutive <i>AtWEE1</i> expressing lines   | 78 |
| 3.5. Results: TBY-2 cells transformed with pTA7002- <i>AtWEE1</i>  | 81 |
| 3.5.1. Mitotic cell area in the inducible <i>AtWEE1</i> cell lines   | 83 |
| 3.5.2. Can dexamethasone affect cell size?   | 86 |
| 3.5.3. Induction of <i>AtWEE1</i> causes tobacco BY-2 cells to divide prematurely  | 88 |
| 3.5.4. Inducible tobacco BY-2 cells divide at smaller mitotic cell areas   | 93 |
| 3.6. Discussion  | 95 |
| 3.7. Summary   | 98 |

**Chapter 4: In tobacco BY-2 cells, induction of *AtWEE1* delays expression of the endogenous *Nicotiana tabacum* WEE1 (*NtWEE1*) gene**

|  |     |
|--|-----|
| 4.1. Introduction  | 100 |
| 4.2. Materials and methods   | 102 |
| 4.3. Results   | 102 |
| 4.3.1. Isolation of a homologue tobacco <i>WEE1</i> ( <i>NtWEE1</i> )                                      | 102 |
| 4.3.2. Identification of the optimum number of PCR cycles  | 104 |
| 4.3.3. Statistical measure of semi-quantitative RT-PCR results   | 106 |
| 4.3.4. Comparison of <i>NtWEE1</i> expression between wild type, WEE1-i-1 non-induced and WEE1-i-1 induced | 107 |
| 4.3.4.1. Expression of <i>NtWEE1</i> in wild type cells during the cell cycle                              | 107 |
| 4.3.4.2. Expression of <i>NtWEE1</i> in non induced WEE1-i-1 cells during the cell cycle                   | 109 |
| 4.3.4.3. Expression of <i>NtWEE1</i> and <i>AtWEE1</i> in induced WEE1-i-1 cells during the cell cycle     | 111 |
| 4.4. Discussion  | 114 |
| 4.5. Summary   | 117 |

**Chapter 5: CDKA and CDKB1 kinase activity through the cell cycle in synchronized TBV-2 cells expressing *AtWEE1***

|  |     |
|--|-----|
| 5.1. Introduction  | 118 |
| 5.2. Materials and methods   | 120 |
| 5.2.1. Protein extraction  | 120 |
| 5.2. 2. Kinase Activity  | 122 |
| 5.3. Results   | 123 |
| 5.3.1. Kinase activity of CDKs in TBV-2 cells expressing <i>AtWEE1</i>   | 123 |
| 5.3.1.1. Kinase activity of CDKA and CDKB1 in non-induced WEE1-i-1 cells | 124 |
| 5.3.1.2. CDKA and CDKB1 kinase activity in induced WEE1-i-1 cells        | 127 |
| 5.4. Discussion  | 130 |
| 5.5. Summary   | 133 |

**Chapter 6: Over expression of *AtWEE1* alters root growth and development in *Arabidopsis thaliana***

|  |     |
|--|-----|
| 6.1. Introduction  | 134 |
| 6.2. Materials and methods   | 135 |
| 6.2.1. Isolation of transgenic lines   | 136 |
| 6.2.2. RNA extraction, cDNA synthesis and RT-PCR   | 139 |
| 6.2.3. Seed sterilization, growth conditions and methods of fixation   | 140 |
| 6.2.4. <i>AtWEE1</i> lines: growth conditions and sampling methods   | 140 |
| 6.2.5. Analysis of lateral root development and primordia size   | 141 |
| 6.3. Results   | 142 |
| 6.3.1. Constitutive and inducible over-expression of <i>AtWEE1</i> in transgenic <i>Arabidopsis</i> plants leads to a smaller length of primary root per unit time | 146 |
| 6.3.2. BIN HYG TX- <i>AtWEE1</i> plants initiate primordia at a slower rate per day than wild type   | 149 |
| 6.3.3. <i>AtWEE1</i> over-expression results in a reduction in the number of lateral roots   | 152 |
| 6.4. Discussion  | 153 |
| 6.5. Summary   | 157 |

**Chapter 7: General Discussion** 158

|              |     |
|--------------|-----|
| Further Work | 165 |
|--------------|-----|

**References** 169

**Appendix I: Sequences of *AtWEE1* ORF** 189

**Appendix II: Tobacco BY-2 cell area raw data** 195

**Appendix III: Tobacco BY-2 mitotic index (MI %)** 217

**Appendix IV: 18S rRNA, H4, *AtWEE1* and *NtWEE1* expression raw data** 220

**Appendix V: CDKA and CDKB1 kinase activity raw data** 223

**Appendix VI: *Arabidopsis* roots measurements raw data** 224

**Appendix VII: The yeast two hybrid technique was employed to investigate possible interaction between AtWEE1 and proteins that regulate its activity** 229

## List of Abbreviations

|                                 |  |
|---------------------------------|--|
| ATP                             | adenosine triphosphate                             |
| <i>AtWEE1</i>                   | <i>Arabidopsis thaliana WEE1</i>                   |
| bp                              | base pair  |
| °C                              | degrees centigrade                                 |
| cdc                             | cell division cycle                                |
| cDNA                            | complementary deoxyribonucleic acid                |
| CDK                             | cyclin dependent kinase                            |
| Cyc                             | cyclin   |
| C-terminus                      | carboxy-terminus                                   |
| 2,4-D                           | 2,4-dinitrophenoxyacetic acid                      |
| DEX or dex                      | dexamethasone                                      |
| dH <sub>2</sub> O               | distilled water                                    |
| DMSO                            | dimethylsulfoxide                                  |
| DNA                             | deoxyribonucleic acid                              |
| Dnase                           | deoxyribonuclease                                  |
| dNTP                            | deoxy nucleotide triphosphate                      |
| DTT                             | dithiothreitol                                     |
| <i>E. coli</i>                  | <i>Escherichia coli</i>                            |
| EDTA                            | ethylene diamine tetraacetic acid                  |
| EGTA                            | ethylenedis (oxyethylenenitrilo) tetra-acetic acid |
| EtBr                            | ethidium bromide                                   |
| g                               | gravity force                                      |
| G1                              | gap 1  |
| G2                              | gap 2  |
| Hyg                             | hygromycin   |
| KH <sub>2</sub> PO <sub>4</sub> | potassium dihydrogen orthophosphate                |
| M-phase                         | mitosis  |
| M                               | molar  |
| MI                              | mitotic index                                      |

|                      |  |
|----------------------|--|
| MPF                  | Maturation promoting factor                              |
| M & S                | Murashige and Skoog medium                               |
| mRNA                 | messenger ribonucleic acid                               |
| N-terminus           | Amino-terminus   |
| OD                   | optical density  |
| ORF                  | open reading frame                                       |
| PAGE                 | polyacrylamide gel electrophoresis                       |
| PBS                  | phosphate buffered saline solution                       |
| PCR                  | polymerase chain reaction                                |
| Rb                   | retinoblastoma protein                                   |
| RNA                  | ribonucleic acid   |
| rpm                  | revolutions per minute                                   |
| RT-PCR               | reverse transcriptase polymerase chain reaction          |
| S                    | DNA synthesis phase                                      |
| <i>S. cerevisiae</i> | <i>Saccharomyces cerevisiae</i>                          |
| SDS                  | sodium dodecyl sulphate                                  |
| SE                   | standard error   |
| <i>S. pombe</i>      | <i>Schizosaccharomyces pombe</i>                         |
| T                    | threonine  |
| TBY-2                | tobacco bright yellow var. 2 cell line                   |
| Thr                  | threonine  |
| Tris                 | 2-amino-2-hydroxymethyl-1-3-aminomethane                 |
| Tris-HCl             | 2-amino -2-hydroxymethyl-1-3- aminomethane hydrochloride |
| Tyr                  | tyrosine   |
| UHP                  | ultra-high purity water                                  |
| WT                   | wild type  |
| Xgal                 | 5-bromo-4-chloro-3-indoyl- $\beta$ -D- galactopyranoside |
| Y                    | tyrosine   |

## **Abstract**

In eukaryotes the regulatory cell cycle gene, *WEE1*, encodes a protein kinase. In late G2, it inactivates cyclin-dependent kinase (CDKs) in the CDK-cyclinA/B complexes, by phosphorylating the CDK on tyrosine 15. This can result in a delay in mitosis. Expression of *Arabidopsis thaliana* homologue of *WEE1* (*AtWEE1*) in fission yeast resulted in an elongated cell length phenotype in the same way as over expression of fission yeast *wee1*. I have tested whether *AtWEE1* could also induce this effect in tobacco cells and *Arabidopsis* plant roots.

The tobacco BY-2 cells have been transformed with *AtWEE1*, both under constitutive and inducible promoters. Phenotypic characteristics observed compared with the control are premature entry into mitosis and a reduced cell size through a shortening of the G2 phase with a compensatory increase in the duration of G1 phase. Hence, the phenotype and cell cycle response is the exact opposite of the known effect of expression of this gene in fission yeast. *NtWEE1* expression data revealed that the endogenous *WEE1* expression is delayed in transgenic lines, this results in a non-inhibition of CDKA and CDKB1 which are already active in early S-phase.

*AtWEE1* was also employed to transform *Arabidopsis thaliana* plants, both under constitutive and inducible promoters. The effect of *AtWEE1* over expression was investigated on primary root growth and lateral root development. In particular, *AtWEE1* over expression lead to less primary root growth and a reduction in the frequency of lateral root primordia initiated when compared with wild type. *Arabidopsis* transgenic plants initiated fewer primordia both per unit time and per cm of primary root.

# **Chapter 1: General introduction**


## **1.1. The eukaryotic cell cycle and its regulation**

One of the key events in the life history of a cell is its division into two identical daughter cells, accomplished during a phase in its life known as mitosis (M phase). In proliferative cells interphase is a preparation for mitosis. The two periods, mitosis and interphase, constitute the well-organized sequence of events known as the cell cycle (Fig. 1.1).

The mitotic cell cycle consists of alternating rounds of DNA replication (which occurs during the S phase) and chromatid separation (which occurs during the M phase) interrupted by gaps known as G1 (the interval before S phase) and G2 (the interval after S phase) (Howard and Pelc, 1953; Taylor, 1958). G1 and G2 may be alternatively considered as periods of general cell growth: G1 represents a phase in which there is a great increase in the rate at which new components (except DNA) are made and G2 represents a period during which a small amount of further growth takes place. Events that occur in each phase are regulated to ensure that the DNA is replicated only once in each cell cycle and that each daughter cell arrives at G2/M with a complete complement of the genome (Eckardt, 2001).

Basic features of cell cycle control are remarkably conserved in all eukaryotes and principal control points at the G1/S and G2/M have been identified in yeast, animals, and plants (Van't Hof, 1966, 1974; Pines, 1995; Huntley and Murray, 1999). Progression through the successive phases of the cell cycle (S, G2, M and G1) in species as diverse as

yeast and humans is driven by a common class of heterodimeric protein kinases. These heterodimers consist of a cyclin-dependent kinase (CDK) and a non-catalytic cyclin (Evans *et al.*, 1983; Nigg, 1995).



**Fig. 1.1:** The cell cycle and its component phases.

Studies of the cell cycle present complicated and confusing nomenclature. This is largely because fission yeast geneticists invented a unique nomenclature whereby the gene is



italic lower case while the protein is normal font but denoted by a capital letter beginning the description. The rest of the cell cycle community uses the conventional upper case italic for the gene and upper case normal font for the protein (i.e. Table 1.1).

### 1.1.1. Cyclin-dependent kinase (CDK)

In fission yeast (*Schizosaccharomyces pombe*) a single CDK, Cdc2 regulates the transition from G1 to S, and G2 to M. The budding yeast (*Saccharomyces cerevisiae*) homologue, *CDC28* is expressed in G1 and drives the cell into S-phase. Because of the poorly defined G2, in budding yeast, it is accepted that *CDC28* expression commits the cell to both S-phase and mitosis (Hartwell *et al.*, 1974; Nurse, 1975; Nasmyth, 1993) (see section 1.3.1. for more details).

In animals, several distinct CDKs have been isolated and function at different stages in the cell cycle (Morgan, 1997). The presence of multiple types of CDKs in higher eukaryotes may reflect the increased regulation necessary to carry out the more complex mitogenic instructions that are required during multicellular development (see section 1.3.2. for more details).

CDKs are not conserved between animals and plants and have variant sequences in their cyclin-binding domain. In animals the nomenclature CDK1 to CDK7 has been adopted (Pines, 1995), whereas in plants the lack of direct equivalents (except between CDK1 and CDKA) has led to the adoption of an alphabetical suffix (CDKA to CDKF) (Joubes *et al.*, 2000), which replaces an earlier confused nomenclature based on *cdc2* (Table 1.1).

At present, 46 CDKs have been identified in 23 different plant species that can be grouped into seven types and can be divided into two main different classes on the basis of their sequences (Segers *et al.*, 1997; Joubes *et al.*, 2000; Stals *et al.*, 2000). All of them possess a specific motif required for ATP and cyclin binding, as first described for the human CyclinA/CDK2 (Jeffrey *et al.*, 1995), as well as conserved sites of regulatory phosphorylation (Segers *et al.*, 1998).

The best characterized plant CDKs belong to the type A class. This class comprises kinases most closely related to the prototypical CDKs, yeast *cdc2/CDC28* and animal CDK1 and CDK2, which share the conserved PSTAIRE (single letter code for amino acid) motif in the cyclin binding domain (Serges *et al.*, 1998; Huntley and Murray, 1999; Mironov *et al.*, 1999). These CDKs are also known as CDKA type (Table 1.1). In addition to this large group of CDKs, several non-PSTAIRE CDKs have been identified in plants, which are characterized by the variant sequences PPTALRE or PPTTLRE, and are known as CDKB types (Table 1.1) (Serges *et al.*, 1998; Huntley and Murray, 1999; Mironov *et al.*, 1999).

**Table 1.1:** Classification of CDKs in Plants based on their cyclin binding motif

| Types of CDKs | Cyclin binding motif | New gene name         | Old gene name | Species                        | Characteristics            |
|---------------|----------------------|-----------------------|---------------|--------------------------------|----------------------------|
| CDKA          | PSTAIRE              | Allce; <i>CDKA;1</i>  | Cdc2          | <i>Allium cepa</i>             | Involved in G1/S and G2/M. |
|               |                      | Antma; <i>CDKA;1</i>  | Amdc2a        | <i>Antirrhinum majus</i>       |                            |
|               |                      | Antma; <i>CDKA;2</i>  | Amdc2b        | <i>Antirrhinum majus</i>       |                            |
|               |                      | Arath; <i>CDKA;1</i>  | Atcdc2a       | <i>Arabidopsis thaliana</i>    |                            |
|               |                      | Betvu; <i>CDKA;1</i>  | cdc2          | <i>Beta vulgaris</i>           |                            |
|               |                      | Brana; <i>CDKA;1</i>  | cdc2          | <i>Brassica napus</i>          |                            |
|               |                      | Cheru; <i>CDKA;1</i>  | cdc2          | <i>Chenopodium rubrum</i>      |                            |
|               |                      | Glyma; <i>CDKA;1</i>  | cdc2-S6       | <i>Glycine max</i>             |                            |
|               |                      | Glyma; <i>CDKA;2</i>  | cdc2-S5       | <i>Glycine max</i>             |                            |
|               |                      | Lycles; <i>CDKA;1</i> | Lecdc2A-1     | <i>Lycopersicon esculentum</i> |                            |
|               |                      | Lycles; <i>CDKA;2</i> | Lecdc2A-2     | <i>Lycopersicon esculentum</i> |                            |
|               |                      | Medsa; <i>CDKA;1</i>  | cdc2MsA       | <i>Medicago sativa</i>         |                            |
|               |                      | Medsa; <i>CDKA;2</i>  | cdc2MsB       | <i>Medicago sativa</i>         |                            |
|               |                      | Nicta; <i>CDKA;1</i>  | Ntcdc2-1      | <i>Nicotiana tabacum</i>       |                            |
|               |                      | Nicta; <i>CDKA;2</i>  | Ntcdc2-2      | <i>Nicotiana tabacum</i>       |                            |
|               |                      | Nicta; <i>CDKA;3</i>  | cdc2Nt1       | <i>Nicotiana tabacum</i>       |                            |
|               |                      | Orysa; <i>CDKA;1</i>  | cdc2Os-1      | <i>Oryza sativa</i>            |                            |
|               |                      | Orysa; <i>CDKA;2</i>  | cdc2Os-2      | <i>Oryza sativa</i>            |                            |
|               |                      | Peter; <i>CDKA;1</i>  | cdc2          | <i>Petroselinum crispum</i>    |                            |
|               |                      | Pethy; <i>CDKA;1</i>  | cdc2          | <i>Petunia hybrida</i>         |                            |
|               |                      | Picab; <i>CDKA;1</i>  | cdc2Pa        | <i>Picea abies</i>             |                            |
|               |                      | Pinco; <i>CDKA;1</i>  | cdc2Pnc       | <i>Pinus contorta</i>          |                            |
|               |                      | Pissa; <i>CDKA;2</i>  | cdc2          | <i>Pisum sativum</i>           |                            |
|               |                      | Soltu; <i>CDKA;2</i>  | cdc2          | <i>Solanum tuberosum</i>       |                            |
|               |                      | Sesro; <i>CDKA;1</i>  | Srcdc21       | <i>Sesbania rostrata</i>       |                            |
|               |                      | Triae; <i>CDKA;1</i>  | cdc2TaA       | <i>Triticum aestivum</i>       |                            |
|               |                      | Triae; <i>CDKA;2</i>  | cdc2TaB       | <i>Triticum aestivum</i>       |                            |
|               |                      | Vigac; <i>CDKA;1</i>  | cdc2          | <i>Vigna acontifolia</i>       |                            |
|               |                      | Vigum; <i>CDKA;1</i>  | cdc2          | <i>Vigna unguiculata</i>       |                            |
|               |                      | Zeama; <i>CDKA;1</i>  | cdc2A         | <i>Zea mays</i>                |                            |
|               |                      | Zeama; <i>CDKA;2</i>  | cdc2B         | <i>Zea mays</i>                |                            |

|      |                     |   |  |  |                             |
|------|---------------------|---|--|--|-----------------------------|
| CDKB | PPTALRE;<br>PPTTLRE | Atma; <i>CDKB1;1</i><br>Atma; <i>CDKB2;1</i><br>Arath; <i>CDKB1;1</i><br>Dunte; <i>CDKB;1</i><br>Medsa; <i>CDKB1;1</i><br>Arath; <i>CDKB2;1</i><br>Medsa; <i>CDKB2;1</i><br>Meser; <i>CDKB2;1</i><br><br>Orysa; <i>CDKB;1</i> | Amdc2c<br>Amdc2d<br>Atcdc2b<br>cdc2<br>cdc2MsD<br>Atcdc2d<br>cdc2MsF<br>cdc2<br><br>cdc2Os-3 | <i>Antirrhinum majus</i><br><i>Antirrhinum majus</i><br><i>Arabidopsis thaliana</i><br><i>Dunaliella tertiolecta</i><br><i>Medicago sativa</i><br><i>Arabidopsis thaliana</i><br><i>Medicago sativa</i><br><i>Mesembryanthemum crystallinum</i><br><i>Oryza sativa</i> | Involved in G2/M.           |
| CDKC | PITAIRE             | Arath; <i>CDKC;1</i><br>Arath; <i>CDKC;2</i><br>Medsa; <i>CDKC;1</i><br>Pissa; <i>CDKC;1</i>  | cdc2At<br>cdc2At<br>cdc2MsC<br>Ps2cdc2   | <i>Arabidopsis thaliana</i><br><i>Arabidopsis thaliana</i><br><i>Medicago sativa</i><br><i>Pisum sativum</i>   | Function is not understood. |
| CDKD | N(IF)TALRE          | Arath; <i>CDKD;1</i><br>Orysa; <i>CDKD;1</i>  | CAK2At<br>R2   | <i>Arabidopsis thaliana</i><br><i>Oryza sativa</i>   | Involved in G1.             |
| CDKE | SPTAIRE             | Medsa; <i>CDKE;1</i><br>Arth; <i>CDKE;1</i>   | cdc2MsE  | <i>Medicago sativa</i><br><i>Arabidopsis thaliana</i>  | Function is not understood. |
| CDKF |                     | Arth; <i>CDKF;1</i>   |  | <i>Arabidopsis thaliana</i>  | Function is not understood. |
| CDKG | PLTSLRE             | Arth; <i>CDKG;1</i>   |  | <i>Arabidopsis thaliana</i>  | Function is not understood. |

The CDKB proteins fall into two subgroups on the basis of sequence relationships (Huntley and Murray, 1999; Umeda *et al.*, 1999; Joubes *et al.*, 2000) (Table 1.1). One group contains *Arabidopsis cdc2b*, *Antirrhinum majus cdc2c* and *Medicago sativa cdc2MsD*, which all contain the sequence PPTALRE and for which the name CDKB1 subgroup has been proposed (Hirayama *et al.*, 1991; Imajuku *et al.*, 1992; Hirt *et al.*, 1993; Forbert *et al.*, 1996; Segers *et al.*, 1996; Magyar *et al.*, 1997; Huntley and Murray, 1999). The other subgroup named CDKB2 contains *Antirrhinum majus cdc2d*, *Medicago sativa cdc2MsF*, *Oryza sativa cdc2Os3* and *Arabidopsis cdc2dAt* (Hirt *et al.*, 1993; Kidou *et al.*, 1994; Forbert *et al.*, 1996; Magyar *et al.*, 1997; Umeda *et al.*, 1999; Huntley

and Murray, 1999). The genes from these two groups differ slightly in the timing of their expression during the cell cycle. CDKB1 transcripts accumulate during S, G2 and M phases, whereas CDKB2 expression is specific to G2 and M phase (Fobert *et al.*, 1996; Segers *et al.*, 1996; Magyar *et al.*, 1997; Umeda *et al.*, 1999). The uniqueness of this CDK has led to the view that CDKB is the kinase that drives plant cells into mitosis. Part of my thesis sheds new light on the timing of Nicta; CDKB1 enzyme activity.

A small group of four plant CDKs was characterized by the presence of the PITAIRE motif (Table 1.1). However, Mironov *et al.* (1999) argued against the involvement of Arath; CDKC in cell cycle control since an *in situ* hybridization signal for an *Arabidopsis* member of the CDKC family could not be obtained in actively dividing cells.

The CDKD proteins (Table 1.1) have a conserved N(I/F)TALRE motif closely related to the equivalent motif of CDK7 kinases from animals. CDKD proteins are considered to be bifunctional, involved in phosphorylation-dependent activation of other CDKs during the cell cycle, and in phosphorylation-dependent regulation of the activity of RNA polymerase II (Harper and Elledge, 1998). In synchronized suspension cells (Sauter, 1997), and during adventitious root growth (Lorbiecke and Sauter, 1999), a preferential expression of the rice *CDKD;1* gene was recorded in G1 and S phases.

Finally, the alfalfa Medsa; *CDKE;1* and Arth; *CDKE;1*, appear unrelated to any other plant sequence as they have a SPTAIRE motif; this CDK was named CDKE (Table 1.1). The involvement of CDKE genes in the plant cell cycle has yet to be proven because they display at the mRNA level a weak constitutive signal during a synchronized cell cycle (Magyar *et al.*, 1997). *Arabidopsis* CDKE might play a role in the specification of stamen and carpel identities and for the proper termination of stem cells in the floral meristem. It

might have a function similar to the differentiation function played by CDK8 in mammals (Wang and Chen, 2004).

In Arabidopsis, *Arath;CDKF;1* encodes a CAKAK (Cyclin dependent kinase activating kinase) (Shimotohno *et al.*, 2004).

CDKG is a relatively new-comer and has a PLTSLRE motif and shows homology to the human protein kinase p58/GTA, a member of the CDC2 kinase subfamily (Menges *et al.* 2005).

CDK activity is dependent on binding with a partner cyclin (see section 1.1.2 for more details) which determines the substrate specificity and the subcellular localization of the CDK complex (Pines, 1995). However, it also depend on a CDK-Activating Kinases (CAK) (see section 1.1.3 for more details) and on phosphorylation of T14 and Y15 residues of the CDK protein (Kugmagai and Dunphy, 1991) (see section 1.4 for more details). Negative regulation by cyclin-dependent kinase inhibitors (CKIs) deactivates the CDK complex (see section 1.1.4 for more details).

### **1.1.2. Cyclins**

Cyclins are a diverse group of proteins with low overall homology that share a large, rather poorly conserved region responsible for their interaction with the CDK; this region is referred to as the cyclin core (Evans *et al.*, 1983). The cyclin core covers about 250 amino acid residues and is organized in two folds of five helices. The first fold is the cyclin box and comprises about 100 amino acid residues (Noble *et al.*, 1997), representing the region of highest conservation, although it contains only five absolutely

invariant positions. The crystal structure reveals the cyclin box as the face of interaction with the cognate CDK (Jeffrey *et al.*, 1995).

In budding yeast seven different cyclins (CLN and CLB proteins) were identified to play important roles at specific phases of the cell cycle (Nasmyth, 1993; Stuart and Wittenberg, 1995) (Table 1.2) (see section 1.2.1 for more details).

**Table 1.2:** Classifications of Cyclins in Budding Yeast

| Cyclins | Characteristics                     |
|---------|-------------------------------------|
| Cln3    | Expressed from early G1 phase       |
| Cln1    | Expressed from G1 to beginning of S |
| Cln2    | Expressed from G1 to beginning of S |
| Cln5    | Expressed from S to G2 phase        |
| Cln6    | Expressed from S to G2 phase        |
| Clb1    | Expressed in M phase                |
| Clb2    | Expressed in M phase                |

In fission yeast six different cyclins (Cig 1 – 6) operate in a similar way (Martin-Castellanos *et al.*, 1996).

A classification based on sequence organization indicates that five types of cyclins exist in plants: A, B, C, D and H types. The first four classes of cyclins (A, B, C and D) are divided into subgroups (Table 1.3) (Renaudin *et al.*, 1996; Vandepoele *et al.*, 2002).

**Table 1.3:** Classifications of Cyclins in Plants

| Cyclins | Species  | Characteristics                     |
|---------|--|-------------------------------------|
| CycA1   | <i>Z. mays</i> ;<br><i>N. tabacum</i> ;<br><i>O. sativa</i> .                          | Expressed from S to M phase.        |
| CycA2   | <i>A. thaliana</i> ;<br><i>Z. mays</i> ;<br><i>N. tabacum</i> ;<br><i>M. sativa</i> .  | Expressed from S to M phase.        |
| CycA3   | <i>C. roseus</i> ;<br><i>A. thaliana</i> ;<br><i>N. tabacum</i> ;<br><i>A. majus</i> . | Expressed from S to early G2 phase. |
| CycB1   | <i>A. thaliana</i> .   | Expressed from G2 to M phase.       |
| CycB2   | <i>M. sativa</i> .   | Expressed from G2 to M phase.       |
| CycB3   | <i>A. thaliana</i>   | Expressed from G2 to M phase.       |
| CycD1   | <i>A. thaliana</i> ;<br><i>A. majus</i> ;<br><i>H. tuberosus</i> .                     | Unknown.                            |
| CycD2   | <i>A. thaliana</i> ;<br><i>N. tabacum</i> .  | Expressed from G1 to S phase.       |
| CycD3   | Widely identified.   | Expressed from G1 to S phase.       |
| CycD4   | <i>A. thaliana</i> .   | Unknown.                            |
| CycH    | <i>A. thaliana</i><br><i>O. sativa</i>   | Unknown.                            |

In plants, the major classes of cyclins have homology to animal groups, including the plant cyclins A, B and D (indicated in the Table 1.3 as CycA, CycB and CycD), and numerous examples continue to be isolated and analyzed (Hsieh and Wolniak, 1998; Day and Reddy, 1998; Nakagami *et al.*, 1999; Sorrell *et al.*, 1999).

In mammals, a single type of cyclin A is sufficient to promote CDK activity in S and G2 phase. The CycA-CDK2 complex plays a role in DNA replication and in transcriptional



regulation during S phase. In contrast, multiple A-type cyclins exist in plants. They can be grouped into three subgroups: CycA1, CycA2 and CycA3 (Table 1.3), according to sequence (Renaudin *et al.*, 1996; Chaubet-Gigot, 2000). The functional significance of their complex expression patterns remains unclear (Bursdens *et al.*, 2000; Roudier *et al.*, 2000). In *Arabidopsis* 10 sequences (*CYCA1;1*, *CYCA1;2*, *CYCA2;1*, *CYCA2;2*, *CYCA2;3*, *CYCA2;4*, *CYCA3;1*, *CYCA3;2*, *CYCA3;3*, *CYCA3;4*) encoding A-type cyclins have been identified (Vandepole *et al.*, 2002). In most plants screened, members of all three subclasses of A-type cyclins have been found (Dewitte and Murray, 2003).

B-type cyclins (Table 1.3) are expressed specifically in late G2 and early M phase of the cell cycle (Ito, 2000). B type cyclins are distinguished from A-type cyclins, not only by their sequence differences, but also by their later expression pattern during the cell cycle. All identified B-type cyclins were subdivided into two subclasses, CycB1 and CycB2 (Renaudin *et al.*, 1996). However, recently a further B-type cyclin gene was discovered in the *Arabidopsis* genome; it encodes for a B-type cyclin-like protein without the typical B-type destruction box, and has been assigned to a third class, CycB3 (Vandepole *et al.*, 2002).

D-type cyclins (Table 1.3) were defined on the basis of a low sequence homology to animal D-type cyclins and the presence of the conserved LxCxE amino acid motif (single letter amino acid code, where x represents any residue), which is responsible for their interaction with retinoblastoma proteins (Rb) (Soni *et al.*, 1995). The LxCxE amino acid motif is conserved in both animals and plants and is located near the N-terminus of the protein (Ach *et al.*, 1997; Huntley *et al.*, 1998). D-type cyclins have been identified in a variety of plant species (Meijer and Murray, 2000). The recently completed *Arabidopsis*

genome sequence reveals the presence of 10 D-type cyclins sequences as well as several more distantly related cyclin-like genes (Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*, 2000; Vandepole *et al.*, 2002).

On the basis of phylogenetic analysis, four groups of D-type cyclins exist in plants: CycD1, CycD2, CycD3 and CycD4 (Renaudin *et al.*, 1996; Meijer and Murray, 2000; Vandepole *et al.*, 2002, Boucheron *et al.*, 2005). Multiple pathways control D-type cyclin activity both transcriptionally and post-transcriptionally. *CycD3* transcription is regulated by cytokinins, but brassinosteroids also target *CycD3* expression (see section 1.2.3 Table 1.4 for more details) (Riou-Khamilichi *et al.*, 1999; Hu *et al.*, 2000). Moreover, sucrose has been shown to differentially regulate *Arabidopsis* D-type cyclin (CycD2 and CycD3) expression (Riou-Khamilichi *et al.*, 2000). The complex regulation of D-type cyclins further strengthens the notion that these molecules play a role throughout the plant cell cycle and are not only restricted to controlling the G1/S transition (Riou-Khamilichi *et al.*, 2000).

Genes encoding H-type cyclins (Table 1.3) have been identified in *Arabidopsis* and rice, but it is still unknown whether these proteins have a regulatory role in the plant cell cycle (Yamaguchi *et al.*, 2000; Vandepole *et al.*, 2002).

*CycA* and *CycB* cyclin genes are expressed in a cell cycle dependent manner, peaking transcriptionally at or near to the G2/M transition (A type cyclins are expressed somewhat earlier than B-type), and differences in the timing of expression exist even between genes encoding different sub-types of *CycA* cyclins (Reicheld *et al.*, 1996). *CycD* cyclin genes like their animal counterparts, show cell cycle independent expression, their transcription being induced by the presence of mitogens. The genes

encoding CycD cyclins are induced at specific times during cell cycle re-entry but generally remain expressed at a constant level in actively dividing cells (Soni *et al.*, 1995; Fuerst *et al.*, 1996; Sorrell *et al.*, 1999). In animals, E-type cyclins are strongly regulated transcriptionally at the G1/S boundary, but a direct equivalent has yet to be identified in plant cells (Dewitte and Murray, 2003).


The levels of cyclins are generally determined by specific protein-turnover mechanisms as well as highly regulated transcription (Ito *et al.*, 1998). The destruction of CycA and CycB cyclins at specific points during M phase depends on a destruction box motif which mediates their ubiquitin-dependent proteolysis (Glotzer *et al.*, 1991; Renaudin *et al.*, 1998). In contrast, most CycD cyclins contain PEST sequences, regions rich in proline, glutamate, serine and threonine, which are thought to be a signal for the rapid proteolysis of many proteins (Rechsteiner, 1990; Soni *et al.*, 1995; Sorrell *et al.*, 1999).

### **1.1.3. CDK-Activating Kinases**

The basic mechanism and logic of cell cycle control, with some exceptions, is highly conserved in all eukaryotes, and so are the key genes that mediate cell cycle progression (Nasmyth, 1996; Novak *et al.*, 1998; Dewitte and Murray, 2003). Cyclin dependent kinases play a central role in mediating cell cycle progression (see section 1.1.2 for more details). CDK activity is regulated by association with cyclin subunits, reversible phosphorylation and association with other regulatory proteins. CDK activity is controlled by phosphorylation at three conserved sites, and many of the enzymes that act on these sites have been identified in many eukaryote organisms including plant

species (Lew and Kornbluth, 1996; Dewitte and Murray, 2003; Sorrell *et al.*, 2002, 2003). Cyclin dependent kinase activity not only requires binding of the CDK to a cyclin but also, phosphorylation of the CDK at a conserved threonine residue within the “T-loop” (residue Thr-160 or Thr-161). The T-loop can mask the catalytic site to prevent substrate binding, or swing open to permit substrate phosphorylation (Solomon, 1993).

CDK-activating kinase (CAK), the enzyme responsible for the phosphorylation of the Thr-160 or 161 residues of CDKs (Fig. 1.2), has been identified in several organisms.



**Fig. 1.2:** CAK activates CDK by phosphorylation of threonine 160 or 161.

The most plausible candidate for the higher eukaryotic CAK is the CDK7/cyclinH/Mat1 complex, which was purified from frog, starfish, and mammalian cells (Nigg, 1996). An *Arabidopsis* cDNA, designated *CAK1Arath*, was isolated as a suppressor of a CAK mutation in budding yeast and fission yeast (Umeda *et al.*, 1998). The amino acid sequence of ArathCAK1 is related to animal CAKs, but similarities are restricted to the conserved kinase domain. ArathCAK1 can phosphorylate human CDK2 at the Thr-160

residue in the T loop (Umeda *et al.*, 1998). The rice R2, closely related to CDK7 of animals (Hata *et al.*, 1991), is structurally similar to the CAKs of metazoans and fission yeast, but is distinct from ArathCAK1 (Yamaguchi *et al.*, 1998).

In Arabidopsis, CAK comprises CyclinD and CDKD although CDKF might function equally well here. Note that in alfalfa, CDK;C1-cyclin T phosphorylates RNA polymerase II at its C-terminal (Fulop *et al.*, 2005).

In higher eukaryotes, the activated cyclin-bound CDK can be inhibited by phosphorylation of two conserved residues within the catalytic cleft at residues Y15 and T14 (Lew and Kornbluth, 1996). However, in fission yeast only Y15 is phosphoregulated (Russell and Nurse, 1987) (see section 1.4 for more details). The catalytic activity of CDK/cyclin complex can also be negatively regulated by interaction with CDK inhibitor (CKIs), which cause extensive structural changes through binding (Martin-Castellanos and Moreno, 1997) (see section 1.1.5 for more details).

#### **1.1.4. Cyclin-dependent kinase inhibitors (CKIs)**

CKIs are low molecular mass proteins that bind and negatively regulate the catalytic activity of the CDK/Cyclin complex by causing extensive structural changes through their binding (Martin-Castellanos and Moreno, 1997). In mammals, several inhibitors have been isolated, grouped into the Ink4 or the Cip/Kip1 family, according to functional and sequence similarity (Sherr and Roberts, 1995; Burssens *et al.*, 1998). In fission yeast and budding yeast cells, CDK inhibitors prevent premature initiation of DNA replication by inhibiting the CDK/Cyclin complex in early G1 phase (Sanchez-Diaz


*et al.*, 1998). In *S. cerevisiae* degradation of the CDK inhibitor is required for initiation of S phase (Schwob *et al.*, 1994). Plant CDK/cyclin complex inhibitory proteins are known as ICK (inhibitor of Cdc2 kinase) or KRP (Kip-related proteins) and bind both CDK and cyclin subunits (Wang *et al.*, 1997, Zhou *et al.*, 2003, Verkest *et al.*, 2005). KRPs can inhibit both CYCD2/CDKA and CYCD2/CDKB (Nakai *et al.*, 2006).

Using the yeast two hybrid system, the first plant CDK inhibitor gene was identified in *A. thaliana* (Wang *et al.*, 1997, Weinl *et al.*, 2005), where the CKI genes constitute a small gene family (Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*, 2000). The plant CKI protein restricts CDK activity *in vitro* and *in vivo* (Wang *et al.*, 1998, 2000).

## **1.2. Comparison between the CDK/cyclin complexes during the different phases of the cell cycle in yeast, animals and plants**

### **1.2.1. Budding yeast cell cycle**

In early interphase (G1) of the budding yeast cell cycle, cells rapidly increase in volume. This cell growth is accompanied by a gradual increase of G1-phase Cln3 cyclin (a “non-cycling” cyclin) that promotes the accumulation of cyclins Cln1 and Cln2 at the transcriptional level (Nasmyth, 1993; Stuart and Wittenberg, 1995) (Fig. 1.3 a). These associate with CDC28 kinase, the budding yeast homologue to Cdc2, to regulate the transition through the G1 checkpoint known as “START” (Nasmyth, 1993) (Fig. 1.3 b).




**Fig.1.3:** Schematic model of budding yeast cell cycle which shows interactions between CDC28 and different cyclins during G<sub>1</sub>, S, G<sub>2</sub> and M phase.

Phosphorylation of Cln1 and Cln2 at G1 phase targets them for ubiquitin-mediated proteolysis (Deshaies *et al.*, 1995; Kirshi and Yamao, 1998) (Fig. 1.3 c). Proteolysis of Cln1 and Cln2 enables budding yeast cells to produce the S-phase cyclins Cln5 and Cln6 and to enter S-phase (Fig. 1.3 d). Synthesis of the mitotic cyclins Clb1 and Clb2 is initiated in late G2 phase, these cyclins in association with CDC28 stimulate entry into mitosis (Nasmyth, 1993; Stuart and Wittenberg, 1995) (Fig. 1.3 e).

### **1.2.2. Animal cell cycle**

Quiescent animal cells are stimulated to divide by growth factors like serum that induces passage from G0 into G1 in a process requiring D-type cyclins, whose transcription is dependent on the presence of serum (Kohn, 1999). Growth factors, also lead to the accumulation of D-type and E-type cyclins in G1 phase (Fig. 1.4 a), which associate with CDKs (Kohn, 1999) (Fig. 1.4 b).





**Fig. 1.4:** Schematic model of animal cell cycle which shows interactions between different CDK and cyclins during G<sub>1</sub>, S, G<sub>2</sub> and M phase.

The progression from G1 phase to S phase involves CDK4-CDK6/CycD and CDK2/CycE (Kohn, 1999) (Fig. 1.6 b). CDK2 is also implicated in regulating S phase, when it is associated with CycA (Kohn, 1999) (Fig. 1.4 c). In G2 phase, CDK2 is associated with CycB (Kohn, 1999) (Fig. 1.4 d). In M phase, both CDK1 and CDK2 are associated with CycB: CDK1/CycB and CDK2/CycB complexes (Kohn, 1999) (Fig. 1.4 e). In telophase, mitotic cyclin is degraded by the APC (Anaphase Promoting Factor) pathway, which resets the cell cycle to G1 phase (Kohn, 1999) (Fig. 1.4 f).

### 1.2.3. Plant cell cycle


In plant cells, homologues to CDKs, cyclins, ICKs, CAKs have been isolated (Lew and Kornbluth, 1996; Renaudin *et al.*, 1996; Wang *et al.*, 1997; Huntley and Murray, 1999; Joubes *et al.*, 2000; Vandepoele *et al.*, 2002; Dewitte and Murray, 2003; Zhou *et al.*, 2003, Beemster *et al.*, 2005). These plant genes are regulated by plant growth regulator (PGRs) signals (Table 1.4). However, these mechanisms are completely different from animal (and yeast) growth factors, which induce calcium signals that control cell proliferation (Berridge, 1995). Differences in growth factors between animals and plants are consistent with evolutionary divergence of the plant genes from the animal ones. Moreover, the uniqueness of plant cytokinesis argues for evolutionary divergence of plant genes that are necessary for cytokinesis.

**Table 1.4:** Effects of exogenous PGRs and sucrose on the plant cell cycle

| PGRs   | Characteristics   | Published examples   |
|--|---|--|
| Cytokinins (i.e. kinetin, zeatin, benzyladenine) | Promote plant cell division, can cause non-cycling cells to divide. Can act at the G1/S transition by stimulating CycD3 and at the G2/M transition by stimulating activation of b-type CDK. | Riou-Kamlichi <i>et al.</i> , 1999.<br>Zhang <i>et al.</i> , 2005.<br>Menges <i>et al.</i> , 2006. |
| Brassinolides                                    | Activate cell proliferation.  | Miyazawa <i>et al.</i> , 2003.   |
| Gibberellins (GAs)                               | Induce cell cycle activation at the G2/M transition by increasing the level of CDK kinase.  | Sauter <i>et al.</i> , 1997.   |
| Auxin  | Required to initiate cell division in the pericycle by acting on D-type cyclin.   | De Veylder <i>et al.</i> , 1999.   |
| Abscisic acid (ABA)                              | Prevents DNA replication by keeping the cells in the G1 phase inducing ICK1 activity.   | Wang <i>et al.</i> , 1998;<br>Swiatek <i>et al.</i> , 2002.  |
| Ethylene   | Delays the entry of cells into mitosis and induces cell death at the G2/M transition.   | Herbert <i>et al.</i> , 2001.  |
| Sucrose  | Stimulates G0 to G1 progression by inducing CycD2. Enables cells arrested in G1 to enter S phase or cells arrested in G2 to enter M phase.  | Soni <i>et al.</i> , 1995;<br>Sorrell <i>et al.</i> , 1999.  |

Cytokinins, brassinolides and sucrose are molecules that regulate plant cell division. In G1 phase CycD, the partner of CDKA, is rapidly induced in response to the accumulation of the above regulators (Fig. 1.5 a). During S-phase, CycA begins to accumulate as a partner for CDKA (Fig. 1.5 b). In G2 phase and in the transition from G2 to M phase, CycA is replaced by CycB as partner of both CDKA and CDKB (Fig. 1.5 c).

### 1.3. Retinoblastoma protein and the E2F transcription factor regulate



**Fig. 1.5:** Schematic model of plant cell cycle. Various cell cycle regulatory proteins are degraded through the ubiquitin/proteasome pathway at specific cell cycle stages in plant cells.


### 1.3. Retinoblastoma protein and the E2F transcription factor regulate G1/S transition

A major control of the G1/S transition in animals and plants is exerted by the retinoblastoma protein (Rb). Progression into S phase requires the activity of E2F transcription factors. In mammals, six E2Fs have been identified and classified into three groups: the activating E2Fs (E2F1, E2F2 and E2F3), the repressive E2Fs (E2F4 and E2F5) and E2F6 (Trimarchi and Lees, 2002). In *Arabidopsis* three E2F genes have been identified (*E2Fa*, *E2Fb* and *E2Fc*) (Vandepoele *et al.*, 2002; He *et al.*, 2004).

During G1 phase, E2F is bound to the Retinoblastoma (Rb) protein, which recruits histone deacetylases to the promoters of E2F-regulated genes thereby switching off their transcription (Fig. 1.6 a). Phosphorylation of Rb by CDKA/cyclinD complex kinase results in hyperphosphorylation of Rb and release of E2F that can activate E2F-responsive genes (Huntley and Murray, 1999) (Fig. 1.6 b). Later in S phase, the CDKB-cyclinA complex phosphorylates E2F proteins, abolishing their ability to bind DNA (Fig. 1.6 c) (Huntley and Murray, 1999).

In plants Rb protein regulates cell proliferation, differentiation and endoreduplication (Park *et al.*, 2005). In tobacco, Rb binds and suppresses E2F but Cyclin D is part of a mechanism that releases E2F (Uemukai *et al.*, 2005). Several plant CYCD genes might bind with Arath; CDKA and replace animal cyclin E in hyperphosphorylation of Rbs. (Menges *et al.*, 2005).





**Fig. 1.6:** Schematic model for the control of G1/S transition in plants, involving Retinoblastoma protein (Rb) and E2F transcription factor (adapted from Huntley and Murray, 1999).


## 1.4. Cell-cycle regulation at the G2/M transition in fission yeast, animals and plants

In all eukaryotes, transition from G2 to M phase is regulated by phosphorylation/dephosphorylation of CDK at either tyrosine 15 or threonine 14 and tyrosine 15. In fission yeast, phosphorylation of Cdc2 at tyrosine15 is mediated by the related kinases Wee1 and Mik1 (Russell and Nurse, 1987; Millar *et al.*, 1992; Parker *et al.*, 1992; Atherton-Fessler *et al.*, 1993; Dunphy, 1994). The first of these to be identified was Wee1 from *Schizosaccharomyces pombe*, which causes a delay in mitosis by phosphorylating G2/M CDKs (Russell and Nurse, 1987), whereas the phosphatases Cdc25, Pyp3 and Stp1 dephosphorylate Y15 releasing cells into mitosis (Lew and Kornbluth, 1996) (Fig. 1.7 a). Wee1 and Cdc25 are primarily responsible for regulation of Y15 phosphorylation, whereas Mik1 and Pyp3 are minor partners whose role becomes apparent only when the major partners are compromised (Lew and Kornbluth, 1996). However, Stp1 was not completely linked to cell cycle control. In fission yeast, a G2 phase size control operates under rapid growth conditions, such that the onset of mitosis is dependent on reaching a critical cell size. Cell size at mitosis is likely to be dependent on *wee1* and *cdc25* gene dosage; as little as a threefold increase in gene dosage, has significant effects on the timing of mitosis (Wee1 delays mitosis, whereas Cdc25 advances mitosis) (Russell and Nurse, 1986, 1987). These observations demonstrated that the timing of entry into mitosis is controlled by the *Cdc25: Wee1* ratio. As cells grow during the G2 phase, the levels of Cdc25 mRNA and protein increase until mitosis is triggered, after which there is a sharp drop in both mRNA and protein abundance

(Moreno *et al.*, 1990). It has been proposed that this accumulation pattern results in a gradually increasing *cdc25: wee1* ratio during the G2 phase, and that the ratio becomes sufficient to trigger mitosis when cells reach a critical size (Lew and Kornbluth, 1996).


In multicellular eukaryotes, phosphorylation and dephosphorylation of both threonine 14 (animals) and tyrosine 15 (animals and plants) of the catalytic subunit of CDKs regulate their activity and determine the timing of G2 phase and mitosis (Dunphy, 1994). Phosphorylation on tyrosine 15 and threonine 14 inactivates CDK, whereas dephosphorylation by CDC25 activates the enzyme, triggering the G2 to M phase transition (Kugmagai and Dunphy, 1991). Threonine 14 and tyrosine 15 are buried beneath the T loop structure, and cyclin binding induces a conformational change that makes these residues accessible for phosphorylation (De Bondt *et al.*, 1993) (Figure 1.7 b, c).





**a**


Fig. 1.7: CDK regulation by phosphorylation/dephosphorylation of threonine 14 and tyrosine 15 in a) *Saccharomyces cerevisiae*, b) animals and c) higher eukaryotes in plants.



**b**

Fig. 1.7: CDK regulation by phosphorylation/dephosphorylation of threonine 14 and tyrosine 15 in a) *Saccharomyces cerevisiae*, b) animals and c) higher eukaryotes in plants.

In *S. pombe*, the Cdc25 gene is a mitotic inducer controlling the G2-M transition (Emsell and Nurse, 1985). In humans three homologues of CDC25 have been detected.



**c** The role of WEE1 during the G2/M transition

**Fig. 1.7:** CDK regulation by phosphorylation/dephosphorylation of threonine 14 and tyrosine 15 in **a)** fission yeast, **b)** animals and **c)** putative mechanism in plants.

MYT1, a membrane-bound tyrosine, threonine protein kinase from *Xenopus* (Mueller *et al.*, 1995) and humans (Liu *et al.*, 1997), preferentially phosphorylates threonine 14 and, to a lesser extent, tyrosine 15 (Fig. 1.7 b). WEE1/MYT1 type kinases have been identified in many different eukaryotes. Most of the sequence variation in WEE1 between species occurs in the amino-terminal domain, whereas the carboxyl-terminal region, where the protein kinase domain resides, is more conserved (Mueller *et al.*, 1995; Alique *et al.*, 1997). In plants, CDK regulation by phosphorylation/dephosphorylation of tyrosine 15 is likely mediated by WEE1 and CDC25 respectively (Sun *et al.*, 1999; Sorrell *et al.*, 2002, 2005) (Fig. 1.7 c).

In *S. pombe*, the *Cdc25* gene is a mitotic inducer controlling the G2-M transition (Russell and Nurse, 1986). In humans three homologues of CDC25 have been detected:

HsCDC25A, B and C. HsCDC25C is functionally homologous to SpCdc25 (Kumagai and Dunphy, 1991). *Cdc25* encodes a phosphatase that removes phosphate groups from threonine 14 and tyrosine 15 and thus activates Cdc2 (Russell and Nurse, 1986). When the *Cdc25* gene from *S. pombe* was used to transform tobacco cells, the mitotic cells were smaller than wild type (McKibbin *et al.*, 1998, Orchard *et al.*, 2005). A small plant *CDC25* gene which comprises only the catalytic domain of the protein was isolated in *Arabidopsis thaliana* (Landrieu *et al.*, 2004). This protein can induce small mitotic cell size in fission yeast (Sorrell *et al.*, 2005).

## **1.5. The role of *WEE1* during the G2/M transition**

The focus of my thesis was to gain an understanding of *WEE1* function in plant cells. The following sections from 1.5.1 to 1.5.4 will provide a more thorough introduction to the role of *WEE1* kinase and its regulation during the cell cycle.

### **1.5.1. *WEE1* plays an important role in checkpoint control**

To maintain a specific size, cells must coordinate their growth and division. Yeast cells are thought to use cell size checkpoints to coordinate these two processes (Nurse, 1975; Fantes and Nurse, 1977; Hartwell and Unger, 1977; Johnston *et al.*, 1977; Rupes, 2002). Cell size checkpoints prevent passage through key cell cycle transitions until cells have reached a critical size. In fission yeast, the critical size requirement is exerted primarily at the G2/M transition, whereas in budding yeast it is exerted primarily at the

G1/S transition (Rupes, 2002). Although the existence of cell size checkpoints was proposed over 20 years ago, the underlying molecular mechanisms have remained elusive. Moreover, it is not understood how a cell monitors its size/growth. Genetic analysis of cell cycle checkpoints is difficult because mutations that accelerate or delay cell cycle progression may have indirect effects on cell size (Kellogg, 2003). Despite these difficulties, early work in fission yeast suggested that the Wee1 kinase plays an important role in a checkpoint that coordinates cell growth and cell division at the G2/M transition (Nurse, 1975; Fantes and Nurse, 1978; Thuriaux *et al.*, 1978). Thus cloning of a *WEE1* in *Arabidopsis* (Sorrell *et al.*, 2002) became an extremely interesting tool as a putative cell size controller in plants.


### **1.5.2. The WEE1 protein kinase is regulated at multiple levels**

Early work in fission yeast suggested that Wee1 is part of a cell-size checkpoint that prevents entry into mitosis until cells have reached a critical size (Russell and Nurse, 1987). Recent experiments on *Swe1*, the budding yeast homologue of *wee1*, have provided new support for this idea (Jorgensen *et al.*, 2002; Harvey and Kellog, 2003). During the cell cycle, WEE1 protein kinases can be negatively regulated by phosphorylation; in addition to phosphorylation WEE1 is also regulated by interaction with 14-3-3 proteins (Honda *et al.*, 1997; Lee *et al.*, 2001).

In budding yeast intricate signaling networks are required for regulation of *Swe1* (Harvey and Kellogg, 2003). WEE1 kinase from *Xenopus*, fission yeast and budding yeast undergo extensive hyperphosphorylation during mitosis. Moreover, the

hyperphosphorylated form of *Xenopus* WEE1, isolated from mitotic extracts, has reduced kinase activity (Tang *et al.*, 1993; Muller *et al.*, 1995; Sreenivasan and Kellogg 1999; Harvey and Kellogg, 2003). Hyperphosphorylation of WEE1 in *Xenopus* extracts is dependent upon mitotic CDK1 activity, that can phosphorylate WEE1 *in vitro* (Tang *et al.*, 1993; Muller *et al.*, 1995). Experiments with fission yeast and budding yeast have identified several kinases required for regulation of WEE1 kinase *in vivo*. In fission yeast, Wee1 is phosphorylated and inactivated by the Cdr/Nim1 protein kinase complex (Coleman *et al.*, 1993; Parker *et al.*, 1993). Wee1 can also be phosphorylated by the Chk1 and Cds1 checkpoint kinases (O'Connell *et al.*, 1997; Boddy *et al.*, 1998). In budding yeast, a complex signaling network is required for regulation of Swel and for coordination of cell growth and cell division at G2/M. This network includes the kinases, Gin4, Hsl1, Cla4 and Elm1. In addition, a number of proteins required for regulation of these kinases have been identified, including Nap1, Hsl7 and a family of proteins called the septins (Fig. 1.8) (Kellogg and Murray, 1995; Ma *et al.*, 1996; Altaman and Kellogg, 1997; Carroll *et al.*, 1998; Tjandra *et al.*, 1998; Barral *et al.*, 1999; Edgington *et al.*, 1999; Shulewitz *et al.*, 1999; Sreenivans and Kellogg, 1999; Longitane *et al.*, 2000).





**Fig. 1.8:** Signaling network required for regulation of Swe1 in budding yeast. Kinases are shown in yellow, proteins required for regulation of kinases are shown in blue and phosphatase is shown in green (adapted from Kellog, 2003).

However, it is unclear how the kinase signaling network regulates Swe1 because none of these proteins has been found to phosphorylate Swe1 directly. In addition the physiological signals that regulate the network are poorly understood, although there is some evidence that components of the network respond to nutritional cues (Fig. 1.8) (Garrett, 1997; Cullen and Sprague, 2000; La Valle and Wittemberg, 2001). Many of the proteins that function in the network are highly conserved, suggesting a similar network

in all eukaryotes. For example, budding yeast Gin4 and Hsl1, are related to fission yeast Cdr1/Nim1 and Cdr2 (Kellogg, 2003).

The human and *Xenopus* WEE1 kinases are negatively regulated by phosphorylation in a cell cycle specific manner (Tang *et al.*, 1993; McGowan *et al.*, 1995; Muller *et al.*, 1995; Parker *et al.*, 1995; Watanabe *et al.*, 1995). Studies carried out in vertebrate cells, have also shown that 14-3-3 proteins can bind and positively regulate WEE1 kinases (Wang *et al.*, 2000; Lee *et al.*, 2001; Rothblum-Oviatt *et al.*, 2001). *Xenopus* WEE1 is regulated by association with 14-3-3 proteins; this association increases the kinase activity of XWEE1 and also affects its distribution inside the cell (Lee *et al.*, 2001). During interphase, WEE1 is found almost exclusively in the nucleus; when the cell enters mitosis, WEE1 is relocalised into the cytoplasm (Baldin and Ducommun, 1995). Binding of 14-3-3 to XWEE1 occurs during interphase, but not M-phase and requires phosphorylation of XWEE1 on Ser-549. The *Xenopus* checkpoint kinase CHK1 can phosphorylate the critical Ser-549 in the 14-3-3 binding site of XWEE1 (Lee *et al.*, 2001). Similarly in humans, phosphorylation of WEE1 on SER-642 is essential for the binding of 14-3-3 proteins to WEE1 (Wang *et al.*, 2000). Like *Xenopus* WEE1, interactions of 14-3-3 proteins with human WEE1 are reduced during mitosis, but not during interphase when interaction between 14-3-3 proteins and WEE1 increases the enzymatic activity of WEE1; this indicates that 14-3-3 proteins function as positive regulators of the human WEE1 protein kinase (Rothblum-Oviatt *et al.*, 2001)

### **1.5.3. WEE1 protein levels fluctuated during the cell cycle**

In addition to phosphorylation and interaction with 14-3-3 proteins, human WEE1 is also regulated at the level of protein synthesis and stability (Watanabe *et al.*, 1995). WEE1 protein levels rise during the S and G2 phases of the cell cycle because of increased synthesis, and WEE1 protein levels fall during M phase because of decreased synthesis combined with proteolysis (Rothblum-Oviatt *et al.*, 2001). Experiments using budding yeast have shown that the Swe1 protein is stable during G2/M and is not degraded until exit from mitosis (Sreenivasan and Kellogg, 1999; Harvey and Kellogg, 2003). It was also concluded that Swe1 is targeted for destruction by the SCF ubiquitin ligase complex (Sia *et al.*, 1998; McMillan *et al.*, 1999, 2002).

### **1.6. Tobacco BY-2 cells**


Dispersed plant cell suspension cultures allow the study of cell division free of developmental constraints, by providing a homogeneous population of near-identical cells (Gould, 1984). Synchronization of such cultures can provide material representative of specific cell cycle phases, and this may be achieved either by removal and subsequent re-supply of a compound required for growth, such as phosphate, nitrate, hormones or sucrose (King *et al.*, 1973; Amino *et al.*, 1983; Kodama *et al.*, 1991; Nishida *et al.*, 1992; Riou-Khamlichi *et al.*, 1999; Riou-Khamlichi *et al.*, 2000), or by applying reversible blocks at different stages of the cell cycle using specific inhibitors (Gould, 1984; Nagata *et al.*, 1992; Magyar *et al.*, 1993; Perennes *et al.*, 1993; Fukuda *et al.*, 1994; Glab *et al.*,



1994; Lucretti and Dolezel, 1995; Ito *et al.*, 1997; Planchais *et al.*, 1997, 2000; Binarova *et al.*, 1998; Roudier *et al.*, 2000). However, in such plant cell suspensions, a synchronous population rarely reaches a mitotic index >10%. The notable exception is the tobacco BY-2 cell line, which is widely used in cell cycle studies (Nagata *et al.*, 1992; Samuels *et al.*, 1998).

The TBV-2 cell line was established from callus induced from seedlings of *Nicotina tabacum* L. cv. Bright Yellow 2 in the Central Research Institute of the Japan Tobacco and Salt Public Corporation (now the Tobacco Science Research Laboratory, Japan Tobacco, Inc.) (Kato *et al.*, 1972). It is propagated in the medium of Linsmaier and Skoog (1964) supplemented with sucrose and 2, 4-dichlorophenoxyacetic acid (2, 4-D). According to Kato *et al.* (1972) the TBV-2 cell line was the most proliferative among the lines examined created from 40 species of *Nicotina* and three species of *Populus*, which suggests that this cultivar of tobacco had unique characteristics. Also, unlike unstable cultures that exhibit mixoploid cells, the TBV-2 line is remarkably stable.

TBV-2 cells exhibit rapid growth in one week of batch culture at 27°C (Fig. 1.9) and can be easily synchronised using the reversible DNA polymerase  $\alpha$  inhibitor, aphidicolin (Sala *et al.*, 1980). Following the synchrony with aphidicolin, TBV-2 cells are the only plant cell line that can attain mitotic indices of 40 to 50% (% frequency of cells in division) (Sorrell *et al.*, 2001).



**Fig. 1.9:** TBY-2 batch cultures at day 0 and day 7

A wide variety of cell cycle processes, including phase specific gene expression (Reichheld *et al.*, 1995; Combettes *et al.*, 1999; Sorrell *et al.*, 1999), CDK activity (Porceddu *et al.*, 2001; Sorrell *et al.*, 2001) and microtubule rearrangements (Hasezawa and Nagata, 1991) have been studied using this model cell line.

## 1.7. Aims

In reviewing the plant cell cycle, much is now known about plant CDKs and plant cyclins. However, until fairly recently very little was known about the phosphoregulation of CDK complexes at G2/M transition. The first report of a partial plant *WEE1* cDNA from *Zea mays* (*ZmWEE1*) appeared seven years ago (Sun *et al.*, 1999), and a full length clone of *Arabidopsis thaliana WEE1* (*AtWEE1*) was first presented by Sorrell *et al.* (2002). Moreover, the first report about a putative plant *CDC25* has only been published in the last 12 months (Sorrell *et al.*, 2005). Interestingly, expression of *AtWEE1* and *AtCDC25* in fission yeast induced a long and short cell length respectively. These data indicate cell cycle function for both genes.

My work began with a central hypothesis that *AtWEE1* regulates plant cell size. I transformed the TBY-2 cell line and *Arabidopsis* plants with *AtWEE1* which then enabled me to test this hypothesis by analyzing the effect of this *transgene* on the:

- TBY-2 cell cycle
- Mitotic cell area

As well as exploiting the benefit of TBY-2 cells for cell cycle work, this strategy also prevented gene silencing that might have occurred by over-expressing *AtWEE1* in *Arabidopsis* cells.

- I also cloned a portion of *Nicotiana tabacum WEE1* (*NtWEE1*) thereby enabling an examination of the effects of *AtWEE1* expression on *NtWEE1* expression
- Tobacco CDKA and CDKB1 kinase activity were investigated in TBY-2 cells expressing *AtWEE1*


- Primary root growth and primordial formation were investigated in *Arabidopsis* plants over-expressing *AtWEE1*.

## **Chapter 2: Materials and Methods**

### **2.1. BIN HYG TX and pTA7002 plasmids**


#### **2.1.1. Construction of plasmids**

The *Arabidopsis thaliana* *WEE1* gene (*AtWEE1*) was used to make constructs for transformation into tobacco BY-2 cells. Two different constructs (BIN HYG TX-*AtWEE1* and pTA7002-*AtWEE1*) were made and used independently to transform tobacco cells, alongside the empty vectors (BIN HYG TX and pTA7002). BIN HYG TX is a constitutive system (Fig. 2.1), originally designed for use in a tetracycline inducible system (Gatz *et al.*, 1991, 1992; Böhner *et al.*, 1999), but in this work only the constitutive component was used. BIN HYG TX is composed of a promoter derived from the cauliflower mosaic virus 35S gene promoter (35S TX promoter), which is attenuated by approximately 20X. The *AtWEE1* open reading frame (*AtWEE1* ORF), is inserted between the *Sma*I and *Sal*I restriction sites, and its transcription is terminated by a terminator (ocs) (Table 2.1). The selectable marker for plant transformation is hygromycin (hpt) which has its own promoter (nos) and terminator (Ag7).



**Fig. 2.1:** Constitutive promoter BIN HYG TX-*WEE1*

Plasmid pTA7002 is based on an inducible system developed by Prof. N.H. Chua (Aoyama and Chua 1997). (Fig. 2.2). This system has been developed using the regulatory mechanism of vertebrate steroid hormone receptors. It is composed of a transcription factor (GVG), consisting of the DNA-binding domain of the yeast transcription factor (GAL4), the transactivating domain of the herpes viral protein (VP16) and the receptor domain of the rat glucocorticoid receptor (GR). GR is not only a receptor molecule but also a transcription factor which, in the presence of a glucocorticoid like dexametasone (DEX), activates transcription from promoters containing glucocorticoid response elements. The 35S promoter drives the *GVG* gene, which is terminated by E9. In the second transcription unit six copies of the GAL4 upstream activating sequence (GAL4 UAS) are fused to the promoter (UASc promoter) to control transcription of the target gene (*AtWEE1*) inserted between the *XhoI* and *SpeI* restriction sites. Transcription of *AtWEE1* is terminated by 3A. These two transcription units are separated by a selectable marker for plant transformation (hygromycin (hpt)) which has its own promoter (nos) and terminator (nos) (Table 2.1).



**Fig. 2.2:** Inducible promoter pTA7002-*WEE1*

Both plasmids BIN HYG TX and pTA7002 also contain a kanamycin resistance gene active in bacteria (Table 2.1).

**Table 2.1:** Details of the transgene vector and antibiotic selection

| Vector                       | Transgene     | Antibiotic selection:<br><i>Agrobacterium tumefaciens</i> and<br><i>Escherichia coli</i> | Antibiotic selection:<br>Tobacco TBY-2 | Notes on Vector  |
|------------------------------|---------------|--|--|--|
| BIN HYG TX-<br><i>AtWEE1</i> | <i>AtWEE1</i> | Kanamycin  | Hygromycin                             | It has an attenuated form of the standard CaMV35S promoter |
| BIN HYG TX                   | none          | Kanamycin  | Hygromycin                             | It has an attenuated form of the standard CaMV35S promoter |
| pTA7002-<br><i>AtWEE1</i>    | <i>AtWEE1</i> | Kanamycin  | Hygromycin                             | CaMV35S  |
| pTA7002                      | none          | Kanamycin  | Hygromycin                             | CaMV35S  |

To construct BIN HYG TX-*AtWEE1*, the *AtWEE1* gene was amplified from *Arabidopsis thaliana* cDNA by PCR (see section 2.4.1, 2.4.2 and 2.4.3 for details

about RNA extraction, DNase treatment of cDNA and cDNA synthesis respectively; also see section 2.4.3 for details about cDNA amplification). The fragment corresponding to *AtWEE1* was purified and the gene was isolated using the QIAquick Gel Extraction Kit Protocol (QIAGEN). *AtWEE1* was amplified using primers P35SX (5'-AGGCCCCGGGCTCGAGATGTTTCGAGAAGAACGG-3') and P36SS (5'-GCACACTAGTCGACTCAACCTCGAATCCTAT-3'), that included *SmaI* and *SalI* sites respectively. PCR reactions were made as follows: 1 µl of cDNA was amplified with 0.625 U of Qiagen Taq polymerase, Qiagen buffer, 1.5 mM MgCl<sub>2</sub>, 0.2 mM dNTP and 100 ng of each of the primers. Thermocycle conditions were: 15 min 95°C hot start followed by 35 cycles of 95°C (1 min), 60°C (1 min) and 72°C (1 min). The PCR machine used was an MJ Research PTC-100™.

The *AtWEE1* (2 µg) PCR product was then digested with *SmaI* and *SalI* restriction enzymes (Promega) using Buffer D (10%) (Promega) in a total volume of 50 µl for 2 hours at 37°C. Digests were purified using the QIAquick Purification Kit Protocol (QIAGEN).

BIN HYG TX vector was extracted from *E. coli* cells (see paragraph 2.2.1 for details about vector extraction) and 2 µg were restriction digested in two separate reactions using either *SmaI* or *SalI* restriction enzymes (Promega) to ensure that both enzymes were able to cut the vector. The reactions were performed in a total volume of 50 µl using MCx10 (10%) (Promega) and incubated at 37°C for 2 hours. Digests were restricted with the second enzyme *SmaI* or *SalI* respectively and then purified using the QIAquick Purification Kit Protocol (QIAGEN).

The restricted vector (100 ng) was then ligated (using T4 DNA ligase enzyme (Promega)) to the PCR-amplified and restricted *AtWEE1* (40 ng). The reaction was



performed in a total volume of 10  $\mu$ l using ligase buffer (10%) and incubated at 4°C over-night.

To construct pTA7002-*AtWEE1*, the vector was extracted from *E. coli* cells (see paragraph 2.2.1 for details about vector extraction), cut with *Xho*I and *Spe*I restriction enzymes and ligated to the PCR-amplified ORF of *AtWEE1* (see above for details). The same primers were used to amplify the *AtWEE1* ORF, P35SX (5'-AGGCCCGGGCTCGAGATGTTTCGAGAAGAACGG-3') and P36SS (5'-GCACACTAGTCGACTCAACCTCGAATCCTAT-3'), which included *Xho*I and *Spe*I restriction sites respectively.

## **2.2. Transformation into bacterial cells and tobacco cells**

### **2.2.1. *Escherichia coli* DH5 $\alpha$ transformation**

Plasmids BIN HYG TX-*AtWEE1*, BIN HYG TX-empty vector, pTA7002-*AtWEE1* and pTA7002-empty vector were used to transform *Escherichia coli* DH5 $\alpha$  competent cells.

*E. coli* DH5 $\alpha$  cells were made competent by growing them into 5 ml of 2xYT-medium (per litre: 16 g bacto-tryptone; 10 g bacto-yeast extract; 5 g NaCl; pH=7) over-night. Four ml of the over-night culture were inoculated in 1 l of 2xYT and grown to OD<sub>600</sub>=0.5, cells were then centrifuged at 6000 g for 15 min at 1°C (Beckman Coulter J-E centrifuge, rotor JA-14) and resuspend in 10 ml of cold 0.1 M CaCl<sub>2</sub> plus 2.5 ml of 80% glycerol. Cells were then aliquoted into microfuge tubes, incubated on ice for 30 min and stored at -80°C.

One hundred  $\mu$ l of *E. coli* DH5 $\alpha$  competent cells were added to 2  $\mu$ l of the ligation mixture in a 14 ml sterile tube and left on ice for 20 min, then heat shocked (42°C for

45 seconds) and returned to ice for a further 2 min. Subsequently, the cells were added to 900  $\mu$ l of LB medium (per litre: 10 g bacto-tryptone; 5 g bacto-yeast extract; 10 g NaCl; pH=7) and incubated at 37°C gently shaking for 1 hour, 200  $\mu$ l of cells were spread onto Petri dishes containing LB-medium (0.8%) agar plus kanamycin (final concentration 50  $\mu$ g/ml) and incubated at 37°C over-night.

To extract plasmids from clones, the cells were inoculated in 10 ml LB with kanamycin (final concentration 50  $\mu$ g/ml), and incubated over-night at 37°C with shaking at 200 rpm. Bacterial cells were harvested by centrifugation at 6000 g for 15 min at 4°C (Beckman Coulter J-E centrifuge, rotor JA-14). DNA was extracted using the Qiagen midiprep kit (QIAGEN). This kit is designed for preparation of up to 100  $\mu$ g of high- or low-copy plasmid DNA. The kit provides lysis buffers to disrupt the bacterial cell wall and membrane and ribonuclease (RNase A) to remove the RNA molecules. Cell debris and proteins are separated from the plasmid DNA by centrifugation. The supernatant containing the plasmid DNA is applied to the QIAGEN-tip 100 column. This column contains a resin able to bind the plasmid DNA, but not other elements that contaminate the DNA, which are washed through. Elution DNA buffer is then applied to the column to release the plasmid DNA from the resin. The DNA is then precipitated by adding isopropanol to the eluted DNA and centrifuged. The pellet is washed with 70% ethanol and centrifuged. Ethanol is then removed and the pellet air-dried for 30 min, and redissolved in 50  $\mu$ l Buffer EB (10 mM Tris-Cl, pH=8.5).

### **2.2.2. *Agrobacterium tumefaciens* LBA4404 transformation**

One  $\mu\text{g}$  of plasmid DNA was added to 100  $\mu\text{l}$  of competent *Agrobacterium tumefaciens* LBA4404 cells (kindly provided by Dr. David Sorrell) and frozen in liquid nitrogen for 10 sec before being thawed for 5 min at 37°C and added to 1 ml of 2xYT-medium. The cells were incubated for 4 h at 30°C with shaking, centrifuged at 1000 rpm for 1 min in a microcentrifuge (Eppendorf MiniSpin) and resuspended in 100  $\mu\text{l}$  of 2xYT. One hundred  $\mu\text{l}$  of cells were plated on a Petri dish containing 2xYT-media and kanamycin (final concentration 50  $\mu\text{g}/\text{ml}$ ). Transformed *Agrobacterium tumefaciens* LBA4404 cells grew in 3-4 days at 30°C.

### **2.2.3. Transformation of tobacco BY-2 cells with *AtWEE1***

Stable transformation of TBV-2 cells was achieved using a modified version of the method described by An (1985). Isolated colonies of *Agrobacterium tumefaciens* LBA4404 containing either BIN HYG TX-*AtWEE1*, BIN HYG TX-empty vector, pTA7002-*AtWEE1* or pTA7002-empty vector were picked from fresh 2xYT-kanamycin (50  $\mu\text{g}/\text{ml}$ ) (Table 2.1) plates and cultured over-night in 7 ml 2xYT, without antibiotic, in 50 ml conical flasks at 30°C with shaking. Four ml aliquots of 6 day old stationary phase TBV-2 cells containing 20  $\mu\text{M}$  of freshly added Acetosyringon (Sigma-Aldrich) were co-cultivated with 100  $\mu\text{l}$  of *Agrobacterium* culture in 90 mm Petri dishes sealed with Nescofilm, for two days at 27°C in the dark without shaking. Cells were washed with 1 litre of BY-2 medium (per litre: 4.3 g MS basal salt medium; 30 g sucrose; 10 ml of 1% myo-inositol, 10 ml of 2%  $\text{KH}_2\text{PO}_4$ ; 1 ml of 0.1 % thiamine HCl; 2 ml of 0.01 % of 2.4-D; pH=5.8)


using a cell dissociation sieve fitted with a 100  $\mu\text{m}$  mesh (Sigma-Aldrich) and resuspended in 5 ml BY-2 medium containing 250  $\mu\text{g/ml}$  Timenten (Melford Laboratories), 2.5 ml aliquots were then plated onto solidified BY-2 medium (0.8% agar) supplemented with 250  $\mu\text{g/ml}$  Timenten and 80  $\mu\text{g/ml}$  hygromycin (Table 2.1). Plates were sealed with micropore tape and incubated at 27°C in the dark. Hygromycin resistant calli were isolated after 2-4 weeks. Each individual callus was considered as an independent clone, and grown for a further two weeks on fresh plates. Calli were then transferred to 95 ml BY-2 medium supplemented with 250  $\mu\text{g/ml}$  Timenten and 80  $\mu\text{g/ml}$  hygromycin and incubated at 27°C and 130 rpm in the dark until the cultures reached stationary phase (3-5 weeks). Cultures were subsequently maintained as described below section 2.3.1. Cultures were subjected to at least four rounds of subculturing before being used in synchrony experiments.

## **2.3. Tobacco BY-2 cells synchronization, mitotic index and cell area measurements**

### **2.3.1. Tobacco BY-2 cell culture propagation**

The tobacco BY-2 cell line (Nagata et al., 1992; Nagata and Kumagai, 1999) is the most highly synchronizable plant cell system known and is thus ideal for studies of the plant cell cycle. The tobacco BY-2 cell line is cultured at 27°C in darkness in modified Linsmaier and Skoog (1964) medium and subcultured at 7 day intervals by transferring 3 ml of the 7-d-old culture to 95 ml of a new modified Linsmaier and Skoog medium (Fig. 2.3).

### 2.3.2. Synchronization of tobacco BY-2 cells




**Fig. 2.3:** The tobacco BY-2 batch culture method. From each mother culture two new sub-cultures are made every week. After 7 days of growth one culture is used to make a new subculture, whereas the other culture is kept for one more week to be used in case of contamination of the new sub-culture.

### **2.3.2. Synchronization of tobacco BY-2 cells**

Aphidicolin, a specific and reversible inhibitor of DNA polymerase  $\alpha$ , was used to synchronize cells. Twenty ml of 7 day old TBV-2 cells are transferred to 95 ml of fresh modified Linsmaier and Skoog medium in which aphidicolin (final concentration 5 mg/l) was added. Aphidicolin was dissolved in dimethyl sulfoxide (DMSO; 5 mg/ml, Sigma) and stored at 4°C in the dark. In the presence of this inhibitor, any cell replicating its nuclear DNA is arrested and all other cells are unable to enter S phase. Hence, a 24 h exposure to, and subsequent removal of, aphidicolin will cause the majority of cycling cells to accumulate in late G1 and then to be released into S-phase upon removal of aphidicolin (Nagata and Kumagai, 1999; Herbert et al., 2001). The drug was removed from the culture by washing with 1 l of fresh medium: the cell suspension was poured into a cylindrical funnel with a fused-in sintered glass filter (17 G2 Hario glass, Tokyo). The flow rate of the washing medium was controlled with a Hoffman clamp placed on a rubber tube connected to the bottom of the funnel (Fig. 2.4).

### 2.3.3. Mitotic index measurement



### 2.3.4. Cell area

**Fig. 2.4** TBY-2 cells can be synchronised in mitosis, which allows for easy study of the plant cell cycle, using the reversible DNA polymerase inhibitor aphidicolin which blocks the cells in S-phase of the mitotic cell cycle.

A total washing time of 15 minutes was optimal for obtaining high levels of synchrony, and the cells were subsequently resuspended in the same volume of fresh medium.

### **2.3.3. Mitotic index measurements**

Tobacco BY-2 cell transgenic lines were used for synchronization experiments. Aphidicolin was used to synchronize the cells. Following release from aphidicolin, 100  $\mu$ l of cell suspension were taken at hourly intervals for 24 hours, and mixed immediately with 5  $\mu$ l Hoechst stain (Bisbenzimidazole Sigma, 100  $\mu$ g/ml in 2% (v/v) Triton X-100). The mitotic index, the sum of prophase, metaphase, anaphase and telophase mitotic figures as a percentage of all cells, was measured for a minimum of 300 cells per slide in a random transect. Cells and mitotic figures were visualized using a fluorescence microscope (Olympus BH2, UV,  $\lambda=420$ nm).

### **2.3.4. Cell area**

Digital images of TB-2 cells were taken using a Fujifilm HC-300Z digital camera attachment for the microscope and saved as JPEGs. Digital images of mitotic cells, captured as described above, were measured using SigmaScan® (Sigma), an image analysis program. To take a mitotic cell area measurement, a line was traced around the perimeter of cells undergoing mitosis. To ensure measurements were accurate, a graticule slide was used as a distance reference point from which SigmaScan® could calculate the area of a cell. For each experiment undertaken in this project, mitotic cell area was measured. On average, 150 measurements were made per experiment and the data obtained was stored in Microsoft® Excel.



### **2.3.5. Growth rate measurement**

Tobacco BY-2 cell density was measured for a period of 7 days (from the day at which the new subculture started to the day at which cells reached stationary phase) using a spectrophotometer  $\lambda=550\text{nm}$  (Smartspec<sup>TM</sup>3000, Bio-Rad laboratories Ltd). Where absorbance readings were in excess of 0.8, samples were diluted to ensure an accurate measurement.

### **2.3.6. Tobacco BY-2 cells containing pTA7002**

Dexamethasone (final concentration 1  $\mu\text{M}$ ) was added three days after subculturing to the TBV-2 cells transformed with pTA7002-*WEE1* vector. Every 12 h, 100  $\mu\text{l}$  of cell suspension were taken and mixed immediately with 5  $\mu\text{l}$  Hoechst stain (Bisbenzamide Sigma, 100  $\mu\text{g cm}^{-3}$  in 2% (v/v) Triton X-100). Samples were collected for a 48 h period and the images were captured and measured using SigmaScan.

## **2.4. RT-PCR and semi quantitative RT-PCR**

### **2.4.1. RNA extraction**

Tobacco BY-2 cells were harvested by centrifugation at 3500 rpm (MSE Centaur 2) for 10 minutes; the pellet was then collected in aluminum foil and frozen in liquid nitrogen, cells were stored at  $-80^{\circ}\text{C}$  until required. The frozen cells were ground to a fine powder using liquid nitrogen and a pestle and mortar precooled to  $-20^{\circ}\text{C}$ . TRI reagent (2 ml) (Sigma-Aldrich, Gillingham, UK) was added to TBV-2 cells

and ground until a homogenous paste was formed. Equal amounts of paste were transferred to two 1.5 ml microcentrifuge tubes, vortexed and left at room temperature for 5 minutes. Samples were subsequently centrifuged at 12000 rpm (Beckman Coulter Allegra™ 21R, rotor F2402H) at 4°C for 10 minutes. The supernatant was transferred to two new microcentrifuge tubes leaving the solid plant material behind. Two hundred µl of chloroform were added to each tube, vortexed for 15 seconds and left at room temperature for 5 minutes. Samples were then centrifuged at 12000 rpm (Beckman Coulter Allegra™ 21R, rotor F2402H) at 4°C for 15 minutes. The top layer containing RNA was transferred to new 1.5 ml microcentrifuge tubes and 0.5 ml of isopropanol were added. The samples were mixed and left at room temperature for 15 minutes and then centrifuged at 12000 rpm (Beckman Coulter Allegra™ 21R, rotor F2402H) at 4°C for 10 minutes. The supernatant was removed and 1 ml of ethanol was added to wash the pellet. The samples were vortexed for 15 seconds and centrifuged at 12000 rpm (Beckman Coulter Allegra™ 21R, rotor F2402H) at 4°C for 1 minute. The supernatant was carefully removed and the pellet air dried for 30 minutes. The pellet was resuspended in 50 µl UHP H<sub>2</sub>O. Samples were stored at -80°C.

#### **2.4.2. DNase treatment of cDNA**

Residual genomic DNA was removed from total RNA extracted, by adding 10 µl 10x DNase buffer and 3 µl DNase (Ambion Inc, Austin, USA). Preparations were incubated at 37°C for 1 hour. DNase deactivation reagent (0.2 ml) was added and samples incubated at room temperature for 2 minutes before centrifugation at 13000 rpm (Eppendorf® MiniSpin) for 1 minute. The supernatant was transferred to a new microcentrifuge tube and stored at -80°C.

Quantity and quality of RNA was estimated by running on an agarose gel. Five  $\mu\text{l}$  of extracted RNA were mixed with 1  $\mu\text{l}$  of loading buffer (150 mM Tris HCl pH=7.6; 50% glycerol; 0.4% bromophenol blue and  $\text{dH}_2\text{O}$ ) and run on an electrophoresis gel (1% agarose, 1  $\mu\text{g}/\text{ml}$  EtBr) in TAE buffer (1 l of 50xTAE buffer contains: 242 g of Tris base; 57.1 ml of glacial acetic acid and 100 ml of 0.5 M EDTA). To minimize RNA degradation the gel tank, tray and comb were treated with 0.1 M NaOH and then washed copiously with distilled water. The RNA was visualized under UV light and photographed using a Gene Genius Bioimaging System and Gene Snap® software package (Syngene). RNA concentration was measured using a Gene Quant (Pharmacia) spectrophotometer.

#### **2.4.3. cDNA synthesis**

To ensure that comparison between samples could occur, equal amounts of RNA from each sample were reverse transcribed. RNA (5  $\mu\text{g}$ ) in a 0.5 ml microcentrifuge tube was made up to 20  $\mu\text{l}$  with sterile  $\text{dH}_2\text{O}$ . One  $\mu\text{l}$  of oligo (dt)-15 (50  $\mu\text{g}/\text{ml}$ ; Promega) was added and samples incubated at 70°C for 10 minutes. Samples were then cooled at 4°C for 10 minutes. Six  $\mu\text{l}$  5x 1st stand buffer (Invitrogen), 2  $\mu\text{l}$  0.1M DTT (dithiothreitol), and 1  $\mu\text{l}$  10 mM dNTPs were added and incubated at 42°C for 2 minutes. Superscript II reverse transcriptase (1  $\mu\text{l}$ ) (GibcoBRL, Paisley, UK) was added and the samples were incubated for 50 minutes at 42°C. The reactions were heated at 70°C for 15 minutes to inactivate the reverse transcriptase and stored at -20°C. Mock reactions, important to ensure no genomic DNA contamination, were also carried out following the above procedure, but Superscript II reverse transcriptase was not added.

#### 2.4.4. PCR amplification of cDNA

cDNA was subjected to PCR (Polymerase Chain Reaction) amplification using the Reddymix PCR master Mix system (ABgene). This contains all reagents necessary for DNA amplification including dNTPs (0.2 mM); thermoprime plus DNA polymerase (1.25 u); MgCl<sub>2</sub> (1.5 mM) and loading buffer for gel electrophoresis. For a total volume of 25 µl, 22 µl of Reddymix, 0.5 µl of forward primer (10 µM), 0.5 µl of reverse primer (10 µM), 1 µl of sterile distilled water and 1 µl of DNA was used. Where possible a master mix was used to reduce pipetting error and a negative control (sterile dH<sub>2</sub>O substitute to cDNA) was always used to ensure no water contamination, also a positive control was used when available.

Table 2.2 shows the primers used for each gene of interest and the annealing temperatures used. Thermocycle conditions were: 35 cycles of 95°C (1 min), annealing temperature (1 min) and 72°C (1 min) in a PTC100 thermocycler. PCR products were run out on an agarose gel (1 %) with EtBr (1 µg/ml) and visualized under UV light.

#### 2.4.5. Design of degenerate and non-degenerate primers

Available amino acid sequences of homologous *WEE1* plant genes from *Zea mays* (*ZmWEE1*) and *Arabidopsis thaliana* (*AtWEE1*) were aligned and highly conserved regions of at least 6 amino acids were found. Selected from these conserved regions were those with at least the last two amino acids at the 3' end conserved across all genes and coded by as few codons as possible, for example methionine which is encoded by only one codon (best 3' amino acids are M and W,

worst are R, L and S). If there were discrepancies between homologues, degeneracies (wobbles) were introduced but sequences with as few wobbles as possible were selected. Regions that required wobbles at the 3' end were avoided. Melting temperatures (T<sub>m</sub>) of the selected regions were calculated using

$$T_m = 2^{\circ}\text{C} \times (\text{A}+\text{T}) + 4^{\circ}\text{C} \times (\text{C}+\text{G})$$

and equalized by the deletion or addition of extra bases if necessary.

Degenerate primers were used to isolate a portion of *Nicotiana tabacum WEE1* (*NtWEE1*). The *NtWEE1* gene fragment was then sequenced and specific (non-degenerate) primers were designed on the basis of its sequence (Table 2.2).

**Table 2.2:** Primer pairs used for DNA amplification (PCR)

| Primer pair        | Target gene  | Oligonucleotide sequence                            | Product size (bp) | T <sub>m</sub> (°C) |
|--------------------|--|---|-------------------|---------------------|
| AtWEE1F<br>AtWEE1R | <i>AtWEE1</i>  | 5'-AGCTTGTCAGCTTTGCCT<br>5'-CGTGCATCCCTCCTTCTTCTACT | 229               | 55                  |
| WEE1F<br>WEE1R     | <i>NtWEE1</i><br>(degenerate primers used to isolate a portion of the <i>WEE1</i> gene from tobacco) | 5'-TCKTGgTtYgARAAyGARCA<br>5'-AGAGAAGATrTCNACyTTRTC | 339               | 60                  |
| WEE1TSF<br>WEE1TSR | <i>NtWEE1</i><br>(used to amplify <i>N. tabacum WEE1</i> )   | 5'-GTGACCACAGCTTATCCAAC<br>5'-GCATCCCCCTCTTCGATC    | 228               | 60                  |

#### **2.4.6. Estimation of optimum PCR cycles for semi-quantitative RT-PCR**

The optimum number of PCR cycles for semi-quantitative PCR was determined for each of the four specific primer pairs by carrying out PCRs on a set of standards over a range of cycles (Table 2.3). The standards were produced from dilutions of a mixed cDNA made using 1 µl of each cDNA sample. The products were run on electrophoresis gels, stained with ethidium bromide and the intensity of each band measured on a Gene Genius Bioimaging System (Syngene Ltd.).

An appropriate number of PCR cycles, which gave a steep linear plot of band strength against concentration, with an intercept close to zero, was selected for each primer pair for use in semi-quantitative work.

#### **2.4.7. Semi-quantitative RT-PCR.**

Samples of cells (5 ml) were taken during synchronization experiments every hour after removal of aphidicolin, and immediately frozen at -80°C. At the same time, measurement of the mitotic index was carried out. RNA was extracted from each sample and used to synthesize cDNA.

PCRs were carried out on the cDNA samples prepared using each of the specific primer pairs at the optimum cycle number as described in section 2.4.4 except that cycles were limited as shown in table 2.3. All PCRs were repeated at least three times to provide replicate results. A set of the standards were included in each PCR run. PCR products were run on ethidium bromide (1 µg/ml) stained agarose (1 %) gels and the band strengths measured using a Gene Genius Bioimaging System and Gene Tools software package (Syngene Ltd.).

The measurement of gel band strength, for all genes of interest, was normalized by dividing the EtBr fluorescence obtained from primers to the gene of interest by the EtBr fluorescence obtained from 18S ribosomal primer set (PUV2 and PUV4).

**Table 2.3:** Primer details for Gene Expression Analysis

| Primer pair        | Target gene                 | Oligonucleotide sequence   | Product size<br>(bp) | T <sub>m</sub> (°C) | PCR<br>cycles<br>used |
|--------------------|-----------------------------|--|----------------------|---------------------|-----------------------|
| PUV2<br>PUV4       | <i>18 ribosomal<br/>RNA</i> | 5'-TTCCATGCTAATGTATTCAGAG<br>5'-ATGGTGGTGACGGGTGAC                     | 488                  | 60                  | 36                    |
| H4F<br>H4R         | <i>Histone H4</i>           | 5'-GGCACAGGAAGGTTCTGAGGGA<br>TAACA<br>5'-TAACCGCCGAAACCGTAGAGAG<br>TCC | 320                  | 60                  | 30                    |
| AtWEE1F<br>AtWEE1R | <i>AtWEE1</i>               | 5'-AGCTTGTCAGCTTTGCCT<br>5'-CGTGCATCCCTCCTTCTTCTACT                    | 229                  | 55                  | 30                    |
| WEE1TSF<br>WEE1TSR | <i>NtWEE1</i>               | 5'-GTGACCACAGCTTATCCAAC<br>5'-GCATCCCCCTCTTCGATC                       | 228                  | 60                  | 30                    |

## **Chapter 3: Expression of *Arabidopsis thaliana* WEE1 (*AtWEE1*) in TBY-2 cells induces a shortened G2 phase, a premature entry into mitosis and a smaller mitotic cell area**

### **3.1. Introduction**

WEE1 kinases function in a highly conserved mechanism that controls the timing of entry into mitosis. Loss of Wee1 and Swe1 function causes fission yeast and budding yeast cells to enter mitosis before sufficient growth has occurred, leading to formation of daughter cells that are smaller than normal (Nurse, 1975; Harvey and Kellog, 2003). Conversely, when WEE1 is over expressed, mitosis is delayed until cells grow to a larger size (Russell and Nurse, 1987). Early work in fission yeast suggested that Wee1 kinase is part of a cell size checkpoint that prevents entry into mitosis before cells have reached a critical size (Nurse, 1975; Fantes and Nurse, 1978). In fission yeast cells, *wee1* is the main genetic element in cell size control (Nurse, 1975; Fantes and Nurse, 1978), Wee1 inhibits CDK1 (encoded by the *cdc2* gene) by phosphorylating a highly conserved tyrosine (Tyr 15) residue near to the N-terminus which results in an inhibition of mitosis (Russell and Nurse, 1987), whereas Cdc25 promotes mitosis by removing the inhibitory phosphate (Russell and Nurse, 1986).

In our lab, the fission yeast *Schizosaccharomyces pombe* was transformed with *AtWEE1* using an inducible promoter (pREP1-*AtWEE1*). Expression of *AtWEE1* in fission yeast cells induced a significant increase in cell length compared with wild type,



yeast cells transformed with uninduced pREP1-*AtWEE1* and pREP1-empty vector (Sorrell *et al.*, 2002). Sorrell and colleagues have also demonstrated that colony formation by cells containing induced pREP1-*AtWEE1* was substantially reduced compared to empty vector. The substantial reduction of colony formation suggested that *AtWEE1* expression was inhibiting cell division (Sorrell *et al.*, 2002). Different results came from the very recent expression of the putative mitotic inducer gene *AtCDC25* in fission yeast using an inducible promoter (pREP1-*AtCDC25*) (Sorrell *et al.*, 2005). Expression of *AtCDC25* in fission yeast induced a reduction in mitotic cell length compared with wild type, empty vector and uninduced pREP1-*AtCDC25* fission yeast cells (Sorrell *et al.*, 2005). The data reported by Sorrell and colleagues indicated that *AtCDC25* can function as a mitotic accelerator in fission yeast (Sorrell *et al.*, 2005). So, over expression of *SpWEE1* and *SpCDC25* in fission yeast resulted in a long and short cell length phenotype, respectively (Russell and Nurse, 1986, 1987), as did expression of *AtWEE1* and *AtCDC25* (Sorrell *et al.*, 2002, 2005).

One of the biggest problems in cell cycle research is how cell size is regulated at mitosis. In this chapter data are reported about tobacco BY-2 cell size at division using *AtWEE1* gene as a tool. To obviate the possibility of gene silencing, or mitotic catastrophe, *AtWEE1* was expressed in the TBY-2 cell line using an inducible system in addition to a constitutive system. The hypothesis was tested that *AtWEE1* induces large cell size in transformed TBY-2 cells.

## 3.2. Materials and methods

Experimental details are fully described in chapter 2 from section 2.1.1 to 2.4.3.

### 3.3. Results: Production of tobacco BY-2 cell lines transformed with *AtWEE1* constructs and empty vector controls

Tobacco BY-2 cells were transformed using two different plasmids: 1) a constitutive plasmid BIN HYG TX-*AtWEE1* and the respective empty vector (BIN HYG TX-EV) (see chapter 2 section 2.1.1 for more details) and 2) an inducible plasmid pTA7002-*AtWEE1* and the respective empty vector (pTA7002-EV) (see chapter 2 section 2.1.1 for more details). pTA7002-*AtWEE1* was induced by adding a synthetic glucocorticoid (dexamethasone).

All plasmids were cloned and selected in *Escherichia coli* DH5 $\alpha$  (see chapter 2, section 2.2.1). For the constitutive plasmid, one clone (1S) was isolated and sequenced (Appendix I), and there were no base changes. For the inducible construct it was necessary to isolate and sequence three different clones 1X, 2X and 3X (Appendix I), to find a clone with perfect amino acid sequence (clone 2X). The plasmids were then extracted and used to transform *Agrobacterium tumefaciens* LBA4404 cells (see chapter 2, section 2.2.2). *Agrobacterium tumefaciens* was then used to transform the tobacco cells (see chapter 2, section 2.2.3.). For each transformation 12 different clones were isolated and allowed to grow over a period of 3-4 weeks after which just three clones per

transformation were chosen (Table 3.1). The clones used for the experiments were maintained in subculture as described in chapter 2 section 2.3.1.

**Table 3.1:** tobacco BY-2 cell lines transformed with *AtWEE1* using the constitutive (BIN HYG TX-*AtWEE1*) or the inducible (pTA7002-*AtWEE1*) promoter

| Plasmids:                  | Clones: | Code used in this thesis |
|----------------------------|---------|--------------------------|
| BIN HYG TX - <i>AtWEE1</i> | 2       | WEE1-c-2                 |
|                            | 10      | WEE1-c-10                |
|                            | 12      | WEE1-c-12                |
| BIN HYG TX-empty vector    | 10      | EV-c-10                  |
| pTA7002- <i>AtWEE1</i>     | 1       | WEE1-i-1                 |
|                            | 3       | WEE1-i-3                 |
|                            | 6       | WEE1-i-6                 |
| pTA 7002-empty vector      | 1       | EV-i-1                   |
|                            | 3       | EV-i-3                   |
|                            | 4       | EV-i-4                   |

Transformation of tobacco BY-2 cells was repeated after one and a half years because TBY-2 cells from the previous transformation began to die. Table 3.2 shows the clones obtained from the new transformation:

**Table 3.2:** tobacco BY-2 cell lines transformed with BIN HYG TX-*AtWEE1* and pTA7002-*AtWEE1* new transformation


| Plasmids:                  | Clones: | Code used in this manuscript |
|----------------------------|---------|------------------------------|
| BIN HYG TX - <i>AtWEE1</i> | 1       | WEE1-c-1+                    |
|                            | 2       | WEE1-c-2+                    |
|                            | 3       | WEE1-c-3+                    |
| BIN HYG TX-empty vector    | 1       | EV-c-1+                      |
|                            | 2       | EV-c-2+                      |
|                            | 3       | EV-c-3+                      |
| pTA7002- <i>AtWEE1</i>     | 1       | WEE1-i-1+                    |
|                            | 2       | WEE1-i-2+                    |
|                            | 3       | WEE1-i-3+                    |
| pTA 7002-empty vector      | 1       | EV-i-1+                      |
|                            | 2       | EV-i-2+                      |
|                            | 3       | EV-i-3+                      |

### 3.4. Results: Analysis of tobacco BY-2 cells transformed with BIN HYG TX-*AtWEE1*

The three independent cell lines carrying *AtWEE1* under the control of an attenuated version of the 35-S cauliflower mosaic virus (CaMV) promoter, BIN HYG TX (Gatz *et al.*, 1992) were investigated in relation to cell size and cell cycle. In conjunction with the *AtWEE1* expressing lines (denoted as shown in table 3.1) one empty vector line (EV-c-10) was used as experimental control.

Forward and reverse *AtWEE1* primers were designed specifically to amplify *Arabidopsis thaliana WEE1* (see chapter 2 section 2.4.5 for more details), and were used to confirm *AtWEE1* expression in WEE1-c-2, WEE1-c-10 and WEE1-c-12 (Fig. 3.1 a). Wild type


TBY-2 cDNA was also used as negative control (Fig. 3.1 b), the quality of wild type cDNA was tested using primers PUV2 and PUV4 directed against *18S rRNA* (see chapter 2 section 2.4.5 for more details) (Fig. 3.1 c).



**Fig. 3.1:** RT-PCR analysis of *AtWEE1* expression in a) three independent tobacco BY-2 cell lines expressing *AtWEE1* constitutively (WEE1-c-2, WEE1-c-10 and WEE1-c-12) using *AtWEE1* primers, distilled water was used as the control; b) wild type (WT) cDNA was also used as negative control with the *AtWEE1* primers; c) the quality of wild type cDNA was tested using PUV2 and PUV4 primers.

### **3.4.1. Comparison of growth rate between tobacco BY-2 cell lines expressing *AtWEE1*, wild type and empty vector**

To test whether expression of *AtWEE1* under the control of the constitutive promoter or the presence of the promoter itself perturbed the normal growth rate of TB Y-2 cells, the density of the cultures was measured over the normal 7 day batch culture of TB Y-2 cells. Measures of culture density were taken every 24 hours for: WEE1-c-2, WEE1-c-10, WEE1-c-12, EV-c-10 and wild type (see chapter 2 section 2.3.5; see also Appendix II for raw data). Temporal changes in absorbance were interpreted as rate of growth over the 7 day culture period. Expression of *AtWEE1* did not alter cellular growth rate. Indeed, with the exception of WEE1-c-2, tobacco BY-2 lines expressing *AtWEE1*, wild type and empty vector all had the same pattern of growth rate. This was flat for the first three days (lag phase), raised for the next three days (exponential phase) until it reached a point where the cells were not dividing any more (stationary phase) (Fig. 3.2).



**Fig. 3.2:** Temporal changes in absorbance at 550 nm in WEE1-c-2, WEE1-c-10, WEE1-c-12, EV-c-10 and wild type.

### 3.4.2. Mitotic cell area is smaller in TBY-2 cells expressing *AtWEE1* over a 6 day period


If *AtWEE1* kinase is a negative regulator at the G2/M transition (see Introduction), then increased expression of *WEE1* in tobacco BY-2 cells should cause cells to divide at a large cell size. This hypothesis was investigated by measuring the mitotic cell area of WEE1-c-10, EV-c-10 and wild type cultures every day over a 6 day period (see Appendix II for raw data); on day 7 mitotic cells were not found. The mitotic WEE1-c-10 cells divided, except on day 2, at a constantly smaller cell size circa 2000-2500  $\mu\text{m}^2$  (Table 3.3, Fig. 3.3 a, b), compared with corresponding data for the wild type

and EV-c-10 that fluctuated between large, circa 3000  $\mu\text{m}^2$  on days 1 and 2 for wild type and 4600 and 2800  $\mu\text{m}^2$  for EV-c-10 on day 1 and 2, respectively (Table 3.3, Fig. 3.3 a, b), to small 1700-2000  $\mu\text{m}^2$  for wild type on days 3 and 4 and 2300-2800  $\mu\text{m}^2$  for EV-c-10 on days 3, 4 and 5, to large again (circa 4000  $\mu\text{m}^2$  for both wild type and EV-c-10) on day 6 (Table 3.3, Fig. 3.3 a, b). To explain the fluctuation in mitotic cell area for both wild type and EV-c-10, it is necessary to consider that the cells sub-cultured on day 1 came from a culture that was in stationary phase, where nutrients were limiting. Seemingly, cells needed a one day recovery from nutrient deficit to nutrient surplus before starting to divide at smaller cell size. However, on day 6 the nutrient deficit occurred and cells were larger again. Interestingly, mitotic cell area for WEE1-c-10 remained relatively constant from day 1 to day 6 so that significant differences were evident compared with wild type and EV-c-10. This result showed that either shortage of nutrients or expression of *AtWEE1* were not resulting in an increased mitotic cell area.




**Table 3.3:** The mean number and standard error (in parenthesis) of WT, EV-c-10 and WEE1-c-10 cell area over 6 days experiment. Levels of significance are indicated by Student's t-test.

| day | WT                | EV-c-10           | WEE1-c-10         | P (WT/WEE1-c-10) | P (EV-c-10/WEE1-c-10) |
|-----|-------------------|-------------------|-------------------|------------------|-----------------------|
| 1   | 3254 ( $\pm$ 153) | 4658 ( $\pm$ 277) | 2442 ( $\pm$ 234) | = 0.007          | = 0.000               |
| 2   | 2948 ( $\pm$ 115) | 2809 ( $\pm$ 162) | 3228 ( $\pm$ 204) | = 0.239          | = 0.115               |
| 3   | 1777 ( $\pm$ 60)  | 2345 ( $\pm$ 97)  | 2267 ( $\pm$ 153) | = 0.005          | = 0.670               |
| 4   | 1971 ( $\pm$ 64)  | 2747 ( $\pm$ 155) | 2179 ( $\pm$ 130) | = 0.156          | = 0.006               |
| 5   | 3244 ( $\pm$ 120) | 2532 ( $\pm$ 91)  | 2020 ( $\pm$ 110) | = 0.000          | = 0.001               |
| 6   | 3922 ( $\pm$ 314) | 3991 ( $\pm$ 225) | 2150 ( $\pm$ 91)  | = 0.000          | = 0.000               |



a

**Fig. 3.3:** a) Comparison between the average mitotic cell area of WEE1-c-10, wild type (WT) and EV-c-10 over a 6 day period ( $\pm$  SE) ( $n \geq 15$ ).




**Fig. 3.3: b)** Pictures of TBY-2 cells expressing *AtWEE1* compared with wild type and empty vector taken during the 6 day experiment (bars = 100  $\mu$ m).


### **3.4.3. Expression of *AtWEE1* in TBY-2 cell lines induces a premature entry into mitosis through a shortening of G2 phase**

Synchronization experiments were conducted on all transformed cell lines (WEE1-c-2, WEE1-c-10 and WEE1-c-12). Following synchronization with aphidicolin for 24 hours, aphidicolin was removed (see chapter 2 sections 2.3.2 and 2.3.3) and the mitotic index was measured every hour for 23-24 h. Notably, compared with wild type and empty vector, the mitotic index began to rise sooner in the constitutive *AtWEE1* cell lines (Fig. 3.4; Appendix III).






**a**




**b**




c

Fig. 3.4. Mitotic index curves obtained following synchronization (with aphidicolin for 24 h) of each of the cell lines, WEHI-231 cell lines, with aphidicolin (100 ng/ml) for 24 h. (a) WEHI-231, (b) WEHI-231-10, (c) WEHI-231-2, (d) WEHI-231-10.



d





e

**Fig. 3.4:** Mitotic index curves obtained following synchronization (with aphidicolin for 24 h) of each of the constitutive *AtWEE1* cell lines, wild type and empty vector. **a)** wild type (WT); **b)** EV-c-10; **c)** WEE1-c-2; **d)** WEE1-c-10 and **e)** WEE1-c-12.


For the constitutive lines the mitotic index rose between 2-3 h and peaked between 3-4 hours for WEE1-c-2, at 4 hour for WEE1-c-10 and at 5 hours for WEE1-c-12 following the release from the aphidicolin block (Fig. 3.4 c, d and e). However, for both wild type and empty vector the mitotic index rose between 6 and 8 hours and peaked at 8 hours for the wild type and 9 hours for EV-c-10 (Fig. 3.4 a and b).

The duration of the cell cycle phases were estimated using different methods as indicated in Table 3.4.

**Table 3.4:** Different methods were used to calculate the duration of the single phases of the cell cycle

| Phases of the cell cycle | Methods used to calculate the duration   |
|--------------------------|--|
| <b>G2</b>                | Using the method of Quastler and Sherman (1959), an estimate of G2 phase duration is obtained from the interval between zero and the 50% intercept of the ascending limb of the mitotic index curve (G2+½M).   |
| <b>M</b>                 | <p>M-phase was calculated using the Nachtwey and Cameron method (1968):</p> $\text{Duration of Mitosis} = \frac{\text{Cell cycle duration} \times \ln(\text{average Mitotic Index as a proportion} + 1)}{\ln 2}$ <p>Formula explanation:</p> <ul style="list-style-type: none"> <li>• Cell cycle duration/ln 2: since the cell cycle time is measured in a population of exponentially growing cells, the cell cycle duration occurs with respect to ln 2 (because an exponential curve increase by the exponent of 2)</li> <li>• ln (Mitotic Index as a proportion+1): Mitotic Index is from exponentially growing population represented by (1). The measurement is as a population (1) that doubles during exponential growth. So, the Mitotic Index value is with respect to a population that is doubling from 1 to 2.</li> </ul> |
| <b>S</b>                 | S-phase was measured as the 50% intercept of the ascending and descending limbs of the first peak of the mitotic index curve (Quastler and Sherman, 1959).   |
| <b>G1</b>                | G1-phase was calculated as difference (ie. C-(S+G2+M)).  |

The more rapid rise in the mitotic index in the *AtWEE1* expressing lines is consistent with a much shorter G2 phase compared with the EV (Fig. 3.5). However, M phase and S phase are almost unaltered in the *AtWEE1* lines compared with the empty vector or wild type lines (Fig. 3.5). The overall duration of the cell cycle calculated as the interval between peaks is marginally longer in the *AtWEE1* lines compared with empty vector, by 1-2 hours. Since G1 is obtained by difference, the dramatic shortening of G2 phase in the *AtWEE1* lines is compensated by a long G1 phase (Fig. 3.5).




**Fig. 3.5:** The duration of the cell cycle (C) and its component phases in the constitutive *AtWEE1* cell lines together with the corresponding empty vector following synchronization, wild type is also represented. The duration of the cell cycle is: 14 hours wild type, 12 hours EV-c-10, 12.5 hours WEE1-c-2, 12 hours WEE1-c-10 and 14 hours WEE1-c-12.

Hence, constitutive expression of *AtWEE1* in the tobacco BY-2 cell line clearly results in premature cell division through a shortening of the G2 phase. Remarkably, this is the exact opposite of the expression of *SpWEE1* and *AtWEE1* in fission yeast, where a delay in mitosis is due to a lengthening of the G2 phase. Clearly, constitutive *AtWEE1* expression led to an unusual and unpredicted cell cycle response in the TBY-2 cell line.




### 3.4.4. Can hygromycin affect the cell cycle?

To test whether hygromycin was causing any of the reported cell cycle changes, synchronization experiments were carried out for the WEE1-c-10, WEE1-c-12 and EV-c-10 without addition of hygromycin after aphidicolin was removed. These data were then compared with the data from experiments where hygromycin was added after removal of aphidicolin (Fig. 3.6 a, b and c; Appendix III).




a

Fig. 4b. Mitotic index curves obtained following aphidicolin (100  $\mu$ M) treatment for 24 hours of the



b




c



**Fig. 3.6:** Mitotic index curves obtained following synchronization (with aphidicolin for 24 hours) of the constitutive *AtWEE1* cell lines and empty vector. a) WEE1-c-10  $\pm$ Hyg; b) WEE1-c-12  $\pm$ Hyg; c) EV-c-10  $\pm$ Hyg.

EV-c-10 was also subcultured for four weeks in absence of hygromycin (Fig. 3.7; Appendix III).



**Fig. 3.7:** Mitotic index curve of the empty vector obtained following synchronization after subculturing for four weeks without Hyg.


Overall, there was very little difference in the mitotic index curves  $\pm$ Hyg, WEE1-c-12 was the only line showing a one hour difference between the mitotic peak  $\pm$ Hyg, although the first peak was lower in the +Hyg treatment. This result shows that Hyg treatment does not have an effect on the timing of the component phases of the cell cycle, but it did result in a slight reduction in the percentage of cycling cells.

### 3.4.5. Mitotic cell area is smaller in the constitutive *AtWEE1* expressing lines


The shortened G2 phase obtained in the *AtWEE1* expressing lines would suggest premature cell division at a reduced cell size. This was confirmed when I analyzed mitotic cell area following synchronization in the various lines by image analysis (see chapter 2 section 2.3.4) (Table 3.5, Fig. 3.8 a, b; see also Appendix II for raw data). Thus expression of *AtWEE1* in tobacco resulted in a decrease in mitotic cell size. WEE1-c-10 cells showed the smallest mitotic cell area ( $\sim 2100 \mu\text{m}^2$ ), followed by WEE1-c-2 and WEE1-c-12 ( $\sim 2600 \mu\text{m}^2$ ). Wild type mitotic cell area ( $\sim 3000 \mu\text{m}^2$ ) was bigger than constitutive *AtWEE1* cell lines, but it was smaller than EV-c-10 ( $\sim 3300 \mu\text{m}^2$ ).

**Table 3.5:** The mean mitotic cell area ( $\pm$  standard error (in parenthesis)) of WT, EV-c-10, WEE1-c-2, WEE1-c-10 and WEE1-c-12 cell area, following synchronization. Levels of significance are indicated by Student's t-test.

|                                      | <b>WEE1-c-2</b><br>2598 ( $\pm 76$ ) | <b>WEE1-c-10</b><br>2128 ( $\pm 61$ ) | <b>WEE1-c-12</b><br>2630 ( $\pm 86$ ) |
|--------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------|
| <b>WT</b><br>3034 ( $\pm 58$ )       | P = 0.000                            | P = 0.000                             | P = 0.000                             |
| <b>EV-c-10</b><br>3350 ( $\pm 112$ ) | P = 0.000                            | P = 0.000                             | P = 0.000                             |



a



b


**Fig. 3.8:** a) Mean ( $\pm$  SE) mitotic cell area in the wild type, constitutive *AtWEE1* cell lines and the corresponding EV. Measurements were taken following release from aphidicolin ( $n \geq 150$ ). b) Examples of mitotic cells in wild type (WT), EV-c-10 and one representative *AtWEE1* expressing line (WEE1-c-10) (bars = 100  $\mu$ m).

The effect of hygromycin on mitotic cell area was investigated using EV-c-10 cell line. In the constitutive *AtWEE1* cell lines, mitotic cell area was significantly smaller compared with the EV if hygromycin was present continuously (Table 3.5, Fig. 3.8 a, b; see also Appendix II for raw data). If EV-c-10 was subcultured for 1 week without hygromycin, mitotic cell area was smaller ( $\sim 2500 \mu\text{m}^2$ ) (Table 3.6) compared with the same clone subcultured with the antibiotic ( $\sim 3500 \mu\text{m}^2$ ). However, over 4 weeks of subculture without hygromycin, mitotic cell area became progressively larger ( $\sim 2700 \mu\text{m}^2$ ) (Table 3.6, Fig. 3.9; see Appendix II for raw data), but did not reach the same cell area as the + Hyg culture. To obviate the possibility that small mitotic cell area was due to the lack of hygromycin in the culture medium, hygromycin was added every week to the fresh subculture.

**Table 3.6:** The mean mitotic cell area ( $\pm$  standard error (in parenthesis)) of EV-c-10 subcultured with and without hygromycin over 4 weeks.

| (+) Hyg            | 1 week (-) Hyg     | 2 weeks (-) Hyg   | 3 weeks (-) Hyg   | 4 weeks (-) Hyg   |
|--------------------|--------------------|-------------------|-------------------|-------------------|
| 3537 ( $\pm 103$ ) | 2515 ( $\pm 118$ ) | 2045 ( $\pm 69$ ) | 1916 ( $\pm 81$ ) | 2687 ( $\pm 67$ ) |






**Fig. 3.9:** Mean ( $\pm$  SE) mitotic cell area for EV-c-10 subcultured with hygromycin ((+) Hyg) or without hygromycin ((-) Hyg) ( $n \geq 50$ ) for 1,2 ,3 and 4 weeks.

### 3.5. Results: TBY-2 cells transformed with pTA7002-*AtWEE1*

To examine whether TBY-2 cells “adapted” to the transgene over several generations or whether the changes to the cell cycle were rapid, tobacco BY-2 cells were transformed with *AtWEE1* using a chemical induction system (pTA7002). To induce expression of *AtWEE1*, dexamethasone (DEX) a strong synthetic glucocorticoid is required.

Three independent cell lines carrying *AtWEE1* under the control of the DEX-inducible promoter (Aoyama and Chua., 1997) were investigated in relation to cell size and cell

cycle. In conjunction with the *AtWEE1* expressing lines (denoted as showed in table 3.1) three empty vector lines (see table 3.1) were used as experimental controls. Preliminary experiments on the inducible lines and the empty vector cell lines confirmed that the promoter was working. Dexamethasone (final concentration 1 $\mu$ M) was added to WEE1-i-1, WEE1-i-3 and WEE1-i-6, three days after the fresh subcultures were prepared (i.e. when the cells were in exponential phase) and cells sampled every 12 hours for a 48 hour period. Using RT-PCR, with specific primers to amplify *AtWEE1* (*AtWEE1* forward and reverse primers), the expression of *AtWEE1* at the mRNA level was already detectable within 15 min of the addition of DEX (Fig. 3.10).




**Fig. 3.10:** RT-PCR analysis of *AtWEE1* expression in three independent tobacco BY-2 cell lines induced to express *AtWEE1* (WEE1-i-1, WEE1-i-3 and WEE1-i-6) distilled water (dH<sub>2</sub>O) was used as the control. Cells were induced at time 0 h (within 15 min of the addition of DEX) and sampled every 12 hours for a 48 hour period.

*AtWEE1* expression was also checked by RT-PCR in non-induced pTA7002-*AtWEE1* cell lines in the same experimental conditions except for the absence of the inducer, and in this case *AtWEE1* expression was not detected (Fig 3.11 a). However, to ensure that cDNA was synthesized successfully and the lack of expression of *AtWEE1* was due to the



lack of induction of the pTA7002-*AtWEE1* promoter, the same cDNA was used in PCR with PUV primers directed against *18S rRNA* (Fig 3.11 b). PCR product of the correct size was obtained for all the lines indicating that mRNA extraction and cDNA synthesis were successful.



**Fig. 3.11:** a) cDNA from three independent inducible tobacco BY-2 cell lines without addition of the DEX inducer showing lack of expression of *AtWEE1* compared to the induced WEE1-i-1 cell line (positive control). The arrow is pointing to the *AtWEE1* band, the lower band is primer dimers. b) not induced WEE1-i-1, WEE1-i-3 and WEE1-i-6 cDNA amplified using primers directed against 18S rRNA, WEE1-i-1 induced cDNA was used as a positive control. In both cases the negative control is distilled water.


### 3.5.1. Mitotic cell area in the inducible *AtWEE1* cell lines

Measurements of mitotic cell area were made in tobacco BY-2 cell lines transformed with *AtWEE1* under the control of the inducible promoter. Three days after


the fresh subcultures were made; mitotic cell size was measured every 12 hours for a 48 hour period following induction (see Appendix II for raw data). The 0 h sample for induced pTA7002-*AtWEE1* cells was collected 15 min after addition of DEX. Mitotic cell area for induced lines (WEE1-i-1 and WEE1-i-6) was compared with the same cell lines which had not been induced (Table 3.7, Fig. 3.12 a, b). At 0 h the average mitotic cell area for induced WEE1-i-1 was  $\sim 1200 \mu\text{m}^2$  (Table 3.7), whereas the average mitotic cell area for not induced WEE1-i-1 was  $\sim 2000 \mu\text{m}^2$  (Table 3.7, Fig 3.12 a). Differences in mitotic cell area dimension between treatments were observed at 12, 24, 36 and 48 hours in both WEE1-i-1 and WEE1-i-6 cell lines (Table 3.7, Fig 3.12 a and b). The only exception is WEE1-i-6 induced and non-induced at 36 h that did not show a significant difference between treatments ( $P = 0.643$ ) (Table 3.7, Fig 3.12 a). The image analysis (Table 3.7, Fig. 3.12 a, b) shows clearly that the mitotic cell size is smaller for WEE1-i-1 and WEE1-i-6 induced compared to the non-induced control.

**Table 3.7:** The mean number ( $\pm$  standard error (in parenthesis)) of WEE1-i-1 and WEE1-i-6 cell area, induced and non-induced. Levels of significance are indicated by Student's t-test ( $n \geq 15$ ).

| <b>Time (h)</b> | <b>WEE1-i-1<br/>induced</b> | <b>WEE1-i-1<br/>non-induced</b> | <b>P</b> | <b>WEE1-i-6<br/>induced</b> | <b>WEE1-i-6<br/>non-induced</b> | <b>P</b> |
|-----------------|-----------------------------|---------------------------------|----------|-----------------------------|---------------------------------|----------|
| <b>0</b>        | 1239 ( $\pm 95$ )           | 2048 ( $\pm 82$ )               | = 0.000  | 1575 ( $\pm 66$ )           | 1886 ( $\pm 72$ )               | = 0.000  |
| <b>12</b>       | 1446 ( $\pm 85$ )           | 2085 ( $\pm 59$ )               | = 0.000  | 1627 ( $\pm 101$ )          | 2088 ( $\pm 91$ )               | = 0.000  |
| <b>24</b>       | 1463 ( $\pm 91$ )           | 2069 ( $\pm 86$ )               | = 0.000  | 1667 ( $\pm 73$ )           | 2029 ( $\pm 90$ )               | = 0.001  |
| <b>36</b>       | 1838 ( $\pm 113$ )          | 2271 ( $\pm 60$ )               | = 0.002  | 2029 ( $\pm 512$ )          | 2367 ( $\pm 90$ )               | = 0.643  |
| <b>48</b>       | 1650 ( $\pm 89$ )           | 2613 ( $\pm 75$ )               | = 0.000  | 2167 ( $\pm 171$ )          | 2694 ( $\pm 62$ )               | = 0.023  |



**a**





**b**

### 3.5.2. Can dexamethasone affect cell size?

To test whether DEX was causing tobacco BY-2 cells to divide at smaller cell size, DEX (final concentration 1 $\mu$ M) was added to TBV-2 cells carrying the pTA7002-empty vector three days after the new subculture was made. Measurements of mitotic cell area were taken every 12 hours for a 48 hour period comparing EV-i-1 and EV-i-3 with and without DEX (Table 3.8, Fig. 3.13 a, b; see Appendix II for raw data).

**Table 3.8:** The mean number ( $\pm$  standard error (in parenthesis)) of EV-i-1 and EV-i-3 cell area ( $\pm$ ) DEX. Levels of significance are indicated by Student's t-test ( $n \geq 20$ ).

| Time (h) | EV-i-1<br>(+) DEX | EV-i-1<br>(-) DEX  | P       | EV-i-3<br>(+) DEX  | EV-i-3<br>(-) DEX | P       |
|----------|-------------------|--------------------|---------|--------------------|-------------------|---------|
| 0        | 2068 ( $\pm 81$ ) | 2181 ( $\pm 67$ )  | = 0.285 | 2167 ( $\pm 80$ )  | 2239 ( $\pm 83$ ) | = 0.534 |
| 12       | 2315 ( $\pm 69$ ) | 2499 ( $\pm 90$ )  | = 0.109 | 1829 ( $\pm 58$ )  | 1830 ( $\pm 56$ ) | = 0.990 |
| 24       | 2253 ( $\pm 72$ ) | 2464 ( $\pm 53$ )  | = 0.021 | 1952 ( $\pm 54$ )  | 1939 ( $\pm 71$ ) | = 0.888 |
| 36       | 2348 ( $\pm 88$ ) | 2325 ( $\pm 82$ )  | = 0.846 | 1929 ( $\pm 75$ )  | 2190 ( $\pm 80$ ) | = 0.020 |
| 48       | 2360 ( $\pm 84$ ) | 2344 ( $\pm 101$ ) | = 0.902 | 2267 ( $\pm 125$ ) | 2241 ( $\pm 86$ ) | = 0.862 |




**Fig. 3.13:** Mean mitotic cell area ( $\pm$  SE) in: **a)** EV-i-1 (+) DEX and (-) DEX and **b)** EV-i-3 (+) DEX and (-) dex ( $n \geq 20$ ).




The image analysis showed clearly that the mitotic cell size in empty vector lines is unaffected by DEX.

### **3.5.3. Induction of *AtWEE1* causes tobacco BY-2 cells to divide prematurely**


pTA-7002-*AtWEE1* cell lines 1 and 6 were synchronized (see chapter 2 section 2.3.2) and following the removal of aphidicolin, DEX (final concentration 1  $\mu$ M) was added. Mitotic index was measured every hour for 21-24 hours. Notably, compared with empty vector (EV-i-1) the mitotic index began to rise sooner in the induced *AtWEE1* cell lines (Fig. 3.14; Appendix III). For WEE1-i-1 and WEE1-i-6 the mitotic index rose between 2-3 hours and peaked at 4-5 hours whereas in the empty vector, the mitotic index began to rise at 5 hours and peaked at 8 hours.



a



b




c


Fig. 3.14: Mitotic index following synchronization with aphidicolin in each of the inducible *AtWEE1* cell lines and empty vector. a) WEE1-i-1 induced; b) WEE1-i-6 induced and c) EV-i-1.

To confirm that the earlier rise of mitotic index was a result of the expression of *AtWEE1*, the WEE1-i-1 was synchronized and the mitotic index was measured with and without DEX. The result clearly shows that in WEE1-i-1 minus DEX the mitotic index peaked at 8 h, which was identical to that in the EV cells (Fig. 3.15; Appendix III).






**a**



**b**

**Fig. 3.15:** The mitotic index following synchronization with aphidicolin in **a)** WEE1-i-1 line  $\pm$  DEX and **b)** EV-i-1  $\pm$  DEX.

Using the methods described in section 3.4.3 of this chapter, an estimation of the four phases of the cell cycle was made (Fig. 3.16).



**Fig. 3.16:** The duration (h) of the cell cycle (C) and its component phases in the inducible *AtWEE1* cell lines  $\pm$  DEX together with the wild type and the corresponding empty vector. The cell cycle duration is 14 hours for wild type; 11 hours for EV-i-1; 13 hours for WEE1-i-1 -DEX; 13 hours for WEE1-i-1 +DEX and 10 hours for WEE1-i-6 +DEX.

As in the constitutive WEE1 cell lines, a rapid rise in the mitotic index occurred in the inducible tobacco cell lines, consistent with a shorter G2 phase compared with the controls. However, S and M phases are almost the same in WEE1-i-1  $\pm$  DEX compared with EV-i-1 and wild type. Although the magnitude of G2 phase reduction was not as great as in the constitutive *AtWEE1* lines, these data are also consistent with a shortened G2 and a premature entry of cells into mitosis compared with the controls. However, a compensatory lengthening of G1 phase was only evident in the induced line 1 but not line

6. Line 6 exhibited one of the fastest cell cycles (10 hours) ever recorded for TBV-2 cells in Cardiff.


### 3.5.4. Inducible tobacco BY-2 cells divide at smaller mitotic cell areas

Following release from aphidicolin block, images of mitotic cells induced to express *AtWEE1* were recorded (as described in chapter 2 section 2.3.4). The image analysis revealed that as in the constitutive lines, the expression of *AtWEE1* resulted in a decrease in mitotic cell size in WEE1-i-1 and WEE1-i-6 compared with wild type, empty vector and non-induced cell lines (Table 3.9, Fig. 3.17 a, b; see also Appendix II for raw data). The data clearly show a reduction in mitotic cell area when DEX was added to the inducible lines (Table 3.9).


**Table 3.9:** Comparison of mean mitotic cell area ( $\mu\text{m}^2$ ) ( $\pm$  standard error (in parenthesis)) in WT, EV-i-1, WEE1-c-1 and WEE1-c-6 cell area (induced and non-induced) following a synchronization experiment. Levels of significance are indicated by Student's t-test ( $n \geq 150$ ).

|  | <b>WEE1-i-1<br/>induced</b><br>2395 ( $\pm 58$ ) | <b>WEE1-i-6<br/>induced</b><br>2266 ( $\pm 67$ ) |
|--|--|--|
| <b>WT</b><br>3034 ( $\pm 58$ )                       | P = 0.000  | P = 0.000  |
| <b>EV-c-10</b><br>2803 ( $\pm 156$ )                 | P = 0.015  | P = 0.002  |
| <b>WEE1-i-1<br/>non-induced</b><br>2903 ( $\pm 53$ ) | P = 0.000  |  |
| <b>WEE1-i-6<br/>non-induced</b><br>2831 ( $\pm 71$ ) |  | P = 0.000  |





a



b

**Fig. 3.17** a) Mean ( $\pm$  SE) mitotic cell area in the wild type; EV-i-1; WEE1-i-1  $\pm$ DEX and WEE1-i-6  $\pm$ DEX. Measurements were taken following release from aphidicolin ( $n \geq 150$ ). b) Examples of mitotic cells that were smaller in WEE1-i-1 +DEX compared with the same line -DEX and the empty vector (bars = 100  $\mu$ m).

### 3.6. Discussion

To maintain a specific size, cells must coordinate their growth and division. Yeast cells are thought to use cell-size checkpoints to coordinate these two processes (Nurse, 1975; Johnston *et al.*, 1977; Fantes and Nurse, 1997; Hartwell and Unger, 1997; Rupes, 2002). In fission yeast, the critical size requirement is exerted primarily at the G2/M transition, whereas in budding yeast it is exerted primarily at the G1/S transition (Rupes, 2002). Early work in fission yeast suggested that the WEE1 kinase plays an important role in a checkpoint that coordinates cell growth and cell division at the G2/M transition (Nurse, 1975; Fantes and Nurse, 1978; Thuriaux *et al.*, 1978).

In *S. pombe*, *wee1* is a negative regulator of mitosis and its over expression causes cells to arrest in G2, to grow but not divide resulting in very elongated cells (Russell and Nurse, 1987). *AtWEE1* was used in a functional assay by cloning the ORF into the fission yeast expression vector, pREP1 under the control of the potent thiamine-repressible *nmt1* promoter (Maudrell, 1993; Sorrell *et al.*, 2002). Colony formation by cells containing pREP1-*AtWEE1* was substantially reduced when the promoter was activated by removal of thiamine from the medium. However, the cells containing empty vector formed colonies regardless of thiamine supply. The substantial reduction of colony formation suggests that *AtWEE1* expression was inhibiting cell division. The cells containing pREP1-*AtWEE1* grown in the absence of thiamine were elongated and exhibited a 3.9-fold increase in length compared with empty vector (Sorrel *et al.*, 2002). Notably, this long cell phenotype was also obtained when human and maize homologues were expressed in *S. pombe* and are indicative of G2 arrest (Sun *et al.*, 1999). Hence,



*AtWEE1* exhibits functional properties that are characteristic of WEE1 kinase expression. In *Arabidopsis* plants, over-expression of *AtWEE1* led to the same results observed previously in other organisms; the average of meristematic epidermis cells size was 1.41 fold greater than wild type (Cardiff cell cycle laboratory, unpublished data). Recent work has shown that Swe1, the budding yeast homolog of WEE1, also delays entry into mitosis and is required for cell size control (Jorgensen *et al.*, 2002; Harvey and Kellogg, 2003).

Interestingly, the experimental results reported in this chapter are in contrast to the data reported above and show that constitutive and inducible expression of *AtWEE1* in tobacco BY-2 cells induces a reduction in mitotic cell size. This reduction was observed in the constitutive *AtWEE1* expressing cell line compared with the wild type and the empty vector on day 1 and day 6 during the normal 7 days batch culture. A significant reduction in mitotic cell size was also obtained in three independent constitutive and two independent inducible *AtWEE1* expressing lines following synchronization.

TBY-2 *AtWEE1* expressing cell lines also exhibited unusual cell morphology. Normally, TBY-2 cells are observed to be 2-6 times longer than wide, in *AtWEE1* expressing cell lines, reduction in cell size is mainly achieved by the reduction in cell length, resulting in isodiametric cells.

To confirm that the small mitotic cell size and the abnormal cell cycle was neither a response to Hyg to select transformed tobacco BY-2 cells nor the glucocorticoid used to induce *AtWEE1*, in the inducible cell lines, different experiments were carried out showing that these chemicals did not play any role in causing a small mitotic cell phenotype and a shortened G2 phase.

To obviate the possibility of non-cell cycle effects as a result of the promoter used, tobacco BY-2 cells were also transformed with *AtWEE1* under the control of a dexamethasone inducible promoter/vector system (Aoyama and Chua, 1997). To investigate whether induction of *AtWEE1* expression would also result in small mitotic cell size, I measured mitotic cell size in inducible *AtWEE1* cell lines that were synchronized using aphidicolin. The results showed that induction of *AtWEE1* results in a small mitotic cell area, confirming the results for the constitutive lines.

In both constitutive and inducible *AtWEE1* expressing cell lines, cell size is approximately two-thirds of that observed in wild type and empty vectors. Notably, these data are very similar to those reported by Russell and Nurse (1986): over expression of *SpCdc25* in fission yeast induces fission yeast cells to divide at approximately two-thirds of that observed in wild type. Experiments carried out in our lab have also confirmed that expression of *SpCdc25* in TBY-2 cell lines reduce cell size at division (Orchard *et al.*, 2005). Taken together, these results indicate that cells do not continue to divide at smaller sizes, resulting in mitotic catastrophe, but instead establish a new threshold size for cell division. Although the existence of cell-size checkpoints was proposed over 20 years ago (Fantes and Nurse, 1977, 1978), the underlying molecular mechanisms have remained elusive, and how exactly cells monitor cell size or cell growth is still unknown.

To test how *AtWEE1* expression resulted in small mitotic cell size, cell cycles were assessed by charting the mitotic index following synchronization with aphidicolin. For both the constitutive and the inducible cell lines the mitotic index curve rose sooner and peaked earlier compared with all the controls (wild type, empty vectors and not induced lines). Hence the data are indeed consistent with a short G2 phase. The

shortening of G2 phase fits with both premature cell division and small cell size at division. Interestingly, the shorter G2 phase was compensated by a longer G1 phase in tobacco BY-2 cells expressing *AtWEE1* compared with the controls, so that the total duration of the cell cycle was not shorter than 12 hours. An exception was induced WEE1-i-6 that had a very unusual short cell cycle (10 hours). The data about a compensatory variation in duration of G1 phase are supported by the data reported by Russell and Nurse (1986) who demonstrated that fission yeast exhibited a compensatory decrease in duration of G1 phase due to the longer G2 phase. Compared with the empty vector, cell cycle duration was 0.5 hour longer for WEE1-c-2 and 2 hours longer for WEE1-c-12. Compared with wild type, cell cycle duration in EV-c-10 and EV-i-1 was reduced by 2 hours and 3 hours, respectively. However, G2 is longer in the empty vector compared with *AtWEE1* transformed cell lines, whereas G1 is shorter. Note that the cell cycle was two hours shorter in the EV-i-1 compared with the WEE1-i-1 cell line. The only exception was WEE1-i-6, which was one hour shorter, compared with the empty vector line.

### **3.7. Summary**

The data reported in this chapter are exactly the opposite of what is known about *AtWEE1* expression in *S. pombe* (Sorrell *et al.*, 2002). Expression of *AtWEE1* in tobacco BY-2 cells induces small mitotic cell area and accelerates the entry into mitosis. Premature cell division occurs through a reduction in the duration of G2 phase. However,



cell cycle duration was very similar to that in wild type with a lengthening of G1 phase compensating for the shortening of G2 phase.

**Chapter 4: In tobacco BY-2 cells, induction of *AtWEE1* delays expression of the endogenous *Nicotiana tabacum* *WEE1* (*NtWEE1*) gene**

**4.1. Introduction**

Transgenic techniques have become a powerful tool to address important biological problems in the study of the cell cycle. Many approaches that were impossible to implement by traditional genetics can now be achieved by transgenic techniques, including the introduction of homologous or heterologous genes into cells. The ability to introduce foreign genes into the nuclear genome of plants has provided the methodology to analyse molecular mechanisms that lead to co-ordinate expression of genes (Schell, 1987). It also serves to express alien genes or to modulate the expression of endogenous ones (Sonnewald *et al.*, 1991). Usually, in most experiments, the transgenes are transcribed via a strong promoter, such as the 35S promoter from the cauliflower mosaic virus (CaMV). However, a more flexible gene expression system is needed to extract greater benefits from transgenic technology. Good inducible transcription systems are desired because transgenic cells with an inducible phenotype are as useful as conditional mutants isolated by traditional genetics. Also a regulated promoter is often desirable for example to induce expression at defined time points during the cell cycle, in addition, a tightly repressed promoter is required if the expression of the gene product of interest interferes with the regeneration process.

Several induction systems have been reported and successfully used. One of the most commonly used is the glucocorticoid receptor (GR) which is a member of the family of vertebrate steroid hormone receptors. The GR is not only a receptor molecule but also a transcription factor which in the presence of a glucocorticoid such as dexamethasone, activates transcription from promoters containing glucocorticoid response elements (GREs) (Beato, 1989; Picard, 1993). The GR system was chosen as a good induction system in plant cells because it is simple, and the glucocorticoid itself does not cause any pleiotropic effects in plant cells (Scheda *et al.*, 1991).

My research has focused on the effect of *AtWEE1* on the plant cell cycle and in particular its effect on cell size in BY-2 cells. To obviate the possibility of gene-silencing, a heterologous approach was taken by expressing *AtWEE1* in the tobacco BY-2 cells. Clearly, the induction of a small cell size was surprising, so it was hypothesised that *AtWEE1* perturbed the native tobacco *WEE1* (*NtWEE1*). To test this hypothesis, the pattern and timing of expression of *AtWEE1* and *NtWEE1* genes were studied in synchronized tobacco BY-2 cells in which *AtWEE1* could be induced by dexamethasone. Cells were sampled every hour for a 12-14 hour period (see chapter 2 sections 2.3.2 and 2.3.6). RNA was extracted and cDNA synthesized from each sample (see chapter 2 sections 2.4.1, 2.4.2 and 2.4.3). Expression was estimated in a semi-quantitative manner by measuring the ethidium bromide fluorescence after gel electrophoresis of the products from limited cycle PCRs using specific primer pairs designed from *18S rRNA*, *H4*, *AtWEE1* and *NtWEE1* genes (see chapter 2 sections 2.4.4 and 2.4.5).

Hence the aims of the work reported here were to examine whether expression of *AtWEE1* results in a change in pattern and timing of *NtWEE1* expression during the

tobacco BY-2 cell cycle. To do this, a portion of the *NtWEE1* gene was cloned by RT-PCR using degenerate primers.

## 4.2. Materials and methods

Details of Materials and Methods are fully described in chapter 2 from section 2.3.2 to section 2.4.5.

## 4.3. Results

Following synchronization with and after removal of aphidicolin, samples were collected and RNA extraction was successfully carried out from wild type cells and the *WEE1-i-1* ± DEX cells. This enabled an analysis of *NtWEE1* expression in its native background, or in the presence of *AtWEE1* transcripts. Semi-quantitative RT-PCR was carried out to investigate the expression of *AtWEE1* and *NtWEE1* in the transgenic TBV-2 cell lines. These results were normalized against *18S rRNA*, the expression of *H4* was investigated as a control since histone H4 expression is a sensitive marker of S phase (Sorrell *et al.*, 1999, Herbert *et al.*, 2001).

### 4.3.1. Isolation of a homologue tobacco *WEE1* (*NtWEE1*)

Using RT-PCR and two degenerate primers, it was possible to amplify a 348 bp region of the tobacco *WEE1* (*NtWEE1*) open reading frame. From a comparison of the amino acid sequence of *NtWEE1* with *AtWEE1* and *ZmWEE1* kinase proteins

(Fig. 4.1) the percentage of homology at the amino acid level was 70.8% and 61.8%, respectively, indicating that a tobacco homologue had been amplified.

The *NtWEE1* sequence was then used to design specific primers which were used for semi-quantitative RT-PCR investigation of the native tobacco *WEE1* gene.

|        |   |     |
|--------|---|-----|
| Zmwee1 | -----   | 1   |
| Atwee1 | MF EKNGR TLLAKRKTQGTIKTRASKKIRKMEGTLERHSL LQFGQLSKI SFENRPSSNVA S       | 60  |
| NtWEE1 | -----   | 1   |
| Zmwee1 | -----CTPDYITPEMPQVANEFDDDD  | 21  |
| Atwee1 | SAFQGLLDS SSELRNQLGSADSDANC GEKDFILSQDFFCTPDYITPDNQNLM SGLDIS-          | 119 |
| NtWEE1 | -----   | 1   |
| Zmwee1 | KENIPC PKSPEKSANPRSKRYRTDCSPKAREV TDF SFDHQITPVLFD SLTRDDSEEEQPK        | 81  |
| Atwee1 | KDHSP CPRSPVKLNTV KSKRCRQESFTGNH SNSTWSSKHRVDEQENDD IDTDEVMGDK-L        | 178 |
| NtWEE1 | -----   | 1   |
| Zmwee1 | QPAL EKRGGYVSQSAVALRCRVMP P CVKNPYLNTDPCIDA AVYGGRCNSAVFSP-SIG          | 140 |
| Atwee1 | QANQTERTGYVSQAAVALRCRAM P PCLKNPYVLNQSETATDPFGHQ RSKCASFLPVSTS          | 238 |
| NtWEE1 | -----   | 1   |
| Zmwee1 | GNGLSRYRTDFHEIEKIGYGNF SVFVKV LNRIDGCLYAVKRSIKQLHNDMERRQAVKEVQ          | 200 |
| Atwee1 | GDGLSRYL TDFHEIRQIGAGHFSRVFVKLKRMDGCLYAVKHSTRKLYLDSERRKAMMEVQ           | 298 |
| NtWEE1 | -----   | 1   |
| Zmwee1 | AMAALGSHENIVRYFT <b>ISWFENEOLYIQMELCDRCLSM</b> ---NRNQPVKRGEALEL LYOICK | 257 |
| Atwee1 | AL AALGFHENIVGYYS <b>ISWFENEOLYIQLELCDHSL SALPKKSSLKVSEREILVIMHQIAK</b> | 358 |
| NtWEE1 | ----- <b>ISWFENEHLYIQMELCDHSLSN</b> ---KKY SKLSSEVAVLEAMYQVAK           | 43  |
| Zmwee1 | <b>GLDFMHERGIAHL DVKPDNIYVRNGIYKLGDFGCATLVNRS LATEDGDSRYMPPEMLNDK</b>   | 317 |
| Atwee1 | <b>ALHFVHEKGI AHL DVKPDNIYKNGVCKL GDFGCATRLDKSLPVEEGDARYMPOEILNED</b>   | 418 |
| NtWEE1 | <b>ALQYI HORGVAHL DVKPDNIYVKS EYKLGDFGCATLLDKSQPIEEGDARYMPOEILNEN</b>   | 103 |
| Zmwee1 | <b>YEHLDKVDIFSLGAAVYELIRG TPLPESGSHFTSIREGKIALLP GCPMQFQSLIKSMMDP</b>   | 377 |
| Atwee1 | <b>YEHLDKVDIFSLGVTVYELIKG SPLTESRNQSLNIKEGKL PLLPGHSLQLQLLKTMMDR</b>    | 478 |
| NtWEE1 | <b>YDHLDKVDIFSNH</b> -----  | 116 |
| Zmwee1 | DPVRRPSAKEILRHPSFDK LHKASSK   | 403 |
| Atwee1 | DPKRRPSARELLDHPMFDRIRG----  | 500 |
| NtWEE1 | -----   | 116 |

a

| Percent Similarity |        |        |        |
|--------------------|--------|--------|--------|
|                    | AtWEE1 | ZmWEE1 | NtWEE1 |
| AtWEE1             |        | 62.7   | 70.8   |
| ZmWEE1             | 34.9   |        | 61.8   |
| NtWEE1             | 25.5   | 34.3   |        |

Percent Divergence

**b**

**Fig. 4.1:** a) Alignment of the cloned fragment of NtWEE1 to AtWEE1 and ZmWEE1 protein kinase, showing conserved (black) and similar (grey) residues. b) Percentage identity between the cloned fragment of NtWEE1 and the corresponding regions of AtWEE1 and ZmWEE1 kinase proteins.

#### 4.3.2. Identification of the optimum number of PCR cycles

To estimate the optimum number of cycles to use for semi-quantitative RT-PCR, mix cDNA was made using 1  $\mu$ l of each cDNA sample and RT-PCR was performed. Eight reactions were then set up and every 2 cycles (from 24 to 38) one reaction was stopped. The band strength was measured and plotted against the number of cycles to identify the linear phase of the RT-PCR reaction which is the correct number of cycles to use to ensure that product amount is proportional to starting template amount in the PCR reaction (Fig. 4.2). This method has been used successfully to measure changes in gene expression in several other systems (Wagstaff *et al.*, 2002; Parfitt *et al.*, 2004).




Fig. 4.2: Calibration plot using dilutions of *WEE1-i-1* cDNA to estimate optimum number of cycles for histone H4 primer pair.

**Fig. 4.2:** Example of estimation of optimum cycle number to use for semi-quantitative RT-PCR for histone H4 from *WEE1-i-1* cDNA. Arrow indicates cycle number chosen for further experiments.

#### 4.3.3. Statistical analysis of semi-quantitative RT-PCR results

To ensure that semi-quantitative RT-PCR was performed to produce a linear relationship between product band strength and template concentration, each of the primers for *18S rRNA*, *H4*, *AtWEE1* and *NtWEE1* genes were tested on dilutions of a mix cDNA. These reactions were included in each semi-quantitative RT-PCR experiment performed. An example of a calibration plot using dilutions of the mix cDNA from *WEE1-i-1* induced using an optimum number of cycles for the *H4* primer pair is shown in Figure 4.3.

The ratio of the target gene to *18S rRNA* was calculated for each hour, divided by the corresponding mean for *18S rRNA*. The ratio of the target gene to *18S rRNA* was multiplied by 1000 to provide a more easily readable scale.



#### 4.3.4. Comparison of WEE1 expression in wild-type WEE1 and histone H4




Fig. 4.3: Calibration plots using dilutions of WEE1-i-1 mix cDNA sample and histone H4 primers at an optimum number of cycles.


#### 4.3.3. Statistical measure of semi-quantitative RT-PCR results

To obtain a formal statistical measure of the significance of results obtained by semi-quantitative RT-PCR, means of three replicate PCRs, were used. The means were normalized against two correction methods: assuming constant *18S rRNA* expression and assuming constant total RNA concentration. The results for each putative gene are displayed as histograms (Figs 4.4 to 4.10). Each column in the histogram represents a mean value for the target gene product for each hour, divided by the corresponding mean for *18S rRNA*. The ratio of the target gene to *18S rRNA* was multiplied by 1000 to provide a more easily readable scale.

#### **4.3.4. Comparison of *NtWEE1* expression in wild type, WEE1-i-1 non-induced and induced.**

##### **4.3.4.1. Expression of *NtWEE1* in wild type cells during the cell cycle**

The pattern of expression for *histone H4* in wild type TBY-2 cells is high during S phase with a peak of expression at 0 hours (corresponding approximately to 15 minutes after removal of aphidicolin). Its level of expression fell away 3-4 hours following the release from aphidicolin (Fig.4.4; see Appendix IV for raw data). This result fits well with the measure of S phase (3.5 hours) obtained from the mitotic index curve (calculated as described in chapter 3 section 3.4.3). Notably, the pattern of *histone H4* expression is similar to that previously reported in synchronized tobacco BY-2 cells using northern blot analysis which also showed an S phase of 3.5 hours (Herbert *et al.*, 2001).




**Fig. 4.4:** Semi-quantitative RT-PCR analysis of *H4* expression (normalised mean  $\pm$  standard error) during the cell cycle of synchronized wild type TBY-2 cells following release from aphidicolin. The duration of the cell cycle component phases are reported below the histogram (see chapter 3 section 3.4.3) (n=3).

The level of *NtWEE1* expression fluctuated during the cell cycle (Fig. 4.5; see Appendix IV for raw data). Maximum expression was observed at 4 hours after removal of aphidicolin (corresponding to the transition point between S and G2 phase), its expression then decreased to reach its minimal level in mid G2 phase. Low levels of *NtWEE1* expression were also detected throughout mitosis and early G1 phase. In late G1 phase of the subsequent cell cycle *NtWEE1* expression rose again.

in agreement with the measure of S phase (3 hours) obtained from the mitotic index curve (calculated as described in chapter 3 section 3.4.3, see also section 3.3.3).






**Fig. 4.5:** Semi-quantitative RT-PCR analysis of *NtWEE1* expression (normalised mean  $\pm$  standard error) during the cell cycle of synchronized wild type TBY-2 cells following release from aphidicolin. Below the histogram is reported the duration of the TBY-2 cell cycle phases derived from the mitotic index (see chapter 3 section 3.4.3) (n=3).

#### 4.3.4.2. Expression of *NtWEE1* in non induced WEE1-i-1 cells during the cell cycle

Expression of *H4* in non induced WEE1-i-1 cells rose to a maximum 1 to 2 hours following the release from aphidicolin (Fig. 4.6; see Appendix IV for raw data). The expression level decreased after 3 hours when the cells were going into G2-phase in agreement with the measure of S phase (3 hours) obtained from the mitotic index curve (calculated as described in chapter 3 section 3.4.3, see also section 3.5.3).



**Fig. 4.6:** Semi-quantitative RT-PCR analysis of *H4* expression (normalised mean  $\pm$  standard error) during the cell cycle in synchronized non induced WEE1-i-1 cells following release from aphidicolin. The duration of the cell cycle component phases derived from the mitotic index are reported below the histogram (see chapter 3 section 3.5.3) (n=3).

The pattern of expression of *NtWEE1* for non induced WEE1-i-1 was very similar to that in wild type tobacco BY-2 cells with a maximum of expression in S-phase; the only difference is that the peak of *NtWEE1* expression in wild type was at hour 4 (Fig. 4.5) whereas the peak of expression for *NtWEE1* in non induced WEE1-i-1 was at hour 3 (Fig. 4.7; see Appendix IV for raw data). Nevertheless, hour 3 represents the border point between S and G2 phase, as was found in the wild type TBV-2 cells.







Fig. 4.7: Semi-quantitative RT-PCR analysis of *NtWEE1* expression (normalised mean  $\pm$  standard error) during the cell cycle in synchronized non induced WEE1-i-1 cells following release from aphidicolin. The duration of the cell cycle component phases derived from the mitotic index are reported below the histogram (see chapter 3 section 3.5.3) (n=3).

#### 4.3.4.3. Expression of *NtWEE1* and *AtWEE1* in induced WEE1-i-1 cells during the cell cycle

The pattern of expression for *H4* in the induced WEE1-i-1 cell line is high during S phase with a peak at 1 hour following the release from aphidicolin. Its level of expression decreases by  $\sim 35\%$  at 4 hours following the release from aphidicolin and remains low thereafter (Fig. 4.8; see Appendix IV for raw data). This result fits well with the measure of S phase (3 hours) obtained from the mitotic index curve (calculated as described in chapter 3 section 3.4.3, see also section 3.5.3).



**Fig. 4.8:** Semi-quantitative RT-PCR analysis of *H4* expression (normalised mean  $\pm$  standard error) during the cell cycle in synchronized induced WEE-1-i-1 cells following release from aphidicolin. The duration of the cell cycle component phases derived from the mitotic index are reported below the histogram (see chapter 3 section 3.5.3) (n=3).

However, the pattern of expression of *NtWEE1* in induced WEE1-i-1 cells was different from the expression of the same gene in wild type and non induced WEE1-i-1 cells. In contrast to these controls, the expression of *NtWEE1* in induced WEE1-i-1 began to rise in late G2 phase and peaked in the middle of M-phase (at 6 hours following release from aphidicolin) decreasing again in late mitosis (Fig.4.9; see Appendix IV for raw data).







Fig. 4.9: Semi-quantitative RT-PCR analysis of *NtWEE1* expression (normalised mean  $\pm$  standard error) during the cell cycle in synchronized and induced WEE-1-i-1 cells following release from aphidicolin. The duration of the component phases derived from the mitotic index are reported below the histogram (see chapter 3 section 3.5.3) (n=3).

Induction of the pTA7002 promoter resulted in an increased level of *AtWEE1* at 1 hour following release from aphidicolin with a transcriptional peak at 3 h which, equates with late S-phase/beginning of G2 phase. In G2, M and G1 phases the expression of *AtWEE1* decreased to a more or less constant level. The expression of *AtWEE1* increased again in G1/S-phase of the next cell cycle (Fig. 4.10, see Appendix IV for raw data).



**Fig. 4.10:** Semi-quantitative RT-PCR analysis of *AtWEE1* expression (normalised mean  $\pm$  standard error) during the cell cycle in synchronized induced WEE-1-i-1 cells following release from aphidicolin. The duration of the cell cycle component phases derived from the mitotic index are reported below the histogram (see chapter 3 section 3.5.3) (n=3).

#### 4.4. Discussion

Calibration plots were used to demonstrate that a linear relationship exists between agarose gel band strengths of semi quantitative RT-PCR products and starting cDNA concentrations. This is important since PCR reactions need to be stopped during the linear phase of amplification if the assumption is made that PCR product yield is proportional to starting cDNA concentration. Normalization against a standard such as *18S rRNA* is also important to ensure that variations in total levels of cDNA in the different samples is taken into account when estimating the relative amounts of a particular message in different samples.

The pattern of expression of *histone H4* which typically peaks in S-phase was investigated for all cell lines studied, as a control for the quality of cDNA synthesized and for the method of normalization used. For all the cell lines investigated, the data reported in this chapter for the expression of *histone H4* show clearly that the level of expression fell in S-phase. These data are remarkably similar to those previously published on synchronized TBY-2 cells (Sorrell *et al.*, 1999; Herbert *et al.*, 2001).

PCR primers, specific for *NtWEE1* were used to analyse the expression of this gene in synchronised wild type, WEE1-i-1 non induced and induced cell lines. Expression of *NtWEE1* in both wild type and non induced WEE1-i-1 peaked during S-phase or S/G2 boundary. Notably, the data related to *NtWEE1* expression in wild type and WEE1-i-1 non induced are in complete agreement with each other. These data are also in complete agreement with the recently published analysis of *NtWEE1* expression during the TBY-2 cell cycle by Gonzalez and colleagues (2004). In their studies the authors investigated by semi-quantitative RT-PCR the pattern of expression of *NtWEE1*, and showed that *NtWEE1* was also strongly expressed during the S-phase and its expression decreased during mitosis and early G1-phase (Gonzalez *et al.*, 2004).

In molecular terms, the premature entry into mitosis at reduced cell size, observed for TBY-2 cell lines expressing *AtWEE1*, could be regulated at the transcriptional, translational or protein level (or all three). My hypothesis was that transcription of the *Nicotiana tabacum WEE1* was inhibited or suppressed in the presence of *AtWEE1* transcripts. Remarkably, when *AtWEE1* was induced in WEE1-i-1 cell line the major peak of *NtWEE1* expression was shifted by 4 hours from S phase to middle M phase. The delayed increase in the level of expression of *NtWEE1* in the induced WEE1-i-1 cell line, compared to non-induced and wild type,

did not confirm the initial hypothesis that transcription of the *Nicotiana tabacum* *WEE1* was inhibited. The timing of its expression was perturbed but not inhibited in the presence of *AtWEE1* transcripts so that a negative regulation of CDKs, normally imposed by NtWEE1, could be removed, enabling *AtWEE1*-expressing cells to enter into mitosis at a smaller cell size. Loss of *wee1* activity in fission yeast causes cells to enter mitosis before sufficient growth has occurred and therefore results in two abnormally small daughter cells (Nurse, 1975). Conversely, increasing the gene dosage of *wee1* causes delayed entry into mitosis and an increase in cell size, indicating that the levels of Wee1 kinase activity determine the timing of entry into mitosis and can have major effects on cell size (Russell and Nurse, 1987). Recent work has shown that Swe1, the budding yeast homologue of WEE1, also delays entry into mitosis and is required for cell size control (Jorgensen *et al.*, 2002; Harvey and Kellog, 2003). *Swe1* $\Delta$  cells undergo premature entry into mitosis before sufficient growth of the daughter bud has occurred; producing abnormally small cells (Harvey and Kellogg, 2003). Loss of function of WEE1 kinases in *Xenopus* and *Drosophila* also causes premature entry into mitosis; however, a requirement for WEE1 itself in cell size control has not yet demonstrated in animal cells (Walter *et al.*, 2000).

In addition, PCR primers specific for *AtWEE1* enabled comparative measurements of *AtWEE1* and *NtWEE1* expression in the inducible WEE1-i-1 cell line. Hence this experimental design enabled a comparison of *NtWEE1* expression alone or in the presence of *AtWEE1* transcripts in the different genetic backgrounds of wild type, induced and non-induced WEE1-i-1. Induction of *AtWEE1* resulted in a transcriptional peak of *AtWEE1* at 3 hours which corresponds to late S-phase/beginning of G2 phase. Transcription of the transgene depends on the activation of the promoter used in the construct. In the experiments reported in this

chapter the promoter was activated by the addition of DEX soon after removal of aphidicolin when the cells were entering into S phase (Aoyama and Chua 1997). Note that the transcriptional peak for *AtWEE1* occurred at the same time as the *NtWEE1* peak in non-induced WEE1-i-1. Thus, these data suggested that rather than a transcriptional inhibition of *NtWEE1 per se*, *AtWEE1* transcript affected the timing of *NtWEE1* transcription. The *AtWEE1* data suggest that replacement of *NtWEE1* expression in S phase by *AtWEE1* expression is insufficient to regulate cell size at division, suggesting that *AtWEE1* is not fully recognized in the tobacco cells.

#### **4.5. Summary**

Expression of *AtWEE1* in the induced WEE1-i-1 TBY-2 cell line results in an alteration of the timing of the endogenous *WEE1* gene during the cell cycle. In wild type the expression profile of *NtWEE1* revealed a major peak of transcription in late S-phase/beginning of G2-phase. In the non-induced *AtWEE1* cell line, *NtWEE1* transcript also peaked in mid-to-late S-phase. Remarkably, when *AtWEE1* was induced, the major peak of *NtWEE1* expression was shifted to M phase. Induction of *AtWEE1* resulted in a transcriptional peak at late S-phase.

## **Chapter 5: CDKA and CDKB1 kinase activity through the cell cycle in synchronized TBY-2 cells expressing *AtWEE1***

### **5.1. Introduction**

The events of the cell cycle usually occur in a fixed sequence, and if an early event such as S-phase is incomplete then a later event such as mitosis becomes blocked. There are two general types of mechanism that could link S-phase and mitosis. There could be a hard wiring of the two events such that the later event is unable to occur physically or chemically without the early event. Alternatively the two events could be linked by a regulatory loop that inhibits the second until the first is complete. Hartwell and Weinert (1989) developed the second of these two events into the idea of checkpoints where the cell monitors or “checks” cell cycle progression at certain points during the cell cycle, and if the events prior to that point are incomplete, further progression is delayed (Hartwell and Weinert, 1989). Cyclin dependent kinases (CDKs) are recognized as key players in checkpoint controls of the eukaryotic cell cycle. One of the most important roles for CDKs is to ensure that there is only one S phase in the normal cell cycle. When a proliferative cell completes S phase and enters G2, another S phase does not take place until the mitosis of that cell cycle is complete (Broeck *et al.*, 1991).

In fission yeast, there is a dependency of mitosis upon completion of S phase, this dependency is lost in cells with specific *cdc2* mutations or in mutants with a high level of

the CDK activator, Cdc25 (Enoch and Nurse, 1990) (for more details see chapter 1 section 1.4).

In contrast to the yeast cell which has only one type of CDK gene (*CDC28* for *Saccharomyces cerevisiae* and *cdc2* for *Schizosaccharomyces pombe*), higher eukaryotic cells have many different types of CDKs (for more details see chapter 1 section 1.1.2). Recently, the tobacco BY-2 cell line was used for studies of plant CDKs and their kinase activity during the cell cycle. A cDNA of a PSTAIRE (CDKA) gene *cdc2Nt1* (renamed Nicta;*CdkA*;3 by Joubes *et al.*, 2000) has been cloned from tobacco, and RNA gel-blot analysis showed this gene to be preferentially expressed in dividing TBV-2 cells but did not show significant cell cycle regulation of transcript abundance (Setiady *et al.*, 1996). Sorrell *et al.* (2001) reported the isolation, from a TBV-2 cell cDNA library, of a PSTAIRE CDKA (Nicta;*CdkA*;4) closely related to *cdc2Nt1* (Setiady *et al.*, 1996) and to other novel tobacco CDKs (Nicta;*CdkB1*;1 and Nicta;*CdkB1*;2) containing the PPTALRE sequence and belonging to the CDK-b1 subgroup. Sorrell *et al.* (2001) demonstrated that CDKA and CDKB1 have a different level of expression during the tobacco BY-2 cell cycle. *CDKA* transcript levels remained at an approximately constant level during the cell cycle, whereas *CDKB1* transcript levels varied with the lowest levels in G1 and the highest levels in S, G2 and M phases (Sorrell *et al.*, 2001). *CDKA* protein levels, like the RNA levels, were relatively constant throughout the cell cycle except for a decline during G1 phase, but the *CDKA* kinase activity was higher during the S and G2 phases of the cell cycle (Sorrell *et al.*, 2001). *CDKB1* protein levels showed a gradual accumulation from S phase until mitosis, followed by a gradual decline, but the *CDKB1* kinase activity showed a sharp peak at the G2/M boundary, followed by an abrupt decline



(Sorrell *et al.*, 2001). These data suggested a differential role for CDKB1 relative to CDKA during the tobacco cell cycle (Sorrell *et al.*, 2001).

In fission yeast, negative (Wee1/Mik1) and positive (Cdc25) regulation of Cdc2 interact to regulate cell size at mitosis (for more details see chapter 1 section 1.4). In data reported in chapter 3, *AtWEE1* expression induced a small cell size in TBY-2 cells. This was surprising given that *AtWEE1* expression induced a long cell length in fission yeast (Sorrell *et al.*, 2002). The *AtWEE1* induction of a small cell size in TBY-2 cells (for more details see chapter 3) was a result of a premature onset of mitosis through a shortening of G2 phase. The data led to the hypothesis that the biochemical basis of this response would be increased CDKA or CDKB (or both) kinase activity, to drive cells into an “early” mitosis.


The aims of the work reported in this chapter were to examine whether expression of *AtWEE1* in transformed TBY-2 cells resulted in a change in CDKA and CDKB1 kinase activity during the cell cycle.

## **5.2. Materials and methods**

### **5.2.1. Protein extraction**

Following synchronization and release from aphidicolin, 5 ml of TBY-2 cell cultures were collected into polypropylene tubes (Falcon) every hour for a 12 h period. The samples were centrifuged at 3500 rpm (MSE Centaur 2) for 5 minutes to form a pellet. The supernatant was discarded and the pellet ground to a fine powder in liquid

nitrogen using a pestle and a mortar precooled at  $-20^{\circ}\text{C}$ . Powdered cells were transferred to 1.5 ml microcentrifuge tube and resuspended in 1 ml of lysis buffer (0.5 M Tris-HCl pH 7.5, 0.75 M NaCl, 0.15 M  $\text{MgCl}_2$ , 1 M dithiothreitol, 0.1% Tween 20, 1x complete Tm protease inhibitors (Roche), 50 mM NaF, 10 mM NaV, 100 mM Na-pyrophosphate, 0.6 M  $\beta$ -glycerophosphate) on ice. The suspension was homogenized by vortexing for 4 x 30 seconds with 1 min on ice between homogenizations and centrifuged at 13000 rpm (Beckman Coulter Allegra <sup>TM</sup> 21R, rotor F2402H) for 30 minutes at  $4^{\circ}\text{C}$ ; the supernatant was transferred to a new microcentrifuge tube on ice. In order to measure the concentration of total proteins extracted using a spectrophotometer  $\lambda=595\text{nm}$  (Smartspec<sup>TM</sup>3000, Bio-Rad laboratories Ltd) one  $\mu\text{l}$  of extract was transferred to 1 ml Coomassie<sup>®</sup> Protein Assay Reagent (Bradford assay; Sigma) and the absorbance was read against a standard curve (Fig. 2.5). Samples were stored at  $-80^{\circ}\text{C}$  until further use.



**Fig. 2.5:** Standard curve for proteins concentration measured against absorbance using known amounts of bovine serum albumen (BSA).

### 5.2. 2. Kinase Activity

For immunoprecipitations and kinase assays, 250 µg of proteins extracted from TBY-2 cell cultures were pre-incubated with 20 µl of protein A-Sepharose and an appropriate volume of lysis buffer to a final volume of 250 µl. The proteins were rotated for 30 min at 4°C and centrifuged at 13000 rpm (Beckman Coulter Allegra™ 21R, rotor F2402H) for 10 minutes at 4°C. The supernatant was transferred into a new microcentrifuge tube and incubated with 2 µl of specific antiserum (NtCDKA or NtCDKB) for 2 h on ice. Twenty µl of protein A-Sepharose and 150 µl of lysis buffer were added to a final volume of 400 µl. The samples were rotated at 4°C for 1 h and centrifuged at 3000 rpm (Beckman Coulter Allegra™ 21R, rotor F2402H) for 1 minute. The pellet was washed 4 times with 1 ml of Suc1-buffer (0.5 M Tris-HCl pH=7.5; 0.75 M NaCl; 0.5 M EDTA; 0.5 M NaF; 0.1% Tween 20) and 2 times with 1 ml of kinase buffer (0.5M Tris-HCl pH=7.5; 0.75 M NaCl; 0.15 M EGTA; 0.1 M DTT); the pellet was centrifuged each time at 3000 rpm for 1 minute using a MiniSpin (Eppendorf).

The pellet was resuspended in 15 µl assay buffer (0.5 M Tris-HCl pH 7.5, 0.75 M NaCl, 0.15 M EGTA, 0.15 M MgCl<sub>2</sub>, 1 M dithiothreitol, 50 mM NaF, 10 mM NaV, 100 mM Na-pyrophosphate, 0.6 M β-glycerophosphate, 5 µg/µl histone H1 and 74 kBq (γ-P<sup>32</sup>) ATP per 15 µl reaction), and incubated at room temperature for 30 min. The reaction was stopped by adding 2x Laemmli buffer (20% glycerol; 8% β-mercaptoethanol; 3.2 % SDS; 0.1 M Tris-HCl pH=6.8; 0.0125% bromophenol blue). Probes were loaded on a 10% SDS-PAGE minigel composed of two phases: 1) resolving gel (where the proteins are resolved: 1.25 ml 10% acrylamide; 0.81 ml resolving gel buffer (1.5 M Tris-HCl pH=8.8;

0.4% SDS); 1.61 ml UHP water; 37.5 µl 12% ammonium persulphate; 3.75 µl TEMED) and 2) stacking gel (where samples are loaded: 0.325 ml 10% acrylamid; 0.5 ml stacking buffer (0.5 M Tris-HCl pH=6.8; 0.4% SDS); 2 ml UHP water; 12.5 µl 12% ammonium persulphate; 2.5 µl TEMED). Samples were run at 150 mV with 1x running buffer (1.5% Tris-base; 7.2% glycine; 0.5% SDS) until the dye front was running over the bottom of the gel. For detection of CDK activity the SDS-PAGE gel was wrapped in Saran Wrap™ (DOW Chemical Company) and exposed to radiograph film (Hyperfilm ECL; Amersham) in the dark room. The orientation of the film compared to the SDS-PAGE gel was marked. After exposure, the film was developed and fixed using the developer machine Curix 60 (Agfa). Samples were analyzed and quantified using a Gene Genius Bioimaging System and Gene Tools software package (Syngene Ltd.).

Since the total number of samples was 11 for non-induced WEE1-i-1, and the SDS-gel contained only eight lanes, it was necessary to run samples on different SDS-gels. However, each sampling time comprised at least two replicates.

## **5.3. Results**


### **5.3.1. Kinase activity of CDKs in TBY-2 cells expressing *AtWEE1***

CDKA and CDKB1 kinase activity were investigated during the cell cycle in synchronously cycling WEE1-i-1 induced and non-induced cells. The cells were synchronized by blocking the cell cycle in early S-phase with aphidicolin. After the aphidicolin block was released, dexamethasone (DEX) was added only to one flask of the

WEE1-i-1 cell line to induce the expression of *AtWEE1*, whereas a second culture of WEE1-i-1 was not induced and served as the control. The progress of cells through the cell cycle was followed by monitoring changes in mitotic index, and cell samples were harvested every hour for 13 hours.

#### **5.3.1.1. Kinase activity of CDKA and CDKB1 in non-induced WEE1-i-1 cells**


Following a synchronization experiment, proteins were extracted from non-induced WEE1-i-1 cells choosing specific time points during the cell cycle (1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 12 hours). Based on the mitotic index curve, these sampling times represented S, G2, M and G1-phase of the non-induced *AtWEE1* line cell cycle (Fig. 5.1).



**Fig. 5.1:** Mitotic index curve, of synchronized non-induced WEE1-i-1 cells, generated from the same experiment used to collect cell samples for protein extractions used for CDK kinase activity assays. The corresponding cell cycle component phases are shown below the graph.

The level of CDKA kinase activity was maximal 1 hour following the release from aphidicolin (early S-phase, Fig. 5.2; see Appendix V for raw data). Subsequently its activity dropped at S/G2 but never fell below 50% compared with the maximal kinase activity level. However, CDKA kinase activity then increased in late G2/beginning of M-phase.






**Fig. 5.2:** Mean ( $\pm$  S.E.) of CDKA kinase activity in synchronized non-induced WEE1-i-1 cells as percentages of maximum activity ( $n \geq 2$ ). The corresponding cell cycle component phases are shown below the graph.

In contrast to CDKA, the kinase activity of CDKB1 peaked at 7 hours (late G2 phase) (Fig. 5.3; see Appendix V for raw data); it started to increase in early G2-phase and peaked in the late G2-phase. The activity then decreased during mitosis and the following G1 phase back to 50% of maximal levels of activity. This pattern of CDKB1 activity is very similar to that for wild type cells (Sorrell *et al.*, 2001).






**Fig.5.3:** Mean ( $\pm$  S.E.) of CDKB1 kinase activity in synchronized non-induced WEE1-i-1 cells as percentages of maximum activity ( $n \geq 2$ ). The corresponding cell cycle component phases are shown below the graph.

#### 5.3.1.2. CDKA and CDKB1 kinase activity in induced WEE1-i-1 cells


Following synchronization, *AtWEE1* expression was induced in the WEE1-i-1 cell line by adding dexamethasone immediately after removal of aphidicolin. Every hour for 13 hours, cell samples were harvested and proteins extracted at specific time points during the cell cycle (1, 3, 4, 5, 6, 7 and 9 hours). Based on the mitotic index curve, these samples represented S, G2, M and G1-phase of the induced *AtWEE1* line cell cycle (Fig. 5.4).



**Fig. 5.4:** Mitotic index curve, of synchronized induced WEE1-i-1 cells, generated from the same experiment made to collect cell samples for protein extractions used for CDK kinase activity assays. The corresponding cell cycle component phases are shown below the graph.


The highest CDKA kinase activity for induced WEE1-i-1 cells was 1 hour following the release from aphidicolin which corresponds to early S-phase (Fig. 5.5; see Appendix V for raw data). Subsequently its activity dropped reaching a more or less constant level during S and G2 phase, without falling significantly below 60% compared to the maximum activity level in S phase.





**Fig.5.5:** Mean ( $\pm$  S.E.) of CDKA kinase activity in synchronized induced WEE1-i-1 cells as percentages of maximum activity ( $n \geq 2$ ). The corresponding cell cycle component phases are shown below the graph.

CDKB1 kinase activity showed a very similar pattern to that of CDKA (Fig 5.5). CDKB1 was, indeed, very high 1 hour following the release from aphidicolin. Its activity, like CDKA, then dropped in mid S-phase and remained low at 50-60% of maximal level in late S, G2 and M phase (Fig. 5.6; see Appendix V for raw data).



**Fig.5.6:** Mean ( $\pm$  S.E.) of CDKB1 kinase activity in synchronized induced WEE1-i-1 cells as percentages of maximum activity ( $n \geq 2$ ). The corresponding cell cycle component phases are shown below the graph.

#### 5.4. Discussion

Previous publications (Mironov *et al.*, 1999; Joubes *et al.*, 2000) have reported CDK kinase activity in plant cells during the cell cycle. According to these publications CDKA kinase activity starts to rise in late G1 phase and peaks at the G1/S border (Table 5.1). In late S phase the level of CDKA activity decreases until late G2 phase and a second peak occurs in mitosis (Table 5.1). Interestingly, tobacco CDKB1 kinase has a constantly flat level of activity until late G2 phase then its activity starts to increase peaking in mitosis (Table 5.1). Recently, Sorrell *et al.* (2001) carried out the first analysis

of tobacco CDKB1 expression, protein abundance, and kinase activity during the TBV-2 cell cycle. CDKB1 is indicated as a key regulator of mitosis, also they confirmed the potential role of CDKA kinase activity in the G2/M transition. Sorrell *et al.* (2001) proposed that the functions of CDKA and CDKB1 may be distinct due to the different timing of their kinase activity. They also reported that in synchronous TBV-2 cells CDKA activity is relatively constant throughout the cell cycle except for a slight decline during G1 phase, whereas CDKB1 kinase activity showed a sharp peak at the G2/M boundary (Table 5.1).

Since CDK kinase activity is negatively regulated by Wee1 kinase (Russell and Nurse, 1987), it was interesting to investigate if expression of *AtWEE1* in tobacco BY-2 cell lines altered the normal level of CDK kinase activity.

Previous experiments (described in chapter 4) had shown that expression of *AtWEE1* in the tobacco BY-2 cell line results in a delayed expression of the endogenous *NtWEE1* gene. This might enable tobacco CDKs to exhibit an earlier kinase activity. This hypothesis was tested by measuring kinase activity of both CDKA and CDKB1 in the induced WEE1-i-1 cell line. As a control CDKA and CDKB1 kinase activity were tested in non-induced WEE1-i-1 cell line.

In the induced WEE1-i-1 cell line both CDKA and CDKB1 kinase activity peaked 1 hour after removal of aphidicolin which corresponds to early S phase (Figs 5.5 and 5.6) (Table 5.1). Subsequently their activities dropped at the S/G2 border, but never went below 50% compared to maximum. In non-induced WEE1-i-1 cell line, CDKA kinase activity, in agreement with Mironov *et al.* (1999) and Joubes *et al.* (2000), was high at 1 hour following release from aphidicolin (early S phase) and exhibited a second peak at

the G2/M boundary (Table 5.1). However, CDKB1 kinase activity was low from S phase to the beginning of G2 phase, where its level of activity started to increase until it peaked at 7 hour after removal of aphidicolin, which corresponds to late G2 phase (Figs 5.2 and 5.3) (Table 5.1). These data are remarkably similar to those from Sorrell *et al.* (2001) who also reported a peak of CDKB1 kinase activity at G2 phase. These results demonstrate clearly that expression of *AtWEE1* delays expression of *NtWEE1*, but does not affect CDKA kinase activity, in the induced WEE1-i-1 cell line. However, the expression of *AtWEE1* had a profound effect on CDKB1 kinase activity which peaks in M phase for non-induced WEE1-i-1 but in early S phase for induced WEE1-i-1 cells.

In tobacco BY-2 cells expressing *AtWEE1*, the earlier entry into mitosis can be interpreted as the response to an earlier activation of CDKB1 in the cell cycle. Interestingly, expression of *Spdc25* in the tobacco BY-2 cell line lead tobacco cells to divide prematurely as a direct result of increased Cdc25 phosphatase activity. This resulted in a higher activity of CDKB1 at earlier samples time compared with the control (empty vector). CDKB1 kinase activity was consistently high in S and early G2 phase (Orchard *et al.*, 2005).

As already reported, CDKA and CDKB1 have distinct roles during the cell cycle due to the different timing of their activity (Sorrell *et al.*, 2001). CDKB1 is thought to be more important at G2/M transition (Sorrell *et al.*, 2001). The data reported in this chapter confirm an involvement of CDKB1 in regulating entry into mitosis.



**Table 5.1:** Summary of peaks of CDKs activity in the data presented in this chapter compared with published data in plants. Note the premature CDKB1 activity in the transgene expressing lines.

| CDKA                                       |      | CDKB1 |      | Source  |
|--|------|-------|------|---|
| G1/S                                       | G2/M | G1/S  | G2/M |   |
| +  | +    | -     | +    | Plants<br>(Mironov <i>et al.</i> ,<br>1999; Joubes <i>et al.</i> ,<br>2000) |
| <b>Constant through<br/>the cell cycle</b> |      | -     | +    | Tobacco BY-2 cells<br>(Sorrell <i>et al.</i> ,<br>2001)                     |
| +  | -    | +     | -    | WEE1-i-1 ind cells  |
| +  | +    | -     | +    | WEE1-i-1 non-ind<br>cells   |

## 5.5. Summary

Expression of *AtWEE1* in tobacco BY-2 cells results in an alteration of the timing of *NtWEE1* expression, and as a consequence unlocks the negative regulation activity normally imposed by *NtWEE1* kinase on CDKB1. The net result is a premature increase in CDKB1 kinase activity.



## **Chapter 6: Over expression of *AtWEE1* alters root growth and development in *Arabidopsis thaliana***

### **6.1. Introduction**

Experimental evidence suggests that regulatory proteins such as WEE1 play a very important role during the cell cycle and they might have an important part in regulating plant growth and development. Altered cell cycle duration can influence growth in a direct manner. For example, in plant roots cell division takes place in the root apical meristem (RAM). Cell division in the RAM contributes to primary root growth by providing the root with the new cells it needs in order to grow. In theory, the faster the rate of cell division, the faster the rate of primary root elongation. Investigations into the primary root growth rates of several ecotypes of *Arabidopsis* confirmed this hypothesis: the ecotypes with the fastest growth rates exhibited short cell cycle duration times and elevated rates of cell division (Beemster *et al.*, 2002). In addition, CDKs and other cell cycle control factors such as cyclins are clearly capable of altering plant growth rates by modulating the cell cycle duration (Doerner *et al.*, 1996; Doonan, 1996; Cockcroft *et al.*, 2000; De Veylder *et al.*, 2001a-b; Beemster *et al.*, 2002). However, cell elongation is also very important in controlling plant growth rate; both elongation and cell division make equally important contributions to growth rate (Beemster *et al.*, 2000, 2003). The cell cycle is not only important in controlling growth, it may also play a complex role in the control of plant organ development. For example, lateral roots form from the pericycle, the outer-most layer of the vascular cylinder. Clusters of pericycle cells, act as founder cells of lateral root morphogenesis. In order for primordia to form, the founder cells of the pericycle must divide first transversely and then longitudinally (Dubrovsky *et al.*, 2001). Thus, the


formation of root organs is highly dependent on the rate, site and plane of cell division (Meyerowitz, 1997). Interference with cell cycle control often leads to changes in plant development. The over-expression of ICK1 (a CDK inhibitor) in *Arabidopsis* altered root, leaf and petal shape (Wang *et al.*, 2003). In addition, the expression of the fission yeast homologue to *cdc25* (*Spcdc25*) in tobacco plants led to small mitotic cell size and an increased frequency in lateral root initiation. In these experiments, the expression of *Spcdc25* led to an increase in the number of lateral roots formed per unit length of primary root (McKibbin *et al.*, 1998). *Spcdc25* expression is also capable of stimulating tobacco shoot development, even in the absence of the plant growth regulators and in high auxin and cytokinin treatment that should favour root formation (Suchomelova *et al.*, 2004). Control genes such as *AtWEE1* may also have an important role to play in the development of plant organs.

The aims of the work reported in this chapter, were to check the possible role of *AtWEE1* in the control of primary root growth and lateral root initiation in *Arabidopsis* plants.

## 6.2. Materials and methods

*Arabidopsis thaliana* (L. Heynh. ecotype Columbia) was independently transformed using two different constructs containing respectively a constitutive promoter (BIN HYG TX) and an inducible promoter (pER8) both driving the *AtWEE1* gene. The BIN HYG TX vector was already discussed in chapter 2 section 2.2.1. The inducible promoter vector (pER8) allowed the expression of *AtWEE1* only when oestradiol was added to the culture (Fig. 6.1). The *AtWEE1* ORF was amplified using PCR primers incorporating *XhoI* and *SpeI* restriction sites. The PCR product was

cloned into pER8 by cutting and ligating at the *XhoI* and *SpeI* restriction sites in the vector (see chapter 2 section 2.1.1). Both vectors also contained a hygromycin resistance gene, so that plants containing the constructs could be selected.



**Fig. 6.1:** The pER8-*AtWEE1* construct was based on the system developed by Professor N.H. Chua (Zuo *et al.*, 2000), consisting of a synthetic promoter driving the transcription of an estradiol-inducible transcriptional factor, composed by the DNA binding domain of the bacterial repressor LexA, the transcription activation domain of VP16 and the regulatory region of the human estrogen receptor (hER). The terminator sequence E9, encodes a poly-A tail addition site that terminates the production of mRNA by the transcription factor. The *hpt* gene encodes hygromycin phosphotransferase II – required for hygromycin resistance. This gene was under the control of its own promoter and terminator. The *AtWEE1* open reading frame (ORF) was inserted under the control of the minimal 35S promoter fused with 8 copies of the LexA operator sequence. Translation of this sequence was terminated by the 3A terminator.

### 6.2.1. Isolation of transgenic lines

Both BIN HYG TX-*AtWEE1* and pER8-*AtWEE1* constructs were transformed first into *Escherichia coli* (see chapter 2 section 2.2.1) and then into *Agrobacterium tumefaciens* GV1301 (see chapter 2 section 2.2.2). The presence of the plasmids in *Agrobacterium* was checked by PCR before they were used as vectors to transform *Arabidopsis* plants.

To transform *Arabidopsis thaliana* plants, *Agrobacterium* cells were cultured overnight in 500 ml of 2xYT-medium (per litre: 16 g bacto-tryptone, 10 g bacto-yeast extract, 5 g NaCl, pH=7) at 30°C with gentle shaking. The bacterial cells were then centrifuged at 5500 g for 20 min at room temperature (Beckman Coulter J-E centrifuge, rotor JA-14). *Agrobacterium* cells were then resuspended in a solution of 5% sucrose and 0.05% Silwet L-77 (Lehle Seeds) in 500 ml distilled water. The *Agrobacterium* cells were placed in a metal bowl and *Arabidopsis* plants were inverted and dipped into the bacterial suspension for approximately 10 seconds with agitation. The plants were then covered with a plastic bag overnight to retain humidity.

A few weeks following the transformation, seeds were collected from the transformed plants and stored for two weeks at room temperature. Seeds were then sterilized by first immersing them in bleach/water mix (1:10 with water) for 5 minutes, followed by a 5 minute immersion in an ethanol/water/bleach mix (7:2:1 parts respectively). The seeds were then washed three times with sterile distilled water for 5 minutes each time. After the third wash, the seeds were added to a solution of water and agar (0.8%) and sown into Petri dishes containing Murashige and Skoog medium (1962) (M&S, basal salt mixture, Duchefa Biochemie, Netherlands) supplemented with 3% (w/v) sucrose, 1% agar and 30 mg/l hygromycin at a density of approx 10,000 seeds per plate. Transformants were clearly identifiable as they grew on and were green, while untransformed seeds bleached at the cotyledon stage and did not grow any further. At the stage of 3-4 small leaves, the plants were transferred to soil to form the first generation. At this stage, the presence of the construct in the plants was confirmed by PCR. Approximately 200 mg of leaf tissue were placed in a 1.5 ml microfuge and ground using a microfuge grinder; 200 µl of DNA extraction buffer

(0.5% sodium dodecyl sulphate, 250 mM NaCl, 100 mM Tris-HCl (at pH=8) and 25 mM EDTA) were added to the homogenate and the mix was centrifuged for 5 minutes at 12,000 rpm in a microcentrifuge (Eppendorf MiniSpin). The resulting supernatant (150  $\mu$ l) was removed from the microfuge tube and transferred to a fresh one containing an equal volume of isopropanol. This mix was incubated on ice for 5 minutes before another centrifugation at 12000 rpm for 10 minutes at 4°C. After centrifugation, the supernatant was removed and the pellet was air dried at room temperature for 10 minutes and then resuspended in 100  $\mu$ l of Tris-EDTA (10 mM Tris , 1 mM EDTA, pH=8) buffer.

The plant DNA extracts were used in PCR reactions with primers designed to target the BIN HYG TX-*AtWEE1* construct (BHTXF 5'-GTTAACGGTACCCGGGCTCG and P39 5'-CATGGGATGGTCCAGTAATTC) or the pER8-*AtWEE1* construct (primers 35STRS 5'-ACGCTGAAGCTAGTCGACTC and P61 - GTAATGCCTTTGCTATC) respectively. The PCR reactions were as follows: 1  $\mu$ l of DNA extract was added to 24  $\mu$ l of PCR mix, composed of 1  $\mu$ l of the respective primers (10  $\mu$ M of forward, 10  $\mu$ M of reverse primer), and 22  $\mu$ l of Reddy Mix™. Each mix was subjected to 28 cycles in a PTC100 thermocycler consisting of 95°C for 1 minute, then 55°C (for BHTXF and P39 primers) or 60°C (for 35STRS and P61 primers) for 1 minute and finally 72°C for 1 minute. A final incubation of 72°C was carried out for 15 minutes before the PCR mixtures were cooled to 15°C. Five  $\mu$ l of each PCR product were run on a 1% agarose gel containing 1 $\mu$ g/ml ethidium bromide. Positive bands indicated the presence of the constructs in the transgenic plants which were selected for further studies. Subsequently, seeds from the first generation of successfully transformed plants were collected and used to form the second generation, which was used for the

experiments described in this chapter.

### 6.2.2. RNA extraction, cDNA synthesis and RT-PCR

DNA extractions were performed to confirm the successful incorporation of the construct into *Arabidopsis* plants, but they did not prove that the transgenes were being expressed. RT-PCR was used to demonstrate that the transgenic lines were expressing *AtWEE1* at the mRNA level. Total RNA was extracted from two to three leaves of the second plant generation. Leaf material (equivalent to about 200 mg), was collected from plants, wrapped in aluminium foil and immediately frozen in liquid nitrogen. Leaves were then removed from the aluminium foil and ground to a powder in liquid nitrogen using a sterile pestle and mortar (pre-cooled to -20°C). Details of mRNA extractions, cDNA synthesis and cDNA amplification are fully described in chapter 2 sections 2.4.1, 2.4.2 and 2.4.3.

BIN HYG TX-*AtWEE1* and pER8-*AtWEE1* cDNA samples were subjected to PCR using the primers described in section 6.2.1 of this chapter. Since BHTXF and 35STRS were designed to be directed against the vectors BIN HYG TX and pER8 respectively, they can be used to discriminate between the transgene and the endogenous *AtWEE1*. Each of the cDNA samples was used in a RT-PCR reaction mix consisting of 1 µl cDNA, 22 µl of Reddy Mix™, 1 µl of 10 µM forward primer and 1 µl of 10 µM reverse primer. The total number of cycles in a PTC100 thermocycler were 40 consisting of 95°C for 1 minute, 55°C for 1 minute and finally 72°C for 1 minute. Five µl of each of the PCR reaction products were run on a 1% agarose gel containing 1 µg/ml ethidium bromide. Positive bands indicated the presence of *AtWEE1* mRNA in leaf material, where it is not normally expressed at high levels.



### **6.2.3. Seed sterilization, growth conditions and methods of fixation**

Seeds from both BIN HYG TX-*AtWEE1* and pER8-*AtWEE1* transgenic lines were planted alongside the wild type seeds for comparison. All seeds were sterilised (as described in section 6.2.1 of this chapter) and plated individually onto round Petri dishes containing a suitable growth medium. In the case of the wild type, the growth medium consisted of just the basal media: 1% Murashige and Skoog medium at pH 5.7, supplemented with 30 g/l of sucrose and 10 g/l of agar. Transgenic lines were grown on the same basal medium (M&S), which contained 30 mg/l hygromycin used to select for recombinant seedlings. All seeds were arranged in rows, with approximately 20-25 seeds per Petri dish placed more or less 1 cm apart. The plates containing the seedlings were then sealed using porous tape (Micropore™) and incubated at 4°C for 48 hours to coordinate and stimulate germination. All seedlings were then grown in 18 hours light / 6 hours dark per day at 22°C, usually for 2 weeks, but different experiments required different incubation times. The plates were placed at an angle of approximately 45°.

### **6.2.4. *AtWEE1* lines: growth conditions and sampling methods**

Samples, consisting of germinated seedlings, were taken at time points ranging from 15 to 19 days post-incubation, typically every two days (i.e. day 15, 17 and 19). All samples were removed from the agar and fixed in 70 % ethanol:glacial acetic acid (3:1).

Fixed samples were stored at 4°C until removed and stained with Feulgen as follows: each sample was transferred from the ethanol:glacial acetic acid fixative to a

fresh 1.5 ml microfuge tube and rinsed in approximately 1 ml of distilled water for 5 minutes at room temperature. The samples were then added to 1 ml of 5 M HCl for 25 minutes at 25°C and then subjected to three separate 5 minute rinses with distilled water at 4°C. All samples were then immersed in 1 ml of Feulgen stain (Schiff's reagent, Fisher Scientific, U.K.) for 60 minutes.

pER8-*AtWEE1* seedlings were grown for 14 days under the conditions described above on M&S/hygromycin supplemented with 5 µM oestradiol (to induce *AtWEE1* expression) at 22°C. Wild type seedlings were grown on M&S medium for the entire duration of the experiment. Every two days (15, 17 and 19) of growth, the length of the primary root was measured by removing the samples from the medium, fixing and staining them as described above.

#### **6.2.5. Analysis of lateral root development and primordia size**

To analyse root development, a dissecting microscope was used. The primary root was taken as the root to have been the first root to emerge from the seedling. This was also the longest root. Lateral roots were categorized as all of the branch roots not classed as the primary.


In addition to the lateral root development analysis, I estimated average primordium volume. This was achieved by selecting randomly, five primordia from each plant. The primordia were then photographed using a Fujix Digital Camera attached to an Olympus BH-2 microscope for image analysis, using the program SigmaScan Pro (Version 5). The height and width of each primordium was measured and used to estimate primordia volume (assuming that a primordium was a cone) using the equation:

$$V = (\pi r^2 h) / 3$$

Where V = volume, r = radius (half of the width) and h = height (McKibbin *et al.*, 1998). Wild type primordium volume was compared with BIN HYG TX-*AtWEE1* primordium volume after 15, 17 and 19 days of growth, respectively.

### 6.3. Results

DNA extractions and PCR analyses were performed in order to ensure that the BIN HYG TX-*AtWEE1* and pER8-*AtWEE1* constructs were successfully incorporated into *Arabidopsis* plants. Only three plants transformed with BIN HYG TX-*AtWEE1* construct (line: 30, 58 and 61 see table 6.1) were found positive out of 16 analysed (Fig 6.2.a), whereas 4 plants (line: 5, 6, 36 and 40 see table 6.1) containing pER8-*AtWEE1* were positive out of 23 different plants analysed (Fig. 6.2.b). Lines 5, 6 and 40 were chosen for further studies because they showed the strongest product signals.



**Fig. 6.2:** PCR was performed on DNA extracted from *Arabidopsis* transgenic plants. **a)** BIN HYG TX *-AtWEE1*. The primers (BHTXF and P39) used were directed against the BIN HYG TX *-AtWEE1* construct. The numbers above the gel correspond to different transgenic lines; **b)** pER8-*AtWEE1*. The primers (35STRS and P61) used were directed against pER8-*AtWEE1* construct. Each number above the gel corresponds to a different transgenic line. Wild type (WT) cDNA was also used as negative control using **c)** primers BHTXF and P39, **d)** primers 35STRS and P61; **e)** the quality of wild type cDNA was tested using PUV2 and PUV4 primers.

**Table 6.1:** *Arabidopsis thaliana* transgenic plants

| Plasmids:                  | Lines              | Code used in this thesis   |
|----------------------------|--------------------|--|
| BIN HYG TX - <i>AtWEE1</i> | 30<br>58<br>61     | c- <i>AtWEE1</i> referred only to lines 58 and 61, line 30 was not used as explained in the text |
| pER8- <i>AtWEE1</i>        | 5<br>6<br>36<br>40 | i- <i>AtWEE1</i> referred to lines 5, 6 and 40, line 36 was not used as explained in the text.   |


RT-PCR was also performed to ensure *AtWEE1* expression in the transgenic lines. When RT-PCR was performed on BIN HYG TX-*AtWEE1* lines, only lines 58 and 61 out of the three tested exhibited positive results; line 30 was negative (Fig. 6.3 a). However, the cDNA from line 30 produced positive PCR product bands when subjected to RT-PCR using primers targeting *18S rRNA*, confirming that the RNA extraction and cDNA synthesis were successfully (Fig. 6.3 b).

RT-PCR was also performed on induced and non-induced pER8-*AtWEE1* cell lines. In order to induce pER8-*AtWEE1*, seeds from *Arabidopsis* plants lines 5, 6 and 40, were sown on MS medium supplemented with 30 mg/l of hygromycin and 5  $\mu$ M oestradiol (Zuo *et al.*, 2000). Induced seedlings were incubated in a growth room for two weeks as specified above. Seeds from the same *Arabidopsis* inducible lines (5, 6 and 40) were grown on MS medium supplemented with 30 mg/l hygromycin, but to prevent promoter induction oestradiol was not added to the culture medium. Seeds were incubated under the same conditions described above for the induced seeds. When the seedlings reached the stage of 3-4 leaves 1 – 1.5 cm in length, samples were collected for total RNA extraction and cDNA synthesis. Figure 6.3 c shows expression of *AtWEE1* in induced *Arabidopsis* plants compared to the negative mock

reactions, whereas figure 6.3 d shows expression of *AtWEE1* in the same non-induced lines, presumably because of leakiness in the inducible system. This could be because oestradiol is a very common estrogen hormone in nature and present in many different plastic materials including containers for distilled water and Petri dishes (Raloff, 1993, 1999).

### 6.3.1. Constitutive and inducible expression of *AtWEE1* in transgenic *Arabidopsis*

Figure 6.3 shows RT-PCR results for *AtWEE1* expression in transgenic *Arabidopsis* lines.



**Fig. 6.3:** RT-PCR was performed on *Arabidopsis* transgenic plants. The primers used BHTXF and P39 were directed against BIN HYG TX-*AtWEE1* cDNA. **a)** BIN HYG TX-*AtWEE1* RT-PCR. Positive bands were detected only for lines 58 and 61, but not for line 30. M30, M58 and M61 represent the respective mock reactions for each line (mocks contain all cDNA synthesis reagents except the reverse transcriptase). **b)** BIN HYG TX-*AtWEE1* RT-PCR. When the expression of *18S rRNA* was

investigated, positive bands were detected for all three lines, demonstrating that cDNA synthesis was successful. c) pER8-*AtWEE1* RT-PCR (using primers 35STRS and P61), *AtWEE1* expression in the induced lines (5 i, 6 i and 40 i) and corresponding mock reactions (M5, M6 and M40) used as negative controls. d) RT-PCR on non-induced transgenic plants (lines: 5, 6, 40) which showed a leakiness in *AtWEE1* expression, cDNA from line 5 induced was used as positive control. e) RT-PCR was also performed on *Arabidopsis* wild type (WT) cDNA (total RNA was extracted from leaf). The negative control in all cases was distilled water.

### **6.3.1. Constitutive and inducible over-expression of *AtWEE1* in transgenic *Arabidopsis* plants leads to a smaller length of primary root per unit time**

The length of primary root was measured in *Arabidopsis* plants constitutively over-expressing *AtWEE1*. As controls, wild type plants were analysed alongside the transgenic plants.


A difference in primary root length, between transgenic and wild type plants, was already noticeable after 15 days of growth, post-incubation (Table 6.2, Fig. 6.4; see Appendix VI for raw data); at this stage the length of wild type root was 3-fold longer ( $P=0.000$  Table 6.2) than in the transgenic plants and the difference was even greater on days 17 and 19 (in both cases the means were significantly different  $P=0.000$  Table 6.2), demonstrating that the rate of primary root elongation was significantly slower in *AtWEE1* over-expressing plants compared with the wild type control.

*AtWEE1* over expression inhibited growth to such an extent that by day 19 the primary root of the transformed plants rarely exceeded 1.5 cm, whereas by this time, wild type primary roots were 4.5 cm in length on the same day. Hence wild type roots were at least 3-fold longer than *AtWEE1* over-expressing plants.



**Table 6.2:** The mean  $\pm$  standard error (in parenthesis) of primary root length of WT and c-AtWEE1 on days 15, 17 and 19. Levels of significance are indicated by Student's t-test ( $n \geq 15$ ).

| Day | WT                   | c-AtWEE1              | P       |
|-----|----------------------|-----------------------|---------|
| 15  | 2.967 ( $\pm 0.21$ ) | 0.947 ( $\pm 0.15$ )  | = 0.000 |
| 17  | 3.51 ( $\pm 0.33$ )  | 1.080 ( $\pm 0.082$ ) | = 0.000 |
| 19  | 4.45 ( $\pm 0.42$ )  | 1.47 ( $\pm 0.28$ )   | = 0.000 |




**Fig. 6.4:** Comparison of mean primary root length ( $\pm$ SE) between WT and c-AtWEE1 on days 15, 17 and 19. ( $n \geq 15$ ).

*Arabidopsis* plants were also transformed with an inducible promoter to ensure the expression of *AtWEE1* gene in a controllable fashion. The choice of non-induced plants as control was not made because of the leakiness of *AtWEE1* expression in non-induced transgenic *Arabidopsis* plants, noted in the RT-PCR results; instead wild type plants were preferred as controls. *Arabidopsis* plants carrying pER8-*AtWEE1* were induced in order to investigate the development of primary roots compared to wild type on days 15, 17 and 19 post-incubation (Table 6.3, Fig. 6.5; see Appendix VI for raw data). On days 15 and 17 the difference in primary root length between wild type and transgenic plants is  $\sim 1.6$ -fold ( $P=0.000$ , Table 6.3, Fig. 6.5), whereas on day 19

the difference is ~1.4-fold ( $P=0.000$ , Table 6.3, Fig. 6.5).

**Table 6.3:** The mean  $\pm$  standard error (in parenthesis) of primary root length of WT and *i-AtWEE1* on days 15, 17 and 19. Levels of significance are indicated by Student's t-test. ( $n \geq 21$ ).

| Day | WT                   | <i>i-AtWEE1</i>      | P      |
|-----|----------------------|----------------------|--------|
| 15  | 3.04 ( $\pm 0.25$ )  | 1.857 ( $\pm 0.14$ ) | =0.000 |
| 17  | 4.022 ( $\pm 0.21$ ) | 2.343 ( $\pm 0.14$ ) | =0.000 |
| 19  | 4.767 ( $\pm 0.21$ ) | 3.23 ( $\pm 0.12$ )  | =0.000 |



**Fig. 6.5:** Mean primary root length ( $\pm$ SE) of *pER8-AtWEE1* (*i-AtWEE1*) transgenic plants compared to wild type (WT) on days 15, 17 and 19 post-germination. ( $n \geq 21$ ).

### 6.3.2. BIN HYG TX-*AtWEE1* plants initiate primordia at a slower rate per day than wild type

The number of primordia measured on the primary root of transgenic plants was compared to the number obtained from the analysis of wild type plants (Table 6.4). A significant decrease in number of primordia was observed in *AtWEE1* transgenic plants compared with wild type (Table 6.4, Fig. 6.6 a; see Appendix VI for raw data).

The number of primordia initiated in the wild type increases linearly per unit root length as time progresses, but this relationship was not observed in the transgenic plants. Due to a lack of primary root growth in the transgenic *Arabidopsis* plants, primordia appear to be initiated much closer to the root tip (Table 6.5, Fig. 6.6 b; see Appendix VI for raw data).

The youngest initiated primordium is significantly closer ( $P \leq 0.002$ , Table 6.5) to the primary root tip after 17 and 19 days of growth in the *AtWEE1* transformed plants when compared to wild type plants, by 1.5- and 3.5-fold respectively (Table 6.5, Fig 6.6 b). This may be because the transgenic primary roots grow slower than the wild type, leading to primordia formation nearer the root tip.



**Table 6.4:** The mean number ( $\pm$  standard error (in parenthesis)) of root primordia of WT and c-*AtWEE1* on days 15, 17 and 19. Levels of significance are indicated by Student's t-test ( $n \geq 15$ ).

| Day | WT                   | c- <i>AtWEE1</i>     | P       |
|-----|----------------------|----------------------|---------|
| 15  | 2.73 ( $\pm 0.33$ )  | 1.80 ( $\pm 0.20$ )  | = 0.024 |
| 17  | 2.933 ( $\pm 0.21$ ) | 1.867 ( $\pm 0.19$ ) | = 0.001 |
| 19  | 3.07 ( $\pm 0.28$ )  | 1.857 ( $\pm 0.21$ ) | = 0.002 |



**Table 6.5:** The mean distance ( $\pm$  standard error (in parenthesis)) of youngest primordia to root tip in WT and c-AtWEE1 on days 15, 17 and 19. Levels of significance are indicated by Student's t-test ( $n \geq 15$ ).

| Day | WT                    | c-AtWEE1              | P       |
|-----|-----------------------|-----------------------|---------|
| 15  | 0.407 ( $\pm 0.034$ ) | 0.321 ( $\pm 0.058$ ) | = 0.218 |
| 17  | 0.540 ( $\pm 0.038$ ) | 0.232 ( $\pm 0.034$ ) | = 0.000 |
| 19  | 0.907 ( $\pm 0.091$ ) | 0.185 ( $\pm 0.023$ ) | = 0.000 |




**Fig. 6.6:** The effects of constitutive *AtWEE1* (c-*AtWEE1*) expression in *Arabidopsis* plants compared to the wild type (WT) on **a**) the number of primordia initiated from the primary root ( $n \geq 15$ ) and **b**) the distance between the youngest primordia to the tip of primary root ( $n \geq 15$ ) ( $\pm$  SE).

However, the mean volume of the primordium was not affected by the over expression of *AtWEE1* (Table 6.6, Fig. 6.7). The average primordium volume measured on day 15, 17 and 19 was not significantly different ( $P > 0.1$ , Table 6.6) between wild type and transgenic plants (Fig. 6.7; see Appendix VI for raw data).

**Table 6.6:** The mean number ( $\pm$  standard error (in parenthesis)) of primordium volume of WT and c-*AtWEE1* on days 15, 17 and 19. Levels of significance are indicated by Student's t-test ( $n \geq 20$ ).

| Day | WT                    | c- <i>AtWEE1</i>      | P       |
|-----|-----------------------|-----------------------|---------|
| 15  | 88850 ( $\pm$ 9558)   | 81825 ( $\pm$ 7352)   | = 0.564 |
| 17  | 127658 ( $\pm$ 10960) | 120202 ( $\pm$ 18326) | = 0.729 |
| 19  | 191015 ( $\pm$ 19196) | 187342 ( $\pm$ 19042) | = 0.893 |



**Fig. 6.7:** Primordium volume comparison between wild type and transgenic *Arabidopsis* plants ( $\pm$  SE) ( $n \geq 20$ ).


### **6.3.3. *AtWEE1* over-expression results in a reduction in the number of lateral roots**

The number of lateral roots initiated from the primary roots on days 15, 17 and 19 after germination was investigated in both wild type and constitutively expressing *AtWEE1* plants (Table 6.7, Fig. 6.8; see Appendix VI for raw data). The number of lateral roots detected in wild type plants for each day of investigation was 2-fold of that in the transgenic plants (Fig. 6.8). This result was easily predictable because the number of primary root primordia was higher in the wild type than *AtWEE1* over-expressing plants.



**Table 6.7:** The mean number ( $\pm$  standard error (in parenthesis)) of lateral roots of WT and c-AtWEE1 on days 15, 17 and 19. Levels of significance are indicated by Student's t-test ( $n \geq 14$ ).

| Day | WT                   | c-AtWEE1            | P       |
|-----|----------------------|---------------------|---------|
| 15  | 12.07 ( $\pm 0.95$ ) | 5.86 ( $\pm 0.67$ ) | = 0.000 |
| 17  | 12.13 ( $\pm 0.87$ ) | 7.07 ( $\pm 0.74$ ) | = 0.000 |
| 19  | 15.87 ( $\pm 0.74$ ) | 7.71 ( $\pm 0.76$ ) | = 0.000 |



**Fig. 6.8:** The correlation between age in days and the mean ( $\pm$  SE) of lateral roots initiated on both wild type and transgenic plants ( $n \geq 14$ ).

## 6.4. Discussion

The experiments reported in this chapter involved the transformation of *Arabidopsis thaliana* plants with vectors carrying *AtWEE1*. Subsequently, root phenotypes were characterized and compared with wild type *Arabidopsis thaliana* plants. In this study, particular emphasis was placed on the effects of *AtWEE1* over-expression on root growth and development over a five-day period. The investigation



of primary root growth in both constitutive and inducible transgenic lines showed a significantly shorter primary root length than the wild type. The wild type primary root appeared to elongate over the course of five days from 3 cm up to 4.5 cm in length, compared with the primary roots of the transgenic lines that did not grow more than 1 cm in length. In the inducible lines, the primary root length was shorter compared to the wild type on days 15, 17 and 19. However, the difference in primary root length in the inducible plant lines compared with wild type was not as great as in the constitutive plant lines. A hypothesis to explain why the growth of primary root in inducible transgenic plants was longer than the constitutive transgenic plants could be that inducible plants were no longer under a strong influence of the inducible system, perhaps oestradiol was partially degraded so its concentration was not enough to induce high levels of *AtWEE1* expression. Interestingly, experiments on transgenic *Arabidopsis* plants transformed with pER8-*GFP*, carried out by Zuo and colleagues, showed that a prolonged incubation (2-3 weeks) appeared to lead to a reduced transcript level (Zuo *et al.*, 2000). This time-dependent reduction was also observed with all other previously reported inducible systems (Aoyama and Chua, 1997; Martinez *et al.*, 1999). For this reason and since expression of *AtWEE1* was also found in non-induced plants, transgenic *Arabidopsis* plants carrying the inducible system pER8-*AtWEE1* were not used in further experiments.

In the experiments with transformed TBV-2 cells the non-induced cultures provided the ideal controls. Here the use of non-induced plants could not be used due to the leakiness of the inducible system. Unfortunately, empty vector lines were not available, so wild type was used as the control. Further experiments using an empty vector or a neutral transgenic line such as pER8-*GUS* would be important to confirm the results presented here.

The most plausible reason to explain the phenomena of shorter primary root length in the transgenic lines is that the over expression of *AtWEE1* leads the meristematic cells, responsible for root growth, to stop in G2 phase and not engage further in division. The over expression of *AtWEE1* is expected to increase the protein kinase level which subsequently phosphorylates CDKB-types. This would result in an inactivation of the CDK/cyclin complex, which normally leads cells to go into mitosis, causing a delayed primary root growth, since its rate of growth is in part dependent on the rate of meristematic cell production (Beemster and Baskin, 1998; Beemster *et al.*, 2002). This hypothesis is supported by many studies. The maize homologue to *WEE1* (*ZmWEE1*) is capable of phosphorylating and subsequently inactivating plant CDKs leading to an inhibition of mitosis (Sun *et al.*, 1999). Conversely, increased cyclin expression, which is required for CDK activity, increases the rate of mitosis in the primary root resulting in a faster rate of root elongation (Doerner *et al.*, 1996). Also, factors that lengthen the cell cycle reduce elongation. The *SUC1/CSK1 Arabidopsis* homologue *CKS1At*, significantly reduces primary root growth compared to wild type when over expressed in transgenic *Arabidopsis* plants by lengthening the duration of the cell cycle (De Veylder *et al.*, 2001b).

The over expression of *AtWEE1* clearly had significant effects on the rate of lateral root formation from the primary root. In particular, it resulted in a reduced number of lateral roots initiated per unit time in comparison with the wild type. The number of lateral roots and root primordia present on the wild type increased from day 15 to day 19, whereas they did not in the transgenic plants. These results imply that the transgenic plants are initiating primordia but a much reduced frequency that eventually falls to zero. Himanen and colleagues (2002) have demonstrated that

pericycle cell cycle arrest led to a reduction in lateral root primordium initiation. *AtWEE1* over expression in pericycle cells may delay their passage into division leading to a shorter root and a subsequent reduction in the rate of primordium.

*AtWEE1* over-expression did not have an effect on the average primordium volume of transgenic plants, presumably because the smaller number of cells due to a reduced number of divisions is compensated by an increase in volume of cells stopped in G2. This implies that there were fewer large cells in primordia of transgenic plants compared with wild type. This hypothesis is based on the fact that the over-expression of *AtWEE1* in fission yeast cells leads to a long cell phenotype which probably reflects a delay in mitosis (Sorrell *et al.*, 2002). Also, very recently, epidermal cells in the primary root meristem of *Arabidopsis* plants transformed with BIN HYG TX-*AtWEE1* were found to be larger than corresponding cells in the wild type (Cardiff cell cycle laboratory, unpublished data). Those experiments carried out with and without hygromycin excluded any effects of this selective agent (Cardiff cell cycle laboratory, unpublished data).

Transgenic *Arabidopsis* plants over expressing *AtWEE1* initiated approximately half as many lateral root primordia than the wild type. In addition, the c-*AtWEE1* lines formed primordia significantly closer to the root tip than wild type plants. Lateral roots are initiated from pericycle cells thought to have arrested in the G2 phase of the cell cycle after leaving the primary root meristem (Blakel and Evans, 1979). Thus lateral root initiation could be controlled via cells re-entering, and passing through, the G2-mitosis checkpoint (Casimiro *et al.*, 2003). Since *AtWEE1* kinase is associated with blocking the cell in G2 it may play a role in determining which pericycle cells go on to form lateral root primordia and which do not. However, it should also be noted that a recent work on *Arabidopsis* has suggested that many

pericycle cells arrest in G1 phase, whilst only those nearest the developing lateral root are found to be in G2 (Beeckman *et al.*, 2001).

Cyclins like CYCD are well known to stimulate cell cycle progression. CYCD4;1 is generally expressed during lateral root formation, suggesting that the expression of cell cycle genes is a key step in controlling lateral root formation (De Veylder *et al.*, 1999). Cell cycle phosphoregulators have also been shown to have an influence on the rate and patterning of lateral root initiation. *Spcdc25* expression led to an increase in the frequency of lateral roots in tobacco (McKibbin *et al.*, 1998). Since positive cell cycle regulators are capable of influencing where and when root development takes place, a negative cell cycle regulator gene *AtWEE1*, also plays a role in the control of lateral root development.

## 6.5. Summary

The over expression of *AtWEE1* in transgenic *Arabidopsis thaliana* has significant effects on root growth and development, resulting in an inhibition of primary root growth and in a reduction in the number of primordia initiated. Transgenic plants also initiated primordia near the primary root tip possibly as result of a short length of the primary root.

*AtWEE1* might exert its influence on root growth and development by inactivating CDKs which results in an inhibition of the onset of mitosis in dividing cells. This modulates root development by inhibiting mitotic progression and subsequently increasing cell size, confirming that *AtWEE1* is a cell cycle regulator.

## **Chapter 7: General Discussion**

My aim was to gain a better understanding of *WEE1* function as regulator of the plant cell cycle, through the study of *AtWEE1* expression in the tobacco BY-2 cells and in the development of *Arabidopsis thaliana* roots. It became important to identify mechanisms in the higher plant that are either homologous or distinct from other eukaryotic systems such as yeast and animal. Different approaches based on molecular and microscopic techniques were employed to carry out this study.

Based on the experimental evidence that *AtWEE1* expression in *Schizosaccharomyces pombe* results in a long cell phenotype, expression of *AtWEE1* in tobacco cells should result in a large cell size phenotype and in an extended G2 phase. I tested this hypothesis by expressing *AtWEE1* both under constitutive and inducible promoters in tobacco BY-2 cells. Unexpectedly, the experimental evidence negates the hypothesis of increased cell size and extended G2 phase. In fact the exact opposite was found. TBY-2 cell size phenotype was smaller and G2 phase was shortened.

In an attempt to explain why the data found during my experimental work were so different from results previously published (Sorrell *et al.*, 2002), I investigated *NtWEE1* expression in TBY-2 cells transformed with *AtWEE1* under the control of an inducible promoter. Hence the expression of *NtWEE1* could be studied in the presence or absence of *AtWEE1* transcripts. My hypothesis was that expression of a foreign *WEE1* gene in tobacco cells inhibits or affects the normal expression of the endogenous *WEE1*. My experimental data do not support an inhibitory mechanism but instead indicate that induction of *AtWEE1* transcripts alters the timing of *NtWEE1* expression. Note that loss

of *WEE1* function in fission yeast is not lethal but causes cells to enter into mitosis prematurely at reduced cell size (Nurse and Thuriaux, 1980). The same result was found in the budding yeast when deletion of *Swe1*, the homologue of *wee1* in *S. cerevisiae*, led cells to divide at reduced cell size (Harvey and Kellogg, 2003). So too did loss of function of *WEE1* in transgenic *Xenopus* and *Drosophila* (Walter *et al.*, 2000).

Expression of *NtWEE1* peaks in S-phase, this is also confirmed by an independent study in TBY-2 cells (Gonzalez *et al.*, 2004). When *NtWEE1* expression peaked in S-phase the mitotic index peaked at 9 hours, whereas when the timing of *NtWEE1* expression was delayed, the mitotic index peak occurred at 4-5 hours and a smaller mitotic cell size was detected. In effect, these cells suffered loss of function of *WEE1*. In other words, the alteration in the timing of native *WEE1* expression resulted in a reduced cell size at division. From the observation of my data one question was then formulated: why is it that *AtWEE1* does not replace *NtWEE1* kinase activity? If it did, then perturbation of cell size would not be predicted. Two hypotheses were formulated: 1) perhaps *AtWEE1* is transcribed but not translated effectively, so that, the normal inhibition of tobacco CDK activity is removed; 2) alternatively, the *AtWEE1* kinase may have insufficient affinity with tobacco CDKs to regulate them negatively. In support of this hypothesis is the extent of variation between the regulatory (N-terminal) components of *AtWEE1* and *NtWEE1* proteins, which is just 38% compared with 70% homology between catalytic domains (C-terminal) (Cardiff cell cycle laboratory, unpublished data). Resolving this issue will require a highly sensitive *WEE1* kinase assay, able to distinguish *NtWEE1* from *AtWEE1* kinase activity.



In animals, 14-3-3 proteins play an important role in regulating WEE1 kinase activity (Honda *et al.*, 1997; Wang *et al.*, 2000). In *Xenopus* WEE1 (XWEE1), phosphorylation of a serine residue (Ser-549) that resides in a region that contains the RXXS sequence from the consensus motif is required for interaction with 14-3-3 proteins (Lee *et al.*, 2001). In humans, WEE1 binds to 14-3-3 proteins, and phosphorylation of WEE1 on Ser-642, inside the RXXS region is essential for the binding of 14-3-3 proteins to WEE1 (Wang *et al.*, 2000). In AtWEE1, there are two serines (Ser 450 and Ser 485) that reside in regions containing the consensus motif sequences RXXS for binding to 14-3-3 proteins. However, they are not in the same relative positions as those in human and *Xenopus* WEE1, because the predicted plant WEE1 proteins are over 50 amino acids shorter than the animal ones. Interestingly, only one of the potential serine residues (Ser-485) is conserved between *Arabidopsis* and tobacco (Cardiff cell cycle laboratory, unpublished data). Thus, despite a high overall homology between the predicted ORFs in *Arabidopsis* and tobacco, there may be significant divergence in specific regulatory amino acid residues.

In the TBY-2 cell lines, expression of *AtWEE1* induces a small mitotic cell size and in synchronized cells these data are in accord with a shortened G2 phase. The cell size data for weekly sub-cultured wild type TBY-2 cells also show a drop in mitotic cell area during exponential phase but an increase at stationary phase. Interestingly, during normal batch culture of TBY-2 cell lines expressing *AtWEE1*, cell size remained constantly small throughout batch culture. In fact, transformed TBY-2 cells were constantly dividing at smaller cell size. Following synchronization experiments, comparisons between controls and TBY-2 transformed cells revealed differences in

mitotic cell size; the control mitotic cells were always bigger than TBY-2 cells expressing *AtWEE1*. A smaller cell size and a shorter G2 phase were also observed when TBY-2 cells were transformed with a different cell cycle regulator *Cdc25* gene from *Schizosaccharomyces pombe* (*SpCdc25*) (Orchard *et al.*, 2005). Notably, in both *AtWEE1* and *SpCdc25* transformed TBY-2 cells, the shorter G2 phase is compensated by a longer G1 phase. As such, cell cycle length in TBY-2 cells remained remarkably stable regardless of the presence of a foreign cell cycle gene.

As outlined in the Introduction, transition of a plant cell from G2 to M is regulated by two CDKs. Whilst CDKA activity peaks at G1/S and G2/M, *CDKB1* expression is cell cycle regulated and kinase activity is restricted to late G2. In an attempt to explain why transformed TBY-2 cells were undergoing mitosis prematurely, the kinase activity of CDKA and CDKB1 were assayed to examine whether either or both enzymes were precociously activated in cells expressing *AtWEE1*.

In transformed TBY-2 cells, the kinase activity of both CDKA and CDKB1 were investigated in the presence or absence of *AtWEE1* expression. In non-induced cells, CDKA activity was high at both G1/S and G2/M in accordance with published data (Joubes *et al.*, 2000). In non-induced cells, CDKB1 activity peaked only at G2/M in complete agreement with published reports of its activity in wild type cells (Mironov *et al.*, 1999; Sorrell *et al.*, 2001). However, in *AtWEE1* expressing cells, CDKB1 activity was precociously high in early S-phase whereas CDKA activity peaked only at G1/S. Expression of *Spcdc25* in TBY-2 cells also generated a very early and persistent CDKB1 activity which also led to a shortened G2 and a small mitotic cell size (Orchard *et al.*, 2005). Note that in both cases, there was a differential effect on CDKB1 compared with

CDKA activity. From this it can be concluded that CDKB1 kinase activity drives cells into mitosis whereas CDKA may be linked more to the development of mitotic competence. Note that CDKB1 activity, but not CDKA activity, is restricted to proliferative regions of the plant (Magyar *et al.*, 1997; Mironv *et al.*, 1999; Umeda *et al.*, 1999; Joubes *et al.*, 2000; Sorrell *et al.*, 2001).

In fission yeast and animal cells, Wee1/Mik1/Myt1 competes with Cdc25 for the tyrosine 15 residue of Cdc2 (CDK) (Rhind and Russell, 2000). Recently a small CDC25 protein that lacks a regulatory domain has been cloned and is expressed at low levels in all *Arabidopsis* tissues (Sorrell *et al.*, 2005). In higher plants, *AtCdc25* seems to be the obvious counter player to *AtWEE1*. In plants, there is no evidence of a Mik1 or Myt1 so, as far as we know, WEE1 kinase is the only putative negative regulator of B-type CDKs. Normal TBY-2 cell lines exhibit a transcriptional peak of *NtWEE1* at S/G2 that presumably leads to *NtWEE1* kinase activity that represses CDKB1 activity until the cell reaches the optimum conditions for division. However, in TBY-2 cell lines expressing *AtWEE1*, the timing of *NtWEE1* expression is shifted. This may result in the loss of the negative regulator of *NtCDKB1*, which results in a premature cell division. Notably, in TBY-2 cell lines when the timing of *NtWEE1* expression was perturbed, CDKB1 activity was high one hour following the induction of *AtWEE1* transcripts and represents the key difference in CDK activity between treatments. Since *AtWEE1* transcript began to be detected within 15 minutes of supplying DEX to the medium, it must start to perturb native *NtWEE1* regulation of CDKB1 very rapidly. In non-induced *AtWEE1*-transformed TBY-2 cells, *NtCDKB1* activity peaks in mid G2 just as it does in wild type TBY-2 cells (Porceddu *et al.*, 2001; Sorrell *et al.*, 2001). However, *NtCDKA* peaks in early S-phase

regardless of the presence of *AtWEE1* transcript. Indeed, NtCDKA activity is much more constitutive, as found previously for wild type TBY-2 cells (Porceddu *et al.*, 2001; Sorrell *et al.*, 2001), and this could be part of a mechanism that regulates G1/S much like one of its suggested roles in *Arabidopsis* (Joubes *et al.*, 2000). It was previously demonstrated that CDK activity is dependent on native Cdc25 activity (Zhang *et al.*, 2005). Interestingly, in TBY-2 cells expressing *SpCdc25*, CDKB1 kinase activity was persistently high from early S-phase. However, CDKA activity was relatively constant. Collectively these results can show that premature and sustained CDKB1 activity is a key feature of cells that divide prematurely at a small size.

In plants, meristems are responsible for the formation of the entire postembryonic plant body. In *Arabidopsis* a small number of stem cells at the tip of the root generate all the cell types of the root through divisions followed by cell differentiation and regulated cell expansion (Scheres *et al.*, 2002). The primary root meristem of *Arabidopsis* is composed of three layers of initials, L1, L2 and L3 (Steeves and Sussex, 1989; Dolan *et al.*, 1993). L1 (the outermost layer) generates the root cap and epidermis, L2 produces the cortex, and the innermost layer L3 produces the vascular tissue (Scheres *et al.*, 2002). To gain a better understanding of *AtWEE1* function during primary root development, transgenic *Arabidopsis* plants were produced that over-expressed *Arabidopsis WEE1*.

The effect of *AtWEE1* over expression was studied on *Arabidopsis* plant roots. Lateral root primordia arise from the pericycle, the outermost layer of the vascular stele (McCully, 1975). Typically, primordia are initiated in a cluster which is then spatially and temporally separated from the next cluster (Mallory 1970). In the development of a lateral root, two distinct stages were identified: 1) the formation of a primordium, 2) the

subsequent formation of a meristem that is capable of emerging as a lateral root (Laskowski *et al.*, 1995). In *Arabidopsis* plants over-expressing *AtWEE1*, primary root elongation was severely inhibited and there was a lower frequency of lateral root primordia detected. Hence, *AtWEE1* over-expression induces a negative regulation of primary root growth and lateral root formation. It can be hypothesised in its negative setting, *WEE1* is a negative regulator of the G2/M transition in plants and when over-expressed causes yet more negative regulation reflected in slow root growth and fewer lateral root primordia.

The expression of *AtWEE1* did not completely inhibit lateral root development, but was associated with fewer lateral root primordia forming per unit length of primary root tissue. Interestingly, this result is the exact converse in transgenic tobacco plants where induction of *SpCdc25* expression resulted in more lateral root primordia forming per unit length of primary root (McKibbin *et al.*, 1998). The authors suggested that *SpCdc25* expression directly altered the control of cell division in the pericycle giving rise to the effects on cell size and subsequent development of lateral root primordia (McKibbin *et al.*, 1998). In the same way, the effect of *AtWEE1* expression on *Arabidopsis* root development can be explained by the fact that *AtWEE1* altered the control of cell division, delaying cells from entering into mitosis, which resulted in shorter primary roots and in fewer primordia. The molecular model to explain the experimental results is that over expression of *AtWEE1* maintained the native CDK in a phosphorylated inactive state that resulted in a delayed cell division.

The data reported here, show clearly that over-expression of *AtWEE1* in the native genetic background of *Arabidopsis* resulted in a predictable slow-growing phenotype.

Also, very recently, epidermal cells in the primary meristem of *AtWEE1* over-expressed *Arabidopsis* plants were found to be larger than corresponding cells in the wild type (Cardiff cell cycle laboratory, unpublished data). Again, this is a predictable negative effect of *WEE1* expression. Hence, it can be concluded that *AtWEE1* expression in tobacco cells is a transcriptional effect of the transgene on native *NtWEE1* and not, in itself, a typical *AtWEE1* cell cycle effect. Also, to my knowledge, this is the first demonstration of transcriptional regulation of mitotic cell size in higher plants.

The table below summarizes the effect of *AtWEE1* and *Spcdc25* expression on the cell size of fission yeast, *Arabidopsis* roots, tobacco roots and tobacco BY-2 cells:

| Organism                 | <i>AtWEE1</i><br>(cell size) | <i>Spcdc25</i><br>(cell size) |
|--------------------------|------------------------------|-------------------------------|
| Fission yeast            | large                        | small                         |
| <i>Arabidopsis</i> roots | large                        | small                         |
| Tobacco roots            |                              | small                         |
| TBY-2 cells              | small                        | small                         |

### Further Work

*Nicotiana tabacum WEE1 (NtWEE1)* was recently sequenced in our laboratory and a pTA7002 inducible vector was used to make a construct carrying *NtWEE1* gene.

pTA7002-*NtWEE1* plasmid was employed to transform tobacco BY-2 cells. Microscopic analysis on the newly established TBV-2 cell line over-expressing *NtWEE1* should be carried out to investigate the cell phenotype. Synchronization experiments can also give new information about the effect of *NtWEE1* on the tobacco cell cycle. It would be of particular interest to transform *Arabidopsis thaliana* cells with *NtWEE1* to investigate cell phenotype and cell cycle compared with the tobacco cell line over-expressing *NtWEE1*. *Arabidopsis thaliana* plants have also recently been transformed with *NtWEE1* to investigate its action on primary root development and primordium formation. These results will provide an important comparison to the effects of *AtWEE1* over expression in *Arabidopsis* plants and *AtWEE1* expression in TBV-2 cells.

Antibodies able to discriminate between *AtWEE1* and *NtWEE1* kinases, should be made to prove whether *AtWEE1* is translated in transgenic TBV-2 cells. Immunoprecipitation could also be employed to study *AtWEE1* and *NtWEE1* protein abundance during the cell cycle of transgenic tobacco cells, and this could be compared with *NtWEE1* protein abundance of wild type TBV-2 cells.

A WEE1 kinase assay that uses as substrate CDK proteins, which contain tyrosine 15 residues that can be phosphorylated by WEE1, should be developed. The antibody should specifically detect only the phosphorylated form of the tyrosine 15 residue on the CDK. To investigate whether *AtWEE1* is active in transgenic tobacco cells, an *AtWEE1* kinase assay could be performed. An *NtWEE1* kinase assay could also be performed on wild type and transgenic TBV-2 cells.

Gene silencing is, without doubt, one of the best approaches to study the function of a target gene in a living organism. To gain a better understanding of *WEE1* activity



during the plant cell cycle, the recently developed RNA interference (RNAi) technique is currently being employed in the Cardiff cell cycle laboratory to silence the expression of *NtWEE1* in wild type tobacco BY-2 cells. This system was successfully used in a variety of organisms and cell types (e.g., worms, fruit flies and plants). It is based on the utilization of long double-stranded RNAs (dsRNAs) to silence the expression of a target gene (as described in the mechanism of RNA interference (RNAi), Ambion's online appendix) the principal steps are: 1) the long dsRNAs enter a cellular pathway that is commonly referred to as the RNA interference (RNAi) pathway; 2) the dsRNAs are processed into 20-25 nucleotide (nt) small interfering RNAs (siRNAs) by an RNase III-like enzyme called Dicer (initiation step); 3) the siRNAs are then assembled into endoribonuclease-containing complexes known as RNA-induced silencing complexes (RISCs), unwinding in the process. The siRNA strands subsequently guide the RISCs to complementary RNA molecules, where they cleave and destroy the cognate RNA (effector step). Cleavage of cognate RNA takes place near the middle of the region bound by the siRNA strand ([www.ambion.com/techlib/append/RNAi\\_mechanism.html](http://www.ambion.com/techlib/append/RNAi_mechanism.html)) (Tang *et al.*, 2003).

Cell phenotype and the cell cycle can be studied using wild type tobacco BY-2 cells that have been subjected to RNAi treatment and are lacking *NtWEE1* kinase. CDKA and CDKB1 kinase activity should be also investigated in order to determine the effect of silencing *NtWEE1* on these two very important regulators of entry into mitosis.

The yeast two-hybrid assay, employed to find interactions between *AtWEE1* and other proteins, gave putative positive interactions (see Appendix VII for more details). However, these results must be further tested to check that they are not falsely positive.

This can be done by transforming *Arabidopsis thaliana* plants with plasmids carrying the gene encoding for the putative positive proteins fused to non-fluorescent domains of the yellow fluorescent protein. The two non-fluorescent domains become fluorescent when brought together by association of AtWEE1 with the interacting protein (Hu *et al.*, 2000, Walter *et al.*, 2004).

The experiments in which *Arabidopsis thaliana* plants are transformed with *AtWEE1* using an inducible promoter (induced by oestradiol) and a constitutive promoter are described in chapter 6. RT-PCR analysis relative to non-induced transgenic *Arabidopsis* plants showed leakiness of *AtWEE1* expression. This could be due to the fact that oestradiol is a very common estrogen hormone in nature and present in many different plastic materials including containers for distilled water and Petri dishes. This inconvenience can be avoided by using glass, not plastic, Petri dishes for seed germination and seedling growth, and storing the distilled water required for all aspects of the technique in a glass container. RT-PCR could be then repeated to check if under these conditions *AtWEE1* is still expressed in non-induced transgenic *Arabidopsis* plants. Eliminating *AtWEE1* expression in non-induced plants presents two advantages: 1) the effect of *AtWEE1* expression can be efficiently studied only when it is required (induced); 2) non-induced transgenic plants can be used as the control.

It would also be instructive to measure the duration of the cell cycle in transgenic *Arabidopsis* root meristem cells using flow cytometric analysis to investigate at which stage of the cell cycle the majority of root meristem cells are delayed. These results can be then compared with wild type.

## References

### A

Ach R.A., Durfee T., Miller A.B., Zambryski P.C., Hanley-Bowdoin L., Gruisem W. (1997). *RRB1* and *RRB2* encode maize retinoblastoma related proteins that interact with a plant D-type cyclin and geminivirus replication protein. *Mol. Cell. Biol.* **17**, 5077-86.

Aligue, R., Wu, L., Russell, P. (1997). Regulation of *Schizosaccharomyces pombe* Wee1 Tyrosine Kinase. *J. Biol. Chem.* **272**, 13320-13325.

Altman, R. and Kellogg, D. R. (1997). Control of mitotic events by Nap1 and the Gin4 kinase. *J. Cell Biol.* **138**, 119-130.

Amino S., Fujimura T. and Komamine A. (1983). Synchrony induced by double phosphate starvation in a suspension culture of *Catharanthus roseus*. *Plant Physiol.* **59**, 393-396.

An G.H. (1985). High-efficiency transformation of cultured tobacco cells. *Plant Physiol.* **79**, 568-570.

Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. (2000), *Nature* **408**, 796-815.

Aoyama T. and Chua N.H. (1997). A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J.* **11**, 605-612.

Atherton-Fessler S., Hannig G. and Piwnica-Worms H. (1993). Reversible tyrosine phosphorylation and cell cycle control. *Semin. Cell Biol.* **4**, 433-442.

### B

Baldin V. and Ducommun B. (1995). Subcellular localization of human wee1 is regulated during the cell cycle. *J. Cell Sci.* **108**, 2425-2432.

Barral, Y., Parra, M., Bidlingmaier, S. and Snyder, M. (1999). Nim1-related kinases coordinate cell cycle progression with the organization of the peripheral cytoskeleton in yeast. *Genes Dev.* **13**, 176-187.

Beato M. (1989). Gene regulation by steroid hormones. *Cell* **56**, 335-344.

Beeckman T., Burssens S. and Inze D. (2001). The peri-cell-cycle in *Arabidopsis*. *J. Exp. Bot.* **52**, 403-411.

Beemster G.T.S. and Baskin T.I. (1998). Analysis of Cell Division and Elongation Underlying the Developmental Acceleration of Root Growth in *Arabidopsis thaliana*. *Plant Physiol.* **116**, 1515-1526.

Beemster G.T.S., De Vusser K., De Tavernier E., De Bock K. and Inze D. (2002). Variation in Growth Rate between *Arabidopsis* Ecotypes Is Correlated with Cell Division and A-Type Cyclin-Dependent Kinase Activity. *Plant Physiol.* **129**, 854-864.

Beemster G.T.S., Fiorani F. and Inze D. (2003). Cell cycle: the key to plant growth control? *Trends Plant Sci.* **8**, 154-158.

Beemster G.T.S., De Veylder L, Vercruyssen S, West G, Rombaut D, Van Hummelen P, Galichet A, Gruijsem W, Inze D. Vuylsteke M. (2005). Genome-wide analysis of gene expression profiles associated with cell cycle transitions in growing organs of *Arabidopsis*. *Plant Physiol.* **138**, 734-743.

Berridge M.J. (1995). Capacitative calcium entry. *Biochem. J.* **312**, 1-11.

Binarova P., Dolezel J., Draber P., Herberle-Bors E., Strnad M. and Bogre L. (1998). Treatment of *Vicia faba* root tip cells with specific inhibitors to cyclin dependent kinases leads to abnormal spindle formation. *Plant J.* **16**, 697-707.

Blakel, L.M. and Evans T.A. (1979). Cell dynamics studies on the pericycle of radish seedling roots. *Plant Sci. Lett.* **14**, 79-83.

Boddy M. N., Furnari B., Mondesert O. and Russell P. (1998 ). Replication checkpoint enforced by kinases Cds1 and Chk1. *Science (Wash. DC)*, **280**, 909-912.

Böhner S., Lenk I., Rieping M., Herold M. and Gatz C. (1999). Transcriptional activator TGV mediates dexamethasone inducible and tetracycline-inactivatable gene expression, *Plant J.* **19**, 87-95.

Bolle C. (2004). The role of GRAS proteins in plant signal transduction and development. *Planta* **218**, 683-692.

Boucheron E, Healy JH, Bajon C, Sauvanet A, Rembur J, Noin M, Sekine M, Riou Khamlichi C, Murray JA, Van Onckelen H. (2005). Ectopic expression of *Arabidopsis* CYCD2 and CYCD3 in tobacco has distinct effects on the structural organization of the shoot apical meristem. *J. Exp. Bot.* **56**, 123-134.

Broek D., Barlett R. et al. (1991). Involvement of p34<sup>cdc2</sup> in establishing the dependency of S phase on mitosis. *Nature* **349**, 388-393.

Burssens S., Van Montagu M. and Inzé D. (1998). The cell cycle in *Arabidopsis*, *Plant Physiol. Biochem.* **36**, 9-12.

Burssens, de Almeida Engler J., Beeckman T., Richard C., Shaul O., Ferreira P., Van Montagu M. and Inze D. (2000). Developmental expression of the *Arabidopsis thaliana* *CycA2;1* gene. *Planta* **211**, 623-631.

## C

Carroll C., Altman R., Schieltz D., Yates J. and Kellogg D. R. (1998). The septins are required for the mitosis-specific activation of the Gin4 kinase. *J. Cell Biol.* **143**, 709-717.

Casimiro I., Beeckman T., Graham N., Bhalerao R., Zhang H., Caser, P., Sandberg G. and Bennet, M.J. (2003). Dissecting *Arabidopsis* lateral root development. *Trends Plant Sci.* **8**, 165-171.

Chaubet-Gigot N. (2000). Plant A-type cyclins. *Plant Mol. Biol.* **43**, 659-675.

Coleman T. R., Tang Z. and Dunphy W. G (1993). Negative regulation of the wee1 protein kinase by direct action of the nim1/cdr1 mitotic inducer. *Cell* **72**, 919-929.

Combettes B., Reichheld J.P., Chauboute M.E., Philipps G., Shen W.H. and Chaubet-Gigot N. (1999). Study of phase-specific gene expression in synchronized tobacco cells. *Methods Cell Sci.* **21**, 109-121.

Cockcroft C.E., den Boer B.G., Healy J.M. and Murray J.A. (2000). Cyclin D control of growth rate in plants. *Nature* **405**, 575-9.

Cullen, P. J. and Sprague, G. F., Jr (2000). Glucose depletion causes haploid invasive growth in yeast. *Proc. Natl. Acad. Sci. USA* **97**, 13619-13624.

## D

Day I.S. and Reddy A.S.N. (1998). Isolation and characterization of two cyclin-like cDNAs from *Arabidopsis*. *Plant Mol. Biol.* **36**, 451-461.

Davies B., Egea-Cortines M., de Andrade Silva E., Saedler H. and Sammer H. (1996). Multiple interactions amongst floral homeotic proteins. *EMBO J.* **15**, 4330-4343.

De Bondt H.L., Rosenblatt J., Jancarik J., Jones H.D., Morgan D.O. and Kim S.H. (1993). Crystal structure of cyclin-dependent kinase 2. *Nature* **363**, 595-602.

De Veylder L., De Almeida J., Burssens E.S., Manevski A., Lescure B., Van Montagu M., Engler G. and Inze D. (1999). A new D-type cyclin of *Arabidopsis thaliana* expressed during lateral root primordia formation. *Planta* **208**, 453-462.

De Veylder L., Beeckman T., Beemster G.T.S., Krols L., Terras F., Landrieu I., Van Der Schueren E., Maes S., Naudts M. and Inze D. (2001a). Functional Analysis of Cyclin-

Dependent Kinase Inhibitors of Arabidopsis. *Plant Cell* **13**, 1653-1668.

De Veylder L., Beemster G.T., Beeckman T. and Inze D. (2001b). CKS1At overexpression in *Arabidopsis thaliana* inhibits growth by reducing meristem size and inhibiting cell-cycle progression. *Plant J.* **25**, 617-26.

Deshaies R.J. (1995). The self-destructive personality of a cell cycle in transition, *Curr. Opin. Cell Biol.* **7**, 781-789.

Dewitte W. and Murray J.A.H. (2003). The Plant Cell Cycle. *Annu. Rev. Plant Biol.* **54**, 235-264.

Doerner P., Jorgensen J.E., You R., Steppuhn J. and Lamb C. (1996). Control of root growth and development by cyclin expression. *Nature* **380**, 520-3.

Doonan J. (1996). Plant growth: roots in the cell cycle. *Curr. Biol.* **6**, 788-9.

Dubrovsky J.G., Rost T.L., Colon-Carmona A. and Doerner P. (2001). Early primordium morphogenesis during lateral root initiation in *Arabidopsis thaliana*. *Planta* **214**, 30-36.

Dunphy W.G. (1994). The decision to enter mitosis. *Trends Cell Biol.* **4**, 202-207.

## E

Eckardt N.A. (2001). A brief tour of the cell cycle. *Plant Cell* **13**, 449-451.

Edgington, N. P., Blacketer, M. J., Bierwagen, T. A. and Myers, A. M. (1999). Control of *Saccharomyces cerevisiae* filamentous growth by cyclin-dependent kinase Cdc28. *Mol. Cell. Biol.* **19**, 1369-1380.

Eisenhaber F., Wechselberger C. and Kreil G. (2001). The Brix domain protein family – a key to the ribosomal biogenesis pathway? *Trends Biochem. Sci.* **26**, 345-347.

Enoch T. and Nurse P. (1990). Mutation of fission yeast cell cycle control genes abolishes dependence of mitosis on DNA replication. *Cell* **60**, 665-673.

Evans T., Rosenthal E., Youngbloom J., Distel D. and Hunt T. (1983). Cyclin: a protein specified by maternal mRNA in sea-urchin eggs that is destroyed at each cleavage division. *Cell* **33**, 389-396.

## F

Fantes P. and Nurse P. (1977). Control of cell size in fission yeast by a growth modulated size control over nuclear division. *Exp. Cell Res.* **107**, 377-386.

Fantes P. and Nurse P. (1978). Control of the timing of cell division in fission yeast. Cell size mutants reveal a second control pathway. *Exp. Cell Res.* **115**, 317-329.

Feiler H. S., Desprez T., Santoni V., Kronenberger J., Caboche M. and Traas, J. (1995). The higher plant *Arabidopsis thaliana* encodes a functional CDC48 homologue which is highly expressed in dividing and expanding cells. *EMBO J.* **14**, 5626-5637.

Fields S. and Song O. K. (1989). A novel genetic system to detect protein-protein interactions. *Nature* **340**, 245-246.

Fobert P.R., Gaudin V., Lunness P., Coen E.S. and Doonan J.H. (1996). Distinct classes of cdc2-related genes are differentially expressed during the cell division cycle in plants. *Plant Cell* **8**, 1465-1476.

Fuerst R.A.U.A., Soni R., Murray J.A.H. and Lindsey K. (1996). Modulation of cyclin transcript levels in cultured cells of *Arabidopsis thaliana*. *Plant Physiol.* **112**, 1023-1033.

Fulop, K., Pettko-Szandtner, A., Magyar, Z., Miskolczi, P., Kondorosi, E., Dudits, D. and Bako, L. (2005). The Medicago CDKC;1-CYCLINT;1 kinase complex phosphorylates the carboxy-terminal domain of RNA polymerase II and promotes transcription. *Plant J.* **42**, 810–820.

Fukuda H., Ito M., Sugiyama M. and Komamine A. (1994). Mechanisms of the proliferation and differentiation of plant cells in cell culture systems. *Int. J. Dev. Biol.* **38**, 287-299.

## G

Garrett, J. M. (1997). The control of morphogenesis in *Saccharomyces cerevisiae* by Elm1 kinase is responsive to RAS/cAMP pathway activity and tryptophan availability. *Mol. Microbiol.* **26**, 809-820.

Gatz C, Kaiser A. and Wenderburg R. (1991). Regulation of a modified CaMV 35s promoter by the Tn10-encoded Tet repressor in transgenic tobacco. *Mol. Gen. Genet* **227**, 229-237.

Gatz C., Frohberg C. and Wenderburg R. (1992). Stringent repression and homogenous derepression by tetracycline of a modified CaMV 35S promoter in intact transgenic tobacco plants. *Plant J.* **2**, 397-404.

Glab N., Labidi B., Qin L-X., Trehin C., Bergonioux C. and Meijer L. (1994). Olomoucine, an inhibitor of the cdc2/cdk2 kinases activity, blocks plant cells at the G1 to S and G2 to M cell cycle transitions. *FEBS Lett.* **353**, 207-211.

Glotzer M., Murray A.W. and Kirschner M.W. (1991). Cyclin is degraded by the ubiquitin pathway. *Nature* **349**, 132-138.



Gonzalez N., Hernould M., Delmas F., Gevaudant F., Duffe P., Causse M., Mouras A., and Chevalier C. (2004). Molecular characterization of a *WEE1* gene homologue in tomato (*Lycopersicon esculentum* Mill.) *Plant Mol. Biol.* **56**, 849-861.

Gould A.R. (1984). Control of the cell cycle in cultured plant cells. *CRC Crit. Rev. Plant Sci.* **1**, 315-344.

## H

Harper J.W. and Elledge S.J. (1998). The role of Cdk7 in CAK function, a retrospective. *Genes Dev.* **12**, 285-289.

Hartwell L. H., Culotti J., Pringle J. R., and Reid B. J. (1974). Genetic control of the cell division cycle in yeast. *Science* **183**, 46-51.

Hartwell L. H. and Unger, M. W. (1977). Unequal division in *Saccharomyces cerevisiae* and its implications for the control of cell division. *J. Cell Biol.* **75**, 422-435.

Hartwell L. H and Weinter T., (1989). Checkpoints: controls that ensure the order of cell cycle events. *Science* **246**, 629-634.

Harvey, S. L. and Kellogg, D. R. (2003). Conservation of mechanisms controlling entry into mitosis: budding yeast *wee1* delays entry into mitosis and is required for cell size control. *Curr. Biol.* **13**, 264-275.

Hasezawa S. and Nagata T. (1991). Dynamic organization of plant microtubules at the 3 distinct transition points during the cell-cycle progression of synchronized tobacco BY-2 cells. *Bot. Acta.* **104**, 206-211.

Hata S., Kouchi H., Suzuka I. and Ishii T. (1991). Isolation and characterization of cDNA clones for plant cyclins. *EMBO J.* **10**, 2681-2688.

He SS, Liu J, Xie Z, O'Neill D, Dotson S. (2004). *Arabidopsis* E2Fa plays a bimodal role in regulating cell division and cell growth. *Plant Mol. Bio.* **56**, 171-184.

Herbert R.J., Vilhar B., Evett C., Orchard C.B., Rogers H.J., Davies M.S. and Francis D. (2001). Ethylene induces cell death at particular phases of the cell cycle in the tobacco TBV-2 cell line. *J. Exp. Bot.* **52**, 1615-1623.

Himanen K., Boucheron E., Vanneste S., de Almeida Engler J., Inze D. and Beeckman T. (2002). Auxin-Mediated Cell Cycle Activation during Early Lateral Root Initiation. *Plant Cell* **14**, 2339-2351.

Hirayama T., Imajuku Y., Anai T., Matsui M. and Oka A. (1991). Identification of two cell-cycle-controlling *cdc2* gene homologs in *Arabidopsis thaliana*. *Gene* **105**, 159-165.

Hirt H., Páy A., Bögre L., Meskiene I. and Herberle-Bors E. (1993). *cdc2MsB*, a cognate *cdc2* gene from alfalfa, complements the G1/S but not the G2/M transition of budding yeast *cdc28* mutants. *Plant J.* **4**, 61-69.

Honda R., Ohba Y. and Yasuda H. (1997). 14-3-3  $\zeta$  protein binds to the carboxyl half of mouse wee1 kinase. *Biochem. Biophys. Res. Commun.* **230**, 262-265.

Howard A. and Pelc S.R. (1953). Synthesis of deoxyribonucleic acid in normal and irradiated cells and its relation to chromosome breakage. *Hevedily (suppl.)* **6**, 261-273.

Hsieh W.L. and Wolniak S.M. (1998). Isolation and characterization of a functional A-type cyclin from maize. *Plant Mol. Biol.* **37**, 121-129.

Hu C.D., Chinenov Y. and Kerpola T.K. (2000). Visualization of interactions among bZIP and Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol. Cell* **9**, 789-798.

Hu Y., Bao F. and Li J. (2000). Promotive effect of brassinosteroids on cell division involves a distinct CycD3-induction pathway in *Arabidopsis*. *Plant J.* **24**, 693-701.

Huntley R.P., Healy S., Freeman D., Lavender P. and de Jager S. (1998). The maize retinoblastoma protein homologue ZmRb-1 is regulated during leaf development and displays conserved interactions with G1/S regulators and plant cyclin D (Cyc D) proteins. *Plant Mol. Biol.* **37**, 155-169.

Huntley R.P. and Murray J.A.H.M. (1999). The plant cell cycle. *Curr. Opin. Plant Biol.* **2**, 440-446.

## I

Imajuku Y., Hirayama T., Endoh H. and Oka A. (1992). Exon-intron organization of the *Arabidopsis thaliana* protein kinase genes *CDC2a* and *CDC2b*. *FEBS Lett* **304**, 73-77.

Ito M., Marieclaire C., Sakabe M., Ohno T., Hata S., Kouchi H., Hashimoto J., Fukuda H., Komamine A. and Watanabe A. (1997). Cell cycle regulated transcription of A- and B- type plant cyclin genes in synchronous cultures. *Plant J.* **11**, 983-992.

Ito M., Iwase M., Kodama H., Lavis P., Komamine A., Nishihama R., Machida Y. and Watanabe A. (1998). A novel *cis*-acting element in promoters of plant B-type cyclin genes activates M phase-specific transcription. *Plant Cell* **10**, 331-341.

Ito M. (2000). Factors controlling cyclin B expression. *Plant Mol. Biol.* **43**, 677-690.

## J

Jakoby M., Weisshaar B., Droge-Laser W., Vicente-Carbajosa J., Tiedemann J., Kroj T. and Parcy, F. (2002). bZIP Transcription Factors in Arabidopsis. *Trends Plant Sci.* **7**, 106-111.

Jarillo J.A., Capel J., Tang R.H., Yang H.Q., Alonso J.M., Ecker J.R. and Cashmore A. R. (2001). An *Arabidopsis* circadian clock component interacts with both CRY1 and phyB. *Nature* **410**, 487-490.

Jeffrey P.D., Russo A.A., Polyak K., Gibbs E., Hurwitz J., Massague J. and Pavletich N.P. (1995). Mechanism of CDK activation revealed by the structure of a cyclinA-CDK2 complex. *Nature* **376**, 313-320.

Johnston, G. C., Pringle, J. R. and Hartwell, L. H. (1977). Coordination of growth with cell division in the yeast *Saccharomyces cerevisiae*. *Exp. Cell. Res.* **105**, 79-98.

Jorgensen, P., Nishikawa, J. L., Breikreutz, B. J. and Tyers, M. (2002). Systematic identification of pathways that couple cell growth and cell division in yeast. *Science* **297**, 395-400.

Joubès J., Chevalier C., Dudits D., Herberle-Bors E., Inze D., Umeda M. and Renaudin J.P. (2000). CDK-related protein kinases in plants. *Plant Mol. Biol.* **43**, 607-620.

## K

Kato K., Matusomoto T., Koiwai A., Mizusaki S., Nishida K., Noguchi M. and Tamaki E. (1972). In "Fermentation Technology Today" (G. Terui, ed.), 689-685. Soc. Ferment. Technol., Osaka, Japan.

Kellogg, D. R. (2003). Wee1-dependent mechanisms required for coordination of cell growth and cell division. *J. Cell Sci.* **116**, 4883-4890.

Kidou S., Umeda M. and Uchimiya H. (1994). Nucleotide sequence of rice (*Oryza sativa* L.) cDNA homologous to *cdc2* gene. *DNA Seq* **5**, 125-129.

King P.J., Mansfield K.J. and Street H.E. (1973). Control of growth and cell division in plant cell suspension cultures. *Can. J. Bot.* **51**, 1807-1823.

Kirshi T. and Yamao F. (1998). An essential function of Grr1 for the degradation of Cln2 is to act as binding core that links Cln2 to Skp1. *J. Cell Sci.* **111**, 3655-3661.

Kodama H., Ito M., Hattori T., Nakamura K. and Komamine A. (1991). Isolation of genes that are preferentially expressed at the G1/S boundary during the cell cycle in synchronized cultures of *Catharanthus roseus* cells. *Plant Physiol.* **95**, 406-411.

Kohn K.W. (1999). Molecular interaction map of the mammalian cell cycle control and DNA repair systems. *Mol. Biol. Cell.* **10**, 2703-2734.

Kumagai A. and Dunphy W.G. (1991). The cdc25 protein controls tyrosine dephosphorylation of the cdc2 protein in a cell-free system. *Cell* **64**, 903-914.

## L

Landrieu I, da Costa M, De Veylder L, Dewitte, F, Vandepoele K, Hassan S, Wieruszski J-M, Faure J-D, Van Montague M, Inze D, Lippens G. (2004). A small CDC25 dual-specificity tyrosine-phosphatase isoform in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the USA* **101**, 13380–13385.

Laskowski M.J., Williams M.E., Nusbaum H.C. and Sussex I.M. (1995). Formation of lateral root meristems is a two stage process. *Development* **121**, 3303-3310.

La Valle, R. and Wittenberg, C. (2001). A role for the Swe1 checkpoint kinase during filamentous growth of *Saccharomyces cerevisiae*. *Genetics* **158**, 549-562.

Lee, J., Kumagai, A. and Dunphy, W. G. (2001). Positive regulation of Wee1 by Chk1 and 14-3-3 proteins. *Mol. Biol. Cell* **12**, 551-563.

Lew D.J. and Kornbluth S. (1996). Regulatory roles of cyclin dependent kinase phosphorylation in cell cycle control. *Curr. Opin. Cell Biol.* **8**, 795-804.

Linsmaire E.M. and Skoog F. (1964). Organic growth factor requirement of tobacco tissue, *Physiol. Plant* **18**, 100-127.

Liu F., Stanton J.J., Wu Z. and Piwinica-Worms H. (1997). *Mol. Cell. Biol.* **17**, 571-583.

Liu P., Koeller D.M. and Kaler S.G. (2003). Genomic organization of ATOX1, a human copper chaperone. *BMC Genetics* **4**:4 doi:10.1186/1471-2156-4-4.

Longtine, M. S., Theesfeld, C. L., McMillan, J. N., Weaver, E., Pringle, J. R. and Lew, D. J. (2000). Septin-dependent assembly of a cell cycle-regulatory module in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **20**, 4049-4061.

Lorbiecke R. and Sauter M. (1999). Adventitious root growth and cell-cycle induction in deepwater rice. *Plant Physiol.* **119**, 21-29.

Lucretti S. and Dolezel J. (1995). Cell cycle synchronization, Chromosome isolation, and flow sorting in plants. In: *Methodos in Plant Cell Biology (PartB)*. (Galbraith D.W., Borque D.P. and Bohnert H.J., eds). San Diego: Academic Press, pp. 61-83.

## M

- Ma, J. X., Lu, Q. and Grunstein, M. (1996). A search for proteins that interact genetically with histone H3 and H4 amino termini uncovers novel regulators of the Swe1 kinase in *Saccharomyces cerevisiae*. *Genes Dev.* **10**, 1327-1340.
- Magyar Z., Bako' L., Bogre L., Dedeoglu D., Kapros T. and Dudits D. (1993). Active *cdc2* genes and cell cycle phase-specific *cdc2*-related kinase complexes in hormone stimulated alfalfa cells. *Plant J.* **4**, 151-161.
- Magyar Z., Meszaros T., Miskolczi P., Deak M., Feher A., Brown S., Kondorosi E., Athanasiadis A., Pongor S. and Bilgin M. (1997). Cell cycle phase specificity of putative cyclin-dependent kinase variants in synchronized alfalfa cells. *Plant Cell* **9**, 223-235.
- Mallory T.E., Chingan S., Cutter E.G. and Gifford E.M. (1970). Sequence and pattern of lateral root formation in five selected species. *Am. J. Bot.* **57**, 800-809.
- Martin-Castellanos C., Labib K. and Moreno S. (1996). B-type cyclins regulate G1 progression in fission yeast in opposition to the p25<sup>rum1</sup> cdk inhibitor. *EMBO J.* **15**(4), 839-49.
- Martin-Castellanos C. and Moreno S. (1997). Recent advances on cyclins, CDKs and CDK inhibitors. *Trends Cell Biol.* **7**.
- Martinez A., Sparks C., Hart C.A., Thompson J. and Jepson I. (1999). Ecdysone agonist inducible transcription in transgenic tobacco plants. *Plant J.* **19**, 97-106.
- Maundrell K. (1993). Thiamine-repressible expression vectors pREP and pRIP for fission yeast. *Gene* **123**, 127-130.
- Mazzurco M., Sulaman W., Elina H., Cock J.M. and Goring D.R. (2001). Further analysis of the interactions between the Brassica S receptor kinase and three interacting proteins (ARC1, THL1 and THL2) in the yeast two-hybrid system. *Plant Mol. Biol.* **45**, 365-376.
- McCully M.E. (1975). The developmental of lateral roots. In: Torrey J.G. and Clarkson D.T. (eds). *The Development and Function of Roots*, pp. 105-124. Academic Press, London.
- McGowan C. H. and Russell P (1995). Cell cycle regulation of human WEE1. *EMBO J.* **14**, 2166-2175.
- McKibbin R.S., Halford N.G. and Francis D. (1998). Expression of fission yeast *cdc25* alters the frequency of lateral root formation in transgenic tobacco, *Plant Mol. Biol.* **36**, 601-612.

- McMillan, J. N., Longtine, M. S., Sia, R. A. L., Theesfeld, C. L., Bardes, E. S. G., Pringle, J. R. and Lew, D. J. (1999). The morphogenesis checkpoint in *Saccharomyces cerevisiae*: cell cycle control of Swe1p degradation by Hsl1p and Hsl7p. *Mol. Cell. Biol.* **19**, 6929-6939.
- McMillan, J. N., Theesfeld, C. L., Harrison, J. C., Bardes, E. S. and Lew, D. J. (2002). Determinants of Swe1p degradation in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **13**, 3560-3575.
- Meijer M. and Murray J.A.H. (2000). The role and regulation of D-type cyclins in the plant cell cycle. *Plant Mol. Biol.* **43**, 621-633.
- Menges M, de Jager SM, Gruissem W, Murray JA. (2005). Global analysis of the core cell cycle regulators of *Arabidopsis* identifies novel genes, reveals multiple and highly specific profiles of expression and provides a coherent model for plant cell cycle control. *Plant J.* **41**, 546–566.
- Menges M, Samland AK, Planchais S, Murray JA. (2006). The D-type cyclin CYCD3;1 is limiting for the G1-to-S-phase transition in *Arabidopsis*. *Plant Cell* **18**, 893–906.
- Meyerowitz E.M. (1997). Genetic control of cell division patterns in developing plants. *Cell* **88**, 299-308.
- Miyazawa Y., Nakajima N., Abe T., Sakai A., Fujoka S., Kawano S., Kuroiwa T. and Yoshida S. (2003). Activation of cell proliferation by brassinolide application in tobacco BY-2 cells: effects of brassinolide on cell multiplication, cell cycle related gene expression and organellar DNA contents. *J. Exp. Bot.* **393**, 2669-2678.
- Millar J.B., Lenaers G. and Russell P. (1992). Pyp3 PTPase acts as a mitotic inducer in fission yeast. *EMBO J.* **11**, 4933-4941.
- Mironov V., De Veylder L., Van Montagu M. and Inzé D. (1999). Cyclin-dependent kinases and cell division in plants -- the Nexus. *Plant Cell* **11**, 509-521.
- Moreno S., Nurse P. and Russell P. (1990). Regulation of mitosis by cyclic accumulation of p80<sup>cdc25</sup> mitotic inducer in fission yeast. *Nature* **344**, 549-552.
- Morgan D.O. (1997). Cyclin-dependent kinases: engines, clocks and microprocessors. *Annu. Rev. Cell Dev. Biol.* **13**, 261-291.
- Mueller P.R., Coleman T.R., Kumagai A. and Dunphy W.G. (1995). Myt1: a membrane-associated inhibitory kinase that phosphorylates Cdc2 on both threonine-14 and tyrosine-15 of special interest. *Science* **270**, 86-90.
- Murashige T. and skoog F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant* **15**, 473-497.

## N

- Nachtwey D.S. and Cameron J.L. (1968). Cell cycle analysis. In: *Methods in Cell Physiology*, vol. 3 (Prescott D.M. ed.). New York-London: Academic Press.
- Nagata T., Nemoto Y. and Hasezawa S. (1992). Tobacco BY-2 cell line as the "HeLa" cell line in the cell biology of higher plants. *Int. Rev. Cytol.* **132**, 1-30.
- Nagata T. and Kumagai F. (1999). Plant cell biology through the window of the highly synchronized tobacco BY-2 cell line. *Methods Cell Sci.* **21**, 123-127.
- Nakagami H., Sekine M., Murakami H. and Shinmyo A. (1999). Tobacco retinoblastoma-related protein phosphorylated by a distinct cyclin-dependent kinase complex with Cdc2/cyclin D *in vitro*. *Plant J.* **18**, 243-252.
- Nakai T, Kato K, Shinmyo A, Sekine M. (2006). *Arabidopsis* KRPs have distinct inhibitory activity toward cyclin D2-associated kinases, including plant-specific B type cyclin-dependent kinase. *FEBS Lett* **580**, 336-340.
- Nasmyth K. (1993). Control of the yeast cell cycle by the Cdc28 protein kinase. *Curr. Opin. Cell Biol.* **5**, 166-179.
- Nasmyth K. (1996). At the heart of the budding yeast cell cycle. *Trends Genet.* **12**, 405-412.
- Nigg E.A. (1995). Cyclin-dependent protein kinases: Key regulators of the eukaryotic cell cycle. *Bioessays* **17**, 471-480.
- Nigg E.A. (1996). Cyclin-dependent kinase7: at the cross road of transcription , DNA repair and cell cycle control? *Curr. Opin. Cell Biol.* **8**, 312-317.
- Nishida T., Ohnishi N., Kodama H. and Komamine A. (1992). Establishment of synchrony by starvation and readition of auxin in suspension cultures of *Catharanthus roseus* cells. *Plant Cell Tissue Organ Cult.* **28**, 37-43.
- Noble M.E., Endicott J.A., Brown N.R. and Johnson L.N. (1997). The cyclin box fold: protein recognition in cell cycle and transcription control. *Trends Biochem. Sci.* **22**, 482-487.
- Novak B., Csikasz -Nagy A., Gyorffy B., Nasmyth K. and Tyson J. (1998). Model scenarios for evolution of the eukaryotic cell cycle. *Philos. Trans. R Soc London B Biol. Sci.* **353**, 2063-2076.
- Nurse, P. (1975). Genetic control of cell size at cell division in yeast. *Nature* **256**, 547-551.



Nurse, P. and Thuriaux P. (1980). Regulatory genes controlling mitosis in the fission yeast *Schizosaccharomyces pombe*. *Genetics* **96**, 627-637.

## O

O'Connell M. J., Raleigh J. M., Verkade H. M. and Nurse P. (1997). Chk1 is a wee1 kinase in the G2 DNA damage checkpoint inhibiting cdc2 by Y15 phosphorylation. *EMBO J.* **16**, 545-554.

Orchard C., Siciliano I., Sorrel D., Marchbank A., Rogers H., Francis D., Herbert R., Suchomelova P., Lipavska H., Azmi A. and van Onckelen H. (2005). Tobacco BY-2 cells expressing fission yeast *cdc25* bypass G2/M block on the cell cycle. *Plant J.* **44**, 290-299.

Oullet F., Overvoorde P.J. and Theologis A. (2001). IAA17/AXR3: Biochemical insight into an auxin mutant phenotype. *Plant Cell* **13**, 829-841.

## P

Park JA, Ahn JW, Kim YK, Kim SJ, Kim JK, Kim WT, Pai, HS. (2005). Retinoblastoma protein regulates cell proliferation, differentiation, and endoreduplication in plants. *Plant J.* **42**, 153-163.

Parker L.L. and Piwnica-Worms H. (1992). Inactivation of the p34cdc2--cyclin B complex by the human WEE1 tyrosine kinase. *Science* **257**, 1955-1957.

Parker L.L., Walter S. A. and Piwnica-Worms H. (1993). Phosphorylation and inactivation of the mitotic inhibitor Wee1 by the nim1/cdr1 kinase. *Nature* **363**, 736-738.

Parker L. L., Sylvestre P. J., Byrnes M. J., Liu F. and Piwnica-Worms H. (1995) Identification of a 95-kDa WEE1-like tyrosine kinase in HeLa cells. *Proc. Natl. Acad. Sci. USA.* **92**, 9638-9642.

Parfit D., Herbert R.J., Rogers H., Francis D. (2004). Differential expression of putative floral genes in *Pharbitis nil* shoot apices cultured on glucose compared with sucrose. *J. Exp. Bot.* **406**, 2169-2177.

Perennes C., Qin L.X., Glab N. and Bergonioux C. (1993). *Pertunia* p34 (Cdc2) protein kinase activity in G2-M cells obtained with a reversible cell cycle inhibitor, mimosine. *FEBS Lett.* **333**, 141-145.

Picard D. (1993). Steroid-binding domains for regulating the functions of heterologous proteins in *cis*. *Trends Cell Biol.* **3**, 278-280.

Pines J. (1995). Cyclins and cyclin-dependent kinases: a biochemical view. *Biochem J.* **308**, 697-711.

Planchais S., Glab N., Trehin C., Perennes C., Bureau J.M., Meijer L. and Bergounioux C. (1997). Roscovitine, a novel cyclin dependent kinase inhibitor, characterizes restriction point and G2/M transition in tobacco BY-2 cell suspension. *Plant J.* **12**, 191-202.

Planchais S., Glab N., Inze D. and Bergounioux C. (2000). Chemical inhibitors: a tool for plant cell cycle studies. *FEBS Lett.* **476**, 78-83.

Porceddu A., Stals H., Reichheld J.P., Serges G., De Veylder L., de Pinho Barroco R., Casteels P., Van Montagu M., Inze D. and Mironov V. (2001). A plant specific cyclin dependent kinase is involved in the control of G2/M progression in plants. *J. Biol. Chem.* **276**, 36354-36360.

## Q

Quastler H. and Sherman F.G. (1959). Cell population kinetics in the intestinal epithelium of the mouse. *Exp. Cell Res.* **17**, 749-758.

## R

Raloff, J. (1993). Plastics may shed chemical estrogens. *Science News* **144** (July 3):12.

Raloff, J. (1999). What's coming out of baby's bottle. *Science News Online* (Sept. 4).

Rechsteiner M.P.E.S.T. (1990). Sequences are signals for rapid intracellular proteolysis. *Semin. Cell. Biol.* **1**, 433-440.

Reichheld J.P., Sonobe S., Clement B., Chaubet N. and Gigot C. (1995). Cell cycle regulated histone gene expression in synchronized plant cells. *Plant J.* **7**, 245-252.

Reichheld J.P., Chaubet N., Shen W.H., Renaudin J.P. and Gigot C. (1996). Multiple A-type cyclins express sequentially during the cell cycle in *Nicotiana tabacum* BY-2 cells. *Proc. Natl. Acad. Sci. USA* **93**, 13819-13824.

Renaudin J.P., Doonan J.H., Freeman D., Hashimoto J., Hirt H., et al. (1996). Plant cyclins: a unified nomenclature for plant A-, B- and D-type cyclins based on sequence organization. *Plant Mol. Biol.* **32**, 10031018.

Renaudin J.P., Savouré A., Philippe H., Van Montagu M., Inzé D. and Rouzé P. (1998). Characterization and classification of plant cyclin sequences related to A- and B-type cyclins. In: D. Francis, D. Dudits and D. Inzé Editors, *Plant Cell Division* Portland Press, London, 67-98.

Rhind N. and Russell P. (2000). Chk1 and cds1: linchpins of the DNA damage and replication checkpoint pathways. *J. Cell Sci.* **113**, 3896-3899.

Riou-Khamlichi C., Huntley R., Jacquard A. and Murray J. (1999). Cytokinin activation of *Arabidopsis* cell division through a D-type cyclin. *Science* **283**, 1541-1544.

Riou-Khamlichi C., Menges M., Healy J. and Murray J. (2000). Sugar control of the plant cell cycle: differential regulation of *Arabidopsis* D-type cyclin gene expression. *Mol. Cell Biol.* **20**, 4513-4521.

Rosenquist M., Sehnke P.C., Ferl R.J., Sommarin M. and Larsson C. (2000). Evolution of the 14-3-3 protein family: does the large number of isoforms in multicellular organisms reflect functional specificity? *J. Mol. Evol.* **51**, 446-458.

Rosenquist M., Asterfjord M., Larsson C. and Sommarin M. (2001). Data mining the *Arabidopsis* genome reveals fifteen 14-3-3 genes. Expression is demonstrated for two out of fifteen genes. *Plant Physiol.* **127**, 142-149.

Rothblum-Oviatt, C. J., Ryan, C. E. and Piwnica-Worms, H. (2001). 14-3-3 binding regulates catalytic activity of human Wee1 kinase. *Cell Growth Differ.* **12**, 581-589.

Roudier F., Fedorova E., Gyorgyey J., Feher A., Brown S., Kondorosi A. and Kondorosi E. (2000). Cell cycle function of a *Medicago sativa* A2-type cyclin interacting with a PSTAIRE-type cyclin-dependent kinase and a retinoblastoma protein. *Plant J.* **23**, 73-83.

Rupes, I. (2002). Checking cell size in yeast. *Trends Genet.* **18**, 479-485.

Russell P. and Nurse P. (1986). *cdc25*<sup>+</sup> functions as an inducer in the mitotic control of fission yeast. *Cell* **45**, 145-153.

Russell P. and Nurse P. (1987). Negative regulation of mitosis by *wee1*<sup>+</sup>, a gene encoding a protein kinase homolog. *Cell* **45**, 145-153.

## S

Sala F., Parisi B., Burrioni D., Amileni A.R., Pedrall-Noy G. and Spadari S. (1980). Specific and reversible inhibition by aphidicolin in the alpha like DNA polymerase of plant cells. *FEBS Lett.* **177**, 93-98.

Samuels A.L., Meehl J., Lipe M. and Stahelin L.A. (1998). Optimizing condition for tobacco BY-2 cell cycle Synchronization rapid communication. *Protoplasma* **202**, 232-236.

Sanchez-Diaz A., Gonzalez I., Arellano M. and Moreno S. (1998). The Cdk inhibitors p25<sup>rum1</sup> and p40<sup>sic1</sup> are functional homologues that play similar roles in the regulation of the cell cycle in fission yeast and budding yeast. *J. Cell Sci.* **111**, 843-851.

Sauter M. (1997). Differential expression of a CAK (*cdc2* activating kinase) like protein kinase, cyclins and *cdc2* genes from rice during the cell cycle and in response to gibberellin. *Plant J.* **11**, 181-190.

- Schell J. (1987). Transgenic plants as tools to study the molecular organization of plant genes. *Science*, **237**, 1176-1183.
- Schena M., Lloyd A.M. and Davis R.W. (1991). A steroid inducible gene expression system for plant cells. *Proc. Natl. Acad. Sci. USA* **88**, 10421-10425.
- Scheres B., Benfey P. and Dolan L. (2002). Root development. The *Arabidopsis* Book. American Society of Plant Biologists.
- Schwob E., Böhm T., Mendenhall M.D. and Nasmyth K. (1994). The B-type cyclin kinase inhibitor p40SICI controls the G1 to S transition in *S. cerevisiae*, *Cell* **79**, 233-234.
- Segers G., Gadisseur I., Bergounioux C., de Almeida Engler J., Jacquard A., van Montagu M. and Inzé D. (1996). The *Arabidopsis* cyclin-dependent kinase gene *cdc2bAt* is preferentially expressed during S and G2 phases of the cell cycle. *Plant J.* **10**, 601-612.
- Segers G., Rouzé P., Van Montagu M., and Inzé D. (1997). Cyclin-dependent kinases in plants. In *Plant Cell Proliferation and Its Regulation in Growth and Development*, J.A. Bryant and D. Chiatante, eds (Chichester, UK: John Wiley and Sons), 1-19.
- Segers G., Rouzé P., Van Montagu M. and Inzé D. (1998). Cyclin dependent kinases in plants. In: J.A. Bryant and D. Chiatante Editors, *Plant Cell Proliferation and its Regulation in Growth and Development* Wiley, Chichester, 1-19.
- Setiady Y.Y., Sekine M., Hariguchi N., Kouchi H. and Shinmyo A. (1996). Molecular cloning of a cDNA clone that encodes a *cdc2* homolog from *Nicotiana tabacum*. *Plant Cell Physiol.* **37**, 369-376.
- Sherr C.J. and Roberts J.M. (1995). Inhibitors of mammalian G1 cyclin-dependent kinase. *Genes Dev.* **9**, 1149-1163.
- Shimotohno A, Umeda-Hara C, Bisova K, Uchimiya H, Umeda M. (2004). The plant-specific kinase CDKF;1 is involved in activating phosphorylation of cyclin dependent kinase-activating kinases in *Arabidopsis*. *Plant Cell* **16**, 2954-2966.
- Shulewitz, M. J., Inouye, C. J. and Thorner, J. (1999). Hsl7 localizes to a septin ring and serves as an adapter in regulatory pathway that relieves tyrosine phosphorylation of Cdc28 protein kinase in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **19**, 7123-7137.
- Sia, R.A.L., Bardes, E.S.G. and Lew, D.J. (1998). Control of Swe1p degradation by the morphogenesis checkpoint. *EMBO J.* **17**, 6678-6688.
- Solomon M.J. (1993). Activation of the various cyclin/cdc2 protein kinases. *Curr Opin Cell Biol.* **5**, 180-186.

Soni R., Carmichael J.P., Shah Z.H. and Murray J.A.H. (1995). A family of cyclin D homologs from plants differentially controlled by growth regulators and containing the conserved retinoblastoma protein interaction motif. *Plant Cell* **7**, 85-103.

Sonnewald U., brauer M., von Schaeven A., Stiff M. and Willmitzer L. (1991). Transgenic tobacco plants expressing yeast-derived invertase in either the cytosol, vacuole or apoplast: a powerful tool for studying sucrose metabolism and sink/source interactions. *Plant J.* **1**, 95-106.

Sorrell D.A., Combettes B., Chaubet-Gigot N., Gigot C. and Murray J.A.H (1999). Distinct cyclin D genes show mitotic accumulation or constant levels of transcripts in tobacco Bright Yellow-2 cells. *Plant Physiol.* **119**, 343-35.

Sorrell D.A., Menges M., Healy J.M.S., Deveaux Y., Amano C., Su Y., Nakagami H., Shinmyo A., Doonan J.H., Sekine M. and Murray J.A.H. (2001). Cell cycle regulation of cyclin dependent kinase in tobacco Bright Yellow-2 (BY-2) cells. *Plant Physiol.* **126**, 1214-1223.

Sorrell D.A., Marchbank A., McMahon K., Dickinson J.R., Rogers H.J. and Francis D. (2002). A WEE1 homologue from *Arabidopsis thaliana*. *Planta* **215**, 518-522.

Sorrell D.A., Marchbank A., Chrimes D., Dickinson J.R., Rogers H.J., Francis D., Grierson C.S. and Halford N.G. (2003). The *Arabidopsis* 14-3-3 protein, GF14 $\omega$ , binds to the *Schizosaccharomyces pombe* Cdc25 phosphatase and rescues checkpoint defects in the rad24<sup>-</sup> mutant. *Planta* **218**, 50-57.

Sorrell D.A., Chrimes D., Dickinson J.R., Rogers H.J. and Francis D. (2005). The *Arabidopsis* CDC25 induces a short cell length when overexpressed in fission yeast: evidence for cell cycle function. *New Phytologist* **165**, 425-428.

Sreenivasan, A. and Kellogg, D. (1999). The Elm1 kinase functions in a mitotic signaling network in budding yeast. *Mol. Cell. Biol.* **19**, 7983-7994.

Stals H., Casteels P., Van Montagu M. and Inze D. (2000). Regulation of cyclin-dependent kinases in *Arabidopsis thaliana*. *Plant Mol. Biol.* **43**, 583-593.

Steeves T.A. and Sussex I.M. (1989). Patterns in Plant development, 2<sup>nd</sup> Edition. Cambridge. New York: Cambridge University Press.

Stuart D. and Wittenberg C. (1995). CLN3, not positive feedback, determines the timing of CLN2 transcription in cycling cells. *Genes and Development* **9**, 2780-2794.

Suchomelova P., Velgova D., Masek T., Francis D., Rogers H.J., Marchbank, A.M. and Lipavska H. (2004). Expression of the fission yeast cell cycle regulator cdc25 induces de novo shoot formation in tobacco: evidence of a cytokinin-like effect by this mitotic activator. *Plant Physiol. Biochem.* **42**, 49-55.

Sun Y., Dilkes B.P., Zhang C., Dante R.A., Carneiro N.P., Lowe K.S., Jung R., Gordon-Kamm W.J. and Larkins B.A. (1999). Characterisation of maize (*Zea mays* L.) Wee1 and its activity in developing endosperm. *Proc. Natl. Acad. Sci. USA* **96**, 4180-4185.

Swiatek A., Lenjour M., Van Bockstaele D., Inze D. and Van Onckelen H. (2002). Differential effect of jasmaine acid and abscisic acid on cell cycle progression in tobacco BY-2 cells. *Plant Phy.* **128**, 201-211.

## T

Tang Z., Coleman T. R. and Dunphy W. G. (1993). Two distinct mechanisms for negative regulation of the Wee1 protein kinase. *EMBO J.* **12**, 3427-3436.

Tang G., Reinhardt B.J., Bartel D.P. and Zamore, P.D. (2003). A Biochemical Framework for RNA Silencing in Plants. *Genes Dev.* **17**, 49-63.

Taylor J.H. (1958). The mode of chromosome duplication in *Crepis capillaries*. *Exp. Cell Res.* **15**, 350-357.

Thuriaux, P., Nurse, P. and Carter, B. (1978). Mutants altered in the control co-ordinating cell division with cell growth in the fission yeast *Schizosaccharomyces pombe*. *Mol. Gen. Genet.* **161**, 215-220.

Tjandra, H., Compton, J. and Kellogg, D. R. (1998). Control of mitotic events by the Cdc42 GTPase, the Clb2 cyclin and a member of the PAK kinase family. *Curr. Biol.* **8**, 991-1000.

Treimer J.F. and Zenk M.H. (1979). Purification and properties of strictosidine synthase, the key enzyme in indole alkaloid formation. *European Journal of Biochemistry.* **101**, 225-233.

Trimarchi M. and Lees J.A. (2002). Sibling rivalry in the E2F family. *Nat. Rev. Mol. Cell Biol.* **3**, 11-20.

## U

Uemukai K, Iwakawa H, Kosugi S, de Uemukai S, Kato K, Kondorosi E, Murray JA, Ito M, Shinmyo A, Sekine M. (2005). Transcriptional activation of tobacco E2F is repressed by co-transfection with the retinoblastoma-related protein: cyclin D expression overcomes this repressor activity. *Plant Mol. Bio.* **57**, 83-100.

Umeda M., Bhalerao R.P., Schell J., Uchimiya H. and Konez C. (1998). A distinct cyclin-dependent kinase-activating kinase of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **95**, 5021-5026.

Umeda M., Umeda-Hara C., Yamaguchi M., Hashimoto J. and Uchimiya H. (1999). Differential expression of genes for cyclin-dependent protein kinases in rice plants. *Plant Physiol.* **119**, 31-40.

## V

Vandepoele K., Raes J., De Veylder L., Rouze P., Rombauts S., and Inze D. (2002). Genome-wide analysis of core cell cycle genes in *Arabidopsis*. *Plant Cell* **14**, 903-916.

Van't Hof, J. (1966). Experimental control of cell division in higher plants. *Am. J. Bot.* **53**, 970-976.

Van't Hof J. (1974). Control of the cell cycle in higher plants. Cell cycle controls (ed. Padilla G.M., Cameron I.H. and Zimmerman A.M.) Academic Press, New York.

Verkest A, Weint C, Inze D, De Veylder L, Schnittger A. 2005. Switching the cell cycle. Kip-related proteins in plant cell cycle control. *Plant Phy.* **139**, 1099–1106.

## W

Wagstaff C., Leverentz M.K., Griffiths G., Thomas ., Chanasut U., Stead A.D. and Rogers H.J.(2002). Cysteine protease gene expression and proteolytic activity during senescence of *Alstroemeria* petals. *J.Exp. Bot.* **53**, 233-240.

Walter M., Chaban C., Schutze K., Batistic O., Weckerman K., Nake C, Blazevic D., Grefen C., Schumacher K., Oecking C., Harter K. and Kudla J. (2004). Visualization of protein interactions in living plant cells using biomolecular fluorescence complementation. *Plant J.* **40**, 428-438.

Walter, S. A., Guadagno, S. N. and Ferrell, J. E. J. (2000). Activation of Wee1 by p42 MAPK in vitro and in cycling *Xenopus* egg extracts. *Mol. Biol. Cell* **11**, 887-896.

Wang H., Fowke L.C. and Crosby W.L. (1997). A plant cyclin-dependent kinase inhibitor gene. *Nature* **386**, 451-452.

Wang H., Qi Q., Schrr P., Culter A., Crosby W.L. and Fowke L.C. (1998). ICK1, a cyclin dependent protein kinase inhibitor from *Arabidopsis thaliana* interacts with both Cdc2a and CycD3, and its expression is induced by abscisic acid. *Plant J.* **24**, 613-623.

Wang, Y., Jacobs, C., Hook, K. E., Duan, H., Booher, R. N. and Sun, Y. (2000). Binding of 14-3-3beta to the carboxyl terminus of Wee1 increases Wee1 stability, kinase activity, and G2-M cell population. *Cell Growth Differ.* **11**, 211-219.

Wang, H., Zhou, Y., Gilmer, S., Cleary, A., John, P., Whitwill, S. and Fowke, L. (2003). Modifying plant growth and development using the CDK inhibitor ICK1. *Cell Biol. Int.*



27, 297-299.

Wang W. and Chen X. (2004). HUA ENHANCER3 reveals a role for a cyclin-dependent protein kinase in the specification of floral organ identity in *Arabidopsis*. *Development* **131**, 3147-3156.

Wagner U., Edwards R., Dixon D. P. and Mauch F. (2002). Probing the Diversity of the *Arabidopsis* glutathione *S*-Transferase Gene Family. *Plant Mol. Biol.* **49**, 515–532.

Watanabe N., Broome M. and Hunter T. (1995). Regulation of the human WEE1Hu CDK tyrosine 15-kinase during the cell cycle. *EMBO J.* **14**. 1878-1891.

Weinl C, Marquardt S, Kuijt SJ, Nowack MK, Jakoby MJ, Hulskamp M, Schnittger A. (2005). Novel functions of plant cyclin-dependent kinase inhibitors, ICK1/KRP1, can act non-cell-autonomously and inhibit entry into mitosis. *Plant Cell* **17**, 1704–1722.

## Y

Yamaguchi M., Umeda M. and Uchimya H. (1998). A rice homolog of CDK7/Mo15 phosphorylates both cyclin-dependent protein kinases and the carboxy terminal domain of RNA polymerase II. *Plant J.* **16**, 613-619.

Yamaguchi M., Fabian T., Sauter M., Bhalerao R.P., Schrader J. et al. (2000). Activation of CDK-activating kinase is dependent on interaction with H-type cyclins in plants. *Plant* **24**, 11-20.

## Z

Zhang K., Diderich L. and John P.C.L. (2005). The cytokinin requirement for cell division in cultured *Nicotiana glauca* cells can be satisfied by yeast Cdc25 tyrosine phosphatase. Implications for mechanisms of cytokinin response and plant development. *Plant Physiol.* **137**, 308-316.

Zhou Y., Brandizzi F., Fowke L.C. and Wang H. (2003). The plant cyclin-dependent kinase inhibitor ICK1 has distinct functional domains for in vivo kinase inhibition, protein instability and nuclear localization. *Plant J.* **35**, 476-489.

Zuo J., Niu Q. and Chua N. (2000). An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. *Plant J.* **24** (2), 265-273.

## Appendix I: Sequences of *AtWEE1* ORF

### BIN HYG TX-*AtWEE1* clone 1S

1s contig 1 ATGTTTCGAGAAGAACGGAAGAACACTGTTGGCGAAGAGGAAAACCCAAGG  
athweel ORF 1 ATGTTTCGAGAAGAACGGAAGAACACTGTTGGCGAAGAGGAAAACCCAAGG

1s contig 51 GACAATCAAACCAGGGCATCGAAGAAGATTTCGGAAGATGGAAGGGACAT  
athweel ORF 51 GACAATCAAACCAGGGCATCGAAGAAGATTTCGGAAGATGGAAGGGACAT

1s contig 101 TGGAGCGTCACTCTCTGCTTCAATTCCGGTCAATTGTCGAAGATTTCTTTC  
athweel ORF 101 TGGAGCGTCACTCTCTGCTTCAATTCCGGTCAATTGTCGAAGATTTCTTTC

1s contig 151 GAAAATCGTCCGTCGTCGAATGTTGCTTCATCGGCGTTTCAGGGTCTCCT  
athweel ORF 151 GAAAATCGTCCGTCGTCGAATGTTGCTTCATCGGCGTTTCAGGGTCTCCT

1s contig 201 GGATTCCGATTCTTCGGAATCCGAAATCAGTTGGGTCCGCTGATTCAG  
athweel ORF 201 GGATTCCGATTCTTCGGAATCCGAAATCAGTTGGGTCCGCTGATTCAG

1s contig 251 ATGCCAATTCGGGAGAGAAGGACTTTATTCTTAGCCAAGACTTCTTCTGC  
athweel ORF 251 ATGCCAATTCGGGAGAGAAGGACTTTATTCTTAGCCAAGACTTCTTCTGC

1s contig 301 ACACCTGACTATATAACCCCGGACAATCAGAACTTGATGAGCGGGCTAGA  
athweel ORF 301 ACACCTGACTATATAACCCCGGACAATCAGAACTTGATGAGCGGGCTAGA

1s contig 351 TATCAGCAAGGATCATTCTCCATGTCCTAGGTCTCCTGTTAAACTAAATA  
athweel ORF 351 TATCAGCAAGGATCATTCTCCATGTCCTAGGTCTCCTGTTAAACTAAATA

1s contig 401 CAGTTAAAAGCAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATTCA  
athweel ORF 401 CAGTTAAAAGCAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATTCA

1s contig 451 AATTCTACCTGGTCTTCAAACATAGAGTAGATGAACAAGAGAATGATGA  
athweel ORF 451 AATTCTACCTGGTCTTCAAACATAGAGTAGATGAACAAGAGAATGATGA

1s contig 501 TATTGACACAGATGAGGTGATGGGGATAAACTCCAAGCTAATCAAACAG  
athweel ORF 501 TATTGACACAGATGAGGTGATGGGGATAAACTCCAAGCTAATCAAACAG

1s contig 551 AGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTCGGGCT  
athweel ORF 551 AGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTCGGGCT

1s contig 601 ATGCCACCCCTTGCCTCAAGAAATCCGTATGTGTTGAATCAGTCTGAGAC  
athweel ORF 601 ATGCCACCCCTTGCCTCAAGAAATCCGTATGTGTTGAATCAGTCTGAGAC

1s contig 651 TGCTACTGACCCTTTTGGACATCAGAGATCAAATGTGCAAGTTTTCTCC  
athweel ORF 651 TGCTACTGACCCTTTTGGACATCAGAGATCAAATGTGCAAGTTTTCTCC

1s contig 701 CTGTAAGTACAAGTGGGGATGGCTTGTCAGATATCTCACGGACTTTCAT  
athweel ORF 701 CTGTAAGTACAAGTGGGGATGGCTTGTCAGATATCTCACGGACTTTCAT

1s contig 751 GAAATCCGGCAAATGGTGTGACATTTTCAGTCCGGTATTTAAGGTGTT  
athweel ORF 751 GAAATCCGGCAAATGGTGTGACATTTTCAGTCCGGTATTTAAGGTGTT

1s contig 801 GAAGAGAATGGACGGTTGCCATATATGCTGTGAAACACAGCACAAAGAAAGT  
athweel ORF 801 GAAGAGAATGGACGGTTGCCATATATGCTGTGAAACACAGCACAAAGAAAGT

1s contig 851 TGTATCTAGATTTCAGAGAGACGTAAGCTATGATGGAAGTGCAGCTCTT  
athweel ORF 851 TGTATCTAGATTTCAGAGAGACGTAAGCTATGATGGAAGTGCAGCTCTT

1s contig 901 GCTGCTCTTGGGTTTCATGAAAATATAGTAGGATATTACTCCTCGTGTT  
athweel ORF 901 GCTGCTCTTGGGTTTCATGAAAATATAGTAGGATATTACTCCTCGTGTT

1s contig 951 TGAAAATGAGCAATTATACATTCAACTGGAATCTGCGATCACAGCTTGT  
athweel ORF 951 TGAAAATGAGCAATTATACATTCAACTGGAATCTGCGATCACAGCTTGT

1s contig 1001 CAGCTTTGCCTAAGAAATCTTCTCTTAAAGTCTCAGAAAGAGAGATCTTG  
athweel ORF 1001 CAGCTTTGCCTAAGAAATCTTCTCTTAAAGTCTCAGAAAGAGAGATCTTG

1s contig 1051 GTGATTATGCATCAGATAGCAAAGGCATTACATTTTGTGCATGAGAAAGG  
athweel ORF 1051 GTGATTATGCATCAGATAGCAAAGGCATTACATTTTGTGCATGAGAAAGG

1s contig 1101 AATAGCTCATTAGATGTAACCTGACAATATTTACATTAAGAACGGTG

*athweel ORF* 1101 AATAGCTCATTAGATGTAACCTGACAATATTTACATTAAGAACGGTG  
*1s contig* 1151 TTTGCAAGCTTGGTGACTTTGGTTGTGCCACACGATTGGACAAAAGCTTA  
*athweel ORF* 1151 TTTGCAAGCTTGGTGACTTTGGTTGTGCCACACGATTGGACAAAAGCTTA  
*1s contig* 1201 CCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAAATTTCTAAATGA  
*athweel ORF* 1201 CCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAAATTTCTAAATGA  
*1s contig* 1251 AGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTTAGGTGTGACGG  
*athweel ORF* 1251 AGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTTAGGTGTGACGG  
*1s contig* 1301 TTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAAGAAACCAGTCC  
*athweel ORF* 1301 TTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAAGAAACCAGTCC  
*1s contig* 1351 CTCAATATCAAAGAAGGAAAACCTCCTCTCCTTCTGGCCATTCGTTGCA  
*athweel ORF* 1351 CTCAATATCAAAGAAGGAAAACCTCCTCTCCTTCTGGCCATTCGTTGCA  
*1s contig* 1401 GTTACAACAACCTTCTTAAGACAATGATGGATCGTGATCCGAAGCGTCGGC  
*athweel ORF* 1401 GTTACAACAACCTTCTTAAGACAATGATGGATCGTGATCCGAAGCGTCGGC  
*1s contig* 1451 CTTCTGCTAGAGAATTACTGGACCATCCCATGTTTGATAGGATTTCAGGT  
*athweel ORF* 1451 CTTCTGCTAGAGAATTACTGGACCATCCCATGTTTGATAGGATTTCAGGT  
*1s contig* 1501 TGA  
*athweel ORF* 1501 TGA

### pTA7002-*AtWEE1* clone 1X

*athweel ORF* 1 ATGTTGAGAGAAGAACGGAAGAACAACACTGTTGGCGAAGAGGAAAACCCAAGG  
*1x contig* 1 ATGTTGAGAGAAGAACGGAAGAACAACACTGTTGGCGAAGAGGAAAACCCAAGG  
*athweel ORF* 51 GACAAATCAAACCAGGGCATCGAAGAAGATTCGGAAGATGGAAGGGACAT  
*1x contig* 51 GACAAATCAAACCAGGGCATCGAAGAAGATTCGGAAGATGGAAGGGACAT  
*athweel ORF* 101 TGGAGCGTCACTCTCTGCTTCAATTCGGTCAATTGTGCAAGATTTCTTTC  
*1x contig* 101 TGGAGCGTCACTCTCTGCTTCAATTCGGTCAATTGTGCAAGATTTCTTTC  
*athweel ORF* 151 GAAAATCSTCCGTCGTGGAATGTTGCTTCATCGGCGTTTCAGGGTCTCCT  
*1x contig* 151 GAAAATCSTCCGTCGTGGAATGTTGCTTCATCGGCGTTTCAGGGTCTCCT  
*athweel ORF* 201 GGATTCGGATTCTTCGGAACCTCGAAATCAGTTGGGTTCCGCTGATTTCAG  
*1x contig* 201 GGATTCGGATTCTTCGGAACCTCGAAATCAGTTGGGTTCCGCTGATTTCAG  
*athweel ORF* 251 ATGCCAATTGCGGAGAGAAGGACTTTATTCTTAGCCAAGACTTCTTCTGC  
*1x contig* 251 ATGCCAATTGCGGAGAGAAGGACTTTATTCTTAGCCAAGACTTCTTCTGC  
*athweel ORF* 301 ACACCTGACTATATAACCCCGACAATCAGAA-CTTGATGAGCGGCTTAG  
*1x contig* 301 ACACCTGACTATATAACCCCGACAATCAGAA-CTTGATGAGCGGCTTAG  
*athweel ORF* 350 ATATCAGCAAGGATCATTCTCCATGTCTTAGGTCTCCTGTAAACTAAAT  
*1x contig* 351 ATATCAGCAAGGATCATTCTCCATGTCTTAGGTCTCCTGTAAACTAAAT  
*athweel ORF* 400 ACAGTTAAAAGCAAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATTTC  
*1x contig* 401 ACAGTTAAAAGCAAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATTTC  
*athweel ORF* 450 AAATCTACCTGGTCTTCAAAAACATAGAGTAGATGAACAAAGAGAAATGATG  
*1x contig* 451 AAATCTACCTGGTCTTCAAAAACATAGAGTAGATGAACAAAGAGAAATGATG  
*athweel ORF* 500 ATATTGACACAGATGAGGTGATGGGGGATAAACTCCAAGCTAATCAAACA  
*1x contig* 501 ATATTGACACAGATGAGGTGATGGGGGATAAACTCCAAGCTAATCAAACA  
*athweel ORF* 550 GAGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTCGGGC  
*1x contig* 551 GAGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTCGGGC  
*athweel ORF* 600 TATGCCA-CCCCCTTGCCCTCAAGAATCCGTATGTGTTGAATCAGTCTGAGA  
*1x contig* 601 TATGCCMCCCCCTTGCCCTCAAGAATCCGTATGTGTTGAATCAGTCTGAGA  
*athweel ORF* 650 CTGCTACTGACCCTTTTGGACATCAGAGATCAAATGTGCAAGTTTCTC  
*1x contig* 651 CTGCTACTGACCCTTTTGGACATCAGAGATCAAATGTGCAAGTTTCTC

*athwee1* ORF 700 CCTGTAAGTACAAGTGGGGATGGCTTGTCAAGATATCTCACGGACTTCA  
*lx contig* 701 CCTGTAAGTACAAGTGGGGATGGCTTGTCAAGATATCTCACGGACTTCA  
  
*athwee1* ORF 750 TGAATCCGGCAAATTGGTGCTGGACATTTTCAGTCGGGTATTTAAGGTGT  
*lx contig* 751 TGAATCCGGCAAATTGGTGCTGGACATTTTCAGTCGGGTATTTAAGGTGT  
  
*athwee1* ORF 800 TGAAGAGAATGGACGGTTCCTATATGCTGTGAAACACAGCACAAAGAAAG  
*lx contig* 801 TGAAGAGAATGGACGGTTCCTATATGCTGTGAAACACAGCACAAAGAAAG  
  
*athwee1* ORF 850 TTGTATCTAGATTCAGAGAGACGTAAAGCTATGATGGAAGTCAAGCTCT  
*lx contig* 851 TTGTATCTAGATTCAGAGAGACGTAAAGCTATGATGGAAGTCAAGCTCT  
  
*athwee1* ORF 900 TGCTGCTCTTGGGTTTCATGAAAATATAGTAGGATATTACTCCTCGTGGT  
*lx contig* 901 TGCTGCTCTTGGGTTTCATGAAAATATAGTAGGATATTACTCCTCGTGGT  
  
*athwee1* ORF 950 TTGAAAATGAGCAATTATACATTCAACTGGAACCTCGCGATCACAGCTTG  
*lx contig* 951 TTGAAAATGAGCAATTATACATTCAACTGGAACCTCGCGATCACAGCTTG  
  
*athwee1* ORF 1000 TCAGCTTTGCCTAAGAAATCTT-CTCTTAAAGTCTCAGAAAGAGAGATCT  
*lx contig* 1001 TCAGCTTTGCCTAAGAAATCTT-CTCTTAAAGTCTCAGAAAGAGAGATCT  
  
*athwee1* ORF 1049 TGGTGATTATGCATCAGATAGCAAAGGCATTACAT-TTTGTCATGAGAA  
*lx contig* 1051 TGGTGATTATGCATCAGATAGCAAAGGCATTACAT-TTTGTCATGAGAA  
  
*athwee1* ORF 1098 AGGAATAGCTCATTTAGATGTAAAACCTGACAATATTTACATTAAGAACC  
*lx contig* 1101 AGGAATAGCTCATTTAGATGTAAAACCTGACAATATTTACATTAAGAACC  
  
*athwee1* ORF 1148 GTGTTTGCAAGCTTGGTGACTTTGGTTGTGCCACACGATTGGACAAAAGC  
*lx contig* 1151 GTGTTTGCAAGCTTGGTGACTTTGGTTGTGCCACACGATTGGACAAAAGC  
  
*athwee1* ORF 1198 TTACCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAAATTCATAA  
*lx contig* 1201 TTACCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAAATTCATAA  
  
*athwee1* ORF 1248 TGAAGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTTAGGTGTGA  
*lx contig* 1251 TGAAGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTTAGGTGTGA  
  
*athwee1* ORF 1298 CGGTTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAAGAAACCAG  
*lx contig* 1301 CGGTTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAAGAAACCAG  
  
*athwee1* ORF 1348 TCGCTCAATATCAAAGA-AGGAAAACCTCCTCTCCTCCTGGCCATTCGT  
*lx contig* 1351 TCGCTCAATATCAAAGA-AGGAAAACCTCCTCTCCTCCTGGCCATTCGT  
  
*athwee1* ORF 1397 TGCAGTTACAACAACCTCTTAAGACAATGATGGATCGTGATCCGAAGCGT  
*lx contig* 1401 TGCAGTTACAACAACCTCTTAAGACAATGATGGATCGTGATCCGAAGCGT  
  
*athwee1* ORF 1447 CGGCCTTCTCCTAGAGAATTACTGGACCATCCCATGTTTGATAGGATTCCG  
*lx contig* 1451 CGGCCTTCTCCTAGAGAATTACTGGACCATCCCATGTTTGATAGGATTCCG  
  
*athwee1* ORF 1497 AGGTTGA  
*lx contig* 1501 AGGTTGA

## pTA7002-*AtWEE1* clone 2X

*athwee1* ORF 1 ATGTTTCGAGAAGAACGGAAGAACAACACTGTTGGCGAAGAGGAAAACCCAAGG  
*2x contig* 1 ATGTTTCGAGAAGAACGGAAGAACAACACTGTTGGCGAAGAGGAAAACCCAAGG  
  
*athwee1* ORF 51 GACAAATCAAAAACGAGGCATCGAAGAAGATTCGGAAGATGGAAGGGACAT  
*2x contig* 51 GACAAATCAAAAACGAGGCATCGAAGAAGATTCGGAAGATGGAAGGGACAT  
  
*athwee1* ORF 101 TGGAGCGTCACTCTCTGCTTCAATTCGGTCAATTGTGCGAAGATTCTTTT  
*2x contig* 101 TGGAGCGTCACTCTCTGCTTCAATTCGGTCAATTGTGCGAAGATTCTTTT  
  
*athwee1* ORF 151 GAAAATCGTCCGTCGTCGAATGTGCTTCATCGGCGTTTCAGGGTCTCCT  
*2x contig* 151 GAAAATCGTCCGTCGTCGAATGTGCTTCATCGGCGTTTCAGGGTCTCCT  
  
*athwee1* ORF 201 GGATTCGGATTCTTCGGAACCTCGGAAATCAGTTGGGTTCCGCTGATTTCAG  
*2x contig* 201 GGATTCGGATTCTTCGGAACCTCGGAAATCAGTTGGGTTCCGCTGATTTCAG  
  
*athwee1* ORF 251 ATGCCAATTGCGGAGAGAAGGACTTTATTCTTAGCCAAGACTTCTTCTGC  
*2x contig* 251 ATGCCAATTGCGGAGAGAAGGACTTTATTCTTAGCCAAGACTTCTTCTGC

**athwee1 ORF** 301 ACACCTGACTATATAACCCCGGACAATCAGAA-CTTGATGAGCGGCTTAC  
**2x contig** 301 ACACCTGACTATATAACCCCGGACAATCAGAA-CTTGATGAGCGGCTTAC

**athwee1 ORF** 350 ATATCAGCAAGGATCATTCTCCATGTCCTAGGTCTCCTGTAAACTAAAT  
**2x contig** 350 ATATCAGCAAGGATCATTCTCCATGTCCTAGGTCTCCTGTAAACTAAAT

**athwee1 ORF** 400 ACAGTTAAAAGCAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATT  
**2x contig** 400 ACAGTTAAAAGCAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATT

**athwee1 ORF** 450 AAATTCTACCTGGTCTTCAAAACATAGAGTAGATGAACAAGAGAATGATG  
**2x contig** 450 AAATTCTACCTGGTCTTCAAAACATAGAGTAGATGAACAAGAGAATGATG

**athwee1 ORF** 500 ATATTGACACAGATGAGGTGATGGGGATAAACTCCAAGCTAATCAAACA  
**2x contig** 500 ATATTGACACAGATGAGGTGATGGGGATAAACTCCAAGCTAATCAAACA

**athwee1 ORF** 550 GAGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTGCGGC  
**2x contig** 550 GAGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTGCGGC

**athwee1 ORF** 600 TATGCCACCCCTTGCCTCAAGAATCCGTATGTGTTGAATCAGTCTGAGA  
**2x contig** 600 TATGCCACCCCTTGCCTCAAGAATCCGTATGTGTTGAATCAGTCTGAGA

**athwee1 ORF** 650 CTGCTACTGACCCTTTTGGACATCAGAGATCAAAATGTGCAAGTTTCTC  
**2x contig** 650 CTGCTACTGACCCTTTTGGACATCAGAGATCAAAATGTGCAAGTTTCTC

**athwee1 ORF** 700 CCTGTAAGTACAAGTGGGGATGGCTTGTCAAGATATCTCAGGACTTTCA  
**2x contig** 700 CCTGTAAGTACAAGTGGGGATGGCTTGTCAAGATATCTCAGGACTTTCA

**athwee1 ORF** 750 TGAATCCGGCAAATGGTGTGACATTCAGTCCGGTATTTAAGGTGT  
**2x contig** 750 TGAATCCGGCAAATGGTGTGACATTCAGTCCGGTATTTAAGGTGT

**athwee1 ORF** 800 TGAAGAGAATGGACGGTGCCTATATGCTGTGAAACACAGCACAAGAAAG  
**2x contig** 800 TGAAGAGAATGGACGGTGCCTATATGCTGTGAAACACAGCACAAGAAAG

**athwee1 ORF** 850 TTGTATCTAGATTCAGAGAGACGTAAGCTATGATGGAAGTGCAAGCTCT  
**2x contig** 850 TTGTATCTAGATTCAGAGAGACGTAAGCTATGATGGAAGTGCAAGCTCT

**athwee1 ORF** 900 TGCTGCTCTTGGGTTTCATGAAAATATAGTAGGATATTACTCCTCGTGGT  
**2x contig** 900 TGCTGCTCTTGGGTTTCATGAAAATATAGTAGGATATTACTCCTCGTGGT

**athwee1 ORF** 950 TTGAAAATGAGCAATTATACATTCACCTGGAACCTCGCATCACAGCTTG  
**2x contig** 950 TTGAAAATGAGCAATTATACATTCACCTGGAACCTCGCATCACAGCTTG

**athwee1 ORF** 1000 TCAGCTTTGCCCTAAGAAATCTT-CTCTTAAAGTCTCAGAAAGAGAGATCT  
**2x contig** 1000 TCAGCTTTGCCCTAAGAAATCTT-CTCTTAAAGTCTCAGAAAGAGAGATCT

**athwee1 ORF** 1049 TGGTGATTATGCATCAGATAGCAAAGGCATTACAT-TTTGTGCATGAGAA  
**2x contig** 1049 TGGTGATTATGCATCAGATAGCAAAGGCATTACAT-TTTGTGCATGAGAA

**athwee1 ORF** 1098 AGGAATAGCTCATTTAGATGTAAAACCTGACAATATTTACATTAAGAACC  
**2x contig** 1098 AGGAATAGCTCATTTAGATGTAAAACCTGACAATATTTACATTAAGAACC

**athwee1 ORF** 1148 GTGTTTGCAAGCTTGGTGACTTTGGTTGTGCCACACGATTGGACAAAAGC  
**2x contig** 1148 GTGTTTGCAAGCTTGGTGACTTTGGTTGTGCCACACGATTGGACAAAAGC

**athwee1 ORF** 1198 TTACCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAAATCTAAA  
**2x contig** 1198 TTACCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAAATCTAAA

**athwee1 ORF** 1248 TGAAGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTTAGGTGTGA  
**2x contig** 1248 TGAAGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTTAGGTGTGA

**athwee1 ORF** 1298 CGGTTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAAGAAACCAG  
**2x contig** 1298 CGGTTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAAGAAACCAG

**athwee1 ORF** 1348 TCGCTCAATATCAAAGA-AGGAAAACCTCCTCTCCTTCTGGCCATTCTG  
**2x contig** 1348 TCGCTCAATATCAAAGA-AGGAAAACCTCCTCTCCTTCTGGCCATTCTG

**athwee1 ORF** 1397 TGCAGTTACAACAACCTCTTAAGACAATGATGGATCGTGATCCGAAGCGT  
**2x contig** 1397 TGCAGTTACAACAACCTCTTAAGACAATGATGGATCGTGATCCGAAGCGT

**athwee1 ORF** 1447 CGGCCTTCTGCTAGAGAATTACTGGACCATCCCATGTTTGATAGGATTCCG  
**2x contig** 1447 CGGCCTTCTGCTAGAGAATTACTGGACCATCCCATGTTTGATAGGATTCCG

**athwee1 ORF** 1497 AGGTTGA  
**2x contig** 1497 AGGTTGA

**pTA7002-*AtWEE1* clone 3X**

*athwee1* ORF 1 ATGTTTCGAGAAGAACCGGAAGAACA  
3x contig 1 ATGTTTCGAGAAGAACCGGAAGAACA

*athwee1* ORF 51 GACAATCAAACCAGGGCATCGAAGAAGATTCGGAAGATGGAAGGGACAT  
3x contig 51 GACAATCAAACCAGGGCATCGAAGAAGATTCGGAAGATGGAAGGGACAT

*athwee1* ORF 101 TGGAGCGTCACTCTCTGCTTCAATTCGGTCAATTGTCGAAGATTTCTTTC  
3x contig 101 TGGAGCGTCACTCTCTGCTTCAATTCGGTCAATTGTCGAAGATTTCTTTC

*athwee1* ORF 151 GAAAATCGTCCGTCGTCGAATGTTGCTTCATCGGCGTTTCAGGGTCTCCT  
3x contig 151 GAAAATCGTCCGTCGTCGAATGTTGCTTCATCGGCGTTTCAGGGTCTCCT

*athwee1* ORF 201 GGATTCGGATTCTTCGGAACTCCGAAATCAGTTGGGTTCGGCTGATTCAG  
3x contig 201 GGATTCGGATTCTTCGGAACTCCGAAATCAGTTGGGTTCGGCTGATTCAG

*athwee1* ORF 251 ATGCCAATTGCGGAGAGAAGGACTTTATTTCTTAGCCAAGACTTCTTCTGC  
3x contig 251 ATGCCAATTGCGGAGAGAAGGACTTTATTTCTTAGCCAAGACTTCTTCTGC

*athwee1* ORF 301 ACACCTGACTATATAACCCCGGACAATCAGAA-CTTGATGAGCGGCTTAG  
3x contig 301 ACACCTGACTATATAACCCCGGACAATCAGAA-CTTGATGAGCGGCTTAG

*athwee1* ORF 350 ATATCAGCAAGGATCATTCTCCATGTCTTAGGTCTCTGTTAAACTAAAT  
3x contig 350 ATATCAGCAAGGATCATTCTCCATGTCTTAGGTCTCTGTTAAACTAAAT

*athwee1* ORF 400 ACAGTTAAAAGCAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATT  
3x contig 400 ACAGTTAAAAGCAAAGATGTCGCCAGGAGAGTTTCACAGGAAATCATT

*athwee1* ORF 450 AAATTCTACCTGGTCTTCAAACATAGAGTAGATGAACAAGAGAATGATG  
3x contig 450 AAATTCTACCTGGTCTTCAAACATAGAGTAGATGAACAAGAGAATGATG

*athwee1* ORF 500 ATATTGACACAGATGAGGTGATGGGGATAAACTCCAAGCTAATCAAACA  
3x contig 500 ATATTGACACAGATGAGGTGATGGGGATAAACTCCAAGCTAATCAAACA

*athwee1* ORF 550 GAGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTCGGGC  
3x contig 550 GAGAGGACTGGATACGTTTCACAGGCTGCAGTTGCTCTGCGGTGTCGGGC

*athwee1* ORF 600 TATGCCACCCCTTGCCCTCAAGAATCCGTATGTGTTGAATCAGTCTGAGA  
3x contig 600 TATGCCACCCCTTGCCCTCAAGAATCCGTATGTGTTGAATCAGTCTGAGA

*athwee1* ORF 650 CTGCTACTGACCCTTTTGGACATCAGAGATCAAATGTGCAAGTTTCTC  
3x contig 650 CTGCTACTGACCCTTTTGGACATCAGAGATCAAATGTGCAAGTTTCTC

*athwee1* ORF 700 CCTGTAAGTACAAGTGGGGATGGCTTGTCAGATATCTCACGGACTTTCA  
3x contig 700 CCTGTAAGTACAAGTGGGGATGGCTTGTCAGATATCTCACGGACTTTCA

*athwee1* ORF 750 TGAATCCGGCAAATGGTGTGACATTTAGTCCGGTATTTAAGGKTGT  
3x contig 750 TGAATCCGGCAAATGGTGTGACATTTAGTCCGGTATTTAAGGKTGT

*athwee1* ORF 800 T-GAAGAGAATGGAC-GGTTGCCTATATGCTGTGAAAC-ACAGCACAAGA  
3x contig 800 TTGAAGAGAATGGACCGGTTGCCTATATGCTGTGAAAKCACAGCACAAGA

*athwee1* ORF 847 AAGTTGTATCTAGATTGAC-AGAGACCTAAAGCTATGATGGAAGTCAAG  
3x contig 850 AAGTTGTATCTAGATTGACAGAGACCTAAAGCTATGATGGAAGTCAAG

*athwee1* ORF 896 CTCTTGCTGCTCTTGGGTTT-CATGAAAATATAGTAGGATATTACTCC-T  
3x contig 900 CTCTTGCTGCTCTTGGGTTT-CATGAAAATATAGTAGGATATTACTCCCT

*athwee1* ORF 944 CGTGGTTTGAATGAGCAATTATACATTCACCTGGAACCTGCGGATCAC  
3x contig 950 CGTGGTTTGAATGAGCAATTATACATTCACCTGGAACCTGCGGATCACXX

*athwee1* ORF 994 A--GCTTGTGACCTTTGCCTAAGAAATCTT-CTCTTAAAGTCTCAGAAAG  
3x contig 1000 ACAGCTTGTGACCTTTGCCTAAGAAATCTT-CTCTTAAAGTCTCAGAAAG

*athwee1* ORF 1041 AGAGATCTTGGTGATTATGCATCAGATAGCAAAGGCATTACAT-TTTGTG  
3x contig 1049 AGAGATCTTGGTGATTATGCATCAGATAGCAAAGGCATTACAT-TTTGTG

*athwee1* ORF 1090 CATGAGAAAGGAATAGCTCATTAGATGTAAAACCTGACAATATTTACAT  
3x contig 1098 CATGAGAAAGGAATAGCTCATTAGATGTAAAACCTGACAATATTTACAT

*athwee1* ORF 1140 TAAGAACGGTGTGTTGCAAGCTTGGTGACTTTGGTGTGCCACACGATTGG  
3x contig 1148 TAAGAACGGTGTGTTGCAAGCTTGGTGACTTTGGTGTGCCACACGATTGG

**athweel ORF** 1190 **ACAAAAGCTTACCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAA**  
**3x contig** 1198 **ACAAAAGCTTACCAGTAGAAGAAGGAGATGCACGTTACATGCCTCAAGAA**

**athweel ORF** 1240 **ATTCTAAATGAAGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTT**  
**3x contig** 1248 **ATTCTAAATGAAGACTACGAACACCTTGATAAAGTCGATATCTTCTCTTT**

**athweel ORF** 1290 **AGGTGTGACGGTTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAA**  
**3x contig** 1298 **AGGTGTGACGGTTTATGAGCTGATTAAGGGATCTCCTCTTACAGAATCAA**

**athweel ORF** 1340 **GAAACCAGTCGCTCAATATCAAAGA-AGGAAAACCTCCTCCTTCCTGG**  
**3x contig** 1348 **GAAACCAGTCGCTCAATATCAAAGA-AGGAAAACCTCCTCCTTCCTGG**

**athweel ORF** 1389 **CCATTTCGTTGCAGTTACAACAACCTCTTAAGACAATGATGGATCGTGATC**  
**3x contig** 1397 **CCATTTCGTTGCAGTTACAACAACCTCTTAAGACAATGATGGATCGTGATC**

**athweel ORF** 1439 **CGAAGCGTCGGCCTTCTGCTAGAGAATTACTGGACCATCCCATGTTTGAT**  
**3x contig** 1447 **CGAAGCGTCGGCCTTCTGCTAGAGAATTACTGGACCATCCCATGTTTGAT**

**athweel ORF** 1489 **AGGATTCGAGGTTGA**  
**3x contig** 1497 **AGGATTCGAGGTTGA**



## **Appendix II: Tobacco BY-2 cell area raw data**

### **II.I. Absorbance at 550 nm**

| WEE1-c-2 | WEE1-c-10 | WEE1-c-12 | WT   | EV-c-10 |
|----------|-----------|-----------|------|---------|
| 0.18     | 0.18      | 0.24      | 0.3  | 0.12    |
| 0.23     | 0.24      | 0.32      | 0.38 | 0.2     |
| 0.3      | 0.32      | 0.41      | 0.53 | 0.34    |
| 0.45     | 0.52      | 0.56      | 0.78 | 0.58    |
| 0.9      | 1.36      | 1.22      | 1.68 | 1.28    |
| 1.74     | 2.54      | 2.32      | 2.61 | 2.32    |
| 2.4      | 2.65      | 2.46      | 2.73 | 2.49    |

## II.II. Mitotic cell area ( $\mu\text{m}^2$ ) over a 6 day period

### Wild type

| day 1    | day 2 | day 3    | day 4    | day 5    | day 6    |
|----------|-------|----------|----------|----------|----------|
| 2728.809 | 23271 | 1104.147 | 1627.609 | 2487.158 | 5310.475 |
| 3454.631 | 43357 | 1393.582 | 2093.58  | 2977.876 | 4610.181 |
| 3159.186 | 29448 | 916.0443 | 1749.683 | 3418.77  | 4006.447 |
| 2233.383 | 37656 | 1045.074 | 1713.432 | 4033.447 | 2919.304 |
| 2732.96  | 28219 | 2187.803 | 1624.127 | 2166.912 | 2419.246 |
| 1916.657 | 34153 | 3001.764 | 2407.508 | 3347.721 | 4145.916 |
| 3241.82  | 21813 | 2235.466 | 2056.799 | 2830.285 | 2459.58  |
| 4113.514 | 27418 | 1453.331 | 2219.969 | 2922.257 | 3685.662 |
| 2570.711 | 26646 | 1922.837 | 2084.347 | 2609.16  | 2763.49  |
| 2516.074 | 32882 | 1049.127 | 1924.583 | 2495.061 | 5221.254 |
| 2737.186 | 39532 | 1872.847 | 1956.597 | 3153.316 | 3658.344 |
| 1797.422 | 41535 | 1971.702 | 1890.148 | 4043.231 | 6001.461 |
| 4338.098 | 21098 | 1774.367 | 2842.98  | 3902.338 | 3782.753 |
| 4154.114 | 34135 | 1746.969 | 2386.847 | 4684.326 |          |
| 3624.804 | 46274 | 1936.573 | 2502.866 | 2829.908 |          |
| 3904.628 | 20409 | 1504.372 | 2204.605 | 3101.911 |          |
| 3565.715 | 25538 | 1762.732 | 1671.807 | 2700.982 |          |
| 3578.921 | 33015 | 1759.279 | 1596.882 | 2761.418 |          |
| 4100.157 | 31166 | 1696.153 | 1924.432 | 3079.557 |          |
| 3044.102 | 24788 | 1678.664 | 1718.578 | 3079.407 |          |
| 4782.812 | 30506 | 1776.393 | 2363.991 | 3853.492 |          |
| 5627.415 | 25973 | 2353.462 | 2055.21  | 3061.645 |          |
| 3165.147 | 26283 | 1799.587 | 2022.061 | 3428.404 |          |
| 3667.668 | 38904 | 2394.521 | 1753.619 | 4565.033 |          |
| 2662.627 | 23658 | 1542.203 | 1718.578 | 3013.175 |          |
| 3210.879 | 33907 | 2105.461 | 1625.565 | 3352.162 |          |
| 3028.48  | 39469 | 1706.512 | 1273.04  | 3686.934 |          |
| 3509.418 | 25930 | 1735.86  | 1937.601 |          |          |
| 2151.805 | 30990 | 1241.659 | 2570.072 |          |          |
| 2361.824 | 28682 | 1980.784 | 1619.586 |          |          |
| 3197.371 | 23129 | 1293.526 |          |          |          |
|          | 33260 | 1451.304 |          |          |          |
|          | 31621 | 1493.789 |          |          |          |
|          | 20184 | 2200.563 |          |          |          |
|          | 19573 | 1735.035 |          |          |          |
|          | 19102 | 2106.512 |          |          |          |
|          | 21938 | 1939.276 |          |          |          |
|          | 24662 | 1903.321 |          |          |          |
|          |       | 2581.197 |          |          |          |
|          |       | 1995.872 |          |          |          |
|          |       | 2020.717 |          |          |          |
|          |       | 1577.482 |          |          |          |
|          |       | 1807.994 |          |          |          |
|          |       | 1648.865 |          |          |          |
|          |       | 1689.548 |          |          |          |
|          |       | 1659.974 |          |          |          |

**EV-c-10**

| day 1    | day 2    | day 3    | day 4    | day 5    | day 6    |
|----------|----------|----------|----------|----------|----------|
| 4833.476 | 2291.097 | 1978.136 | 3115.138 | 1322.453 | 3173.991 |
| 4835.127 | 1836.656 | 2389.384 | 2569.084 | 1558.124 | 2872.782 |
| 4391.687 | 2032.341 | 1732.856 | 1722.459 | 2325.758 | 3043.708 |
| 3571.384 | 2347.996 | 1857.199 | 2248.233 | 1916.323 | 3767.003 |
| 4346.503 | 1504.969 | 2086.813 | 2349.331 | 2212.237 | 3133.123 |
| 5087.77  | 1917.489 | 1993.272 | 1759.463 | 2815.718 | 5069.611 |
| 2532.364 | 2689.391 | 1908.283 | 2288.036 | 3129.568 | 3948.617 |
| 5805.32  | 1899.425 | 2895.762 | 2060.716 | 3022.025 | 4598.603 |
| 5833.691 | 4052.112 | 1793.022 | 2369.838 | 2757.065 | 4542.757 |
| 2668.368 | 3254.997 | 2230.909 | 2250.427 | 3119.427 | 5031.15  |
| 1656.218 | 5643.56  | 4155.009 | 4138.231 | 2589.204 | 2852.837 |
| 8372.735 | 3683.547 | 2389.384 | 3626.533 | 3003.786 | 4889.878 |
| 3950.875 | 3816.914 | 2497.455 | 4631.238 | 2014.255 | 4728.209 |
| 4911.16  | 2208.683 | 2439.56  | 3310.449 | 2014.709 | 4219.645 |
| 5085.594 | 2997.595 | 2370.993 | 3949.806 | 3451.894 |          |
| 4763.147 | 2955.598 | 2254.445 | 2372.563 | 3431.839 |          |
| 5162.678 | 2221.628 | 2189.663 | 2572.868 | 2070.788 |          |
| 4369.095 | 2320.525 | 3838.513 | 3794.829 | 3119.048 |          |
| 4065.938 | 2021.879 | 2523.187 | 3370.987 | 2298.589 |          |
| 5137.834 | 2478.051 | 2141.454 | 4162.522 | 3214.18  |          |
|          | 2632.868 | 2377.426 | 2119.892 | 2637.338 |          |
|          | 5798.151 | 1537.6   | 1948.645 | 3044.275 |          |
|          | 3053.591 | 2884.031 | 4507.059 | 2234.865 |          |
|          | 3852.514 | 2348.289 | 1793.062 | 2062.615 |          |
|          | 2087.81  | 2373.037 | 1781.711 | 4058.555 |          |
|          | 3194.409 | 1839.641 | 2029.917 | 3277.903 |          |
|          | 3411.168 | 2281.388 | 3373.862 | 2483.402 |          |
|          | 2372.607 | 2262.997 | 4479.893 | 3313.019 |          |
|          | 2777.525 | 2523.187 | 3876.177 | 2806.334 |          |
|          | 2840.52  | 2010.149 | 1833.093 | 2229.189 |          |
|          | 2208.608 | 2590.996 | 2040.965 | 2096.066 |          |
|          | 1863.525 |          | 1993.821 | 2154.265 |          |
|          | 2748.925 |          | 2389.135 | 2339.835 |          |
|          | 2760.967 |          | 2113.081 | 2917.509 |          |
|          | 2547.52  |          | 1842.779 | 2537.741 |          |
|          |          |          | 2117.168 | 2351.868 |          |
|          |          |          |          | 2345.738 |          |
|          |          |          |          | 1558.351 |          |
|          |          |          |          | 2316.752 |          |
|          |          |          |          | 2374.043 |          |
|          |          |          |          | 1845.183 |          |
|          |          |          |          | 1730.526 |          |
|          |          |          |          | 1535.344 |          |
|          |          |          |          | 3112.843 |          |
|          |          |          |          | 3189.281 |          |

**WEE1-c-10**

| day 1    | day 2    | day 3    | day 4    | day 5    | day 6    |
|----------|----------|----------|----------|----------|----------|
| 5643.89  | 3443.532 | 2172.483 | 1470.318 | 4406.234 | 1911.44  |
| 3326.143 | 2491.298 | 1457.454 | 1875.848 | 1818.013 | 2149.86  |
| 1843.235 | 3560.416 | 3533.672 | 4299.792 | 1367.862 | 2188.88  |
| 1730.112 | 2363.801 | 1766.912 | 1820.077 | 1446.419 | 2397.035 |
| 1801.764 | 5560.243 | 1793.627 | 2310.435 | 1662.791 | 1859.74  |
| 2364.519 | 5184.527 | 1853.793 | 2433.176 | 2769.401 | 1824.872 |
| 2191.861 | 3576.071 | 1445.194 | 1905.284 | 2265.062 | 2403.526 |
| 2180.195 | 3266.136 | 1926.976 | 2563.712 | 2290.415 | 1466.524 |
| 2189.152 | 2386.155 | 2335.272 | 2129.729 | 1209.537 | 1686.831 |
| 1865.588 | 1623.735 | 2604.922 | 1425.596 | 1964.456 | 2028.65  |
| 2308.07  | 1938.186 | 2152.277 | 2054.586 | 1839.053 | 2910.632 |
| 2612.215 | 3282.618 | 1749.808 | 2312.1   | 3664.483 | 2450.32  |
| 2306.188 | 3100.255 | 2054.649 | 1303.006 | 1963.321 | 2124.35  |
| 2318.908 | 3404.846 | 2488.676 | 3100.609 | 2424.749 | 1280.935 |
| 1962.304 | 3053.215 | 1910.402 | 3580.675 | 1871.066 | 1520.714 |
| 2424.731 | 4210.392 | 1397.137 | 1170.731 | 2056.712 | 1850.608 |
|          | 2776.095 | 1825.413 | 1925.867 | 1608.982 | 2665.796 |
|          | 3286.381 | 2442.209 | 3596.188 | 1991.247 | 1997.328 |
|          | 3412.749 | 1488.408 | 1333.578 | 1489.33  | 1527.582 |
|          | 2117.991 | 2367.739 | 2568.479 | 1255.702 | 2475.226 |
|          | 2462.547 | 7155.361 | 1916.257 | 1768.745 | 2447.603 |
|          | 2204.619 | 1241.462 | 1467.443 | 2395.915 | 2166.162 |
|          | 3959.011 | 2945.862 | 2671.394 | 3848.009 | 2079.594 |
|          | 4806.78  | 2110.955 | 4309.251 | 1750.43  | 1892.496 |
|          |          | 2131.086 | 1736.232 | 1912.615 | 3203.846 |
|          |          | 2203.739 | 1859.502 | 1668.467 | 3084.9   |
|          |          | 2326.115 | 1644.819 | 2172.277 | 2922.104 |
|          |          | 1563.785 | 1673.348 | 1837.312 | 1987.819 |
|          |          | 1956.491 | 1924.354 | 1491.676 | 2760.062 |
|          |          | 2410.044 | 3444.843 | 2260.9   | 2493.717 |
|          |          | 2113.301 | 3299.628 | 1873.109 | 1934.006 |
|          |          | 1698.572 | 2159.771 | 2101.212 | 1108.402 |
|          |          | 3117.203 | 1930.029 | 1971.495 |          |
|          |          | 1525.869 | 2079.407 | 1998.816 |          |
|          |          | 2926.715 | 1388.214 | 1535.571 |          |
|          |          | 2761.504 | 1460.33  | 1554.34  |          |
|          |          | 2045.492 | 3312.265 | 1617.837 |          |
|          |          | 2500.785 | 1740.848 | 1648.942 |          |
|          |          | 2915.665 | 1396.613 |          |          |
|          |          |          | 1109.284 |          |          |
|          |          |          | 1645.5   |          |          |

## II.III. Mitotic cell area ( $\mu\text{m}^2$ ) following synchronization

### Wild type

|          |          |          |          |          |          |
|----------|----------|----------|----------|----------|----------|
| 2938.952 | 2801.709 | 2799.265 | 3921.299 | 3311.524 | 3664.139 |
| 3486.39  | 3637.401 | 2866.189 | 3431.628 | 2212.048 | 2203.063 |
| 2857.107 | 3724.45  | 2095.968 | 3276.592 | 2856.629 | 2349.259 |
| 2770.659 | 3324.533 | 3100.209 | 3106.168 | 2644.086 | 3805.702 |
| 3675.291 | 3169.785 | 2155.697 | 3170.784 | 2811.7   | 2906.511 |
| 3320.573 | 3381.595 | 2109.984 | 2850.441 | 2859.066 | 3830.46  |
| 2607.429 | 3651     | 2721     | 3581.698 | 3557.116 |          |
| 3620.295 | 2813.073 | 3097.047 | 2667.158 | 2368.521 |          |
| 3458.215 | 2814.862 | 2760.46  | 2738.315 | 3872.147 |          |
| 2655.658 | 3132.769 | 2572.002 | 3493.937 | 3226.495 |          |
| 2894.932 | 3152.678 | 2370.749 | 3334.948 | 3029.915 |          |
| 3412.646 | 2813.856 | 2234.689 | 2832.982 | 2579.836 |          |
| 3318.273 | 2980.033 | 2192.066 | 3651.13  | 2302.108 |          |
| 3555.319 | 3245.47  | 3198.32  | 3133.344 | 2728.331 |          |
| 3498.465 | 2280.474 | 3031.28  | 3114.153 | 3706.043 |          |
| 2502.354 | 1868.769 | 2344.155 | 3588.667 | 3410.712 |          |
| 3159.931 | 2870.279 | 3153.325 | 3739.289 | 2870.782 |          |
| 3567.538 | 3024.955 | 3829.983 | 3357.021 | 2837.647 |          |
| 2677.364 | 3425.95  | 3533.627 | 2960.719 | 2542.094 |          |
| 2815.222 | 3139.956 | 3378.304 | 3810.622 | 3257.976 |          |

### Constitutive lines

| EV-c-10  | EV-c-10<br>(-)Hyg | WEE1-c-2 | WEE1-c-10 | WEE1-c-10<br>(-)Hyg | WEE1-c-12 | WEE1-c-12<br>(-)Hyg |
|----------|-------------------|----------|-----------|---------------------|-----------|---------------------|
| 3734.324 | 5442.708          | 1403.238 | 4110.473  | 1903.277            | 2301.642  | 3124.107            |
| 2040.267 | 2513.281          | 1723.695 | 1771.888  | 1331.985            | 2935.352  | 5040.923            |
| 2645.394 | 3486.143          | 2153.839 | 1900.333  | 1565.686            | 1615.287  | 2953.240            |
| 4435.438 | 2286.724          | 1852.335 | 1808.289  | 1517.366            | 2797.007  | 2703.227            |
| 3265.23  | 2247.798          | 1739.223 | 1623.339  | 1597.779            | 2760.397  | 2587.859            |
| 3794.874 | 2757.833          | 3535.840 | 2400.115  | 1218.976            | 2806.235  | 3663.969            |
| 5234.25  | 1635.214          | 2492.869 | 2707.049  | 1651.053            | 2989.358  | 2259.981            |
| 3331.129 | 2321.359          | 1491.991 | 2588.727  | 1699.157            | 3870.259  | 3096.395            |
| 3321.62  | 2914.292          | 2360.499 | 1248.485  | 2512.119            | 1928.057  | 2780.339            |
| 4289.153 | 1630.847          | 1778.119 | 4957.682  | 1888.128            | 6535.653  | 3644.541            |
| 2904.012 | 2492.425          | 1338.233 | 2666.555  | 2495.247            | 3348.950  | 3752.15             |
| 3662.63  | 3952.583          | 2017.130 | 2717.818  | 1781.652            | 2931.343  | 3774.743            |
| 2613.299 | 2826.274          | 1773.324 | 2350.575  | 1844.259            | 4231.136  | 7107.145            |
| 2427.935 | 3451.583          | 1292.638 | 1705.690  | 1919.575            | 2726.132  | 5194.772            |
| 3045.394 | 3890.014          | 1370.735 | 1437.455  | 2092.248            | 3143.966  | 3641.905            |
| 3007.652 | 3559.553          | 1760.536 | 1935.729  | 3549.591            | 3942.419  | 3448.522            |
| 3646.285 | 3471.837          | 2127.349 | 1799.386  | 2289.834            | 2622.884  | 2494.105            |
| 4750.743 | 3495.855          | 2191.289 | 2133.459  | 3852.218            | 2145.143  | 5033.092            |
| 4469.762 | 1985.703          | 1986.303 | 2717.603  | 1433.363            | 2757.523  | 3917.145            |
| 3265.527 | 1737.763          | 1911.326 | 1586.507  | 1341.606            | 1527.923  | 2486.047            |
| 4125.26  | 2211.507          | 2218.92  | 2742.014  | 1298.958            | 3624.052  | 2154.780            |
| 4094.502 | 3160.350          | 1997.949 | 2467.174  | 1750.779            | 1299.491  | 1960.191            |
| 2553.566 | 2027.490          | 1401.944 | 1731.107  | 1718.111            | 3493.876  | 1835.110            |
| 1817.979 | 2394.770          | 1751.021 | 2294.214  | 2074.801            | 2620.312  | 3376.982            |

|          |          |          |          |          |          |          |
|----------|----------|----------|----------|----------|----------|----------|
| 1545.914 | 3895.285 | 1520.079 | 2448.937 | 1508.104 | 1739.033 | 2845.328 |
| 2791.753 | 3125.941 | 1388.623 | 3447.495 | 1433.363 | 3478.521 | 2179.630 |
| 3570.059 | 5063.984 | 1770.279 | 2216.385 | 1712.152 | 2540.663 | 2312.393 |
| 4874.889 | 5069.329 | 1547.634 | 1988.141 | 1679.054 | 2822.346 | 2267.361 |
| 2374.666 | 3643.430 | 2318.102 | 3277.623 | 3716.377 | 2289.237 | 2784.180 |
| 5251.932 | 3641.397 | 2343.829 | 8076.633 | 3257.017 | 6010.108 | 5434.543 |
| 3882.318 | 5050.355 | 1810.394 | 2150.547 | 3231.744 | 2190.300 | 7868.479 |
| 2957.355 | 5289.26  | 2233.839 | 2016.860 | 1750.995 | 1785.476 | 5016.299 |
| 2539.302 | 3134.599 | 3274.907 | 2470.907 | 1078.756 | 1915.173 | 1529.371 |
| 2410.03  | 2890.273 | 1927.311 | 2015.999 | 1665.771 | 2209.137 | 5215.255 |
| 3088.187 | 2122.360 | 1677.643 | 2397.458 | 1344.621 | 2380.604 | 2502.012 |
| 3136.924 | 3671.062 | 3390.530 | 3068.764 | 1680.418 | 1775.282 | 1901.228 |
| 4457.058 | 4765.296 | 3031.557 | 1628.149 | 3561.151 | 1929.804 | 2758.651 |
| 2898.96  | 3875.257 | 3513.689 | 1939.319 | 2672.155 | 3231.817 | 2195.294 |
| 3484.25  | 2648.206 | 2426.798 | 2868.162 | 2755.512 | 1502.742 | 1901.303 |
| 2359.435 | 2429.706 | 1973.210 | 1688.962 | 2371.683 | 2696.889 | 2298.989 |
| 3248.663 | 2477.216 | 1922.972 | 1931.063 | 3158.511 | 3607.671 | 1673.053 |
| 3744.577 | 3491.564 | 1240.573 | 2640.923 | 2867.229 | 1204.000 | 2097.699 |
| 1843.016 | 3534.104 | 1482.933 | 1794.647 | 1832.700 | 1135.323 | 2331.822 |
| 2547.994 | 4390.638 | 2776.104 | 1576.599 | 2384.750 | 2541.546 | 2424.598 |
| 3530.906 | 5264.941 | 1608.071 | 2320.994 | 1397.392 | 2014.455 | 1451.581 |
| 2629.792 | 2554.241 | 1975.418 | 2747.255 | 2184.005 | 2516.091 | 1510.168 |
| 3101.263 | 2322.865 | 2363.087 | 2188.743 | 3020.804 | 1922.862 | 3026.135 |
| 3309.435 | 3014.356 | 2131.231 | 1412.183 | 2596.050 | 1979.296 | 2162.536 |
| 2944.279 | 2282.734 | 2638.512 | 1871.901 | 2996.321 | 3008.395 | 2203.352 |
| 3637.147 | 3222.015 | 4765.7   | 1836.936 | 1347.780 | 2948.154 | 1854.840 |
| 4294.874 | 3197.469 | 5092.877 | 2083.776 | 1731.753 | 2670.463 | 1846.857 |
| 3983.581 | 4292.907 | 2852.123 | 3183.137 | 2078.032 | 2342.459 | 3182.920 |
| 4818.722 | 2638.343 | 5209.820 | 1789.119 | 2040.841 | 2099.330 | 3167.558 |
| 1944.056 | 6278.912 | 3708.618 | 1848.783 | 1451.169 | 2037.148 | 2166.979 |
| 1605.126 | 4144.731 | 2397.639 | 2550.746 | 2078.391 | 1918.233 | 2117.579 |
| 1803.789 | 2688.78  | 3023.2   | 1575.306 | 1581.337 | 1816.563 | 1656.486 |
| 3723.328 | 2811.517 | 2488.156 | 2566.111 | 1504.873 | 1847.392 | 1986.096 |
| 3723.328 | 3031.071 | 1639.255 | 2379.150 | 1787.539 | 2530.274 | 1599.405 |
| 3450.149 | 6251.204 | 2554.829 | 2574.726 | 1172.451 | 4081.985 | 2208.623 |
| 4815.973 | 5274.051 | 1752.249 | 2844.756 | 2133.316 | 1847.467 | 1764.925 |
| 2412.11  | 5275.030 | 1958.039 | 1906.508 | 2683.715 | 2215.482 | 2672.728 |
| 4959.435 | 3515.281 | 1875.192 | 1891.861 | 2121.397 | 2340.444 | 1637.660 |
| 3130.163 | 2095.555 | 2876.271 | 1880.374 | 1797.591 | 2745.708 | 1806.192 |
| 3626.226 | 3306.719 | 1896.378 | 2108.546 | 4503.348 | 2813.564 | 1588.636 |
| 3790.49  | 2439.41  | 1837.375 | 3151.618 | 2626.492 | 2918.519 | 1959.212 |
| 2685.587 | 3185.874 | 2609.428 | 1671.156 | 2149.686 | 2570.435 | 1845.652 |
| 3452.897 | 3095.221 | 3332.57  | 1950.017 | 1808.720 | 3018.697 | 2360.287 |
| 3568.945 | 2067.923 | 2489.979 | 2344.113 | 2015.568 | 1709.248 | 2507.961 |
| 4833.432 | 3733.330 | 3198.473 | 2300.460 | 3188.738 | 1990.172 | 2191.077 |
| 2622.66  | 5676.49  | 3382.848 | 1464.810 | 2459.420 | 2989.817 | 2428.213 |
| 2777.117 | 3393.532 | 2537.136 | 1256.454 | 2570.634 | 1968.750 | 2948.194 |
| 3706.166 | 3142.882 | 2504.787 | 1580.476 | 2293.783 | 2509.480 | 2954.294 |
| 3361.144 | 3352.573 | 3641.490 | 1330.477 | 1284.814 | 2164.212 | 2530.326 |
| 4382.095 | 2832.223 | 3219.052 | 1790.627 | 5283.714 | 1399.988 | 2013.206 |
| 3307.801 | 5049.753 | 2001.855 | 1891.143 | 7957.449 | 2288.493 | 2021.113 |
| 5960.327 | 2419.61  | 1838.970 | 1607.831 | 1891.574 | 1414.877 | 2064.338 |
| 4202.452 | 2256.833 | 2896.243 | 1951.812 | 2309.866 | 2461.393 | 2373.692 |
| 2982.764 | 3910.569 | 3301.747 | 1805.991 | 1937.093 | 2506.214 | 2297.634 |

|          |          |          |          |          |          |          |
|----------|----------|----------|----------|----------|----------|----------|
| 3454.235 | 2040.215 | 2335.978 | 1843.039 | 2784.805 | 1454.760 | 4355.797 |
| 3107.281 | 3630.404 | 1938.22  | 2455.184 | 2666.196 | 1319.615 | 2451.783 |
| 4175.78  | 2788.025 | 2444.948 | 1618.887 | 1642.796 | 2138.839 | 1550.456 |
| 2844.428 | 2731.782 | 2068.201 | 2074.442 | 2237.996 | 2459.114 | 2502.991 |
| 2397.548 | 3129.555 | 2481.069 | 3313.952 | 2581.619 | 1832.010 | 1640.973 |
| 2895.914 | 2676.517 | 1801.747 | 1750.995 | 2507.811 | 2928.740 | 2079.023 |
| 2022.363 | 4246.226 | 2893.860 | 1377.863 | 1766.216 | 2522.243 | 2832.676 |
| 4024.22  | 3299.416 | 2015.608 | 1905.790 | 1606.969 | 2122.430 | 947.865  |
| 4663.596 | 3477.032 | 1276.114 | 2337.579 | 2458.702 | 2057.85  | 2814.151 |
| 4660.03  | 2445.743 | 1953.920 | 2127.644 | 1995.824 | 3431.33  | 2515.190 |
| 3383.21  | 2449.132 | 2441.358 | 1387.843 | 1644.304 | 2089.157 | 4579.228 |
| 4386.627 | 5297.166 | 2205.672 | 2177.256 | 2254.438 | 2756.448 | 3932.206 |
| 4476.969 | 3583.572 | 2003.861 | 1359.771 | 1480.247 | 2422.650 | 3224.639 |
| 2440.639 | 3635.675 | 7702.462 | 1638.344 | 1458.420 | 1283.075 | 2752.025 |
| 4774.591 | 2751.734 | 2910.533 | 1468.831 | 2438.239 | 1940.946 | 1915.084 |
| 2702.675 | 2436.031 | 4777.833 | 1726.009 | 1993.311 | 2000.808 | 2310.511 |
| 3284.101 | 2906.687 | 3657.908 | 1639.493 | 2147.388 | 2418.700 | 2670.017 |
| 3474.74  | 2554.994 | 5003.971 | 2908.153 | 1793.642 | 2650.626 | 1482.531 |
| 6383.061 | 4880.193 | 2376.033 | 2353.447 | 2047.949 | 2306.345 | 2065.468 |
| 6769.539 | 3872.019 | 2784.354 | 1590.527 | 1481.539 | 1552.225 | 2283.401 |
| 3623.254 | 2492.274 | 3436.242 | 2124.269 | 2700.946 | 2407.761 | 2274.440 |
| 3483.73  | 1686.639 | 2885.752 | 1834.567 | 1661.176 | 1991.996 | 1739.849 |
| 6467.311 | 7169.855 | 2481.524 | 1973.423 | 1324.159 | 3287.302 | 1670.794 |
| 3068.351 |          | 2988.362 | 2058.359 | 2067.190 | 3409.380 | 1597.974 |
| 4258.841 |          | 3250.345 | 1850.434 | 2576.449 | 2836.972 | 1558.439 |
| 6090.267 |          | 3158.799 | 1661.750 | 1683.505 | 1609.808 | 3343.471 |
|          |          | 2604.141 | 1930.704 | 1689.680 | 2094.019 | 2841.487 |
|          |          | 2476.825 | 1822.648 | 1882.025 | 4036.865 | 1959.288 |
|          |          | 2413.015 | 2317.620 | 2520.878 | 1889.365 | 2177.145 |
|          |          | 1730.511 | 1870.107 | 2105.530 | 1529.435 | 2511.801 |
|          |          | 2363.605 | 1616.518 | 1857.757 | 1829.275 | 1728.026 |
|          |          | 3170.924 | 1550.321 | 1873.696 | 1766.071 | 3216.883 |
|          |          | 3384.860 | 3101.647 | 1949.084 | 1928.488 | 2001.383 |
|          |          | 2886.661 | 2037.610 | 1888.989 | 1548.655 | 1787.818 |
|          |          | 2805.042 | 2215.811 | 1964.089 | 2827.856 | 1641.500 |
|          |          | 2601.185 | 2533.012 | 2270.880 | 1405.230 | 1697.603 |
|          |          | 2165.204 | 2906.358 | 1815.828 | 2806.889 | 2343.268 |
|          |          | 2376.943 | 2112.710 | 1472.492 | 1964.040 | 2047.244 |
|          |          | 3598.873 | 1941.832 | 1720.337 | 2911.267 | 2396.509 |
|          |          | 4001.585 | 1457.343 | 1295.871 | 3221.971 | 2037.756 |
|          |          | 3453.293 | 1865.870 | 1958.058 | 2401.684 | 2241.456 |
|          |          | 2644.912 | 1829.326 | 1838.587 | 2133.825 | 2054.925 |
|          |          | 3627.215 | 2737.850 | 1649.688 | 1521.762 | 4520.490 |
|          |          | 1684.738 | 3053.184 | 3135.033 | 1675.063 | 2119.085 |
|          |          | 3509.903 | 2099.068 | 3561.797 | 2934.639 | 1500.604 |
|          |          | 1975.822 | 1615.872 | 2327.959 | 3147.602 | 2517.299 |
|          |          | 1927.245 | 1810.730 | 1883.245 | 2116.374 | 2662.562 |
|          |          | 2748.281 | 1911.175 | 2043.497 | 2490.791 | 1512.879 |
|          |          | 3618.046 | 1571.50  | 2666.771 | 2507.810 | 1496.011 |
|          |          | 3495.428 | 1838.157 | 2467.748 | 2189.947 | 1990.313 |
|          |          | 3571.818 | 1551.182 | 2118.238 | 1870.428 | 3265.304 |
|          |          | 1500.206 | 1551.541 | 2153.060 | 3566.675 | 3737.015 |
|          |          | 4521.989 | 1547.521 | 1395.310 | 2135.426 | 2632.214 |
|          |          | 2941.907 | 1879.727 | 1708.778 | 2494.481 | 2504.045 |



|          |          |          |          |          |
|----------|----------|----------|----------|----------|
| 2462.275 | 1512.699 | 2059.652 | 1712.136 | 2494.406 |
| 3153.646 | 1417.783 | 1827.674 | 2361.417 | 2651.794 |
| 2607.930 | 1140.573 | 2014.635 | 2126.616 | 2756.543 |
| 2298.28  | 2040.195 | 2198.292 | 3033.063 | 2079.701 |
| 2686.821 | 2195.492 | 2012.912 | 2821.155 | 2955.951 |
| 2685.532 | 2375.991 | 1969.115 | 2646.221 | 1625.234 |
| 3753.925 | 2031.363 | 2088.945 | 2932.682 | 4008.792 |
| 2787.309 | 2092.75  | 1520.166 | 2638.841 | 2502.463 |
| 2707.055 | 2139.562 | 2475.143 | 2351.401 | 2785.234 |
| 3696.557 | 2236.058 | 2117.377 | 2834.785 | 1641.425 |
| 2328.517 | 1629.657 | 2224.427 | 2279.334 | 2147.400 |
| 2436.129 | 1677.330 | 1810.443 | 3080.581 | 2260.056 |
| 2425.596 | 1563.173 | 1971.987 | 3729.259 | 1996.790 |
| 3049.595 | 2769.513 | 1436.307 | 2733.725 | 2079.776 |
| 3387.892 | 1541.203 | 1561.449 | 2441.240 | 3233.601 |
| 4078.278 | 2171.799 | 1712.296 | 2487.779 | 2589.290 |
|          | 1769.231 | 5862.329 | 3163.341 | 3072.674 |
|          | 3219.89  | 1347.565 | 3459.215 | 2930.121 |
|          | 2330.615 |          | 1687.662 | 4347.062 |
|          |          |          | 1708.974 | 5687.794 |
|          |          |          | 1822.910 |          |
|          |          |          | 1925.626 |          |
|          |          |          | 3117.330 |          |
|          |          |          | 1811.238 |          |
|          |          |          | 1943.398 |          |
|          |          |          | 2084.445 |          |
|          |          |          | 2215.626 |          |
|          |          |          | 1807.548 |          |
|          |          |          | 2371.583 |          |
|          |          |          | 2864.229 |          |
|          |          |          | 2917.093 |          |
|          |          |          | 1772.531 |          |
|          |          |          | 3144.967 |          |
|          |          |          | 3101.666 |          |
|          |          |          | 2178.350 |          |
|          |          |          | 2480.248 |          |
|          |          |          | 2777.930 |          |
|          |          |          | 1723.056 |          |
|          |          |          | 2443.650 |          |
|          |          |          | 1800.620 |          |
|          |          |          | 2443.424 |          |
|          |          |          | 1988.431 |          |
|          |          |          | 2668.511 |          |
|          |          |          | 2483.788 |          |
|          |          |          | 2026.159 |          |
|          |          |          | 1947.088 |          |
|          |          |          | 3391.892 |          |
|          |          |          | 2589.215 |          |
|          |          |          | 2984.567 |          |
|          |          |          | 3116.049 |          |
|          |          |          | 3394.980 |          |
|          |          |          | 3103.323 |          |
|          |          |          | 2447.039 |          |
|          |          |          | 1844.222 |          |

2906.701  
 2636.582  
 2027.213  
 1725.390

**Inducible lines**

| EV-i-1   | WEE1-i-1 (-)dex | WEE1-i-1 (+)dex | WEE1-i-6 (-)dex | WEE1-i-6 (+)dex |
|----------|-----------------|-----------------|-----------------|-----------------|
| 1615.854 | 2759.679        | 2495.094        | 2538.462        | 3238.2875       |
| 2594.865 | 1654.265        | 2796.774        | 5458.596        | 1986.730        |
| 1509.486 | 2753.481        | 2096.338        | 1857.061        | 2757.028        |
| 1545.158 | 3611.854        | 1848.997        | 6189.480        | 1482.204        |
| 1999.46  | 2976.884        | 2270.744        | 3308.180        | 1546.677        |
| 2004.72  | 2733.951        | 1977.928        | 2688.798        | 1380.053        |
| 2623.547 | 4897.288        | 2014.756        | 2569.688        | 1574.673        |
| 2558.977 | 2907.125        | 2077.095        | 2195.262        | 2364.261        |
| 2940.42  | 1980.650        | 1954.002        | 1796.927        | 1639.896        |
| 2478.768 | 2218.827        | 4776.073        | 3324.772        | 1701.892        |
| 1581.263 | 1912.116        | 1616.287        | 4294.165        | 1032.694        |
| 2788.506 | 3215.782        | 2861.276        | 3540.385        | 2402.690        |
| 2910.729 | 3577.695        | 2631.160        | 3592.478        | 1388.234        |
| 1925.808 | 3579.569        | 1900.238        | 3487.206        | 1359.637        |
| 2397.622 | 4979.803        | 1610.377        | 3107.636        | 1280.378        |
| 2035.348 | 1875.506        | 1703.202        | 3122.561        | 1638.020        |
| 1971.642 | 2912.890        | 1476.474        | 3061.123        | 1707.822        |
| 2035.853 | 2721.556        | 1444.043        | 2103.394        | 1127.264        |
| 2499.523 | 2163.552        | 1961.497        | 2103.032        | 1497.515        |
| 1717.539 | 2351.355        | 1497.734        | 2643.516        | 1894.336        |
| 1501.342 | 2595.369        | 2759.803        | 3528.721        | 1808.172        |
| 1776.272 | 2539.663        | 2188.730        | 2754.366        | 1565.366        |
| 2392.793 | 3185.226        | 2600.171        | 2507.236        | 2101.640        |
| 2680.623 | 2807.386        | 2427.710        | 3236.961        | 2333.113        |
| 1879.326 | 3727.880        | 3455.628        | 3287.025        | 1541.423        |
| 1850.716 | 3614.953        | 2092.518        | 2484.413        | 3654.697        |
| 2044.717 | 4244.374        | 1781.181        | 2705.171        | 4279.988        |
| 2490.298 | 3147.896        | 1765.470        | 2133.968        | 2198.087        |
| 2204.198 | 3607.098        | 2172.298        | 4186.285        | 3339.838        |
| 2030.015 | 2156.418        | 3225.079        | 4416.100        | 2231.562        |
| 2893.505 | 2004.215        | 1824.638        | 2628.229        | 1319.633        |
| 2441.51  | 2289.307        | 2382.379        | 3477.788        | 2641.292        |
| 1703.198 | 2833.114        | 2392.685        | 3279.127        | 2295.435        |
| 2639.69  | 2714.350        | 2821.206        | 2673.221        | 1496.239        |
| 2388.83  | 4519.520        | 6119.075        | 2384.649        | 2384.301        |
| 2769.336 | 2642.644        | 3345.939        | 2482.747        | 1742.573        |
| 1863.256 | 3289.433        | 2357.155        | 2826.020        | 1592.686        |
| 2180.056 | 3680.965        | 2133.021        | 3299.269        | 1945.299        |
| 2480.281 | 3283.812        | 1375.361        | 3502.131        | 3047.795        |
| 2382.993 | 4152.490        | 2309.877        | 2609.898        | 2308.269        |
| 1943.537 | 3488.262        | 2913.526        | 1879.376        | 1707.747        |
| 2170.687 | 2419.169        | 3399.990        | 1817.938        | 1737.319        |
| 2293.631 | 2406.702        | 3181.838        | 2411.166        | 2373.493        |
| 2246.14  | 2586.577        | 1892.022        | 2597.002        | 1740.021        |
| 2495.775 | 2292.261        | 2376.325        | 4463.555        | 2251.152        |
| 2240.807 | 2407.062        | 2267.645        | 3313.831        | 2453.953        |

|          |          |          |          |          |
|----------|----------|----------|----------|----------|
| 2316.548 | 2471.056 | 3217.801 | 2745.961 | 2080.925 |
| 1802     | 3619.998 | 2946.245 | 3529.083 | 2361.484 |
| 1687.343 | 3552.400 | 2908.697 | 3446.851 | 2134.139 |
| 1627.457 | 2641.780 | 3034.746 | 2860.506 | 1643.574 |
| 1599.856 | 2758.021 | 1655.925 | 3484.598 | 2407.118 |
| 1842.573 | 2549.247 | 1974.109 | 2775.956 | 2817.824 |
| 1854.608 | 3847.365 | 2286.239 | 4568.536 | 2336.865 |
| 2302.207 | 3327.051 | 2972.983 | 2141.576 | 2495.008 |
| 2213.278 | 3449.418 | 2748.705 | 2564.254 | 2649.398 |
| 2205.567 | 3459.796 | 2194.063 | 3806.932 | 2601.738 |
| 2035.708 | 3989.910 | 2406.378 | 1764.034 | 2364.936 |
| 1855.761 | 2218.106 | 2692.707 | 3690.649 | 2193.208 |
| 2133.646 | 2034.195 | 2767.226 | 2681.190 | 2821.952 |
| 2927.952 | 2071.885 | 2961.740 | 2346.105 | 2698.185 |
| 2432.502 | 2866.408 | 1588.252 | 4076.015 | 2193.959 |
| 2195.982 | 2423.853 | 2099.220 | 2897.601 | 2625.230 |
| 2718.314 | 1700.963 | 1526.994 | 3575.741 | 3478.466 |
| 2808.107 | 2216.160 | 2200.261 | 3278.331 | 3430.655 |
| 1958.598 | 2890.982 | 1483.752 | 3978.569 | 1984.928 |
| 2077.507 | 2596.378 | 1620.107 | 2835.655 | 1974.946 |
| 1739.159 | 2549.103 | 1718.625 | 2096.222 | 2175.645 |
| 2398.703 | 2546.437 | 1947.876 | 1558.781 | 1959.034 |
| 2386.524 | 2004.215 | 3710.031 | 1850.541 | 2420.478 |
| 2533.105 | 2729.555 | 1709.400 | 2249.165 | 1989.807 |
| 2168.525 | 3014.142 | 1813.828 | 2187.220 | 2431.361 |
| 2472.426 | 3828.267 | 2224.476 | 2010.802 | 2673.791 |
| 2475.309 | 2890.118 | 2920.589 | 2056.808 | 3745.440 |
| 4070.264 | 3361.715 | 2613.143 | 2130.926 | 2433.763 |
| 2191.875 | 3613.007 | 2690.257 | 2729.587 | 2879.144 |
| 2179.263 | 3938.600 | 2813.278 | 2389.503 | 3834.756 |
| 2766.67  | 2895.955 | 2769.893 | 1850.468 | 3491.075 |
| 1650.734 | 2264.084 | 3563.875 | 1640.071 | 1824.984 |
| 2328.511 | 2936.312 | 1913.715 | 1907.632 | 1674.797 |
| 2187.046 | 2672.047 | 2967.073 | 1852.134 | 1896.212 |
| 2277.272 | 2733.159 | 2636.710 | 2568.022 | 1620.982 |
| 2432.79  | 2870.588 | 1458.528 | 2639.386 | 2785.174 |
| 2514.44  | 3357.391 | 1446.925 | 2183.308 | 2322.155 |
| 3754.401 | 2326.132 | 1793.144 | 1747.008 | 2004.368 |
| 2640.771 | 2374.416 | 1165.929 | 1891.330 | 1878.499 |
| 2566.688 | 2354.526 | 2875.618 | 1622.538 | 2017.427 |
| 1967.102 | 1655.058 | 3372.676 | 1870.247 | 2693.231 |
| 3134.492 | 1763.660 | 2877.275 | 1666.515 | 2549.874 |
| 2778.849 | 1701.684 | 2564.713 | 1891.620 | 2430.986 |
| 3978.669 | 1529.375 | 2669.933 | 1902.778 | 2915.096 |
| 2007.819 | 2104.386 | 2498.482 | 3017.870 | 3339.237 |
| 2630.394 | 2476.173 | 2973.199 | 2692.348 | 4921.341 |
| 2707.432 | 3035.834 | 2323.859 | 2495.933 | 3874.761 |
| 3135.069 | 2882.623 | 2370.776 | 2453.549 | 2751.850 |
| 2197.063 | 2179.911 | 2285.374 | 2170.049 | 1784.829 |
| 3629.295 | 617.313  | 2084.806 | 2148.821 | 1572.721 |
| 3415.26  | 2449.292 | 2562.623 | 2044.492 | 2165.438 |
| 2829.006 | 1998.882 | 3541.750 | 1952.696 | 1660.462 |
| 3101.054 | 2456.427 | 3515.084 | 2339.874 | 1778.750 |
| 2575.191 | 1986.920 | 3185.225 | 1527.844 | 1631.940 |

|          |          |          |          |          |
|----------|----------|----------|----------|----------|
| 3545.915 | 2657.850 | 3105.157 | 2374.143 | 2092.483 |
| 3285.902 | 3300.531 | 4189.432 | 1876.623 | 1755.557 |
| 2357.625 | 1553.878 | 3228.178 | 1713.029 | 1769.893 |
| 2666.787 | 1996.432 | 2689.320 | 3258.334 | 2823.978 |
| 2709.594 | 1495.432 | 1826.151 | 3224.644 | 2763.934 |
| 2260.769 | 2628.303 | 2504.103 | 2830.367 | 2563.309 |
| 1946.708 | 2449.581 | 2149.380 | 3495.828 | 2172.268 |
| 3449.347 | 3548.869 | 2482.843 | 2871.881 |          |
| 3832.448 | 3176.722 | 2694.220 | 2962.445 |          |
| 3815.296 | 1962.057 | 2286.815 | 1859.162 |          |
| 4041.726 | 2271.507 | 2404.071 | 2008.339 |          |
| 2840.393 | 2639.618 | 2013.747 | 1912.848 |          |
| 3622.953 | 2305.954 | 2568.244 | 2141.431 |          |
| 1998.018 | 2129.465 | 2601.684 | 4550.206 |          |
| 3124.908 | 2383.785 | 2163.650 | 4799.872 |          |
| 2632.051 | 3709.359 | 2170.208 | 2540.563 |          |
| 2793.046 | 4407.963 | 1984.198 | 2387.547 |          |
| 2731.574 | 2918.367 | 2341.372 | 2135.345 |          |
| 3154.815 | 1816.845 | 1948.596 | 2944.332 |          |
| 2952.311 | 1899.288 | 2533.507 | 2221.851 |          |
| 2900.64  | 2701.234 | 2114.211 | 2403.703 |          |
| 3467.075 | 2115.413 | 1220.557 | 2552.083 |          |
| 3754.112 | 2249.815 | 1623.998 | 2517.958 |          |
| 3367.552 | 2142.149 | 1620.323 | 2832.613 |          |
| 3423.259 | 3231.13  | 1786.586 | 2945.419 |          |
| 3350.185 | 2601.35  | 1822.332 | 3851.924 |          |
| 4657.31  | 3349.824 | 1297.238 | 5233.998 |          |
| 3516.224 | 2720.187 | 1925.462 | 2649.602 |          |
|          | 3099.036 | 1516.544 | 3693.474 |          |
|          | 2307.756 | 1520.940 | 2994.97  |          |
|          | 2740.509 | 2084.879 | 2405.152 |          |
|          | 1998.450 | 2666.546 | 4221.207 |          |
|          | 2187.838 | 3281.509 | 3437.867 |          |
|          | 2844.140 | 1552.506 | 3261.884 |          |
|          | 2330.024 | 2290.203 | 2567.732 |          |
|          | 2646.824 | 3196.180 | 2586.787 |          |
|          | 2199.081 | 2532.930 | 2447.464 |          |
|          | 2633.060 | 2675.555 | 5147.564 |          |
|          |          | 2350.668 | 4045.441 |          |
|          |          | 2030.322 | 4269.459 |          |
|          |          | 2583.811 | 2926.944 |          |
|          |          | 2901.707 | 2980.050 |          |
|          |          | 2536.390 | 3836.637 |          |
|          |          |          | 3099.884 |          |
|          |          |          | 2639.821 |          |
|          |          |          | 3177.914 |          |
|          |          |          | 3097.638 |          |

**II.IV. Mitotic cell area ( $\mu\text{m}^2$ ) following synchronization of EV-c-10 subcultured for 4 weeks without hygromycin**

|          |          |          |          |          |          |          |          |
|----------|----------|----------|----------|----------|----------|----------|----------|
| 2616.374 | 2668.523 | 2308.57  | 3211.868 | 3115.946 | 2315.1   | 2949.773 | 3392.077 |
| 3434.634 | 2774.292 | 3973.836 | 1807.797 | 2226.609 | 3069.862 | 2630.71  | 2227.509 |
| 2875.843 | 2585.076 | 2973.866 | 1832.565 | 2292.508 | 1739.346 | 3053.875 | 2879.445 |
| 2388.955 | 4085.519 | 3145.068 | 2206.269 | 2221.355 | 1586.832 | 2782.698 | 5287.614 |
| 3044.718 | 3493.853 | 3737.785 | 1809.748 | 2079.875 | 1681.853 | 2835.312 | 5637.075 |
| 2553.627 | 2073.044 | 2457.106 | 2179.474 | 2528.334 | 1969.092 | 2113.95  | 3271.762 |
| 1071.798 | 3398.307 | 2219.178 | 2967.411 | 2174.22  | 1930.513 | 2766.486 | 1792.26  |
| 2684.825 | 1685.08  | 5710.404 | 3950.193 | 3535.809 | 2808.067 | 2712.296 | 2983.698 |
| 2730.759 | 1635.468 | 2250.101 | 4796.899 | 2943.018 | 3327.154 | 2656.004 | 2163.712 |
| 3243.016 | 2759.431 | 1501.794 | 5994.641 | 3142.516 | 2569.239 | 3041.416 | 2573.968 |
| 2411.622 | 2514.223 | 2240.119 | 3058.303 | 2377.471 | 2543.57  | 3894.877 | 2399.763 |
| 2040.695 | 3234.76  | 4170.482 | 2926.655 | 2140.144 | 2795.308 | 3195.055 | 2097.663 |
| 4144.588 | 3253.899 | 3722.173 | 3593.602 | 2736.989 | 2663.284 | 2018.028 | 3340.138 |
| 2926.355 | 1880.226 | 2753.952 | 2437.441 | 1912.95  | 1793.836 | 1841.647 | 2212.123 |
| 2096.087 | 1555.234 | 3813.291 | 3419.247 | 3409.64  | 2107.795 | 2649.849 | 2742.618 |
| 2247.775 | 2100.59  | 3797.079 | 2467.088 | 2665.536 | 2537.565 | 2164.538 |          |
| 2943.093 | 2083.027 | 2064.938 | 2189.081 | 1650.93  | 2678.971 | 2519.327 |          |
| 2116.502 | 1603.57  | 3909.137 | 1867.316 | 2889.503 | 4659.246 | 2872.465 |          |
| 2341.069 | 1568.519 | 2672.891 | 1608.749 | 2556.254 | 2119.954 | 2315.1   |          |
| 1824.759 | 1921.957 | 3527.178 | 2531.636 | 2502.064 | 2086.78  | 2080.325 |          |

**II.V. Mitotic cell area ( $\mu\text{m}^2$ ) of EV-c-10 subcultured for 3 weeks without hygromycin**

| 1 week   | 2 week   | 3 week   |
|----------|----------|----------|
| 2196.059 | 1719.747 | 2226.083 |
| 2490.276 | 1383.615 | 2182.326 |
| 2404.972 | 2066.907 | 2325.608 |
| 2920.821 | 3696.469 | 1361.664 |
| 2250.892 | 2329.96  | 3057.928 |
| 1229.328 | 1775.494 | 1228.44  |
| 4449.395 | 1589.721 | 3290.526 |
| 2463.183 | 2259.208 | 2178.123 |
| 2393.761 | 2627.676 | 2066.515 |
| 3987.23  | 1820.736 | 1501.869 |
| 2505.296 | 1887.737 | 1455.935 |
| 2436.52  | 2323.881 | 1913.551 |
| 1855.993 | 1895.988 | 1918.354 |
| 3013.671 | 1794.287 | 1747.002 |
| 3247.664 | 3214.12  | 1806.296 |
| 3224.02  | 2182.701 | 1962.187 |
| 2405.341 | 2238.993 | 1434.844 |
| 1498.032 | 1598.541 | 3067.685 |
| 2635.765 | 1990.858 | 1579.552 |
| 3467.267 | 1861.687 | 1111.728 |
| 2456.256 | 1933.29  | 2331.237 |
| 2689.184 | 2499.587 | 3341.79  |
| 2234.987 | 2622.979 | 1169.897 |
| 2455.662 | 3011.619 | 1383.205 |
| 2231.508 | 1905.144 | 2468.814 |

|          |          |          |
|----------|----------|----------|
| 3462.302 | 1970.443 | 2126.86  |
| 2369.98  | 4027.726 | 1397.992 |
| 1544.926 | 3840.386 | 1485.432 |
| 2345.432 | 1667.292 | 1787.081 |
| 2254.99  | 1386.433 | 1279.178 |
| 1845.098 | 1415.329 | 1643.199 |
| 1738.478 | 1029.467 | 1313.103 |
| 2298.09  | 1052.209 | 1777.474 |
|          | 1287.509 | 1715.553 |
|          | 2324.857 | 2267.214 |
|          | 2114.55  | 2137.893 |
|          | 1203.296 | 1486.783 |
|          | 1405.197 | 1733.266 |
|          | 1796.238 | 2807.392 |
|          | 2031.914 | 1649.279 |
|          | 2008.947 | 2676.269 |
|          | 1290.962 | 2394.809 |
|          | 1402.87  | 1691.235 |
|          | 2425.132 | 1453.983 |
|          | 1693.637 | 2202.216 |
|          | 2329.51  | 2351.652 |
|          | 1816.203 | 1895.912 |
|          | 1643.95  | 851.4343 |
|          | 1600.417 | 1676.299 |
|          | 2243.196 |          |
|          | 1801.192 |          |
|          | 2105.168 |          |
|          | 1948.902 |          |
|          | 1683.729 |          |
|          | 2070.192 |          |
|          | 2692.406 |          |
|          | 2552.201 |          |
|          | 1998.364 |          |
|          | 2243.571 |          |
|          | 1668.718 |          |
|          | 2152.379 |          |
|          | 1934.641 |          |
|          | 2350.901 |          |
|          | 2405.017 |          |
|          | 1726.061 |          |
|          | 2291.532 |          |
|          | 1940.946 |          |
|          | 1906.42  |          |
|          | 1657.235 |          |
|          | 3352.523 |          |
|          | 3095.306 |          |
|          | 1949.502 |          |
|          | 3357.551 |          |
|          | 896.693  |          |
|          | 1290.737 |          |
|          | 1866.566 |          |
|          | 1330.216 |          |
|          | 1691.685 |          |
|          | 2413.723 |          |

**II.VI. Mitotic cell area ( $\mu\text{m}^2$ ) of WEE1-i-1, WEE1-1-6, EV-i-1 and EV-i-3 ( $\pm$ ) dexametasone at 0, 12, 24, 36 and 48 hour**

**WEE1-i-1 (+) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1390.031 | 1307.101 | 1909.789 | 1395.294 | 1281.538 |
| 721.2551 | 1776.485 | 1707.539 | 1870.617 | 955.7589 |
| 813.0567 | 1160.414 | 1189.962 | 2262.183 | 1247.329 |
| 1652.353 | 1293.868 | 1224.322 | 1490.704 | 2441.576 |
| 1096.506 | 1634.309 | 1363.791 | 1719.795 | 1859.64  |
| 623.3635 | 1282.29  | 2223.688 | 1855.805 | 1575.363 |
| 998.3891 | 1989.034 | 1892.12  | 1482.283 | 1592.731 |
| 1575.739 | 1138.911 | 1084.777 | 2071.513 | 1561.529 |
| 1066.582 | 1543.71  | 1220.111 | 1249.058 | 1482.509 |
| 922.6773 | 1810.393 | 1436.947 | 1567.995 | 1571.604 |
| 2421.201 | 1122.459 | 1592.957 | 1383.414 | 1698.216 |
| 2196.17  | 1298.372 | 1422.135 | 1552.732 | 962.4504 |
| 1268.381 |          | 1571.153 | 3162.38  | 1348.303 |
| 864.2581 |          | 975.4576 | 2642.848 | 2456.012 |
| 890.1971 |          | 1127.408 | 2099.858 | 2054.446 |
| 720.8792 |          |          | 1790.92  | 2258.499 |
| 944.1052 |          |          | 1832.056 | 1697.84  |
| 961.3979 |          |          | 1645.906 | 1999.56  |
| 1392.512 |          |          |          | 1559.574 |
| 1729.644 |          |          |          | 1161.692 |
| 1923.848 |          |          |          | 1850.693 |
| 1731.524 |          |          |          | 1692.202 |
| 1104.702 |          |          |          |          |
| 657.573  |          |          |          |          |
| 1008.614 |          |          |          |          |
| 1549.424 |          |          |          |          |



**WEE1-i-1 (-) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1725.386 | 2558.959 | 1963.991 | 2575.606 | 2722.202 |
| 1957.112 | 2484.367 | 2073.169 | 2050.313 | 2724.861 |
| 1905.865 | 2068.21  | 1859.771 | 2514.698 | 2948.968 |
| 2689.664 | 1950.277 | 2427.011 | 2862     | 2760.224 |
| 2343.707 | 2187.235 | 1697.19  | 1843.815 | 2867.102 |
| 2665.708 | 1846.978 | 2240.063 | 2380.637 | 2015.453 |
| 2017.322 | 1307.051 | 2590.153 | 2099.116 | 3478.832 |
| 2280.745 | 2104.363 | 2289.657 | 2681.88  | 2329.404 |
| 2247.754 | 2001.869 | 2565.141 | 2457.342 | 2765.313 |
| 1840.581 | 1742.399 | 1086.897 | 2076.331 | 1972.629 |
| 2034.716 | 2224.754 | 2143.679 | 1857.256 | 2879.429 |
| 1794.796 | 1813.34  | 1940.2   | 2631.999 | 2387.048 |
| 1224.754 | 2121.182 | 2330.482 | 1614.318 | 1948.321 |
| 1985.941 | 1583.842 | 1694.818 | 2380.076 | 2941.637 |
| 1242.866 | 2468.555 | 2345.432 | 2403.436 | 2474.304 |
| 2048.588 | 1944.009 | 1471.501 | 2589.24  | 1854.812 |
| 1880.472 | 2405.808 | 1927.334 | 2463.164 | 3003.234 |
| 2699.202 | 2432.617 | 2277.941 | 2299.145 | 1994.537 |
| 1840.221 | 2073.96  | 1599.296 | 2429.023 | 3105.800 |
| 2265.651 | 1453.317 | 2783.44  | 2300.87  | 2690.936 |
| 1828.29  | 2045.353 | 1299.792 | 2091.282 | 2569.395 |
| 2647.452 | 1706.821 | 2338.964 | 1784.662 | 2083.662 |
| 1700.496 | 2198.807 | 2526.055 | 2247.898 | 2057.859 |
| 2293.898 | 1685.618 | 2412.923 | 2321.21  | 2284.769 |
|          | 2568.821 | 1835.549 | 2563.214 | 2816.286 |
|          | 1932.94  |          | 2568.389 | 2054.912 |
|          | 2413.139 |          | 1648.027 | 3019.334 |
|          | 2648.89  |          | 2229.929 | 3164.881 |
|          | 2347.876 |          | 1798.318 | 2231.006 |
|          | 2279.379 |          | 1768.202 | 2777.258 |
|          | 2058.722 |          | 1802.199 | 2759.074 |
|          | 2067.635 |          | 2296.507 | 2950.837 |
|          |          |          | 2310.422 | 3316.829 |
|          |          |          | 3145.26  | 2898.152 |
|          |          |          | 2407.82  |          |

**WEE1-i-6 (+) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1671.45  | 1043.049 | 2654.201 | 1534.253 | 1530.252 |
| 1155.151 | 2990.205 | 1445.518 | 1793.852 | 1881.594 |
| 1116.506 | 2602.022 | 1518.222 | 1634.035 | 1649.346 |
| 1339.356 | 1049.741 | 1686.863 | 1209.36  | 2137.526 |
| 1428.601 | 1758.515 | 1595.889 | 1864.151 | 1595.513 |
| 1659.045 | 1804.078 | 1542.883 | 1475.899 | 1803.777 |
| 1741.899 | 1813.025 | 1185.3   | 1352.748 | 2012.116 |
| 1560.476 | 1195.375 | 1147.257 | 1179.511 | 1736.486 |
| 1487.321 | 1065.154 | 1610.926 | 1426.12  | 1924.525 |
| 1637.842 | 1942.945 | 1736.035 | 1497.621 | 1703.404 |
| 1307.627 | 1710.396 | 1649.421 | 1536.267 | 2249.627 |
| 1518.072 | 2430.073 | 1437.473 | 1574.072 | 2398.72  |
| 1861.82  | 948.5411 | 2012.417 | 1742.638 | 2171.885 |
| 1916.987 | 1467.322 | 2091.587 | 1660.038 | 1176.053 |
| 1754.098 | 1612.054 | 2328.046 | 1600.642 | 2853.743 |
| 2043.467 | 1292.515 | 2269.251 | 1464.765 | 2158.878 |
|          | 961.9241 | 1487.095 | 1528.735 | 2530.145 |
|          | 2078.956 | 1408.526 | 1317.101 | 2171.434 |
|          | 1550.251 | 1128.46  | 1034.102 | 2302.257 |
|          | 1365.295 | 1131.693 | 1350.784 | 5738.99  |
|          | 929.5192 | 1910.24  | 1563.264 | 1947.908 |
|          | 932.2258 | 1442.736 | 1545.986 | 2167.473 |
|          | 1980.388 | 1774.079 | 1750.743 | 1935.902 |
|          | 1735.374 | 1591.829 | 1764.133 | 2223.478 |
|          | 1658.936 | 1496.569 | 15296.46 |          |
|          | 2234.154 | 1628.745 | 1698.305 |          |
|          | 1556.305 | 2090.986 | 1398.473 |          |
|          | 1856.362 |          |          |          |

**WEE1-i-6 (-) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 2188.888 | 1498.465 | 1257.107 | 2666.727 | 2212.048 |
| 2413.714 | 2502.354 | 1770.659 | 2265.164 | 2856.629 |
| 2152.519 | 1159.931 | 1275.291 | 2903.054 | 2644.086 |
| 1784.59  | 1567.538 | 1320.573 | 3464.037 | 2811.7   |
| 1834.399 | 1558.122 | 1607.429 | 2899.963 | 2859.066 |
| 1670.093 | 2349.259 | 1620.295 | 2101.503 | 2557.116 |
| 1275.282 | 2305.702 | 1458.215 | 2632.226 | 3368.521 |
| 2346.946 | 2906.511 | 1905.857 | 2283.205 | 3072.147 |
| 1920.362 | 1830.46  | 1721.783 | 1701.73  | 2226.495 |
| 3633.221 | 1799.265 | 2631.515 | 2370.965 | 3029.915 |
| 1874.937 | 2012.808 | 2559.351 | 2484.737 | 2579.836 |
| 1878.171 | 1934.032 | 2047.74  | 1629.279 | 3208.67  |
| 1818.228 | 2061.828 | 1665.217 | 2328.127 | 2893.063 |
| 1343.492 | 2333.805 | 1811.987 | 1697.346 | 2665.145 |
| 1472.436 | 2009.143 | 1874.304 | 2443.337 | 1996.277 |
| 1402.429 | 2332.368 | 2047.884 | 2111.925 | 1926.989 |
| 1258.607 | 2280.474 | 1709.493 | 2370.462 | 2738.818 |
| 2240.063 | 1868.769 | 1816.659 | 2456.857 | 2866.189 |
| 1700.855 | 1870.279 | 1337.895 | 1419.546 | 3095.968 |
| 1386.832 | 2024.955 | 2517.16  | 1280.394 | 2100.209 |
| 2448.322 | 2425.95  | 2209.173 | 2229.435 | 2155.697 |
| 1254.654 | 2139.956 | 3533.627 | 2675.646 | 2109.984 |
| 1681.233 | 2311.524 | 2670.831 | 2599.099 | 2721     |
| 1234.816 | 1998.217 | 2302.108 | 1905.857 | 2097.047 |
| 1901.387 | 1675.639 | 2728.331 | 1768.718 | 2760.46  |
| 2093.797 | 1761.099 | 1706.043 | 1671.614 | 2572.002 |
| 2363.114 | 1815.222 | 2203.063 | 2678.018 | 2370.749 |
| 1951.7   | 1743.49  | 2166.838 | 2924.552 | 2234.689 |
| 1486.02  | 1878.185 | 2296.214 | 1944.526 | 3192.066 |
| 2151.888 | 2196.091 | 2313.824 | 1600.529 | 3198.32  |
| 1851.232 | 2516.945 | 2472.454 | 2237.923 | 3031.28  |
| 1987.22  | 4154.633 | 1135.924 | 3074.988 | 2344.155 |
| 2535.201 |          | 1716.536 | 2165.4   | 3153.325 |
| 2097.981 |          | 3378.304 | 1637.401 | 2767.568 |
| 1732.708 |          | 2360.831 | 2724.45  | 2938.848 |
| 1799.769 |          | 2152.032 | 2324.533 | 2784.387 |
| 2093.812 |          | 1803.865 | 3169.785 | 2943.161 |
| 1381.236 |          |          | 3381.595 | 3095.609 |
| 1912.829 |          |          | 3651     | 3175.679 |
| 1902.407 |          |          | 2813.073 | 2155.769 |
|          |          |          |          | 2954.023 |

**EV-i-1 (+) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1783.74  | 2396.768 | 1340.483 | 2772.679 | 2046.662 |
| 2463.972 | 2687.146 | 1896.01  | 1942.945 | 1998.002 |
| 1850.441 | 1807.531 | 2092.518 | 1845.985 | 1557.404 |
| 2513.351 | 2020.499 | 1558.482 | 2001.021 | 1453.974 |
| 2246.692 | 2980.184 | 1968.964 | 2124.288 | 2148.582 |
| 2808.257 | 1912.038 | 1847.135 | 2086.122 | 2228.723 |
| 1898.31  | 2333.374 | 2030.921 | 2356.015 | 2295.208 |
| 3234.265 | 2081.45  | 2802.435 | 1551.078 | 2475.832 |
| 1211.537 | 2316.34  | 1460.659 | 2231.382 | 2523.414 |
| 1429.177 | 2388.215 | 2317.849 | 2063.696 | 2951.865 |
| 1660.545 | 3341.288 | 2603.699 | 2540.736 | 2101.215 |
| 1527.647 | 2293.411 | 2861.805 | 3418.698 | 2335.315 |
| 1407.255 | 2564.814 | 3215.433 | 2832.192 | 2258.12  |
| 1492.787 | 1681.605 | 2355.656 | 2753.416 | 2552.451 |
| 1138.009 | 2007.274 | 2748.025 | 2008.783 | 2676.94  |
| 1212.256 | 2332.152 | 2013.527 | 2389.868 | 2867.626 |
| 1288.732 | 1957.392 | 2993.912 | 2406.903 | 3585.593 |
| 2377.65  | 2149.3   | 2050.04  | 2366.293 | 2445.213 |
| 2542.532 | 2316.699 | 2489.776 | 2257.833 | 1737.093 |
| 2537.271 | 2025.53  | 2435.941 | 3652.869 | 1749.456 |
| 2659.043 | 1959.98  | 1794.234 | 2107.612 | 3184.598 |
| 1862.229 | 2748.672 | 2733.722 | 3112.22  | 3380.891 |
| 2518.095 | 2220.314 | 2427.819 | 2288.667 | 3223.914 |
| 2099.49  | 1589.101 | 2331.074 | 2239.289 | 2617.786 |
| 2227.07  | 2993.337 | 2350.265 | 2357.021 | 2264.805 |
| 2149.228 | 2826.585 | 1716.752 | 1860.719 | 1924.76  |
| 2741.413 | 2172.516 | 2104.881 | 1810.622 | 2223.476 |
| 1826.291 | 1960.123 | 2770.522 | 2410.712 | 2396.912 |
| 1703.311 | 2095.681 | 2306.205 | 1813.281 | 1900.179 |
| 1635.101 | 2085.618 | 2270.411 | 2838.804 | 1820.469 |
| 2060.893 | 2716.903 | 1399.924 |          | 2207.52  |
| 2554.967 | 2502.498 | 2655.809 |          | 2195.588 |
| 2367.946 | 2376.212 | 2159.794 |          | 2189.623 |
| 2049.896 | 2856.486 | 2524.132 |          | 2335.171 |
| 3010.731 |          | 2107.828 |          | 2745.869 |
| 1773.318 |          | 2130.325 |          |          |
| 1850.01  |          | 2597.805 |          |          |
| 1705.324 |          | 2165.76  |          |          |
| 2101     |          |          |          |          |
| 2174.026 |          |          |          |          |
| 2303.977 |          |          |          |          |
| 2851.311 |          |          |          |          |

**EV-i-1 (-) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1761.315 | 2079.437 | 2084.468 | 2576.53  | 3670.335 |
| 1881.204 | 2699.725 | 1921.526 | 1942.442 | 1572.857 |
| 2133.56  | 2081.953 | 2263.583 | 2708.35  | 1765.556 |
| 2192.354 | 2443.128 | 2368.378 | 1969.97  | 2469.147 |
| 2341.006 | 3195.739 | 2752.266 | 2058.377 | 2574.373 |
| 1445.637 | 3009.437 | 2434.359 | 2232.101 | 2490.854 |
| 2743.928 | 3259.062 | 2733.075 | 2401.225 | 3037.109 |
| 1665.145 | 1529.803 | 2649.699 | 2347.534 | 2773.613 |
| 2456.497 | 2997.003 | 2117.747 | 2356.949 | 2125.941 |
| 2218.013 | 2475.329 | 2244.464 | 3115.813 | 1921.526 |
| 1448.009 | 2554.464 | 2095.839 | 3408.635 | 2417.037 |
| 2412.509 | 2727.253 | 2642.296 | 2199.613 | 2228.435 |
| 2640.068 | 1500.406 | 2549.289 | 1844.619 | 2229.37  |
| 1248.985 | 2131.331 | 2128.959 | 2009.789 | 2337.615 |
| 1878.257 | 2502.641 | 1968.978 | 2971.774 | 2561.508 |
| 2083.462 | 2354.434 | 2895.155 | 2616.708 | 2319.574 |
| 2319.071 | 2684.631 | 2394.476 | 1942.514 | 2062.762 |
| 2174.529 | 3259.709 | 2026.903 | 2083.606 | 1865.319 |
| 2217.798 | 2727.253 | 2366.868 | 2102.94  | 1865.535 |
| 2215.067 | 3089.076 | 2845.92  | 2744.503 | 2430.191 |
| 2651.352 | 3128.895 | 2709.715 | 1867.332 | 2497.97  |
| 1990.239 | 1984.345 | 2616.996 | 2198.679 |          |
| 2479.138 | 2435.725 | 2314.119 | 2333.662 |          |
| 3067.8   | 2518.382 | 2065.206 | 1948.551 |          |
| 2551.373 | 1833.622 | 2537.645 | 2131.763 |          |
| 1846.703 | 3169.72  | 2487.195 |          |          |
| 1633.232 | 2335.674 | 2493.729 |          |          |
| 2286.727 | 1489.193 | 2289.17  |          |          |
| 2257.689 | 1882.066 | 2938.208 |          |          |
| 2008.136 | 2215.857 | 2299.312 |          |          |
| 2433.425 | 1824.35  | 2848.148 |          |          |
| 2344.371 | 1978.739 | 2152.039 |          |          |
| 2457.863 | 2565.892 | 3177.195 |          |          |
| 1722.79  | 2544.617 | 2726.75  |          |          |
| 1610.807 | 2140.603 | 3334.316 |          |          |
| 1887.888 | 2679.312 | 2711.8   |          |          |
| 1606.279 | 2167.269 | 2679.528 |          |          |
| 3208.964 | 4330.585 | 2282.206 |          |          |
| 2618.433 | 2698.934 | 2140.754 |          |          |
| 2918.083 | 2727.469 | 2045.735 |          |          |
| 2081.665 |          | 2702.463 |          |          |
| 1888.463 |          |          |          |          |
| 2031.568 |          |          |          |          |
| 2891.633 |          |          |          |          |

**EV-i-3 (+) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1794.796 | 1307.051 | 2405.808 | 2565.141 | 2553.367 |
| 2427.011 | 1599.296 | 1700.496 | 1086.897 | 1583.267 |
| 1828.29  | 2783.44  | 2293.898 | 2143.679 | 1781.571 |
| 2279.379 | 1299.792 | 1932.94  | 1940.2   | 4418.817 |
| 1058.722 | 2048.588 | 1413.139 | 2330.482 | 2489.614 |
| 2067.635 | 1880.472 | 1648.89  | 1694.818 | 1860.49  |
| 2045.353 | 2699.202 | 2347.876 | 1625.386 | 1919.572 |
| 1706.821 | 1840.221 | 2121.182 | 2345.432 | 1846.69  |
| 2198.807 | 2001.869 | 1963.991 | 1471.501 | 2460.936 |
| 1685.618 | 1742.399 | 2073.169 | 1583.842 | 2423.776 |
| 2568.821 | 2224.754 | 1859.771 | 1468.555 | 1932.294 |
| 1976.497 | 2027.6   | 1765.974 | 1474.161 | 3136.85  |
| 2523.395 | 2039.819 | 1927.19  | 1627.902 | 3051.103 |
| 2147.488 | 2533.602 | 1475.958 | 2898.153 | 2190.829 |
| 1846.69  | 1484.655 | 1619.421 | 1502.192 | 2971.106 |
| 1627.543 | 2176.382 | 1420.973 | 2082.153 | 2614.893 |
| 2903.04  | 1975.994 | 1913.462 | 1446.417 | 2324.876 |
| 2613.096 | 1765.184 | 1863.15  | 1830.159 | 1719.974 |
| 2836.34  | 2366.779 | 2703.443 | 2300.007 | 2188.888 |
| 2417.236 | 1847.193 | 1415.439 | 2293.035 | 2413.714 |
| 2583.699 | 2524.258 | 2072.45  | 1722.202 | 2152.519 |
| 2361.748 | 1882.7   | 1974.053 | 1724.862 | 1784.59  |
| 1892.69  | 1579.099 | 1809.89  | 1948.969 | 1834.399 |
| 1992.453 | 1201.035 | 1785.884 | 2760.224 | 1670.093 |
| 2669.661 | 1135.341 | 2282.038 | 1867.103 | 1275.282 |
| 2500.18  | 1176.022 | 1681.88  | 1015.453 | 2346.946 |
| 3106.447 | 1591.317 | 2257.313 | 1478.833 |          |
| 2104.003 | 1654.783 | 2389.391 | 1105.8   |          |
| 1948.969 | 1735.571 | 1647.524 | 1690.937 |          |
| 1813.124 | 1368.504 | 1927.909 | 2569.396 |          |
| 1649.465 | 1379.429 | 2099.835 | 2083.663 |          |
|          | 1387.048 | 2505.189 | 2057.86  |          |
|          | 1948.322 | 2082.01  | 2284.77  |          |
|          | 1941.637 | 2099.116 | 2575.606 |          |
|          | 1474.305 | 2681.88  | 2050.313 |          |
|          | 1854.812 | 1457.342 | 2514.698 |          |
|          | 2003.234 | 1378.567 | 1862     |          |
|          | 1994.537 | 2248.688 | 1843.815 |          |
|          | 1568.389 | 2371.02  | 2380.637 |          |
|          | 1648.027 | 2287.213 |          |          |
|          | 2229.929 | 1415.295 |          |          |
|          | 1798.318 | 2202.113 |          |          |
|          | 1823.187 | 1588.227 |          |          |
|          | 1738.087 | 1768.993 |          |          |
|          | 2040.538 |          |          |          |

**EV-i-3 (-) dex**

| 0 h      | 12 h     | 24 h     | 36 h     | 48 h     |
|----------|----------|----------|----------|----------|
| 1944.009 | 1647.452 | 1813.34  | 2292.554 | 2294.76  |
| 1857.112 | 1104.363 | 1432.617 | 2420.283 | 1637.03  |
| 1805.865 | 1224.754 | 2073.96  | 2241.716 | 1771.868 |
| 2589.664 | 1185.941 | 1453.317 | 2666.571 | 1474.951 |
| 2343.707 | 1242.866 | 1697.19  | 1980.809 | 1365.917 |
| 2665.708 | 2558.959 | 2240.063 | 2325.739 | 2041.256 |
| 2017.322 | 1484.367 | 2590.153 | 1952.778 | 2998.203 |
| 2280.745 | 1068.21  | 2289.657 | 2147.344 | 2832.028 |
| 2247.754 | 1550.277 | 2265.651 | 2120.535 | 2603.393 |
| 1840.581 | 2187.235 | 1927.334 | 1540.574 | 2337.239 |
| 2034.716 | 1846.978 | 2277.941 | 1393.229 | 2335.298 |
| 2698.627 | 1175.735 | 2866.025 | 1801.409 | 3367.929 |
| 1886.078 | 1810.178 | 2245.813 | 3589.161 | 2457.126 |
| 2355.207 | 1087.544 | 2294.401 | 2440.811 | 2076.044 |
| 2976.497 | 1082.225 | 2616.689 | 1814.059 | 1900.884 |
| 1888.953 | 2445.303 | 1811.543 | 2303.529 | 2417.02  |
| 2170.776 | 1651.621 | 2224.323 | 1799.396 | 1924.819 |
| 2066.269 | 1782.721 | 1647.524 | 2103.428 | 2273.341 |
| 2511.033 | 1665.205 | 1472.436 | 2018.759 | 1943.65  |
| 3171.279 | 1909.222 | 1402.429 | 1937.612 | 1592.036 |
| 2248.76  | 2144.182 | 1258.607 | 1898.728 | 1389.132 |
| 2484.942 | 1732.48  | 2240.063 | 2482.225 | 1707.18  |
| 2254.726 | 1920.362 | 1700.855 | 2591.246 | 2764.752 |
| 2047.078 | 2076.331 | 1386.832 | 2338.964 | 2246.316 |
| 2899.734 | 1857.256 | 2448.322 | 2526.055 | 2575.936 |
| 3298.57  | 2631.999 | 1254.654 | 2412.923 | 1877.74  |
| 2724.574 | 1614.318 | 1681.233 | 1835.549 | 2403.436 |
| 1405.233 | 1380.076 | 1234.816 | 1777.259 | 2589.24  |
| 1704.665 | 1797.599 | 1901.387 | 2759.074 | 2463.164 |
| 1461.151 | 1587.436 | 2093.797 |          | 2299.145 |
| 2130.525 | 2192.554 | 2363.114 |          | 2429.023 |
| 1642.277 | 2240.998 | 1951.7   |          | 3145.26  |
|          | 1733.415 | 1486.02  |          | 2407.82  |
|          | 1308.488 | 1950.837 |          |          |
|          | 1294.904 | 2316.826 |          |          |
|          | 1971.537 | 1898.153 |          |          |
|          | 2455.761 |          |          |          |
|          | 1769.856 |          |          |          |
|          | 2191.835 |          |          |          |
|          | 2225.041 |          |          |          |
|          | 2046.503 |          |          |          |
|          | 1723.209 |          |          |          |
|          | 1869.978 |          |          |          |
|          | 1734.996 |          |          |          |
|          | 2300.87  |          |          |          |
|          | 2091.282 |          |          |          |
|          | 1784.662 |          |          |          |
|          | 2247.898 |          |          |          |
|          | 2321.21  |          |          |          |
|          | 2563.214 |          |          |          |
|          | 1965.356 |          |          |          |

1838.496  
1768.202  
1802.199  
2296.507  
2310.422



## **Appendix III: Tobacco BY-2 mitotic index (MI %)**

### **Wild type**

| Time (h) | MI (%) |
|----------|--------|
| 0        | 0      |
| 1        | 0      |
| 2        | 0      |
| 3        | 0.66   |
| 4        | 1.6    |
| 5        | 2.6    |
| 6        | 13.1   |
| 7        | 28.1   |
| 8        | 34.7   |
| 9        | 15.2   |
| 10       | 7.7    |
| 11       | 3.6    |
| 12       | 4.3    |
| 13       | 1.3    |
| 14       | 1      |
| 15       | 2.3    |
| 16       | 2      |
| 17       | 3.8    |
| 18       | 4.1    |
| 19       | 7.1    |
| 20       | 8.8    |
| 21       | 15.4   |
| 22       | 20.1   |
| 23       | 13.4   |
| 24       | 5.8    |

**Constitutive lines**

| Time (h) | EV-c-10 (+)hyg MI (%) | EV-c-10 (-)hyg MI (%) | EV-c-10 (-)hyg 4 week MI (%) | WEE1-c-2 (+)hyg MI (%) | WEE1-c-10 (+)hyg MI (%) | WEE1-c-10 (-)hyg MI (%) | WEE1-c-12 (+)hyg MI (%) | WEE1-c-12 (-)hyg MI (%) |
|----------|-----------------------|-----------------------|------------------------------|------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 0        | 0                     | 0                     | 0                            | 1                      | 0.33                    | 1                       | 0.94                    | 0                       |
| 1        | 0                     | 0                     | 0                            | 1.75                   | 2.2                     | 1.7                     | 1.8                     | 6                       |
| 2        | 0                     | 0                     | 0.33                         | 5.3                    | 10.3                    | 2                       | 1.3                     | 8.6                     |
| 3        | 0                     | 0                     | 0                            | 20                     | 17.4                    | 14.6                    | 1.9                     | 12.2                    |
| 4        | 0                     | 0                     | 0.33                         | 20                     | 22                      | 31.6                    | 8.7                     | 21.2                    |
| 5        | 0                     | 0.66                  | 5.2                          | 13                     | 10.3                    | 18                      | 13.8                    | 12.6                    |
| 6        | 0.33                  | 1.66                  | 6                            | 6.4                    | 9.3                     | 12.4                    | 10.2                    | 8.8                     |
| 7        | 1.99                  | 7.95                  | 15.7                         | 6.1                    | 9.4                     | 9.3                     | 8.6                     | 9.8                     |
| 8        | 14.33                 | 24.43                 | 23.3                         | 3                      | 6.4                     | 6.5                     | 9.4                     | 7.4                     |
| 9        | 18.99                 | 27.66                 | 29.7                         | 3.5                    | 4.3                     | 5.5                     | 10.8                    | 6.2                     |
| 10       | 15.66                 | 17.33                 | 15.5                         | 4.4                    | 2                       | 2.3                     | 8.6                     | 4.3                     |
| 11       | 10.96                 | 7.33                  | 5                            | 3.7                    | 2.3                     | 2                       | 4.6                     | 6.6                     |
| 12       | 7.8                   | 4.65                  | 2.3                          | 2.6                    | 2.3                     | 2.3                     | 2.5                     | 5.3                     |
| 13       | 7.3                   | 1.33                  | 2.6                          | 6.8                    | 2.9                     | 3.3                     | 4.3                     | 7                       |
| 14       | 4.8                   | 2.6                   | 6.3                          | 11.35                  | 6                       | 5.8                     | 3                       | 5.4                     |
| 15       | 2.56                  | 3.3                   | 1.2                          | 15.4                   | 11.7                    | 9.9                     | 2.4                     | 6.8                     |
| 16       | 2.17                  | 0.92                  | 1.3                          | 15.6                   | 11.8                    | 15.5                    | 3.6                     | 6.6                     |
| 17       | 1                     | 2.6                   | 3.2                          | 14.3                   | 14.4                    | 14.9                    | 5.2                     | 7.8                     |
| 18       | 2.3                   | 2.3                   | 5.5                          | 7.6                    | 11.5                    | 10.7                    | 9.8                     | 14.7                    |
| 19       | 2                     | 2                     | 5.3                          | 7                      | 10.3                    | 11.4                    | 5.2                     | 7.3                     |
| 20       | 8                     | 7.3                   | 11.4                         | 6.2                    | 8                       | 9.2                     | 4.7                     | 8.3                     |
| 21       | 1.97                  | 12.3                  | 6.7                          | 5.6                    | 6.5                     | 8.5                     | 4                       | 4.8                     |
| 22       | 2.8                   | 5.3                   | 3.3                          | 3.1                    | 5.6                     | 6.7                     | 3.6                     | 4.7                     |
| 23       | 1.33                  | 3.5                   |                              |                        | 4.2                     | 5.7                     | 3.3                     | 3.5                     |

## **Inducible lines**

| Time<br>(h) | EV-i-1<br>(-)dex<br>MI (%) | EV-i-1<br>(+)dex<br>MI (%) | WEE1-i-1<br>(+)dex<br>MI (%) | WEE1-i-1<br>(-) dex<br>MI (%) | WEE1-i-6<br>(+) dex<br>MI (%) |
|-------------|----------------------------|----------------------------|------------------------------|-------------------------------|-------------------------------|
| 0           | 0                          | 0                          | 1.3                          | 2.2                           | 0.66                          |
| 1           | 0                          | 0.3                        | 0.3                          | 0.6                           | 3.6                           |
| 2           | 0.3                        | 0.65                       | 1.3                          | 1.66                          | 11.9                          |
| 3           | 0.6                        | 0.68                       | 7                            | 2.3                           | 14                            |
| 4           | 1.26                       | 1.33                       | 14.7                         | 3.5                           | 28.3                          |
| 5           | 4.7                        | 7.6                        | 26.2                         | 11.44                         | 10.3                          |
| 6           | 13.3                       | 17.3                       | 15.9                         | 11.8                          | 6.4                           |
| 7           | 20.44                      | 16.07                      | 12.8                         | 25.5                          | 4                             |
| 8           | 30.7                       | 23.5                       | 9.9                          | 18.83                         | 1.3                           |
| 9           | 23.4                       | 8.7                        | 10.2                         | 16.3                          | 2.5                           |
| 10          | 21.24                      | 5.1                        | 8                            | 12                            | 1.2                           |
| 11          | 23.3                       | 4.16                       | 5.1                          | 9.2                           | 3                             |
| 12          | 16.4                       | 3                          | 2.5                          | 6.09                          | 1.5                           |
| 13          | 7.9                        | 2.24                       | 2.6                          | 5.23                          | 11.5                          |
| 14          | 3.5                        | 1.6                        | 1.8                          | 3.2                           | 12.6                          |
| 15          | 1.6                        | 2.23                       | 1.8                          | 2.7                           | 9.3                           |
| 16          | 2.15                       | 3.7                        | 2.8                          | 3.3                           | 4.6                           |
| 17          | 4.3                        | 9.24                       | 5.3                          | 2.65                          | 4.8                           |
| 18          | 1.9                        | 13.16                      | 12.5                         | 5                             | 1.1                           |
| 19          | 4.4                        | 14                         | 4.2                          | 7.3                           | 0.93                          |
| 20          | 14.24                      | 9.4                        | 6.7                          | 15.6                          | 1                             |
| 21          | 6.3                        | 9.1                        | 3.9                          | 12.54                         |                               |
| 22          | 9.65                       | 5.23                       |                              | 8.33                          |                               |
| 23          | 10.8                       | 2.7                        |                              | 4                             |                               |
| 24          | 7.23                       |                            |                              |                               |                               |

**Appendix IV: 18S rRNA, H4, AtWEE1 and NtWEE1 expression raw data**

**Wild type**

| Time (h) | 18S     | 18S     | 18S     |         |         | NtWEE1    | NtWEE1   | NtWEE1    |
|----------|---------|---------|---------|---------|---------|-----------|----------|-----------|
|          | rRNA 1  | rRNA 2  | rRNA 3  | H4 1    | H4 2    | 1         | 2        | 3         |
| 0        | 1678.96 | 2500.36 | 2805.84 | 2519.95 | 2832.32 | 123.98345 | 99.87329 | 122.98264 |
| 1        | 316.063 | 19173.8 | 26880.9 | 14143.8 | 14098.5 | 3119.67   | 3469     | 3565.83   |
| 2        | 16628.9 | 9227.1  | 21715.4 | 5951.18 | 5297.09 | 3190.1    | 2974.95  | 3287.22   |
| 3        | 29762.8 | 10363.4 | 13932.7 | 5150.96 | 5314.65 | 5088.82   | 5276.13  | 4976.27   |
| 4        | 31001.7 | 10603.9 | 13952.6 | 5049.62 | 5087.25 | 6651.29   | 6531.62  | 6746.44   |
| 5        | 26638.5 | 8927.72 | 32126.8 | 392.403 | 262.624 | 2432.999  | 2369.44  | 2325.95   |
| 6        | 40526.3 | 21083.1 | 42430.2 | 638.129 | 201.759 | 529.819   | 571.115  | 537.855   |
| 7        | 14417.8 | 12632   | 17556.4 | 727.067 | 517.635 | 515.295   | 552.901  | 577.787   |
| 8        | 27049.7 | 25112.5 | 21523.4 | 974.275 | 858.236 | 1501.54   | 1574.5   | 1414.01   |
| 9        | 26831.8 | 2868.53 | 3466.41 | 421.222 | 422.618 | 615.002   | 575.911  | 599.057   |
| 10       | 21555.7 | 2685.29 | 25499.9 | 550.827 | 465.19  | 694.11    | 720.858  | 643.055   |
| 11       | 20280.3 | 36744.4 | 38743.5 | 427.655 | 190.474 | 722.979   | 734.446  | 786.066   |
| 12       | 27984.1 | 1284.77 | 1254.65 | 762.246 | 413.764 | 622.664   | 660.884  | 645.232   |
| 13       | 9545.08 | 840.591 | 9305.24 | 866.615 | 410.083 | 1575.2    | 1482.61  | 1579.25   |
| 14       | 984.009 | 10582.6 | 10726.7 | 575.967 | 246.416 | 860.998   | 886.396  | 893.399   |
| Mix 25%  | 27145.5 | 36766.9 | 567.938 | 1848.42 | 2094.84 | 656.779   | 805.031  | 939.569   |
| Mix 50%  | 2483.11 | 9364.79 | 13627.2 | 3187.67 | 3426.05 | 2019      | 2273.7   | 2349.57   |
| Mix 100% | 1440.65 | 36139.2 | 45205.3 | 7417.57 | 8334.92 | 4004.66   | 6459.02  | 4875.22   |

**WEE1-i-1 non-induced**

| Time (h) | NtWEE1 1 | NtWEE1 2 | NtWEE1 3 | NtWEE1 4 | NtWEE1 5 |
|----------|----------|----------|----------|----------|----------|
| 0        | 495.469  | 12169.5  | 333.182  | 2648.08  | 13323.7  |
| 1        | 2688.3   | 6589     | 1627.3   | 9987.4   | 9057.5   |
| 2        | 5330     | 5078.6   | 2296.5   | 7045.6   | 7594     |
| 3        | 21321.19 | 15814.5  | 17308.7  | 12731.7  | 25901.9  |
| 4        | 35739.3  | 49037.1  | 43622.9  | 33166.3  | 49141.2  |
| 5        | 7259.12  | 8301.44  | 1532.87  | 6573.92  | 3182.95  |
| 6        | 13859.9  | 15916.6  | 825.609  | 25127.3  | 28241.7  |
| 7        | 31439.9  | 20528.9  | 20158.6  | 20241.4  | 23657.6  |
| 8        | 34497.7  | 12761.5  | 18581.6  | 25065.3  | 16251.4  |
| 9        | 37264.5  | 42651.8  | 29210.9  | 25690.9  | 43869.5  |
| 10       | 33081.7  | 42963.1  | 23256.3  | 19112.2  | 44673.1  |
| 11       | 27297.3  | 44971    | 36292.1  | 4720.7   | 47016.5  |
| 12       | 30503.6  | 14792.5  | 17388.2  | 24900.8  | 18495.6  |
| 13       | 36982.8  | 10068.6  | 23479    | 28323.7  | 13381.7  |
| 14       | 29699.9  | 42068.5  | 11248.6  | 31138.5  | 47182.5  |
| Mix 25%  | 17745.8  | 38862.8  | 18048.8  | 31806.6  | 31159.6  |
| Mix 50%  | 21270.1  | 41168.7  | 21087.7  | 31434.7  | 34231.6  |
| Mix 100% | 39134.3  | 49860.5  | 18808.2  | 33999.1  | 46264.8  |

| Time (h) | H4 1    | H4 2    | H4 3    | NtWEE1 1 | NtWEE1 2 | NtWEE1 3 |
|----------|---------|---------|---------|----------|----------|----------|
| 0        | 459.023 | 403.41  | 341.103 | 579.999  | 599.5    | 583.873  |
| 1        | 4592.3  | 4963.08 | 4034.19 | 1113.44  | 1072.35  | 1123.05  |
| 2        | 4117.79 | 4652.56 | 4436.19 | 2894.02  | 3185.23  | 2998.12  |
| 3        | 11862.2 | 11659.5 | 10918.6 | 11221.8  | 11832    | 11150.2  |
| 4        | 8684.95 | 8869.32 | 8288.69 | 8683.93  | 9581.7   | 8300.45  |
| 5        | 1269.09 | 1994.96 | 1325    | 1115.35  | 1166.302 | 1012.07  |
| 6        | 2007.26 | 2593.14 | 2288.93 | 3275.82  | 3529.5   | 2790.85  |
| 7        | 2546.10 | 2633.36 | 2373.56 | 292.68   | 238.792  | 215.907  |
| 8        | 2374.24 | 2446.82 | 2611.08 | 1394.49  | 2247.16  | 1430.95  |
| 9        | 1540.54 | 1785.88 | 1366.95 | 1523.34  | 1648.26  | 1186.42  |
| 10       | 1820.02 | 1627.89 | 1726.38 | 491.986  | 490.781  | 420.791  |
| 11       | 1864.44 | 1948.3  | 1834.90 | 4905.1   | 4136.49  | 4331.91  |
| 12       | 1953.61 | 1856.86 | 1981.91 | 739.086  | 929.952  | 630.717  |
| 13       | 1601.55 | 1704.62 | 1995.13 | 723.85   | 898.951  | 564.839  |
| 14       | 2118.34 | 2423.02 | 2160.3  | 859.15   | 854.153  | 851.6    |
| Mix 25%  | 815.084 | 977     | 1154.21 | 389.032  | 215.963  | 733.338  |
| Mix 50%  | 1837.65 | 2146.55 | 2077.51 | 2193.7   | 2014.64  | 2833.62  |
| Time (h) | 3223.12 | 7229.29 | 3278.89 | 6437.23  | 4606.74  | 8224.3   |

### **WEE1-i-1 induced**

| Time (h) | 18S     |         |         | H4 1    | H4 2    | H4 3    | NtWEE1  |         |         |
|----------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
|          | rRNA 1  | rRNA 2  | rRNA 3  |         |         |         | 1       | 2       | 3       |
| 0        | 11803.9 | 22835.2 | 11187.4 | 983.092 | 973.127 | 979.382 | 278.984 | 235.359 | 288.865 |
| 1        | 11187.5 | 17571.8 | 15805.4 | 7487.04 | 7732.61 | 7539.94 | 1570.55 | 1340.39 | 832.673 |
| 2        | 10411.9 | 12083.3 | 6429.79 | 2865.54 | 2821.24 | 2961.02 | 590.213 | 670.411 | 274     |
| 3        | 5784.67 | 8101.52 | 7729.67 | 1804.21 | 1679.86 | 1235.62 | 1474.12 | 953.229 | 1522.93 |
| 4        | 27484.1 | 43924.6 | 24316.5 | 2863.07 | 2538.14 | 2152.61 | 8491.16 | 7573.75 | 7742.24 |
| 5        | 9334.71 | 13549.5 | 9611.84 | 1162.07 | 1871.3  | 1199.36 | 2523.09 | 3384.89 | 3823.87 |
| 6        | 7955.58 | 8679.52 | 7916.78 | 358.359 | 212.934 | 249.637 | 8227.3  | 8884.89 | 8309.6  |
| 7        | 7168.45 | 6098    | 5354.24 | 920.042 | 1027    | 684.375 | 3219.73 | 2610.2  | 2486.98 |
| 8        | 3574.86 | 4477.44 | 2714    | 390.894 | 485.573 | 500.441 | 1432.33 | 1446.79 | 1933.56 |
| 9        | 5848.82 | 8213.13 | 6597.89 | 593.158 | 454.379 | 445.071 | 1565.45 | 648.49  | 915.741 |
| 10       | 2528.4  | 3704.22 | 2447.69 | 209.564 | 501.111 | 307.434 | 741.371 | 564.717 | 151.201 |
| 11       | 16257.6 | 23567.4 | 15355.8 | 1073.97 | 879.143 | 1368.15 | 603.109 | 614.046 | 607.48  |
| 12       | 2496.26 | 3277.24 | 2774.32 | 611.102 | 495.428 | 727.056 | 654.986 | 387.182 | 892.84  |
| Mix 25%  | 7658.93 | 4505.89 | 4799.01 | 1486.49 | 1183.66 | 1035.84 | 11861   | 17346.6 | 22675.8 |
| Mix 50%  | 21515   | 14056   | 15188.9 | 2976.64 | 2247.85 | 2829.29 | 17572.8 | 24814.7 | 34549.9 |
| Mix 100% | 36511.2 | 27775.7 | 34514.6 | 4517.91 | 3688.21 | 4551.46 | 27087.3 | 35007.5 | 37649.5 |

| Time (h) | AtWEE1<br>1 | AtWEE1<br>2 | AtWEE1<br>3 |
|----------|-------------|-------------|-------------|
| 0        | 240.981     | 180.564     | 129.098     |
| 1        | 1249.19     | 1122.30     | 1481.81     |
| 2        | 1279.09     | 1347.47     | 1298.05     |
| 3        | 1950.38     | 2188.94     | 2219.59     |
| 4        | 3278.19     | 3266.96     | 3377.34     |
| 5        | 1236.64     | 1130.48     | 1371.76     |
| 6        | 440.263     | 259.323     | 1099.2      |
| 7        | 827.35      | 407.589     | 940.232     |
| 8        | 463.024     | 212.948     | 491.977     |
| 9        | 615.6       | 624.381     | 922.031     |
| 10       | 307.974     | 493.804     | 371.705     |
| 11       | 873.56      | 1176.80     | 1047.72     |
| 12       | 750.33      | 589.192     | 366.92      |
| Mix 25%  | 542.081     | 413.654     | 262.301     |
| Mix 50%  | 6670.1      | 6077.28     | 4754.64     |
| Mix 100% | 15675       | 13710.3     | 10341.1     |

## Appendix V: CDKA and CDKB1 kinase activity raw data as percentage of maximum

### WEE1-i-1 non induced

| Time (h) | CDKA 1    | CDKA 2     | CDKA 3    | CDKA 4    | CDKA 5   |
|----------|-----------|------------|-----------|-----------|----------|
| 1        | 100       | 100        |           |           |          |
| 2        | 67.309014 | 65.5573503 |           |           |          |
| 3        | 55.219807 | 57.4142102 |           |           |          |
| 4        | 57.409191 | 57.7932567 |           |           |          |
| 5        | 65.878511 | 57.4523634 | 90.932333 | 36.465628 | 60.05409 |
| 6        | 62.513976 | 54.8004999 | 75.599645 | 53.795134 | 80.38205 |
| 7        | 65.528723 | 70.8865126 | 100       | 62.109355 | 50.5921  |
| 8        |           |            | 81.641436 | 72.02204  | 99.33326 |
| 9        |           |            | 83.738376 | 100       | 100      |
| 10       |           |            | 75.529407 | 56.378313 | 92.77344 |
| 12       |           |            | 78.932017 | 60.916056 | 72.38456 |

| Time (h) | CDKB1 1    | CDKB1 2     | CDKB1 3     | CDKB1 4     | CDKB1 5    |
|----------|------------|-------------|-------------|-------------|------------|
| 1        | 69.0226538 | 34.27793961 |             |             |            |
| 2        | 54.6464205 | 33.07220874 |             |             |            |
| 3        | 55.1390765 | 40.1736447  |             |             |            |
| 4        | 53.2202205 | 24.74461312 |             |             |            |
| 5        | 60.4503629 | 65.28348997 | 91.31264962 | 100         | 94.0466846 |
| 6        | 68.5456619 | 70.0396686  | 100         | 99.19674386 | 100        |
| 7        | 100        | 100         | 84.0908945  | 90.37903514 | 85.6250143 |
| 8        |            |             | 62.65219149 | 79.23067636 | 66.5381303 |
| 9        |            |             | 69.4984821  | 66.02394678 | 46.6933646 |
| 10       |            |             | 50.92165307 | 58.35722693 | 51.3564902 |
| 12       |            |             | 34.02671254 | 52.35069913 | 55.4553006 |

### WEE1-i-1 induced

| Time (h) | CDKA 1   | CDKA 2   | CDKA 3   |
|----------|----------|----------|----------|
| 1        | 100      | 100      | 100      |
| 3        | 60.24659 | 94.36019 | 53.43055 |
| 4        | 64.81426 | 95.01102 | 63.25103 |
| 5        | 58.03517 | 97.53319 | 49.73508 |
| 6        | 45.45365 | 96.99092 | 56.70823 |
| 7        | 33.3446  | 95.98753 | 47.4936  |
| 9        | 42.85049 | 94.81163 | 38.94798 |

| Time (h) | CDKB1 1  | CDKB1 2  | CDKB1 3  |
|----------|----------|----------|----------|
| 1        | 100      | 100      | 100      |
| 3        | 58.78956 | 63.15473 | 51.95804 |
| 4        | 68.41233 | 49.33648 | 34.51408 |
| 5        | 56.42295 | 60.84588 | 47.06691 |
| 6        | 54.99347 | 64.55488 | 44.09816 |
| 7        | 51.89385 | 53.14281 | 41.26649 |
| 9        | 73.26231 | 64.16574 | 48.82118 |

## Appendix VI: *Arabidopsis* roots measurements raw data

### Wild Type

| Day                                      | 15  | 17   | 19  |                         | 15 | 17 | 19 |
|--|-----|------|-----|-------------------------|----|----|----|
|  |     |      |     | Number of primordia     |    |    |    |
| Primary root length (cm)                 |     |      |     |                         |    |    |    |
|  | 2.4 | 2.6  | 3.2 |                         | 2  | 1  | 2  |
|  | 2.2 | 2.8  | 5.7 |                         | 2  | 2  | 4  |
|  | 2.4 | 1.25 | 4.5 |                         | 3  | 3  | 4  |
|  | 5.2 | 2.9  | 3.5 |                         | 4  | 3  | 3  |
|  | 3.5 | 3.5  | 5.8 |                         | 3  | 3  | 3  |
|  | 2.8 | 3    | 6   |                         | 6  | 4  | 2  |
|  | 3   | 3.6  | 4.5 |                         | 2  | 3  | 4  |
|  | 3.1 | 4.7  | 3   |                         | 3  | 3  | 5  |
|  | 4.2 | 2.1  | 8   |                         | 3  | 2  | 2  |
|  | 2.2 | 5    | 2   |                         | 2  | 3  | 1  |
|  | 3.1 | 4    | 2.8 |                         | 4  | 3  | 4  |
|  | 2.3 | 5.4  | 3   |                         | 1  | 4  | 4  |
|  | 2.7 | 3    | 4.1 |                         | 2  | 3  | 3  |
|  | 2.3 | 6    | 4.6 |                         | 3  | 3  | 3  |
|  | 3.1 | 2.8  | 6   |                         | 1  | 4  | 2  |
| Last primordia to root tip distance (cm) |     |      |     | Number of lateral roots |    |    |    |
|  | 0.5 | 0.2  | 0.7 |                         | 11 | 10 | 16 |
|  | 0.3 | 0.5  | 1.2 |                         | 8  | 18 | 21 |
|  | 0.5 | 0.6  | 1.2 |                         | 10 | 7  | 15 |
|  | 0.6 | 0.6  | 0.4 |                         | 19 | 11 | 15 |
|  | 0.3 | 0.7  | 1   |                         | 11 | 14 | 17 |
|  | 0.5 | 0.6  | 1.1 |                         | 14 | 11 | 16 |
|  | 0.2 | 0.5  | 1.4 |                         | 12 | 12 | 18 |
|  | 0.3 | 0.4  | 0.6 |                         | 17 | 6  | 12 |
|  | 0.4 | 0.7  | 0.9 |                         | 8  | 14 | 9  |
|  | 0.4 | 0.6  | 1   |                         | 5  | 17 | 14 |
|  | 0.5 | 0.6  | 0.9 |                         | 16 | 11 | 19 |
|  | 0.3 | 0.5  | 0.4 |                         | 11 | 15 | 16 |
|  | 0.6 | 0.3  | 1.2 |                         | 12 | 9  | 15 |
|  | 0.5 | 0.7  | 1.3 |                         | 12 | 13 | 17 |
|  | 0.2 | 0.6  | 0.3 |                         | 15 | 14 | 18 |



| Day       | 15        |           |           | 17        |           |           | 19        |           |  |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|--|
| Primordia | Primordia | Primordia | Primordia | Primordia | Primordia | Primordia | Primordia | Primordia |  |
| radius    | height    | volume    | radius    | height    | volume    | radius    | height    | volume    |  |
| 41.75     | 27.25     | 49715     | 55.22     | 82.47     | 263207    | 60.67     | 56.56     | 217904    |  |
| 42.62     | 40.21     | 76448.6   | 50.14     | 62.72     | 165038    | 60.6      | 56.04     | 215403    |  |
| 44.77     | 47.89     | 100468    | 48.27     | 53.19     | 129716    | 59.36     | 47.59     | 175514    |  |
| 50.66     | 42.71     | 114728    | 52.57     | 65.31     | 188914    | 51.19     | 51.53     | 141331    |  |
| 47.82     | 54.68     | 130875    | 41.5      | 46.59     | 83984.1   | 45.85     | 44.47     | 97848.5   |  |
| 30.89     | 35.66     | 35614.4   | 40.71     | 51.69     | 89663.8   | 37.69     | 46.13     | 68587.4   |  |
| 30.5      | 37.21     | 36230     | 43.76     | 68.6      | 137495    | 68.19     | 21.57     | 104978    |  |
| 36.74     | 47.36     | 66911.1   | 42.37     | 36.79     | 69128.2   | 57.96     | 59.28     | 208436    |  |
| 52        | 42.96     | 121585    | 54.11     | 51.9      | 159049    | 45.32     | 45.37     | 97534.2   |  |
| 50.73     | 81.04     | 218292    | 43.11     | 55.55     | 108056    | 53.48     | 63.75     | 190841    |  |
| 42.61     | 45.7      | 86845.6   | 47.11     | 37.34     | 86737.9   | 58.79     | 66.45     | 240387    |  |
| 35        | 32.56     | 41747.3   | 49.14     | 22.43     | 56690.2   | 55.36     | 86.25     | 276668    |  |
| 39.78     | 45.89     | 76007.4   | 49.58     | 44.49     | 114468    | 56.19     | 95.84     | 316718    |  |
| 40.3      | 54        | 91793.6   | 51.22     | 46.78     | 128454    | 58.23     | 110       | 390422    |  |
| 35.72     | 48.34     | 64556.2   | 50        | 57.8      | 151243    | 49        | 99.81     | 250827    |  |
| 33.25     | 73        | 84472.3   | 38.98     | 80        | 127228    | 61.23     | 73.32     | 287713    |  |
| 42.12     | 54.32     | 100866    | 50        | 72.84     | 190598    | 53.98     | 53.98     | 164629    |  |
| 49        | 55        | 138218    | 42.7      | 58.43     | 111506    | 34.56     | 68.78     | 85984.1   |  |
| 35.67     | 62.54     | 83286.1   | 41.2      | 48.34     | 85883.4   | 48.32     | 70.32     | 171847    |  |
| 38.3      | 38        | 58343.1   | 39.21     | 65.93     | 106093    | 54.9      | 37        | 116723    |  |

**Plants transformed with BIN HYG TX-*AtWEE1***

| Day                                      | 15  | 17  | 19                      | 15                  | 17 | 19 |
|--|-----|-----|-------------------------|---------------------|----|----|
| Primary root length (cm)                 |     |     |                         | Number of primordia |    |    |
| 0.8                                      | 0.6 | 0.7 | 3                       | 1                   | 2  |    |
| 2.8                                      | 0.8 | 1.1 | 2                       | 3                   | 2  |    |
| 1.4                                      | 0.9 | 0.6 | 1                       | 2                   | 2  |    |
| 0.7                                      | 1   | 0.9 | 1                       | 2                   | 1  |    |
| 0.6                                      | 1.4 | 1.2 | 1                       | 2                   |    |    |
| 1.1                                      | 1.7 | 1   | 2                       | 2                   | 1  |    |
| 0.9                                      | 1.3 | 1.3 | 2                       | 1                   | 1  |    |
| 0.7                                      | 1.1 | 0.9 | 2                       | 3                   | 1  |    |
| 0.4                                      | 0.9 | 1.7 | 3                       | 1                   | 3  |    |
| 1.2                                      | 0.6 | 1.9 | 1                       | 1                   | 2  |    |
| 0.4                                      | 1.2 | 2   | 1                       | 2                   | 2  |    |
| 0.4                                      | 1   | 0.9 | 2                       | 2                   | 3  |    |
| 0.8                                      | 1.3 | 1   | 3                       | 3                   | 2  |    |
| 1  | 0.9 | 5   | 2                       | 2                   | 3  |    |
| 1  | 1.5 | 1.8 | 1                       | 1                   | 1  |    |
| Last primordia to root tip distance (cm) |     |     | Number of lateral roots |                     |    |    |
| 0.1                                      | 0.2 | 0.1 | 5                       | 8                   | 9  |    |
| 0.3                                      | 0.3 | 0.1 | 12                      | 6                   | 10 |    |
| 0.7                                      | 0.5 | 0.1 | 3                       | 7                   | 7  |    |
| 0.2                                      | 0.5 | 0.2 | 8                       | 14                  | 6  |    |
| 0.6                                      | 0.1 | 0.3 | 5                       | 7                   | 8  |    |
| 0.1                                      | 0.2 | 0.1 | 5                       | 5                   | 6  |    |
| 0.7                                      | 0.3 | 0.2 | 6                       | 7                   | 6  |    |
| 0.2                                      | 0.1 | 0.2 | 4                       | 7                   | 9  |    |
| 0.1                                      | 0.3 | 0.2 | 7                       | 11                  | 14 |    |
| 0.3                                      | 0.2 | 0.1 | 3                       | 4                   | 11 |    |
| 0.5                                      | 0.2 | 0.3 | 8                       | 7                   | 4  |    |
| 0.3                                      | 0.3 | 0.1 | 8                       | 3                   | 7  |    |
| 0.2                                      | 0.1 | 0.3 | 4                       | 5                   | 8  |    |
| 0.2                                      | 0.4 | 0.3 | 4                       | 8                   | 3  |    |

| Day       | 15        |           |           | 17        |           |           | 19        |           |  |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|--|
| Primordia | Primordia | Primordia | Primordia | Primordia | Primordia | Primordia | Primordia | Primordia |  |
| radius    | height    | volume    | radius    | height    | volume    | radius    | height    | volume    |  |
| 39.98     | 41.54     | 69496.1   | 50.05     | 44.28     | 116098    | 64.52     | 74.93     | 326477    |  |
| 47.44     | 60.21     | 141829    | 37.62     | 52.74     | 78124.3   | 56.17     | 45.71     | 150948    |  |
| 36.71     | 55        | 77578.2   | 43.51     | 50.04     | 99152.5   | 45.29     | 41.81     | 89762.1   |  |
| 41.09     | 52.42     | 92635.6   | 40.89     | 61.31     | 107294    | 42.88     | 46.71     | 89893.4   |  |
| 30.69     | 46.47     | 45811.5   | 30.61     | 44.19     | 43337     | 68.18     | 75.48     | 367244    |  |
| 39.94     | 55.29     | 92314.8   | 35.09     | 54.2      | 69851.3   | 41.62     | 55.57     | 100752    |  |
| 48.8      | 55.15     | 137465    | 41.81     | 42.48     | 77723.7   | 44.41     | 63.65     | 131392    |  |
| 45.37     | 62.99     | 135712    | 33.65     | 49.97     | 59222.7   | 52.69     | 46.91     | 136311    |  |
| 38.3      | 49.17     | 75492.9   | 33.37     | 62.75     | 73136.6   | 51.02     | 79.51     | 216626    |  |
| 44.53     | 47.84     | 99289.9   | 63.03     | 47.15     | 196058    | 67.89     | 44.77     | 215977    |  |
| 33.33     | 55.39     | 64403.6   | 59.77     | 55.04     | 205804    | 79.62     | 30.77     | 204165    |  |
| 29.87     | 42.98     | 40137     | 45.63     | 25.95     | 56551.8   | 44.79     | 50.91     | 106899    |  |
| 24.39     | 35.42     | 22053.7   | 34.43     | 43.01     | 53364.4   | 54.54     | 37.82     | 117750    |  |
| 42.1      | 45.98     | 85298.5   | 61.29     | 98.61     | 387711    | 63.5      | 52.57     | 221868    |  |
| 35.34     | 55.87     | 73033.2   | 53.66     | 33.06     | 99635.2   | 55.94     | 51.39     | 168319    |  |
| 41        | 50.32     | 88535.4   | 55.71     | 49.31     | 160181    | 53.14     | 73.32     | 216708    |  |
| 33.65     | 47.92     | 56793.1   | 49.32     | 55.34     | 140894    | 62.11     | 88.99     | 359313    |  |
| 40.35     | 69.21     | 117941    | 55.55     | 68.31     | 220628    | 47.09     | 98.43     | 228451    |  |
| 30.85     | 48.29     | 48103.4   | 39.43     | 51.98     | 84585.9   | 44.32     | 90        | 185034    |  |
| 32.1      | 67.3      | 72582.8   | 37.78     | 50        | 74696.9   | 48.55     | 45.78     | 112944    |  |

**Plants transformed with pER8-*AtWEE1***


| Day                      | 15  | 17  | 19  |
|--------------------------|-----|-----|-----|
| Primary root length (cm) |     |     |     |
|                          | 1.2 | 1.5 | 2.8 |
|                          | 2   | 2.4 | 3   |
|                          | 1.6 | 2.2 | 3.6 |
|                          | 0.8 | 1.4 | 4   |
|                          | 1.4 | 1.9 | 3.5 |
|                          | 2.8 | 3.2 | 2.6 |
|                          | 2.4 | 3.6 | 2.9 |
|                          | 1.9 | 2.6 | 3.5 |
|                          | 0.8 | 1.4 | 3.9 |
|                          | 2.3 | 1.8 | 2.6 |
|                          | 1.4 | 2.4 | 4   |
|                          | 1.6 | 3   | 4.2 |
|                          | 2.5 | 2.2 | 2.3 |
|                          | 3   | 3.2 | 2.5 |
|                          | 2   | 2.4 | 2.8 |
|                          | 2.7 | 2.5 | 3.5 |
|                          | 1.4 | 1.9 | 3.8 |
|                          | 2.4 | 2.1 | 3.7 |
|                          | 1.1 | 2.4 | 3.1 |
|                          | 1.3 | 3.2 | 2.9 |
|                          | 2.4 | 1.9 | 2.8 |

**Appendix VII: The yeast two hybrid technique was employed to investigate possible interaction between AtWEE1 and proteins that regulate its activity**

**VII.I. Introduction**


One technique used to study protein-protein interactions is the "yeast two hybrid" system. The first two-hybrid system was described by Fields and Song in 1989. It is based on the fact that proteins are composed of modules, or domains, which are units within the same polypeptide (protein) chain. The presence of these individual domains allows the same protein to perform different functions. The yeast two-hybrid technique uses GAL4 protein which is a transcriptional activator with two separate domains: 1) DNA-binding domain (BD) that is capable of binding to DNA, and 2) activation domain (AD) that is capable of activating transcription of the DNA. Both of these domains came from the same gene *GAL4* of *Saccharomyces cerevisiae* and are required for transcription. Transcriptional activators (TA), like the GAL4 protein, use their binding domain to interact (bind) to the promoter, a region situated upstream from the gene (coding region of the DNA) (Fig.VII.I). Once the TA has bound to the promoter, it is then able to activate transcription via its activation domain, which interacts with RNA polymerase. Hence, the activity of a TA requires both a DNA binding domain and an

activation domain. If either of these domains is absent, then transcription of the gene will fail.



**Fig. VII.I:** Normal transcription requires both the DNA-binding domain (BD) and the activation domain (AD) of a transcriptional activator (TA).

The binding domain and the activating domain do not necessarily have to be on the same protein. In fact, a protein with a DNA binding domain can activate transcription when bound to another protein containing an activation domain. This principle is used in the yeast two-hybrid technique, where the *GAL4* gene is divided in two parts: 1) the DNA binding domain fused to the protein of interest (X), known as the bait and 2) the activation domain which is fused to a potential binding partner (Y). If protein X interacts with protein Y, the binding of these two will form an intact and functional transcriptional activator. This newly formed transcriptional activator will then go on to transcribe a reporter gene, which is a gene whose protein product can be easily detected or can guarantee the survival of yeast cells growing on a medium lacking one or more amino acids (Fig. VII.II).



**Fig. VII.II:** The yeast two-hybrid technique measures protein-protein interactions by detecting properties of a reporter gene. If protein X and protein Y interact, then their DNA-binding domain and activating domain will combine to form a functional transcriptional activator (TA). The TA will then proceed to transcribe the reporter gene that is downstream of its promoter.

For the two-hybrid analysis of protein interaction used in this work, the bait protein AtWEE1 was fused to the GAL4 binding domain (GAL4 BD), while the potential binding partner was fused to the GAL4 activation domain (GAL4 AD). For a screen to find unknown interacting proteins, the GAL4 AD was fused to an *Arabidopsis* seedling root cDNA library. The two plasmids carrying these fusion proteins also contain selection genes, or genes encoding proteins that are required for cell survival on a particular medium. Yeast two-hybrid assays typically use selection genes encoding proteins capable of synthesizing amino acids such as histidine, leucine and tryptophan.

In the original system, only one reporter gene (*LacZ*) was used (Fields and Song, 1989). However, as the technology developed, yeast strains containing a number of reporter genes were generated, such as *HIS3* in the case of the GAL4 system, and used as a reporter gene in conjunction with *LacZ* to provide a more stringent assay for protein-protein interactions.

The yeast two-hybrid system has been used extensively in plant science research to analyze known interactions and to isolate new interacting partners. It has been used to study protein-protein interactions involved in a number of diverse processes such as flower development (Davies et al. 1996), the self incompatibility mechanism (Mazzurco et al. 2001), the circadian clock (Jarillo et al. 2001), and phytohormone signaling (Oullet et al. 2001).

In non-plant systems the yeast two-hybrid screen has been used successfully to find proteins interacting with WEE1. In mouse and human the yeast two-hybrid system was employed to isolate proteins that may interact with WEE1. Interestingly, in both mouse and human the results obtained through the yeast two-hybrid technique showed interaction between WEE1 and a member of the 14-3-3 family. In the case of mouse, WEE1 interacted with 14-3-3  $\zeta$  (Honda *et al.*, 1997), whereas in the case of the human gene, WEE1 interacted with 14-3-3  $\beta$  (Wang *et al.*, 2000).

The aims of the work reported in this chapter were to examine whether AtWEE1 protein interacts with members of the 14-3-3 family and other proteins. The study of proteins interacting with AtWEE1 could lead to the discovery of protein regulators of AtWEE1 kinase activity.




## VII.II. Materials and Methods

### VII.II.I. pBD-GAL4 Cam and pAD-GAL4-2.1 plasmids

In the yeast two-hybrid system, used in this work two different plasmids were constructed:

1. pBD-GAL4 Cam (Fig. VII.III a): encoding the protein of interest fused in frame with the DNA-binding domain of the transcription factor GAL4
2. pAD-GAL4-2.1 excised from the HybridZap-2.1 vector (Fig. VII.III.b): encoding the potential binding protein fused in frame with the GAL4 transcription activation domain.



**Fig. VII.III:** **a)** DNA encoding the bait protein is inserted into the region of insertion (MCS) of pBD-GAL4 Cam vector close to a segment of GAL4 BD; **b)** The HybriZap-2.1 contains lambda genes *A* through *J* in the left arm and *att*, *int*, *xis* and *cl857* in the right arm. The *fl* initiator (I) and terminator (T) allow efficient in vivo excision of the pAD-GAL4-2.1 phagemid vector from the HybriZap-2.1 vector. DNA inserts are ligated into the HybriZap-2.1 vector to generate the primary lambda library. This primary lambda library is amplified and converted by in vivo mass excision to a pAD-GAL4-2.1 vector library. Both pBD-GAL4 Cam and pAD-GAL4-2.1 vectors contain the pUC origin for replication in *E.coli*, an *fl* origin for production of single stranded DNA in *E.coli*, and 2 micron origin for replication in yeast. The pBD-GAL4 Cam vector contains the chloramphenicol resistance gene and promoter for selection with chloramphenicol in *E.coli*. The pAD-GAL4-2.1 vector contains the ampicillin resistance  $\beta$ -lactamase gene for selection with ampicillin in *E.coli*. For selection in yeast the pAD-GAL4-2.1 vector contains the *LEU2* gene and the pBD-GAL4 Cam vector contains the *TRP1* gene. Hybrid proteins are expressed in yeast from the ADH1 promoter and terminated by the ADH1 terminator. (Extracted from <http://www.stratagene.com/products/showProduct.aspx?pid=256>).

### VII.II.II. Mass library in vivo excision

RNA isolated from *Arabidopsis* seedling root was used, by Stratagene, to construct a yeast two hybrid cDNA library into pAD-GAL4-2.1 phagemid vector within the HybriZAP-2.1 lambda vector. To excise the pAD-GAL4-2.1 phagemid vector for use in yeast transformation, a single colony of both *E. coli* strain XL1-Blue MRF' and strain XL0LR were transferred from a fresh bacterial streak plate containing LB (2% agar)-medium (per litre: 10 g bacto-tryptone; 5 g bacto-yeast extract; 10 g NaCl; pH=7) plus tetracycline (15  $\mu$ g/ml) medium) to the appropriate liquid medium, 10 ml NZY-medium (per litre: 5 g bacto-yeast extract; 5 g NaCl; pH=7.5) supplemented with 0.2% maltose and 10 mM MgSO<sub>4</sub> for XL1-Blue MRF' and LB-medium for XL0LR. The two strains were left to grow over-night in 50 ml Falcon tubes with vigorous shaking at 30°C. Cells were then centrifuged at 3500 rpm (MSE Centaur 2) for 10 minutes and the pellet was resuspended in 10 mM MgSO<sub>4</sub> at an OD<sub>600</sub> of 1.0. The cell suspension (0.25 ml) was

added to 50 ml of NZY broth supplemented with 0.2% (w/v) maltose and 10 mM MgSO<sub>4</sub> in a 250 ml flask. Cell cultures were incubated at 37°C with shaking until an OD<sub>600</sub> of 0.3-0.4 was reached and then centrifuged at 3500 rpm (MSE Centaur 2) for 10 minutes. XL1-Blue MRF<sup>r</sup> cells were resuspended in 10 mM MgSO<sub>4</sub> (4 ml) to an OD<sub>600</sub> of 5.0 (assuming an OD<sub>600</sub> of 5.0 is equal to a final cell concentration of 4x10<sup>9</sup> cells). XL0LR cells were resuspended in 10 mM MgSO<sub>4</sub> (10 ml) to an OD<sub>600</sub> of 1.0.

To mass excise an entire library, a portion of the amplified library stock was combined with the XL1-Blue MRF<sup>r</sup> cells at a multiplicity of infection (MOI) of 1:10 lambda phage to cell ratio in a 250 ml flask. ExAssist helper phage (Stratagene) at a 10:1 helper phage to cells ratio was added to ensure that every cell was coinfecting with lambda phage and helper phage. The cells were incubated for 15 minutes at 37°C before adding 20 ml of LB broth and the culture was incubated with gentle agitation for a further 3 hours at 37°C. Cultures were incubated for 20 minutes at 70°C to lyse the lambda phage particles and the XL1-Blue MRF<sup>r</sup> cells. The cells were then centrifuged at 3500 rpm (MSE Centaur 2) for 10 minutes and the supernatant was transferred to a fresh tube. The titer of Amp<sup>r</sup> colonies per milliliter (or colony forming units per milliliter (cfu/ml)) was determined. Each Amp<sup>r</sup> colony represents a single excised phagemid. The supernatant (containing the excised phagemid particles) was diluted in different volumes of TE buffer (10 mM Tris-Cl, 1 mM EDTA, pH=7.5) and mixed with 200 µl of XL0LR cells (OD<sub>600</sub> of 1.0). The cells were then incubated at 37°C for 15 minutes and spread onto an LB agar plate containing 50 µg/ml of ampicillin. The plate was incubated over-night at 37°C. The titer of excised phagemid (in cfu/ml) was determined as follows:

$$\frac{\text{Number of colonies (cfu)} \times \text{dilution factor} \times 1000 \mu\text{l/ml}}{\text{Volume of phagemid plated } (\mu\text{l})}$$

### **VII.II.III. Amplification of the excised phagemid library**

A culture of XLOLR cells was transferred from a fresh bacterial streak plate (LB-tetracycline medium) to LB medium and incubated over-night with vigorous shaking at 30°C in a 50 ml Falcon tube. Cells were then centrifuged at 3500 rpm (MSE Centaur 2) for 10 minutes; the pellet was resuspended in 10 mM MgSO<sub>4</sub> at an OD<sub>600</sub> of 1.0.

To amplify the mass excised library, XLOLR cells were combined with a portion of the excision supernatant in a 2 l flask at minimum cells to phagemid ratio of 10:1, assuming an OD<sub>600</sub> of 1.0 equals a cell concentration of 8x10<sup>8</sup> cells/ml. The phagemid and the cells were incubated at 37°C for 15 minutes, and then 500 ml of LB broth containing 50 µg/ml of ampicillin was added. The cells were incubated with shaking at 37°C for 3 hours to an OD<sub>600</sub> of 0.4. The cells were then centrifuged at 3500 rpm (MSE Centaur 2) for 10 minutes.

### **VI.II.IV. Isolation of DNA plasmid library**

The plasmid DNA was extracted from XLOLR pelleted cells using a QIAfilter Plasmid Maxi Protocol (QIAGEN). The bacterial cells were harvested by centrifugation at 3500 rpm (MSE Centaur 2) for 10 minutes. The kit is designed for preparation of up to 500 µg of high- or low-copy plasmid DNA. The kit provides lysis buffers to disrupt the bacterial cell wall and membrane and ribonuclease (RNase A) to remove the RNA

molecules. Cell debris and proteins are separated from the plasmid DNA by centrifugation. The supernatant containing the plasmid DNA is applied to the QIAGEN-tip 500 column. This column contains a resin able to bind the plasmid DNA, but not other elements that contaminate the DNA, which are washed through. Elution DNA buffer is then applied to the column to remove the plasmid DNA from the resin. The DNA is then precipitated by adding isopropanol to the eluted DNA and centrifuged. The pellet was washed with 70% ethanol and centrifuged at 15000 g for 10 min (Beckman Coulter J-E centrifuge, rotor JA-14). The ethanol was then removed and the pellet was air-dried for 30 min, and redissolved in 300 µl distilled water.

#### **VII.II.V. Insert preparation and ligation**

The bait used in the two hybrid screen was constructed by PCR amplifying the ORF of the *Arabidopsis thaliana WEE1 (AtWEE1)*, using forward primer (5'-CATGGAGAATTCATGTTCGAGAAGAACG-3') and reverse primer (5'-TAAGCGGTCTGACTCAACCTCGAATCCTATATC-3'). One µl of cDNA was amplified with 0.625 U of Qiagen Taq polymerase, Qiagen buffer, 1.5 mM MgCl<sub>2</sub>, 0.2 mM dNTPs and 1 µg of each of the primers. Thermocycle conditions were: 35 cycles of 95°C (1 min), 55°C (1 min) and 72°C (1 min) in a PTC100 thermocycler (see chapter 2 sections 2.4.1, 2.4.2 and 2.4.3 for details about RNA extraction, DNase treatment of cDNA and cDNA synthesis respectively; also see section 2.4.4 for details about cDNA amplification, except that a hot start step of 15 min at 95°C was used with this Taq polymerase). The primers introduced *EcoRI* (5'-

CATGGAGAATTCATGTTTCGAGAAGAACG-3') and *SalI* (5'-TAAGCGGTCGACTCAACCTCGAATCCTATATC-3') restriction sites (see chapter 2 section 2.1.1 for more details about restriction and ligation reactions). The product was then cut with *EcoRI* and *SalI* restriction enzymes, purified using the "QIAquick Purification Kit Protocol" (QIAGEN), and cloned into the *EcoRI* and *SalI* sites of Gal4 BD (binding domain) plasmid vector pBD-GAL4Cam (Stratagene) giving pBD-GAL4Cam-*AtWEE1*.

#### VII.II.VI. Transformation into yeast YRG-2 cells

To transform YRG-2 cells with pBD-GAL4Cam-*AtWEE1*, 50 ml of YPD broth (per litre: 10 g yeast extract; 20 g bacteriological peptone; 20 g glucose; 0.1 g adenine; 0.1 g uracil) were inoculated with YRG-2 cells harvested from a plate (YPD-medium) using a flamed loop. The cells were incubated at 200 rpm over-night at 30°C to a cell density of  $1-5 \times 10^7 \text{ ml}^{-1}$  (OD<sub>600</sub> 1-2). Cells were then harvested by centrifugation at 8000 rpm (Beckman Coulter J-E centrifuge, rotor JA-14) for 1 second; the pellet was resuspended in 20 ml TE buffer and spun down at 8000 rpm for 1 second. This was repeated and the pellet was then resuspended in 10 ml LiAc (0.1 M) and incubated at 200 rpm for 1 hour at 30°C. Competent YRG-2 cells (150 µl) were transferred into two Eppendorf tubes and 5 µl of pBD-GAL4Cam-*AtWEE1* (1 or 2 µg) were added to each tube. As a control 5 µl TE buffer were also added to a different Eppendorf tube containing YRG-2 competent cells. PEG4000 (350µl) was added to each tube of competent cells. The transformation mixtures were then incubated at 30°C for 1 hour

without shaking. The cells were heated at 42°C for 5 minutes and quickly cooled down on ice for 3 minutes. The transformed cells (200 µl) were then plated on minimal medium lacking tryptophan and incubated at 30°C for 3-4 days. Single colonies were then patched onto minimal medium lacking histidine and containing 0, 1, 5, 10, 25, 30, 35, 40, 45 and 50 mM 3-AT (3-amino-1, 2, 4-triazole, Sigma). 3-AT is an inhibitor of the *HIS3* gene (*HIS3*-encoded IGP-dehydratase, an enzyme required for histidine biosynthesis) product and reduces the background due to basal *HIS3* expression.

#### **VII.II.VII. Double transformation into YRG-2 cells**

For the two hybrid screen the pAD-GAL4-2.1 library was transformed into YRG-2 cells containing pBD-GAL4Cam-*AtWEE1*. YRG-2 cells carrying pBD-GAL4Cam-*AtWEE1* were inoculated in 10 ml minimal medium lacking tryptophan and incubated at 30°C on a shaker at 200 rpm over-night. The titre of the over-night culture was determined by measuring the OD<sub>600</sub> of a 1 in 10 dilution into water, considering an OD<sub>600</sub> of 0.1 to correspond to approximately 1x10<sup>6</sup> cells/ml. The volume of the over-night culture containing 2.5x10<sup>8</sup> cells was then calculated and added to prewarmed (30°C) YPD medium, to give a final volume of 50 ml in a 250 ml flask. The diluted culture was then incubated at 30°C on a shaker at 200 rpm until the cells reached approximately a concentration 2x10<sup>7</sup> cells/ml. The cells were harvested in a sterile 50 ml disposable centrifuge tube (Falcon) by centrifugation at 3500 rpm (MSE Centaur 2) for 5 minutes. Cell pellets were resuspended in 25 ml of sterile water to wash the cells before pelleting them again by centrifugation as described above. The pellet was then resuspended in 900

μl of sterile water and transferred to a 1.5 ml microcentrifuge tube. Cells were centrifuged again at 13000 rpm (MiniSpin, Eppendorf) for 1 minute; the pellet was resuspended in 100 mM LiAc to a final volume of 1 ml and incubated at 30°C for 10 minutes. For each 1x transformation reaction aliquots of 100 μl LiAc cell suspension were transferred into a new 1.5 ml microcentrifuge tube. The cells were spun down at top speed (MiniSpin, Eppendorf) for 1 minute and the following reagents were added: 50% PEG (240 μl); 1 mM LiAc (36 μl); ss-DNA 2 mg/ml (50 μl); plasmid DNA from the mass excision and isolation of the plasmid library 1, 2 and 5 μg, sterile water to a final volume of 360 μl. The transformation mixtures were incubated at 30°C for 30 minutes, heat shocked at 42°C for 30 minutes and spun down at top speed (MiniSpin, Eppendorf) for 1 minute. Sterile water (400 μl) was added to resuspend the cell pellet and cells were plated onto minimal medium (per litre: 1.62 g yeast nitrogen base; 5 g ammonium sulfate; 20 g glucose; 20 g agar) lacking tryptophan and leucine (0.2 ml of appropriate “drop-out” solution were spread onto each Petri dishes immediately before use). Petri dishes containing the transformed cells were incubated at 30°C until colonies appeared.

### **VII.III. Results**

#### **VII.III.I. Preparation of yeast two-hybrid construct**


The yeast two hybrid screen was performed to identify *Arabidopsis thaliana* proteins that bind *Arabidopsis thaliana* WEE1 kinase (AtWEE1). The *AtWEE1* open reading frame was amplified by PCR from the BIN HYG TX-*AtWEE1* plasmid and



ligated in frame with the binding domain of the pBD-GAL4 Cam plasmid. The construct was sequenced to confirm that no PCR errors had been generated and that the test protein was in frame with its fusion partner. AtWEE1 was then used as a bait to screen an *Arabidopsis thaliana* seedling root cDNA library in the GAL4 transcriptional activation domain vector pAD-GAL4-2.1.

### VII.III.II. YRG-2 cells transformed with pBD-GAL4 Cam-*AtWEE1* plasmid

The Stratagene-2-hybrid yeast strain YRG-2 was transformed with *AtWEE1* in the pBD-GAL4 Cam vector. Following transformations, positive colonies growing on selective medium (minimal medium lacking tryptophan) were tested by PCR (Fig. VII.IV) to confirm *AtWEE1* presence inside the cells.



**Fig. VII.IV:** Independent colonies (from 1 to 9) of YRG-2 cells transformed with pBD-GAL4 Cam-*AtWEE1* were amplified by PCR using primers *AtWEE1* (forward and reverse) to confirm *AtWEE1* presence. As a positive control ((+)ve) BIN HYG TX-*AtWEE1* plasmid was used, whereas sterile distilled water was used as the negative control ((-)ve). Product size is 229 bp.

### VII.III.III. Testing the pBD-GAL4Cam-*AtWEE1* plasmid for auto-activation


Auto-activation of reporter gene expression can be a problem with some proteins fused to the DNA-binding domain. To test that the *AtWEE1* bait protein itself was unable

to activate the transcription of the *HIS3* and *lacZ* reporter genes, the YRG-2 cells transformed with pBD-GAL4 Cam-*AtWEE1* were plated onto minimal medium lacking histidine with different concentrations of 3-AT. Yeast cells containing the AtWEE1 bait protein were able to grow on minimal medium lacking histidine up to a concentration of 35 mM 3-AT, but were unable to grow on minimal medium lacking histidine with 40 mM 3-AT.

These results show that the AtWEE1 bait protein could auto-activate transcription of the *HIS3* reporter gene, but the growth of yeast cells on histidine free minimal medium could be blocked by the addition of 40 mM 3-AT. The two-hybrid screen could therefore be performed on minimal medium lacking leucine, tryptophan and histidine with 40 mM 3-AT. Leucine and tryptophan were used to select yeast cells containing the two plasmids. The *lacZ* filter-lift assay was also performed showing that yeast cells containing pBD-GAL4 Cam-*AtWEE1* plasmid did not have *lacZ* activity. This suggests that the AtWEE1 bait protein did not auto-activate the *lacZ* reporter gene.

#### **VII.III.IV. Transformation of YRG-2 cells containing AtWEE1 bait protein with pAD-GAL4-2.1 cDNA library**


To transform YRG-2 cells with pAD-GAL4-2.1 cDNA library, yeast cells already transformed with *AtWEE1* were used in the exponential phase of growth. Cell growth was thus measured every 2 hours for a 24 hour period (Fig. VII.V).



**Fig. VII.V:** Growth of YRG-2 cells transformed with pBD-GAL4 Cam-*AtWEE1*. YRG-2 cells containing the bait protein grow exponentially for 12 hours.

Aproximately  $1 \times 10^7$  YRG-2 cells transformed with both pBD-GAL4 Cam-*AtWEE1* and pAD-GAL4-2.1 cDNA library were selected on minimal medium lacking tryptophan, leucine and histidine with 40 mM 3-AT. After 4 days of incubation, the primary two-hybrid screening plates were covered with a background layer of yeast cell growth. However, after further incubation, larger distinct colonies appeared to grow out from the background. These fast-growing colonies are putatively positive for a protein-protein interaction. Only 700 colonies larger than the overall background of colonies became apparent out of  $1 \times 10^7$  transformants screened on the selective medium (minimal medium lacking histidine, leucine and tryptophan with 40mM 3-AT) (Fig. VII.VI a). The 700 colonies were harvested and re-selected onto a fresh selective medium for the selection of


protein-protein interaction (Fig. VII.IV b, c). Of the original 700 colonies only 442 grew on the fresh selective medium when retested.



**Fig. VII.VI:** Example of the method used to screen  $1 \times 10^7$  transformants: **a)** large Petri dishes (13.5 cm in diameter) containing minimal medium lacking tryptophan, leucine and histidine with 40mM 3AT, were used to screen YRG-2 cells carrying both the bait protein and the activating domain. Large distinct colonies, which appeared to grow out from the background, were harvested and transferred to **b)** small Petri dishes (8.5 cm in diameter) containing the same selective medium. Positive colonies growing on the fresh selective medium were then transferred to **c)** new small Petri dishes containing the same selective minimal medium. These colonies were then used for the LacZ assay.

The presence of the pAD-GAL4-2.1 cDNA library was confirmed by PCR on randomly chosen colonies (Fig. VII.VII).

## VII.VI. Results



**Fig. VII.VII:** Colonies of YRG-2 cells (from 1 to 8), containing both plasmid pBD-GAL4 Cam-*AtWEE1* and pAD-GAL4-2.1 cDNA library, were randomly chosen to check presence of the activating domain in the cell. The different band size reflects the different inserts contained in the pAD-GAL4-2.1 plasmid. Some of the colonies appeared to give a negative result which may be due to a low quantity of YRG-2 cells in the PCR reaction. Distilled water was used as negative control ((-)ve).

The 442 colonies selected were replica-plated onto minimal medium lacking tryptophan, leucine and histidine with 40mM 3AT for the second library screen which was performed using the lacZ assay to reduce the number of false positives. Surprisingly, not one of these colonies gave a positive blue color signal. YRG-2 cells containing Spcdc25 as bait co-transformed with 14-3-3  $\kappa$ ,  $\lambda$  and  $\omega$  respectively were chosen as positive controls in the LacZ assay. As previously demonstrated by Sorrell and colleagues using the two-hybrid screen, Spcdc25 protein can interact with three *Arabidopsis* 14-3-3 proteins: GF14  $\kappa$ ,  $\lambda$ , and  $\omega$  (Sorrell *et al.*, 2003). The LacZ assay was repeated three times on the 442 colonies and the control, each time making sure that: 1) the LacZ buffer with X-gal were freshly prepared, 2) the colonies transferred on fresh plates were between 4-5 days old at the time of screening, 3) the colonies were transferred from the plate to the filter paper by visual inspection of both the filter paper and the colonies left on the plate.

## VII.IV. Discussion

Eukaryotic cell cycle regulation is controlled by the activity of CDKs. WEE1, as a key regulator of CDK, plays a critical role in the G2/M transition of the cell cycle. Previous work indicates that protein phosphorylation may play an important role in the control of WEE1 activity in yeast (Russell and Nurse, 1987; Coleman *et al.*, 1993; Parker *et al.*, 1993; O'Connell *et al.*, 1997; Boddy *et al.*, 1998). However, little is known about how WEE1 activity is regulated in plants.

In an attempt to gain a better understanding of how WEE1 activity is regulated in plant cells during the cell cycle, the yeast two-hybrid screen approach was used to identify proteins that interact with WEE1. To investigate AtWEE1 interactions with other proteins, the pBD-GAL4-Cam *AtWEE1* plasmid was successfully transformed into the Stratagene two hybrid yeast strain YRG-2 as demonstrated by the growth on selective minimal medium lacking tryptophan and by PCR. Positive colonies carrying the binding domain were re-transformed using the pAD-GAL4-2.1 cDNA library. The presence of the activating domain inside YRG-2 cells already containing pBD-GAL4-Cam *AtWEE1* was confirmed by PCR and growth on selective minimal medium lacking histidine, leucine and tryptophan with 40mM 3-AT.

The yeast two hybrid system was used in previous work on mouse and human cells to identify proteins which bind to WEE1 kinase. In mouse cells, the yeast two-hybrid technique was performed. Using the carboxyl half of WEE1 kinase the 14-3-3  $\zeta$  protein was isolated. Recombinant 14-3-3  $\zeta$  binds to WEE1 kinase *in vitro*, but the functional significance of this protein-protein interaction was not addressed in the study (Honda *et*

*al.*, 1997). 14-3-3 proteins are a family of acidic proteins of low molecular weight found in all eukaryotic cells. There are 7 known 14-3-3 isoforms in humans and 15 in plants (Rosenquist *et al.*, 2000, 2001). In human cells, the yeast two hybrid screen demonstrated that WEE1 activity in mammalian cells can be regulated by 14-3-3  $\beta$ . The 14-3-3  $\beta$  protein binds directly to WEE1 at the COOH-terminal RSVSLT motif. Phosphorylation of a serine residue within this motif is required for the binding. The interaction of WEE1 with 14-3-3  $\beta$  protein may change WEE1 protein conformation and block the degradation motif contained in the NH<sub>2</sub>-terminal domain (Wang *et al.*, 2000; Rothblum-Oviatt *et al.*, 2001).

In my study the yeast two hybrid technique was performed using the whole AtWEE1 ORF as the bait domain to screen an *Arabidopsis* root cDNA activation domain library. Interactions between *AtWEE1* and other proteins were not detected using the LacZ assay when glucose (2%) was used as the carbon source to prepare the minimal media. Interestingly, subsequently in the Cardiff cell cycle laboratory when Anne Lentz (unpublished data) changed the composition of the minimal medium adding galactose (2%) and raffinose (1%) instead of glucose (2%) as specified in the Clontech Yeast Protocol Handbook (published the 14<sup>th</sup> March 2001), 82 positive yeast patches were obtained from the LacZ assay. The 82 positive results were submitted for sequencing and 77 of those gave a good quality sequence. In the table below (Table VII.I) there is a list of proteins that were detected as interacting with AtWEE1 via the two-hybrid screen in more than one case.



**Table VII.I:** proteins interacting with AtWEE1 detected more than once using the yeast two-hybrid technique (Anne Lentz, Cardiff cell cycle laboratory, unpublished data)

| Protein                        | Number of hits | Function   | Reference                       |
|--------------------------------|----------------|--|---------------------------------|
| bZIP Transcription factor GBF5 | 2              | This transcription factor regulates processes including pathogen defense, light and stress signaling, seed maturation and flower development. The <i>Arabidopsis</i> genome sequence contains 75 distinct members of the bZIP family.  | Jakoby <i>et al.</i> , 2002     |
| SCL6                           | 3              | Member of GRAS proteins. Although the <i>Arabidopsis</i> genome encodes at least 33 GRAS protein family members only a few GRAS proteins have been characterized so far. However, it is becoming clear that GRAS proteins exert important roles in very diverse processes such as signal transduction, meristem maintenance and development. | Bolle C., 2004                  |
| Chitinase AtCTL1               | 2              | Chitinase-like proteins have been proposed to play roles in normal plant growth and development and lignin deposition.   | Zhong <i>et al.</i> , 2002      |
| Strictosidine synthase         | 3              | Strictosidine synthase (STR1) is a central enzyme that participates in the biosynthesis of almost all plant monoterpenoid indole alkaloids.  | Treimer and Zenk, 1979          |
| Cdc48A                         | 2              | Cell cycle gene involved in the spindle pole body separation.  | Feiler <i>et al.</i> , 1995     |
| Brix domain protein            | 3              | Role in ribosome biogenesis and rRNA binding.  | Eisenhaber <i>et al.</i> , 2001 |
| Copper chaperone ATX1          | 3              | Bind copper ions and deliver them to specific cellular pathways.   | Liu <i>et al.</i> , 2003        |
| GSTs AtGSTF9<br>Phi class GST  | 2              | Glutathione S-transferases (GSTs) appear to be ubiquitous in plants and have defined roles in herbicide detoxification. In contrast, little is known about their roles in normal plant physiology and during responses to biotic and abiotic stress.   | Wagner <i>et al.</i> , 2002     |

To confirm that the putative positive interactions occur *in planta* and are not false positives further tests need to be done. The specificity of the interaction between AtWEE1 and the proteins listed in Table VII.I has to be verified *in vivo*. One method for doing this is to transform *Arabidopsis thaliana* plants with plasmids carrying the gene encoding for the above proteins fused to non-fluorescent domains of the yellow



fluorescent protein. The two non-fluorescent domains become fluorescent when brought together by association of AtWEE1 with the interacting protein (Hu *et al.*, 2000, Walter *et al.*, 2004).

## VII.V. Summary

To gain a better understanding of WEE1 kinase regulation in plants, the yeast two hybrid screen was used to isolate proteins that may interact with AtWEE1. The construction of pBD-GAL4 Cam-*AtWEE1* plasmid and its transformation into yeast YRG-2 cells were successfully performed. Co-transformation of YRG-2 cells carrying AtWEE1 bait with pAD-GAL4-2.1 cDNA library from *Arabidopsis* root was also successful. The first screening on medium HIS- and 40 mM 3-AT gave 442 positive colonies, which were all negative when re-screened using the LacZ assay to identify blue (positive) colonies.

This work is now taken on by Anne Lentz; she found very exiting interactions between AtWEE1 and other proteins. These interactions need to be further tested to confirm that they are not false positive.

