

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/65530/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Siani, Harsha and Maillard, Jean-Yves 2015. Best practice in healthcare environment decontamination. *European Journal of Clinical Microbiology & Infectious Diseases* 34 (1), pp. 1-11. 10.1007/s10096-014-2205-9

Publishers page: <http://dx.doi.org/10.1007/s10096-014-2205-9>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Harsha Siani and Jean-Yves Maillard*

Best practice in healthcare environment decontamination.

Cardiff School of Pharmacy and Pharmaceutical Sciences, Cardiff University, Wales

*Address for correspondence:

College of Biomedical and Life Sciences

Cardiff School of Pharmacy and Pharmaceutical Sciences

Cardiff University

Redwood Building

King Edward VII Avenue

Cardiff CF10 3NB

United Kingdom

Tel: +442920879088

Fax: +442920874149

Email: maillardj@cardiff.ac.uk

Abstract

There is now strong evidence that surface contamination is linked to healthcare associated infections. Cleaning and disinfection should be sufficient to decrease microbial bioburden from surfaces in healthcare settings, and overall help in decreasing infections. It is however not necessarily the case. Evidence suggests there is a link between educational interventions and a reduction in infections. To improve the overall efficacy and appropriate usage of disinfectants, manufacturers need to engage with the end users in providing clear claim information and product usage instructions. This review provides a clear analysis of scientific evidence supporting the role of surfaces in healthcare associated infections, and the role of education in decreasing such infections. It is also looking at the debate opposing the use of cleaning vs. disinfection in healthcare settings.

Keywords: cleaning, disinfection, antibiotic resistance, surface, HCAI

Introduction

Healthcare-associated infections (HCAIs) are defined as infections associated with interventions, devices or procedures carried out in healthcare facilities occurring in patients at the time of hospital admission or within 48 hours of admission [1,2]. In 2011-2012 the European Centre for Disease Prevention and Control (ECDC) coordinated a Point Prevalence Survey (PPS) of HCAIs in acute care hospitals in Europe, the study revealed 6.0% of patients were infected with at least one HCAI, of which 54% were associated with a previous stay in the same hospital. It is estimated on any given day 81,089 patients have a HCAI in Europe, with the most common HCAI associated with respiratory tract infections [2]. Non-device related infections might account for a significant proportion of HCAIs [2]. The most frequently reported microorganisms in HCAIs are *Escherichia coli* (15.9%), followed by *Staphylococcus aureus* (12.3%), *Enterococcus* spp. (9.6%), *Pseudomonas aeruginosa* (8.9%), *Klebsiella* spp. (8.7%), coagulase-negative staphylococci (7.5%) (see [2] for more details). While *Clostridium difficile* accounts only for 5.4%, it is responsible for 48% of all gastrointestinal infections.

It is conservatively estimated that HCAs cost the NHS £1 billion annually (£3,154 per patient) [1,3-5]. Significantly it is believed that 20-30% of HCAs could be avoided with better application of existing knowledge and realistic infection control practices [4]. Enhanced cleaning practices are reported to save hospitals between £30,000–£70,000 [6]. With this in mind, infection prevention and control should be a priority at the forefront for all healthcare professionals and users, with a high standard of cleanliness being an intrinsic part of infection prevention. With HCAs, such as methicillin-resistant *Staphylococcus aureus* (MRSA), *Clostridium difficile* and Norovirus, frequently reported in the media, infection control policies are subject to increased public scrutiny. Yet there appears to be a distinct lack of investment in the field of infection control, both from a research and product application perspective. This is also concurrent with a lack of understanding in disinfectant (biocidal product) efficacy and usage, which are often associated with, or lead to, poor practice. This review aims to analyse in more details the issues faced by infection control professionals and the industry.

The unjustified controversy of cleaning/disinfection failing to impact on HCAs

Until relatively recently, there was a belief that the hospital environment was not a source of transmission for HCAs. Indeed, early studies in the 1970s and 1980s indicated endemic transmission of pathogens via the hospital environment was negligible [7,8]. Since then a number of investigators have highlighted the importance of environmental contamination in the transmission of clinically relevant pathogens, such as *C. difficile* and MRSA [9-12] as well as the role of surface disinfection for controlling pathogenic microorganisms [13]. The importance of surface disinfection is further emphasised by its inclusion in several national and international infection control policies, including the epic3: national evidence-based guidelines for preventing healthcare-associated infections in NHS hospitals in England [14].

Hospital setting, environmental persistence and transmission

The most common source of microorganisms in a hospital setting are the patients themselves, infected and colonised patients (and hospital staff) shed bacteria, viruses and spores into the hospital environment. Whilst a direct link between HCAs and the presence of a microorganism on a hospital surface has not been established

[10,15-19], studies have reported many organisms responsible for HCAs, including MRSA, *C. difficile*, norovirus and vancomycin-resistant enterococci, survive and persist on hospital surfaces at concentrations sufficient for transmission and transference to the hands of healthcare workers. Given that the infectious dose for most potential pathogens appears to be low [20-22] coupled with the persistence of these organisms on hospital surfaces and medical equipment for prolonged periods (Table 1)[23], the presence of a pathogen on a surface does pose a transmission and/or infection risk (Table 2)[10-24].

In the hospital environment, areas near the patient and high-touch surfaces have been found to harbour microorganisms (Figure 1)[10,13,15,50,51]. A number of studies highlighted the transference of microorganisms from surfaces to hands (Table 3). Kampf and Kramer [61] reported the percentage of pathogens on healthcare workers hands as rhinovirus (65%), and rotavirus (19.5-78.6%), vancomycin-resistant *enterococci* (41%), *Clostridium difficile* (14-59%), *Klebsiella* spp. (17%), MRSA (16.9%), *Serratia marcescens* (15.4-24%), *Pseudomonas* spp (1.3-25%) and *Acinetobacter* spp. (3-15%). Adequate cleaning and/or disinfection of these surfaces (bedrails, commodes, doorknobs, light switches, patient call button, surfaces and equipment in close proximity to the patient) have been shown to be of particular importance [62-66]. It has been suggested that cleaning specifications do not adequately address high touch surfaces [10,50,67], with increased frequency and intensity of cleaning recommended for pathogens with an intestinal reservoir (*C. difficile* and norovirus)[68,69]. More recently, it has been suggested that cleaning and/or disinfection protocols should be ward specific and hence tailored to prevent ward-specific transmission routes. In addition to focusing on near patient surfaces, staff (medical chart, drug locker, staff toilet,) and patient (paper towel dispensers, bin lids) contact surfaces should also be considered as reservoirs of infection [70].

Evidence that surface decontamination eliminates transmission and lowers infection rates

There is an increasing body of knowledge which highlights improved infection control practices can help break the chain of transmission [20,71,72]. A review was undertaken by Rutala and Weber [72] who recommended routine cleaning and disinfection of surfaces following a comprehensive review of epidemiological and

microbiological data following surface disinfection. Studies which show a positive impact of environmental cleaning have focussed predominantly on MRSA, *C. difficile* and norovirus, which is not surprising, given the infection rates and the ability of these organisms to persist in the environment (Table 4).

Roles, responsibilities and education of healthcare workers

The document compiled by the Comptroller and Auditor General on behalf of the National Audit Office [4] identified three staffing groups with cleaning responsibilities: a) dedicated cleaning staff, b) nursing, ambulance staff and departmental staff and c) estates staff. The division of cleaning responsibilities has often resulted in confusion, resulting in a number of objects (ward-based equipment) which ‘fall through the gaps’ in the cleaning schedule [106,107]. With this in mind it is apt to refer to the Matron’s Charter which specifies that cleanliness is everyone’s responsibility, not just the cleaner’s [108]. Nonetheless it is evident that regular teaching of microbiological principles and infection control policies is beneficial [107,109].

Cleaning and disinfection form a fundamental part of infection control and prevention, integral to this is the appropriate education and training of all NHS personnel (medical and non-medical staff) and NHS users (patients and visitors). However, there appears to be a disparity in the provision of education and training provided to key healthcare personnel in the NHS. Nurses and healthcare assistants were provided with induction training on infection control in 90% of NHS Trusts, whilst only 16% of senior doctors received training [4]. The importance of education and training is reinforced by evidence that they can contribute to reductions in HCAs (Table 5).

The education of healthcare workers may be hampered by the lack of general guidelines on cleaning standard and evaluation, and by conflicting information between the need for cleaning and/or disinfection and the evaluation of disinfectants/cleaning agents. There are no guidelines or standardised methods for monitoring of environmental cleaning. Visual assessment is the most generally accepted measure of cleanliness [115,116], despite being an unreliable indicator of microbial contamination. Currently the UK guidelines for surfaces in wards, is that they are “visually clean” [16]. A surface may be visibly free of soil however; this may

not reflect that the surface is free of microbial load. Visual assessments are the cheapest and quickest means of assessing cleanliness, providing an indication of personal performance and cleaning efficiencies. However, subjective visual inspections have been reported to be poor indicators of cleanliness in comparison to fluorescent markers and adenosine triphosphate (ATP) assays [117].

UK guidelines do not currently advise on the routine sampling of floors, walls, surfaces and air [118]. Given that high touch surfaces are implicated in the transmission of HCAs, validating and assessing the thoroughness of cleaning would be justified, serving as an additional training and educational tool. If sampling is to be undertaken the number of organisms per unit area or volume should be reported. Despite the time and resources required for microbial culturing, it represents the most accurate indication of the potential infection risk. The presence of indicator organisms, such as *S. aureus*, *C. difficile*, VRE or *Acinetobacter* spp., is indicative of a requirement for increased cleaning [119]. It has been proposed that aerobic colony counts on hand-touch sites should not exceed 2.5 CFU/cm² [55,119-121].

Cleaning or disinfection?

The choice of decontamination procedure will depend on the infection risk associated with the surface and the type of microorganism likely to have contaminated the surface [122,123]. An inherent consideration of all disinfection strategies is the elimination of the most resistant microbial sub-population. Yet there are disagreements about when and where a cleaning agent (removing of a bioburden from surfaces) or a disinfectant (killing microorganisms on surfaces) should be used (Table 6). This is further complicated by the fact that many disinfectant products will have a detergent (cleaning) ability too. In addition, there are many factors that will affect the efficacy of a disinfectants; these include factors related to the disinfectant such as concentration, pH and overall formulation, factors related to the target microorganisms and factors related to the product usage such as contact time, organic load, type of surface and temperature [124]. Failure to understand the effect of these factors on antimicrobial activity will result in the failure of the disinfectant. To assess the efficacy of a disinfectant a number of standard efficacy tests can be performed. These efficacy tests are key to product development and are the basis for regulatory clearance, labelling and use [125]. The type of test method employed and the requirements will depend on the type and intended

purpose of the microbicide (disinfection, preservation, and antiseptics) and its application (medical, agricultural, industrial). Although data from standardised efficacy test methods (e.g. European Norm tests) are required for a product to be commercialised and for a producer to make a product claim, the parameters used in these standard tests may not reflect realistic in-use conditions. For example, disinfectants used in the healthcare settings generally have a contact time of 10 minutes, i.e. the surface must stay wet after cleaning for 10 minutes to achieve a 3 log reduction [13], however such a long exposure time is not practical and will require re-application of the product. Generally the contact time specified on the label of a product is too long to reflect realistic in-use conditions, thus the efficacy of some products may be grossly overestimated [125,126]. The Centres for Disease Control and Prevention (CDC) specifies a contact time of 3-5 minutes based on the evaporation of the product, however a 1 minute contact time is more realistic of in-use conditions, indeed contact times of 30 – 60 seconds have been reported for a number of disinfectants [127-129]. For antimicrobial wipes, there is no international or national guidance on wipe selection and use [130,131]. Without an accepted standard test for wipes, information on the effectiveness of a product can only be gleaned from laboratory tests. This can lead to the use of wipes that might not be appropriate for applications in the health care environment [132,133].

The choice of disinfectant will depend on its intended use, thus the manufacturer's instructions should be followed to ensure correct application [124]. Incorrect selection and use of a formulated disinfectant can result in transference of microorganisms to clean surfaces [65,132,134-138]. In laboratory simulated conditions, studies have demonstrated the transference of microorganisms from contaminated cleaning cloths (commercial wipes and microfiber cloths) to clean surfaces [132,139,140]. Nine of the ten commercially available wipes tested demonstrated the repeated transfer of *C. difficile* spores [132]. The changing and/or cleaning of cloths and the wiping of surfaces from clean to dirty is crucial to limiting microbial transference [133].

Conclusions

A valid infection control intervention will reduce the microbial burden in the environment and hence the persistence of the organism, which can only be achieved with appropriate cleaning and disinfection programmes. As such surface disinfection

should be included in local, national and international infection control policies. The current debate as to whether or not cleaning only (i.e. without a disinfection step) is sufficient to eliminate microbial pathogens from surfaces in the healthcare environment needs to be addressed and supported by practical evidence.

It is clear that better education together with better compliance [141] of end users is needed. A number of considerations can easily be taken into account prior to choosing a disinfectant (Table 7). A product label will state the name of the test method used to assess the efficacy of the product. Information relating to the test, the laboratory in which it was undertaken and the test results should be available from the manufacturer.

Concurrently, manufacturers need to have clear instructions about standard efficacy tests that need to be performed not only to make a product claim but also to represent better the usage of a product in practice. If no standard test is available, manufacturers should be encouraged to provide evidence of the activity of their products under in use conditions. Unfortunately, it is increasingly clear that a product that passes a standard efficacy test (such as European Norm tests) will meet its label claim but it might not necessarily mean that the product will be efficacious in practice; two of the most documented examples are the use of antimicrobial wipes¹³⁰⁻¹³³ or the testing of products against *Clostridium difficile* [142-144]. Manufacturers need also to provide clear product use instructions. Decreasing microbial bioburden on surfaces through cleaning and disinfection should be easily achievable with most of the disinfectant formulations available at present. More efforts need to be done to educate and motivate the end users to use the purposefully designed disinfectant appropriately. Decreasing HCAs remain a multifactorial approach [145] in which surface decontamination is central [14].

Conflict of interest

None to declare

Funding and sponsorship

H Siani is funded by Cardiff University and recipient of a Knowledge Transfer Partnership with GAMA Healthcare.

References

1. Health Protection Agency (2012) English national point prevalence survey on healthcare associated infections and antimicrobial use, 2011: Preliminary data. Health Protection Agency, London
2. European Centre for Disease Prevention and Control (2011) Point prevalence survey of healthcare-associated infections and antimicrobial use in European acute care hospitals. ECDC, Stockholm
3. Plowman R, Graves N, Griffin MAS et al (2001) The rate and cost of hospital-acquired infections occurring in patients admitted to selected specialties of a district general hospital in England and the national burden imposed. *J Hosp Infect* 47:198-209
4. National Audit Office (2009) The management and control of hospital acquired infection in acute NHS trusts in England. Report from the Comptroller and Auditor General. The Stationary Office, London
5. The International Federation of Infection Control (2011) IFIC basic concepts of infection control, 2nd edn. Friedman C, Newson W, Eds. IFIC, Portadown
6. Dancer SJ, White L, Lamb J, Girvan EK, Robertson C (2009) Measuring the effect of enhanced cleaning in a UK hospital: a prospective cross-over study. *BMC Med* 7:28
7. Weber D, Gooch JJ, Wood WR, Britt EM, Kraft RO (1976) Influence of operating room surface contamination on surgical wounds: A prospective study. *Arch Surg* 111:484-488
8. Maki DG, Alvarado CJ, Hassemer CA, Zilz MA (1982) Relation of the inanimate hospital environment to endemic nosocomial infection. *New Engl J Med* 307:1562-1566
9. Weber DJ, Rutala WA, Miller MB, Huslage K, Sickbert-Bennett E (2010) Role of hospital surfaces in the transmission of emerging health care-associated pathogens: Norovirus, *Clostridium difficile*, and *Acinetobacter* species. *Am J Infect Control* 38:S25-S33
10. Dancer SJ (2009) The role of environmental cleaning in the control of hospital-acquired infection. *J Hosp Infect* 73:378-385
11. Manangan LP, Pugliese G, Jackson M et al (2001) Infection control dogma: top 10 suspects. *Infect Control Hosp Epidemiol* 22:243-247

12. Landelle C, Verachten M, Legrand P, Girou E, Barbut F, Brun Buisson C (2014) Contamination of healthcare workers' hands with *Clostridium difficile* spores after caring for patients with *C. difficile* infection. *Infect Control Hosp Epidemiol* 35:10-15
13. Gebel J, Exner M, French G et al (2013) The role of surface disinfection in infection prevention. *GMS Hyg Infect Control* 8:1-12
14. Loveday H, Wilson J, Pratt R et al (2014) epic3: National evidence-based guidelines for preventing healthcare-associated infections in NHS hospitals in England. *J Hosp Infect* 86:S1-S70
15. Rutala WA, Weber DJ, Weinstein RA et al (2008) Guideline for disinfection and sterilization in healthcare facilities. Centers for Disease Control and Prevention, Atlanta, GA
16. Pratt RJ, Pellowe CM, Wilson JA et al (2007) epic2: National evidence-based guidelines for preventing healthcare-associated infections in NHS hospitals in England. *J Hosp Infect* 65(Suppl. 1):S1-S64
17. Dettenkofer M, Wenzler S, Amthor S, Antes G, Motschall E, Daschner FD (2004) Does disinfection of environmental surfaces influence nosocomial infection rates? A systematic review. *Am J Infect Control* 32:84-89
18. Health Protection Scotland (HPS)(2012) Standard Infection Control Precautions (SICPs) literature review: Routine cleaning of the environment in the hospital setting. HPS; 2012. <http://www.documents.hps.scot.nhs.uk/hai/infection-control/ic-manual/sicp-lr-cleaning-v1.0.pdf> Accessed 24 March 2014
19. Sehulster L, Chinn RY (2003) Guidelines for environmental infection control in health-care facilities: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *Morbidity and Mortality Weekly Report - Recommendations and Reports* 52:1-42
20. Otter JA, Yezli S, French GL (2011) The role played by contaminated surfaces in the transmission of nosocomial pathogens. *Infect Control Hosp Epidemiol* 32:687-699
21. Lawley TD, Clare S, Deakin LJ et al (2010) Use of purified *Clostridium difficile* spores to facilitate evaluation of health care disinfection regimens. *Appl Environ Microbiol* 76:6895-6900
22. Teunis PF, Moe CL, Liu P et al (2008) Norwalk virus: How infectious is it? *J Med Virol* 80:1468-1476

23. Kramer A, Schwebke I, Kampf, G (2006) How long do nosocomial pathogens persist on inanimate surfaces? A systematic review. *BMC Infect Dis* 6:130
24. Guerrero DM, Nerandzic MM, Jury LA, Jinno S, Chang S, Donskey CJ (2012) Acquisition of spores on gloved hands after contact with the skin of patients with *Clostridium difficile* infection and with environmental surfaces in their rooms. *Am J Infect Control* 40:556-558
25. Chen K-H, Chen L-R, Wang Y-K (2014) Contamination of medical charts: An important source of potential infection in hospitals. *PLoS ONE* 9:e78512
26. Tan TY, Tan JSM, Tay H, Chua GH, Ng LSY, Syahidah N (2013) Multidrug-resistant organisms in a routine ward environment: Differential propensity for environmental dissemination and implications for infection control. *J Med Microbiol* 62:766-772
27. Jury LA, Sitzlar B, Kundrapu S et al (2013) Outpatient healthcare settings and transmission of *Clostridium difficile*. *PLoS ONE* 8:e70175
28. Kiedrowski LM, Perisetti A, Looock MH, Khaitza ML, Guerrero DM (2013) Disinfection of iPad to reduce contamination with *Clostridium difficile* and methicillin-resistant *Staphylococcus aureus*. *Am J Infect Control* 41:1136-1137
29. Faires MC, Pearl DL, Olaf Berke O, Reid-Smith RJ, Weese JS (2013) The identification and epidemiology of methicillin-resistant *Staphylococcus aureus* and *Clostridium difficile* in patient rooms and the ward environment. *BMC Infect Dis* 13:342
30. Sigler V, Hensley S (2013) Persistence of mixed Staphylococci assemblages following disinfection of hospital room surfaces. *J Hosp Infect* 83:253-256
31. Muzslay M, Moore G, Turton JF, Wilson AP (2012) Dissemination of antibiotic-resistant enterococci within the ward environment: The role of airborne bacteria and the risk posed by unrecognized carriers. *Am J Infect Control* 41:283
32. Best E, Sandoe J, Wilcox M (2012) Potential for aerosolization of *Clostridium difficile* after flushing toilets: the role of toilet lids in reducing environmental contamination risk. *J Hosp Infect* 80:1-5
33. Shaughnessy MK, Micielli RL, DePestel DD et al (2011) Evaluation of hospital room assignment and acquisition of *Clostridium difficile* infection. *Infect Control Hosp Epidemiol* 32:201-206

34. Ferreira AM, Andrade D, Rigotti, MA, Almeida MTG (2011) Methicillin-resistant *Staphylococcus aureus* on surfaces of an Intensive Care Unit. *Acta Paulista de Enfermagem* 24:453-458
35. Bhatta D, Gokhale S, Ansari M et al (2011) Stethoscopes: A possible mode for transmission of nosocomial pathogens. *J Clin Diagn Res* 5:S1173-1176
36. Dumford III DM, Nerandzic MM, Eckstein BC, Donskey CJ (2009) What is on that keyboard? Detecting hidden environmental reservoirs of *Clostridium difficile* during an outbreak associated with North American pulsed-field gel electrophoresis type 1 strains. *Am J Infect Control* 37:15-19
37. Drees M, Snyderman D, Schmid C et al (2008) Prior environmental contamination increases the risk of acquisition of vancomycin-resistant enterococci. *Clin Infect Dis* 46:678-685
38. Zanetti G, Blanc D, Federli I et al (2007) Importation of *Acinetobacter baumannii* into a burn unit: A recurrent outbreak of infection associated with widespread environmental contamination. *Infect Control Hosp Epidemiol* 28:723-725
39. Huang SS, Datta R, Platt R (2006) Risk of acquiring antibiotic-resistant bacteria from prior room occupants. *Arch Int Med* 166:1945-1951
40. Hayden MK, Bonten MJM, Blom DW, Lyle EA, van de Vijver DAMC, Weinstein RA (2006) Reduction in acquisition of vancomycin-resistant *Enterococcus* after enforcement of routine environmental cleaning measures. *Clin Infect Dis* 42:1552-1560
41. van der Mee-Marquet N, Girard S, Lagarrigue F et al (2006) Multiresistant *Enterobacter cloacae* outbreak in an intensive care unit associated with therapeutic beds. *Critical Care* 10:405
42. Grabsch EA, Burrell LJ, Padiglione A, O'Keefe JM, Ballard S, Grayson ML (2006) Risk of environmental and healthcare worker contamination with vancomycin-resistant enterococci during outpatient procedures and hemodialysis. *Infect Control Hosp Epidemiol* 27:287-293
43. Rampling A, Wiseman S, Davis L et al (2001) Evidence that hospital hygiene is important in the control of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 49:109-116

44. Cohen SH, Tang YJ, Rahmani D, Silva Jr, J (2000) Persistence of an endemic (toxigenic) isolate of *Clostridium difficile* in the environment of a general medicine ward. Clin Infect Dis 30:952-954
45. Wagenvoort J, Sluijsmans W, Penders R (2000) Better environmental survival of outbreak vs. sporadic MRSA isolates. J Hosp Infect 45:231-234
46. Jawad A, Seifert H, Snelling A, Heritage J, Hawkey P (1998) Survival of *Acinetobacter baumannii* on dry surfaces: comparison of outbreak and sporadic isolates. J Clin Microbiol 36:1938-1941
47. Hirai Y (1991) Survival of bacteria under dry conditions; from a viewpoint of nosocomial infection. J Hosp Infect 19:191-200
48. Duckworth GJ, Jordens JZ (1990) Adherence and survival properties of an epidemic methicillin-resistant strain of *Staphylococcus aureus* compared with those of methicillin-sensitive strains. J Med Microbiol 32:195-200
49. Kim KH, Fekety R, Batts D et al (1981) Isolation of *Clostridium difficile* from the environment and contacts of patients with antibiotic-associated colitis. J Infect Dis 143:42-50
50. Dancer SJ (2008) Importance of the environment in methicillin-resistant *Staphylococcus aureus* acquisition: the case for hospital cleaning. Lancet Infect Dis 8:101-113
51. Smith SJ, Young V, Robertson C, Dancer SJ (2012) Where do hands go? An audit of sequential hand-touch events on a hospital ward. J Hosp Infect 80:206-211
52. Istenes N, Bingham J, Hazelett S et al (2013) Patients' potential role in the transmission of health care-associated infections: Prevalence of contamination with bacterial pathogens and patient attitudes toward hand hygiene. Am J Infect Control 41:793-798
53. Kundrapu S, Sunkesula V, Jury LA, Sitzlar BM, Donskey CJ (2012) Daily disinfection of high-touch surfaces in isolation rooms to reduce contamination of healthcare workers' hands. Infect Control Hosp 33:1039-1042
54. Hayden MK, Blom DW, Lyle EA, Moore CG, Weinstein RA (2008) Risk of hand or glove contamination after contact with patients colonized with vancomycin-resistant *Enterococcus* or the colonized patients' environment. Infect Control Hosp Epidemiol 29:149-154

55. White LF, Dancer SJ, Robertson C, McDonald J (2008) Are hygiene standards useful in assessing infection risk? *Am J Infect Control* 36:381-384
56. Duckro AN, Blom DW, Lyle EA, Weinstein RA., Hayden MK (2005) Transfer of vancomycin-resistant enterococci via health care worker hands. *Arch Int Med* 165:302-307
57. Bhalla A, Pultz NJ, Gries DM et al (2004) Acquisition of nosocomial pathogens on hands after contact with environmental surfaces near hospitalized patients. *Infect Control Hosp Epidemiol* 25:164-167
58. Weinstein RA, Bridges CB, Kuehnert MJ et al (2003) Transmission of influenza: implications for control in health care settings. *Clin Infect Dis* 37:1094-1101
59. Ray AJ, Hoyen CK, Taub TF, Eckstein EC, Donskey CJ (2002) Nosocomial transmission of vancomycin-resistant enterococci from surfaces. *J Amer Med Assoc* 287:1400-1401
60. Boyce JM, Potter-Bynoe G, Chenevert C, King T (1997) Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: possible infection control implications. *Infect Control Hosp Epidemiol* 18:622-627
61. Kampf G, Kramer A (2004) Epidemiologic background of hand hygiene and evaluation of the most important agents for scrubs and rubs. *Clin Microbiol Rev* 17:863-893
62. Huslage K, Rutala WA, Weber DJ (2010) A quantitative approach to defining "high-touch" surfaces in hospitals. *Infect Control Hosp Epidemiol* 31:850-853
63. Casewell M, Phillips I (1977) Hands as route of transmission for *Klebsiella* species. *Br Med J* 2(6098):1315-1317
64. Sanderson PJ, Weissler S (1992) Recovery of coliforms from the hands of nurses and patients: activities leading to contamination. *J Hosp Infect* 21:85-93
65. Barker J, Vipond I, Bloomfield S (2004) Effects of cleaning and disinfection in reducing the spread of norovirus contamination via environmental surfaces. *J Hosp Infect* 58:42-49
66. Roberts K, Smith C, Snelling A et al. (2008) Aerial dissemination of *Clostridium difficile* spores. *BMC Infect Dis* 8:7
67. Siegel JD, Rhinehart E, Jackson M, Chiarello L (2006) Management of multidrug-resistant organisms in healthcare settings. CDC, Atlanta, GA

68. Siegel JD, Rhinehart E, Jackson M, Chiarello L (2007) Guideline for isolation precautions: Preventing transmission of infectious agents in health care settings. *Am J Infect Control* 35(Suppl. 2):S65-S164
69. Donskey CJ (2004) The role of the intestinal tract as a reservoir and source for transmission of nosocomial pathogens. *Clin Infect Dis* 39:219-226
70. Moore G, Muzslay M, Wilson A et al (2013) The type, level, and distribution of microorganisms within the ward environment: A zonal analysis of an intensive care unit and a gastrointestinal surgical ward. *Infect Control Hosp Epidemiol* 34:500-506
71. Boyce JM (2007) Environmental contamination makes an important contribution to hospital infection. *J Hosp Infect* 65:50-54
72. Rutala WA, Weber DJ (2001) Surface disinfection: should we do it? *J Hosp Infect* 48:S64-S68
73. Friedman ND, Walton AL, Boyd S et al (2013) The effectiveness of a single-stage versus traditional three-staged protocol of hospital disinfection at eradicating vancomycin-resistant Enterococci from frequently touched surfaces. *Am J Infect Control* 41:227-231
74. Jadhav S, Sahasrabudhe T, Kalley V, Gandham N (2013) The microbial colonization profile of respiratory devices and the significance of the role of disinfection: A blinded study. *J Clin Diagn Res* 7:1021-1026
75. Sitzlar B, Deshpande A, Fertelli D, Kundrapu S, Sethi AK, Donskey CJ (2013) An environmental disinfection odyssey: Evaluation of sequential interventions to improve disinfection of *Clostridium difficile* isolation rooms. *Infect Control Hosp Epidemiol* 34:459-465
76. Chuan A, Tiong C, Maley M, Descallar J, Ziochos H (2013) Decontamination of ultrasound equipment used for peripheral ultrasound-guided regional anaesthesia. *Anaesth Intensive Care* 41:529-534
77. Boyce J, Havill N (2013) Evaluation of a new hydrogen peroxide wipe disinfectant. *Infect Control Hosp Epidemiol* 34:521-523
78. Manian FA, Griesnauer S, Bryant A (2013) Implementation of hospital-wide enhanced terminal cleaning of targeted patient rooms and its impact on endemic *Clostridium difficile* infection rates. *Am J Infect Control* 41:537-541

79. Passaretti CL, Otter JA, Reich NG et al (2013) An evaluation of environmental decontamination with hydrogen peroxide vapor for reducing the risk of patient acquisition of multidrug-resistant organisms. *Clin Infect Dis* 56:27-35
80. Chmielarczyk A, Higgins PG, Wojkowska-Mach J et al (2012) Control of an outbreak of *Acinetobacter baumannii* infections using vaporized hydrogen peroxide. *J Hosp Infect* 81:239-245
81. Ali S, Moore G, Wilson APR (2012) Effect of surface coating and finish upon the cleanability of bed rails and the spread of *Staphylococcus aureus*. *J Hosp Infect* 80:192-198
82. Munoz-Price LS, Hayden MK, Lolans K, Won S, Calvert K, Lin M, Stemer A, Weinstein RA (2010) Successful control of an outbreak of *Klebsiella pneumoniae* carbapenemase-producing *K. pneumoniae* at a long-term acute care hospital. *Infect Control Hosp Epidemiol* 31:341-347
83. Attaway III HH, Fairey S, Steed LL, Salgado CD, Michels HT, Schmidt MG (2012) Intrinsic bacterial burden associated with intensive care unit hospital beds: Effects of disinfection on population recovery and mitigation of potential infection risk. *Am J Infect Control* 40:907-912
84. Iosifidis E, Karakoula K, Protonotariou E et al (2012) Polyclonal outbreak of vancomycin-resistant *Enterococcus faecium* in a pediatric oncology department. *J Ped Hematol/Oncol* 34:511-516
85. Thom KA, Johnson JK, Lee MS, Harris AD (2011) Environmental contamination because of multidrug-resistant *Acinetobacter baumannii* surrounding colonized or infected patients. *Am J Infect Control* 39:711-715
86. Orenstein R, Aronhalt KC, McManus JE, Fedraw LA (2011) A targeted strategy to wipe out *Clostridium difficile*. *Infect Control Hosp Epidemiol* 32:1137-1139
87. Ahmad R, Tham J, Naqvi SGA, Butt U, Dixon J (2011) Supports used for positioning of patients in hip arthroplasty: is there an infection risk? *Ann Royal Coll Surg Engl* 93:130-132
88. Datta R, Platt R, Yokoe DS et al (2011) Environmental cleaning intervention and risk of acquiring multidrug-resistant organisms from prior room occupants. *Arch Int Med* 171:491-494
89. Ray A, Hoeny CK, Taub TF, Eckstein EC, Donskey CJ (2010) Use of vaporized hydrogen peroxide decontamination during an outbreak of multidrug-resistant

- Acinetobacter baumannii* infection at a long-term acute care hospital. Infect Control Hosp Epidemiol 31:1236-1241
90. Boyce JM, Havill NL, Otter JA et al (2008) Impact of hydrogen peroxide vapor room decontamination on *Clostridium difficile* environmental contamination and transmission in a healthcare setting. Infect Control Hosp Epidemiol 29:723-729
 91. Gallimore CI, Taylor C, Gennery AR, Cant AJ, Galloway A, Xerry J, Adigwe J, Gray JJ (2008) Contamination of the hospital environment with gastroenteric viruses: comparison of two pediatric wards over a winter season. J Clin Microbiol 46:3112-3115
 92. Dryden M, Parnaby R, Dailly S, Lewis T, Davis-Blues K, Otter J, Kearns A (2008) Hydrogen peroxide vapour decontamination in the control of a polyclonal methicillin-resistant *Staphylococcus aureus* outbreak on a surgical ward. J Hosp Infect 68:190-192
 93. Patel SS, Pevalin DJ, Prosser R, Couchman A (2007) Comparison of detergent-based cleaning, disinfectant-based cleaning, and detergent-based cleaning after enhanced domestic staff training within a source isolation facility. Br J Infect Control 8:20-25
 94. McMullen KM, Zack J, Coopersmith CM, Kollef M, Dubberke E, Warren DK (2007) Use of hypochlorite solution to decrease rates of *Clostridium difficile*-associated diarrhea. Infect Control Hosp Epidemiol 28: 205-207
 95. Muto CA, Blank MK, Marsh JW et al (2007) Control of an outbreak of infection with the hypervirulent *Clostridium difficile* BI strain in a university hospital using a comprehensive "bundle" approach. Clin Infect Dis 45:1266-1273
 96. Bates C, Pearse R (2005) Use of hydrogen peroxide vapour for environmental control during a *Serratia* outbreak in a neonatal intensive care unit. J Hosp Infect 61:364-366
 97. Denton M, Wilcox M, Parnell P, Green D, Keer V, Hawkey P, Evans I, Murphy P (2005) Role of environmental cleaning in controlling an outbreak of *Acinetobacter baumannii* on a neurosurgical intensive care unit. J Hosp Infect 56:106-110
 98. Wright MO, Hebden JN, Harris AD et al (2004) Aggressive control measures for resistant *Acinetobacter baumannii* and the impact on acquisition of methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus* in a medical intensive care unit. Infect Control Hosp Epidemiol 25:67-168

99. Wilcox M, Fawley W, Wigglesworth N, Parnell P, Verity P, Freeman J (2003) Comparison of the effect of detergent versus hypochlorite cleaning on environmental contamination and incidence of *Clostridium difficile* infection. *J Hosp Infect* 54:109-114
100. Sample ML, Gravel D, Oxley CA, Toye B, Garber G, Ramotar K (2002) An outbreak of Vancomycin-resistant Enterococci in a hematology–oncology unit: Control by patient cohorting and terminal cleaning of the environment. *Infect Control Hosp Epidemiol* 23:468-470
101. Makris AT, Morgan L, Gaber DJ, Richter A, Rubino JR (2000) Effect of a comprehensive infection control program on the incidence of infections in long-term care facilities. *Am J Infect Control* 28:3-7
102. Falk PS, Winnike, J, Woodmansee C, Desai M, Mayhall CG (2000) Outbreak of vancomycin-resistant Enterococci in a burn unit. *Infect Control Hosp Epidemiol* 21:575-582
103. Mayfield JL, Leet T, Miller J, Mundy LM (2000) Environmental control to reduce transmission of *Clostridium difficile*. *Clin Infect Dis* 31:995-1000
104. Fitzpatrick F, Murphy OM, Brady A, Prout S, Fenelon LE (2000) A purpose built MRSA cohort unit. *J Hosp Infect* 46:271-279
105. Zafar AB, Gaydos LA, Furlong WB, Nguyen MH, Mennonna PA (1998) Effectiveness of infection control program in controlling nosocomial *Clostridium difficile*. *Am J Infect Control* 26:588-593
106. National Patient Safety Agency (2007) The national specifications for cleanliness in the NHS: a framework for setting and measuring performance outcomes. NPSA, London
107. Dancer S (2011) Hospital cleaning in the 21st century. *Eur J Clin Microbiol Infect Dis* 30:1473-1481
108. Department of Health (2004) A matron's charter: An action plan for cleaner hospitals. London
109. Davies S (2009) Making the connections: contract cleaning and infection control. UNISON, London
110. Albrecht U-V, von Jan U, Sedlacek L, Groos S, Suerbaum S, Vonberg R-B (2013) Standardized, app-based disinfection of iPads in a clinical and nonclinical setting: Comparative analysis. *J Med Internet Res* 15:e176

111. Boyce JM, Havill NL, Havill HL, Mangione E, Dumigan DG, Moore BA (2011) Comparison of fluorescent marker systems with 2 quantitative methods of assessing terminal cleaning practices. *Infect Control Hosp Epidemiol* 32:1187-1193
112. Goodman ER, Platt R, Bass R, Onderdonk AB, Yokoe DS, Huang SS (2008) Impact of an environmental cleaning intervention on the presence of methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant enterococci on surfaces in intensive care unit rooms. *Infect Control Hosp Epidemiol* 29:593-599
113. Eckstein B, Adams D, Eckstein E, Rao A, Sethi A, Yadavalli G, Donskey C (2007) Reduction of *Clostridium difficile* and vancomycin-resistant *Enterococcus* contamination of environmental surfaces after an intervention to improve cleaning methods. *BMC Infect Dis* 7:61
114. al Barrak A, Embilm J, Dyck B (1999) An outbreak of toxin negative, toxin positive *Clostridium difficile*-associated diarrhea in a Canadian tertiary-care hospital. *Can Commun Dis Rep* 25:65-69
115. United Lincolnshire Hospitals NHS Trust (2009) Cleaning manual and the national specifications for cleanliness in the NHS, Lincolnshire
116. Guh A, Carling P (2010) Options for evaluating environmental cleaning. CDC, Atlanta GA
117. Luick L, Thompson PA, Looock MH, Vetter SL, Cook J, Guerrero DM (2013) Diagnostic assessment of different environmental cleaning monitoring methods. *Am J Infect Control* 41:751-752
118. Ayliffe GAJ, Fraise AP, Geddes AM, Mitchell K (2000) Control of hospital infection: a practical handbook, 4th edn. Arnold, London.
119. Dancer SJ (2004) How do we assess hospital cleaning? A proposal for microbiological standards for surface hygiene in hospitals. *J Hosp Infect* 56:10-15
120. Mulvey D, Redding P, Robertson C et al (2011) Finding a benchmark for monitoring hospital cleanliness. *J Hosp Infect* 77:25-30
121. Griffith CJ, Cooper RA, Gilmore J, Davies C, Lewis M (2000) An evaluation of hospital cleaning regimes and standards. *J Hosp Infect* 45:19-28

122. Microbiology Advisory Committee (2010) Sterilization, disinfection and cleaning of medical equipment: guidance on decontamination from the Microbiology Advisory Committee (the MAC manual). MHRA, London.
123. Maillard J-Y, Bloomfield S, Rosado Coelho J et al (2013) Does microbicide use in consumer products promote antimicrobial resistance? A critical review and recommendations for a cohesive approach to risk assessment. *Microb Drug Res* 19:344-354
124. Maillard J-Y, McDonnell G (2012) Use and Abuse of Disinfectants. *In Practice* 34:292-299
125. Fraise A (2011) Currently available sporicides for use in healthcare, and their limitations. *J Hosp Infect* 77:210-212
126. Maillard J-Y (2005) Antimicrobial biocides in the healthcare environment: efficacy, usage, policies, and perceived problems. *Ther Clin Risk Manag* 1:307-320
127. Rutala WA, Barbee SL, Aguiar NC, Sobsey MD, Weber DJ (2000) Antimicrobial activity of home disinfectants and natural products against potential human pathogens. *Infect Control Hosp Epidemiol* 21:33-38
128. Best M, Sattar S, Springthorpe VS, Kennedy M (1990) Efficacies of selected disinfectants against *Mycobacterium tuberculosis*. *J Clin Microbiol* 28:2234-2239
129. Best M, Kennedy M, Coates F (1990) Efficacy of a variety of disinfectants against *Listeria* spp. *Appl Environ Microbiol* 56:377-380
130. Royal College of Nursing (2011) The selection and use of disinfectant wipes. RCN, London
131. Sattar SA, Maillard J-Y (2013) The crucial role of wiping in decontamination of high-touch environmental surfaces: review of current status and directions for the future. *Am J Infect Control* 41:S97-S104
132. Siani H, Cooper C, Maillard J-Y (2011) Efficacy of “sporicidal” wipes against *Clostridium difficile*. *Am J Infect Control* 39:212-218
133. Williams GJ, Denyer SP, Hosein IK, Hill DW, Maillard J-Y (2009) Limitations of the efficacy of surface disinfection in the healthcare setting. *Infect Control Hosp Epidemiol* 30:570-573

134. Exner M, Vacata V, Hornei B, Dietlein E, Gebel J (2004) Household cleaning and surface disinfection: new insights and strategies. *J Hosp Infect* 56(Suppl. 2):S70-S75
135. Dharan S, Mourouga P, Copin P, Bessmer G, Tschanz B, Pittet D (1999) Routine disinfection of patients' environmental surfaces. Myth or reality? *J Hosp Infect* 42:113-117
136. Moore G, Hall TJ, Wilson APR, Gant VA (2008) The efficacy of the inorganic copper-based biocide CuWB50 is compromised by hard water. *Lett Appl Microbiol* 46:655-660
137. Whitby JL, Rampling A (1972) *Pseudomonas aeruginosa* contamination in domestic and hospital environments. *The Lancet* 299(7740):15-17
138. Bergen LK, Meyer M, Høg M, Rubenhagen B, Andersen LP (2009) Spread of bacteria on surfaces when cleaning with microfibre cloths. *J Hosp Infect* 71:132-137
139. Scott E, Bloomfield SF (1990) The survival and transfer of microbial contamination via cloths, hands and utensils. *J Appl Microbiol* 68:271-278
140. Moore G, Griffith C (2006) A laboratory evaluation of the decontamination properties of microfibre cloths. *J Hosp Infect* 64:379-385
141. Anderson DJ, Weber DJ, Sickbert-Bennett E (2014) On Contact Precautions: The Good, the Bad, and the Ugly. *Infect Control Hosp Epidemiol* 35:222-224
142. Maillard J-Y (2011) Innate resistance to sporicides and potential failure to decontaminate. *J Hosp Infect* 77:204-209
143. Wilcox MH, Fraise AP, Bradley CR, Walker J, Finch RG (2011) Sporicides for *Clostridium difficile*: the devil is in the detail. *J Hosp Infect* 77:187-188
144. Speight S, Moy A, Macken S et al (2011) Evaluation of the sporicidal activity of different chemical disinfectants used in hospitals against *Clostridium difficile*. *J Hosp Infect* 79:18-22
145. Tacconelli E, Cataldo MA, Dancer SJ et al (2014) ESCMID guidelines for the management of the infection control measures to reduce transmission of multidrug-resistant Gram-negative bacteria in hospitalized patients. *Clin Microbiol Infect* 20:1-55

Table 1 Persistence of microorganisms on dry surfaces (based on [23])

Organism	Persistence (range)
<i>Acinetobacter</i> spp	3 days to 5 months
<i>Clostridium difficile</i> (spores)	5 months
<i>Enterococcus</i> spp. including Vancomycin Resistant enterococci	5 days – 4 months
<i>Escherichia coli</i>	1.5 hours – 16 months
<i>Klebsiella</i> spp.	2 hours to > 30 months
<i>Mycobacterium tuberculosis</i>	1 day – 4 months
<i>Pseudomonas aeruginosa</i>	6 hours – 16 months
<i>Salmonella Typhimurium</i>	10 days – 4.2 years
<i>Shigella</i> spp.	2 days – 5 months
<i>Staphylococcus aureus</i> , including MRSA	7 days – 7 months
<i>Haemophilus influenzae</i>	12 days
Adenovirus	7 days – 3 months
Influenza virus	1 – 2 days
Norovirus and feline calici virus (FCV)	8 hours – 7 days

Table 2 Evidence of persistence of microorganisms on surfaces and/or acquisition of infection from contaminated environment

Evidence	Organisms	Reference
Plastic cover of medical charts are frequently contaminated with pathogens and may serve as source of infection	Coagulase-negative staphylococci, MRSA, <i>E. coli</i> K. <i>pneumoniae</i> and <i>A. baumannii</i>	[25]
24% of HCWs hands contaminated with <i>C. difficile</i> spores after routine care of CDI patient. 44% of the HCWs with contaminated hands provided at least one episode of direct patient care without use of gloves.	<i>C. difficile</i>	[12]
79% of sampled surfaces were positive for MDROs. Molecular typing identified related strains from patients, the environment and hands of healthcare workers	MRSA, VRE, <i>E. coli</i> and <i>K. pneumoniae</i> resistant to extended-spectrum cephalosporins,	[26]

	and carbapenem-resistant (CR) <i>A. baumannii</i> .	
14% of clinical and emergency department rooms had ≥ 1 surface contaminated with <i>C. difficile</i> . Outpatient clinics maybe an important source of community-associated <i>Clostridium difficile</i> Infection (CDI)	<i>C. difficile</i> .	[27]
15% of iPads sampled were positive for <i>S. aureus</i>	<i>S. aureus</i>	[28]
3% and 6% of hospital surfaces were contaminated with MRSA or <i>C. difficile</i> , respectively	MRSA, <i>C. difficile</i>	[29]
The persistence of potentially pathogenic staphylococci on hospital surfaces represents an infection threat	<i>Staphylococci</i> spp.	[30]
Unrecognized colonization and/or the aerosolization of Enterococci together with inadequate cleaning can lead to widespread persistence in environmental contamination	<i>Enterococci</i> spp.	[31]
Environmental contamination due to <i>C. difficile</i> aerosolisation can occur when a lidless toilet is flushed	<i>C. difficile</i>	[32]
A prior room occupant with CDI is a significant risk factor for CDI acquisition. Of the patients who acquired CDI after admission 11% had a prior occupant with CDI	<i>C. difficile</i>	[33]
60% of surfaces (gowns, bed rail/cranks, table and infusion pumps) in close proximity to patient were positive for MRSA and may serve as reservoirs for infection	MRSA	[34]
Bacterial contamination of stethoscopes ranges between 66-90% depending on site sampled (bells, earpieces and diaphragms). The presence of pathogenic and non-pathogenic bacteria on stethoscopes may pose a potential transmission risk	<i>Micrococcus</i> spp., Coagulase negative staph, MRSA, MSSA, <i>Pseudomonas</i> spp, <i>Enterobacter</i> spp., <i>E. coli</i> , Streptococcus viridans Group	[35]

Toxin-producing <i>C. difficile</i> present in non-isolation rooms (16%), physician work areas (31%), nurses work station (10%) and portable equipment (21%)	<i>C. difficile</i>	[36]
Acquisition of VRE from prior environmental contamination of ICU	VRE	[37]
Environmental contamination responsible for outbreak of <i>A. baumannii</i>	<i>A. baumannii</i>	[38]
Increased risk of acquiring MRSA and VRE from rooms previously occupied by MRSA-positive and VRE-positive patients	MRSA, VRE	[39]
Enforced environmental cleaning reduces surface contamination with VRE	VRE	[40]
Epidemiological link found between outbreak strains of <i>Enterobacter cloacae</i> and strains isolated from therapeutic beds	<i>Enterobacter cloacae</i>	[41]
Widespread VRE contamination of surfaces and hands	VRE	[42]
Epidemiological link between hospital dust and transmission of MRSA	MRSA	[43]
Presence of two toxigenic <i>C. difficile</i> in the environment accounted for 45.3% of CDAD cases	<i>C. difficile</i>	[44]
Outbreak strains survive longer than in the environment than non-outbreak strains	MRSA	[45]
Survival and persistence of <i>A. baumannii</i> on dry surfaces	<i>A. baumannii</i>	[46]
Survival and persistence of non-sporulating bacteria on dry surfaces	<i>M. bovis</i> , <i>S. aureus</i> , <i>E. faecalis</i> , <i>S. thyphi</i> , <i>Ps. aeruginosa</i>	[47]
Outbreak strains survive longer than in the environment than non-outbreak strains	MRSA	[48]
Patient, HCW and environment implicated as source of <i>C. difficile</i> contamination	<i>C. difficile</i>	[49]

Table 3 Evidence of transference of microorganisms onto hands of healthcare workers

Comments	Organisms	Reference
24% of HCWs hands contaminated with <i>C. difficile</i> spores after routine care of CDI patient	<i>C. difficile</i>	[12]
39% of patients hands were contaminated with at least 1 pathogenic organism. Pathogenic organisms can be frequently detected on hands of acute care patients.	<i>Acinetobacter</i> spp., MRSA, <i>C. difficile</i> , VRE	[52]
Molecular typing identified related strains from patients, the environment and hands of HCWs	MRSA, VRE, <i>E. coli</i> and <i>K. pneumoniae</i> resistant to extended-spectrum cephalosporins, and carbapenem-resistant (CR) <i>A. baumannii</i> .	[26]
Acquisition of <i>C. difficile</i> spores on gloved hand following contact with contaminated surfaces	<i>C. difficile</i>	[24]
Daily disinfection of high-touch surfaces was associated with a significant reduction on pathogen acquisition on hands	MRSA <i>C. difficile</i>	[53]
A 10% risk of acquiring VRE is associated with each contact with VRE colonised patient and environment	VRE	[54]
23% of samples analysed did not meet hygiene standards, with hand touch sites found to display significantly more failures than non-hand touch sites	<i>S. aureus</i> Aerobic colony counts	[55]
Transfer of VRE from inanimate objects and patient skin via hands of HCWs	VRE	[56]
Environmental contamination is an important source of MDRO transmission	MRSA, <i>P. aeruginosa</i> , Enterobacteriaceae, and <i>A. baumannii</i>	[57]
Evidence of transmission of influenza virus from objects to hands of healthcare workers	Influenza virus	[58]
Transfer of VRE onto gloved hands after contact with contaminated surfaces.	VRE	[59]
Surfaces in close proximity to patients are frequently contaminated with MRSA. The contaminated surfaces may serve as a reservoir	MRSA	[60]

of MRSA		
Patient, HCW and environment implicated as source of <i>C. difficile</i> contamination	<i>C. difficile</i>	[49]

Table 4 Evidence that cleaning and disinfection eliminates transmission and lowers colonisation/infection rates

Comments	Organisms	Reference
Surface disinfection reduced environmental contamination with VRE by 9%	VRE	[73]
Cleaning and disinfection of respiratory equipment with 70% ethanol wipe reduced fungal and bacterial contamination by 60% and 75%, respectively	<i>Pseudomonas spp.</i> , <i>Acinetobacter spp.</i> , <i>Klebsella pneumoniae</i> , <i>E.coli</i> and <i>Stenotrophomonas maltophilia</i> . <i>Candida</i> spp, <i>Streptomyces</i> spp, <i>Aspergillus</i>	[74]
Daily disinfection of high touch surfaces and a dedicated housekeeping team resulted in a 60% reduction in the number of <i>C. difficile</i> positive cultures	<i>C. difficile</i>	[75]
Disinfection of portable ultrasound machines with isopropanol reduced contamination by 85%	Coagulase negative <i>staphylococcus</i> spp., <i>Neisseria</i> spp., <i>Streptococcus</i> spp.	[76]
Cleaning with a hydrogen peroxide disinfectant wipe yielded <2.5 CFU/cm ² on 99% of surfaces	Aerobic colony count	[77]
37% reduction in CDAD rate was observed following HPV decontamination	<i>C. difficile</i>	[78]
Hydrogen peroxide vapour (VHP) decontamination of rooms reduced the likelihood of MDROs and VRE acquisition by 64% and 80% respectively	VRE, multidrug resistant Gram-negatives	[79]
Environmental cleaning, education, hand hygiene and VHP decontamination successfully controlled MRAB in an intensive therapy unit	<i>A. baumannii</i>	[80]
Antibacterial wipes reduce the numbers of bacteria	<i>S. aureus</i>	[81]

near to the patient		
Early intensification of infection control practices (disinfection, hand hygiene and education) interrupts the transmission of carbapenemase-producing <i>Klebsiella pneumoniae</i> outbreak	<i>K. pneumoniae</i>	[82]
Disinfection of bed rails reduced the intrinsic bacterial burden by up to 99%	<i>Staphylococci</i> spp. VRE	[83]
Active surveillance and adherence to infection control procedures required to prevent transmission of VRE	VRE	[84]
Patients with MDR-AB, frequently contaminate the environment. Surfaces often touched by health care workers are commonly contaminated and may facilitate transmission	<i>A. baumannii</i>	[85]
Daily disinfection with a germicidal bleach wipe reduced hospital acquired-CDAD by 85%	<i>C. difficile</i>	[86]
Use of disinfectant wipes on supports used in hip arthroplasty may reduce infection rates	Coagulase negative staphylococci, coryforms, <i>Bacilli</i> spp.	[87]
Enhanced ICU cleaning may reduce VRE and MRSA transmission and acquisition	VRE, MRSA	[88]
Environmental decontamination using VHP halted the transmission of MDR <i>A. baumannii</i> in a long term acute care hospital	MDR <i>A. baumannii</i>	[89]
Enhanced cleaning reduced microbial contamination of high-risk hand-touch sites by 32.5% and MRSA infections by 26.6%	MRSA	[10]
39% reduction in CDAD rate was observed following HPV decontamination. When adjusted for presence of epidemic NAP1 strain a 53% reduction in CDAD rate.	<i>C. difficile</i>	[90]
Changes to cleaning protocols reduced environmental contamination with gastroenteric viruses	Norovirus, astrovirus, rotavirus	[91]
Patient and staff decolonisation combined with HPV decontamination terminated MRSA outbreak on surgical ward	MRSA	[92]
Cleaning with water and detergent followed by cleaning with hypochlorite helps to achieve a greater reduction in	Aerobic count, MRSA	[93]

the total bacterial bio-burden on hand touch sites in isolation rooms		
Environmental cleaning with sodium hypochlorite solution reduced rate of CDAD	<i>C. difficile</i>	[94]
Implementation of appropriate control measures controlled <i>C. difficile</i> outbreak	<i>C. difficile</i>	[95]
Epidemiological link found between outbreak strains of <i>Enterobacter cloacae</i> and strains isolated from therapeutic beds	<i>Enterobacter cloacae</i>	[41]
Increased cleaning reduced environmental contamination of VRE	VRE	[40]
Thorough cleaning and HPV disinfection eradicated <i>Serratia marcescens</i> from a neonatal intensive care unit (NICU)	<i>Serratia marcescens</i>	[96]
Transfer of VRE from inanimate objects and patient skin via hands of HCWs	VRE	[56]
Environmental contamination is an important source of transmission of nosocomial pathogens	MRSA, <i>P. aeruginosa</i> , Enterobacteriaceae, and <i>A. baumannii</i>	[57]
Significant correlation between environmental contamination and patient colonisation/infection with <i>A. baumannii</i>	<i>A. baumannii</i>	[97]
Thorough environmental cleaning and education can reduce transmission of <i>A. baumannii</i>	<i>A. baumannii</i>	[98]
Cleaning with hypochlorite significantly reduced incidence of CDI on one ward	<i>C. difficile</i>	[99]
Environmental decontamination with 0.5% sodium hypochlorite attributed to control of VRE outbreak	VRE	[100]
Routine and thorough cleaning contributed to the control of MRSA outbreak	MRSA	[43]
Hand washing, environmental cleaning and disinfecting may help reduce infection rate in long-term care facilities	-	[101]
Barrier precautions and environmental decontamination	VRE	[102]

eradicated VRE outbreak		
Environmental disinfection with sodium hypochlorite reduced rates of CDAD	<i>C. difficile</i>	[103]
Environmental contamination by MRSA can be controlled by supervised cleaning and education	MRSA	[104]
Aggressive infection control measures (included environmental disinfection, hand washing, education) resulted in sustained decrease in CDI over 7 year period	<i>C. difficile</i>	[105]

Table 5 Evidence that education and training reduces environmental contamination

Comments	References
The use of fluorescent markers resulted in a 10% reduction in the number of positive CDI cultures after disinfection	[76]
Daily disinfection of iPads with isopropanol wipes following app based instructions reduced microbial load	[110]
Gram staining of environmental cultures and use of UV markers was successful at improving cleaning in operating rooms	[111]
Improved cleaning practices, staff education, and monitoring cleanliness reduced environmental prevalence of MRSA and VRE in ICU rooms.	[112]
Educational interventions directed at housekeeping staff reduced <i>C. difficile</i> and VRE contamination of surfaces	[113]
Implementation of appropriate control measures controlled CDI outbreak	[95]
Educating health-care workers and families of patients, and all head nurses contributed to controlling outbreak of <i>C. difficile</i>	[114]

Table 6 Summary for and against the use of detergents and disinfectants (modified from Rutala and Weber [72])

Justification for detergent use	Justification for disinfection use
Surfaces contribute minimally to endemic nosocomial infections	Surfaces may contribute to the transmission of epidemiologically important microbes (e.g., VRE, MRSA, <i>C. difficile</i> , viruses)
There is no difference in infection rates of floors cleaned with detergent versus disinfectant	Disinfectants are needed for surfaces contaminated by blood and other potentially infective material
No environmental impact associated with disposal of detergents	Disinfectants are more effective than detergents in reducing microbial load on floors
Lower costs	Detergents become contaminated and result in seeding the patient's environment with bacteria
No occupational health exposure issues	Some newer disinfectants have persistent antimicrobial activity
Use of antiseptics/disinfectants may select for antibiotic-resistant bacteria, especially where a residual activity is present	Advantage of using a single product for decontamination of floors and equipment
More aesthetically pleasing floor	Formulations can achieve a combination of cleaning and disinfection, while resulting in aesthetically pleasing floor
	Disinfectants may reduce the risk of emerging bacterial resistance

Table 7 Evaluating efficacy testing data for surface disinfection

Questions	Comments
What standardised test was used?	Test used appropriate to make a claim for healthcare application? A phase 2 step 2 test should be used [#] (i.e. surface test)
Does it support the application claim?	Bactericidal, sporicidal, fungicidal For sporicidal claim: what bacterial species was used?
Was the exposure time realistic?	A contact time of 5, 10 and >10 min for surface disinfection is not realistic (see text)
Was the test conducted in clean (0.3 g/L bovine serum albumin) or dirty conditions (3.0 g/L bovine serum albumen [#])?	Absence of test with organic load limits the practicability of the test.
Was a neutralisation step used?	Absence of neutralisation increases the apparent activity of a disinfectant [*]
Was <i>C. difficile</i> used?	Need to have information on spore production method and purity level of the preparation (>90% spores). Need to have assurance test laboratory has access to anaerobic facility.
Were controls performed?	For specific activity, for example sporicidal activity, a hypochlorite control can be used to validate the appropriateness of the test method.

[#] for European Norm test

^{*} European Norm test include neutraliser, and controls of neutraliser toxicity and efficacy.

Figure 1: Examples of high touch surfaces found to harbour microorganisms in the healthcare setting. (1) bed frame and cot sides, (2) bed controls, (3) light switch, (4) patient chair, (5) mattress, (6) tray table, (7) bedside table, (8) IV pole, (9) IV pump, (10) patient entertainment system and nurse call button

