

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/69287/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Sergouniotis, Panagiotis I., Perveen, Rahat, Thiselton, Dawn L., Giannopoulos, Konstantinos, Sarros, Marios, Davies, Jennifer R., Biswas, Susmito, Ansons, Alec M., Ashworth, Jane L., Lloyd, I. Christopher, Black, Graeme C. and Votruba, Marcela ORCID: <https://orcid.org/0000-0002-7680-9135> 2015. Clinical and molecular genetic findings in autosomal dominant OPA3-related optic neuropathy. *Neurogenetics* 16 (1) , pp. 69-75. 10.1007/s10048-014-0416-y file

Publishers page: <http://dx.doi.org/10.1007/s10048-014-0416-y>
<<http://dx.doi.org/10.1007/s10048-014-0416-y>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Clinical and molecular genetic findings in autosomal dominant *OPA3*-related optic**
2 **neuropathy**

3

4 Panagiotis I. Sergouniotis,^{1,2} Rahat Perveen,¹ Dawn L. Thiselton,^{3,4} Konstantinos
5 Giannopoulos,⁴ Marios Sarros,⁴ Jennifer R. Davies,⁴ Susmito Biswas,^{1,2} Alec M. Ansons,²
6 Jane L. Ashworth,^{1,2} I. Christopher Lloyd,^{1,2} Graeme C. Black,^{1,5} Marcela Votruba^{4,6}

7

8 ¹ Institute of Human Development, Faculty of Medical and Human Sciences, University of
9 Manchester, Manchester, UK

10 ² Manchester Royal Eye Hospital, Central Manchester University Hospitals NHS Foundation
11 Trust, Manchester M13 9WH, UK

12 ³ Virginia Institute of Psychiatric and Behavioural Genetics, Virginia Commonwealth
13 University, Richmond VA 23298-0424, USA

14 ⁴ School of Optometry and Vision Sciences, Cardiff University, Cardiff CF24 4LU, UK

15 ⁵ St Mary's Hospital, Central Manchester University Hospitals NHS Foundation Trust,
16 Manchester Academic Health Science Centre, Manchester M13 9WL, UK

17 ⁶ Cardiff Eye Unit, University Hospital Wales, Heath Park, Cardiff CF14 4XW, UK

18

19

20 Correspondence: Graeme C. Black,
21 St. Mary's Hospital, Oxford Road, Manchester M13 9WL, UK;
22 Tel: +44 (0)161 276 6269
23 Fax: +44 (0)161 276 6145
24 graeme.black@manchester.ac.uk

25

26 Key words: *OPA3*; optic atrophy; inherited optic neuropathy; congenital cataract.

27

28 | Word count: 17~~5308~~

29

30

31 **ABSTRACT**

32

33 Leber hereditary optic neuropathy and autosomal dominant optic atrophy are the two most
34 common inherited optic neuropathies. The latter has been associated with mutations in the
35 *OPA1* and *OPA3* genes. To date, only six families with *OPA3*-associated dominant optic
36 atrophy have been reported. In order to identify additional families we performed Sanger
37 sequencing of the *OPA3* gene in 75 unrelated optic neuropathy patients. Affected individuals
38 from two families were found to harbour the c.313C>G, p.(Gln105Glu) change in
39 heterozygous state; this genetic defect has been previously reported in four dominant optic
40 atrophy families. Intra- and inter-familial variability in age of onset and presenting symptoms
41 was observed. Although dominant *OPA3* mutations are typically associated with optic
42 atrophy and cataracts, the former can be observed in isolation; we report a case with no lens
43 opacities at age 38. Conversely, it is important to consider *OPA3*-related disease in
44 individuals with bilateral infantile onset cataracts and to assess optic nerve health in those
45 whose vision fail to improve following lens surgery. The papillomacular bundle is primarily
46 affected and vision is typically worse than 20/40. Notably, we describe one subject who
47 retained normal acuities into the fifth decade of life. The condition can be associated with
48 extraocular clinical features: two affected individuals in the present study had sensorineural
49 hearing loss. The clinical heterogeneity observed in the individuals reported here (all having
50 the same genetic defect in *OPA3*) suggests that the molecular pathology of the disorder is
51 likely to be complex.

52

53

54

55 **KEYWORDS**

56

57 *OPA3*; optic atrophy; inherited optic neuropathy; 3-methylglutaconic aciduria type III;
58 congenital cataract; genetic ophthalmology

59

60 INTRODUCTION

61

62 Inherited optic neuropathies are a clinically and genetically heterogenous group of disorders
63 associated with selective loss of retinal ganglion cells. Clinically, they are characterised by
64 colour vision deficits, visual field defects and, typically, bilateral, symmetrical and
65 irreversible visual loss [1]. Mitochondrial dysfunction appears to play a central role in the
66 pathophysiology of these disorders and inheritance can be mitochondrial [MIM #535000] or
67 monogenic; autosomal dominant (for example [MIM #165500] and [MIM #165300]),
68 autosomal recessive (for example [MIM #612989]) and X-linked [MIM #311050] subtypes
69 have been described. Notably, certain types of inherited optic neuropathy form part of clinical
70 syndromes that include additional ocular or non-ocular features [2].

71

72 Defects in the *OPA3* gene [MIM *606580] have been previously associated with both
73 recessive and dominant optic neuropathy. Biallelic *OPA3* mutations cause 3-
74 methylglutaconic aciduria type III [MIM #258501], a recessive neuro-ophthalmological
75 syndrome, most prevalent amongst individuals of Iraqi-Jewish origin and classically
76 characterized by the following triad: (i) bilateral optic atrophy diagnosed in the first decade of
77 life; (ii) a movement disorder (ataxia or extrapyramidal dysfunction) of variable severity
78 beginning in the first or second decade of life; (iii) increased urinary excretion of 3-
79 methylglutaconic acid [3-8]. Disease-causing *OPA3* variants inherited in an autosomal
80 dominant fashion can also cause optic neuropathy. This is often associated with
81 congenital/infantile lenticular opacities; hearing loss and neurological symptoms can also be
82 features of the disorder. Autosomal dominant *OPA3*-related disease is less common than the
83 recessive form and only six families have been identified to date. The following dominant
84 disease-associated variants have been reported [NCBI Reference Sequence: NM_025136.3]:
85 c.277G>A, p.(Gly93Ser); c.313C>G, p.(Gln105Glu) (recurrent mutation);
86 c.10_11insCGCCCG, p.(Val3_Gly4insAlaPro) [9,10].

87

88 The *OPA3* gene is composed of at least 3 exons that are alternatively spliced to produce two
89 major transcripts: *OPA3A* (exon 1 plus exon 2a; encodes the 179 amino acid isoform b; NCBI
90 Reference Sequence: NM_025136.3) and *OPA3B* (exon 1 plus exon2b; encodes the 180
91 amino acid isoform a; NCBI Reference Sequence: NM_001017989.2). Although cDNA
92 studies indicate ubiquitous expression of both transcripts, *OPA3A* is much more strongly
93 expressed in most tissues, including the brain [11]. Notably, the *OPA3A* amino acid sequence

94 appears to be more conserved in evolution and yet no human disease has been associated with
95 mutations in the *OPA3B*-specific exon 2b [7,11]. At the subcellular level, *OPA3* localises
96 predominantly to the mitochondrial inner membrane and although its function remains
97 unclear previous studies have suggested involvement in the regulation of mitochondrial
98 morphology [10,12].

99

100 In the present study we report clinical and genetic findings in two families with dominant
101 *OPA3*-related optic neuropathy. The phenotypic spectrum of the disorder is broadened and
102 intra- and inter-familial variability is discussed.

103

104

105 **MATERIALS & METHODS**

106

107 Initially, 74 unrelated individuals with a presumed diagnosis of inherited optic neuropathy
108 (mean age 43 years; range 24 to 66 years) were tested for *OPA3* mutations using Sanger
109 sequencing of the three exons and flanking intron-exon boundaries of the *OPA3* gene
110 (primers and conditions available on request). All these subjects were previously screened
111 and excluded for: (i) defects in the *OPA1* gene ([MIM *605290]; the major cause of
112 dominant optic neuropathy [2,13]) and (ii) the three primary mitochondrial mutations
113 associated with Leber hereditary optic neuropathy ([MIM #535000]; m.11778G>A,
114 m.14484T>C and m.3460G>A). Subsequently, members of an additional family with
115 suspected dominant optic atrophy and cataracts were recruited and tested for *OPA3* mutations
116 in a similar fashion.

117

118 Clinical assessment of individuals with *OPA3*-related disease included detailed history, best
119 corrected Logarithm of the Minimum Angle of Resolution (logMAR) visual acuity, dilated
120 fundus examination and optic disc imaging. Optical coherence tomographs of the optic nerve
121 head were obtained with the spectral-domain Cirrus platform (Carl Zeiss Meditec, Dublin,
122 CA, USA) in two cases. An audiogram was performed in three patients. Informed consent
123 was obtained from all participants and all investigations were conducted in accordance with
124 the principles of the Declaration of Helsinki. Institutional Review Board (IRB)/Ethics
125 Committee approval was obtained from the Multicentre Research Ethics Committee
126 (MREC).

127

128

129

130 RESULTS

131

132 Three *OPA3* coding variants that have not been previously reported in publicly available
133 databases (1000 genomes project database, National Heart, Lung, and Blood Institute Exome
134 Sequencing Project or NHLBI ESP, dbSNP Build 139, accessed Jun 2014) were identified in
135 the 180 amino acid isoform a (*OPA3B*; [NCBI Reference Sequence: NM_001017989.2]:
136 c.227C>T, p.(Ala76Val); c.379G>A, p.(Gly127Ser) and c.389G>A, p.(Gly130Glu)
137 (Supplementary Table S1). Each of these changes was detected in the heterozygous state in a
138 simplex sporadic case and it was not possible to perform segregation analysis to support their
139 pathogenicity. Also, none of these alters the amino acid sequence of isoform b (*OPA3A*) and
140 w~~We~~ therefore, consider them to be variants of unknown significance.

141

142 In addition to these variants, a heterozygous c.313C>G, p.(Gln105Glu) change (only
143 affecting *OPA3A*, the transcript encoding the 179 amino acid isoform b; [NCBI Reference
144 Sequence: NM_025136.3]) was identified in an individual with a diagnosis of dominant optic
145 atrophy (subject C1; Figure 1A). This sequence alteration is the most common *OPA3*
146 mutation associated with dominant disease as it was previously identified in four families
147 segregating optic atrophy [9,10]; therefore no further evidence was needed to confirm that
148 this is a functional variant. The proband as well as his affected sister (subject C2) and
149 daughter (subject C3) presented in the first years of life with nystagmus and abnormal optic
150 disc appearance. The clinical findings are detailed in Table 1.

151

152 The same disease-associated variant, c.313C>G, p.(Gln105Glu), was detected in all three
153 affected members of a family (Figure 1B) that were recruited and tested for *OPA3* mutations
154 at a later time. Colour disc images and clinical findings are presented in Figure 1C and Table
155 1 respectively.

156

157

158 DISCUSSION

159

160 Dominant *OPA3*-related disease is a clinically heterogeneous disorder (Table 1); some
161 patients present with poor visual behaviour and nystagmus from birth (for example subject

162 C1) while others remain asymptomatic until later in life (for example subject M2). The optic
163 neuropathy is characterised by primary involvement of the papillomacular bundle and has
164 similarities with that observed in individuals with *OPAI* mutations [14]. Age of onset is in
165 the first two decades of life (Table 1) and, although comprehensive longitudinal data are
166 lacking, patients typically experience a slowly progressive, symmetrical decrease in vision
167 [9,10]. Visual acuity is usually worse than 0.3 logMAR and it is of interest that one subject in
168 the present series had preserved acuity and no visual complaints at age 41 (subject M2; 0.1
169 logMAR right, 0.2 logMAR left; Table 1 and Figure 1D). Previous reports have shown that
170 peripheral visual fields are spared and that colour vision is impaired [9,10]. Nevertheless, loss
171 of colour discrimination is variable and without a systemic axis [10]. Subject M3 reported
172 dyschromatopsia only in the right eye, in keeping with findings from colour vision testing
173 with the Ishihara plates (right eye 4 of 17 plates; left eye 16 of 17 plates).

174

175 On fundus examination, temporal optic disc pallor was the most common finding in the
176 present cohort although diffuse pallor involving the whole neuroretinal rim and optic disc
177 excavation are not unusual [9,10]. Notably, subject M1 was referred to the clinic after a
178 routine eye test at age 60 revealed an enlarged cup-to-disc ratio (0.8) in the left eye (Figure
179 1C); an erroneous initial diagnosis of glaucoma was made as consideration was not given to
180 the fact that she had infantile-onset cataracts operated at age 35. Furthermore, subject M4 had
181 a healthy optic disc appearance in the right eye and only subtle pallor in the left eye at age 14.
182 In such equivocal cases measurement of retinal nerve fibre layer thickness with optical
183 coherence tomography can be helpful (for example Figure 1D). Future imaging studies in
184 patients with *OPA3*-related disease are expected to provide further insights.

185

186 Crystalline lens opacities (i.e. cataracts) are observed in most affected individuals [9,10].
187 Nevertheless, a patient with a heterozygous p.(Val3_Gly4ins2) mutation and no cataracts at
188 age 52 has been reported by Grau and colleagues [10]. Only one of the seven patients
189 reported here had no clinical history of cataract: subject C3 was examined at age 38 and no
190 lens opacity was detected. It is worth highlighting that a number of patients with *OPA3*-
191 related disease presented with bilateral lens opacities at birth or in early childhood (including
192 subject M4 who was only noted to have optic atrophy after lens extraction at age 15);
193 therefore, *OPA3* mutations should be in the differential in cases of bilateral infantile-onset
194 cataract and it is important to assess optic nerve function and health before concluding about
195 the likely amblyogenicity of the lens opacities.

196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229

Ocular *OPA3*-related disease often occurs in parallel with extraocular clinical features. Reynier and colleagues reported affected members of a family with a heterozygous p.(Gly93Ser) change in *OPA3* to have mild neurological signs including spasticity and extrapyramidal dysfunction [9]. Subsequently, Grau and colleagues reported four individuals with a heterozygous p.(Val3_Gly4ins2) change to have hearing loss. Expression analysis of *OPA3* in murine cochlear tissue was consistent with the notion that auditory neuropathy can be an extraocular feature of *OPA3*-related disease [10]. Notably, two affected individuals from the present cohort had sensorineural hearing loss (Table 1); hearing impairment has been previously described in a patient with the same genetic defect [10]. Clinicians should be vigilant to the development of such a complication and audiograms should be performed in all patients with *OPA3*-related disease at least once.

All *OPA3* families with dominant disease reported to date have affected individuals in two or more generations and disease-associated variants appear to be highly penetrant (Figure 1A-B, [9,10]). The p.(Gln105Glu) change identified in both families reported here has been previously shown to be a recurrent mutation [10]; it was not possible to perform haplotype analysis in subjects from the present cohort and thus, the possibility of a recent common ancestor in the two families cannot be excluded. The p.(Gln105Glu) change affects an amino acid in a predicted coiled-coil structure (Uniprot) but it is unclear how this common cause of *OPA3*-related disease affects mitochondrial function and/or structure.

Findings from this and other studies [9,10] suggest that dominant *OPA3*-related disease is associated with significant intra- and inter-familial phenotypic variability. Any attempt to draw genotype-phenotype correlations would therefore be highly speculative and secondary genetic factors are likely to be the basis for the observed clinical heterogeneity; a similar conclusion has been reached for the much commoner *OPA1*-associated optic neuropathy [14]. Expansion of the phenotypic spectrum of *OPA3*-related disease has led to greater understanding of the condition and the risks of developing extraocular complications. However, it is still unclear what disease mechanism explains multi-system tissue involvement and future studies on the pathogenesis of the disorder are expected to provide important insights.

230 **ACKNOWLEDGEMENTS**

231

232 Supported by the Medical Research Council UK (Grant G0500790), the Manchester

233 Biomedical Research Centre and NIHR Greater Manchester: Clinical Research Network.

234

235 **REFERENCES**

236

- 237 1. Votruba M (2004). Molecular genetic basis of primary inherited optic neuropathies. *Eye*
238 (*Lond*) 18:1126-1132.
- 239 2. Yu-Wai-Man P, Griffiths PG, Chinnery PF (2011). Mitochondrial optic neuropathies -
240 disease mechanisms and therapeutic strategies. *Prog Retin Eye Res* 30:81-114.
- 241 3. Costeff H, Gadoth N, Apter N, Prialnic M, Savir H (1989). A familial syndrome of
242 infantile optic atrophy, movement disorder, and spastic paraplegia. *Neurology* 39:595-597.
- 243 4. Anikster Y, Kleta R, Shaag A, Gahl WA, Elpeleg O (2001). Type III 3-methylglutaconic
244 aciduria (optic atrophy plus syndrome, or Costeff optic atrophy syndrome): identification of
245 the OPA3 gene and its founder mutation in Iraqi Jews. *Am J Hum Genet* 69:1218-24.
- 246 5. Kleta R, Skovby F, Christensen E, Rosenberg T, Gahl WA, Anikster Y (2002). 3-
247 Methylglutaconic aciduria type III in a non-Iraqi-Jewish kindred: clinical and molecular
248 findings. *Mol Genet Metab* 76:201-206.
- 249 6. Ho G, Walter JH, Christodoulou J (2008). Costeff optic atrophy syndrome: new clinical
250 case and novel molecular findings. *J Inherit Metab Dis* 31:S419-423.
- 251 7. Arif B, Kumar KR, Seibler P, Vulinovic F, Fatima A, Winkler S, Nürnberg G, Thiele H,
252 Nürnberg P, Jamil AZ, Brüggemann A, Abbas G, Klein C, Naz S, Lohmann K (2013). A
253 novel OPA3 mutation revealed by exome sequencing: an example of reverse phenotyping.
254 *JAMA Neurol* 70:783-787.
- 255 8. Wortmann SB, Duran M, Anikster Y, Barth PG, Sperl W, Zschocke J, Morava E, Wevers
256 R (2013). Inborn errors of metabolism with 3-methylglutaconic aciduria as discriminative
257 feature: proper classification and nomenclature. *J Inherit Metab Dis* 36:923-928.
- 258 9. Reynier P, Amati-Bonneau P, Verny C, Olichon A, Simard G, Guichet A, Bonnemains C,
259 Malecaze F, Malinge MC, Pelletier JB, Calvas P, Dollfus H, Belenguer P, Malthièry Y,
260 Lenaers G, Bonneau D (2004). OPA3 gene mutations responsible for autosomal dominant
261 optic atrophy and cataract. *J Med Genet* 41:e110
- 262 10. Grau T, Burbulla LF, Engl G, Delettre C, Delprat B, Oexle K, Leo-Kottler B, Roscioli T,
263 Krüger R, Rapaport D, Wissinger B, Schimpf-Linzenbold S (2013). A novel heterozygous
264 OPA3 mutation located in the mitochondrial target sequence results in altered steady-state
265 levels and fragmented mitochondrial network. *J Med Genet* 50:848-858.
- 266 11. Huizing M, Dorward H, Ly L, Klootwijk E, Kleta R, Skovby F, Pei W, Feldman B, Gahl
267 WA, Anikster Y 2010. OPA3, mutated in 3-methylglutaconic aciduria type III, encodes two
268 transcripts targeted primarily to mitochondria. *Mol Genet Metab* 100:149-154.

- 269 12. Ryu SW, Jeong HJ, Choi M, Karbowski M, Choi C (2010). Optic atrophy 3 as a protein
270 of the mitochondrial outer membrane induces mitochondrial fragmentation. *Cell Mol Life Sci*
271 67:2839-2850.
- 272 13. Yu-Wai-Man P, Shankar SP, Biousse V, Miller NR, Bean LJ, Coffee B, Hegde M,
273 Newman NJ (2011). Genetic screening for OPA1 and OPA3 mutations in patients with
274 suspected inherited optic neuropathies. *Ophthalmology* 118, 558-563.
- 275 14. Yu-Wai-Man P, Griffiths PG, Burke A, Sellar PW, Clarke MP, Gnanaraj L, Ah-Kine D,
276 Hudson G, Czermin B, Taylor RW, Horvath R, Chinnery PF (2010). The prevalence and
277 natural history of dominant optic atrophy due to OPA1 mutations. *Ophthalmology* 117, 1538-
278 1546.
- 279
280
281
282

Table 1. Clinical and genetic findings in subjects with dominant *OPA3*-related disease reported here and elsewhere.

Subject (family ID)	Age; sex	VA (logMAR) right/left	Presentation/onset	Lens	Heterozygous <i>OPA3</i> change identified	Other features
C1 (GCC1)	72; M	1.30/1.30	Nystagmus in infancy	Blue-dot cataracts diagnosed at age 2	p.(Gln105Glu)	-
C2 (GCC1)	62; F	1.00/1.00	Nystagmus in infancy	Cataracts operated at age 4	p.(Gln105Glu)	-
C3 (GCC1)	38; F	0.80/0.80	Nystagmus in infancy	No significant lens opacity at age 38	p.(Gln105Glu)	-
M1 (G43755)	65; F	0.72/1.04	Cataracts at age 34; re-referred as found by optometrist to have cupped left disc at age 60	Cataracts operated at age 34; bilateral aphakia	p.(Gln105Glu)	Right mild-moderate s/n hearing loss; left moderate-severe s/n hearing loss; bilateral ocular hypertension on topical treatment
M2 (G43755)	41; M	0.10/0.20	Mild cataracts at age 9; presently asymptomatic	Nuclear cataracts operated at age 35	p.(Gln105Glu)	Right and left mild-moderate s/n hearing loss
M3 (G43755)	36; F	0.60/0.30	Decrease in VA from age 19	Blue-dot cataracts operated at age 36&37	p.(Gln105Glu)	Normal audiogram; familial hypercholesterolaemia
M4 (G43755)	13; F	0.08/0.18	Cataracts at age 13	Lamellar cataracts operated at age 15	p.(Gln105Glu)	Congenital AV malformation in frontal lobe
III.3 (#1) [9]	38; F	0.70/0.70	Poor vision from infancy	Posterior cortical cataracts operated at age 51	p.(Gly93Ser)	Tremor of hands; extrapyramidal rigidity of upper limbs; absence of deep tendon reflexes
IV.1 (#1) [9]	15; F	0.30/0.30	Decreased VA before age 10	Posterior cortical cataracts operated at age 47&48	p.(Gly93Ser)	Mild postural tremor & mild rigidity of upper extremities
IV.2 (#1) [9]	57; F	0.60/0.50	Decreased VA before age 10	Anterior cortical cataracts operated at age 45&46	p.(Gly93Ser)	Postural tremor without extrapyramidal signs
V.1 (#1) [9]	29; F	0.70/0.52	Decrease in VA from age 12	Anterior cortical cataracts operated at age 25	p.(Gly93Ser)	Normal neurological examination
VI.1 (#1) [9]	4; F	0.15/0.15	Visual impairment investigated at age 3	Anterior & posterior cortical cataracts operated at age 4	p.(Gly93Ser)	Normal neurological examination
IV.6 (#1) [9]	50; F	NA	Visual impairment from infancy	Cataracts	p.(Gly93Ser)	Normal neurological examination
V.6 (#1) [9]	5; F	NA	Visual impairment from infancy	Cataracts operated at age 5	p.(Gly93Ser)	Normal neurological examination
III.3 (#2) [9]	37; F	1.70/1.70	Decrease in VA from age 12	Posterior capsular cataract diagnosed at age 56	p.(Gln105Glu)	-
III.7 (#2) [9]	49; F	“legally blind”	Decrease in VA from age 10	Cataracts diagnosed at age 45	p.(Gln105Glu)	-
IV.1 (#2) [9]	33; F	1.30/1.30	Decrease in VA from age 6	Cataracts diagnosed at age 10; cerulean cataracts at age 33	p.(Gln105Glu)	Normal neurological examination

IV.2 (#2) [9]	12; M	0.52/0.52	Decreased VA at age 12	Posterior capsular cataracts operated at age 19	p.(Gln105Glu)	-
II.2 (OAK1) [10]	57	1.00/1.00	Diagnosed at age 18	Monocular cataract	p.(Gln105Glu)	-
II.6 (OAK1) [10]	54	0.40/0.50	Onset in infancy	Cataracts	p.(Gln105Glu)	Hearing loss
III.1 (OAK1) [10]	35	0.52/0.30	Onset in infancy	No significant lens opacity at age 19	p.(Gln105Glu)	-
III.2 (OAK1) [10]	31	0.40/0.52	Onset in infancy	Cataracts	p.(Gln105Glu)	-
III.3 (OAK1) [10]	17	0.52/0.40	Onset in infancy	Cataracts	p.(Gln105Glu)	Chiari malformation type I
I.1 (OAK61) [10]	57	1.30/1.40	Diagnosed at age 10	No information	p.(Gln105Glu)	-
II.1 (OAK61) [10]	33	0.40/0.40	Diagnosed at age 19	No significant lens opacity at age 25	p.(Gln105Glu)	-
II.4 (OAK105) [10]	84	1.15/1.30	Onset in infancy	Congenital cataracts	p.(Val3_Gly4ins 2)	Hearing loss
III.2 (OAK105) [10]	61	0.30/0.40	Onset in adolescence	No significant lens opacity at age 52	p.(Val3_Gly4ins 2)	Hearing loss
III.3 (OAK105) [10]	62	0.52/0.40	Diagnosed at age 19	Cataracts	p.(Val3_Gly4ins 2)	Hearing loss
III.5 (OAK105) [10]	57	1.15/1.15	Onset in adolescence	Congenital cataracts	p.(Val3_Gly4ins 2)	Hearing loss
IV.1 (OAK105) [10]	32	1.30/1.30	Diagnosed at age 19	Congenital cataracts	p.(Val3_Gly4ins 2)	Morbus Scheuermann
III.1 (OAK255) [10]	46	NA	Onset in infancy	Cataracts	p.(Gln105Glu)	Intestinal pseudo- obstruction
III.2 (OAK255) [10]	47	NA	Onset in infancy	Cataracts	p.(Gln105Glu)	-

F, female; M, male; VA, visual acuity; logMAR, logarithm of the minimum angle of resolution (equivalent); BE, both eyes; NA, not available; s/n, sensorineural; AV, arteriovenous. The term cataracts is used to denote bilateral crystalline lens opacities. [Variants are annotated according to the NCBI Reference Sequence NM_025136.3 \(OPA3 isoform b\).](#)

285

286

287 **FIGURE LEGEND**

288

289 **Figure 1.**

290 **A.** Pedigree from a family segregating optic atrophy. Subjects C1 (IV:2), C2 (IV:5) and C3
291 (V:3) had a heterozygous *OPA3* change, c.313C>G, p.(Gln105Glu).

292 **B.** Pedigree from a family segregating optic atrophy and cataracts. DNA from subjects M1
293 (II:6), M2 (III:3) and M3 (III:4) was available for testing; a heterozygous c.313C>G,
294 p.(Gln105Glu) change in *OPA3* was identified in all these patients. See text and Table 1 for
295 clinical findings of all tested individuals as well as subject M4 (IV:2).

296 **C.** Right and left optic disc appearance of an individual with a confirmed *OPA3* mutation
297 (subject M1) showing pallor of the neuroretinal rim, which is more marked temporally. Left
298 optic nerve appears more affected than the right and has optic disc excavation in addition to
299 the temporal pallor.

300 **D.** Left optic disc appearance and pattern of retinal nerve fibre layer (RNFL) thinning in an
301 individual with a confirmed *OPA3* mutation (subject M2). Sparing of the nasal, superior and
302 inferior peripapillary quadrants is observed. The RNFL profile for each eye is superimposed
303 on the normal distribution percentiles. The normal distribution indices are colour-coded: (i)
304 red <1%, (ii) yellow <5%, (iii) green <95%, and (iv) white >95%.

305