

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/70609/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Meyer, Jan-Philip, Probst, Katrin C., Trist, Iuni Margaret, McGuigan, Christopher and Westwell, Andrew David 2014. A novel radiochemical approach to 1-(2'-deoxy-2'-[F-18] fluoro-beta-D-arabinofuranosyl)cytosine (F-18-FAC). *Journal of Labelled Compounds and Radiopharmaceuticals* 57 (11), pp. 637-644. 10.1002/jlcr.3233

Publishers page: <http://dx.doi.org/10.1002/jlcr.3233>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# A novel radiochemical approach to 1-(2'-deoxy-2'-<sup>18</sup>F)fluoro-β-D-arabinofuranosyl)cytosine (<sup>18</sup>F-FAC)

Jan-Philip Meyer<sup>[a,b]</sup>, Katrin C. Probst<sup>[b]</sup>, Iuni M. L. Trist<sup>[a]</sup>, Christopher McGuigan<sup>[a]</sup>, Andrew D. Westwell<sup>[a]\*</sup>

<sup>a</sup>*School of Pharmacy and Pharmaceutical Sciences, Cardiff University, Redwood Building, King Edward VII Avenue, Cardiff, CF10 3NB, Wales, U.K.;*

<sup>b</sup>*Wales Research & Diagnostic PET Imaging Centre (PETIC), Institute for Translation, Innovation, Methodology & Engagement (TIME), School of Medicine, Heath Park, Cardiff University, Cardiff, CF14 4XN, Wales, U.K.*

## Abstract

<sup>18</sup>F-FAC (1-(2'-deoxy-2'-<sup>18</sup>F)fluoro-β-D-arabinofuranosyl)-cytosine) is an important 2'-fluoro-nucleoside-based PET tracer that has been used for *in vivo* prediction of response to the widely used cancer chemotherapy drug gemcitabine. Previously reported synthetic routes to <sup>18</sup>F-FAC have relied on early introduction of the <sup>18</sup>F radiolabel prior to attachment to protected cytosine base. Considering the <sup>18</sup>F radiochemical half-life (110 minutes) and the technical challenges of multi-step syntheses on PET radiochemistry modular systems, late-stage radiofluorination is preferred for reproducible and reliable radiosynthesis with *in vivo* applications. Herein, we report the first late-stage radiosynthesis of <sup>18</sup>F-FAC. Cytidine derivatives with leaving groups at the 2'-position are particularly prone to undergo anhydro side product formation upon heating due to their electron density at the 2-carbonyl pyrimidone oxygen. Our rationally developed fluorination precursor showed an improved reactivity-to-stability ratio at elevated temperatures. <sup>18</sup>F-FAC was obtained in radiochemical yields (RCY) of 4.3-5.5% (n = 8, decay-corrected from end of bombardment (EoB)), with purities ≥98% and specific activities ≥63 GBq/μmol. The synthesis time was 168 min.

**Keywords:** nucleosides • Positron Emission Tomography • radiochemistry • radiopharmaceuticals • computational chemistry

## Introduction

Nucleoside derivatives represent an important emerging class of biomarkers for Positron Emission Tomography (PET) imaging.<sup>[1]</sup> For example, the clinically approved proliferation biomarker <sup>18</sup>F-FLT (3'-deoxy-3'-<sup>18</sup>F)fluoro-*L*-thymidine) is accessible via an effective

---


\* Correspondence to: Andrew D. Westwell, School of Pharmacy and Pharmaceutical Sciences, Cardiff University, Redwood Building, King Edward VII Avenue, Cardiff, CF10 3NB, Wales, U.K. E-mail: [WestwellA@cf.ac.uk](mailto:WestwellA@cf.ac.uk)

radiochemical synthesis in high radiochemical yield (RCY) and purity suitable for human production under GMP guidelines.<sup>[2]</sup> Hence, <sup>18</sup>F-FLT is now commonly used for the early detection of various cancers and the evaluation of treatment response.<sup>[3]</sup> Especially in high-grade tumours it appears to be more powerful in predicting treatment response and survival than <sup>18</sup>F-FDG, as measured by standard uptake values of <sup>18</sup>F-FLT directly corresponding to cell proliferation shown by the Ki-67 proliferation index.<sup>[4]</sup>

Cytidine-based radiotracers for PET imaging such as <sup>18</sup>F-FAC (**1**, Figure 1) and 1-(2'-deoxy-2'-[<sup>18</sup>F]fluoroarabinofuranosyl)-5-methylcytosine (<sup>18</sup>F-FMAC, **2**, Figure 1) have been previously shown to be successful as predictors of response for nucleoside analogue based treatment of cancer<sup>[5,6]</sup> as well as local immune activation in mouse models.<sup>[7]</sup> As such, <sup>18</sup>F-FAC has been used for response prediction to the widely used chemotherapy drug gemcitabine.<sup>[5]</sup> Gemcitabine is frequently used for (combination) chemotherapy of a number of solid tumors such as pancreatic, non-small lung, breast and bladder cancer. Importantly, gemcitabine efficacy is compromised by poor cellular uptake by nucleoside transporters, variable activity of the essential deoxycytidine kinase (dCK) activating enzyme, and rapid deactivating metabolism by cytidine deaminase (CDA). In addition, serious dose-limiting toxicities are associated with this chemotherapy drug.<sup>[8]</sup> Prediction of patient response to gemcitabine chemotherapy would therefore be an important milestone towards personalised patient chemotherapy in the cancer clinic.

Given the close structural resemblance to gemcitabine, <sup>18</sup>F-FAC was found to have a comparably high affinity for deoxycytidine kinase (dCK), a rate-limiting enzyme in the nucleoside salvage pathway. Hence, *in vivo* imaging determined dCK-positive and -negative tumours. Furthermore, an additional study showed that <sup>18</sup>F-FMAC PET imaging provides information about cytidine deaminase (CDA) enzymatic activity *in vivo* which is related to gemcitabine resistance.<sup>[6]</sup> Together, cytidine-based PET probes enable further conclusions towards personalised chemotherapy which, considering the low ( $\leq 20\%$ ) response rates and severe side effects, could lead to significantly improved treatment efficacy.<sup>[8]</sup>

2'-[<sup>18</sup>F]arabincytidines such as <sup>18</sup>F-FAC and <sup>18</sup>F-FIAC (**3**, Figure 1) have been synthesized in high purities using an early stage <sup>18</sup>F introduction strategy.<sup>[9]</sup> However, this synthetic approach limits reliable and reproducible production on commercially available synthesis modules for human application.<sup>[10]</sup> For instance, multiple steps after <sup>18</sup>F introduction with an additional purification step due to the formation of the D- and L-isomers, as well as the use of toxic substances such as hydrogen bromide, make the translation to automated synthesizers difficult. These challenges go hand in hand with reproducibility issues when it comes to Quality Control (QC) in clinical practice according to the British/European/American Pharmacopeia (BPh/Ph. Eur./USP). Furthermore, as the number and space in hot cells is limited multiple-step radiosyntheses should preferably be replaced by shorter and less complex synthetic procedures. Lazari et. al. recently presented a newly designed radiosynthesis module system which furnished the nucleoside analogues <sup>18</sup>F-FMAU and <sup>18</sup>F-FAC in good overall radiochemical yields (46% and 31% respectively, decay-corrected) and with acceptable reproducibility following early stage fluoride introduction.<sup>[11]</sup> Overall synthesis times were reported to be 165-170 minutes in each case. The flexible radio-synthesizers developed are described as being suitable for both the development and production of different radiotracers for applications in PET.



**Figure 1.** Chemical structures of the cytidine-based biomarkers  $^{18}\text{F}$ -FAC,  $^{18}\text{F}$ -FMAC and  $^{18}\text{F}$ -FIAC (**1-3**), and the anticancer drug gemcitabine.

The aim of this study was to develop a novel and straightforward synthetic approach towards  $^{18}\text{F}$ -FAC that could be applied to the radiosynthesis of related cytidine-based radiotracers. A more straightforward radiosynthesis similar to the  $^{18}\text{F}$ -FDG synthesis in combination with innovative radio-synthesizers would enable a faster transition from tracer development to clinical application. Late-stage  $^{18}\text{F}$ -introduction at the 2'-arabino position of an intact nucleoside towards nucleoside-based radiotracers was previously considered as difficult.<sup>[12]</sup> However, Alauddin and co-workers recently developed a novel radiosynthesis of the uridine analogue  $^{18}\text{F}$ -FMAU based on a late-stage  $^{18}\text{F}$ -introduction approach in order to increase possibilities for making nucleoside based PET tracer suitable for clinical applications.<sup>[10,12]</sup> The key towards a stable precursor that would be less likely to undergo anhydro side product formation (e.g. Scheme 1) due to neighbouring group participation was the introduction of a good electron withdrawing N-protecting group(s). Since the amino-group at the 4-position of cytidine derivatives likely enhances the nucleophilicity of the 2-carbonyl oxygen, potential precursor molecules containing a leaving group at the 2'-position are even more prone to undergo anhydro formation especially at high temperatures ( $\geq 75\text{ }^\circ\text{C}$ ). Thus, a reasonable radiofluorination precursor must combine sufficient reactivity with stability towards heat as well as basic conditions. In this study we present for the first time a late-stage radiofluorination approach towards the  $^{18}\text{F}$ -labelled cytidine derivative  $^{18}\text{F}$ -FAC.

## Experimental Section

### Radiosynthetic procedures

*N*<sup>4</sup>-bis-Boc-3',5'-O-bis-tetrahydropyranyl-2'-deoxy-2'-[ $^{18}\text{F}$ ]fluoroarabincytidine (**25**). [ $^{18}\text{F}$ ]fluoride was trapped on a QMA cartridge (equilibrated with 5%  $\text{NaHCO}_3$ -solution (5 mL) and water (5 mL)) before it was eluted with an aqueous solution (0.7 mL) of  $\text{KHCO}_3$  (22.6 mg/mL) and kryptofix  $\text{K}_{222}$  (22.6 mg/mL) in MeCN. The resulting  $^{18}\text{F}/\text{KHCO}_3/\text{K}_{222}$ -complex was dried by co-evaporation under reduced pressure and a stream of nitrogen. The drying process was repeated 3 times ( $3 \times 1\text{ mL MeCN}$ ). A solution of precursor (**23**) (10 mg,

14.5  $\mu\text{mol}$ ) in DMF (0.3 mL) was then added and the resulting mixture was stirred at 110 °C for 20 min. The crude mixture was either analyzed directly for determination of  $^{18}\text{F}$ -incorporation (radio-TLC) or passed through an Alumina-Sep-Pak cartridge (equilibrated with water (5 mL)) for characterization (radio-HPLC). The  $^{18}\text{F}$ -labelled intermediate was eluted with EtOAc (2 mL) and an aliquot (0.2 mL) of the mixture was analyzed by analytical HPLC using co-elution with the non-radioactive reference compound (**24**).  $R_t = 17.2$  min, MeCN/H<sub>2</sub>O 9:1, 1 mL min<sup>-1</sup>,  $\geq 87\%$ . Radio-TLC.  $R_f = 0.6$  (4% MeOH/DCM).

*<sup>18</sup>F-FAC (1)*. After elution of the  $^{18}\text{F}$ -labelled intermediate (**25**) with EtOAc into a new reaction V-vial the solvent was removed under reduced pressure and a stream of nitrogen at 80 °C. 2N HCl-solution (0.3 mL) was added to the dried residue followed by stirring at 90 °C for 15 min. The solution was then neutralized with 2N NaOH solution (0.3 mL). The resulting mixture was loaded onto a semi-preparative HPLC column via a 10 mL injection loop. The fraction containing  $^{18}\text{F}$ -FAC was collected after 22.5 min with 3% MeCN in H<sub>2</sub>O and a flow rate of 3.5 mL min<sup>-1</sup>. The appropriate fraction was flushed through a sterile filter and concentrated via evaporation on the module (100 °C under a stream of nitrogen for 10 min.). The residue was taken up in sterile saline. An aliquot (0.2 mL) of the mixture was analyzed by analytical HPLC using co-elution with the non-radioactive reference compound **27**,  $R_t = 13.8$  min, 3% MeCN/H<sub>2</sub>O, 1 mL min<sup>-1</sup>,  $\geq 98\%$ . A carrier-added synthesis was performed according to the above procedure and the product  $^{19}\text{F}/^{18}\text{F}$ -FAC was saved for decay for 20 half-lives ( $\approx 2$  days) and subsequently analyzed by mass spectrometry; MS (70 eV):  $m/z$  (%): 245.4 (5) [(M+H)<sup>+</sup>], 267.7 (95) [(M+Na)<sup>+</sup>].

### Computational Methods


MOE 2010.10 software was used to design each compound and to explore possible conformations. The searching method applied was LowModMD that generates conformations using a short run of Molecular Dynamics (MD) at constant temperature followed by an all-atom energy minimization. Among the conformations, sixteen of them were selected for each compound for quantum mechanics (QM) evaluation. QM calculations were executed for the selected conformations with the GAMESS<sup>[13]</sup> software. A geometry optimisation was performed *in vacuo*, using analytical energy gradients with the Restricted Hartree Fock (RHF) wave function. The 3-21G split valence basis set was used and diffuse sp shell was added to heavy atoms. The initial hessian was guessed and the convergence gradient tolerance was set to 0.0005 Hartree/Bohr. The electrostatic potential was calculated at points determined by Michael Connolly algorithm on the surface of van der Waals fused spheres. Atomic charges were consequently fitted to this potential. The partial net charges of atoms of interest were gathered from each molecule conformation and values were averaged.

## Results and Discussion

### *Precursor design*

In order to develop a new precursor for radiofluorination of the 2'-arabino position of cytidine derivatives the idea of radiofluorination of an intact nucleoside moiety was applied. The main challenge of this new approach is the rational design of precursors that combine reactivity and stability in a way that the  $^{18}\text{F}$ -nucleophile can be incorporated at high temperatures resulting in acceptable conversion rates while heat-mediated anhydro-formation is kept to a minimum. A quantitative method for the characterization of precursor stabilities was desirable prior to synthesis. The electric charge density of the 2-carbonyl oxygen of the pyrimidone moiety was determined using theoretical methods for the potential to undergo an intramolecular nucleophilic attack at the 2'-position. Hence, the net electric charge of the 2-carbonyl oxygen of a family containing  $\text{N}^4$ -substituted cytidine derivatives (**4-8**, Figure 2) was calculated using quantum chemical calculations with Restricted Hartree Fock (RHF) wave function and the 3-21G basis set.

Additionally, selected uridine derivatives (**9-11**, Figure 2) were included in this study in order to obtain both a reference value as well as a measurement of how the substitution of the 4-carbonyl group (uridines) by an amino group (cytidines) impacts the electric charge density at the 2-carbonyl oxygen. The N-substituents were chosen with regard to both their synthetic accessibility and deprotection mechanisms that would be suitable for radiochemical syntheses in terms of subsequent efficient removal. The  $\text{N}^3$ -nitro-2'-(methanesulfonyl)uridine (**11**) was considered as a standard value as a  $\text{N}^3$ -nitro-2'-(trifluoromethanesulfonyl)uridine derivative was previously reported as successfully synthesized and thermal stable.<sup>[14]</sup> However, the radiofluorination method could not be applied to a  $\text{N}^3$ -nitro-uridine-based precursor.




**Figure 2.** Selected cytidine and uridine based nucleosides for electric charge density calculations at the 2-carbonyl oxygen.

The results shown below (Table 1) indicate that the electron density at the 2-carbonyl oxygen is reduced when H-substitution at the N<sup>4</sup> (N<sup>3</sup> for uridine derivatives) with an electron-withdrawing group is performed. Furthermore, it shows that the stronger the electron-withdrawing effect of the substituent(s) the lower the calculated electron density  $\Omega$  at the 2-carbonyl oxygen. For instance, double N<sup>4</sup>-Boc-protection (**7**) decreases the electron density more than single N<sup>4</sup>-Boc-protection (**6**) and N<sup>4</sup>-monoacetylation (**5**), respectively.


**Table 1.** Calculated electron density  $\Omega$  of selected cytidine (**4-8**) and uridine derivatives (**9-11**).

Compounds **5** and **7** were synthesized (see synthesis section and Supporting Information) and subsequently tested for their affinity to undergo heat-mediated decomposition by forming the appropriate anhydro-compounds **12** and **13**. First, a NMR sample of compound **5** in d<sub>6</sub>-DMSO was heated up to 50 °C. <sup>1</sup>H-NMR spectra were recorded hourly. Figure 3 shows that almost full conversion to anhydro compound **12** occurred after 2h (Scheme 1).

Chemical shifts of the amino proton (11.5 to 12.25 ppm) and the ribose protons clearly show the conversion into the cyclic nucleoside compound (Figure 3). Furthermore, the vanishing signal of the mesyl group at 3.5 ppm also indicates decomposition as the free mesylate is expected to have a different chemical shift that indeed could be observed at around 2.3 ppm next to the methyl-signal of the acetamide. <sup>13</sup>C-NMR data after 1h incubation showed 24 instead of 12 signals of pure compound **5**. Additionally, ESI-MS was performed in order to show the change in the molecular mass. The m/z-signal 268 [M + H]<sup>+</sup> further confirmed generation of anhydro compound **12**.




**Scheme 1.** Heat-mediated intramolecular cyclization towards **12** in d<sub>6</sub>-DMSO.




**Figure 3.** Heat-mediated anhydro formation of **12**. After only 2 hours at 50 °C nearly full conversion could be observed by  $^1\text{H}$  NMR analysis.

The double  $\text{N}^4$ -Boc-protected compound (**7**), however, showed significantly increased thermal stability as shown in Scheme 2 and Figure 4. Even though a tiny set of additional peaks appeared after 2 hours at 80 °C the observed stability of compound **7** looked promising with regard to  $^{18}\text{F}$ -radiolabeling reactions.



**Scheme 2.** Increased thermal stability of **7**. No significant formation of side product (**13**) could be observed (in  $\text{d}_4\text{-MeOD}$ ).






**Figure 4.** Recorded <sup>1</sup>H-NMR spectra of the purified compound (**7**) after 0, 60 and 120 min at 80 °C. The absence of additional peaks indicate no or minimal anhydro formation through increased thermal stability.

Hence, the calculated and experimental data lead to 3 major conclusions: 1. The electron density at the 2-carbonyl oxygen of the pyrimidine moiety can be used as a measurement for its ability to undergo intramolecular cyclization resulting in the appropriate 2,2'-anhydro compounds. 2. Electron withdrawing groups have a measurable/quantifiable effect on the electron density at the 2-carbonyl oxygen. 3. Depending on the strength of its electron withdrawing nature a N<sup>4</sup>(N<sup>3</sup>)-protecting group influences the thermal stability of the nucleoside analogue. Finally, an N,N-diBoc fluorination precursor was selected for radiofluorination studies.

### Synthesis

The synthesis of test compound **5** was carried out with the selective 3'- and 5'-protection of N-acetylcytidine (**14**) using TIPDSCl<sub>2</sub> in pyridine yielding compound **15** in good yield which was consecutively mesylated at the 2'-position using an excess of MsCl and TEA in DCM furnishing compound **16**. Subsequent removal of the silica-based protecting group using TBAF in anhydrous THF at low temperatures gave test compound **5** which was purified immediately and characterized due to its high ability to undergo anhydro formation (Scheme 3).




**Scheme 3.** Reagents and conditions: a) TIPDSCl<sub>2</sub>, pyridine, 0 °C – r.t., 12 h, 87 %; b) MsCl, TEA, DCM, 0 °C – r.t., 2 h, 81 %; c) TBAF, THF, -10 °C, 30 min, 53 %.

The synthesis of precursor (**23**) started with commercially available cytidine (**17**) which was first converted into the 3'- and 5'- protected nucleoside **18** using TIPDSCl<sub>2</sub> and pyridine at room temperature as previously described.<sup>[8]</sup> Subsequent TMS-protection furnished **19** which was followed by double N<sup>4</sup>-Boc protection using excess of Boc<sub>2</sub>O (5-6 eq.) and DMAP which gave compound **20** in good yield. Selective removal of the TMS-group using *p*-TsOH was performed at -5 °C to give intermediate **21**. Subsequent mesylation of the free 2'-hydroxy group was performed in high yield to form compound **22**. TBAF-mediated removal of the bidendate protecting group had to be carried out at lower temperatures (~ -10 °C) in order to avoid significant decrease in yield probably due to single Boc-group removal. Purified compound **7** was then converted into fluorination precursor **23** using 3,4-dihydro-6H-pyran in excess. Highly purified precursor **23** was obtained in 9% overall yield from **17** (Scheme 4).

The choice of the leaving group as well as the protecting group is based on a previous study<sup>[10]</sup> that found that 3'/5'-THP-protection of uridine derivatives gave the best radiochemical yields and a 2'-mesyl group an appropriate balance between stability and leaving group abilities. Furthermore, our pilot studies showed that the appropriate tosyl- and


nosyl-containing counterparts already decomposed during purification. Furthermore, even though double THP-protection makes NMR analysis complex (mixture of diastereomers) the later removal of these groups was straightforward.



**Scheme 4.** Reaction conditions: a) TIPDSCl<sub>2</sub>, pyridine, 0 °C – r.t., 18 h, 81 %; b) TMSCl, TEA, DCM, 0 °C – r.t., 2 h, 83 %; c) Boc<sub>2</sub>O, DMAP, DCM, r.t., 5 h, 77 %; d) *p*-TsOH, THF, -5 °C, 66-78 %; e) MsCl, TEA, DCM, 0 °C, 90 %; f) TBAF, THF, -10 °C, 70-80 %, g) DHP, *p*TsOH, DCM, 0 °C - RT, 6-8 h, 45-56 %.

Firstly, cold fluorination reactions were carried out to assess fluoride incorporation for precursor (**23**). Amongst fluorination conditions tested the best incorporation yield was found at 100 °C for 100 min. in DMF using freshly prepared 1N TBAF/THF-solution (Scheme 5). The fluorinated intermediate (**24**) was purified by column chromatography on SiO<sub>2</sub> and its

identity and purity ( $\geq 97\%$ ) confirmed by  $^1\text{H}/^{19}\text{F}$ -NMR, HR-MS and analytical HPLC (see Supporting Information). Notably, the  $^{19}\text{F}$ -spectrum showed a set of 4 signals due to the 4 diastereomers that the THP-protection yields.




**Scheme 5.** Reagents and conditions: a) 1N TBAF/THF, DMF, 100 °C, 100 min, 19 %.

The obtained intermediate **24** was used as a cold reference standard for co-elution on HPLC in order to identify the hot intermediate.


### Radiosynthesis of $^{18}\text{F}$ -FAC

$^{18}\text{F}$ -incorporation to form intermediate (**25**) was carried out using an Eckert & Ziegler modular system and appropriate software for radiosynthesis. The reaction parameters of solvent, temperature and reaction time were varied in order to determine optimal  $^{18}\text{F}$ -incorporation conditions, using  $\text{KHCO}_3$  as base (Scheme 6).  $[^{18}\text{F}]$ Fluoride was eluted from the QMA-cartridge using a  $\text{K}_{222}/\text{KHCO}_3$ -solution. Table 2 shows the calculated incorporation data, and Figure 5 shows  $^{18}\text{F}$ -incorporation plotted against the reaction temperature run in dry DMF as solvent. The highest incorporation values were obtained using DMF at 110 °C for 20 minutes. Although MeCN or *t*BuOH as solvent failed to show any conversion, use of an alternative base such as TBA-OH or TBA- $\text{CO}_3$  in MeCN or *t*BuOH might provide an alternative strategy. However, higher temperatures and longer reaction times led to decreased  $^{18}\text{F}$ -incorporation in the arabino position due to the formation of the anhydro intermediate (Scheme 2).  $^{18}\text{F}$ -incorporation was calculated by peak integration of the crude reaction mixture after radiofluorination using radio-TLC (see Supporting Information).




**Scheme 6.** Radiofluorination was performed using different reaction conditions as outlined in the table below.

**Table 2.** [ $^{18}\text{F}$ ]-incorporation results obtained under different conditions.




**Figure 5.** [ $^{18}\text{F}$ ]-incorporation plotted against the reaction temperature run in dry DMF as solvent.

The  $^{18}\text{F}$ -fluorinated intermediate (**25**) was confirmed by radio-HPLC (Figure 6) and co-injection of the  $^{19}\text{F}$ -containing reference standard (**24**) (see Supporting Information) after trapping the majority of the unreacted  $^{18}\text{F}$ -fluoride on a Sep-Pak alumina cartridge. The intermediate was eluted with 2.5 mL EtOAc into a new reaction vial.




**Figure 6.** Radio-HPLC was performed in order to confirm the formation of **25**. The fluoridation product was co-eluted with reference compound **24** after 17.2 min (MeCN/H<sub>2</sub>O 10:1, flow rate 1 mL min<sup>-1</sup>). The profile of the radio-HPLC diagram is consistent with the radio-TLC diagram shown above.

The solvent was then removed at 90 °C under a stream of nitrogen. 2N HCl solution was added to the dried residue (**25**) which was then stirred at 95 °C for 20 min (Scheme 7). A shorter deprotection time was found to lead to decreased final yields of the product  $^{18}\text{F}$ -FAC.




**Scheme 7.** Subsequent deprotection of the hot intermediate **25** using (a) 2N HCl at 95 °C for 20 min. A shorter deprotection time results in decreased final yields of the product  $^{18}\text{F}$ -FAC (**1**).

After neutralization with 2N NaOH solution the mixture was loaded onto a semi-preparative HPLC column via a 10 mL injection loop. [ $^{18}\text{F}$ ]-FAC was eluted after 22.5 minutes at a flow rate of 3.5 mL/min using 3% MeCN in  $\text{H}_2\text{O}$  as the mobile phase. Following concentration on the module (100 °C under a stream of nitrogen for 10 min.), the sample was taken up in sterile saline and an aliquot was co-eluted on an analytical HPLC in combination with both possible 2'- $^{19}\text{F}$ -isomers FC (**26**) and FAC (**27**) as references (Figures 7 and 8).  $^{18}\text{F}$ -FAC was confirmed and its purity calculated to  $\geq 98\%$  with specific activity of  $\geq 63$  GBq/ $\mu\text{mol}$ . The total synthesis time was 168 minutes, including the concentration and saline formulation time. Finally, 0.75-0.86 GBq of  $^{18}\text{F}$ -FAC were obtained resulting in radiochemical yields (RCY) of 4.3-5.5% ( $n = 8$ , decay-corrected from end of bombardment (EoB)).




**Figure 7.** Radio-HPLC chromatogram of the purified fraction. Co-elution with both cold 2'- $^{19}\text{F}$ -isomers **26** and **27** confirms that the correct stereoisomer was obtained.



**2'-fluoro-2'-deoxy-cytidine  
(FC)**


**26**




**2'-fluoro-2'-deoxy-arabinocytidine  
(FAC)**

**27**

**Figure 8.** Chemical structures of the two different “cold” 2'-fluoro-cytidines FC (**26**) and FAC (**27**).





**Figure 9.** Radio-HPLC traces recorded after radiofluorinations performed at different reaction temperatures.

Additionally, a temperature study was performed to reveal why the  $^{18}\text{F}$ -incorporation decreases when the reaction temperature is increased. According to the mechanism intramolecular anhydro formation is expected at higher temperatures leading to a lower yield of the correct stereoisomer. Figure 9 shows radio-HPLC traces that were recorded after radiofluorination at 110 °C, 120 °C and 130 °C (20 min and 40 min) and co-elution with both 2'-fluoro stereoisomers FC and FAC (bottom panel). The results show that higher temperatures do lead to anhydro formation and radiofluorination of the anhydro intermediate results in  $^{18}\text{F}$ -FC. The  $^{18}\text{F}$ -FAC/FC ratio decreases as the reaction temperature and time increase.

## Conclusions

A novel radiochemical approach to  $^{18}\text{F}$ -FAC is described. For the first time,  $^{18}\text{F}$ -FAC was synthesized via late-stage radiofluorination of the intact pyrimidine nucleoside. Prior to synthesis, a novel fluorination precursor was designed using *in silico* methods and stability studies monitored by NMR. Precursor **23** was found to be a promising candidate for further testing out of the family of nucleosides that were investigated.  $^{18}\text{F}$ -incorporation studies led to optimal reaction conditions (110 °C, 20 minutes) that furnished  $^{18}\text{F}$ -FAC in 4.3-5.5% decay-corrected radiochemical yield. Considering that this new process involves late-stage fluorination and labelling of a cytidine derivative (more challenging than the equivalent uridine derivative), our radiochemical yield is good. Our overall synthesis time (168 minutes) is similar to that reported one for  $^{18}\text{F}$ -FAC.<sup>[11]</sup> Even though early  $^{18}\text{F}$ -introduction gives a higher radiochemical yield, its automation is more difficult and the reliability and reproducibility is less easy to maintain. In addition, the radiochemical yield and overall radiosynthesis time of our methods might be further improvable using other synthesizers; alternatively one could obtain the same amount of labelled nucleoside product by starting with a higher  $^{18}\text{F}$ -activity. The reaction time for the radiolabelling step (20 minutes) is comparable with that used in the related  $^{18}\text{F}$ -FMAU radiosynthesis,<sup>[12]</sup> reflective of the similar steric hindrance around the 2'-position in these nucleoside precursors. Shorter reaction times for this radiolabelling step were found to lead to lower yields of product through incomplete reaction.

Furthermore, it was shown that long reaction times and high reaction temperatures led to intramolecular anhydro-formation identified by the detection of  $^{18}\text{F}$ -FC. This study shows that rational precursor design can at least minimize the issue of side-product formation and hence can offer synthetic access of important PET radiotracer as a single stereoisomer. This



method should be transferable to GMP environments as it offers a quick and reproducible 2-step-route towards  $^{18}\text{F}$ -FAC comparable to the radiosynthesis of  $^{18}\text{F}$ -FDG. However, further optimization is required in order to improve the radiochemical yield and prepare for transfer to production of clinical grade radiotracer.

## Acknowledgements

This work was supported by Cancer Research Wales through PhD studentship funding (to J-PM). We thank Dr. Andrea Brancale (Cardiff University, School of Pharmacy and Pharmaceutical Sciences) for advice and support with DFT calculations. We acknowledge the National Mass Spectrometry Facility (Swansea, Wales, U.K) for provision of HR-MS determinations.

## References

- [1] J-P. Meyer, K. C. Probst, A. D. Westwell. *J. Label. Compd. Radiopharm.* **2014**, *57*, 333-337.
- [2] S. J. Martin, J. A. Eisenbarth, U. Wagner-Utermann, W. Mier, M. Henze, H. Pritzkow, U. Haberkorn, M. Eisenhut. *Nucl. Med. Biol.*, *29*:263–273, 2002.
- [3] A.F. Shields. *J. Nucl. Med.*, *44*:1432–1434, 2003.
- [4] M. Wagner, U. Seitz, A. Buck, B. Neumaier, S. Schultheiß, M. Bangerter, M. Bommer, F. Leithäuser, E. Wawra, G. Munzert, S. N. Reske. *Cancer Res.*, *63*:2681–2687, 2003.
- [5] R. E. Laing, M. A. Walter, D. O. Campbell, H. R. Herschman, N. Satyamurthy, M. E. Phelps. J. Czernin, O. N. Witte, C. G. Radu, *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 2847-2852.
- [6] J. T. Lee, D. O. Campbell, N. Satyamurthy, J. Czernin, C. G. Radu, *J. Nucl. Med.* **2012**, *53*, 275-280.
- [7] C. G. Radu, C. J. Shu, E. Nair-Gill, S. M. Shelly, J. R. Barrio, N. Satyamurthy, M. E. Phelps, O. N. Witte, *Nature Med.* **2008**, *14* (7), 783-788.
- [8] O. S. Tehrani, O. Muzik, L. K. Heilbrun, K. A Douglas, J. M. Lawhorn-Crews, H. Sun, T. J. Mangner, A. F. Shields, *J. Nucl. Med.* **2007**, *48*, 1436-1441.
- [9] C.-Y. Wu, P.-C. Chan, W.-T. Chang, R.-S. Liu, M. M. Alauddin, H.-E. Wang, *Appl. Radiat. Isot.* **2009**, *67*, 1362-1365.
- [10] N. Turkman, V. Paolillo, J. G. Gelovani, M. M. Alauddin, *Tetrahedron* **2012**, *68*, 10326-10332.
- [11] M. Lazari, K. M. Quinn, S. B. Claggett, J. Collins, G. J. Shah, H. E. Herman, B. Maraglia, M. E. Phelps, M. D. Moore, R. M. van Dam, *EJNMMI Research* **2013**, *3*, 52.
- [12] N. Turkman, J. G. Gelovani, M. M. Alauddin, *J. Label. Compd. Radiopharm.* **2010**, *53*, 782-786.
- [13] M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. J. Su, T. L. Windus, M. Dupuis, J. A. Montgomery, *J. Comp. Chem.* **1993**, *14*, 1347-1363.

[14]C. Serra, C. Aragonés, J. Bessa, J. Farras, J. Vilarrasa, *Tetrahedron Lett.* **1998**, 39, 7575–7578.

**Table 1.** Calculated electron density  $\Omega$  of selected cytidine (**4-8**) and uridine derivatives (**9-11**).

#	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	$[\Omega \pm \%]^a$
4	H	H		-0.911±0.09
5	H	Ac		-0.901±0.05
6	H	Boc		-0.860±0.01
7	Boc	Boc		-0.825±0.06
8	phthalimide	phthalimide		-0.814±0.08
9			H	-0.838±0.02
10			Boc	-0.770±0.03
11			NO <sub>2</sub>	-0.589±0.10

[a] Quantum chemically calculated net charge at the 2-carbonyl oxygen.

**Table 2.** [<sup>18</sup>F]-incorporation results obtained under different conditions.

#	solvent	T [°C]	t [min]	[ <sup>18</sup> F] <sup>a</sup> [%]
1	MeCN	90	20	n/a
2	<i>t</i> BuOH	95	20	n/a
3	DMF	90	20	1.3±0.3
4	DMF	95	20	2.6±0.2
5	DMF	100	20	4.9±0.6
6	DMF	105	20	7.7±0.8
7	DMF	110	20	9.4±0.8
8	DMF	110	30	7.9±0.6
9	DMF	115	20	6.9±0.5
10	DMF	120	20	4.1±0.4
11	DMF	125	20	2.3±0.2
12	DMF	130	20	1.8±0.2

[a] [<sup>18</sup>F]-incorporation was calculated using radio-TLC. Each reaction was repeated three times under identical conditions.