

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/72484/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Pereira, Paula, Pedrosa, Sílvia S., Wymant, Jennifer M., Sayers, Edward, Correia, Alexandra, Vilanova, Manuel, Jones, Arwyn Tomos and Gama, Francisco M. 2015. siRNA inhibition of endocytic pathways to characterize the cellular uptake mechanisms of folate functionalized glycol chitosan nanogels. *Molecular Pharmaceutics* 12 (6) , pp. 1970-1979. 10.1021/mp500785t

Publishers page: <http://dx.doi.org/10.1021/mp500785t>


Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.


This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Supplementary Figures



Supplementary Figure 1. Effect of CME inhibition on cellular uptake of the nanogel. Live cell confocal fluorescence microscopy images of the HeLa cells transfected with si-CHC treated with nanogels for 7 h. Each row, from left to the right, represent maximum projection, Z section and DIC images.



Supplementary Figure 2. Effect of si-Cav-1 depletion on cellular uptake of the nanogel. Live cell confocal microscopy observation of the Alexa488-nanogels incubated with si-Cav-1 transfected cells for 7 h. Each row, from left to the right, represent maximum projection, Z section and DIC images.