

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/76318/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Kitamura, Akane, Miyauchi, Naoko, Hamada, Hirotaka, Hiura, Hitoshi, Chiba, Hatsune, Okae, Hiroaki, Sato, Akiko, John, Rosalind Margaret and Arima, Takahiro 2015. Epigenetic alterations in sperm associated with male infertility. *Congenital Anomalies* 55 (3) , pp. 133-144. 10.1111/cga.12113

Publishers page: <http://dx.doi.org/10.1111/cga.12113>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



**Invited Review Article: Congenital anomalies**

**Title: Epigenetic alterations in sperm associated with male infertility**

**Short title: Epigenetics and male infertility**

Akane Kitamura<sup>1</sup>, Naoko Miyauchi<sup>1</sup>, Hirotaka Hamada<sup>1</sup>, Hitoshi Hiura<sup>1</sup>, Hatsune Chiba<sup>1</sup>, Hiroaki Okae<sup>1</sup>, Akiko Sato<sup>1</sup>, Rosalind M. John<sup>2</sup>, Takahiro Arima<sup>1,3</sup>

<sup>1</sup>Department of Informative Genetics, Environment and Genome Research Center, Tohoku University Graduate School of Medicine, 2-1 Seiryō-cho, Aoba-ku, Sendai, 980-8575, Japan.

<sup>2</sup>Cardiff School of Biosciences, Museum Avenue, Cardiff, CF10 3US UK.

<sup>3</sup>Correspondence address

TEL: ++81-22-717-7844;

FAX: ++81 22-717-7063;

Email: tarima@med.tohoku.ac.jp

**Key words:** Male infertility, Genomic imprinting, DNA methylation, Genomic imprinting disorders, Assisted reproductive technologies (ART)

## **Abstract**

The most common form of male infertility is a low sperm count, known as oligozoospermia. Studies suggest that oligozoospermia is associated with epigenetic alterations. Epigenetic alterations in sperm, which may arise due to the exposure of gametes to environmental factors or preexist in the sperm of infertile individuals, may contribute to the increased incidence of normally rare imprinting disorders in babies conceived after assisted reproductive technology using the sperm of infertile men. Genomic imprinting is an important developmental process whereby the allelic activity of certain genes is regulated by DNA methylation established during gametogenesis. The aberrant expression of several imprinted genes has been linked to various diseases, malignant tumors, lifestyle and mental disorders in humans. Understanding how infertility and environmental factors such as reproductive toxicants, certain foods, and drug exposures during gametogenesis contribute to the origins of these disorders via defects in sperm is of paramount importance. In this review, we discuss the association of epigenetic alterations with abnormal spermatogenesis and the evidence that epigenetic processes, including those required for genomic imprinting, may be sensitive to environmental exposures during gametogenesis, fertilization and early embryonic development. In addition, we review imprinting diseases and their relationships with environmental factors. While the plasticity of epigenetic marks may make these more susceptible to modification by the environment, this also suggests that aberrant epigenetic marks may be reversible. A greater understanding of this process and the function of epidrugs may lead to the development of new treatment methods for many

1 adult diseases in the future.

2

## 1    **Introduction**

2            Approximately half of human infertility can be explained by abnormal  
3    spermatogenesis. Disturbingly, the incidence of abnormal spermatogenesis has  
4    increased in developed countries, including Japan (Japan Society of Obstetrics and  
5    Gynecology Registry). Oligozoospermia is the most common disorder of male  
6    infertility characterized by abnormally low concentrations of spermatozoa in the semen.  
7    Although different genetic causes are known, they account only for a fraction of the  
8    cases of aberrant spermatogenesis (Dohle et al. 2002; Fernandes et al. 2002; Gianotten  
9    et al. 2003). Epigenetic factors, including DNA methylation, histone modifications and  
10   chromatin remodeling, have been studied extensively during gametogenesis and germ  
11   cell maturation and it is clear that germ cells undergo extensive epigenetic  
12   reprogramming in a sex-specific manner (Dada et al. 2012; van Montfoort et al. 2012;  
13   Boissonnas et al. 2013). Consequently, aberrant epigenetic alterations may underlie  
14   some cases of oligozoospermia.

15           Indirect evidence for a role for aberrant epigenetic processes in  
16   oligozoospermia comes from studies on human assisted reproductive technology (ART),  
17   in which the eggs and/or sperm are manipulated in the laboratory to help infertile  
18   persons of reproductive age conceive. Recent reports identified an increased incidence  
19   of normally rare imprinting disorders, especially Beckwith-Wiedemann syndrome  
20   (BWS; OMIM 130650), Angelman syndrome (AS; OMIM 105830) and Silver-Russell  
21   syndrome (SRS; OMIM 180860), in babies conceived after ART (DeBaun et al. 2003;  
22   Gosden et al. 2003; Maher 2005). Several reports have suggested that imprint

methylation errors occur during the process of ART, both in *in vitro* fertilization (IVF) and intracytoplasmic sperm injection (ICSI) (Cox et al. 2002; DeBaun et al. 2003; Gicquel et al. 2003; Maher et al. 2003; Moll et al. 2003; Orstavik et al. 2003; Ludwig et al. 2005; Rossignol et al. 2006; Bowdin et al. 2007; Kagami et al. 2007) which may be due to *in vitro* embryo transfer procedures performed at the time of epigenetic fluidity (Lucifero et al. 2004; Niemitz and Feinberg 2004; Thompson and Williams 2005; Horsthemke and Buiting 2006). However, our work and that of others suggests that epigenetic risks linked to ART techniques can also originate in the use of sperm with preexisting epigenetic errors (Kobayashi et al. 2007; Kobayashi et al. 2009). This review provides an overview of the current state of knowledge of human sperm epigenetics and what is known regarding the effects of environmental and nutritional factors on the sperm epigenome.

## **Genomic imprinting**

Genomic imprinting is an epigenetic phenomenon that describes parent-of-origin patterns of monoallelic gene expression reported in mammals and some plant species (Barlow and Bartolomei 2014). The term genomic imprinting was first used to describe the failure of mono-parental embryos to develop appropriately *in utero* despite their diploid DNA content (Barton et al. 1984; McGrath and Solter 1984; Surani et al. 1984). We now know that there are over one hundred genes in mammals that are regulated by genomic imprinting and many of these have critically important roles in early development and also later life process, both metabolic and behavioural (Surani

1998; Tilghman 1999; Cleaton et al. 2014).

Differences in the parental genomes are first established in the germline when the two parental genomes are physically separate. Discrete DNA regions are marked by DNA methylation in one or other germline. After fertilisation these marks are maintained despite the extensive epigenetic reprogramming that takes place early in development (Morgan et al. 2005), to generate regions of the genome that have DNA methylation present on one parental allele and absent on the other allele. These regions are termed gametic differentially methylated regions (gDMR; **Figure 1**). These gDMRs act as the catalyst for a further series of epigenetic changes including both the modification of histones and somatic DNA methylation events, which generate extensive domains of imprinted chromatin some of which span several megabases. Within these domains certain genes are silenced on one parental allele and active on the other parental allele with most imprinted domains containing both paternally- and maternally-expressed genes. While these gDMRs are maintained for the lifetime of the individual, the monoallelic expression status of imprinted genes can vary with tissue type and developmental stage suggesting that functional imprinting is important at different times for different genes. In the mouse female germline, gDMRs acquire DNA methylation after birth during the transition from primordial to antral follicles in the postnatal growth phase (post-pachytene) (Obata and Kono 2002; Lucifero et al. 2004; Hiura et al. 2006). In the human female germline, maternal methylation of gDMRs has already been initiated to some extent in adult non-growing oocytes but not in neonatal oocytes (Sato et al. 2007). In mouse male germline, methylation at three sites

(*H19*, *Rasgrf1* and *Gtl2*) is present prenatally before meiosis and completed by the pachytene phase of postnatal spermatogenesis (Davis et al. 1999; Davis et al. 2000; Ueda et al. 2000; Li et al. 2004) with complete loss of methylation of maternal DMRs. While gDMRs are established in the germline, some imprinted domains also contain somatic DMRs (sDMR) which are not inherited via the germline but which appear during embryogenesis either before or after monoallelic expression is established and which are also important for maintaining monoallelic gene expression (John and Lefebvre 2011). In addition to the establishment and maintenance of allele-specific epigenetic marks, imprints must be erased in the developing germline and reset for the next generation (**Figure 2**). Establishment, maintenance and erasure of imprints all involve dynamic changes in epigenetic marks that take place at different stages of development in males and females. In summary, genomic imprinting is a dynamic epigenetic process both in the germline and during early development. Epigenetic errors at any stage in the process of establishment, maintenance or erasure of imprints can have a catastrophic consequence for the next generation, as evidenced by the genomic imprinting disorders.

## **Genomic Imprinting disorders**

The importance of correct genomic imprinting in humans is best illustrated by a number of rare but striking childhood developmental disorders associated with imprinted loci. Prader-Willi syndrome (PWS; OMIM 176270) and Angelman syndrome (AS; OMIM 105830) are two clinically distinct imprinting disorders linked to the same



1 imprinted region on chromosome 15q11-q1 ([Buiting 2010](#)). PWS is characterized by  
2 endocrine and neural abnormalities and malformation and is mainly associated with  
3 maternal uniparental disomy of 15q11-q1 (70%) and methylation defects (2-5%). In  
4 contrast, AS, which is characterized by global developmental delay, convulsions,  
5 scoliosis, excessive laughter, and movement, balance and sleep disorders, is associated  
6 with loss of function of the maternally expressed *UBE3A* gene either through deletions  
7 (70%), paternal uniparental disomy (0-20%) or aberrant methylation (2-5%) of the  
8 maternal allele. Beckwith-Wiedemann syndrome (BWS; OMIM 130650) and  
9 Silver-Russell syndrome (SRS; OMIM 180860) are similarly clinically distinct  
10 syndromes associated with a single chromosomal region at 11p15.5 ([Jacob et al. 2013](#)).  
11 BWS is a fetal overgrowth disorder characterized by exomphalos, macroglossia,  
12 gigantism and an increased risk of developing embryonal tumors in childhood. BWS is  
13 associated with a number of genetic and epigenetic alterations. The most frequent  
14 alteration observed in BWS is hypomethylation of a gDMR located over the promoter  
15 of a long, non-coding RNA called *LIT1* or *Kcnq1ot1*, which is found in >60% of  
16 sporadic BWS patients. Animal studies suggest that this gDMR regulates expression of  
17 the maternally-expressed *CDKN1C* gene known to play a key role in limiting fetal  
18 growth ([Andrews et al. 2007](#); [Tunster et al. 2011](#)). SRS is a similarly clinically  
19 heterogeneous condition characterized by severe intrauterine growth retardation, poor  
20 postnatal growth, craniofacial features such as a triangular-shaped face and a broad  
21 forehead, body asymmetry, and a variety of minor malformations. The most frequent  
22 alteration in SRS is hypomethylation of the gDMR spanning the promoter of a

non-coding RNA called *H19* apparent in 40% of cases ([Bliek et al. 2006](#)). This gDMR regulates the imprinted expression of the fetal growth factor gene *IGF2* (Insulin like growth factor 2) ([DeChiara et al. 1991](#); [Leighton et al. 1995](#)). However, rare SRS patients have been reported with maternal microduplications spanning *CDKN1C* ([Bonaldi et al. 2011](#)). Furthermore, additional loci on various chromosomes have been implicated as having a role in this syndrome ([Davis et al. 2000](#); [Ueda et al. 2000](#); [Gicquel et al. 2003](#); [Maher et al. 2003](#); [Sato et al. 2007](#)). These disorders highlight the necessity of appropriately regulated gene dosage at imprinted loci mediated by epigenetic processes, which might consequently be subject to external influences acting on the epigenome.

## **ART and congenital imprinting disorders**

A number of publications have suggested an association between ART and genomic imprinting disorders (**Table 1**) ([Chiba et al. 2013](#); [Hiura et al. 2014](#)). The first report linking ART to AS in 2002 highlighted loss of DNA methylation on chromosome 15 ([Cox et al. 2002](#)). In 2004 an increased frequency of BWS after ART was reported, again linked to changes in DNA methylation ([DeBaun et al. 2003](#)). In 2007 SRS was linked to ART and hypermethylation at an imprinted loci ([Kagami et al. 2007](#)). ART does not, however, appear to be a risk factor in PWS ([Gold et al. 2014](#)).

There are several proposed mechanisms which may underlie the increased frequency of imprinting disorders in ART including the exposure of gametes and early embryos to culture conditions, the superovulation of oocytes and the presence of

preexisting imprinting mutations in sperm. Some studies have shown that exposure of mouse embryos to different culture conditions can alter the expression and imprinting of various genes, which could result in abnormal development ([DeBaun et al. 2003](#); [Gicquel et al. 2003](#); [Maher et al. 2003](#); [Lucifero et al. 2004](#)). We, and others, have demonstrated that superovulation (artificial induction of ovulation with high doses of gonadotrophins) affects imprint methylation ([Chang et al. 2005](#); [Ligon 2005](#); [Sato et al. 2007](#)). Embryo freezing may also be an issue as this has been found to have deleterious effects on DNA, embryonic gene expression, telomeres and plasma and nuclear membranes ([Emiliani et al. 2000](#); [Honda et al. 2001](#)). Furthermore, the timing of embryo transfer may be an issue. Case reports of monochorionic dizygotic twins and conjoined twins with BWS resulting from transfer at the blastocyst stage ([Shimizu et al. 2004](#); [Miura and Niikawa 2005](#)) reported demethylation of *LIT1* (*KCNQ1OT1*), suggesting that this demethylation occurs at a critical stage of preimplantation development. In addition to epigenetic errors induced by the process of ART, there is evidence that sperm from men with fertility issues carry preexisting epigenetic errors.

### **Sperm from infertile men and epigenetic errors**

Studies have shown that disturbed spermatogenesis is associated with incorrect DNA methylation at gDMRs (Table 2). In spermatozoa from oligozoospermic men, the occurrence of hypermethylation of several maternally imprinted DMRs or hypomethylation of paternally imprinted DMRs is increased ([Marques et al. 2004](#); [Kobayashi et al. 2007](#); [Marques et al. 2008](#); [Hammoud et al. 2010](#); [Sato et al. 2011](#)).

1 Boissonnas et al. also reported the association between methylation and sperm  
2 concentration in teratozoospermic (TZ) and oligoasthenoteratozoospermic (OAT)  
3 patients ([Boissonnas et al. 2010](#)). In the TZ group, 11 of 19 patients displayed loss of  
4 methylation of the *IGF2* DMR or of both the *IGF2* DMR and the *H19* DMR. In the  
5 OAT group, 16 of 22 patients displayed a severe loss of methylation of the *H19* DMR,  
6 and this closely correlated with sperm concentration. Marques et al. suggested an  
7 association between aberrant epigenetic sperm modifications and oligozoospermia  
8 ([Marques et al. 2004](#)). Normozoospermic individuals (0.13%), Moderate (17%) and  
9 Severe (30%) oligozoospermic patients all showed abnormal methylation of *H19*. We  
10 examined the DNA methylation status of seven imprinted genes in spermatonic DNA  
11 obtained from infertile men and also found abnormal maternal and paternal DNA  
12 methylation at several imprinted loci (**Figure 3**). Samples (10/96 cases) with both  
13 maternal and paternal defects were primarily from men with severe oligospermia.  
14 Importantly, the outcome of ART (fertility rates and implantation rates) with sperm  
15 shown to have an abnormal DNA methylation pattern is generally poor ([Kobayashi et](#)  
16 [al. 2007](#)).

17 As spermatogenesis progresses, the genome undergoes major changes that not  
18 only influence genetic and epigenetic information but also alter the nuclear structure. It  
19 is consequently important to understand how the specific nucleoprotamine/histone  
20 structure of the sperm nucleus conveys epigenetic information and how this might  
21 control early embryonic growth. In most cell types, DNA is wrapped around histone but  
22 in sperm, protamines, which are small arginine-rich nuclear proteins, replace histones

late in the haploid phase of spermatogenesis and these proteins are essential for spermatogenic function (Cho et al. 2001)(**Figure 2**). Both the phosphorylation of protamines and the ratio of the two human protamines, protamine (P1) and protamine 2 (P2), are important for optimal sperm function. The P1/ P2 ratio in fertile men ranges from 0.8 to 1.2 (Carrell and Liu 2001). Perturbation of this ratio, either higher or lower than normal, has been reported to be associated with poor semen quality, increased DNA damage and/or decreased fertility (Chevaillier et al. 1987; Balhorn et al. 1988; Belokopytova et al. 1993; Carrell et al. 1999; Razavi et al. 2003; Aoki et al. 2005). An increasing number of reports now support the hypothesis that sperm DNA is not homogeneously packed with these protamines and that histones are still present at some sites (Rousseaux et al. 2005). While some investigators have suggested that this is due to inefficient protamine replacement, the persistence of histones at certain sites may play a functional role in supporting the epigenetic code in the sperm (Weber et al. 2007). Protamine replacement occurs in the spermatid stage of spermatogenesis after the completion of meiosis (Baarends et al. 1999). The elongating spermatid also undergoes other maturational events that affect motility and fertilization ability during the period of protamine replacement. The association between abnormal protamine replacement and generally diminished semen quality may be a defect in the unique gene regulation system of temporal uncoupling of transcription and translation during spermatogenesis (Carrell et al. 2007).

Alteration of the P1 to P2 ratio generally denotes abnormal spermatogenesis and is a possible direct cause of abnormal methylation of maternal and paternal gDMRs

(Hammoud et al. 2010). Azoospermia caused by anejaculation and secondary inflammatory obstruction is related to an increase of methylation level in maternal DMRs (Marques et al. 2010). Male infertility may also be related to the improper erasure of DNA methylation during spermatogenesis at many non-imprinted genes in addition to abnormal methylation levels at gDMRs (Houshdaran et al. 2007). There are some significant implications for sperm with abnormal protamine replacement, and for the use of such sperm for ICSI. Further research should be done to classify the role of retained histones throughout the spermatogenic genome in mature sperm from men with normozoospermia as well as in patients with known chromatin abnormalities.

## **Teratological environmental factors (endocrine disruptors) and epigenetic modifications**

Abnormal sperm development may originate from exposure of the male germline to environmental factors. Persistent organic pollutants (POPs), which were used intensively worldwide for several decades until the 1980s, have been implicated in reproductive disorders. Because of the stability and bioaccumulation of these compounds in the environment, human populations are simultaneously exposed to a variety of those contaminants through the consumption of food. Several POPs have been shown to have toxic effects on reproductive and endocrine functions in humans (Govarts et al. 2012) and a number of human epidemiological studies have demonstrated the adverse effects of POPs exposure on markers of reproduction,

1 including semen quality (sperm concentration, motility, and morphology) (Guo et al.  
2 2000; Richthoff et al. 2003; Toft and Guillette 2005; Meeker et al. 2010), spermat-  
3 DNA integrity (Bonde et al. 2008; McAuliffe et al. 2012), and circulating reproductive  
4 hormone levels (Richthoff et al. 2003), though some studies found only marginal effects  
5 (Toft et al. 2006; Haugen et al. 2011). In general, however, these reports suggest that  
6 POPs have adverse effects on reproductive health outcomes.

7         Endocrine disruptors are another potential environmental factor driving  
8 abnormal sperm development. Male gonadal development occurs around midgestation  
9 in humans initiated by the differentiation of precursor Sertoli cells in response to the  
10 testis-determining factor SRY. The fetal testis contains steroid receptors and is a target  
11 for endocrine hormones. The androgen receptor and estrogen receptor- $\beta$  are present in  
12 both Sertoli cells and germ cells. Although the testis does not produce steroids at this  
13 stage of development, estrogens and androgens can affect testis cellular functions.  
14 Treatment with endocrine disruptors at a critical time of gonadal sex determination  
15 promotes an adult testis phenotype with decreased spermatogenic capacity in rat and, as  
16 a result, male infertility. External factors could induce an epigenetic transgenerational  
17 phenotype through apparent reprogramming of the male germ line (Anway et al. 2005).  
18 However, it is still unclear whether steroids acting inappropriately during the time of  
19 gonadal sex determination act to reprogram the germ line via epigenetic DNA  
20 methylation to cause this transgenerational transmission of an altered phenotype.

21         Seminal tract infection, one of the most common causes of infertility in men  
22 (Keck et al. 1998), may also contribute to abnormal sperm development. The presence

1 of leukocytes in semen, also known as leukocytospermia, ([Korrovits et al. 2008](#);  
2 [Cumming and Carrell 2009](#)), is an indicator of seminal tract infection although this  
3 correlation remains controversial ([Bezold et al. 2007](#)). Asthenozoospermia is often  
4 associated with the presence of infection or leukocytes in semen although it is not  
5 known whether infection plays a causative role ([Wolff 1995](#)). The association between  
6 epigenetic changes and such sperm abnormalities as asthenozoospermia and  
7 leukocytospermia is unknown. However, there is a precedent for infection inducing  
8 epigenetic alterations in other cell types. In gastric carcinogenesis, *H. pylori* infection  
9 induces aberrant promoter methylation in tumor-suppressor genes, including *p16<sup>INK4A</sup>*,  
10 *LOX*, and *CDH1* ([Kaneda et al. 2004](#); [Ushijima et al. 2006](#)). Further work is required to  
11 establish whether epigenetic alterations in sperm are induced by seminal tracts  
12 infections.

13         Social stress, acting through hormone signalling pathways, is another recent  
14 additions to the group of environmental factors that are known to induce epigenetic  
15 changes. The extent and type of maternal care very early in life in rodents has been  
16 shown to influence epigenetic marks at the glucocorticoid receptor in the neonatal  
17 hippocampus, and this may influence later life stress responses in the offspring ([Weaver](#)  
18 [et al. 2004](#); [Meaney et al. 2007](#)). Furthermore, in another rodent model, Roth et al.  
19 found that psychosocial stress (comparable to human post-traumatic stress disorder  
20 (PTSD)) led to an increase in *Bdnf* methylation in the dorsal hippocampus and  
21 downregulation of *Bdnf* expression in the dorsal and ventral hippocampus, but not in  
22 other PTSD-relevant regions ([Roth et al. 2011](#)). The induction of region-specific



epigenetic changes in response to traumatic stress during adulthood demonstrates that DNA methylation remains an active process that can be shaped by environmental factors even in the adult nervous system. Again, the effect of stress on the sperm epigenome has not been investigated. However, stress is also a cause of male infertility this may occur through epigenetic alterations in the germline (Bale 2014).

## **Nutrition and epigenetic regulation**

Epigenetic marks are tightly regulated, both temporally and spatially, during fetal development and lactation (Lee et al. 2002; Allegrucci et al. 2005; Morgan et al. 2005) but can be influenced at key stages by diet. Agouti viable yellow (Avy) is a fascinating animal model whereby the environmental influences on the epigenome can be monitored via a coat colour phenotype (Wolff et al. 1999). A gene alteration, which involves an intra-cisternal A particle (IAP) retrotransposon insertion upstream of the agouti gene (A), leads to ectopic expression of the agouti protein and a change of hair color from agouti to yellow. The extent of this coat colour change is influenced by the degree of methylation of the IAP element, which can be influenced by methyl donor supplementation of the maternal diet (Waterland and Jirtle 2003). Dietary supplementation with a methyl donor during pregnancy increases the proportion of pups carrying a methylated IAP sequence and thus the number with a yellow coat colour (Rakyan et al. 2003; Waterland and Jirtle 2004). Maternal and post-weaning high fat diets can also alter epigenetic regulation of the hedonic reward pathways and metabolic regulation of the energy balance in mice (Vucetic et al. 2011), and alter methylation of

1 the leptin promoter in rats (Milagro et al. 2009). These data provide compelling  
2 evidence that diet alone can alter the epigenome.

3 Nutrition during early growth and development may influence DNA  
4 methylation because one-carbon metabolism is dependent on dietary methyl donors and  
5 on cofactors such as methionine, choline, folic acid and vitamin B-12 (MacLennan et al.  
6 2004). The limited availability of acetyl-CoA for HAT activity and methyl donors of  
7 SAM (*S* - adenosylmethionine) provided via the folate-methionine pathway may  
8 therefore play a role in the establishment of inappropriate epigenetic patterns.  
9 Conversely, dietary supplementation may provide a route to attenuating inappropriate  
10 epigenetic patterns as the changes in DNA methylation which result from a decrease in  
11 DNMT1 (DNA methyltransferase) activity can be partially prevented by folate  
12 supplementation (Lillycrop et al. 2005; Lillycrop et al. 2007).

13 The influences of poor nutrition on epigenetic marks is not limited to the fetal  
14 stage. Nutrition during postnatal development can permanently alter the epigenetic  
15 regulation of some imprinted genes. Methyl-donor-deficient diet in postnatal life is  
16 associated with altered epigenetic regulation of *IGF2* and growth retardation (Waterland  
17 et al. 2006). In humans, diet has been shown to affect the DNA methylation status of  
18 patients with hyperhomocysteinaemia. This disease is caused by the accumulation of  
19 S-adenosylhomocysteine (an inhibitor of DNA methyltransferases)(Waterland et al.  
20 2006).

21 Given the consequences of altered nutrient availability in a number of situations, it is  
22 possible that changes may be also occur the male germline in response to diet. One very

1 compelling study demonstrated that a low protein diet in male rats results in altered  
2 chromatin packing in sperm and changes in DNA methylation in the offspring (Carone  
3 et al. 2010). These data all suggest that something as seemingly innocuous as a dietary  
4 imbalance can have a detrimental effect on the epigenome at certain critical stages.

5 In addition to the availability of specific nutrients, alterations in the expression,  
6 localization and/or activity of epigenetic modifiers, such as the DNA methyltransferases,  
7 the histone-modification enzymes and their associated proteins, may play a role in  
8 driving abnormalities in the sperm epigenome. Some modifiers are specifically  
9 expressed in germ cells and the crucial roles of germ-cell-specific genes such as  
10 *Dnmt3L* and *Prdm9* has been highlighted in conventional mouse gene knockout studies  
11 (Bourc'his et al. 2001; Hata et al. 2002; Hayashi et al. 2005). We reported DNA  
12 sequence variations in the gene encoding *DNMT3L* associated with imprinting errors and  
13 oligospermia (Kobayashi et al. 2009). A recent report suggests that gestational diet can  
14 alter the expression of histone demethylases and Dnm3L, at least in the exposed placenta  
15 (Gabory et al. 2012). Consequently both poor sperm quality and imprinting errors may  
16 be linked by both genetic and dietary-driven alterations in epigenetic regulators.

## 1   **Conclusions**

2           Mounting evidence from both human studies and animal models suggests that  
3   epigenetic modifications provide a link between the environment and alterations in gene  
4   expression that might lead to disease phenotypes. Importantly, direct evidence from  
5   animal studies supports the role of environmental epigenetics in male infertility and  
6   suggests the possibility that the use of ART to treat male infertility may lead to disease  
7   later in life. However, ART is a relatively recent technology and the longer term  
8   consequences of ART treatments such as ICSI and embryo freezing before transfer have  
9   not yet been manifested due to the young age of the majority of ART children.  
10   Environmental exposures to nutritional, chemical and physical factors all have the  
11   potential to alter gene expression and, therefore, modify sperm quality in various ways  
12   through changes in the epigenome. A summary of the factors known to influence DNA  
13   methylation is presented in **Figure 4**.

14           It is still unknown when imprinting epigenetic errors related to male infertility  
15   arise and what factors may predispose to epigenetic changes. Hormonal stimulation of  
16   oocytes, *in vitro* culture, cryopreservation, and the timing of embryo transfer have all  
17   been shown to influence the proper establishment and maintenance of genomic imprints.  
18   Some infertile males, particularly those with oligozoospermia, carry preexisting  
19   imprinting errors in their sperm. Therefore the process of ART and infertility itself  
20   might increase the risk of imprinting disorders.

21           The developmental origins of health and disease (DOHaD) paradigm, first  
22   proposed by Prof. David Barker, postulates that suboptimal growth early life can

1 program changes which affect life long health, increasing the risks for various diseases.  
2 There is evidence both from human studies and experimental models that this  
3 programming may be mediated via changes in the epigenome. Epigenetic changes likely  
4 occur during the fetal and infant periods but it is clear that oocytes and sperm are also  
5 vulnerable to environmentally-induced epigenetic alterations, and that the newly  
6 fertilised zygote is at a particular susceptible stage.

## 7 **Future perspective**

9 While genomic imprinting disorders are very rare, it is increasingly apparent that the  
10 bulk of common human diseases do not arise solely from genetic or environmental  
11 causes but also have an epigenetic component. Our knowledge that the epigenomes of  
12 gametes and newly fertilized embryos are susceptible stages for  
13 environmentally-induced epigenetic changes has particularly important implications as  
14 changes in lifestyle and modes of reproduction may have long term implications for  
15 human health that are not yet fully appreciated. Recent work identifies advanced  
16 paternal age as a risk factor for autism, depression, epilepsy and prostate cancer in  
17 children (Kondrashov 2012; Sun et al. 2012). While there are a number of possible  
18 explanations for these associations, the accumulation of epigenetic errors in the sperm  
19 may be a contributory factor. As the human population ages and the use of ART  
20 increases worldwide, it will become increasingly important to determine the extent to  
21 which environmentally-induced epigenetic changes contribute to disease. A detailed  
22 characterisation of the normal epigenetic process that take place in the germline and

1 during very early development will be important in achieving this goal. Understanding  
2 how and when environmental factors can influence the epigenome to cause disease,  
3 identifying ways in which to modulate aberrant epigenetic marks, and also determining  
4 the best timeframe to reverse aberrant epigenetic marks all have the potential to lead to  
5 improved human health.

6

## Acknowledgments

We would like to thank all the members of our laboratory for their technical assistance, support and valuable suggestions. In particular, we thank Prof. Kazuyoshi Arishima of Azabu University and Prof. Kentaro Tanemura of Tohoku University for giving us the opportunity to write this review.

## Funding

This work was supported by Grants-in-Aid from the Ministry of Health, Labour and Welfare of the Japanese government (Health Labour Sciences Research Grant, Research on Measures for Intractable Diseases 26310301), Scientific Research (KAKENHI) (Exploratory Research 25670691, Scientific Research (C) 24613001, Grant-in-Aid for Young Scientists (B), 26870036, 26870033), as well as the Takeda Science Foundation (TA).

**Conflict of interest statement:** Disclosure none.

.

## REFERENCES

- Allegrucci, C., Thurston, A., Lucas, E. and Young, L. 2005. Epigenetics and the germline, *Reproduction* 129: 137-49.
- Andrews, S. C., Wood, M. D., Tunster, S. J., Barton, S. C., Surani, M. A. and John, R. M. 2007. *Cdkn1c* (p57Kip2) is the major regulator of embryonic growth within its imprinted domain on mouse distal chromosome 7, *BMC Dev Biol* 7: 53.
- Anway, M. D., Cupp, A. S., Uzumcu, M. and Skinner, M. K. 2005. Epigenetic transgenerational actions of endocrine disruptors and male fertility, *Science* 308: 1466-9.
- Aoki, V. W., Liu, L. and Carrell, D. T. 2005. Identification and evaluation of a novel sperm protamine abnormality in a population of infertile males, *Hum Reprod* 20: 1298-306.
- Baarends, W. M., Hoogerbrugge, J. W., Roest, H. P., Ooms, M., Vreeburg, J., Hoeijmakers, J. H. and Grootegoed, J. A. 1999. Histone ubiquitination and chromatin remodeling in mouse spermatogenesis, *Dev Biol* 207: 322-33.
- Bale, T. L. 2014. Lifetime stress experience: transgenerational epigenetics and germ cell programming, *Dialogues Clin Neurosci* 16: 297-305.
- Balhorn, R., Reed, S. and Tanphaichitr, N. 1988. Aberrant protamine 1/protamine 2 ratios in sperm of infertile human males, *Experientia* 44: 52-5.
- Barlow, D. P. and Bartolomei, M. S. 2014. Genomic imprinting in mammals, *Cold Spring Harb Perspect Biol* 6.
- Barton, S. C., Surani, M. A. and Norris, M. L. 1984. Role of paternal and maternal genomes in mouse development, *Nature* 311: 374-6.
- Belokopytova, I. A., Kostyleva, E. I., Tomilin, A. N. and Vorob'ev, V. I. 1993. Human male infertility may be due to a decrease of the protamine P2 content in sperm chromatin, *Mol Reprod Dev* 34: 53-7.
- Bezold, G., Politch, J. A., Kiviat, N. B., Kuypers, J. M., Wolff, H. and Anderson, D. J. 2007. Prevalence of sexually transmissible pathogens in semen from asymptomatic male infertility patients with and without



leukocytospermia, *Fertil Steril* 87: 1087-97.

Bliek, J., Terhal, P., van den Bogaard, M. J., Maas, S., Hamel, B., Salieb-Beugelaar, G., Simon, M., Letteboer, T., van der Smagt, J., Kroes, H. et al. 2006. Hypomethylation of the H19 gene causes not only Silver-Russell syndrome (SRS) but also isolated asymmetry or an SRS-like phenotype, *Am J Hum Genet* 78: 604-14.

Boissonnas, C. C., Abdalaoui, H. E., Haelewyn, V., Fauque, P., Dupont, J. M., Gut, I., Vaiman, D., Jouannet, P., Tost, J. and Jammes, H. 2010. Specific epigenetic alterations of IGF2-H19 locus in spermatozoa from infertile men, *Eur J Hum Genet* 18: 73-80.

Boissonnas, C. C., Jouannet, P. and Jammes, H. 2013. Epigenetic disorders and male subfertility, *Fertil Steril* 99: 624-31.

Bonaldi, A., Mazzeu, J. F., Costa, S. S., Honjo, R. S., Bertola, D. R., Albano, L. M., Furquim, I. M., Kim, C. A. and Vianna-Morgante, A. M. 2011. Microduplication of the ICR2 domain at chromosome 11p15 and familial Silver-Russell syndrome, *Am J Med Genet A* 155A: 2479-83.

Bonde, J. P., Toft, G., Rylander, L., Rignell-Hydbom, A., Giwercman, A., Spano, M., Manicardi, G. C., Bizzaro, D., Ludwicki, J. K., Zvyezday, V. et al. 2008. Fertility and markers of male reproductive function in Inuit and European populations spanning large contrasts in blood levels of persistent organochlorines, *Environ Health Perspect* 116: 269-77.

Bourc'his, D., Xu, G. L., Lin, C. S., Bollman, B. and Bestor, T. H. 2001. Dnmt3L and the establishment of maternal genomic imprints, *Science* 294: 2536-9.

Bowdin, S., Allen, C., Kirby, G., Brueton, L., Afnan, M., Barratt, C., Kirkman-Brown, J., Harrison, R., Maher, E. R. and Reardon, W. 2007. A survey of assisted reproductive technology births and imprinting disorders, *Hum Reprod* 22: 3237-40.

Buiting, K. 2010. Prader-Willi syndrome and Angelman syndrome, *Am J Med Genet C Semin Med Genet* 154C: 365-76.

Carone, B. R., Fauquier, L., Habib, N., Shea, J. M., Hart, C. E., Li, R., Bock, C., Li, C., Gu, H., Zamore, P. D. et al. 2010. Paternally induced transgenerational environmental reprogramming of metabolic gene

expression in mammals, *Cell* 143: 1084-96.

Carrell, D. T., Emery, B. R. and Hammoud, S. 2007. Altered protamine expression and diminished spermatogenesis: what is the link?, *Hum Reprod Update* 13: 313-27.

Carrell, D. T., Emery, B. R. and Liu, L. 1999. Characterization of aneuploidy rates, protamine levels, ultrastructure, and functional ability of round-headed sperm from two siblings and implications for intracytoplasmic sperm injection, *Fertil Steril* 71: 511-6.

Carrell, D. T. and Liu, L. 2001. Altered protamine 2 expression is uncommon in donors of known fertility, but common among men with poor fertilizing capacity, and may reflect other abnormalities of spermiogenesis, *J Androl* 22: 604-10.

Chang, A. S., Moley, K. H., Wangler, M., Feinberg, A. P. and Debaun, M. R. 2005. Association between Beckwith-Wiedemann syndrome and assisted reproductive technology: a case series of 19 patients, *Fertil Steril* 83: 349-54.

Chevaillier, P., Mauro, N., Feneux, D., Jouannet, P. and David, G. 1987. Anomalous protein complement of sperm nuclei in some infertile men, *Lancet* 2: 806-7.

Chiba, H., Hiura, H., Okae, H., Miyauchi, N., Sato, F., Sato, A. and Arima, T. 2013. DNA methylation errors in imprinting disorders and assisted reproductive technology, *Pediatr Int* 55: 542-9.

Cho, C., Willis, W. D., Goulding, E. H., Jung-Ha, H., Choi, Y. C., Hecht, N. B. and Eddy, E. M. 2001. Haploinsufficiency of protamine-1 or -2 causes infertility in mice, *Nat Genet* 28: 82-6.

Cleaton, M. A., Edwards, C. A. and Ferguson-Smith, A. C. 2014. Phenotypic outcomes of imprinted gene models in mice: elucidation of pre- and postnatal functions of imprinted genes, *Annu Rev Genomics Hum Genet* 15: 93-126.

Cox, G. F., Burger, J., Lip, V., Mau, U. A., Sperling, K., Wu, B. L. and Horsthemke, B. 2002. Intracytoplasmic sperm injection may increase the risk of imprinting defects, *Am J Hum Genet* 71: 162-4.

Cumming, J. A. and Carrell, D. T. 2009. Utility of reflexive semen cultures

for detecting bacterial infections in patients with infertility and leukocytospermia, *Fertil Steril* 91: 1486-8.

Dada, R., Kumar, M., Jesudasan, R., Fernandez, J. L., Gosalvez, J. and Agarwal, A. 2012. Epigenetics and its role in male infertility, *J Assist Reprod Genet* 29: 213-23.

Davis, T. L., Trasler, J. M., Moss, S. B., Yang, G. J. and Bartolomei, M. S. 1999. Acquisition of the H19 methylation imprint occurs differentially on the parental alleles during spermatogenesis, *Genomics* 58: 18-28.

Davis, T. L., Yang, G. J., McCarrey, J. R. and Bartolomei, M. S. 2000. The H19 methylation imprint is erased and re-established differentially on the parental alleles during male germ cell development, *Hum Mol Genet* 9: 2885-94.

DeBaun, M. R., Niemitz, E. L. and Feinberg, A. P. 2003. Association of in vitro fertilization with Beckwith-Wiedemann syndrome and epigenetic alterations of LIT1 and H19, *Am J Hum Genet* 72: 156-60.

DeChiara, T. M., Robertson, E. J. and Efstratiadis, A. 1991. Parental imprinting of the mouse insulin-like growth factor II gene, *Cell* 64: 849-59.

Dohle, G. R., Halley, D. J., Van Hemel, J. O., van den Ouwel, A. M., Pieters, M. H., Weber, R. F. and Govaerts, L. C. 2002. Genetic risk factors in infertile men with severe oligozoospermia and azoospermia, *Hum Reprod* 17: 13-6.

Emiliani, S., Van den Bergh, M., Vannin, A. S., Biramane, J., Verdoodt, M. and Englert, Y. 2000. Increased sperm motility after in-vitro culture of testicular biopsies from obstructive azoospermic patients results in better post-thaw recovery rate, *Hum Reprod* 15: 2371-4.

Fernandes, S., Huellen, K., Goncalves, J., Dukal, H., Zeisler, J., Rajpert De Meyts, E., Skakkebaek, N. E., Habermann, B., Krause, W., Sousa, M. et al. 2002. High frequency of DAZ1/DAZ2 gene deletions in patients with severe oligozoospermia, *Mol Hum Reprod* 8: 286-98.

Gabory, A., Ferry, L., Fajardy, I., Jouneau, L., Gothie, J. D., Vige, A., Fleur, C., Mayeur, S., Gallou-Kabani, C., Gross, M. S. et al. 2012. Maternal diets trigger sex-specific divergent trajectories of gene expression and

epigenetic systems in mouse placenta, PLoS One 7: e47986.

Gianotten, J., van der Veen, F., Alders, M., Leschot, N. J., Tanck, M. W., Land, J. A., Kremer, J. A., Hoefsloot, L. H., Mannens, M. M., Lombardi, M. P. et al. 2003. Chromosomal region 11p15 is associated with male factor subfertility, Mol Hum Reprod 9: 587-92.

Gicquel, C., Gaston, V., Mandelbaum, J., Siffroi, J. P., Flahault, A. and Le Bouc, Y. 2003. In vitro fertilization may increase the risk of Beckwith-Wiedemann syndrome related to the abnormal imprinting of the KCN1OT gene, Am J Hum Genet 72: 1338-41.

Gold, J. A., Ruth, C., Osann, K., Flodman, P., McManus, B., Lee, H. S., Donkervoort, S., Khare, M., Roof, E., Dykens, E. et al. 2014. Frequency of Prader-Willi syndrome in births conceived via assisted reproductive technology, Genet Med 16: 164-9.

Gosden, R., Trasler, J., Lucifero, D. and Faddy, M. 2003. Rare congenital disorders, imprinted genes, and assisted reproductive technology, Lancet 361: 1975-7.

Govarts, E., Nieuwenhuijsen, M., Schoeters, G., Ballester, F., Bloemen, K., de Boer, M., Chevrier, C., Eggesbo, M., Guxens, M., Kramer, U. et al. 2012. Birth weight and prenatal exposure to polychlorinated biphenyls (PCBs) and dichlorodiphenyldichloroethylene (DDE): a meta-analysis within 12 European Birth Cohorts, Environ Health Perspect 120: 162-70.

Guo, Y. L., Hsu, P. C., Hsu, C. C. and Lambert, G. H. 2000. Semen quality after prenatal exposure to polychlorinated biphenyls and dibenzofurans, Lancet 356: 1240-1.

Hammoud, S. S., Purwar, J., Pflueger, C., Cairns, B. R. and Carrell, D. T. 2010. Alterations in sperm DNA methylation patterns at imprinted loci in two classes of infertility, Fertil Steril 94: 1728-33.

Hata, K., Okano, M., Lei, H. and Li, E. 2002. Dnmt3L cooperates with the Dnmt3 family of de novo DNA methyltransferases to establish maternal imprints in mice, Development 129: 1983-93.

Haugen, V., Pechacek, J., Maher, T., Wilde, J., Kula, L. and Powell, J. 2011. Decreasing pressure ulcer risk during hospital procedures: a rapid process improvement workshop, J Wound Ostomy Continence Nurs 38:

155-9.

Hayashi, K., Yoshida, K. and Matsui, Y. 2005. A histone H3 methyltransferase controls epigenetic events required for meiotic prophase, *Nature* 438: 374-8.

Hiura, H., Obata, Y., Komiyama, J., Shirai, M. and Kono, T. 2006. Oocyte growth-dependent progression of maternal imprinting in mice, *Genes Cells* 11: 353-61.

Hiura, H., Okae, H., Chiba, H., Miyauchi, N., Sato, F., Sato, A. and Arima, T. 2014. Imprinting methylation errors in ART, *Reprod Med Biol* 13: 193-202.

Honda, S., Weigel, A., Hjelmeland, L. M. and Handa, J. T. 2001. Induction of telomere shortening and replicative senescence by cryopreservation, *Biochem Biophys Res Commun* 282: 493-8.

Horsthemke, B. and Buiting, K. 2006. Imprinting defects on human chromosome 15, *Cytogenet Genome Res* 113: 292-9.

Houshdaran, S., Cortessis, V. K., Siegmund, K., Yang, A., Laird, P. W. and Sokol, R. Z. 2007. Widespread epigenetic abnormalities suggest a broad DNA methylation erasure defect in abnormal human sperm, *PLoS One* 2.

Jacob, K. J., Robinson, W. P. and Lefebvre, L. 2013. Beckwith-Wiedemann and Silver-Russell syndromes: opposite developmental imbalances in imprinted regulators of placental function and embryonic growth, *Clin Genet* 84: 326-34.

John, R. M. and Lefebvre, L. 2011. Developmental regulation of somatic imprints, *Differentiation* 81: 270-80.

Kagami, M., Nagai, T., Fukami, M., Yamazawa, K. and Ogata, T. 2007. Silver-Russell syndrome in a girl born after in vitro fertilization: partial hypermethylation at the differentially methylated region of PEG1/MEST, *J Assist Reprod Genet* 24: 131-6.

Kaneda, A., Wakazono, K., Tsukamoto, T., Watanabe, N., Yagi, Y., Tatematsu, M., Kaminishi, M., Sugimura, T. and Ushijima, T. 2004. Lysyl oxidase is a tumor suppressor gene inactivated by methylation and loss of heterozygosity in human gastric cancers, *Cancer Res* 64: 6410-5.

- 1 Keck, C., Gerber-Schafer, C., Clad, A., Wilhelm, C. and Breckwoldt, M. 1998.  
2 Seminal tract infections: impact on male fertility and treatment options,  
3 Hum Reprod Update 4: 891-903.
- 4 Kobayashi, H., Hiura, H., John, R. M., Sato, A., Otsu, E., Kobayashi, N.,  
5 Suzuki, R., Suzuki, F., Hayashi, C., Utsunomiya, T. et al. 2009. DNA  
6 methylation errors at imprinted loci after assisted conception originate  
7 in the parental sperm, Eur J Hum Genet 17: 1582-91.
- 8 Kobayashi, H., Sato, A., Otsu, E., Hiura, H., Tomatsu, C., Utsunomiya, T.,  
9 Sasaki, H., Yaegashi, N. and Arima, T. 2007. Aberrant DNA methylation  
10 of imprinted loci in sperm from oligospermic patients, Hum Mol Genet  
11 16: 2542-51.
- 12 Kondrashov, A. 2012. Genetics: The rate of human mutation, Nature 488:  
13 467-8.
- 14 Korrovits, P., Ausmees, K., Mandar, R. and Punab, M. 2008. Prevalence of  
15 asymptomatic inflammatory (National Institutes of Health Category IV)  
16 prostatitis in young men according to semen analysis, Urology 71:  
17 1010-5.
- 18 Lee, J., Inoue, K., Ono, R., Ogonuki, N., Kohda, T., Kaneko-Ishino, T., Ogura,  
19 A. and Ishino, F. 2002. Erasing genomic imprinting memory in mouse  
20 clone embryos produced from day 11.5 primordial germ cells,  
21 Development 129: 1807-17.
- 22 Leighton, P. A., Ingram, R. S., Eggenschwiler, J., Efstratiadis, A. and  
23 Tilghman, S. M. 1995. Disruption of imprinting caused by deletion of the  
24 H19 gene region in mice, Nature 375: 34-9.
- 25 Li, J. Y., Lees-Murdock, D. J., Xu, G. L. and Walsh, C. P. 2004. Timing of  
26 establishment of paternal methylation imprints in the mouse, Genomics  
27 84: 952-60.
- 28 Ligon, B. L. 2005. Albert Ludwig Sigismund Neisser: discoverer of the cause  
29 of gonorrhea, Semin Pediatr Infect Dis 16: 336-41.
- 30 Lillycrop, K. A., Phillips, E. S., Jackson, A. A., Hanson, M. A. and Burdge, G.  
31 C. 2005. Dietary protein restriction of pregnant rats induces and folic  
32 acid supplementation prevents epigenetic modification of hepatic gene  
33 expression in the offspring, J Nutr 135: 1382-6.

- 1 Lillycrop, K. A., Slater-Jefferies, J. L., Hanson, M. A., Godfrey, K. M.,  
2 Jackson, A. A. and Burdge, G. C. 2007. Induction of altered epigenetic  
3 regulation of the hepatic glucocorticoid receptor in the offspring of rats  
4 fed a protein-restricted diet during pregnancy suggests that reduced  
5 DNA methyltransferase-1 expression is involved in impaired DNA  
6 methylation and changes in histone modifications, *Br J Nutr* 97:  
7 1064-73.
- 8 Lucifero, D., Mann, M. R., Bartolomei, M. S. and Trasler, J. M. 2004.  
9 Gene-specific timing and epigenetic memory in oocyte imprinting, *Hum*  
10 *Mol Genet* 13: 839-49.
- 11 Ludwig, M., Katalinic, A., Gross, S., Sutcliffe, A., Varon, R. and Horsthemke,  
12 B. 2005. Increased prevalence of imprinting defects in patients with  
13 Angelman syndrome born to subfertile couples, *J Med Genet* 42: 289-91.
- 14 MacLennan, N. K., James, S. J., Melnyk, S., Piroozzi, A., Jernigan, S., Hsu, J.  
15 L., Janke, S. M., Pham, T. D. and Lane, R. H. 2004. Uteroplacental  
16 insufficiency alters DNA methylation, one-carbon metabolism, and  
17 histone acetylation in IUGR rats, *Physiol Genomics* 18: 43-50.
- 18 Maher, E. R. 2005. Imprinting and assisted reproductive technology, *Hum*  
19 *Mol Genet* 14 Spec No 1: R133-8.
- 20 Maher, E. R., Brueton, L. A., Bowdin, S. C., Luharia, A., Cooper, W., Cole, T.  
21 R., Macdonald, F., Sampson, J. R., Barratt, C. L., Reik, W. et al. 2003.  
22 Beckwith-Wiedemann syndrome and assisted reproduction technology  
23 (ART), *J Med Genet* 40: 62-4.
- 24 Marques, C. J., Carvalho, F., Sousa, M. and Barros, A. 2004. Genomic  
25 imprinting in disruptive spermatogenesis, *Lancet* 363: 1700-2.
- 26 Marques, C. J., Costa, P., Vaz, B., Carvalho, F., Fernandes, S., Barros, A. and  
27 Sousa, M. 2008. Abnormal methylation of imprinted genes in human  
28 sperm is associated with oligozoospermia, *Mol Hum Reprod* 14: 67-74.
- 29 Marques, C. J., Francisco, T., Sousa, S., Carvalho, F., Barros, A. and Sousa,  
30 M. 2010. Methylation defects of imprinted genes in human testicular  
31 spermatozoa, *Fertil Steril* 94: 585-94.
- 32 McAuliffe, M. E., Williams, P. L., Korrick, S. A., Dadd, R. and Perry, M. J.  
33 2012. The association between sperm sex chromosome disomy and

1 semen concentration, motility and morphology, *Hum Reprod* 27:  
2 2918-26.

3 McGrath, J. and Solter, D. 1984. Completion of mouse embryogenesis  
4 requires both the maternal and paternal genomes, *Cell* 37: 179-83.

5 Meaney, M. J., Szyf, M. and Seckl, J. R. 2007. Epigenetic mechanisms of  
6 perinatal programming of hypothalamic-pituitary-adrenal function and  
7 health, *Trends Mol Med* 13: 269-77.

8 Meeker, J. D., Ehrlich, S., Toth, T. L., Wright, D. L., Calafat, A. M., Trisini, A.  
9 T., Ye, X. and Hauser, R. 2010. Semen quality and sperm DNA damage in  
10 relation to urinary bisphenol A among men from an infertility clinic,  
11 *Reprod Toxicol* 30: 532-9.

12 Milagro, F. I., Campion, J., Garcia-Diaz, D. F., Goyenechea, E., Paternain, L.  
13 and Martinez, J. A. 2009. High fat diet-induced obesity modifies the  
14 methylation pattern of leptin promoter in rats, *J Physiol Biochem* 65:  
15 1-9.

16 Miura, K. and Niikawa, N. 2005. Do monozygotic dizygotic twins increase  
17 after pregnancy by assisted reproductive technology?, *J Hum Genet* 50:  
18 1-6.

19 Moll, A. C., Imhof, S. M., Cruysberg, J. R., Schouten-van Meeteren, A. Y.,  
20 Boers, M. and van Leeuwen, F. E. 2003. Incidence of retinoblastoma in  
21 children born after in-vitro fertilisation, *Lancet* 361: 309-10.

22 Morgan, H. D., Santos, F., Green, K., Dean, W. and Reik, W. 2005. Epigenetic  
23 reprogramming in mammals, *Hum Mol Genet* 14 Spec No 1: R47-58.

24 Niemitz, E. L. and Feinberg, A. P. 2004. Epigenetics and assisted  
25 reproductive technology: a call for investigation, *Am J Hum Genet* 74:  
26 599-609.

27 Obata, Y. and Kono, T. 2002. Maternal primary imprinting is established at a  
28 specific time for each gene throughout oocyte growth, *J Biol Chem* 277:  
29 5285-9.

30 Orstavik, K. H., Eiklid, K., van der Hagen, C. B., Spetalen, S., Kierulf, K.,  
31 Skjeldal, O. and Buiting, K. 2003. Another case of imprinting defect in a  
32 girl with Angelman syndrome who was conceived by intracytoplasmic  
33 semen injection, *Am J Hum Genet* 72: 218-9.



- 1 Rakyan, V. K., Chong, S., Champ, M. E., Cuthbert, P. C., Morgan, H. D., Luu,  
2 K. V. and Whitelaw, E. 2003. Transgenerational inheritance of epigenetic  
3 states at the murine Axin(Fu) allele occurs after maternal and paternal  
4 transmission, *Proc Natl Acad Sci U S A* 100: 2538-43.
- 5 Razavi, S., Nasr-Esfahani, M. H., Mardani, M., Mafi, A. and Moghdam, A.  
6 2003. Effect of human sperm chromatin anomalies on fertilization  
7 outcome post-ICSI, *Andrologia* 35: 238-43.
- 8 Richthoff, J., Rylander, L., Jonsson, B. A., Akesson, H., Hagmar, L.,  
9 Nilsson-Ehle, P., Stridsberg, M. and Giwercman, A. 2003. Serum levels  
10 of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) in relation to markers of  
11 reproductive function in young males from the general Swedish  
12 population, *Environ Health Perspect* 111: 409-13.
- 13 Rossignol, S., Steunou, V., Chalas, C., Kerjean, A., Rigolet, M.,  
14 Viegas-Pequignot, E., Jouannet, P., Le Bouc, Y. and Gicquel, C. 2006.  
15 The epigenetic imprinting defect of patients with Beckwith-Wiedemann  
16 syndrome born after assisted reproductive technology is not restricted to  
17 the 11p15 region, *J Med Genet* 43: 902-7.
- 18 Roth, T. L., Zoladz, P. R., Sweatt, J. D. and Diamond, D. M. 2011. Epigenetic  
19 modification of hippocampal Bdnf DNA in adult rats in an animal model  
20 of post-traumatic stress disorder, *J Psychiatr Res* 45: 919-26.
- 21 Rousseaux, S., Caron, C., Govin, J., Lestrat, C., Faure, A. K. and Khochbin, S.  
22 2005. Establishment of male-specific epigenetic information, *Gene* 345:  
23 139-53.
- 24 Sato, A., Hiura, H., Okae, H., Miyauchi, N., Abe, Y., Utsunomiya, T.,  
25 Yaegashi, N. and Arima, T. 2011. Assessing loss of imprint methylation  
26 in sperm from subfertile men using novel methylation polymerase chain  
27 reaction Luminex analysis, *Fertil Steril* 95: 129-34, 134 e1-4.
- 28 Sato, A., Otsu, E., Negishi, H., Utsunomiya, T. and Arima, T. 2007. Aberrant  
29 DNA methylation of imprinted loci in superovulated oocytes, *Hum*  
30 *Reprod* 22: 26-35.
- 31 Shimizu, Y., Fukuda, J., Sato, W., Kumagai, J., Hirano, H. and Tanaka, T.  
32 2004. First-trimester diagnosis of conjoined twins after in-vitro  
33 fertilization-embryo transfer (IVF-ET) at blastocyst stage, *Ultrasound*

Obstet Gynecol 24: 208-9.

Sun, J. X., Helgason, A., Masson, G., Ebenesersdottir, S. S., Li, H., Mallick, S., Gnerre, S., Patterson, N., Kong, A., Reich, D. et al. 2012. A direct characterization of human mutation based on microsatellites, *Nat Genet* 44: 1161-5.

Surani, M. A. 1998. Imprinting and the initiation of gene silencing in the germ line, *Cell* 93: 309-12.

Surani, M. A., Barton, S. C. and Norris, M. L. 1984. Development of reconstituted mouse eggs suggests imprinting of the genome during gametogenesis, *Nature* 308: 548-50.

Thompson, J. R. and Williams, C. J. 2005. Genomic imprinting and assisted reproductive technology: connections and potential risks, *Semin Reprod Med* 23: 285-95.

Tilghman, S. M. 1999. The sins of the fathers and mothers: genomic imprinting in mammalian development, *Cell* 96: 185-93.

Toft, G. and Guillette, L. J., Jr. 2005. Decreased sperm count and sexual behavior in mosquitofish exposed to water from a pesticide-contaminated lake, *Ecotoxicol Environ Saf* 60: 15-20.

Toft, G., Rignell-Hydbom, A., Tyrkiel, E., Shvets, M., Giwercman, A., Lindh, C. H., Pedersen, H. S., Ludwicki, J. K., Lesovoy, V., Hagmar, L. et al. 2006. Semen quality and exposure to persistent organochlorine pollutants, *Epidemiology* 17: 450-8.

Tunster, S. J., Van de Pette, M. and John, R. M. 2011. Fetal overgrowth in the *Cdkn1c* mouse model of Beckwith-Wiedemann syndrome, *Dis Model Mech* 4: 814-21.

Ueda, T., Abe, K., Miura, A., Yuzuriha, M., Zubair, M., Noguchi, M., Niwa, K., Kawase, Y., Kono, T., Matsuda, Y. et al. 2000. The paternal methylation imprint of the mouse H19 locus is acquired in the gonocyte stage during foetal testis development, *Genes Cells* 5: 649-59.

Ushijima, T., Nakajima, T. and Maekita, T. 2006. DNA methylation as a marker for the past and future, *J Gastroenterol* 41: 401-7.

van Montfoort, A. P., Hanssen, L. L., de Sutter, P., Viville, S., Geraedts, J. P. and de Boer, P. 2012. Assisted reproduction treatment and epigenetic

inheritance, *Hum Reprod Update* 18: 171-97.

Vucetic, Z., Kimmel, J. and Reyes, T. M. 2011. Chronic high-fat diet drives postnatal epigenetic regulation of mu-opioid receptor in the brain, *Neuropsychopharmacology* 36: 1199-206.

Waterland, R. A. and Jirtle, R. L. 2003. Transposable elements: targets for early nutritional effects on epigenetic gene regulation, *Mol Cell Biol* 23: 5293-300.

Waterland, R. A. and Jirtle, R. L. 2004. Early nutrition, epigenetic changes at transposons and imprinted genes, and enhanced susceptibility to adult chronic diseases, *Nutrition* 20: 63-8.

Waterland, R. A., Lin, J. R., Smith, C. A. and Jirtle, R. L. 2006. Post-weaning diet affects genomic imprinting at the insulin-like growth factor 2 (Igf2) locus, *Hum Mol Genet* 15: 705-16.

Weaver, I. C., Cervoni, N., Champagne, F. A., D'Alessio, A. C., Sharma, S., Seckl, J. R., Dymov, S., Szyf, M. and Meaney, M. J. 2004. Epigenetic programming by maternal behavior, *Nat Neurosci* 7: 847-54.

Weber, M., Hellmann, I., Stadler, M. B., Ramos, L., Paabo, S., Rebhan, M. and Schubeler, D. 2007. Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome, *Nat Genet* 39: 457-66.

Wolff, G. L., Roberts, D. W. and Mountjoy, K. G. 1999. Physiological consequences of ectopic agouti gene expression: the yellow obese mouse syndrome, *Physiol Genomics* 1: 151-63.

Wolff, H. 1995. The biologic significance of white blood cells in semen, *Fertil Steril* 63: 1143-57.

**Figure legends**

**Figure 1. The regulation of imprinted genes by DNA methylation.** Genomic imprinting describes the differential expression of the two parental alleles in mammals (and some plants). This differential expression is initiated within the germline when discrete regions of the genome acquire DNA methylation in one germline but not the other. These differentially methylated regions (DMRs) are present within all well characterised imprinted loci and are key to establishing and, in some cases, maintaining imprinted gene expression. Paternal: paternal allele; Maternal: maternal allele; ICR: imprint control region; TF: transcriptional factor.

**Figure 2. Imprints in gametogenesis and ART procedure.** (Upper: Oogenesis) During the transition from primordial to antral follicles in the postnatal growth phase (post-pachytene) methylation is acquired asynchronously in a gene-specific manner in mouse oogenesis. In sperm, imprint methylation is initiated prenatally before meiosis and is completed by the pachytene phase of postnatal spermatogenesis. The imprints of gametes are maintained stably in the early embryo despite overall epigenetic reprogramming. (Lower: Spermatogenesis) ART results from the use of sperm with incomplete reprogramming and from in vitro embryo procedures performed at the time of epigenetic reprogramming. IVM: *in vitro* oocyte maturation; GIFT: gamete intrafallopian transfer; ZIFT: zygote intrafallopian transfer; PGD: preimplantation genetic diagnosis; IVF: *in vitro* fertilization; ICSI: intracytoplasmic sperm injection; ROSI: round spermatid injection; PGC: primordial germ cell; Oog: oogonium; POo:

primary oocyte; ProSpg: prospermatogonium; Spg: spermatogonium; PSp: primary spermatocyte; SSp: secondary spermatocyte.

**Figure 3. Aberrant DNA methylation of imprinted loci in sperm from infertile male.** (A) Frequency of imprint methylation errors (B) Abnormal imprinted loci (C) Abnormal methylation imprinting and sperm concentrations, morphology and motility. Methylation errors at maternal and paternal imprinted loci specific to oligozoospermic men. (D) Model comparing oligozoospermia and epigenetic errors (described in detail by Kobayashi et al. HMG 2007).

**Figure 4. Factors influencing DNA methylation.** DNA methylation is influenced by a number of external factors including nutrition, aging and hormones. Preventive and promotive factors are shown.

**Table 1. ART and imprint-associated disorders.**

BWS: Beckwith-Wiedemann syndrome, AS: Angelman syndrome, SRS: Silver-Russell syndrome, RB: Retinoblastoma.

**Table 2. DNA methylation errors in the human spermatozoa**

OAT: patients presenting with combined oligozoospermia, asthenozoospermia and teratozoospermia, ANJ: Anejaculation, OAZI: secondary inflammatory obstructive azoospermia, CBAVD: obstructive azoospermia due to congenital bilateral absence of

1 the vas deferens, HP: secretory azoospermia due to hypospermatogenesis.

2

1    **Abbreviations**

2

3    ANJ: Anejaculation,

4    ART: assisted reproductive technologies

5    AS: Angelman syndrome

6    Avy: Agouti viable yellow

7    BS: Bisulphite PCR sequence method

8    BWS: Beckwith-Wiedemann syndrome

9    CBAVD: obstructive azoospermia due to congenital bilateral absence of the vas  
10    deferens

11    COBRA: combined bisulphite PCR restriction analysis

12    DNMT: DNA methyltransferase

13    DOHaD: The developmental origins of health and disease

14    gDMRs: gametic differentially methylated regions

15    GIFT: gamete intrafallopian transfer

16    HP: secretory azoospermia due to hypospermatogenesis

17    IAP: intra-cisternal A particle

18    ICRs: imprinting control regions

19    ICSI: intracytoplasmic sperm injection

20    IVF: *in vitro* fertilization

21    IVM: *in vitro* oocyte maturation

22    OAT: oligoasthenoteratozoospermic

- 1 OAZI: secondary inflammatory obstructive azoospermia
- 2 Oog: oogonium
- 3 PGC: primordial germ cell
- 4 PGD: preimplantation genetic diagnosis
- 5 POPs: Persistent organic pollutants
- 6 POo: primary oocyte
- 7 PSp: primary spermatocyte
- 8 PTSD: post-traumatic stress disorder
- 9 ProSpg: prospermatogonium
- 10 PWS : Prader-Willi syndrome
- 11 ROSI: round spermatid injection
- 12 SAM: S - adenosylmethionine
- 13 SRS: Silver-Russell syndrome
- 14 SSp: secondary spermatocyte
- 15 Spg: spermatogonium
- 16 TF: transcriptional factor
- 17 TZ: teratozoospermic
- 18 ZIFT: zygote intrafallopian transfer
- 19



Figure.1






Figure.3



D.



(Kobayashi et al. HMG 2007 Revised)




Figure.4




Figure.2

