**ORCA – Online Research @ Cardiff** 



# This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:https://orca.cardiff.ac.uk/id/eprint/86768/

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Hildebrand, Robert, Oertel, Timm and Weismantel, Robert 2015. Note on the complexity of the mixed-integer hull of a polyhedron. Operations Research Letters 43 (3), pp. 279-282. 10.1016/j.orl.2015.03.002

Publishers page: http://dx.doi.org/10.1016/j.orl.2015.03.002

## Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See http://orca.cf.ac.uk/policies.html for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



## Note on the Complexity of the Mixed-Integer Hull of a Polyhedron

Robert Hildebrand

Timm Oertel

Robert Weismantel

December 8, 2014

#### Abstract

We study the complexity of computing the mixed-integer hull  $\operatorname{conv}(P \cap \mathbb{Z}^n \times \mathbb{R}^d)$  of a polyhedron P. Given an inequality description, with one integer variable, the mixed-integer hull can have exponentially many vertices and facets in d. For n, d fixed, we give an algorithm to find the mixed integer hull in polynomial time. Given  $P = \operatorname{conv}(V)$  and n fixed, we compute a vertex description of the mixed-integer hull in polynomial time and give bounds on the number of vertices of the mixed integer hull.

Keywords --- Mixed-integer hull, polyhedron, mixed-integer concave minimization

#### 1 Introduction

Given a polyhedron  $P \subseteq \mathbb{R}^n \times \mathbb{R}^d$ , we focus on computing the mixed-integer hull  $P_{MI} = \operatorname{conv}(P \cap \mathbb{Z}^n \times \mathbb{R}^d)$ . The mixed-integer hull is a fundamental object in mixed-integer linear programming and is well known to be a polyhedron [12]. In 1992, Cook, Kannan, Hartman, and McDiarmid [5] showed that the integer hull of  $P_I = \operatorname{conv}(P \cap \mathbb{Z}^n)$  has at most  $2m^n (6n^2 \varphi)^{n-1}$  many vertices, where m is the number of facets of P and  $\varphi$  is the maximum binary encoding size of a facet. Hartman [8] gave a polynomial time algorithm in fixeddimension to enumerate the vertices of the integer hull of a polyhedron. See [14] for a survey of these results, improvements, and lower bounds and also [4] for a discussion of implementations. As far as we know, no similar results have been shown for the mixed-integer hull.

We will give two algorithms to compute the mixedinteger hull; each algorithm produces a bound for the number of vertices of the mixed-integer hull. In Section 2, we consider P given as an inequality description. We show that even with one integer variable and the restriction of small subdeterminants of the inequality description, the mixed-integer hull can have exponentially many facets and vertices if d varies. Hence, we fix both the number of continuous and integer variables. Through a simple scaling or decretization technique, we can apply the results of [5] and [8] to compute the integer hull of a scaled polyhedron. By scaling back, we obtain the mixed-integer hull. This leads to a bound on the number of vertices that is exponential in n + d.

In Section 3, we consider P given by a list of vertices and extreme rays. In this setting, we can allow dto vary. We reduce the original task of computing the mixed-integer hull of a polyhedron to the special case of polytopes by writing an extended formulation using a Minkowski-Weyl type decomposition of P. Hence we can assume that P is bounded and given the vertices of P, we show how to compute a vertex description of  $P_{MI}$  that runs in polynomial time in the encoding size of the vertices provided that the number of integer variables is fixed. This algorithm implies a better bound on the number of vertices of the mixed-integer hull that depends on the number of vertices of the original polytope and does not depend on the number of continuous variables d. This algorithm also implies an algorithm for concave minimization over the mixed-integer points in a polytope since the solution lies at an extreme point.

**Theorem 1** (Mixed-integer concave minimization). Let  $V \subseteq \mathbb{Q}^{n+d}$ ,  $P = \operatorname{conv}(V)$ , and  $f \colon \mathbb{R}^{n+d} \to \mathbb{R}$ be concave. When n is fixed, the problem  $\min\{f(x,y) \colon (x,y) \in P \cap \mathbb{Z}^n \times \mathbb{R}^d\}$  can be solved in polynomial time in the evaluation time of f, d, |V| and  $\nu$ , where  $\nu$  is the maximum binary encoding size of a point in V.

Concave minimization over polyhedra presented by an inequality description is NP-Hard when the dimension varies, even for the case of minimizing a concave quadratic function over the continuous points in a cube. This is because every extreme point can be a local minimum. Most exact algorithms require in the worst case to enumerate all extreme points of the feasible region [11] which can be of exponential size in the number of inequalities. In order to cope with these complexity results, it appears natural to assume that the underlying polyhedron is presented by means of its vertices. From the viewpoint of optimization, Theorem 1 is the mixed-integer analogue of integer concave minimization based on integer hull computations in fixed dimension for which vertex complexity and inequality complexity of the polyhedron are polynomial time equivalent.

**Notation:** For a set  $Q \subseteq \mathbb{R}^n \times \mathbb{R}^d$ , we define  $Q_{\hat{x}} = \{(x, y) \in Q : x = \hat{x}\} \subseteq \mathbb{R}^n \times \mathbb{R}^d$  and  $\operatorname{proj}_x(Q) \subseteq \mathbb{R}^n$  as the projection of Q onto the first n variables.

## 2 Mixed-integer hull from inequalities

We study the problem of computing the mixed-integer hull of a polyhedron P when we are given an inequality description of P. Unfortunately, in this setting, it may be impossible to give a compact facet or vertex representation of the mixed-integer hull if d is allowed to vary. This is demonstrated in the following example that has only one integer variable. Notice that in the example, the maximum subdeterminant in absolute value of A is two.



Figure 1: Left: The mixed integer hull of the rotated cube in Example 2 with d+1 = 3 and  $b_i = 2^i+1$ . Right: The example with  $b_i = 3$  mentioned in Remark 3. In both cases, every vertex of P has been cut off, resulting in exponentially many facets and vertices.

**Example 2.** Let  $P = R \prod_{i=1}^{d+1} [-b_i, b_i] = \{x \in \mathbb{R}^{d+1} : -b_i \leq A_i \cdot x \leq b_i, \forall i\}$  be the linear transformation of the hypercube with

$$R = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & \dots & \frac{1}{2} \\ 0 & I_d \end{bmatrix}, \quad A = \begin{bmatrix} 2 & -1 & \dots & -1 \\ 0 & I_d \end{bmatrix},$$

where  $I_d$  is the  $d \times d$  identity matrix. Choosing odd numbers  $b_i = 2^i + 1$ , the first coordinates of neighboring vertices of P are contained in the interiors of distinct and separated unit intervals. It follows that the mixedinteger hull  $P_{MI} = \operatorname{conv}(P \cap \mathbb{Z} \times \mathbb{R}^d)$  does not contain any of the vertices of P. In particular, it can be shown that every vertex of P violates a distinct facet defining inequality of  $P_{MI}$ . See Figure 1.

**Remark 3.** In Example 2, if we instead choose  $b_i = 3$ , the mixed integer hull  $P_{MI}$  will still have exponentially many vertices and facets. In this case though, the first coordinate of all vertices is polynomially bounded by d, which allows one to write an extended formulation of  $P_{MI}$  as the convex hull of the union of polynomially many integer fibers. By [2], this leads to a polynomial size representation. Note that it could still be possible that for the case of  $b_i = 2^i + 1$  there is a polynomial size extended formulation for  $P_{MI}$ .

In view of Example 2, we first restrict ourselves to the setting where n, d are fixed. We present a scaling algorithm that can be used to apply previous results on computing the integer hull. This technique is similar to [6] where they show that the mixed-integer split closure of a polyhedron is again a polyhedron by a scaling argument to transform the mixed-integer linear program to an integer linear program.

Let  $P = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^d : (A_1, A_2)(x, y) \leq b\}$ be a polyhedron with  $A_1 \in \mathbb{Z}^{m \times n}, A_2 \in \mathbb{Z}^{m \times d}$  and  $b \in \mathbb{Z}^m$ . We compute the mixed-integer hull  $P_{MI} =$  $\operatorname{conv}(P \cap \mathbb{Z}^n \times \mathbb{R}^d)$  in time polynomial in the binary encoding size of  $A_1, A_2$  and b. The output is either a facet or a vertex description of  $P_{MI}$ . Since n, d are both fixed, the facet and vertex descriptions are polynomial time equivalently computable (see, for instance, [12]).

For a polytope  $Q \subseteq \mathbb{R}^n \times \mathbb{R}^d$ , define  $Q^t := \{(x, ty) \in \mathbb{R}^n \times \mathbb{R}^d : (x, y) \in Q\}$ . Therefore,  $Q^1 = Q$ . Notice that we can write

$$P^{t} = \{ (x, y) \in \mathbb{R}^{n} \times \mathbb{R}^{d} : (A_{1}, A_{2})(x, \frac{1}{t}y) \leq b \}$$
$$= \{ (x, y) \in \mathbb{R}^{n} \times \mathbb{R}^{d} : (A_{1}, \frac{1}{t}A_{2})(x, y) \leq b \}.$$

**Theorem 4** (Mixed-integer hull from inequality description). The mixed-integer hull can be computed in time polynomial in the binary encoding size of  $A_1, A_2, b$ , provided that n and d are fixed. Furthermore, let  $\phi$  be the maximum binary encoding size of a row of  $(A_1, A_2, b)$ . Then  $|\operatorname{vert}(P_{MI})| \leq 2m^{n+d}(6(n + d)^2 \varphi)^{n+d-1}$  where  $\varphi = \phi + n\phi(m+n)^{n+d}$ .

*Proof.* We first show that we can compute in polynomial time an integer t such that  $P_I^t = P_{MI}^t$ . Let  $(\hat{x}, \hat{y})$  be a vertex of  $P_{MI}$ . Notice that  $(\hat{x}, \hat{y})$  is a vertex of the d-dimensional polyhedron  $P_{\hat{x}} = \operatorname{conv}(P \cap \{(x, y) : x = \hat{x}\})$ . We analyze the vertices of  $P_{\hat{x}}$  by considering its inequality description as  $P_{\hat{x}} = \{(x, y) : A(x, y) \leq \bar{b}\}$  where

$$A = \begin{bmatrix} A_1 & A_2 \\ I_n & \mathcal{O}_d \\ -I_n & \mathcal{O}_d \end{bmatrix} \in \mathbb{Z}^{(m+2n) \times (n+d)}$$

and  $\overline{b} = (b, \hat{x}, -\hat{x})$ . Here  $\mathcal{O}_d$  is the  $d \times d$  matrix of all 0's. Therefore, there exists a basis B of the rows of A of size n + d such that  $(\hat{x}, \hat{y}) = (A_B)^{-1} (b, \hat{x}, -\hat{x})_B$ . By

Cramer's rule, for any  $i \in B$ , we have

$$(\hat{x}, \hat{y})_i = \frac{\det(A_B^i)}{\det(A_B)}$$

where  $A_B^i$  is the matrix  $A_B$  where the  $i^{\text{th}}$  column is replaced by the vector  $(b, \hat{x}, -\hat{x})_B$ . Therefore  $\det(A_B^i) \in \mathbb{Z}$ . Hence  $(\hat{x}, \hat{y})_i \cdot \det(A_B) \in \mathbb{Z}$ , that is,  $\hat{y} \cdot \det(A_B) \in \mathbb{Z}^d$ is integral. Note that  $\det(A_B)$  is completely independent of  $(\hat{x}, \hat{y})$ . Now if we let

$$t := \prod_{B \text{ basis}} \det(A_B),$$

then (x, ty) is integral for any basis B of A. There are at most  $\binom{m+n}{n+d} \leq (m+n)^{n+d}$  many bases B. By Hadamard's inequality,  $\det(A_B) \leq \prod_{i=1}^{n} 2^{\phi} = 2^{n\phi}$ . Therefore,  $t \leq (2^{n\phi})^{(m+n)^{(n+d)}} = 2^{n\phi(m+n)^{n+d}}$ .

Since  $P_{MI}^t$  is the convex hull of all polyhedra  $P_{\hat{x}}^t$  for  $\hat{x} \in \mathbb{Z}^n$ , and  $P_{\hat{x}}^t$  has integer vertices by our choice of t, we have that  $P_{MI}^t = P_I^t$ . By computing the integer hull  $P_I^t$  using [8], and then scaling back by computing  $(P_I^t)^{1/t} = (P_{MI}^t)^{1/t} = P_{MI}$ , we find a description of the mixed-integer hull. Note that [8] yields a vertex description of the integer hull. This can be converted to an inequality description in polynomial time since the dimension is fixed. By [5],  $|\operatorname{vert}(P_{MI})| = |\operatorname{vert}(P_I^t)| \leq 2m^{n+d}(6(n+d)^2\varphi)^{n+d-1}$  where  $\varphi = \phi + n\phi(m+n)^{n+d}$ .

## 3 Mixed-integer hull from vertices

We now consider the case where we are given P as a list of vertices and extreme rays and we show how to compute a vertex and extreme ray description of the mixed-integer hull. In this setting, we show that the number of continuous variables d may vary, and we still obtain a polynomial time algorithm to compute the mixed-integer hull. We begin by reducing the problem to finding the mixed-integer hull of a polytope.

#### 3.1 Reduction to bounded polyhedra

We adapt a result of Nemhauser and Wolsey [10, Section I.4, Theorem 6.1] to the mixed-integer case. The proof is essentially the same, but we provide it here to obtain complexity bounds. Recall that by the Minkowski-Weyl theorem, a polyhedron P can be represented as an inequality description or a vertex and extreme ray description, that is,  $P = \{x \in \mathbb{R}^{n+d} : Ax \leq b\} = \operatorname{conv}(V) + \operatorname{cone}(W)$  for some sets  $V \subseteq \mathbb{Q}^{n+d}$ ,  $W \subseteq \mathbb{Z}^{n+d}$ . Here,  $\operatorname{rec}(P) = \{x \in \mathbb{R}^{n+d} : Ax \leq 0\} = \operatorname{cone}(W)$ . In the following, we request that  $P \cap \mathbb{Z}^n \times \mathbb{R}^d$ is non-empty, which can be tested in polynomial time provided that n is fixed using Lenstra's algorithm [9].

**Lemma 5** (Relevant mixed-integer points). Let P be a rational polyhedron given by a vertex/ray representation  $\operatorname{conv}(V) + \operatorname{cone}(W)$  for  $V \subseteq \mathbb{Q}^{n+d}$  and  $W \subseteq \mathbb{Z}^{n+d}$ (resp. an inequality description  $\{x \in \mathbb{R}^{n+d} : Ax \leq b\}$  for  $A \in \mathbb{Q}^{m \times (n+d)}$  and  $b \in \mathbb{Q}^m$ ). Suppose that  $P \cap \mathbb{Z}^n \times \mathbb{R}^d \neq \emptyset$ . In polynomial time, we can compute a vertex description (resp. an inequality description) of a rational polytope Q of polynomial size such that  $P = Q + \operatorname{rec}(P)$  and

$$P_{MI} = \operatorname{conv}(Q \cap \mathbb{Z}^n \times \mathbb{R}^d) + \operatorname{rec}(P).$$

*Proof.* By the Minkowski-Weyl theorem, we can decompose P as  $P = \operatorname{conv}(V) + \operatorname{cone}(W)$  where V is the set of vertices of P and W is the set of minimal integral extreme rays of P, and hence  $\operatorname{cone}(W) = \operatorname{rec}(P)$ . For any  $x \in P \cap \mathbb{Z}^n \times \mathbb{R}^d$ , by this decomposition and Carathéodory's theorem we can write

$$x = \sum_{i \in I} \lambda_i v_i + \sum_{j \in J} \mu_j w_j$$
  
=  $\sum_{i \in I} \lambda_i v_i + \sum_{j \in J} (\mu_j - \lfloor \mu_j \rfloor) w_j + \sum_{j \in J} \lfloor \mu_j \rfloor w_j,$ 

where  $\lambda_i, \mu_j \geq 0$ ,  $\sum_{i \in I} \lambda_i = 1$ ,  $v_i \in V$ ,  $w_j \in W$ ,  $|I| \leq n + d + 1$  and  $|J| \leq n + d$ . Since  $x \in \mathbb{Z}^n \times \mathbb{R}^d$  and  $\sum_{j \in J} \lfloor \mu_j \rfloor w_j \in \mathbb{Z}^{n \times d}$ , we have that  $x - \sum_{j \in J} \lfloor \mu_j \rfloor w_j = \sum_{i \in I} \lambda_i v_i + \sum_{j \in J} (\mu_j - \lfloor \mu_j \rfloor) w_j \in \mathbb{Z}^n \times \mathbb{R}^d$ . Therefore, if we define  $T := \operatorname{conv}(V) + (n + d) \operatorname{conv}(W \cup \{0\})$ , then  $x - \sum_{j \in J} \lfloor \mu_j \rfloor w_j \in T \cap \mathbb{Z}^n \times \mathbb{R}^d$ . It follows that

$$\operatorname{conv}(P \cap \mathbb{Z}^n \times \mathbb{R}^d) = \operatorname{conv}(T \cap \mathbb{Z}^n \times \mathbb{R}^d) + \operatorname{rec}(P).$$

If V and W are given as input, then we are done by setting Q = T for which we obtain a vertex description by taking the Minkowski sum V + (n + d)W. On the other hand, if we are given as input an inequality description of P, then the descriptions of V and W may be exponential in the input. In this case, we instead determine a box that contains T and intersect that box with P to obtain our choice of Q.

More precisely, let  $R \ge 0$  be a bound on the infinity norm of the vertices V and an integral representation of the extreme rays W. By, for instance [12], we can choose R of polynomial encoding size. Setting R' =(n+d+1)R, we have  $T \subseteq B := [-R', R']^{n \times d}$ . Setting  $Q = P \cap B$  finishes the argument.  $\Box$ 

#### 3.2 Mixed-integer hull for polytopes

We begin by showing how to compute the vertices of the integer hull of a polytope  $Q = \operatorname{conv}(V)$  presented by its vertex set  $V \subseteq \mathbb{Q}^n$ . To do so, we employ the algorithm from [8] to compute integer hulls from an inequality description, but this requires some care. If we apply a standard transformation of Q into an inequality description, this description could have many facets, causing the algorithm to require a double exponential complexity in terms of n. Therefore, we instead find a triangulation of V and then compute the integer hull of simplices, which have exactly n + 1 facets, allowing us to procure a single exponential complexity for the number of vertices of  $Q_I$ .

**Lemma 6** (Integer hull from vertex description). Let  $V \subseteq \mathbb{Q}^n$  and let  $Q = \operatorname{conv}(V)$ . When n is fixed, we can compute a vertex description of the integer hull  $Q_I$  in polynomial time in  $\nu$  and |V| where  $\nu$  is the maximum binary encoding size of a vector in V. Furthermore,  $|\operatorname{vert}(Q_I)| \leq \frac{1}{3} 12^n n^{3n-2} \varphi^{n-1} |V|^{n+1}$  where  $\varphi = 2(n + 1)^{n+1}$ 

1)<sup>2</sup>( $\nu + \log(n+1)$ ). If |V| = n+1, we obtain the tighter bound of  $| \operatorname{vert}(Q_I) | \leq \frac{2}{3} 24^n n^{3n-2} \varphi^{n-1}$ .

*Proof.* Let  $n' = \dim(\operatorname{conv}(V))$ . We compute a Delaunay triangulation of the points V, yielding a list of n'-dimensional simplices. As is well known, this can be done by computing the convex hull of the extended point set  $V' = \{(v, \|v\|_2^2) : v \in V\} \subset \mathbb{R}^{n+1}$ . Then facets of  $\operatorname{conv}(V')$  corresponds to a n'-dimensional cells of the triangulation. The convex hull can be computed in polynomial time using [1]. See for instance, [7] for a discussion of algorithms to compute the Delaunay triangulation. Moreover, there are at most  $\binom{|V|}{n'+1} \leq |V|^{n+1}$  cells. If |V| = n+1, we can instead use the bound  $2^{n+1}$  for the number of cells. For general |V|, a tighter asymptotic bound on the number of simplices is  $O(|V|^{\lceil n/2 \rceil})$  [13]. For each simplex C in the triangulation, we compute an inequality description. Since it may be lower dimensional, at most 2n inequalities are needed to describe it. These inequalities can be computed in polynomial time using Gaussian elimination and have binary encoding size bounded by  $\varphi = 2(n+1)^2(\nu + \log(n+1))$  which follows from Cramer's rule and Hadamard's inequality.

Finally, we can apply the result of [5] to see  $|\operatorname{vert}(C_I)| \leq 2m^n (6n^2 \varphi)^{n-1}$  where *m* is the number of facets. Here  $m \leq 2n$ , so we have  $|\operatorname{vert}(C_I)| \leq 2(2n)^n (6n^2 \varphi)^{n-1}$ . Finally, applying this to each cell *C*, we have at most  $|V|^{n+1}2(2n)^n (6n^2 \varphi)^{n-1} = \frac{1}{3}12^n n^{3n-2} \varphi^{n-1} |V|^{n+1}$  many vertices of the integer hull, or only  $\frac{2}{3}24^n n^{3n-2} \varphi^{n-1}$  if we choose |V| = n + 1 and use the improved bound mentioned above. By [8], it follows that these can be computed in polynomial time when *n* is fixed. This creates a superset of the vertices of the integer hull; points that are not vertices can be discovered using linear programming, which can be done in polynomial time.

Although this approach may not produce a tight bound on the number of vertices of the integer hull, the bound in Lemma 6 is in the right order of magnitude in terms of  $\varphi$ . Indeed, in [3], they show that for every dimension  $n \geq 2$ , there exists a simplex  $P \subseteq \mathbb{R}^n$  given as an inequality description with encoding size  $\varphi$ such that  $P_I$  has at least  $c_n \varphi^{n-1}$  many vertices, where  $c_n$  is a constant depending only on n. This shows that even if P has a small number of vertices, the number of vertices of the integer hull can be large.

Our next goal is to make use of Lemma 6 in order to compute the mixed-integer hull. This requires one more ingredient.

**Lemma 7.** Every vertex of  $P_{MI}$  lies in a face F of P with  $\dim(F) \leq n$ . Furthermore, let  $n' = \min(n, \dim(P))$  and let  $\mathcal{F}_{n'}$  denote the set of faces of P of dimension n'. Then

$$\operatorname{vert}(P_{MI}) \subseteq \bigcup_{F \in \mathcal{F}_{n'}} \bigcup_{\hat{x} \in \operatorname{vert}(\operatorname{proj}_x(F)_I)} \operatorname{vert}(F_{\hat{x}}).$$
(1)

Proof. Since  $P_{MI} = \operatorname{conv}(P_{\hat{x}} : \hat{x} \in \mathbb{Z}^n)$ , every vertex  $(\bar{x}, \bar{y})$  of  $P_{MI}$  is a vertex of  $P_{\bar{x}}$ . Since  $P_{\bar{x}} = \{(x, y) : A(x, y) \leq b, Ix = \bar{x}\}$  and any vertex of  $P_{\bar{x}}$  is defined uniquely by n + d tight inequality constraints, at least d of those tight constraints come from the inequalities  $A(x, y) \leq b$ . Hence,  $(\bar{x}, \bar{y})$  satisfies at least d affinely independent tight constraints from P, i.e., it is contained in a face F that is n + d - d = n dimensional at most. Since we choosing a maximal such face F, we only consider F of dimension  $n' = \min(n, \dim(P))$ . Furthremore, there exist d tight inequalities  $(\bar{A}_1, \bar{A}_2)(x, y) \leq \bar{b}$  such that  $\bar{A}_2$  is invertible.

Then, for any  $\hat{x} \in \operatorname{proj}_x(F)$ , the corresponding  $\hat{y}$  such that  $(\hat{x}, \hat{y}) \in F$  is given uniquely as

$$\hat{y} = \bar{A}_2^{-1}\bar{b} - \bar{A}_2^{-1}\bar{A}_1\hat{x}.$$

Now let  $(\bar{x}, \bar{y}) \in \operatorname{vert}(P_{MI})$  and suppose that  $\hat{x} \notin \operatorname{vert}(\operatorname{proj}_x(F)_I)$ . Then  $\bar{x} = \sum \mu_i x^i$  for some  $\mu_i > 0$ ,  $\sum \mu_i = 1$ , and  $x^i \in \operatorname{vert}(\operatorname{proj}_x(F)_I)$ . Then setting,  $y^i := \bar{A}_2^{-1}\bar{b} - \bar{A}_2^{-1}\bar{A}_1 x^i$  it follows that  $y^i \in F \cap \mathbb{Z}^n \times \mathbb{R}^d$  and  $(\bar{x}, \bar{y}) = \sum \mu_i(x^i, y^i)$ ; therefore,  $(\bar{x}, \bar{y})$  is not a vertex of  $P_{MI}$ . This proves (1).

**Theorem 8** (Mixed-integer hull from vertex description). Let  $V \subseteq \mathbb{Q}^{n+d}$  and let  $P = \operatorname{conv}(V)$ . For fixed

 $n \geq 1$  there exists an algorithm to compute vert $(P_{MI})$  that runs in polynomial time in  $d, \nu, |V|$  where  $\nu$  is the maximum binary encoding size of a point in V. Furthermore,

$$|\operatorname{vert}(P_{MI})| \le \frac{4}{3} 48^n n^{3n-2} \varphi^{n-1} |V|^{n+1}$$

where  $\varphi = 2(n+1)^2(\nu + \log(n+1)).$ 

Proof. Let  $n' = \min(n, \dim(P)) \leq |V|$  and let  $(\hat{x}, \hat{y})$ be a vertex of  $P_{MI}$ . By Lemma 7, there exists a face F of P of dimension n' such that  $(\hat{x}, \hat{y}) \in \operatorname{vert}(F_{\hat{x}})$ and  $\hat{x} \in \operatorname{vert}(\operatorname{proj}_{x}(F)_{I})$ . By Carathéodory's theorem, there exist n' + 1 vertices of F such that  $(\hat{x}, \hat{y}) \in \bar{F} = \operatorname{conv}(v^{1}, \ldots, v^{n'+1})$ . Since  $\bar{F} \subseteq F$  and  $(\hat{x}, \hat{y}) \in \bar{F}$ , it follows that  $\hat{x} \in \operatorname{vert}(\operatorname{proj}_{x}(\bar{F})_{I})$  and  $(\hat{x}, \hat{y}) \in \operatorname{vert}(\bar{F}_{\hat{x}})$ . Indeed, if  $\hat{x} \notin \operatorname{vert}(\operatorname{proj}_{x}(\bar{F})_{I})$ , then it can be written as a strict convex combination of points in  $\operatorname{vert}(\operatorname{proj}_{x}(\bar{F})_{I}) \subseteq \operatorname{conv}(\operatorname{vert}(\operatorname{proj}_{x}(F)_{I}))$ , which shows that  $\hat{x} \notin \operatorname{vert}(\operatorname{proj}_{x}(F)_{I})$ .

Therefore, we can compute a superset of the vertices of  $P_{MI}$  by enumerating every (n'+1)-elementary subset  $\{v^1, \ldots, v^{n'+1}\}$  of V, yielding at most  $\binom{|V|}{n'+1} \leq |V|^{n'+1}$ sets to consider. Fix a subset  $\{v^1, \ldots, v^{n'+1}\}$  and set  $\bar{F} = \operatorname{conv}(v^1, \ldots, v^{n'+1})$ . We will next find a vertex description of  $\operatorname{conv}(\bar{F} \cap \mathbb{Z}^n \times \mathbb{R}^d)$ .

To this end, let  $\hat{v}^i = \operatorname{proj}_x(v^i)$ . Applying Lemma 6 with  $Q = \operatorname{conv}(\hat{v}^1, \dots, \hat{v}^{n'+1})$ , we have a vertex description of  $Q_I$  with at most  $\frac{2}{3}24^{n'}\varphi^{n'-1}n'^{3n'-2}$  many vertices. For each vertex  $\hat{x}$  of  $Q_I$ , we compute the vertices of  $\bar{F}_{\hat{x}}$ , which can be written as the set of  $(x, y) \in \mathbb{R}^n \times \mathbb{R}^d$  satisfying

$$(x,y) = \sum_{i=1}^{n'+1} \lambda_i v^i, x = \hat{x}, \sum_{i=1}^{n'+1} \lambda_i = 1, \lambda_i \ge 0.$$

Every vertex of this set corresponds to a unique face of the (n' + 1)-dimensional standard simplex. There are  $2^{n'+1}$  many faces of the n'-dimensional simplex. Hence, this set has at most  $2^{n'+1}$  vertices, which can be enumerated, for instance by [1] in time polynomial in the encoding size  $\varphi$  provided that n is fixed. This number of enumerated points provides us with an upper bound on the number of vertices of  $P_{MI}$ . Since we can test whether points are vertices of  $P_{MI}$  using linear programming, the proof is complete.

Combined with Lemma 5, the above theorem gives a similar result for polyhedra.

**Remark 9.** Consider a polyhedron  $P = \{(x, y) \in \mathbb{R}^{n+d} : A(x, y) \leq b, (x, y) \geq 0\}$ , where  $A \in \mathbb{Q}^{m \times (n+d)}$ ,  $b \in \mathbb{Q}^m$ . It follows that  $|\operatorname{vert}(P)| \leq (n+d)^m$ . Therefore, for fixed m and n, using Theorem 8, we can find a vertex description of the mixed-integer hull in polynomial time. This applies, for instance, to a mixed-integer knapsack problem with no upper bounds on the variables.

#### References

- D. Avis and K. Fukuda. A pivoting algorithm for convex hulls and vertex enumeration of arrangements and polyhedra. *Discrete & Computational Geometry*, 8(1):295–313, 1992.
- [2] E. Balas. Disjunctive programming: Properties of the convex hull of feasible points. *Discrete Applied Mathematics*, 89(13):3 – 44, 1998.
- [3] I. Bárány, R. Howe, and L. Lovász. On integer points in polyhedra: A lower bound. *Combinatorica*, 12(2):135–142, 1992.
- [4] P. J. Charles, J. M. Howe, and A. King. Integer polyhedra for program analysis. In A. V. Goldberg and Y. Zhou, editors, *Algorithmic Aspects* in Information and Management, volume 5564 of *Lecture Notes in Computer Science*, pages 85–99. Springer Berlin Heidelberg, 2009.
- [5] W. Cook, M. Hartmann, R. Kannan, and C. Mc-Diarmid. On integer points in polyhedra. *Combinatorica*, 12(1):27–37, 1992.
- [6] W. Cook, R. Kannan, and A. Schrijver. Chvátal closures for mixed integer programming prob-

lems. Mathematical Programming, 47(1-3):155–174, 1990.

- [7] S. Fortune. Handbook of discrete and computational geometry. chapter Voronoi Diagrams and Delaunay Triangulations, pages 377–388. CRC Press, Inc., Boca Raton, FL, USA, 1997.
- [8] M. E. Hartmann. Cutting Planes and the Complexity of the Integer Hull. Phd thesis, Cornell University, Department of Operations Research and Industrial Engineering, Ithaca, NY, 1989.
- [9] H. W. Lenstra, Jr. Integer programming with a fixed number of variables. *Mathematics of Operations Research*, 8:538–548, 1983.
- [10] G. L. Nemhauser and L. A. Wolsey. Integer and Combinatorial Optimization. Wiley-Interscience, New York, NY, USA, 1988.
- [11] P. M. Pardalos and J. B. Rosen. Methods for global concave minimization: A bibliographic survey. SIAM Review, 28(3):pp. 367–379, 1986.
- [12] A. Schrijver. Theory of Linear and Integer Programming. John Wiley and Sons, New York, 1986.
- [13] R. Seidel. The upper bound theorem for polytopes: an easy proof of its asymptotic version. *Computational Geometry*, 5(2):115 – 116, 1995.
- [14] Z. N. Yu. On the number of vertices in integer linear programming problems. eprint arXiv:math/0611356 [math.CO], 2006.