

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/91063/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Morgan, Maria Zeta , Fairchild, R. and Broughton, David James 2016. A survey of sports drinks consumption amongst adolescents. *British Dental Journal* 220 , pp. 639-643. 10.1038/sj.bdj.2016.449

Publishers page: <http://dx.doi.org/10.1038/sj.bdj.2016.449>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



A survey of sports drinks consumption amongst adolescents

D BROUGHTON BDS (Hons)¹

R M FAIRCHILD BSc (Hons), PhD²

M Z MORGAN BSc (Hons), PGCE, MPH, MPhil, FFPH¹

¹Applied Clinical Research and Public Health, College of Biomedical and Life Sciences,
Cardiff University, School of Dentistry, Heath Park, Cardiff CF14 4XY

²Cardiff Metropolitan University, Department of Healthcare and Food, Cardiff CF5 2YB

Key words: oral health, children, sports drinks

Word count including abstract 2871

Abstract – 249

Corresponding author:

Maria Z Morgan (Mrs)

Applied Clinical Research and Public Health,
College of Biomedical and Life Sciences,
Cardiff University,
School of Dentistry,
Heath Park,
Cardiff CF14 4XY

Abstract

Background: Sports drinks intended to improve performance and hydrate athletes taking part in endurance sport are being marketed to children, for whom these products are not intended. Popularity amongst children has grown exponentially, worryingly they consume them socially, as well as during physical activity. Sports drinks are high in sugar and are acidic. Product marketing ignores the potential harmful effects of dental caries and erosion.

Objective: To investigate the use of sports drinks by children.

Method: 183 self-complete questionnaires were distributed to four schools in South Wales. Children in high school years 8 and 9 (aged 12 - 14) were recruited to take part. Questions focussed on use of sports drinks, type consumed, frequency of and reason for consumption and where drinks were purchased.

Results: 160 children responded (87% response rate). 89.4% (143) claimed to drink sports drinks, half drinking them at least twice a week. Lucozade Sport™ was the most popular brand. The main reason for consuming the drinks was attributed to the “nice taste” (90%, 129/143). Most respondents purchased the drinks from local shops (80.4%, 115) or supermarkets (54.5%, 78). More boys claimed to drink sports drinks during physical activity (77.9% versus 48.6% girls, $p < 0.001$). Whereas more girls claimed to drink them socially (51.4% versus 48.5% boys, NS).

Conclusion: A high proportion of children consumed sports drinks regularly and outside of sporting activity. Dental health professionals should be aware of the popularity of sports drinks with children when giving health education advice or designing health promotion initiatives.

INTRODUCTION

Sports drinks have become more popular over recent years with the younger generation^{1,2}. In the 1970s, marathon runners were discouraged from drinking any fluids as it was thought it would slow the runners down. But, now runners are being encouraged to “drink ahead of thirst”³ with the recognition that hydration is associated with performance. Sports drinks are designed to improve performance and hydrate elite adult athletes taking part in endurance and intense sporting events⁴. There is no evidence of beneficial effects in non-elite athletes or children³. However, these drinks are being consumed by the general population during physical activity and socially. It is the latter which is the most cause for concern⁵, particularly amongst children aged under 16³.

The marketing of sports products has become a multibillion-dollar industry⁶. In 2014, the UK sports drinks market was worth £218 million and the core consumers were 15-24 year olds¹. However, the marketing campaigns ignore the detrimental impact sports drinks have on teeth and the effects they can have on general health. Certain marketing campaigns have even misled consumers into incorrectly concluding that the drinks contained no carbohydrates or additives⁶. In addition, studies have shown that while certain supplements in sports drinks can potentially improve performance, many have not been proven to be beneficial and some have harmful side-effects⁶.

If consumed socially and in large quantities, sports drinks can lead to serious problems, such as obesity, diabetes, heart disease and gout⁷, as well as poor oral health⁴. Studies in the past have shown that non-athletes are consuming these drinks simply because of their nice taste².

Sports drinks contain both free sugars and acids⁴, hence these drinks have the ability to cause both dental caries and erosion. There is a strong relationship between eating foods high in “free” sugars and dental caries⁸⁻¹¹. The term free sugars refers to all mono and disaccharides added to foods by the manufacturer, cook or consumer, plus sugars naturally present in honey, fruit juices and syrups¹². Many sports drinks have a pH below 5.5, the critical pH for the demineralisation of enamel, leading to erosion^{13,14}. Dehydration associated with physical activity increases erosion risk, as the buffering capacity is inhibited due to lower salivary flow⁴. Dehydration also reduces clearance of acids and sugars from the tooth surface, affecting both erosion and caries⁴.

The frequency and duration of sports drinks consumption are also important factors impacting on dental health¹⁵. The general consensus is that the frequency and amount of sugary food and drinks should be reduced and, when consumed, limited to mealtimes¹⁶. Certain sports drinks, e.g. Powerade zero™, are marketed as being sugar free. However, these drinks still have an acid content that can lead to erosion of enamel^{13,14}.

A study conducted in the dental clinic at the Olympic Park during the London 2012 Olympics assessed the general oral hygiene of 278 participating athletes¹⁷. The authors reported that 55% of the athletes treated had dental caries and 45% had dental erosion (as measured by the Basic Erosive Wear Examination). This is a much higher prevalence than the experience of “tooth wear” recorded by the UK Adult Dental Health Survey in 2009¹⁸ for similarly aged adults. Whilst the Olympic athletes came from the five continents¹⁷ the authors reported no significant difference in erosion by continental location or ethnicity. They also reported a link between the frequency of sports drinks use and dental erosion in anterior teeth¹⁷.

Recent surveys have shown that Wales has higher levels of decayed, missing and filled teeth (DMFT/dmft) when compared with other areas in the UK, especially in children. In 2013, 52% of children aged 12 in Wales had at least one DMFT in their adult teeth compared with 32% in England¹⁹. Of concern are the wide inequalities in experience, with children from more deprived areas experiencing higher levels of dental disease^{19, 20}.

There is confusion over the difference between energy and sports drinks. An energy drink is marketed for its mental stimulant effect and contains high levels of substances such as caffeine, taurine and glucoronolactone²¹. Sports drinks do not have a stimulant effect, but concentrate on providing carbohydrates, salts and hydration^{1,22}. For the purpose of this study, the participants were not told what constitutes a sports or energy drink, as their knowledge of sports drinks was also being tested.

According to Mintel (2014), 78% of UK 16-24 year olds have reported consuming a sports drink in the last 12 months, with 39% drinking them at least once per week¹. From 1989–2008, the percentage of American children aged 6 to 11 consuming sports drinks increased significantly, from 2% to 12%. The amount of sports drinks consumed by these American children also increased, from 255 millilitres per day to 289 millilitres per day during the same timeframe².

All school meal policies in the devolved countries of the UK have banned the sale of fizzy/sugary drinks other than fruit juices diluted with carbonated water²³⁻²⁶. Therefore students are accessing sports drinks from outside the school gates.

Few studies have investigated the use of sports drinks in children, their knowledge about the product and the impact they may have on their oral health. The aim of this study was to investigate the use of and knowledge surrounding sports drinks by 12-14 year old school children in South Wales, UK. This paper reports on consumption aspects only; the knowledge component is reported separately.

METHOD

This study consisted of a questionnaire survey of 12-14 year old school children recruited from a convenience sample of four secondary schools in South Wales. Schools were selected to reflect the range of deprivation experienced within the area (according to the Welsh Index of Multiple Deprivation, WIMD 2011²⁷). School A was a private school; School B a comprehensive school, whose catchment area draws from the most deprived areas in Wales; Schools C and D were also comprehensive schools, but their catchment area drew from a more mixed demographic²⁷. Furthermore schools C and D provided male and female education one separate sites.

Prior to the commencement of this study ethical approval was gained from the Dental School Ethical Committee at Cardiff University.

A focus group of eight adolescents (aged 12 – 16) was undertaken to inform the questionnaire design. Participants were informally asked whether they drank sports drinks and why?

With information gained from the focus group, a self-complete anonymous questionnaire was designed containing mainly closed questions, allowing categories to be analysed efficiently and with minimum bias²⁸.

Questions reported in this paper were designed to assess:

1. whether respondents consume sports drinks and if so how often and what types?
2. where and when respondents purchase and consume sports drinks?
3. why respondents consume sports drinks?

Children from school years 8 (12-13 years old) and 9 (13-14 years old) were invited to take part in the study from each of the selected schools. Headmasters from each school were asked to nominate one class per school year to take part. A combination of school consent, parental negative consent and child assent was used in this study. Respondents were told that participation was voluntary before they completed the questionnaire. One researcher (DB) was present at each school to distribute and collect the questionnaires and also to answer any questions about the project.

The data were analysed using appropriate descriptive and inferential statistics, such as frequency distributions and chi-squared test for categorical variations (with an alpha value of 0.05 accepted as significant). Statistical analysis was performed with IBM SPSS Statistics (Version 20) software.

RESULTS

Demographic details of the schools and the number of participants from each school and year are presented in Table 1. 183 questionnaires were distributed amongst the four schools. In total, 160 respondents completed the survey (87% response rate); one Year 8 class from school A was unable to take part because of the timing for exams coinciding with data collection. The majority, 89.4% (143/160), of respondents claimed to drink sports drinks. The analyses below relate to the responses from those (n=143) who claimed to drink sports drinks.

Consumption of sports drinks, frequency and type

Almost half of these respondents drank them more than once a week (48.3%, 69); 14% (20) drinking one or more every day. The modal consumption frequency was 2-3 times a week (Figure 1).

Respondents were asked to indicate which of the four bestselling UK sports drinks (Lucozade Sport™, Powerade™, Gatorade™, LSV®) they consumed and an “other” category was also provided. The most popular drink was Lucozade Sport™ (88.8%, 127) whilst Gatorade™ was the least popular (9.8%. 14, Figure 2).

All but one response to the “Other” category were “energy” drinks, not marketed as “sports” drinks. These energy drinks were Relentless®, Monster®, Red Bull®, Power-up™, Original Energy Drink, Rockstar Energy drink™, Emerge, No Fear™ and Boost energy drink. The only “other” sports drink was an unbranded isotonic drink.

Location of purchase

The purchase locations for the sports drinks are presented in Figure 3; there were 242 locations mentioned by 143 respondents who reported drinking them, allowing multiple purchase location responses.

Most of the sports drinks were being purchased in local shops, with 115 (80.4% of respondents). Supermarkets were also popular with 78 mentions, which equated to 54.5% of respondents. The four mentions of “Other” were; bakery, Home Bargains, Spar and Lidl.

Context of consumption

Respondents were asked in which situation they drank sports drinks limited to 5 categories which were isolated during the focus group discussion (Figure 4).

There was a clear gender difference, far more boys than girls consumed sports drinks during physical activity and at meal times (Figure 4). One respondent failed to state their gender on the questionnaire resulting in n=142 for this analysis. The majority of boys (77.9%, 53 respondents out of 68) reported drinking these products during physical activity compared with only 48.6% (36 respondents out of 74) of girls ($p < 0.001$). However, more girls drank sports drinks at home (41.9% of girls compared with 32.4% of boys) and socially (51.4% of girls compared to 48.5% of boys); neither of these results were statistically significant ($p > 0.05$). These results reflect responses to respondent’s participation in sport. 86.8% (59/68) of male respondents participated in sport compared with 71.6% (53/74) of female respondents.

Reasons for consuming sports drinks

The most popular reason for consuming sports drinks was the taste of the product (90.2% 129/143). Energy and hydration were reasons given by 47.6% (68/143) and 23.1% (33/143) of respondents respectively. The fact that sports drinks enhance performance was only stated by 18.2% (26/143) as a reason for consumption (Figure 5).

Approximately one third (32.9%, 47/143) of respondents stated price as a reason for purchasing sports drinks (Figure 5). When questioned how much they would spend, the majority would spend between 50p and £1 for a sports drink (55.9% of respondents). 38.5% would pay over £1 for a sports drink.

DISCUSSION

The results obtained clearly show a high proportion of adolescents in South Wales are consuming sports drinks.

The reported prevalence of sports drinks consumption for 12-14 year olds participating in this study was high at 89%, with 68% of these children drinking them regularly (1-7 times a week). Both the prevalence and the frequency of consumption in this age group appear to have increased in recent years^{29, 30}.

In the late 1990s a study of 418 14 year old secondary school children in Birmingham, recorded a prevalence of 44%, with 77% of these consuming sports drinks 1-7 times per week²⁹. Whilst a larger UK study in the North West of England, of 2,385 14 year old children in 1999, reported 81% drinking sports drinks occasionally or regularly³⁰.

Mintel market analysis data (52 weeks ending 1 March 2014) indicated that Lucozade™ represents 63% of the sports drink market, followed by Powerade™ taking 14% and own label and others taking 23%¹. This was reflected by the younger population of this study where Lucozade™ and Powerade™ predominated. However just under half of the sample reported drinking LSV® which was not separately identified by the Mintel data - this may be a reflection of the fact that LSV® is labelled as an Energy Drink but marketed as

“a functional beverage, specially developed for periods of increased mental and physical exertion”³¹.

When questioned why they drink sports drinks, 90% claimed the taste was a factor where only 18% of the respondents claimed it was due to the performance enhancing effect. This mirrors findings of the US Healthy Eating Research Review in 2012² and Food Standards Australia New Zealand in 2010³², which found people were consuming these drinks simply because they tasted nice. Taste of these sports drinks was the primary factor given by this group of 12-14 year olds and appears to be the main reason why they appeal to younger consumers. Improvements in physical performance was given as a reason for drinking by a minority of participants, even though over 71.6% of girls and 86.8% of boys in this study claimed to be taking part in exercise.

The results also showed that about a half of the respondents who drink sports drinks claim to drink them socially. This is likely to be linked to the fact that consumer independence is well developed and also the fact that the influence of peers (whilst socialising) is particularly strong

for this age-group³³. This has implications for dental and wider public health, in terms of dental caries, dental erosion and obesity.

Most of the children (80%) purchased sports drinks in local shops, once again highlighting the influence of stores surrounding the school fringe in the availability of “less healthy” foods and beverages, including sports drinks – often available at value prices^{34,35}. Indeed, price itself was the third most recorded reason for purchase. Of concern, 26% and 6% of children cited leisure centres and schools as purchase sources. In the UK in recent years holistic food and health policies^{35,36} have come to the fore, where local government, health and third sector agencies work together to address food, health and exercise. These can involve whole school approaches to food and health and the promotion of healthier choices in leisure centres³⁷. However, these findings appear to suggest that there is room for improvement, similar to experiences in other parts of the UK³⁸.

Whilst the majority of participants claimed to drink sports drinks, it became apparent that there was some confusion over the definition of a sports versus an energy drink. However from a dental and wider health perspective these two drinks types have similar detrimental effects due to their high free sugar (unless sugar free) content and low pH.

In supermarkets and local shops, sports drinks are sold alongside other sugar sweetened beverages. This could indicate to children and parents that they are meant for use by everyone. Where these drinks are sold in shops should be re-assessed, so as to ensure people do not misunderstand the purpose of the product. The prices of these drinks also needs to be considered; the recent lobby by Public Health England³⁹ and others for a sugar tax on sugar sweetened beverages was successful in March 2016⁴⁰.

The fact that sports drinks are so popular with children because of their sweet taste as opposed to reasons associated with sport (which are tenuous) should add weight to the case for an excise duty on sweetened beverages incorporating sports drinks.

CONCLUSION

A high proportion of children consumed sports drinks regularly and outside of sporting activity. Dental health professionals should be aware of the popularity of sports drinks with children when giving health education advice or designing health promotion initiatives.

REFERENCES

1. Mintel. Sports and energy drinks – Executive Summary: UK. July 2014. Mintel Group Ltd: London UK.
2. Healthy Eating Research. Consumption of Sports Drinks by Children and Adolescents. 2012. Online information available at <http://healthyeatingresearch.org/wp-content/uploads/2013/12/HER-Sports-Drinks-Research-Review-6-2012.pdf> (accessed September 2015).
3. Cohen D. The truth about sports drinks. *BMJ*. 2012; **345**: e4737.
4. Noble W, Donovan T, Geissberger M. Sports drinks and dental erosion, *CDA Journal*. 2011; **39** (4): 233-238.
5. Schnieder MB, Benjamin HJ. Sports drinks and energy drinks for children and adolescents: are they appropriate? *Pediatrics*. 2011; **127** (6): 1182-1189.
6. Heneghan C, Howick J, O'Neill B, Gill PJ, Lasserson D, Cohen D, et al. The evidence underpinning sports performance products: a systematic assessment. *BMJ* 2012; **2**:e001702.
7. Department of Nutrition at Harvard School of Public Health. Sugary Drinks and Obesity Fact Sheet. 2012. Online information available at <http://www.hsph.harvard.edu/nutritionsource/sugary-drinks-fact-sheet/> (accessed October 2012).
8. Burt BA, Pai S. Sugar consumption and caries risk: A systematic review. *J Dent Educ*. 2001; **65** (10): 1017-1023.
9. Harris R, Nicoll A, Adair P, Pine C. Risk factors for dental caries in young children: a systematic review of the literature. *CDH*. 2004; **21S**: 71–85.
10. Ruxton CH, Gardner EJ, McNulty HM. Is sugar consumption detrimental to health? A review of the evidence 1995-2006. *Crit Rev Food Sci Nutr*. 2010; **50**(1):1-19.

11. World Sugar Research Organisation. Sugar and Dental Caries. 2011. Online information available at <http://www.wsro.org/Portals/12/Docs/position-statement-sugar-and-dental-caries-2011-11.pdf> (accessed October 2012).]
12. Scientific Advisory Committee on Nutrition. Carbohydrates and Health. 2015 TSO: London.
13. Sirimaharaj V, Brearley Messer L, Morgan MV. Acidic diet and dental erosion among athletes. *Aust Dent J*. 2002; **47**(3): 228-236.
14. Rees J, Loyn T, McAndrew R. The acidic and erosive potential of five sports drinks. *Eur J Prosthodont Restor Dent*. 2005; **13**: 186-190.
15. Burt BA, Eklund SA, Morgan KJ, et al. The effects of sugars intake and frequency of ingestion on dental caries increment in a three-year longitudinal study. *J Dent Res*. 1988. **67** (11): 1422-1429.
16. Moynihan PJ, Kelly SAM. Effect on Caries of restricting sugar intake: systematic review to inform WHO guidelines. *J Dent Res*. 2013; **93**(1): 8-18.
17. Needleman I, Ashley P, Petrie A. et al. Oral health and impact on performance of athletes participating in the London 2012 Olympic Games: a cross-sectional study. *Br J Sports Med*. 2013;**47**:1054–8
18. White D, Pitts N, Steele J, Sadler K, Chadwick B. Disease and related disorders – a report from the Adult Dental Health Survey 2009. London: The Health and Social Care Information Centre, 2011.
19. Pitts N, Chadwick B Anderson T. Children’s Dental Health Survey 2013. Report 2: Dental Disease and Damage in Children, England, Wales and Northern Ireland. London: The Health and Social Care Information Centre, 2015.

20. Watt RG, Listl S, Peres M, Heilmann, A. Social inequalities in oral health: from evidence to action. International Centre for Oral Health Inequalities Research & Policy. 2015. Online information available at http://media.news.health.ufl.edu/misc/cod-oralhealth/docs/posts_frontpage/SocialInequalities.pdf (accessed September 2015).
21. British Soft Drinks Association. 2013. Code of practice on high caffeine soft drinks. Online information available at <http://www.britishsoftdrinks.com/PDF/130924%20high%20caffeine%20soft%20drinks.pdf> (accessed October 2013).
22. International Olympic Committee Report. Sports or energy drinks? 2002. Online information available at http://www.olympic.org/Documents/Reports/EN/en_report_539.pdf (accessed October 2012).
23. Scottish Executive. Hungry for Success: A Whole School Approach to School Meals in Scotland. Edinburgh: Scottish Executive. 2002.
24. School Meals Review Panel. Turning the Tables: Transforming School Food London: Department for Education and Skills. 2005.
25. Welsh Assembly Government. Appetite for Life Action Plan. 2008 Online information available at <http://new.wales.gov.uk/dcells/publications/publications/guidanceandinformation/apetiteforlife/appetiteforlifeactionplane.pdf?lang=en> (accessed September 2015).
26. Health Promotion Agency for Northern Ireland School Food: Top Marks! Initiative. 2009. Online information available at http://www.healthpromotionagency.org.uk/work/Publicrelations/PressReleases/school_food_09.html (accessed September 2015).
27. Cardiff University School of Dentistry. Welsh Index of Multiple Deprivation. 2011. Online information available at <http://wales.gov.uk/statistics-and-research/welsh-index-multiple-deprivation/?lang=en> (accessed October 2013)

28. Babbie E R. *The practice of social research*. 10th Edition. Belmont. California. 2004.
29. Al-Dlaigan YH, Shaw L, Smith A. Dental erosion in a group of British 14-year-old school children Part II: Influence of dietary intake. *BDJ*. 2001; **190** (5): 258-261.
30. Milosevic A, Bardsley P F, Taylor S. Epidemiological studies of tooth wear and dental erosion in 14-year old children in North West England. Part 2: The association of diet and habits. *BDJ* 2004; **197**: 479–483
31. LSV, 2015. What are the effects of LSV® Energy drink? Online information available at http://54chd3v4.wix.com/lsv#!_page-1 (accessed October 2013).
32. Food Standards Australia New Zealand Consumer research investigating the use of formulated supplementary sports foods. 2010. Online information available at <http://www.foodstandards.gov.au/publications/Pages/Consumer-research-investigating-the-use-of-sports-foods.aspx> (accessed August 2015).
33. World Health Organization. Nutrition in adolescence: issues and challenges for the health sector: issues in adolescent health and development. 2005. Online information available at http://apps.who.int/iris/bitstream/10665/43342/1/9241593660_eng.pdf accessed 11th September 2015 (accessed September 2015).
34. Sinclair S, Winkler JT. The School Fringe: what pupils buy and eat from shops surrounding secondary schools. Nutrition Policy Unit, London Metropolitan University. 2008.
35. Caraher M, Lloyd S, Madelin T. The “School Foodshed”: schools and fast-food outlets in a London borough. *BFJ*. 2014; **116** (3): 472-493.
36. Fairchild RM, Morgan MZ. Delivering multi-disciplinary public health in action – the Cardiff food strategy case study. *Public Health Nutr*. 2007; 10: 42-48.
37. Cardiff Food and Health Strategy. Cardiff Health Alliance: Cardiff. 2006. Online information available at

<http://www.physicalactivityandnutritionwales.org.uk/Documents/740/Cardiff%20Food%20and%20Health%20Strategy.pdf> (accessed September 2015).

38. Ennis K, Holt A, Cheater S. Sugar sweetened beverages availability and purchasing behaviour within the school fringe. *Int J Health Prom Educ.* 2014; **52** (5): 300-312
39. Public Health England. Sugar reduction: the evidence for action. 2015. Online information available at https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/470179/Sugar_reduction_The_evidence_for_action.pdf (accessed 21 April 2016)
40. Lords Hansard text for 03 February 2016. Column 1791-1793. Online information available at <http://www.publications.parliament.uk/pa/ld201516/ldhansrd/text/160203-0001.htm> (accessed 21 April 2016)

Table 1 Location and participation by school and year group

School	Location	Number of respondents			
		Year 8		Year 9	
		Total number in class	Number taking part	Total number in class	Number taking part
A	Cardiff	20	0	13	12 (7M, 5F)
B	Rhondda Cynon Taf	30	30 (15M, 14F, 1 MV)	30	30 (15M, 15F)
C	Rhondda Cynon Taf	21	21 (21M)	18	18 (18M)
D	Rhondda Cynon Taf	25	25 (25F)	26	24 (24F)
Total number of respondents		96	76	87	84

*M – male, F- female, MV – missing value

Figure 1 Reported frequency of sports drinks consumption, for the 143 respondents who reported drinking them




Figure 2 Drink type consumed, by the 143 respondents who reported drinking them (respondents could report drinking more than one type)





Figure 3 Location of purchase, identified by the 143 respondents who reported drinking sports drinks (respondents could report more than one location)



Figure 4 Context of consumption, identified by the 142* respondents who reported drinking sports drinks (respondents could report drinking in more than one context).



*1 missing value for gender

		Swhen_drink ^a					Total	
		when_physicalactivity	when_mealtime	when_school	when_home	when_socially		
gender	Male	Count	53	12	17	22	33	68
		% within gender	77.9%	17.6%	25.0%	32.4%	48.5%	
	Female	Count	36	4	18	31	38	74
		% within gender	48.6%	5.4%	24.3%	41.9%	51.4%	
Total		Count	89	16	35	53	71	140

^aPercentages and totals are based on respondents.

Figure 5 Reason for consumption, identified by the 143 respondents who reported drinking sports drinks (respondents could report more than one reason)

